

1 Satellite Passive Microwave Sea-Ice Concentration Data Set Inter- 2 comparison using Landsat data

3
4 Stefan Kern¹, Thomas Lavergne², Leif Toudal Pedersen³, and Rasmus Tage Tonboe^{4a}, Louisa Bell^{1b},
5 Maybritt Meyer^{1c}, and Luise Zeigermann^{1d}

6
7 ¹Integrated Climate Data Center (ICDC), Center for Earth System Research and Sustainability (CEN), University of Hamburg,
8 Hamburg, Germany

9 ²Research and Development Department, Norwegian Meteorological Institute, Oslo, Norway

10 ³Danish Technical University, Lyngby, Denmark

11 ⁴Danish Meteorological Institute, Copenhagen, Denmark

12 ^anow at Danish Technical University, Lyngby, Denmark

13 ^bnow at Climate Service Center Germany (GERICS), Helmholtz-Zentrum Hereon, Hamburg, Germany

14 ^cnow at Federal Maritime and Hydrographic Agency, Hamburg, Germany

15 ^dnow at Faculty of Science for Physics and Physical Oceanography, Memorial University of Newfoundland, St. Johns, Canada

16 *Correspondence to:* Stefan Kern (stefan.kern@uni-hamburg.de)

17 **Abstract.** We report on results of an inter-comparison of 10 global sea-ice concentration (SIC) data products at 12.5 to 50.0
18 km grid resolution from satellite passive microwave (PMW) observations. For this we use SIC estimated from > 350 images
19 acquired in the visible / near-infrared frequency range by joint the National Aeronautics and Space Administration
20 (NASA)/United States Geological Survey (USGS) Landsat sensor during years 2003-2011 and 2013-2015. Conditions covered
21 are late winter / early spring in the Northern Hemisphere and from late winter through fall freeze-up in the Southern
22 Hemisphere. Among the products investigated are the four products of the European Organisation for the Exploitation of
23 Meteorological Satellites (EUMETSAT) Ocean and Sea Ice Satellite Application Facility (OSI SAF) and European Space
24 Agency (ESA) Climate Change Initiative (CCI) algorithms: SICCI-2 and OSI-450. We stress the importance to consider inter-
25 comparison results across the entire SIC range instead of focusing on overall mean differences, and to take in account known
26 biases in PMW SIC products, e.g. for thin ice. We find superior linear agreement between PMW SIC and Landsat SIC for the
27 25 km and the 50 km SICCI-2 products in both hemispheres. We discuss quantitatively various uncertainty sources of the
28 evaluation carried out. First, depending on the number of mixed ocean-ice Landsat pixels classified erroneously as ice only,
29 our Landsat SIC is found to be biased high. This applies to some of our Southern Hemisphere data, promotes an overly large
30 fraction of Landsat SIC under-estimation by PMW SIC products, and renders PMW SIC products overestimating Landsat SIC
31 particularly problematic. Secondly, our main results are based on SIC data truncated to the range 0 % to 100 %. We demonstrate
32 using non-truncated SIC values, where possible, can considerably improve linear agreement between PMW and Landsat SIC.
33 Thirdly, we investigate the impact of filters often used to clean up the final products from spurious SIC over open water due
34 to weather effects and along coastlines due to land spillover. Benefiting from the possibility to switch on or off certain filters
35 in the SICCI-2 and OSI-450 products we quantify the impact land spillover filtering can have on evaluation results as shown
36 in this paper.

37 1 Introduction

38 We carry on the evaluation of sea-ice concentration (SIC) products derived from satellite passive microwave (PMW)
39 observations. In Kern et al. (2019), we presented an evaluation of ten PMW SIC products at 0 % and 100 % SIC, and with
40 respect to sea-ice observations along ship tracks. Another study focused on Arctic summer conditions, investigating the bias
41 between these PMW SIC products and independent SIC and net ice surface fraction estimates based on MODerate resolution
42 Imaging Spectroradiometer (MODIS) observations (Kern et al., 2020). With this study, we shift our focus more towards
43 intermediate SIC and utilize a much larger and, partly, more accurate reference dataset than in the two earlier studies. The

44 evaluation at 0 % SIC in Kern et al. (2019) utilized a few fixed open water locations only. The evaluation at 100 % SIC used
45 near-100 % SIC estimates based on the analysis of freezing-season synthetic aperture radar (SAR) image pairs representing
46 convergent ice motion coinciding with a complete ice coverage and therefore a high probability to encounter near-100 % SIC.
47 Thus, we evaluated the PMW SIC products for one specific set of ice conditions only (winter and near-100 %). Kern et al.
48 (2019) also presented results of an evaluation of PMW SIC against a multi-annual set of standardized manual visual ship-
49 based observations of the ice conditions. These observations are, however, of limited accuracy and of limited representativity
50 because the average accuracy is between 5 % and 10 % and observations mostly represent sea-ice conditions where it is
51 possible to navigate. In addition, to reduce noise, PMW and ship-based SIC were averaged over all observations along a ship-
52 track within one day, representing sea-ice conditions across spatial scales, that – in the worst case – vary by an order of
53 magnitude. The averaging resulted in a reduction of the number of valid data pairs from approximately 15 000 to less than
54 800, i.e. about 400 per hemisphere.

55 Another aspect is that the accuracy of the hemispheric but also the regional sea-ice area (SIA) computed from PMW
56 SIC estimates strongly depends on their accuracy. PMW SIC values biased high yield an overestimation of the SIA whereas
57 PMW SIC biased low results in an underestimation of the SIA. This seems not to be critical as long as the trend is correct (e.g.
58 Ivanova et al., 2014) but limits the use of such SIA estimates for quantitative inter-comparisons of climate-model results
59 against observations (e.g. Burgard et al., 2020). It is definitely important PMW SIC is 100 % where the actual SIC is 100 %
60 to avoid artificially elevated ocean-atmosphere heat fluxes when used as a surface forcing. It is equally important PMW SIC
61 is an accurate estimate of the open water fraction, hence providing 95 % where the actual SIC is 95 % due to leads and openings
62 in the sea-ice cover. In addition, it is desirable to check the performance of PMW SIC products across the entire SIC range in
63 order to have a reliable estimate of the actual ice cover in, for example, the marginal ice zone (MIZ). Here gradients in heat
64 fluxes are often particularly large. A correct definition of and accurate SIC distribution within the MIZ are also crucial should
65 SIC values be used to evaluate numerical models capable to simulate wave-sea ice interaction (e.g. Boutin et al., 2020; Nose
66 et al., 2020). The ship-based SIC observations used in Kern et al. (2019) offer only limited potential to carry out this
67 performance check because of the earlier-mentioned reasons, the small number of observations falling into the relevant SIC
68 range of, e.g. 20 % to 80 %, and the larger observational error in this SIC range.

69 Therefore, in this paper we focus on the evaluation of PMW SIC products against a large number of high-resolution
70 binary sea-ice cover maps estimated from satellite observations acquired in the visible frequency range by NASA/USGS
71 Landsat-5, 7 and 8. Overall, we used over 350 such Landsat-based maps, corresponding to more than 10 000 25 km x 25 km
72 resolution PMW SIC grid cells. We chose Landsat over MODIS because of the substantially finer spatial resolution of the
73 visible channels of Landsat: 30 m compared to MODIS: 250 m. We note in this context that several studies used MODIS
74 visible / near-infrared observations to either evaluate or complement PMW SIC products (e.g. Ludwig et al., 2020; Shi et al.,
75 2021). Another option would have been to use Sentinel-2 MultiSpectral Instrument (MSI) (Drusch et al., 2012). We discarded
76 this option in light of the limited overlap between this satellite mission (Sentinel-2A was launched June 2015) and our PMW
77 SIC data set but it will be very valuable in the future since it will allow extending the dataset to areas much further from land
78 and will likely provide an even more accurate evaluation data set.

79 Utilization of the high-resolution information provided by Landsat as a means for assessing satellite PMW SIC
80 products dates back to the early 1980s when Comiso and Zwally (1982) compared Nimbus-7 Scanning Multichannel
81 Microwave Radiometer (SMMR) SIC with Landsat imagery. Since then a number of studies used a small number of such
82 images for SIC inter-comparison and/or evaluation (e.g. Steffen and Maslanik, 1988; Steffen and Schweiger, 1991; Comiso
83 and Steffen, 2001; Cavalieri et al., 2006; Wiebe et al., 2009; Lu et al., 2018; Zhao et al., 2021). Landsat imagery has also
84 recently been used for quality assessment of SIC estimates from Suomi/NPP VIIRS observations (e.g. Liu et al., 2016).
85 Common to all these studies is they used a comparably small number of Landsat scenes, i.e. less than ten, an order of magnitude
86 smaller than the 368 scenes used in this study (see above).

87 Analysis of visible satellite imagery for SIC estimation is quite straightforward. A threshold based method
88 discriminating between open water and ice is applied at the native spatial resolution (pixel size: 30 m x 30 m) of the Landsat
89 channels in the visible frequency range, assuming that a pixel is covered by either ice or water. Co-locating this high-resolution
90 information of the binary ice-water distribution with the coarse-resolution PMW SIC products and counting ice and water
91 pixels within a PMW SIC product's grid cell provides an adequate independent measure of the SIC. We refer to Section 2.2
92 for more details.

93 For evaluating the PMW SIC products across the SIC range, we prefer to use visible data instead of SAR data. The
94 main advantages of SAR data would be the larger area covered by a single scene compared to Landsat (about 400 km to 500
95 km in SAR wide-swath mode (WSM) vs. 180 km for Landsat) and their independence to daylight and cloud cover. In fact,
96 many PMW SIC inter-comparison studies have already used SAR images (e.g., Comiso et al., 1991; Dokken et al., 2000;
97 Belchansky and Douglas, 2002; Kwok, 2002; Heinrichs et al., 2006; Andersen et al., 2007; Wiebe et al., 2009; Han and Kim,
98 2018). However, despite the past decade's substantial progress in developing and testing methods to translate SAR images into
99 high-resolution SIC maps (e.g.: Cooke and Scott, 2019; Karvonen, 2014, 2017; Komarov and Buehner, 2017, 2019; Leigh et
100 al., 2014; Lohse et al., 2019; Ochilov and Clausi, 2012; Singha et al., 2018; Wang et al., 2016, 2017; Zakhvatkina et al., 2017,
101 Boulze et al., 2020; Malmgren-Hansen et al., 2020; Wang and Li, 2020), some using machine learning approaches, the accuracy
102 of the obtained SIC maps is not always satisfactory. Particularly at intermediate SIC – the main focus of this study – SAR
103 signatures are often ambiguous, resulting in SAR SIC uncertainties too large for our purposes. Furthermore, applications of
104 such methods to derive Southern Ocean SIC from SAR are comparably sparse. Therefore, we do not use SAR-based SIC maps.

105 We note that also Ice charting services (FMI, DMI, MET Norway, CIS, NATICE, AARI) heavily depend on SAR
106 imagery for production of their ice charts. They thus have a large demand to automate processes of classification and are
107 potentially most advanced in testing automated SAR SIC retrieval (e.g. Cheng et al., 2020). However, ice charts provide SIC
108 ranges within polygons that are highly variable and heterogeneous in size and shape. Several studies used such ice charts for
109 various inter-comparison purposes (e.g. Shokr and Markus, 2006; Shokr and Agnew, 2013, Titchner and Rayner, 2014). Some
110 centers providing operational sea-ice information also use such charts for routine quality checking of PMW SIC products.
111 However, for our purpose evaluating PMW SIC CDRs and similar SIC products, the limitations of such charts in terms of
112 precision and accuracy – particularly in the intermediate SIC range (e.g. Cheng et al., 2020), exclude their usage in this study.

113 After this introduction, this paper provides information about the PMW SIC products, the Landsat data set used and
114 the methods applied to derive SIC from the Landsat images (Sect. 2). We present our results in Sections 3 and 4, discuss some
115 additional aspects in Section 5 and conclude the study in Section 6.

116 2 Data & Methodologies

117 2.1 Sea-ice concentration data sets

118 The ten different PMW SIC products considered in our study are summarized briefly in Table 1. We refrain from
119 repeating information about the algorithms themselves, tie point selection, application of weather filters, consideration of land
120 spillover effects and so forth. All this information is provided in detail in Lavergne et al. (2019), Kern et al. (2019, Appendix
121 7.1-7.6), and Kern et al., (2020). The same applies to the fact that four of the products (SICCI-12km, SICCI-25km, SICCI-
122 50km, and OSI-450) allow us to take into account the full SIC distribution at 0 % and 100 %. Such a distribution is the natural
123 result of the SIC retrieval method used in all SIC products considered - except NT2-AMSR. This distribution contains negative
124 as well as above-100 % SIC values that are typically truncated, i.e. set to the exactly 0 % and 100 %. We refer to Lavergne et
125 al. (2019) and Kern et al. (2019) for more information in this regard.

126 In order to extend the time-series of the Comiso Bootstrap (CBT) algorithm and the NASA-Team 2 (NT2) algorithm
127 using Advanced Microwave Scanning Radiometer aboard Earth Observation Satellite (AMSR-E) data beyond AMSR-E's

capabilities to provide daily maps of the polar regions (2011-10-03), we use the respective unified product based on data from the Advanced Microwave Scanning Radiometer aboard GCOM-W1: AMSR2 and of AMSR-E (Meier et al., 2018). With that we use five products based on AMSR-E and AMSR2 data and five products based on Special Sensor Microwave / Imager: SSM/I, and Special Sensor Microwave Imager and Sounder: SSMIS data, of the period 2002 through 2015. We do not use PMW SIC data from the period October 2011 through July 2012 because of the gap between AMSR-E and AMSR2. All PMW SIC data have daily temporal resolution. The grid type and grid resolution of all datasets is shown in Table 1. We estimate the Landsat SIC (see Section 2.2) at the grid resolution of the respective product. We chose the 25 km grid resolution version of the AMSR-E/2 products because this resolution is closer to the footprint sizes of the involved channels, and this is the resolution of the respective SSMI(S) versions of these products. We use version 3 of the NOAA/NSIDC SIC CDR (Peng et al., 2013; Meier et al., 2017) even though version 4 has been released (Meier et al., 2021) because we want to be consistent to the two previous papers (Kern et al., 2019; 2020).

2.2 The Landsat data set

We use Landsat data of the Thematic Mapper TM on Landsat-5, the Enhanced Thematic Mapper (ETM) on Landsat-7, and the Operational Land Imager (OLI) on Landsat-8 obtained in Level 1c GeoTIFF format from <https://earthexplorer.usgs.gov> [last accessed: June 28, 2021] for years 2003-2011 (Landsat-5), 2003 (Landsat-7), and 2013-2015 (Landsat-8). We downloaded only images with a cloud fraction < 30 % provided as a search criterion upfront. In the Northern Hemisphere, we use images of months March, April, May and September, i.e. from late winter to spring and at the onset of fall freeze-up; in the Southern Hemisphere we use images of months October through March, i.e. from late winter over summer to fall freeze-up. The total number of images acquired is 421; these split into 152, 12, and 227 for Landsat-5, 7 and 8, respectively, and partition into 259 images for the Northern Hemisphere and 162 images for the Southern Hemisphere.

2.2.1 Processing

We compute the top of atmosphere (TOA) reflectance for channels 2 to 4 (Landsat-5 and 7) or channels 3 to 5 (Landsat-8) following Chander et al. (2007; 2009) and USGS (2015). Table 2 provides the wavelengths of these channels (e.g. Chander et al., 2009; Barsi et al., 2014). The solar zenith angle and other parameters required for this computation are either included in the Landsat data files or are taken from Chander et al. (2007, 2009) and the Landsat 8 data users handbook (USGS, 2015). To convert the TOA reflectances to surface reflectances or surface albedo we follow the approaches of Koepke (1999) and Knap et al. (1999). They assume that the TOA reflectance (or planetary reflectance) equals the TOA albedo (or planetary albedo) and that the TOA albedo α_{TOA} is related to the surface albedo $\alpha_{surface}$ via the simple linear relationship:

$$\alpha_{TOA} = a + b\alpha_{surface} \quad (1)$$

The coefficients a and b are a function of the atmospheric conditions, the solar zenith angle, and the wavelength. We follow Koepke (1999) and take values for a and b from his figure 1 (KF1) and figure 2 (KF2). KF1 derived for the Advanced Very High Resolution Radiometer (AVHRR) channel 1 we use for Landsat channels in the wavelength range 500-700 nm. KF2 derived for AVHRR channel 2 we use for Landsat channels in the wavelength range 700-900 nm. We choose those atmospheric conditions that are appropriate for a polar marine atmosphere. For aerosol optical depth we use 0.05, for ozone content we use 0.24 cm[NTP] (NTP stands for normal temperature and pressure) corresponding to 240 Dobson Units, and for water vapor content we used 0.5 g/cm². Using Eq. (1) we convert TOA albedo into surface albedo values separately for the three channels of the respective Landsat instrument. Subsequently, we compute from these surface albedo values an estimate of the surface broadband shortwave albedo (e.g. Brandt et al., 2005) using the bandwidths of the channels as weights (see Table 2).

For every broadband surface albedo map, we perform a supervised visual classification into open water, bare / thin ice and snow covered / thick ice. For that, we assume the respective surface class covers a Landsat pixel entirely. We assign all dark pixels (with an albedo of, on average, smaller than 0.06) to the open water class. We assign all bright pixels (with an

170 albedo of, on average, larger than 0.45) to the class snow covered / thick ice; all remaining pixels fall into the class bare / thin
171 ice. We pay more attention separating open water from ice very accurately than to distinguish between bare / thin ice and
172 snow-covered / thick ice. In every Landsat albedo map we search for leads or openings, zoom into these and perform histogram-
173 equalized slicing to visually identify – based on albedo values and spatial structures – whether the leads or openings selected
174 contain open water. The threshold value chosen to separate open water from ice we take from Pegau and Paulsen (2001). The
175 threshold value chosen to distinguish between bare / thin ice and snow covered / thick ice is based on Brandt et al. (2005) and
176 Zatko and Warren (2015). They found an albedo of around 0.33 for bare thin ice less than 30 cm thick and of around 0.42 for
177 snow covered thin ice (5 - 10 cm thick) with a thin (< 3 cm) snow cover. Note that the actual threshold values chosen for a
178 particular Landsat image varies between 0.03 and 0.08 for the open water – ice discrimination and between 0.35 and 0.55 for
179 the bare / thin ice – snow covered / thick ice discrimination. This variation results from the varying illumination conditions
180 encountered – despite our limitation to Landsat scenes acquired at solar zenith angles < 65°.

181 Usage of a three-class distribution is motivated by the fact that it has been shown that PMW SIC is often biased low
182 over thin sea ice (e.g. Wensnahan et al., 1993; Cavalieri, 1994; Ivanova et al., 2015). Therefore, in addition to using the Landsat
183 images just for a high-resolution ice-water discrimination we also use them to derive the fraction of thin ice with the aim to
184 discuss differences between Landsat SIC and PMW SIC in the light of a potential impact by thin ice. However, we discarded
185 this aim – but kept the classification results – because during analyses of the Landsat images we encountered ambiguities in
186 surface albedos between snow-covered thin ice and bare thick ice. While there is little ambiguity between open water and ice,
187 except for very thin dark nilas or ice rind (e.g. Zatko and Warren, 2015), resulting in high confidence of pixels classified as
188 either open water or ice, the confidence of pixels classified as bare/thin or snow covered/thick ice is considerably worse.

190 **2.2.2 Co-location and comparison**

191 For the co-location, we first select a rectangular area within the PMW SIC grid, EASE-2 for the SICCI-2 and OSI-
192 450 products (EPSG: 6930 and 6931) and polar-stereographic true at 70 degrees northern or southern latitude (known as
193 NSIDC grid, EPSG: 3411) for the other six products, which encloses the Landsat SIC map. For this we take the geographic
194 corner coordinates of the Landsat SIC map (still at 30 m grid resolution), convert these into Cartesian Coordinates and find
195 those PMW SIC grid cells which centers have minimum distance (in meters) to these corner coordinates. Beforehand, we also
196 convert PMW SIC grid cell coordinates into Cartesian coordinates and rotate the grid for the Northern Hemisphere PMW SIC
197 products on the NSIDC grid clockwise by 45 degrees; this is not required for the respective Southern Hemisphere PMW SIC
198 products.

199 Subsequently, we compute the Landsat SIC by summing over all 30 m pixels classified as ice that fall into the PMW
200 SIC grid cells within the above-defined rectangular area. Because we do this is at the grid resolution of the PMW products, we
201 obtain Landsat SIC maps at 12.5 km, 25.0 km, and 50.0 km grid resolution. We compare the resulting gridded Landsat SIC
202 with the respective co-located PMW SIC by computing the mean difference PMW SIC minus Landsat SIC and its standard
203 deviation, the median difference, and deriving a linear regression line and computing the linear correlation coefficient.

204 Based on a visual quality check of the obtained Landsat SIC maps we discard ~50 of the processed Landsat scenes
205 from further analysis – mainly because of cloud artifacts but also because a few scenes we obtained twice. Therefore, the
206 resulting lower final number of Landsat SIC maps used is 368: 234 for the Arctic, partitioning into Landsat-5: 134, Landsat-
207 7: 12, and Landsat-8: 88, and 134 for the Antarctic. The spatial distribution of the Landsat scenes is illustrated in Fig. 1. Note
208 that we focus on data of Landsat-5 and Landsat-8 in this paper.

209 **2.2.3 Sensitivity analysis**

211 In order to estimate how Landsat SIC depends on the choice of the albedo thresholds used to discriminate open water
212 from ice and bare / thin ice from snow covered / thick ice, we repeat the classification into the three surface classes using

modified thresholds. We vary the albedo value for the open water – ice discrimination by ± 0.03 , i.e. for an actual albedo value of 0.06 we employ additional threshold values of 0.03 and 0.09. We vary the albedo value for the bare / thin ice – snow covered / thick ice discrimination by ± 0.1 , i.e. for an actual albedo value of 0.45 we employ additional threshold values of 0.35 and 0.55. The range of albedo threshold values we choose is motivated by our experience with the supervised classification of the many Landsat scenes under varying illumination conditions. We randomly select 12 Landsat 8 scenes for the Northern Hemisphere, and 15 scenes for the Southern Hemisphere. For every image we perform the classification into the three surface classes with the above-mentioned four additional albedo threshold value combinations, compute Landsat SIC on the 25 km and 50 km EASE grid and derive a Landsat scene mean SIC value (Tables 3 and 4). We find that changing the albedo value of the open water – ice discrimination by ± 0.03 changes the average Landsat SIC by between 0.7 % and 1.2 % in the Northern Hemisphere and by between 0.8 % and 1.5 % in the Southern Hemisphere. Thus, the sensitivity appears to be independent of the overall SIC which is close to 100 % for the Northern Hemisphere cases (Table 3) but 55 – 60 % for the Southern Hemisphere cases (Table 4). The difference in the sensitivity between grid resolutions of 25 km and 50 km is less than 0.2 %.

As expected, changing the albedo value of the bare / thin ice – snow-covered / thick ice discrimination by ± 0.1 does not influence the Landsat SIC. However, it influences the Landsat SIC computed at the respective grid resolutions when using Landsat pixels classified as snow-covered / thick ice only (Tables S02 and S03 in the Supplementary Material). We find Landsat SIC of thick ice to vary by between 1.4 % and 2.4 % in the Northern Hemisphere and by between 2.1 % and 2.7 % in the Southern Hemisphere with little difference between the grid resolutions.

2.2.4 Potential biases in Landsat SIC

In our approach, we assume either ice or water to cover a Landsat pixel (30 m x 30 m) entirely, not taking into account that ice floes or leads / openings might be smaller than the pixel size, resulting in a mixed ocean-ice pixel. This can introduce a positive bias in the Landsat SIC computed at the grid resolution of the PMW SIC products. For instance, for a Landsat pixel covered just half by snow covered / thick sea ice, which exhibits a surface albedo of 0.8 under cold conditions, the resulting pixel average albedo would be $0.5 \times 0.06 + 0.5 \times 0.8 = 0.43$. With that, such a pixel is classified as bare / thin ice and counts as a pixel with 100 % instead of 50 % sea-ice concentration. Depending on the albedo of the ice, an ice-cover fraction of 0.04 in one Landsat pixel could be sufficient to increase the pixel average albedo above the upper open water – ice discrimination threshold value of 0.09 (see Tables 3, 4), causing the respective pixel to be classified as 100 % ice.

In order to quantify this positive bias better, it is useful to distinguish between sea-ice conditions during summer and winter, between pack ice and the MIZ, and to take into account the dimensions of leads / openings and ice floes. Distributions of lead width and floe size both follow a power law. Leads / openings and ice floes with dimensions smaller than the Landsat pixel size are orders of magnitude more abundant than wide leads / openings (e.g. Tschudi et al., 2002; Marcq and Weiss, 2012) and large ice floes (e.g. Steer et al., 2008; Toyota et al., 2011; Perovich and Jones, 2014).

Based on airborne digital camera visible imagery captured along several thousands' of kilometers long tracks of Operation Icebridge (OIB) flights in the Arctic in April 2010 and in the Antarctic in October 2009 analyzed by Onana et al. (2013) with respect to the lead and open water fraction, we find a SIC bias of less than 0.2 %. This value derived for an open water fraction of ~ 1 % falls into the uncertainty range of our approach (see Tables 3, 4) and represents winter conditions in the pack ice. Based on manual visual analysis of airborne visible imagery obtained in the MIZ in the Greenland Sea in March 1997, we find a SIC bias of the order of 5 to 10 %. This value is clearly outside the uncertainty range of our approach. The images used here represent an ice cover of ~ 70 % SIC comprising closely packed but also broken bands of a few thicker ice floes, pancake ice, brash and grease ice with little or no new ice formation in the openings – a typical situation at an ice edge located in comparably warm water.

Next, we again take the results of Onana et al. (2013) but assume that the thin ice identified in the OIB digital camera imagery adds to the open water fraction thereby simulating a summer situation. For an open water fraction of then ~ 5 %, we

256 estimate a SIC bias of less than 0.8 %, which is still within the uncertainty range of our approach. However, this low positive
257 bias during summer would only apply to a situation where ice floes are still packed closely together, e.g. by herding of ice
258 floes (e.g. Toyota et al., 2016), and where gaps between the ice floes from additional openings created by the melt process are
259 filled by brash ice and/or slush. While this is a situation that might be encountered during summer (Steer et al., 2008; Lu et al.,
260 2008), it is not necessarily typical. In summer, it can be more common to encounter isolated floes. Depending on the size of
261 the floes and their distribution across a 25 km grid cell with, e.g., 50 % SIC, we find the bias to range between less than 2 %
262 to 50 % in the two most extreme cases. We refer to the Supplementary Material to this subsection, where we describe in more
263 detail how we obtain estimates of the positive bias caused by the combination of the finite resolution of the Landsat sensor and
264 our classification approach for both winter and summer conditions at the scale of a 25 km PMW SIC product grid.

265 According to the high-resolution optical images used to infer the floe size distribution (Steer et al., 2008; Toyota et al., 2011; 2016) and similar studies (e.g. Paget et al., 2001; Lu et al., 2008; Zhang and Skjelne, 2015), the ice cover often
266 comprises a large spectrum of floes. The larger and largest floes at the upper end of the floe-size distribution form the major
267 fraction of the sea-ice area (in square kilometers) (e.g. Paget et al., 2001; Steer et al., 2008). A small number of large floes
268 results in a smaller number of mixed ocean-ice Landsat pixels than a large number of smaller floes. Hence, where larger floes
269 dominate, our Landsat SIC estimate is less biased than where small floes dominate. The effect of the ocean swell, the
270 dominating force for fracturing ice floes according to, e.g., Toyota et al. (2016), is larger close to the ice edge than further
271 inside the ice pack. Therefore, a larger number of smaller floes exists along the ice edge, suggesting a larger bias in our Landsat
272 SIC near the ice edge than inside the ice pack. Without further independent information about the actual ice cover, we are not
273 able to quantify this bias accurately.

274 Thus, for high-concentration winter conditions and for those cases during summer where ice floes are closely packed
275 and openings between the floes are covered with brash ice and slush, the bias in Landsat SIC derived at the spatial scale of the
276 PMW SIC products falls within the retrieval uncertainty range of our approach (see Tables 3, 4). The bias could fall outside
277 the uncertainty range near the ice edge during winter when sea ice drifts into comparably warm waters that inhibit ice formation
278 in newly created openings; here biases as high as 10 % in a single PMW grid cell could occur. The bias could also fall outside
279 the uncertainty range during summer; here biases between 5 % and 20 % in single PMW grid cells might occur depending on
280 proximity to the ice edge and hence floe-size distribution and depending on conditions favoring / inhibiting herding of ice floes
281 into bands.

283 3 Results

284 In the following, we present and discuss results obtained in the Northern and Southern Hemisphere. We preferred to
285 not merge the results of Landsat-5 and Landsat-8 in the Northern Hemisphere because with that we have a relatively natural
286 discrimination between cases dominated by first-year ice (Landsat-5) and cases dominated by mixed first-year / multiyear ice
287 or multiyear ice (Landsat-8) (see Fig. 1).

288 3.1 Northern Hemisphere

289 Out of the ten products, SICCI-25km, SICCI-50km, ASI-SSMI, and SICCI-12km offer the best linear agreement with
290 Landsat SIC for first-year ice dominated cases as expressed, e.g., by the location of mean and median PMW SIC (red symbols)
291 in Fig. 2 and the values of slope, intercept and correlation coefficient listed in Table 5. The two CBT products, NOAA-CDR
292 and NT2-AMSRE have the smallest overall mean difference and zero median (Table 5). These four products exhibit, however,
293 a considerable tail of near-100 % PMW SIC values stretching across almost the entire Landsat SIC range, pointing towards
294 over-estimation of Landsat SIC. ASI-SSMI and NT1-SSMI SIC exhibit the overall largest underestimation of Landsat SIC
295 among the ten products (Table 5).

296 For cases with mixed first-year / multiyear or multiyear ice, SICCI-25km and SICCI-50km offer best linear agreement
297 with Landsat SIC (Fig. 3). Most other products have a less convincing linear relationship. Like for first-year ice, the two CBT
298 products, NOAA-CDR and NT2-AMSR have the smallest mean difference for mixed first-year / multiyear or multiyear ice
299 (Fig. 3, Table 6). However, particularly at higher Landsat SIC these products show many data pairs above the identity line and
300 the linear regressions through the mean and median PMW SIC (red dashed and solid lines) are also located above the identity
301 line – in contrast to, e.g. SICCI-25km and SICCI-50km.

302 The linear agreement between PMW SIC and Landsat SIC improves in general for all ten products for mixed first-
303 year / multiyear or multiyear ice cases (Fig. 3, Table 6) compared to first-year ice (Fig. 2, Table 5). This improvement is
304 comparably large for OSI-450: slope increases by ~ 0.10 and NT2-AMSR: slope increases by ~ 0.15 but quite small for SICCI-
305 25km and SICCI-50km because slopes are close to unity already. Hence, despite the larger magnitude of overall mean and
306 median SIC differences, of all ten products SICCI-25km and SICCI-50km provide SIC estimates for first-year ice that are
307 almost as accurate as those for mixed first-year ice / multiyear ice or multiyear ice. This could be one consequence of the self-
308 optimizing hybrid SICCI-2 / OSI-450 algorithm (Lavergne et al., 2019) and of the way ice tie points are chosen in comparison
309 to the other products (e.g., Kern et al. 2020).

310 3.2 Southern Hemisphere

311 In the Southern Hemisphere, slope and location of the linear regression lines as well as of the mean and median PMW
312 SIC values (red symbols) is more similar between the ten products (Fig. 4, Table 7). The linear agreement is fairly good for
313 SICCI-2 products, CBT-AMSR2 and ASI-SSMI. Like in the Northern Hemisphere, SICCI-25km and SICCI-50 km reveal the
314 best linear agreement with Landsat SIC but SICCI-50km appears to be negatively biased. This bias is associated with a large
315 number of PMW SIC values of 0 % at non-zero Landsat SIC which is also reflected by the mean and median PMW SIC
316 (compare Fig. 4c) with Fig. 3c)). We discuss this issue and the observation that all products except CBT-SSMI, NOAA-CDR
317 and CBT-AMSR2 exhibit SIC values below about 10-15 % while these three products lack values in the PMW SIC range
318 between 0 % and ~ 15 % in Section 5.3.

319 Like in the Northern Hemisphere (Table 6), the magnitude of the SIC difference is smallest for NT2-AMSR2, NOAA-
320 CDR and the two CBT products and largest for NT1-SSMI and ASI-SSMI. Of all ten products, NT2-AMSR2 (Fig. 4 j) offers
321 the most asymmetric SIC distribution and a considerable overestimation of Landsat SIC in the range between ~ 40 % and ~ 90
322 %, also expressed by median SIC $>$ mean SIC for all Landsat SIC bins above 25 % (Fig. 4 j). NT2-AMSR2 is the only product
323 with a substantial positive overall mean difference of 3.4 %, even the median difference is > 0 % (Table 7).

324 3.3 Hemispheric Similarities and Differences

325 Overall, agreement between PMW SIC and Landsat SIC differs between the two hemispheres. For all products, we find
326 a substantially larger scatter of SIC values around the identity line in the Southern Hemisphere (section 3.2) than the Northern
327 Hemisphere (section 3.1). One the one hand, this larger scatter in the Southern Hemisphere could be the result of a considerably
328 larger number of Landsat scenes of cases with low SIC, naturally resulting in a larger spread of the SIC. In addition, the
329 majority of the Landsat scenes in the Southern Hemisphere reflect late spring / summer conditions. During such conditions,
330 snow metamorphism due to melt and melt-refreeze cycles substantially change the sea ice surface emissivity on daily time-
331 scales and sub grid-cell size spatial scales (e.g. Willmes et al., 2014) causing a larger scatter in SIC. Another factor impacting
332 the sea ice emissivity is flooding at the snow-ice interface and subsequent snow-ice formation, causing considerable variations
333 in basal snow layer wetness and salinity on similar spatiotemporal scales. On the other hand, we are dealing with an unknown
334 amount of overestimation of the actual sea-ice concentration by our Landsat SIC during summer melt due to mixed ocean-ice
335 Landsat pixels (Subsection 2.2.4). We refer to Sections 4.3, 5.1 and 5.2 for more discussion on this issue.

336 In general, we find the scatter is larger for products with finer grid resolution, e.g. SICCI-12km and ASI-SSMI, than
337 for the coarser grid-resolution products. The larger number of valid SIC pairs of the high-resolution products result in more
338 scatter due to the inherent retrieval noise – even though the capability to resolve smaller-scale SIC variations is better for the
339 fine- than the coarser-resolution products (see section 5.1). In addition, a mismatch in the location of, for example, a 10km-
340 scale patch of ice between a Landsat scene and a PMW SIC product has a substantially larger influence on the SIC difference
341 at 12.5 km than at 25 or 50 km grid resolution. The fact that oversampling is much larger at 12.5 km than at 50 km plays a role
342 here also. Even using simulated brightness temperatures one gets a large spread between a reference SIC and the PMW SIC
343 due to resolution mismatch (see e.g. Tonboe et al., 2016). We discuss the effect of different footprint sizes and grid resolutions
344 (see Table 1) in more detail in Section 5.1.

345 SICCI-2 products and OSI-450 provide access to SIC values above 100 % and below 0 % that are naturally retrieved
346 due to the brightness temperature distribution around ice and water tie points used. Kern et al. (2019) found that incorporation
347 of these so-called off-range or non-truncated SIC values provides a more accurate estimate of accuracy, i.e. difference to the
348 true SIC value, and precision, i.e. standard deviation of this difference. Table 8 reveals that independent of the ice type, the
349 magnitude of the mean difference decreases while the slope of the linear regression increases, becoming closer to unity in most
350 cases, in agreement with Kern et al. (2019). Of particular interest in this regard are high-concentration cases discussed in more
351 detail in Section 4.2 but also the effect of the truncation at 0 % in the context of filters used to mitigate spurious SIC values
352 (see Section 5.3).

353 4 Case Studies

354 In the previous section, we showed results independent of the ice regime (see below) – except for a general discussion
355 of the observed differences between predominantly first-year ice (Landsat-5) and a mixture of first-year / multiyear or
356 multiyear ice (Landsat-8). This section deals with our comparison between PMW SIC and Landsat SIC for the following ice
357 regimes: “ice edge”, “leads and openings” = cases with leads and coastal polynyas or openings, “heterogeneous ice” = cases
358 with irregularly shaped openings in the ice pack, “freeze-up”, “high-concentration ice”, and “melt conditions” (see Table S01
359 in the supplementary material). We show in more detail results of the last three ice regimes. Freeze-up cases are characterized
360 by a comparably large fraction of new and thin ice, an ice type for which some of the SIC products investigated here are already
361 known to be negatively biased from preliminary work based on Soil Moisture and Ocean Salinity (SMOS) thin ice thickness
362 observations (Ivanova et al., 2015). We elaborate on their findings using an alternative data set. Investigating high-
363 concentration cases in more detail aids in a better understanding of saturation effects near 100 % caused by truncating PMW
364 SIC at 100 %, expanding on the work of Kern et al. (2019), refining our knowledge of SIC precision and accuracy for high-
365 concentration regions and hence application potential of the products for surface heat flux computations. Finally, melt
366 conditions – even without melt ponds – represent a multitude of different snow and sea ice conditions causing enhanced
367 variability of the sea ice microwave emissivity (e.g. Willmes et al., 2014), which in turn can result in biases in PMW SIC
368 products of both signs in the Arctic (Kern et al., 2016; 2020). Here we have the opportunity to better quantify such biases
369 especially for the Antarctic. For all remaining regimes, we show examples in Figs. S03 through S08 in the supplementary
370 material and include their results of the statistical comparison into our summary figures (Fig. 11 and 12), but refrain from a
371 detailed discussion. For regimes “ice edge” and “leads and openings” such a discussion would require a comprehensive
372 investigation of open water and land spill-over filters (see Section 5.3) which is beyond the scope of this paper. For regime

373 “heterogeneous ice”, application of a more accurate evaluation SIC data set seems to be advisable given the identified
374 shortcomings of the used one (see Section 2.2.4) before going into more detail.

375 **4.1 Freeze-Up**

376 These are cases where according to the date, geographic location and information in the Landsat scene freeze-up has
377 commenced. We select Landsat scenes containing a considerable fraction of new and thin ice; these are acquired in September
378 and February/March in the Northern and Southern Hemisphere, respectively. We have only few such cases in both hemispheres
379 (see Table S01 in the supplementary material). We expect PMW SIC underestimates Landsat SIC (LSIC) – particularly for
380 young and thin ice with a thickness < 0.2 m (e.g. Ivanova et al., 2015). Figure 5 illustrates the conditions for a Landsat-8 scene
381 close to Greenland in the Fram Strait on September 15 2015. The classified Landsat-8 image (Fig. 5, top left) reveals a mix of
382 large ice floes – presumably second-year or older ice – and meandering patches of smaller floes embedded into a matrix of
383 mostly grey and a few dark pixels; the grey pixels are supposed to represent bare / thin sea ice, the dark pixels open water. All
384 products agree well with Landsat SIC in the topmost part of the scene over high-concentration ice. PMW SIC maps of six of
385 the ten products (SICCI-2 products, OSI-450, NT1-SSMI and ASI-SSMI) reveal an overall SIC distribution similar to Landsat
386 SIC. For the remaining four products, the SIC difference maps show widespread overestimation of LSIC by PMW SIC
387 expressed by positive (red) values. Contrary to expectations, we do not observe negative SIC differences for the entire greyish
388 area of the Landsat-8 scene.

389 The main reason for this observation is the actual ice condition. Very likely the greyish area represents a mixture of
390 sub-pixel size, i.e. less than 30 m x 30 m, ice floes and brash ice formed from disintegrated thicker ice floes and young / new
391 sea ice. On the one hand, the sub-pixel size floes and the brash ice are thicker than young / new sea ice. These forms of sea ice
392 exhibit different surface properties and hence microwave emissivity than young / new thin sea ice. For such a mixture of ice
393 types, it is particularly difficult to retrieve an accurate SIC with any of the algorithms used in the ten products. Ice tie points
394 do not adequately represent these ice conditions. On the other hand, for the greyish area the Landsat SIC could likely be too
395 high because of mixed ocean-ice Landsat pixels (see Subsection 2.2.4 and the respective supplementary material). What
396 appears to be 100 % thin ice might be just 50 % thick ice. However, observations conducted at Henrik Krøyer Holme station
397 (80°38'N 13°43'W, see star in Fig. 5, top left panel) on September 15 2015 and the preceding days indicate freezing conditions
398 with air temperatures between -5°C and -13°C (<https://www.dmk.dk/vejarkiv>, last access: June 29 2021). Therefore, it is quite
399 likely, new / thin ice covers most open water patches and any over-estimation of Landsat SIC due to sub-pixel size open water
400 patches is rather small. Thus, the complex sea-ice conditions encountered appear to be a valid explanation for the observed
401 differences. Contributing factors also are the different footprint sizes and grid resolutions, that cause heterogeneous sub-grid
402 surface conditions to be mixed differently (see the panels for the three SICCI products in Fig. 5), and unaccounted weather
403 influence. An apparent underestimation of the SIC (see e.g. ASI-SSMI) could be caused by actual weather conditions being
404 less severe, i.e. smaller atmospheric water vapor content, than are included inherently in the open tie point used (see also
405 Kaleschke et al., 2001).

406 Figure 6 illustrates a freeze-up case in Pine Island Bay, Amundsen Sea, Southern Ocean, on March 12, 2014. The
407 classified Landsat-8 scene features a predominant coverage with new / young ice, some open water towards the coast and little
408 thick / snow covered ice and icebergs in the open bay. Landsat SIC is mostly around 90 %; only few grid cells with low SIC
409 exist close to the coast at 12.5 km and 25 km grid resolution. Nine of the ten PMW SIC products reveal considerably lower
410 SIC values with SICCI-25km, OSI-450, NT1-SSMI and ASI-SSMI exhibiting particularly large widespread negative
411 differences with magnitude > 40 %. An exception is NT2-AMSR2 exhibiting the highest PMW SIC of all ten products and
412 overall the smallest differences. It is the only product, though, which also exhibits positive differences, i.e. an over-estimation
413 of Landsat SIC by up to 20 %.

414 The widespread under-estimation of Landsat SIC by almost all products agrees very well with the findings of Ivanova
415 et al. (2015), albeit a bit large in magnitude. The new ice encountered in our example comprises a comparably large fraction
416 of frazil / grease / small pancake ice while compared to nilas and grey/grey-white ice in Ivanova et al. (2015). Because Pine
417 Island Glacier Automatic Weather Station (see star in top left map of Fig. 6) reported air temperatures between -11°C and -
418 21°C on March 12, 2014 and the three preceding days (Mojica Moncada et al., 2019), the grey pixels in this Landsat scene
419 very likely represent new/thin sea ice formed locally. However, we cannot fully exclude an over-estimation of Landsat SIC by
420 sub-pixel size open water patches between streaks of new ice formed being classified as thin ice instead of open water (see
421 Subsection 2.2.4 and respective supplementary material); for the conditions encountered this positive bias in Landsat SIC
422 should be less than 5 %, maximum 10 %. The existence of such a positive bias combined with the different ice type encountered
423 compared to Ivanova et al. (2015) could explain why the observed under-estimation of Landsat SIC for most of the PMW SIC
424 products is larger in magnitude than expected.

425 Table 9 summarizes our results of the freeze-up cases for which we expect, overall, an under-estimation of Landsat
426 SIC, i.e. a negative bias, due to a notable fraction of new / thin ice (see Ivanova et al., 2015). In the Northern Hemisphere,
427 performance of the products differs a lot. We find positive biases for the two CBT-products, NOAA-CDR and NT2-AMSR2,
428 large negative biases for the remaining products. SICCI-25km offers the best linear agreement with Landsat SIC. In the
429 Southern Hemisphere, a number of products have a regression line slope close to unity, a small intercept and a squared linear
430 correlation coefficient > 0.8 . Most importantly, however, all products – except NT2-AMSR2 – on average under-estimate
431 Landsat SIC in agreement with Ivanova et al. (2015).

432 4.2 High-concentration ice

433 These are cases where the Landsat scene indicates either a closed ice cover without any leads or openings or an almost
434 closed ice cover with few refrozen leads or openings, resulting in near-100 % Landsat SIC. In the ideal case, we expect PMW
435 SIC is close to 100 %. Figure 7 illustrates such a case for April 4, 2015 in the Beaufort Sea, Arctic Ocean. Landsat SIC is
436 100.0 %. All ten PMW SIC products exhibit quite high sea-ice concentrations – particularly SICCI-50km, NOAA-CDR and
437 NT2-AMSR2. However, the difference maps clearly reveal a (very) small and negative bias for all PMW products. This bias
438 is largest in magnitude for SICCI-12km and ASI-SSMI and smallest in magnitude for NT2-AMSR2.

439 Table 10 summarizes the results obtained for, in total, 40 high-concentration cases in the Northern Hemisphere: 28
440 first-year ice dominated scenes (Landsat-5) and 12 scenes of mixed first-year / multiyear or multiyear ice cases (Landsat-8).
441 We find the largest biases for SICCI-12km and ASI-SSMI independent of ice type. Except for CBT-AMSR and NT2-AMSR,
442 all products exhibit a larger bias for first-year ice cases than mixed first-year / multiyear or multiyear ice cases. We hypothesize
443 that the different biases between PMW and Landsat SIC for these near-100 % cases are caused by the different capabilities of
444 the respective algorithms to derive an accurate SIC independent of ice type – as stated already in Section 3.1. NT1-SSMI and
445 ASI-SSMI appear to have the largest difficulties with the different ice types encountered because their biases vary most. We
446 note the two CBT products and NOAA-CDR (and NT2-AMSR2) have a median difference of 0.0 % independent of ice type
447 – similar to Tables 5 and 6. For SICCI-2 products and OSI-450, median differences are smaller in magnitude than for all ice
448 and approach zero for the mixed first-year / multiyear or multiyear ice cases.

449 Using non-truncated SIC of SICCI-2 products and OSI-450 (see also Table 8), reduces the magnitude of the bias by
450 between 0.5 % for SICCI-50km and 2.1 % for SICCI-12km for the mixed first-year / multiyear or multiyear ice cases (LS8)
451 and less than that for the first-year ice cases. As expected, the standard deviation of the bias increases using non-truncated SIC.
452 The other six PMW products set SIC values > 100 % to 100 % or do not permit a simple retrieval of such SIC values (NT2-
453 AMSR2, but see Ivanova et al., 2015), and would therefore have a different bias and a larger standard deviation than shown in
454 Table 10 (see Kern et al., 2019). Of the SICCI-2 / OSI-450 products, SICCI-50km provides the smallest bias and the smallest

455 standard deviation of this bias: $-0.7\% \pm 2.2\%$ in line with Kern et al. (2019) who reported a bias of $-0.5\% \pm 2.1\%$ for non-
456 truncated SICCI-50km SIC.

457 Figure 8 illustrates a high-concentration case in the Weddell Sea, Southern Ocean, on March 12, 2015. Six of the ten
458 PMW SIC products show almost 100 % sea-ice concentration and almost zero bias. We only find notable deviations from 100
459 % concomitant with a small negative bias for ASI-SSMI, the two CBT-products and SICCI-12km. For our four high-
460 concentration cases in the Southern Ocean (Table 11), we find the largest overall bias for ASI-SSMI. Most products reveal a
461 bias of magnitude 0.3 % or smaller.

462 Using non-truncated SICCI-2 and OSI-450 SIC results in positive biases, ranging between 1.8 % for OSI-450 and 2.7
463 % for SICCI-50km (Table 11, values to the right of the “/”). This amounts to an increase of the mean SICCI-2 / OSI-450 SIC
464 for these cases by $\sim 2.5\%$. This increase is larger than in the Northern Hemisphere (compare Table 10). We explain this with
465 a comparably large fraction of PMW SIC $> 100\%$ for our small high-concentration cases data set of the Southern Hemisphere
466 (four) compared to the Northern Hemisphere (40). Three of the four high-concentration cases identified in the Southern Ocean
467 are from months November / December, a time of the year when melt onset and melt-refreeze cycles cause higher variability
468 of the ice emissivity. One of the likely impacts is a notable fraction of PMW SIC $> 100\%$ (see Fig. S01 in the supplementary
469 material). The same applies in a different way to the case shown in Fig. 8, the only late fall / early winter case out of these four
470 cases. The overall Landsat SIC of this scene is 99.9 %; that of an adjacent scene is 98.9 % (not shown). Sea ice and snow
471 properties in late fall / early winter are often as well quite variable and can cause an elevated spread of the retrieved PMW SIC
472 around 100 %, resulting in a substantial fraction of SIC values $> 100\%$. For instance, the overall SICCI-25km SIC is 101.9 %
473 for the scene shown in Fig. 8 and 103.1 % for the adjacent scene (not shown).

474 4.3 Melt conditions

475 For melt-condition cases, we select Landsat scenes by means of the calendar date. In the Northern Hemisphere, we
476 consider the time-period May 15 to May 31, in the Southern Hemisphere we use the time-period November 15 to February 28.
477 We do not include Landsat scenes subject to melt ponding on sea ice, e.g. during months June through August; this topic is
478 covered in Kern et al. (2020).

479 In the Northern Hemisphere (Table 12), we find positive and comparably small biases for the two CBT products,
480 NOAA-CDR and NT2-AMSR2, negative biases for all other products. We find the best quality of the linear agreement between
481 PMW SIC and Landsat SIC for SICCI-25km, followed by SICCI-50km and SICCI-12km. According to Kern et al. (2020), the
482 second half of May is characterized by an upswing of number and magnitude of SIC values $> 100\%$ for SICCI-2 / OSI-450
483 products (see Fig. S02 in the supplementary material). Using non-truncated SIC of these products reduces the mean bias by
484 1.0 % for SICCI-12km, 0.5 % for SICCI-25km, and 0.3 % for OSI-450 and further improves the already good linear agreement.
485 For SICCI-50km, results remain almost unchanged. We explain the difference in the response between SICCI-50km and
486 SICCI-12km with the larger sensitivity of the higher frequency channels used by SICCI-12km to early stages of melt
487 encountered at that time of the year.

488 Figure 9 illustrates a typical case of late summer melt conditions in the Ross Sea, Southern Ocean. The classified
489 Landsat-8 image shows a heterogeneous mixture of black, grey and white pixels. The grey pixels denote a mixture of open
490 water and thicker ice, possibly brash ice, sea ice with a wet or even flooded snow cover, or bare relatively thick ice from which
491 the snow has been washed off. New/young ice is unlikely according to 6-hourly forecasts of the Antarctic Mesoscale Prediction
492 System (AMPS) revealing near-surface temperatures around -1°C on January 27 2014 and between -3°C and -5°C on January
493 28 and 29 2014 (<http://polarmet.osu.edu/AMPS/>, last access June 29, 2021), indicating that freeze-up has not yet commenced.

494 PMW SIC distributions match well with Landsat SIC, which is $> 70\%$ for a considerable fraction of the map, but for
495 most products PMW SIC is considerably smaller. Negative biases dominate and are widespread 30 % to 50 % in magnitude.
496 Striking is the similarity between LSIC 12.5km and ASI-SSMI, and between LSIC 25 km and SICCI-25km as well as CBT-

497 AMSR2. Striking also is the similarity between OSI-450, NT1-SSMI, CBT-SSMI and NOAA-CDR. These similarities indicate
498 that different native spatial resolutions, TB sampling intervals and grid spacings of, SSMI(S) on the one hand and AMSR-E(2)
499 on the other hand, can cause a substantial difference in the agreement with LSIC especially when ice conditions are as
500 heterogeneous as in this example (see Section 5.1).

501 Overall, we find negative biases for nine of the ten products in the Southern Hemisphere (Table 13). These are smallest
502 in magnitude for CBT-SSMI and NOAA-CDR: < 1 %, and largest for NT1-SSMI, ASI-SSMI and SICCI-50km. NT2-AMSR2
503 stands out as the only product with a positive bias of 5 % (see also Section 5.2). SICCI-25km and SICCI-50km again provide
504 the best linear agreement with Landsat SIC (Table 13). Results for SICCI-2 products and OSI-450 improve when using non-
505 truncated SIC (see also Fig. S01 in the supplementary material). In contrast to the Northern Hemisphere (see Table 12, Fig.
506 S02 in the supplementary material), also SICCI-50km reveals a reduction of the bias and increase in the linear regression line
507 slope. We attribute this to the presence of advanced melt conditions and the different melt-induced snow and ice properties in
508 the Southern Hemisphere comprising a larger fraction of coarse-grained snow due to pro-longed melt-freeze cycles and a
509 generally drier snow surface, at least for the high-concentration parts of the sea-ice cover.

510 On the one hand, the negative biases (Figure 9, Table 13) agree well with results of earlier comparisons between
511 Southern Hemisphere summer PMW SIC estimates and ship-based observations of the sea-ice cover (e.g. Worby and Comiso,
512 2004; Ozsoy-Cicek et al., 2009). These studies hypothesized that under-estimation of the actual sea-ice concentration in PMW
513 SIC products is primarily caused by wet, flooded sea ice exhibiting a similar surface emissivity as open water and hence
514 looking like open water in PMW imagery. On the other hand, an unknown fraction of these negative biases could be caused
515 by our Landsat SIC estimates being biased high because of the reasons laid out in Subsection 2.2.4 and the respective
516 supplementary material.

517 5 Discussion

518 5.1 A note on grid resolutions

519 SICCI-25km and SICCI-50km SIC have a grid resolution close to the actual algorithm resolution largely determined
520 by the native resolution of the lowest-frequency channel used (see field-of-view dimensions in Table 1). This is not the case
521 for, e.g. CBT-SSMI or OSI-450. Actually, we find a relatively poor performance of OSI-450 in comparison to SICCI-25km
522 (see Tables 5 to 7) – albeit the retrieval algorithm is exactly the same. We hypothesize that the coarser native resolution of the
523 satellite data used for OSI-450 provides one of the main explanations for this observation. SICCI-25km uses AMSR-E/2
524 brightness temperatures observed at spatial resolutions (footprint sizes) between 14 km x 8 km (AMSR2: 12 km x 7 km) and
525 27 km x 16 km (AMSR2: 22 km x 14 km) (see Table 1). In contrast, OSI-450 uses SSM/I(S) brightness temperatures observed
526 at footprint sizes between 37 km x 28 km and 69 km x 43 km. In addition, the relevant channels are sampled spatially every
527 10 km for AMSR-E/2 and every 25 km for SSM/I(S). Therefore, spatial brightness temperature variations caused, e.g., by
528 variations in the open water fraction, can be identified at a finer spatial scale by AMSR-E/2 than by SSM/I(S) at the same
529 frequency. The grid spacing at which OSI-450 and other SIC products relying on SSMI(S) 19 / 37 GHz channels are provided
530 is not the actual resolution of the estimated SIC. Surface information is smeared in the SSM/I(S) data much more. A similar
531 observation applies to CBT-SSMI and CBT-AMSR. The latter provides SIC at a grid resolution closer to the algorithm
532 resolution than CBT-SSMI; consequently, CBT-AMSR SIC agree closer to Landsat SIC than CBT-SSMI SIC (see Tables 5,
533 6, and 7 and compare panels e) and g) in Fig. 2, 3 and 4). We are confident that, besides the differences between the algorithms
534 themselves, a substantial fraction of the observed difference in the agreement with Landsat SIC is caused by the spatial
535 representation of the true sea-ice concentration, which differs due to the above-mentioned differences in satellite data used as
536 input.

537 Our results confirm the stated impact of the native spatial resolution on potential biases between PMW SIC and
538 Landsat SIC very well. For instance, out of the ten products, ASI-SSMI and SICCI-12km both incorporating high-frequency,
539 fine spatial resolution imagery channels provide the 3rd and 4th best linear fits in the Northern Hemisphere (Tables 5, 6) and
540 the 3rd and 5th best linear fits in the Southern Hemisphere. SICCI-12km actually performs best out of the four SICCI-2 / OSI-
541 450 products in the Southern Hemisphere (Table 7). Our Landsat data set of the Southern Hemisphere contains more cases of
542 ice regimes (see Section 4) with variable open water fractions such as “heterogeneous ice”, “leads / openings”, “freeze-up”,
543 and “ice edge” than the one of the Northern Hemisphere (see Table S01 in the supplementary material). Because a SIC product
544 at finer spatial resolution is capable to depict such variable open water fractions better and to observe the full SIC range more
545 adequately it seems reasonable to have a better linear agreement between Landsat SIC and, e.g., SICCI-12km SIC in the
546 Southern than the Northern Hemisphere (compare Figs. 3 and 4 with respect to low SIC).

547 However, ASI-SSMI, does not show better results in the Southern than the Northern Hemisphere when compared to,
548 e.g. NT1-SSMI or SICCI-2 products. ASI-SSMI utilizes near-90 GHz brightness temperatures only while SICCI-12km
549 combines near-90 GHz with 19 GHz brightness temperatures. Atmospheric effects known to cause biases in near-90 GHz
550 PMW SIC products (Kern, 2004; Ivanova et al., 2015) therefore have less impact on SICCI-12km than ASI-SSMI SIC. In
551 addition, all SICCI-2 products utilize atmospherically corrected brightness temperatures while ASI-SSMI utilizes uncorrected
552 brightness temperatures. The fact that most of our Landsat scenes in the Southern Hemisphere represent atmospheric conditions
553 during summer melt and hence at a comparably higher water vapor load than in the Northern Hemisphere fits into this picture.
554 While atmospheric effects are efficiently mitigated for SICCI-12km in both hemispheres these are larger for ASI-SSMI in the
555 Southern than the Northern Hemisphere.

556 5.2 Hemispheric differences versus Landsat SIC bias

557 At this point, we look at the difference between the PMW SIC minus Landsat SIC values obtained in the Northern
558 Hemisphere and the Southern Hemisphere from a different perspective. Ice conditions represented by our Landsat SIC data
559 set comprise more cases with melt conditions and at the ice edge in the Southern Hemisphere (see Table S01 in the
560 supplementary material). These conditions are likely particularly subject to the positive bias in Landsat SIC due to mixed
561 pixels described in Subsection 2.2.4 and the respective supplementary material. Therefore, we can expect that the positive SIC
562 difference is, on average, larger in the Southern than the Northern Hemisphere. We compare the differences listed in Tables 5,
563 6 and 7 and find the following. OSI-450, SICCI-12km, and SICCI-25km exhibit small changes in the SIC differences between
564 +0.8 % and -0.8 %. NT2-AMSR reveals a positive change of +2.8 %. All other products show a negative change by between
565 -2.2 % and -3.2 %. This change of ~ 3 % in the SIC difference between the Northern and the Southern Hemisphere is of the
566 correct sign and of an order of magnitude we deem a realistic estimate of the difference in the mentioned positive Landsat SIC
567 bias between the hemispheres. What does this mean? For example, for a PMW grid cell covered by an actual SIC of 95 %, due
568 to the positive bias, Landsat SIC might be 97 % in the Northern Hemisphere and 100 % in the Southern Hemisphere. A PMW
569 SIC algorithm tuned equally well for the ice conditions in the respective hemisphere would provide 95 % in both hemispheres.
570 Compared to Landsat SIC this results in a negative difference of -2 % in the Northern Hemisphere and of -5 % in the Southern
571 Hemisphere, i.e. the difference becomes more negative by ~ 3 %. In contrast, the difference NT2-AMSR SIC minus Landsat
572 SIC becomes more positive, increasing from +0.6 % in the Northern Hemisphere to +3.4 % in the Southern Hemisphere. When
573 only considering the melt-condition cases the overall difference increases from +1.7% to +5.1% (Tables 12, 13). Without
574 further independent evaluation data to better assess the accuracy of our Landsat SIC data we cannot draw a quantitative
575 conclusion here. However, the increase in the positive value of the difference PMW SIC minus Landsat SIC between the
576 Northern and the Southern Hemisphere observed for NT2-AMSR is opposite to our well-motivated suggestion that Landsat
577 SIC values are biased higher in the Southern than the Northern Hemisphere.

578 **5.3 A note on the effect of filters**

579 In this subsection, we comment on the observation that in the scatterplots of the Northern Hemisphere (Figs. 2 and 3)
 580 particularly the SICCI-2 products but also OSI-450, CBT-AMSR and NT2-AMSR exhibit a relatively large number of cases
 581 with PMW SIC = 0 % and Landsat SIC > 0 %. In addition, we find an unexpected large number of comparably low PMW SIC
 582 values (< ~ 50 %) at Landsat SIC > ~ 70 %, especially for SICCI-50km (Fig. 2c, Fig. 3c). In the scatterplots of all products in
 583 the Southern Hemisphere (Fig. 4) we observe a large number of cases with PMW SIC = 0 % and Landsat SIC > 0 %.

584 We hypothesize this observation is linked to the various filters applied. Examples of such filters are the weather or open
 585 water filter (OWF) and the land spill-over filter (LSO). The OWF reduces the number of erroneous SIC values resulting from
 586 unaccounted atmospheric influence, for example high cloud liquid water contents. OWF is effective along the ice edge and the
 587 adjacent open water. One common realization of the OWF is to set PMW SIC = 0 % once brightness temperature gradient
 588 ratios sensitive to the atmospheric influence exceed a certain threshold (e.g. Wensnahan et al., 1993; Spreen et al., 2008;
 589 Lavergne et al., 2019). Such filters might cut off true SIC values (Andersen et al., 2006). The SICCI-2 and OSI-450 algorithm
 590 employs a modified version of such an OWF (Lavergne et al., 2019; Kern et al., 2019). The LSO reduces the number of
 591 erroneous SIC values along coastlines resulting from unaccounted spillover of the (higher) land surface brightness temperature
 592 into the (lower) open water brightness temperature. The LSO is particular effective during summer. It has also an influence
 593 during the freezing season for situations where the coastline is only fringed by a quite narrow sea ice cover, for example,
 594 during fall freeze-up in the Hudson Bay and along the Siberian coast or during winter / spring along the coast of Greenland
 595 facing the Irminger Sea. One realization of the LSO is a statistical approach, where the SIC of grid cells adjacent to the coast
 596 is corrected, i.e. set to 0 % or interpolated to a more adequate value, based on SIC values within a certain neighborhood (e.g.
 597 Cavalieri et al., 1999). The SICCI-2 and OSI-450 algorithm employs a novel attempt. Here the method of Maass and Kaleschke
 598 (2010) is used to correct for the land spillover already at brightness temperature level; the “classical” LSO filtering of Cavalieri
 599 et al. (1999) is still included, though (Lavergne et al., 2019). Note: the OWF sets PMW SIC to zero; the LSO reduces the PMW
 600 SIC to lower values but not necessarily to zero.

601 The SICCI-2 and OSI-450 products offer the full SIC distribution around 0 % and around 100 % SIC and the
 602 opportunity to reverse-engineer the effect of flags, i.e. switch the effect of certain flags on or off. Therefore, we are able to
 603 investigate the impact of the OWF and the LSO on our comparison results, an investigation not possible for the six other
 604 products. We choose ice regime “leads/openings” in the Southern Hemisphere in years 2013-2015 and look, as an example for
 605 such an investigation, at the impact of the two above-mentioned filters on SICCI-50km SIC (Fig. 10). We switch off these
 606 flags together with the near-100 % SIC flag to work with a more realistic SIC distribution at the high-concentration end. We
 607 do not find even one PMW SIC = 0 % case in the fully non-truncated, i.e. no filters applied, SIC scatterplot (Fig. 10b) – in
 608 contrast to the fully truncated SIC (Fig. 10a). Accordingly, the overall SIC difference reduces in magnitude from 7.5 % to 4.3
 609 % when going from fully truncated to fully non-truncated; the standard deviation of the difference reduces from 15.0 % to
 610 11.1 %.

611 If we switch off the OWF, i.e. include the originally retrieved SIC values for those grid cells where the OWF is applied,
 612 we get a number of SIC data pairs concentrated between Landsat SIC: 0 – 20 % and SICCI-50km: 0 – 30 % that can be clearly
 613 associated with the OWF (compare Fig. 10 panel c) with panels a) and d). The magnitude of the difference decreases by only
 614 0.5 % while the standard deviation stays the same. There is still a comparably large number of cases with SICCI-50km SIC =
 615 0 % or at least relatively low: < 30 %, concomitant with Landsat SIC > 50 %. If we instead switch off the LSO, i.e. include
 616 the originally retrieved SIC values for those grid cells where the LSO is applied, we find that almost all of the above-mentioned
 617 cases of low or equal-to-0 % SICCI-50km SIC can be traced back to substantially higher SIC values (Fig. 10d). The magnitude
 618 of the difference changes considerably from 7.5 % (see above) to 4.9 % if keeping only the LSO filtered grid cells; the standard
 619 deviation of the difference reduces from 15.0 % (see above) to 11.2 %. This reduction in the spread of values around the

620 identity line is also evident very well in the respective scatterplots (Fig. 10): the standard deviation of the Landsat SIC 10 %
621 bin average SICCI-50km SIC (red vertical bars) is much smaller in panel d) than panel a).

622 We observe a similar tendency for all other ice regimes where the LSO is applied, e.g. “freeze-up” or “melt conditions”,
623 in the Southern and in the Northern Hemisphere and for SICCI-25km and SICCI-12km as well (see Tables S04 and S05 in the
624 supplementary material). However, we find far fewer SIC data pairs subject to LSO filtering for OSI-450; hence the effect of
625 switching on or off the LSO is comparably small. We hypothesize that this could be explained with the different native
626 resolution of the satellite data used, the different sampling, and the different grid cell size and spacing (see Section 5.1).
627 However, testing this hypothesis is beyond the scope of this paper. For the SICCI-2 SIC products, we can confirm the
628 hypothesis that the comparably large number of PMW SIC = 0 % or <~30 % across basically the entire SIC range (see Figs.
629 2, 3, and 4, panels a) to c) can be explained with the application of an LSO resulting in an elevated number of cases with PMW
630 SIC < Landsat SIC. This provides a viable explanation for unexpectedly large SIC differences observed for SICCI-50km along
631 coastlines, of particularly Greenland or the Eastern Antarctic, reported in Kern et al. (2019, their Fig. 8 c) and Fig. 11 c).
632 Whether this is due to the land spillover correction at the brightness temperature level (Maass and Kaleschke, 2010) or the
633 statistical filtering (Cavalieri et al., 1999) remains to be investigated. We clearly see it as an advantage to be able to switch off
634 filters and in a reverse-engineering way investigate the impact of these filters for SICCI-2 and OSI-450 products. Application
635 of the LSO can produce an artificially large number of SIC values near or at 0 % that agree less well with the Landsat SIC than
636 the originally retrieved SIC values – as we demonstrate for the SICCI-2 and OSI-450 products. As a consequence, results of
637 an evaluation including a considerable number of near-coastal grid cells need to be interpreted carefully. The number of
638 artificially low SIC values resulting from the LSO for the other six PMW SIC products is unknown as is their impact on the
639 evaluation results shown in this paper.

640 6 Summary and Conclusions

641 In this paper, we present results of an evaluation of ten passive microwave (PMW) SIC products against SIC estimates
642 derived from more than 350 clear-sky Landsat visible images acquired in the Northern Hemisphere during mostly late winter
643 / spring (March through May) and in the Southern Hemisphere during spring / summer / early fall (October through March).
644 We estimate Landsat SIC at the grid resolution of the PMW SIC products using results of supervised classification of Landsat
645 broadband albedo maps into ice and water at 30 m pixel resolution. The comparison between PMW and Landsat SIC is carried
646 out based on all valid collocated SIC map pairs but also based on subsets of these pairs defining certain ice regimes. These ice
647 regimes are “high concentration”, “freeze-up”, “ice edge”, “leads/openings”, “heterogeneous ice”, and “melt conditions”. Our
648 comparison uses statistical parameters such as the mean difference between PMW and Landsat SIC and its standard deviation,
649 the median difference, and parameters describing the linear agreement: slope and intercept of a linear regression and the linear
650 regression coefficient. We summarize these parameters in Figures 11 and 12 and make the following conclusions.

- 651 • It is important to take an integrated view of the statistical parameters because, for instance, a small overall bias is not
652 necessarily associated with a good linear agreement across the entire SIC range and a perfect linear agreement with a slope
653 close to unity and a high correlation could be associated with a large overall bias.
- 654 • It is also important to take into account the expected influences of, e.g. melt conditions (section 4.3), fraction of new/thin
655 ice (section 4.1) as well as sub-pixel size ocean-ice mixture (Section 2.2.4) on both PMW SIC and Landsat SIC.
- 656 • SICCI-25km and SICCI-50km SIC offer overall the best linear agreement to Landsat SIC as demonstrated in Fig. 11 and
657 Fig. 12, right column. This is illustrated as well by mean and median PMW SIC values computed for Landsat SIC bins
658 aligned very well along the identity line (Figs. 2 to 4), with exceptions being explainable by filters applied in the products
659 (see Section 5.3). The magnitude of the difference PMW SIC minus Landsat SIC is, however, larger than for the two
660 CBT-products and NOAA-CDR, almost without exception (Fig. 12, left column).

- The two CBT products, NOAA-CDR and NT2-AMSR offer the smallest overall magnitude of the difference PMW SIC minus Landsat SIC (Fig. 12, left column). Except for CBT-AMSR2 in the Southern Hemisphere, mean and median PMW SIC values align less well along the identity line than for SICCI-25km and SICCI-50km in Figs 2 to 4. The linear agreement is considerably worse than for SICCI-25km and SICCI-50km (Fig. 11, Fig. 12, right column).
- NT2-AMSR is the only product over-estimating Landsat SIC in the Southern Hemisphere – overall but also for almost all ice regimes. This is problematic in view of the potential positive bias of Landsat SIC for ice conditions with an elevated number of mixed ocean-ice Landsat pixels (see Subsection 2.2.4), e.g. ice regimes “melt conditions”, “ice edge” and “freeze-up”.

All products provide SIC data truncated to the range 0 % to 100 % albeit all algorithms, except NT2-AMSR, use a SIC retrieval procedure which in principle provides a full SIC distribution around the end-members 0 % and 100 %. Only the SICCI-2 products and OSI-450 allow consideration of the full SIC distribution. While our main results are derived with the truncated SIC distribution, we demonstrate that, without exception, using the full SIC distribution reduces the mean difference and enhances the quality of the linear agreement between PMW SIC and Landsat SIC which is already superior for SICCI-25km and SICCI-50km. It is important to consider this observation when comparing the results obtained with the ten products against each other in order to avoid misinterpretation. While we obtain smallest SIC differences for the two CBT products, NOAA-CDR and NT2-AMSR/2, these are likely to change using the full SIC distribution. This applies in particular to ice regimes “high-concentration” (section 4.2) and “melt conditions”, but also to the full set of SIC data pairs (denoted “all” in Fig. 12). The impact this difference in the comprehensiveness of the SIC products has on our evaluation results prevents us from making a ranking between the SIC products.

This paper is limited to clear-sky visible imagery. It is hence impossible to evaluate the performance of the SIC products under the full set of possible weather conditions influencing SIC retrieval, i.e. surface wind speed and atmospheric water vapor and cloud liquid water content. Our results likely cover a certain range of surface wind speeds and atmospheric water vapor contents which we, however, did not quantify, e.g. by means of atmospheric reanalysis data, to stay focused. Obviously, this would be an issue worth pursuing in a forthcoming study for which SIC estimates based on SAR data have to be used. These might allow us to assess PMW SIC quality also under higher loads of atmospheric water vapor content and, more importantly, clouds. Such a study could then focus in particular on an improved accuracy assessment of the PMW SIC in the marginal ice zone and along the ice edge. In such regions, our approach to derive Landsat SIC likely results in the highest positive biases – between a few to in the worst case 20 % for single PMW grid cells – due to mixed ocean-ice Landsat pixels classified as ice. Such a study would also be an excellent opportunity to evaluate the weather filters currently employed in the SIC products. In order to have a meaningful sample, such a study would require an equally extensive data set of SAR images interpreted into well-evaluated SIC estimates. This calls for continued development of reliable and consistent SIC estimates from SAR and, thorough evaluation of SAR SIC products in both hemispheres.

Data availability. All sea-ice concentration products except SICCI-12km are publicly available from the sources provided in the reference list or in Kern et al. (2019). The SICCI-12km product is available upon request from T. Lavergne. The classified Landsat images are available from <https://doi.org/10.25592/uhhfdm.9181> (last access: July 9 2021).

Author contributions. SK wrote the manuscript. TL, LTP and RT contributed to the concept and work presented in the paper and also assisted in the writing. SK performed the data analysis together with LB, MM, and LZ. SK conducted the interpretation with contributions in the interpretation of the results from TL, LTP and RT.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The work presented here was funded by EUMETSAT (through the 3rd Continuous Developments and Operation Phase of OSI SAF) and ESA (through the Climate Change Initiative Sea_Ice_cci project), and the German Research Foundation (DFG) Excellence Initiative CLISAP under Grant EXC 177/2. The publication contributes to the Cluster of

705 Excellence 'CLICCS – Climate, Climatic Change, and Society' and to the Center for Earth System Research and Sustainability
706 (CEN) of the University of Hamburg.

707 **7 References**

- 708 Andersen, S., Tonboe, R. T., Kern, S., and Schyberg, H.: Improved retrieval of sea ice total concentration from spaceborne
709 passive microwave observations using Numerical Weather Prediction model fields: An intercomparison of nine algorithms.
710 *Rem. Sens. Environ.*, 104(4), 374-392, 2006.
- 711 Andersen, S., Pedersen, L. T., Heygster, G., Tonboe, R. T., and Kaleschke, L.: Intercomparison of passive microwave sea ice
712 concentration retrievals over the high concentration Arctic sea ice. *J. Geophys. Res.*, 112, C08004,
713 <https://doi.org/10.1029/2006JC003543>, 2007.
- 714 Barsi, J. A., Kenton, L., Kvaran, G., Markham, B. L., and Pedelty, J. A.: The spectral response of the Landsat-8 operational
715 land imager. *Rem. Sens.*, 6(10), 10232-10251, <https://doi.org/10.3390/rs61010232>, 2014.
- 716 Belchansky, G. I., and Douglas, D. C.: Seasonal comparisons of sea ice concentration estimates derived from SSM/I, OKEAN,
717 and RADARSAT data. *Rem. Sens. Environ.*, 81, 67-81, 2002.
- 718 Boutin, G., Lique, C., Arduin, F., Rousset, C., Talandier, C., Accensi, M., and Girard-Ardhuin, F.: Towards a coupled model
719 to investigate wave-sea ice interactions in the Arctic marginal ice zone. *The Cryosphere*, 14(2), 709-735,
720 <https://doi.org/10.5194/tc-14-709-2020>, 2020.
- 721 Brandt, R. E., Warren, S. G., Worby, A. P., and Grenfell, T. C.: Surface Albedo of the Antarctic sea ice zone. *J. Climate*, 18,
722 3606-3622, 2005.
- 723 Burgard, C., Notz, D., Pedersen, L. T., and Tonboe, R. T.: The Arctic Ocean Observation Operator for 6.9 GHz (ARC3O) –
724 Part 1: How to obtain sea ice brightness temperatures at 6.9 GHz from climate model output. *The Cryosphere*, 14(7), 2369–
725 2386, <https://doi.org/10.5194/tc-14-2369-2020>, 2020.
- 726 Cavalieri, D. J.: A microwave technique for mapping thin sea ice. *J. Geophys. Res.*, 99(C6), 12561-12572, 1994.
- 727 Cavalieri, D. J., Gloersen, P., and Campbell, W. J.: Determination of Sea Ice Parameters with the NIMBUS 7 SMMR. *J.*
728 *Geophys. Res.*, 89(D4), 5355-5369, 1984.
- 729 Cavalieri, D. J., Crawford, J., Drinkwater, M., Emery, W. J., Eppler, D. T., Farmer, L. D., Goodberlet, M., Jentz, R., Milman,
730 A., Morris, C., Onstott, R., Schweiger, A., Shuchman, R., Steffen, K., Swift, C. T., Wackerman, C., and Weaver, R. L.: NASA
731 sea ice validation program for the DMSP SSM/I: final report. NASA Technical Memorandum 104559. National Aeronautics
732 and Space Administration, Washington, D.C. 126 pages, 1992.
- 733 Cavalieri, D. J., Parkinson, C. L., Gloersen, P., Comiso, J. C., and Zwally, H. J.: Deriving long-term time series of sea ice
734 cover from satellite passive-microwave multisensor data sets. *J. Geophys. Res.*, 104(C7), 15803–15814,
735 <http://doi.org/10.1029/1999JC900081>, 1999.
- 736 Cavalieri, D. J., Markus, T., Hall, D. K., Gasiewski, A. J., Klein, M., and Ivanoff, A.: Assessment of EOS Aqua AMSR-E
737 Arctic sea ice concentrations using Landsat-7 and airborne microwave imagery. *IEEE Trans. Geosci. Rem. Sens.*, 44(11),
738 3057-3069, <https://doi.org/10.1109/TGRS.2006.878445>, 2006.
- 739 Chander, G., Markham, B. L., and Barsi, J. A.: Revised Landsat-5 Thematic Mapper Radiometric Calibration. *IEEE Geosci.*
740 *Rem. Sens. Lett.*, 4(3), 490-494, 2007.
- 741 Chander, G., Markham, B. L., and Helder, D. L.: Summary of current radiometric calibration coefficients for Landsat MSS,
742 TM, ETM+, and EO-1 ALI sensors. *Rem. Sens. Environ.*, 113, 893-903, 2009.
- 743 Cheng, A., Casati, B., Tivy, A., Zagon, T., Lemieux, J.-F., and Tremblay, L. B.: Accuracy and inter-analyst agreement of
744 visually estimated sea ice concentrations in Canadian Ice Service ice charts using single-polarization RADARSAT-2. *The*
745 *Cryosphere*, 14(4), 1289-1310, <https://doi.org/10.5194/tc-14-1289-2020>, 2020.

- 746 Comiso J. C.: Characteristics of arctic winter sea ice from satellite multispectral microwave observations. *J. Geophys. Res.*,
747 91(C1), 975-994, 1986.
- 748 Comiso, J. C.: Enhanced sea ice concentrations and ice extents from AMSR-E data. *J. Rem. Sens. Soc. Japan*, 29(1), 199-215,
749 2009.
- 750 Comiso, J. C., and Zwally, H. J.: Antarctic sea ice concentrations inferred from Nimbus 5 ESMR and Landsat imagery. *J.*
751 *Geophys. Res.*, 87(C8), 5836-5844, <https://doi.org/10.1029/JC087iC08p05836>, 1982.
- 752 Comiso, J. C., and Steffen, K.: Studies of Antarctic sea ice concentrations from satellite data and their applications. *J. Geophys.*
753 *Res.*, 106(C12), 31,361-31,385, 2001.
- 754 Comiso, J. C., and Nishio, F.: Trends in the sea ice cover using enhanced and compatible AMSR-E, SSM/I, and SMMR data. *J.*
755 *Geophys. Res.*, 113, C02S07, <http://doi.org/10.1029/2007JC004257>, 2008.
- 756 Comiso, J. C., Wadhams, P., Krabill, W. B., Swift, R. N., Crawford, J. P., and Tucker III, W. B.: Top/bottom multisensory
757 remote sensing of Arctic sea ice. *J. Geophys. Res.*, 96(C2), 2693-2709, <https://doi.org/10.1029/90JC02466>, 1991.
- 758 Comiso, J. C., Cavalieri, D. J., Parkinson, C. L., and Gloersen, P.: Passive microwave algorithms for sea ice concentration: A
759 comparison of two techniques. *Rem. Sens. Environ.*, 60(3), 357-384, 1997.
- 760 Comiso, J. C., Cavalieri, D. J., and Markus, T.: Sea ice concentration, ice temperature, and snow depth, using AMSR-E data.
761 *IEEE Trans. Geosci. Rem. Sens.*, 41(2), 243-252, <https://doi.org/10.1109/TGRS.2002.808317>, 2003.
- 762 Cooke, C. L. V., and Scott, K. A.: Estimating sea ice concentration from SAR: Training convolutional neural networks with
763 passive microwave data. *IEEE Trans. Geosci. Rem. Sens.*, 57(7), 4735-4747, <https://doi.org/10.1109/TGRS.2019.2892723>,
764 2019.
- 765 Dokken, S. T., Håkansson, B., and Askne, J.: Inter-comparison of Arctic sea ice concentration using RADARSAT, ERS, SSM/I
766 and In-Situ Data. *Can. J. Rem. Sens.*, 26(6), 521-536, <https://doi.org/10.1080/07038992.2000.10874793>, 2000.
- 767 Drusch, M., Del Bello, U., Carlier, S., Colin, O., Fernandez, V., Gascon, F., Hoersch, B., Isola, C., Laberinti, P., Martimort,
768 P., Meygret, A., Spoto, F., Sy, O., Marchese, F., and Bargellini, P.: Sentinel-2: ESA's optical high-resolution mission for
769 GMES operational services. *Rem. Sens. Environ.*, 120, 25-36, <https://doi.org/10.1016/j.res.2011.11.026>, 2012.
- 770 Ezraty, R., Girard-Ardhuin, F., Piollé, J.-F., Kaleschke, L., and Heygster, G.: Arctic and Antarctic sea ice concentration and
771 Arctic sea ice drift estimated from special sensor microwave data – Users's Manual, Version 2.1, IFREMER, Brest, France,
772 February 2007.
- 773 Han, H., and Kim, H.-C.: Evaluation of summer passive microwave sea ice concentrations in the Chukchi Sea based on
774 KOMPSAT-5 SAR and numerical weather prediction data. *Rem. Sens. Environ.*, 209, 343-362,
775 <https://doi.org/10.1016/j.rse.2018.02.058>, 2018.
- 776 Heinrichs, J. F., Cavalieri, D. J., and Markus, T.: Assessment of the AMSR-E sea ice concentration product at the ice edge
777 using RADARSAT-1 and MODIS imagery. *IEEE Trans. Geosci. Rem. Sens.*, 44(11), 3070-3080,
778 <https://doi.org/10.1109/TGRS.2006.880622>, 2006.
- 779 Ivanova, N., Johannessen, O. M., Pedersen, R. T., and Tonboe, R. T.: Retrieval of Arctic sea ice parameters by satellite passive
780 microwave sensors: A comparison of eleven sea ice concentration algorithms. *IEEE Trans. Geosci. Rem. Sens.*, 52(11), 7233-
781 7246, <http://doi.org/10.1109/TGRS.2014.2310136>, 2014.
- 782 Ivanova, N., Pedersen, L. T., Tonboe, R. T., Kern, S., Heygster, G., Lavergne, T., Sørensen, A., Saldo, R., Dybkjær, G.,
783 Brucker, L., and Shokr, M.: Inter-comparison and evaluation of sea ice algorithms: towards further identification of challenges
784 and optimal approach using passive microwave observations. *The Cryosphere*, 9(5), 1797-1817, <http://doi.org/10.5194/tc-9-1797-2015>, 2015.
- 786 Kaleschke, L., Lüpkes, C., Vihma, T., Haarpaintner, J., Bochert, A., Hartmann, J., and Heygster, G.: SSM/I sea ice remote
787 sensing for mesoscale ocean-atmosphere interaction analysis. *Can. J. Rem. Sens.*, 27(5), 526-537, 2001.

- 788 Karvonen, J.: A sea ice concentration estimation algorithm utilizing radiometer and SAR data. *The Cryosphere*, 8(5), 1639-
789 1650, <https://doi.org/10.5194/tc-8-1639-2014>, 2014.
- 790 Karvonen, J.: Baltic sea ice concentration estimation using SENTINEL-1 SAR and AMSR2 microwave radiometer data. *IEEE*
791 *Trans. Geosci. Rem. Sens.*, 55(5), 2871-2883, <https://doi.org/10.1109/TGRS.2017.2655567>, 2017.
- 792 Kern, S.: A new method for medium-resolution sea ice analysis using weather-influence corrected Special Sensor
793 Microwave/Imager 85 GHz data. *Int. J. Rem. Sens.*, 25(21), 4555-4582, 2004.
- 794 Kern, S., Kaleschke, L., and Clausi, D. A.: A comparison of two 85-GHz SSM/I ice concentration algorithms with AVHRR
795 and ERS-2 SAR imagery. *IEEE Trans. Geosci. Rem. Sens.*, 41(10), 2294-2306, <https://doi.org/10.1109/TGRS.2003.817181>,
796 2003.
- 797 Kern, S., Kaleschke, L., Girard-Ardhuin, F., Spreen, G., and Beitsch, A.: Global daily gridded 5-day median-filtered, gap-
798 filled ASI Algorithm SSMI-SSMIS sea ice concentration data, Integrated Climate Date Center (ICDC), CEN, University of
799 Hamburg, Germany, <https://www.cen.uni-hamburg.de/en/icdc/data/cryosphere/seaiceconcentration-asi-ssmi.html> [last access:
800 December 9, 2021], 2020.
- 801 Kern, S., Lavergne, T., Notz, D., Pedersen, L., Tonboe, R., Saldo, R., and Sørensen, A. M.: Satellite passive microwave sea-
802 ice concentration data set intercomparison: closed ice and ship-based observations. *The Cryosphere*, 13(12), 3261-3307,
803 <http://doi.org/10.5194/tc-13-3261-2019>, 2019.
- 804 Kern, S., Lavergne, T., Notz, D., Pedersen, L. T., and Tonboe, R.: Satellite passive microwave sea-ice concentration data set
805 intercomparison for Arctic summer conditions. *The Cryosphere*, 14(7), 2469-2493, <https://doi.org/10.5194/tc-14-2469-2020>,
806 2020.
- 807 Knap, W. H., Brock, B. W., Oerlemans, J., and Willis, I. C.: Comparison of Landsat TM-derived and ground-based albedos of
808 Haut Glacier d'Arolla, Switzerland. *Int. J. Rem. Sens.*, 20(17), 3293-3310, 1999.
- 809 Koepke, P.: Removal of Atmospheric Effects from AVHRR albedos. *J. Appl. Meteorol.*, 28, 1341-1348, 1989.
- 810 Komarov, A. S., and Buehner, M.: Automated detection of ice and open water from dual-polarization RADARSAT-2 images
811 for data assimilation. *IEEE Trans. Geosci. Rem. Sens.*, 55(10), 5755-5769, <https://doi.org/10.1109/TGRS.2017.2713987>,
812 2017.
- 813 Komarov, A. S., and Buehner, M.: Improved retrieval of ice and open water from sequential RADARSAT-2 images. *IEEE*
814 *Trans. Geosci. Rem. Sens.*, 57(6), 3694-3702, <https://doi.org/10.1109/TGRS.2018.2886685>, 2019.
- 815 Kwok, R.: Sea ice concentration estimates from satellite passive microwave radiometry and openings from SAR ice motion.
816 *Geophys. Res. Lett.*, 29(9), 1311, <https://doi.org/10.1029/2002GL014787>, 2002.
- 817 Lavergne, T., Macdonald Sørensen, A., Kern, S., Tonboe, R., Notz, D., Aaboe, S., Bell, L., Dybkjær, Eastwood, S., Gabarro,
818 C., Heygster, G., Killie, M., Brandt Kreiner, M., Lavelle, J., Saldo, R., Sandven, S., and Pedersen, L.: Version 2 of the
819 EUMETSAT OSI SAF and ESA-CCI sea-ice concentration climate data records. *The Cryosphere*, 13(1), 49-78,
820 <http://doi.org/10.5194/tc-13-49-2019>, 2019.
- 821 Leigh, S., Wang, Z., and Clausi, D. A.: Automated ice-water classification using dual polarization SAR satellite imagery. *IEEE*
822 *Trans. Geosci. Rem. Sens.*, 52(9), 5529-5539, <https://doi.org/10.1109/TGRS.2013.2290231>, 2014.
- 823 Liu, Y., Key, J., and Mahoney, R.: Sea and freshwater ice concentration from VIIRS on Suomi NPP and the future JPSS
824 satellites. *Rem. Sens.*, 8(6), 523-542, <https://doi.org/10.3390/rs8060523>, 2016.
- 825 Lohse, J., Doulgeris, A. P., and Dierking, W.: An optimal decision-tree design strategy and its application to sea ice
826 classification from SAR imagery. *Rem. Sens.*, 11(13), 1574-1588, <https://doi.org/10.3390/rs11131574>, 2019.
- 827 Lu, J., Heygster, G., and Spreen, G.: Atmospheric correction of sea-ice concentration retrieval for 89GHz AMSR-E
828 observations. *IEEE J. Sel. Topics Appl. Earth. Obs. Rem. Sens.*, 11(5), 1442-1457,
829 <https://doi.org/10.1109/JSTARS.2018.2805193>, 2018.

- 830 Lu, P., Li, Z. L., Zhang, Z. H., and Dong, X. L.: Aerial observations of floe size distributions in the marginal ice zone of
831 summer Prydz Bay. *J. Geophys. Res.*, 113, C02011, <https://doi.org/10.1029/2006JC003965>, 2008.
- 832 Ludwig, V., Spreen, G., and Pedersen, L. T.: Evaluation of a new merged sea-ice concentration dataset at 1 km resolution from
833 thermal infrared and passive microwave satellite data in the Arctic. *Rem. Sens.*, 12(19), 3183-3210,
834 <https://doi.org/10.3390/rs12193183>, 2020.
- 835 Maass, N., and Kaleschke, L.: Improving passive microwave sea ice concentration algorithms for coastal areas: applications
836 to the Baltic Sea. *Tellus*, 62A(4), 393-410, <https://doi.org/10.1111/j.1600-0870.2010.00452.x>, 2010.
- 837 Malmgren-Hansen, D., Pedersen, L. T., Aasbjerg Nielsen, A., Brandt Kreiner, M., Saldo, R., Skriver, H., Lavelle, J., Buus-
838 Hinkler, J., and Harnvig Krane, K.: A convolutional neural network architecture for sentinel-1 and AMSR2 data fusion. *IEEE*
839 *Trans. Geosci. Rem. Sens.*, 59(3), 1890-1902, <https://doi.org/10.1109/TGRS.2020.3004539>, 2020.
- 840 Marcq, S., and Weiss, J.: Influence of sea ice lead-width distribution on turbulent heat transfer between the ocean and the
841 atmosphere. *The Cryosphere*, 6(1), 143-156, <https://doi.org/10.5194/tc-6-143-2012>, 2012.
- 842 Markus, T., and Cavalieri, D. J.: An enhancement of the NASA Team sea ice algorithm. *IEEE Trans. Geosci. Rem. Sens.*,
843 38(3), 1387-1398, 2000.
- 844 Markus, T., and Cavalieri, D. J.: The AMSR-E NT2 sea ice concentration algorithm: its basis and implementation. *J. Rem.*
845 *Sens. Soc. Japan*, 29(1), 216-225, 2009.
- 846 Markus, T., and Dokken, S. T.: Evaluation of late summer passive microwave Arctic sea ice retrievals. *IEEE Trans. Geosci.*
847 *Rem. Sens.*, 40(2), 348-356, 2002.
- 848 Meier, W. N.: Comparison of passive microwave ice concentration algorithm retrievals with AVHRR imagery in Arctic
849 peripheral seas. *IEEE Trans. Geosci. Rem. Sens.*, 43(6), 1324-1337, <http://doi.org/10.1109/TGRS.2005.846151>, 2005.
- 850 Meier, W. N., and Windnagel, A.: Sea ice concentration – climate algorithm theoretical basis document, NOAA Climate Data
851 Record Program CDRP-ATBD-0107 Rev. 7 (03/06/2018), available at <https://www.ncdc.noaa.gov/cdr/oceanic/sea-ice-concentration>, 2018.
- 852 Meier, W. N., Markus, T., and Comiso, J. C.: AMSR-E/AMSR2 Unified L3 Daily 25.0 km Brightness Temperatures, Sea Ice
853 Concentration, Motion & Snow Depth Polar Grids, Version 1, Boulder, Colorado USA, NASA National Snow and Ice Data
854 Center Distributed Active Archive Center, 2018, <https://doi.org/10.5067/TRUIAL3WPAUP>, [last access: July 13, 2021], 2018.
- 855 Meier, W. N., Fetterer, F., Savoie, M., Mallory, S., Duerr, R., and Stroeve, J.: *NOAA/NSIDC Climate Data Record of Passive*
856 *Microwave Sea Ice Concentration, Version 3*. Boulder, Colorado USA. NSIDC: National Snow and Ice Data Center.
857 <http://doi.org/10.7265/N59P2ZTG>, [last access: July 13, 2021], 2017.
- 858 Meier, W. N., Fetterer, F., Windnagel, A. K., and Stewart, S.: NOAA/NSIDC Climate Data Record of Passive Microwave Sea
859 Ice Concentration, Version 4. Boulder, Colorado USA. NSIDC: National Snow and Ice Data Center
860 <https://doi.org/10.7265/efmz-2t65>. [last access: December 9, 2021], 2021.
- 861 Mojica Moncada, J. F., and Holland, D.: Automatic Weather Station Pine Island Glacier, U.S. Antarctic Program (USAP) Data
862 Center. <https://doi.org/10.15784/601216> [last access: June 29, 2021], 2019.
- 863 Nose, T., Waseda, T., Kodaira, T., and Inoue, J.: Satellite-retrieved sea ice concentration uncertainty and its effect on modelling
864 wave evolution in marginal ice zones. *The Cryosphere*, 14(6), 2029-2052, <https://doi.org/10.5194/tc-14-2029-2020>, 2020.
- 865 Ochilov, S., and Clausi, D. A.: Operational SAR sea-ice image classification. *IEEE Trans. Geosci. Rem. Sens.*, 50(11), 4397-
866 4408, <https://doi.org/10.1109/TGRS.2012.2192278>, 2012.
- 867 Onana, V.-De-P., Kurtz, N. T., Farrell, S. L., Koenig, L. S., Studinger, M., and Harbeck, J. P.: A sea-ice lead detection
868 algorithm for use with high-resolution airborne visible imagery. *IEEE Trans. Geosci. Rem. Sens.*, 51(1), 38-56,
869 <https://doi.org/10.1109/TGRS.2012.2202666>, 2013.
- 870 OSI SAF: Global Sea Ice Concentration Climate Data Record v2.0 - Multimission, EUMETSAT SAF on Ocean and Sea Ice,
871 http://dx.doi.org/10.15770/EUM_SAF_OSI_0008 [last access: December 9, 2021], 2017a. OSI SAF: Global sea ice

- 873 concentration interim climate data record 2016-onwards (v2.0, 2017), OSI-430-b, <https://osi-saf.eumetsat.int/products/osi-430-b-complementing-osi-450> [last access: September 17 2021], 2017b.
- 874 Ozsoy-Cicek, B., Xie, H., Ackley, S. F., and Ye, K.: Antarctic summer sea ice concentration and extent: comparison of ODEN
875 2006 ship observations, satellite passive microwave and NIC sea ice charts. *The Cryosphere*, 3, 1-9, <https://doi.org/10.5194/tc-3-1-2009>, 2009.
- 876 Paget, M. J., Worby, A. P., and Michael, K. J.: Determining the floe-size distribution of East Antarctic sea ice from digital
877 aerial photographs. *Ann. Glaciol.*, 33, 94-100, 2001.
- 878 Pedersen, L. T.; Dybkjær, G.; Eastwood, S.; Heygster, G.; Ivanova, N.; Kern, S.; Lavergne, T.; Saldo, R.; Sandven, S.;
879 Sørensen, A.; Tonboe, R. T.: ESA Sea Ice Climate Change Initiative (Sea_Ice_cci): Sea Ice Concentration Climate Data Record
880 from the AMSR-E and AMSR-2 instruments at 25km grid spacing, version 2.1. Centre for Environmental Data Analysis, 05
881 October 2017. <http://dx.doi.org/10.5285/f17f146a31b14dfd960cde0874236ee5> [last access: December 9, 2021], 2017.
- 882 Pegau, W. S., and Paulson, C. A.: The albedo of Arctic leads in summer. *Ann. Glaciol.*, 33, 221-224, 2001.
- 883 Peng, G., and Meier, W. N.: Temporal and regional variability of Arctic sea-ice coverage from satellite data. *Ann. Glaciol.*,
884 59(76, part 2), 191-200, <http://doi.org/10.1017/aog.2017.32>, 2018.
- 885 Peng, G., Meier, W. N., Scott, D., and Savoie, M.: A long-term and reproducible passive microwave sea ice concentration data
886 record for climate studies and monitoring. *Earth Syst. Sci. Data*, 5, 311-318. <http://doi.org/10.5194/essd-5-311-2013>, 2013.
- 887 Perovich, D. K., and Jones, K. F.: The seasonal evolution of sea ice floe size distribution. *J. Geophys. Res.-Oceans*, 119, 8767-
888 8777, <https://doi.org/10.1002/2014JC010136>, 2014.
- 889 Singha, S., Johansson, M., Hughes, N., Hvidegaard, S. M., and Skourup, H.: Arctic sea ice characterization using spaceborne
890 fully polarimetric L-, C-, and X-band SAR with validation by airborne measurements. *IEEE Trans. Geosci. Rem. Sens.*, 56(7),
891 3715-3734, <https://doi.org/10.1109/TGRS.2018.2809504>, 2018.
- 892 Shi, Q., Su, J., Heygster, G., Shi, J., Wang, L., Zhu, L., Lou, Q., and Ludwig, V.: Step-by-step validation of Antarctic ASI
893 AMSR-E sea-ice concentrations by MODIS and an aerial image. *IEEE Trans. Geosci. Rem. Sens.*, 59(1), 392-403,
894 <https://doi.org/10.1109/TGRS.2020.2989037>, 2021.
- 895 Shokr, M., and Markus, T.: Comparison of NASA Team2 and AES-York ice concentration algorithms against operational ice
896 charts from the Canadian Ice Service. *IEEE Trans. Geosci. Rem. Sens.*, 44(8), 2164-2175,
897 <https://doi.org/10.1109/TGRS.2006.872077>, 2006.
- 898 Shokr, M., and Agnew, T. A.: Validation and potential applications of Environment Canada Ice Concentration Extractor
899 (ECICE) algorithm to Arctic ice by combining AMSR-E and QuikSCAT observations. *Rem. Sens. Environ.*, 128, 315-332,
900 <https://doi.org/10.1016/j.rse.2012.10.016>, 2013.
- 901 Spreen, G., Kaleschke, L., and G. Heygster, G.: Sea ice remote sensing using AMSR-E 89-GHz channels. *J. Geophys. Res.*,
902 113, C02S03, <https://doi.org/10.1029/2005JC003384>, 2008.
- 903 Steer, A., Worby, A. P., and Heil, P.: Observed changes in sea-ice floe size distribution during early summer in the western
904 Weddell Sea. *Deep-Sea Res. II*, 55, 933-942, <https://doi.org/10.1016/j.dsr2.2007.12.016>, 2008.
- 905 Steffen, K., and Maslanik, J. A.: Comparison of Nimbus 7 scanning multichannel microwave radiometer radiance and derived
906 sea ice concentrations with Landsat imagery for the north water area of Baffin Bay. *J. Geophys. Res.*, 93(C9), 10769-10781,
907 <https://doi.org/10.1029/JC093iC09p10769>, 1988.
- 908 Steffen, K. and Schweiger, A.: NASA team algorithm for sea ice concentration retrieval from Defense Meteorological Satellite
909 Program special sensor microwave imager: comparison with Landsat satellite data. *J. Geophys. Res.*, 96(C12), 21,971-21,987,
910 1991.
- 911 Titchner, H. A., and Rayner, N. A.: The Met Office Hadley Centre sea ice and sea surface temperature data set, version 2: 1.
912 Sea Ice concentrations. *J. Geophys. Res. Atmos.*, 119(6), 2864-2889, <https://doi.org/10.1002/2013JD020316>, 2014.

- 915 Tonboe, R. T., Eastwood, S., Lavergne, T., Sørensen, A. M., Rathmann, N., Dybkjær, G., Pedersen, L. T., Høyer, J. L., and
916 Kern, S.: The EUMETSAT sea ice concentration climate data record. *The Cryosphere*, 10(5), 2275-2290,
917 <http://doi.org/10.5194/tc-10-2275-2016>, 2016.
- 918 Torres, R., Snoeij, P., Geudtner, D., Bibby, D., Davidson, M., Attema, E., Potin, P., Rommen, B., Flouy, N., Brown, M.,
919 Traver, I. N., Deghaye, P., Duesmann, B., Rosich, B., Miranda, N., Bruno, C., L'Abbate, M., Croci, R., Pietropaolo, A.,
920 Huchler, M., and Rostan, F.: GMES Sentinel-1 mission. *Rem. Sens. Environ.*, 120, 9-24,
921 <https://doi.org/10.1016/j.rse.2011.05.028>, 2012.
- 922 Toyota, T., Haas, C., and Tamura, T.: Size distribution and shape properties of relatively small sea-ice floes in the Antarctic
923 marginal ice zone in late winter. *Deep-Sea Res. II*, 58, 1182-1193, <https://doi.org/10.1016/j.dsr2.2010.10.034>, 2011.
- 924 Toyota, T., Kohout, A., and Fraser, A. D.: Formation processes of sea ice floe size distribution in the interior pack and its
925 relationship to the marginal ice zone off East Antarctica. *Deep-Sea Res.*, II, 131, 28-40,
926 <https://doi.org/10.1016/j.dsr2.2015.10.003>, 2016.
- 927 Tschudi, M. A., Curry, J. A., and Maslanik, J. A.: Characterization of springtime leads in the Beaufort/Chukchi Seas from
928 airborne and satellite observations during FIRE/SHEBA. *J. Geophys. Res.*, 107(C10), 8034,
929 <https://doi.org/10.1029/2000JC000541>, 2002.
- 930 Wang, L., Scott, K. A., Xu, L., and Clausi, D. A.: Sea ice concentration estimation during melt from dual-pol SAR scenes
931 using deep convolutional neural networks: A case study. *IEEE Trans. Geosci. Rem. Sens.*, 54(8), 4524-4533,
932 <https://doi.org/10.1109/TGRS.2016.2543660>, 2016.
- 933 Wang, L., Scott, K. A., and Clausi, D. A.: Sea ice concentration estimation during freeze-up from SAR imagery using a
934 convolutional neural network. *Rem. Sens.*, 9(5), 408-427, <https://doi.org/10.3390/rs9050408>, 2017.
- 935 Wang, Y.-R. and Li, X.-M.: Arctic sea ice cover data from spaceborne SAR by deep learning, *Earth Syst. Sci. Data Discuss.*
936 [preprint], <https://doi.org/10.5194/essd-2020-332>, in review, 2020.
- 937 Wensnahan, M., Maykut, G. A., Grenfell, T. C., and Winebrenner, D. P.: Passive microwave remote sensing of thin sea ice
938 using principal component analysis. *J. Geophys. Res.*, 98(C7), 12453-12468, <https://doi.org/10.1029/93JC00939>, 1993.
- 939 Wiebe, H., Heygster, G., and Markus, T.: Comparison of the ASI ice concentration algorithm with Landsat-7 ETM+ and SAR
940 imagery. *IEEE Trans. Geosci. Rem. Sens.*, 47(9), 3008-3015, <https://doi.org/10.1109/TGRS.2009.2026367>, 2009.
- 941 Willmes, S., Nicolaus, M., and Haas, C.: The microwave emissivity variability of snow covered first-year sea ice from late
942 winter to early summer: a model study. *The Cryosphere*, 8(3), 891-904, <https://doi.org/10.5194/tc-8-891-2014>, 2014.
- 943 Worby, A. P., and Comiso, J. C.: Studies of the Antarctic sea ice edge and ice extent from satellite and ship observations. *Rem.*
944 *Sens. Environ.*, 92(1), 98-111, <https://doi.org/10.1016/j.rse.2004.05.007>, 2004.
- 945 Zakhvatkina, N., Korosov, A., Muckenhuber, S., Sandven, S., and Babiker, M.: Operational algorithm for ice-water
946 classification on dual-polarized RADARSAT-2 images. *The Cryosphere*, 11(1), 33-46, <https://doi.org/10.5194/tc-11-33-2017>,
947 2017.
- 948 Zatko, M. C., and Warren, S. G.: East Antarctic sea ice in spring: spectral albedo of snow, nilas, frost flowers and slush, and
949 light-absorbing impurities in snow. *Ann. Glaciol.*, 56(69), 53-64, <https://doi.org/10.3189/2015AoG69A574>, 2015.
- 950 Zhang, Q., and Skjetne, R.: Image processing for identification of sea-ice floes and the floe size distributions. *IEEE Trans.*
951 *Geosci. Rem. Sens.*, 53(5), 2913-2924, <https://doi.org/10.1109/TGRS.2014.2366640>, 2015.
- 952 Zhao, X., Chen, Y., Kern, S., Qu, M., Ji, Q., Fan, P., and Liu, Y.: Sea ice concentration derived from FY-3D MWRI and its
953 accuracy assessment. *IEEE Trans. Geosci. Rem. Sens. (Early Access)*, 18 pp., <https://doi.org/10.1109/TGRS.2021.3063272>,
954 2021.
- 955
- 956
- 957

958 **8 Tables**

959

960 **Table 1.** Overview of the investigated PMW SIC products. Column “ID (Algorithm)” holds the identifier we use henceforth
 961 to refer to the data product, and which algorithm it uses. Note that for those algorithms where an AMSR sensor forms part of
 962 the name, we refer to AMSR-E or AMSR2, depending on the respective data used. Column “Input data” refers to the input
 963 satellite data for the data set, together with the frequencies and respective field-of-view dimensions.

964

ID (algorithm)	Input data; frequencies (field-of-views)	Grid resolution & type	Reference
OSI-450 (SICCI2)	SSM/I, SSMIS; 19.35 GHz (69 km x 43 km), 37.0 GHz (37 km x 28 km)	25 km x 25 km EASE2.0	Tonboe et al., 2016; OSI-SAF, 2017a; 2017b; Lavergne et al., 2019
SICCI-12km (SICCI2)	AMSR-E/AMSR2; 18.7 GHz (27 km x 16 km/ 22 km x 14 km), 89.0 GHz (6 km x 4 km/ 5 km x 3 km)	12.5 km x 12.5 km EASE2.0	Pedersen et al., 2017; Lavergne et al., 2019
SICCI-25km (SICCI2)	AMSR-E/AMSR2; 18.7 GHz (27 km x 16 km/ 22 km x 14 km), 36.5 GHz (14 km x 8 km/ 12 km x 7 km)	25 km x 25 km EASE2.0	Pedersen et al., 2017; Lavergne et al., 2019
SICCI-50km (SICCI2)	AMSR-E/AMSR2 6.9 GHz (75 km x 43 km/ 62 km x 35 km), 36.5 GHz (14 km x 8 km/ 12 km x 7 km)	50 km x 50 km EASE2.0	Pedersen et al., 2017; Lavergne et al., 2019
CBT-SSMI (Comiso bootstrap)	SSM/I, SSMIS; 19.35 GHz (69 km x 43 km), 37.0 GHz (37 km x 28 km)	25 km x 25 km PolarStereo	Comiso, 1986; Comiso et al., 1997; Comiso and Nishio, 2008; Meier et al., 2017
NOAA-CDR (NASA Team & Comiso bootstrap)	SSM/I, SSMIS; 19.35 GHz (69 km x 43 km), 37.0 GHz (37 km x 28 km)	25 km x 25 km PolarStereo	Peng et al., 2013; Meier et al., 2017; Meier and Windnagel, 2018
CBT-AMSR (Comiso bootstrap)	AMSR-E/AMSR2; 18.7 GHz (27 km x 16 km/ 22 km x 14 km), 36.5 GHz (14 km x 8 km/ 12 km x 7 km)	25 km x 25 km PolarStereo	Comiso et al., 2003; Comiso and Nishio, 2008; Comiso, 2009; Meier et al., 2018
ASI-SSMI (ASI)	SSM/I, SSMIS; 85.5 GHz (15 km x 13 km)	12.5 km x 12.5 km PolarStereo	Kaleschke et al., 2001; Ezraty et al., 2007; Kern et al., 2020
NT1-SSMI (NASA-Team)	SSM/I, SSMIS; 19.35 GHz (69 km x 43 km), 37.0 GHz (37 km x 28 km)	25 km x 25 km PolarStereo	Cavalieri et al, 1984; 1992; 1999; Meier et al., 2017
NT2-AMSR (NASA-Team-2)	AMSR-E/AMSR2; 18.7 GHz (27 km x 16 km/ 22 km x 14 km), 36.5 GHz (14 km x 8 km/ 12 km x 7 km), 89.0 GHz (6 km x 4 km/ 5 km x 3 km)	25 km x 25 km PolarStereo	Markus and Cavalieri, 2000; 2009; Meier et al., 2018

965

966

967

Table 2. Overview about the wavelengths and bandwidths of the Landsat channels used.

Wavelength [nm] of	Landsat-5	Landsat-7	Landsat-8
Channel 2	528-609	519-601	--
Channel 3	626-693	631-692	533-590
Channel 4	776-904	772-898	636-673
Channel 5	--	--	851-879

968

969

970 **Table 3.** Landsat SIC derived using the actual pair of albedo threshold values (“Actual value”) and the four variations of them
 971 (see text) averaged for 12 Landsat-8 scenes selected for the Northern Hemisphere (NH) at 25 km and 50 km grid resolution.
 972 The number to the right of the \pm denotes one standard deviation. All SIC values are in percent.

$\alpha_{\text{thinice}} \setminus \alpha_{\text{openwater}}$	-0.03	Actual value	+0.03	NH, 25km
-0.1	99.2 \pm 2.1	--	97.3 \pm 3.7	
Actual value	--	98.0 \pm 3.1	--	
+0.1	99.2 \pm 2.1	--	97.3 \pm 3.7	
				NH, 50km
-0.1	98.9 \pm 3.2	--	96.9 \pm 4.5	
Actual value	--	97.7 \pm 4.1	--	
+0.1	98.9 \pm 3.2	--	96.9 \pm 4.5	

973

974 **Table 4.** Landsat SIC derived using the actual pair of albedo threshold values (“Actual value”) and the four variations of them
 975 (see text) averaged for 15 Landsat-8 scenes selected for the Southern Hemisphere (SH) at 25 km and 50 km grid resolution.
 976 The number to the right of the \pm denotes one standard deviation. All SIC values are in percent.

$\alpha_{\text{thinice}} \setminus \alpha_{\text{openwater}}$	-0.03	Actual value	+0.03	SH, 25km
-0.1	63.0 ± 27.0	--	60.5 ± 26.4	
Actual value	--	61.5 ± 26.6	--	
+0.1	63.0 ± 27.0	--	60.5 ± 26.4	
				SH, 50km
-0.1	54.5 ± 34.8	--	52.3 ± 33.8	
Actual value	--	53.1 ± 34.1	--	
+0.1	54.5 ± 34.8	--	52.3 ± 33.8	

977

978 **Table 5.** Summary of the statistical parameters displayed in Fig. 2. Diff, DiffSDEV, and Median (all in percent SIC) are the
 979 mean difference PMW SIC minus Landsat SIC, its standard deviation and the median difference; Slope and Intercept (in
 980 percent SIC) are the coefficients of the linear regression, and R^2 and N are the squared linear correlation coefficient and number
 981 of data pairs, respectively. Numbers in **bold** and **bold italic** font denote the respective “best” and “2nd best” value, respectively,
 982 e.g. largest and 2nd-largest values of R^2 and lowest and 2nd-lowest values of Diff, Intercept and difference unity minus slope.

983

LS5, NH 2003-11	SICCI- 12	SICCI- 25	SICCI- 50	OSI- 450	CBT- SSMI	NOAA- CDR	CBT- AMSRE	NT1- SSMI	ASI- SSMI	NT2- AMSRE
Diff	-5.5	-5.4	-3.5	-4.9	0.6	0.7	-0.3	-8.4	-7.8	0.0
DiffSDEV	9.2	8.3	9.1	8.7	8.2	8.2	7.7	11.7	10.5	7.5
Median	-3.2	-3.4	-1.7	-3.3	0.0	0.0	0.0	-5.7	-6.0	0.0
Slope	0.833	0.963	0.967	0.675	0.515	0.524	0.730	0.665	0.846	0.675
Intercept	10.6	-1.9	-0.3	26.4	47.4	46.6	25.9	23.9	7.0	31.5
R^2	0.57	0.64	0.57	0.50	0.49	0.49	0.54	0.32	0.51	0.55
N	30549	8519	2748	8519	7557	7491	8384	7637	32855	8384

984

985 **Table 6.** Summary of statistical parameters shown in Fig. 3. See Table 5 for an explanation of the parameters given.

LS8, NH 2013-15	SICCI- 12	SICCI- 25	SICCI- 50	OSI- 450	CBT- SSMI	NOAA- CDR	CBT- AMSR2	NT1- SSMI	ASI- SSMI	NT2- AMSR2
Diff	-6.2	-4.7	-3.6	-4.3	1.6	1.6	0.4	-4.8	-6.0	1.2
DiffSDEV	11.0	8.2	9.0	9.8	9.9	9.8	8.0	11.4	12.2	8.1
Median	-2.8	-2.8	-2.0	-2.9	0.0	0.0	0.0	-1.5	-3.8	-1.5
Slope	0.868	0.974	0.997	0.779	0.688	0.704	0.841	0.842	0.919	0.828
Intercept	6.1	-2.4	-3.3	16.2	30.5	29.1	15.2	9.8	1.5	17.2
R^2	0.72	0.84	0.79	0.73	0.72	0.72	0.81	0.67	0.69	0.80
N	23433	6484	2056	6576	5944	5945	5831	6008	22655	5831

986

987

988

989

990

991

992

993

Table 7. Summary of statistical parameters shown in Fig. 4. See Table 5 for an explanation of the parameters given.

LS8, SH 2013-15	SICCI-12	SICCI-25	SICCI-50	OSI-450	CBT-SSMI	NOAA-CDR	CBT-AMSR2	NT1-SSMI	ASI-SSMI	NT2-AMSR2
Diff	-5.1	-5.9	-6.8	-5.3	-1.5	-1.6	-3.0	-9.5	-9.1	3.4
DiffSDEV	13.3	13.5	16.0	13.5	14.6	14.8	14.2	15.5	16.9	13.8
Median	-1.3	-2.1	-1.9	-2.8	0.0	0.0	-0.2	-7.3	-6.5	0.4
Slope	0.915	0.969	1.033	0.827	0.826	0.843	0.915	0.834	0.898	0.821
Intercept	2.1	-3.3	-9.6	9.5	13.4	11.8	4.2	4.7	-0.4	18.7
R ²	0.78	0.77	0.72	0.73	0.70	0.70	0.74	0.68	0.68	0.72
N	34331	9796	3098	9796	9788	9788	10009	9883	34252	10009

996 **Table 8.** Comparison of statistical parameters listed in Tables 5 to 7 in both hemispheres for SICCI-2 and OSI-450 products
997 using truncated or non-truncated (near-100 % SIC) PMW SIC data. See Table 5 for an explanation of the parameters given.
998 Top (LS5, NH 2003-11) is for first-year ice dominated cases, middle (LS8, NH 2013-15) is for mixed first-year / multiyear
999 and multiyear ice cases, both Northern Hemisphere; bottom (LS8, SH 2013-15) is for the Southern Hemisphere. The overall
1000 median differences do not change and are not listed again.

LS5, NH 2003-11	SICCI-12	SICCI-12 non-truncated	SICCI-25	SICCI-25 non-truncated	SICCI-50	SICCI-50 non-truncated	OSI-450	OSI-450 non-truncated
Diff	-5.5	-4.6	-5.4	-5.0	-3.5	-3.0	-4.9	-4.5
DiffSDEV	9.2	10.0	8.3	8.7	9.1	9.3	8.7	9.0
Slope	0.833	0.852	0.963	0.974	0.967	0.979	0.675	0.684
Intercept	10.6	9.6	-1.9	-2.5	-0.3	-1.0	26.4	26.0
R ²	0.57	0.54	0.64	0.63	0.57	0.56	0.50	0.48
LS8, NH 2013-15								
Diff	-6.2	-4.9	-4.7	-4.4	-3.6	-3.4	-4.3	-3.9
DiffSDEV	11.0	12.1	8.2	8.5	9.0	9.1	9.8	9.9
Slope	0.868	0.891	0.974	0.982	0.997	1.000	0.779	0.786
Intercept	6.1	5.2	-2.4	-2.7	-3.3	-3.5	16.2	15.9
R ²	0.72	0.68	0.84	0.83	0.79	0.79	0.73	0.73
LS8, SH 2013-15								
Diff	-5.1	-4.3	-5.9	-5.6	-6.8	-6.5	-5.3	-5.1
DiffSDEV	13.3	13.8	13.5	13.7	16.0	16.2	13.5	13.7
Slope	0.915	0.931	0.969	0.976	1.033	1.040	0.827	0.832
Intercept	2.1	1.6	-3.3	-3.5	-9.6	-9.9	9.5	9.3
R ²	0.78	0.77	0.77	0.77	0.72	0.71	0.73	0.73

1010 **Table 9.** Summary of statistical results obtained for three freeze-up cases in the Northern Hemisphere (NH) and for 11 freeze-
 1011 up cases in the Southern Hemisphere (SH) using Landsat 8 data. See Table 5 for an explanation of the parameters given.

NH	SICCI-12	SICCI-25	SICCI-50	OSI-450	CBT-SSMI	NOAA-CDR	CBT-AMSR2	NT1-SSMI	ASI-SSMI	NT2-AMSR2
Diff	-8.2	-8.9	-10.5	-7.7	5.0	4.6	2.6	-14.1	-12.0	4.3
Diff SDEV	13.5	10.8	17.8	13.9	18.5	18.4	12.9	20.8	21.9	13.8
Slope	0.799	0.960	0.948	0.665	0.655	0.679	0.881	0.673	0.738	0.866
Intercept	7.8	-5.7	-6.4	19.3	31.6	29.4	12.0	11.3	8.6	14.9
R ²	0.77	0.84	0.65	0.70	0.58	0.58	0.77	0.50	0.51	0.74
N	751	208	64	210	191	191	186	196	702	186
SH										
Diff	-11.8	-12.1	-7.4	-12.1	-6.3	-6.1	-6.5	-10.9	-11.4	2.1
Diff SDEV	18.1	15.9	16.1	15.1	12.1	12.1	11.8	15.3	18.1	10.6
Slope	0.839	0.915	1.027	0.861	0.965	0.971	0.977	0.953	0.982	0.943
Intercept	2.0	-4.8	-9.7	0.1	-3.3	-3.7	-4.5	-6.9	-9.8	7.0
R ²	0.66	0.72	0.75	0.73	0.83	0.84	0.84	0.75	0.72	0.86
N	1843	531	169	531	536	536	547	540	1842	547

1012

1013 **Table 10.** Summary of statistical results obtained in the Northern Hemisphere for 28 cases with first-year ice (top, LS5, NH
 1014 2003-11) and for 12 cases with mixed first-year / multiyear or multiyear ice (bottom, LS8, NH 2013-15). See Table 5 for an
 1015 explanation of the parameters shown. For SICCI-2 and OSI-450 products, we include in all rows but “N” values based on non-
 1016 truncated (near 100 %) SIC data to the right of the “/”. We omit slope and intercept because SIC data pairs cluster at 100 %.

LS5, NH 2003-11	SICCI-12	SICCI-25	SICCI-50	OSI-450	CBT-SSMI	NOAA-CDR	CBT-AMSR	NT1-SSMI	ASI-SSMI	NT2-AMSR
Diff	-4.0 / -3.0	-3.7 / -3.4	-1.5 / -1.0	-3.5 / -3.2	-0.8	-0.7	-0.9	-5.8	-6.9	-0.6
Diff SDEV	5.2 / 6.0	4.0 / 4.4	1.8 / 2.5	3.7 / 4.1	1.6	1.4	1.8	6.6	5.6	1.4
Median	-2.6 / -2.6	-2.5 / -2.5	-1.0 / -1.0	-2.4 / -2.4	0.0	0.0	0.0	-3.5	-6.0	0.0
N	7028	1978	677	1978	1940	1940	2104	1940	7633	2104
LS8, NH 2013-15										
Diff	-2.9 / -0.8	-1.5 / -0.5	-0.9 / -0.4	-1.3 / -0.3	-0.5	-0.2	-1.0	-0.3	-2.6	-0.6
Diff SDEV	4.1 / 6.2	2.2 / 3.1	1.2 / 1.7	1.9 / 3.0	1.4	0.9	3.0	0.9	2.6	2.5
Median	-0.2 / -0.2	-0.2 / -0.2	-0.3 / -0.3	-0.2 / -0.2	0.0	0.0	0.0	0.0	-2.1	-0.5
N	2659	764	242	764	714	714	723	714	2571	723

1017

1018 **Table 11.** Summary of statistical results obtained for the four high concentration cases in the Southern Hemisphere. See Table
 1019 5 for an explanation of the parameters shown. For SICCI-2 and OSI-450 products, we include in rows “Diff”, “Diff SDEV”,
 1020 and “Median” values obtained using non-truncated SIC to the right of the “/”.

LS8, SH 2013-15	SICCI-12	SICCI-25	SICCI-50	OSI-450	CBT-SSMI	NOAA-CDR	CBT-AMSR2	NT1-SSMI	ASI-SSMI	NT2-AMSR2
Diff	-0.1 / 2.5	0.0 / 2.4	0.0 / 2.7	-0.3 / 1.8	-0.7	0.1	-1.1	-0.9	-2.9	-0.1
Diff SDEV	1.7 / 2.9	0.8 / 2.3	1.2 / 2.7	2.1 / 3.1	1.7	0.7	2.0	2.6	2.5	1.2
Median	0.0 / 2.8	0.0 / 2.5	0.1 / 2.6	0.0 / 2.2	0.0	0.1	0.0	0.0	-2.4	0.0
N	978	287	93	287	288	288	302	288	973	302

1021

1022

1023 **Table 12.** Summary of statistical results obtained for 15 melt-condition cases (without melt-ponds) in the Northern
 1024 Hemisphere. See Table 5 for an explanation of the parameters shown. Numbers added to the right of the “/” for SICCI-2 and
 1025 OSI-450 products denote the results obtained using non-truncated SIC.

LS8, NH 2013-15	SICCI-12	SICCI-25	SICCI-50	OSI-450	CBT- SSMI	NOAA- CDR	CBT- AMSR2	NT1- SSMI	ASI- SSMI	NT2- AMSR2
Diff	-5.3 / -4.3	-5.1 / -4.6	-4.2 / -4.2	-4.6 / -4.3	2.2	2.4	0.2	-3.5	-4.7	1.7
DiffSDEV	10.5 / 11.2	8.9 / 9.3	9.6 / 9.6	9.5 / 9.7	9.8	9.7	7.4	10.8	12.2	8.3
Slope	0.829/0.852	0.930/0.943	0.898/0.899	0.617/0.626	0.418	0.416	0.727	0.637	0.740	0.564
Intercept	10.5 / 9.4	1.4 / 0.6	5.3 / 5.2	30.9 / 30.4	56.9	57.3	26.1	30.6	19.5	43.0
R ²	0.67 / 0.65	0.72 / 0.71	0.61 / 0.61	0.61 / 0.60	0.54	0.54	0.66	0.48	0.55	0.56
N	2926	817	266	817	817	817	795	823	3117	795

1026
 1027 **Table 13.** Summary of statistical results obtained for 45 melt-conditions cases in the Southern Hemisphere. See caption of
 1028 Table 5 for an explanation of the parameters given. Numbers added to the right of the “/” for SICCI-2 products and OSI-450
 1029 denote results obtained using non-truncated SIC.

LS8, SH 2013-15	SICCI-12	SICCI-25	SICCI-50	OSI-450	CBT- SSMI	NOAA- CDR	CBT- AMSR2	NT1- SSMI	ASI- SSMI	NT2- AMSR2
Diff	-5.0 / -4.3	-5.8 / -5.5	-8.1 / -7.8	-4.9 / -4.6	-0.4	-0.6	-2.8	-8.7	-7.8	5.1
DiffSDEV	13.7 / 14.1	13.9 / 14.1	17.1 / 17.2	14.8 / 14.9	15.6	15.6	15.4	16.4	18.6	15.9
Slope	0.888/0.903	0.951/0.958	0.983/0.991	0.750/0.754	0.772	0.794	0.895	0.791	0.859	0.824
Intercept	4.0 / 3.5	-1.8 / -2.1	-6.7 / -7.1	14.1 / 15.4	18.0	16.0	5.8	8.2	3.6	19.4
R ²	0.79 / 0.78	0.78 / 0.78	0.69 / 0.69	0.71 / 0.71	0.69	0.69	0.72	0.67	0.65	0.69
N	10214	2915	916	2915	2899	2899	2955	2929	10129	2955

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048


1049

1050

1051

9 Figures

1052



1053

1054 **Figure 1.** Location of the Landsat scenes used. Panels a) through c) Arctic; panel d) Antarctic. Note that scenes do overlap.

1055 The total number of scenes shown is 134 (a), 12 (b), 88 (c), and 134 (d).

1056

1057

1058






Figure 2. Scatterplots of PMW SIC (y-axis) versus Landsat SIC (x-axis) for all ten products for the first-year ice dominated cases from 2003-2011 in the Northern Hemisphere (Landsat-5). Black dots are individual data pairs, the black solid line is the linear regression, and the black dashed line is the identity line. Red triangles denote the mean PMW SIC computed for Landsat SIC ranges 0%-5%, 5%-15%, 15%-25%, ..., 85%-95%, 95%-100%, red bars one standard deviation of these mean values and the red dashed line is the respective linear regression line. Red squares denote the median PMW SIC for the same Landsat SIC ranges and the red solid line is the respective linear regression line. The overall mean and median difference PMW SIC minus Landsat SIC, its standard deviation, and the equation of the linear regression through the individual data pairs is shown at the top, the number N of data pairs and the squared linear correlation coefficient at the bottom of each panel.






1085


1086 **Figure 4.** Scatterplots of PMW SIC (y-axis) versus Landsat SIC (x-axis) for all ten products for 2013-2015 in the Southern
1087 Hemisphere. See Fig. 2 for a description of symbols, lines and text.

1088



1089


1090 **Figure 5.** Landsat SIC, PMW SIC, and the difference PMW SIC minus Landsat SIC (LSIC) for all ten products for a freeze-
1091 up case in the Fram Strait on September 15, 2015. The Landsat surface class map at the top left shows white: thick / snow-
1092 covered ice; grey: bare / thin ice; black: open water). The red star marks the location of Henrik Krøyer Holme station (see
1093 text). White and grey pixels are used to compute maps of gridded LSIC at 12.5 km, 25 km and 50 km, respectively (blue:
1094 outside Landsat image). A subset of SICCI-12km SIC grid cells shown at the top right illustrates the array used for the
1095 collocation. Panels in the remaining four rows show PMW SIC and PMW SIC minus LSIC for all ten products. Land is flagged
1096 brown in the SIC panels and black in the SIC difference panels; it differs between the PMW products. The land masks in the
1097 two bigger maps at the top come from the plotting routine used. LSIC maps use the land masks of the SICCI-2 products.
1098



1099

1100 **Figure 6.** Landsat SIC, PMW SIC, and the difference PMW SIC minus Landsat SIC for all ten products for a scene near the
 1101 coast during freeze-up in Pine Island Bay, Amundsen Sea, Southern Ocean, on March 12, 2014. The red star in the top left
 1102 map marks the location of the Pine Island Glacier Automatic Weather Station (see text). Some of the white patches near the
 1103 coast in this map are actually glacier ice not adequately flagged by the land mask. See Fig. 5 for more details.

1104




1105

1106 **Figure 7.** Landsat SIC, PMW SIC, and the difference PMW SIC minus Landsat SIC for all ten products for a high-
 1107 concentration scene in the Beaufort Sea, Arctic Ocean, on April 4, 2015. See Fig. 5 for a description of the maps shown.


1108

1109



1110

1111 **Figure 8.** Landsat SIC, PMW SIC, and the difference PMW SIC minus Landsat SIC for all ten products for a high-
 1112 concentration scene in the Weddell Sea, Southern Ocean, on March 12, 2015. See Fig. 5 for a description of the maps shown.
 1113




1114

1115

Figure 9. Landsat SIC, PMW SIC, and difference PMW SIC minus Landsat SIC for all ten products for a melt-condition case in the Ross Sea, Southern Ocean, on January 29, 2014. See Fig. 5 for more description of the maps shown.

1116

1117



1118
1119 **Figure 10.** Scatterplots of SICCI-50km SIC (y-axis) versus Landsat SIC (x-axis) for ice regime “leads/openings” in the
1120 Southern Hemisphere in years 2013-2015. Black dots are individual data pairs, the black solid line is the linear regression, and
1121 the black dashed line is the identity line. Red triangles denote the mean PMW SIC computed for Landsat SIC ranges 0%-5%,
1122 5%-15%, 15%-25%, ..., 85%-95%, 95%-100%, the red bars one standard deviation of these mean values; the red line is the
1123 respective linear regression line. The overall difference PMW SIC minus Landsat SIC, its standard deviation, and the equation
1124 for the linear regression using the individual data pairs is given at the top, the number N of data pairs and the squared linear
1125 correlation coefficient at the bottom of each panel. Panel a) Fully truncated SIC, all filters applied; panel b) fully non-truncated
1126 SIC, no filters applied; panel c) truncated / non-truncated SIC, GT100 and OWF applied; panel d) truncated / non-truncated
1127 SIC, GT100 and LSO applied. Blue circles mark SICCI-50km SIC values set to 0 % by the OWF; orange circles mark SICCI-
1128 50km SIC values set changed by the LSO (solid circle: SIC set to 0 %, broken circle: SIC reduced).
1129




Figure 11. Summary of all linear regression lines obtained for the comparison between Landsat SIC and PMW SIC for all ice regimes – except high-concentration ice. Columns denote, from left to right, Landsat-5 Arctic (i.e. first-year ice), Landsat-8 Arctic (i.e. mixed first-year / multiyear ice and multiyear ice), and Landsat-8 Antarctic. Ice regimes are sorted per row from top to bottom: “all” cases, “ice edge”, and “freeze-up”. Different colors and line styles denote different products as indicated. The black solid line denotes the identity line.




Figure 11. continued for ice regimes “leads and coastal openings”, “heterogeneous ice”, “melt-conditions”.




Figure 12. Illustration of the statistical parameters of the comparison between Landsat SIC and PMW SIC for all ice regimes. Rows denote, from top to bottom, first-year ice Arctic (Landsat-5), mixed first-year / multiyear ice and multiyear ice Arctic (Landsat-8), and all ice Antarctic (Landsat-8). Columns denote, from left to right, accuracy (difference PMW SIC minus Landsat SIC), precision (standard deviation of the SIC difference), and squared linear correlation coefficient. The uni-colored rows denote cases left out, either because these ice regimes are not populated (topmost row of panels) or because the retrieval of parameters did not make sense (Squared linear correlation for ice regime “high concentration”).