

Causes and Evolution of Winter Polynyas North of Greenland

Younjoo J. Lee¹, Wieslaw Maslowski¹, John J. Cassano^{2,3}, Jaclyn Clement Kinney¹, Anthony P. Craig⁴,
Samy Kamal⁵, Robert Osinski⁶, Mark W. Seefeldt^{2,3}, Julienne Stroeve^{2,7}, Hailong Wang⁸

¹Naval Postgraduate School, Monterey, California, USA

5 ²National Snow and Ice Data Center, Boulder, Cooperative Institute for Research in Environmental
Sciences, University of Colorado, Boulder, Colorado, USA

³Department of Atmospheric and Oceanic Sciences, University of Colorado, Boulder, Colorado, USA

⁴Independent Researcher

⁵RedLine Performance Solutions, College Park, Maryland, USA

10 ⁶Institute of Oceanology of the Polish Academy of Sciences, Sopot, Poland

⁷University of Manitoba, Winnipeg, Manitoba, Canada

⁸Pacific Northwest National Laboratory, Richland, Washington, USA

Correspondence to: Younjoo J. Lee (ylee1@nps.edu)

15 **Abstract.** During the 42-year period (1979-2020) of satellite measurements, **four major winter**
(**December-March**) polynyas have been observed north of Greenland: **one in December 1986 and the**
rest of them in the last decade, i.e. February of 2011, 2017 and 2018. The 2018 polynya was
unparalleled by its magnitude and duration compared to the **three** previous events. Combined with the
limited weather station and remotely-sensed sea ice data, a fully-coupled Regional Arctic System Model
20 (RASM) hindcast simulation was utilized to examine the causality and evolution of these **extreme**
events. We found that neither the accompanying anomalous warm surface air intrusion nor the ocean
below had an impact on the development of these winter open water episodes in the study region (i.e.,
no significant ice melting). Instead, the extreme atmospheric wind forcing resulted in greater sea ice
deformation and transport offshore, accounting for the majority of sea ice loss. Our analysis suggests
25 that strong southerly winds (i.e., northward wind with speeds **greater than 10 m/s**) blowing persistently
for at least 2 days or more, were required over the study region to mechanically redistribute some of the
thickest sea ice out of the region and thus to create open water areas (a latent heat polynya). In order to
assess the role of internal variability versus external forcing of such events, we additionally simulated
and examined results from two RASM ensembles forced with output from the Community Earth
30 System Model (CESM) Decadal Prediction Large Ensemble (DPLE) simulations. Out of 100 winters in
each of the two ensembles, initialized 30 years apart, one in December 1985 and another in December
2015, respectively, **25** and **16** winter polynyas were produced north of Greenland. The frequency of
polynya occurrence **had** no apparent sensitivity to the initial sea ice thickness in the study area pointing
to internal variability of atmospheric forcing as a dominant cause of winter polynyas north of
35 Greenland. We assert that dynamical downscaling using a high-resolution regional climate model offers
a robust tool for process-level examination in space and time, synthesis with limited observations, and
probabilistic forecasts of Arctic events, such as the ones being investigated here and elsewhere.

Comment [MW(2R1): I second that.

Deleted: over

Deleted: only three

Deleted: ever

Deleted: and

Deleted: they all

Deleted: occurred

Deleted: two

Deleted: recent

Deleted: of

Deleted: 17

Deleted: 14

Deleted: over

Deleted: and

1. Introduction

The Arctic has experienced amplified warming, both through the enhancement of global temperature rise, as well as through the reductions in sea ice and snow cover that impact the regional energy budget (Serreze and Francis, 2006). This is commonly referred to as Arctic amplification (AA). On a seasonal basis, Arctic winter warming (AWW) exceeds summer warming by about a factor of four (Bintanja and van der Linden, 2013). In addition to sea ice variations, changes in the atmospheric circulation have been linked to an increase in frequency and duration of winter warming events in the Arctic (Graham et al., 2017). The trends in AWW and winter sea ice extent (SIE) have continued over the satellite record, with the March SIE decline rate of ~2.6% per decade for the period of 1979-2020, including the four lowest winter maxima of SIE in 2015-2018 and the lowest ten during 2005-2019. While SIE reductions in winter are less than ones in summer (Stroeve and Notz, 2018), there is clear supporting evidence that the winter ice pack has thinned and the amount of multi-year sea ice has declined (Kwok et al., 2009; Meier et al., 2014; Kwok, 2018; Ricker et al., 2021). It is expected that the resulting younger and thinner ice is more susceptible to atmospheric wind forcing (Sprenn et al., 2011; Itkin et al., 2017), yielding increased sea ice drift speed, enhanced fracturing and more lead openings (Rampal et al., 2009). Hence, along with the current trend in the Arctic sea ice toward younger and thinner ice, it can be hypothesized that polynyas may become more prevalent in recent years. Since about half of the total atmosphere-ocean heat exchange over the Arctic Ocean may occur through leads and polynyas in winter (Maykut, 1982), the occurrence and formation of polynyas could play a crucial role in the alteration of regional climate (Morales Maqueda et al., 2004).

In late February 2018, satellite imagery revealed an unusual open water polynya north of Greenland between the Lincoln and Wandel seas (Fig. 1a). This event received considerable attention not only because it was claimed to be a one-of-a-kind extreme event involving some of the thickest Arctic sea ice but also because its emergence coincided with anomalous, above freezing, warming of surface air temperature over the region [after the sudden stratospheric warming \(SSW\) event observed in mid-February 2018](#) (Moore et al., 2018). The presumed contribution of this warm surface air to the polynya opening is of interest and so are its causality, evolution and past occurrences. Yet, detailed *in situ* observations of polynyas are limited due to their intermittency and restricted access in winter. Hence, in addition to satellite measurements and weather data, fully coupled climate models become critical tools in studying such events (e.g., Ludwig et al., 2019). However, for the majority of global climate models (GCMs), many coastal polynyas are at sub-grid scales; thus, the GCM utility for comprehensive polynya studies are impeded (Weijer et al., 2017). In addition, GCMs are not intended to represent specific climate events in space and time and they are not suitable for process-level investigation of such events and quantification of their impact at both local and larger scales.

On the other hand, regional climate models (RCMs) used for dynamical downscaling are expected to show improvement in reproducing extreme weather events, given that their atmospheric boundary conditions for simulations of the past to present are derived from global atmospheric reanalysis such as Climate Forecast System (CFS) version 2 (CFSv2; Saha et al., 2011). Therefore, high-resolution RCMs, such as the Regional Arctic System Model (RASAM; e.g. Maslowski et al. 2012), offer unique

Deleted: over

95 capabilities for examining the spatio-temporal development and impact of observed specific events like
a polynya (Fig. 1b), in the context of a fully-coupled climate system model (atmosphere-sea ice-ocean-
land), while CFSv2 is shown to be less skillful in simulating such an event (Fig. 1c). In addition, RCMs
afford ensemble sizes prohibitive to their global fine-resolution counterparts, which is often a
requirement to distinguish the forced response from internal model variability (e.g., Peings et al. 2021).

100 The 2018 winter polynya north of Greenland has been investigated by Moore et al. (2018) using an ice-
ocean model forced by surface atmospheric reanalysis. They have established the dominant role of
surface winds in generating this polynya. Here, by taking advantage of the fully-coupled and high-
resolution RASM hindcast simulation, combined with weather station and satellite sea ice data, we
105 evaluate the capability of a RCM in reproducing the observed natural phenomena (Fig. 2) and
investigate the coupled mechanisms involved in the development of northern Greenland coastal
polynyas within some of the thickest Arctic ice-pack cover. In particular, this study focuses on analysis
of historical winter (December-March) polynya events for the full period of satellite data availability
(1979-2020) to diagnose the relative roles of thermodynamic and dynamic processes and to assess
required forcing changes over the last four decades. Furthermore, by dynamic downscaling of the
110 Community Earth System Model (CESM)-Decadal Prediction Large Ensemble (DPLE), two sets of 30
years apart RASM ensemble (termed RASM-DPLE) simulations are performed to examine the relative
roles of internal variability and external forcing influencing the development of winter polynyas under
two different sea ice regime scenarios: i.e. a thicker ice in the 1980s versus a thinner ice in the 2010s.
We provide details of the satellite and weather station data used for this study in section 2 and the model
115 setup for the hindcast and DPLE simulations are described in section 3. Next, section 4 presents a
synthesis of observed and modeled past winter polynya results and examines the statistics of polynya
occurrence and the required conditions for their generation in the RASM-DPLE simulations. This is
followed by the discussion in section 5 and the study is summarized in section 6.

Deleted: over

2 Data

2.1. Surface air temperature and wind

120 Hourly surface air temperature (3-hourly prior to 2015) data were obtained from the seven weather
stations (World Meteorological Organization station identifier: 04221, 04254, 04285, 04351, 04330,
04312 and 04301) around Greenland (data available at <https://rp5.ru>) and then averaged daily for
January-March of 2011, 2017, and 2018. Since wind data were incomplete at the closest weather station
125 from the center of the polynya in 2018 (i.e., Station 04301, Cape Morris Jessup at 83° 39' N and 33° 22'
W), we used 3-hourly surface wind data from the adjacent Station 04312 (Station Nord at 81°43' N and
17°47' W), which were originally binned for 16 wind directions. Also, ERA-Interim atmospheric
reanalysis of 6-hourly 10 m wind fields (Dee et al., 2011;
<https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era-interim>) was used for comparison
with the RASM hindcast simulation over the study region.

130

2.2. Sea ice concentration and thickness

Daily sea ice concentration (SIC) data were obtained via the National Snow and Ice Data Center (NSIDC): <https://nsidc.org/data/G02202/versions/4> for December 1978– March 2020 (Meier et al. 2021). Satellite-derived SIC used for this study is based on passive microwave measurements using the NASA team (NT) algorithm (Cavalieri et al., 1984); all the data are on a polar stereographic 25 km×25 km grid. The daily SIC was used to examine the occurrence of a polynya in the satellite measurements. We later detected observed polynya events when the daily averaged satellite SIC dropped below 90% over the study region. RASM sea ice thickness (SIT) was also compared with the CryoSat-2/the Soil Moisture Ocean Salinity (SMOS) satellite merged data that are only available in winter months (Ricker et al., 2017) as well as CFSv2 reanalysis (<https://doi.org/10.5065/D61C1TXF>). Due to the lack of persistent SIT observations over the Arctic, the Pan-Arctic Ice Ocean Modeling and Assimilation System (PIOMAS) is often considered as an “observational” proxy (Zhang and Rothrock 2003; Schweiger et al. 2011; Stroeve et al. 2014). The PIOMAS (version 2.1) sea ice data were retrieved from the Polar Science Center at the University of Washington (<http://psc.apl.uw.edu/research/projects/arctic-sea-ice-volume-anomaly/data/>).

Deleted:	https://nsidc.org/data/nsidc-0051
Deleted:	9
Deleted:	19
Deleted:	and https://nsidc.org/data/nsidc-0081 for 2020
Deleted:	mean
Deleted:	detect
Deleted:	found out
Deleted:	it
Deleted:	from 2011
Deleted:	https://www.meereisportal.de ; Grosfeld et al., 2016

3 Method

3.1 Regional Arctic System Model

RASM is a limited-area, fully-coupled climate model consisting of the Weather Research and Forecasting (WRF version 3.7.1) for atmosphere, the Los Alamos National Laboratory Sea Ice Model (CICE version 5.1.2) for sea ice and Parallel Ocean Program (POP version 2.1) for ocean, the Variable Infiltration Capacity (VIC version 4.0.6) land hydrology and routing scheme (RVIC version 1.0.0) for land (see Maslowski et al. 2012; Roberts et al. 2015; DuVivier et al. 2015; Hamman et al. 2016; Hamman et al. 2017; Cassano et al. 2017). All the components are coupled using the Craig et al. (2012) version of CESM flux coupler. RASM is configured over a pan-Arctic domain, including the entire Northern Hemisphere marine cryosphere and all terrestrial drainage basins that drain to the Arctic Ocean. The ocean and sea ice components share a 1/12-degree (~9 km) rotated spherical grid and are configured with 45 levels in the vertical and five ice thickness categories, respectively. The atmosphere and land hydrology components are set up on a 50-km polar stereographic mesh with the vertical resolution of 40 levels and 3 soil layers, respectively. For the hindcast simulation (September 1979–present), CFS Reanalysis (CFSR)/CFSv2 are dynamically downscaled to provide RASM-WRF with atmospheric lateral boundary conditions and for linearly increasing grid nudging of winds and temperature for the top half of the model domain, approximately above 540 hPa. The RASM-POP ocean temperature and salinity along the closed lateral boundaries are restored to monthly Polar Science Center Hydrographic Climatology version 3.0 (PHC 3.0; Steele et al., 2001). The initial conditions at the beginning of the hindcast simulation are derived from the 32-year spin-up of the ocean-sea ice model forced with the Common Ocean-Ice Reference Experiment Inter-Annual Forcing version 2 (CORE2-IAF; Large and Yeager, 2009) atmospheric reanalysis for 1948-1979.

Deleted:	e
Deleted:	ice

The RASM-DPLE simulations are derived by dynamically downscaling global atmospheric output from the initialized CESM-DPLE simulations (Yeager et al., 2018). Output from the two 10-member decadal ensembles, initialized on December 1st of 1985 and 2015, was selected for in-depth analysis under different regimes of the Arctic SIT distribution. Each RASM-DPLE simulation is initialized with the ocean and sea ice conditions, with thinner ice in the latter period (Fig. S1), from the RASM hindcast and integrated for 121 months with CESM-DPLE atmospheric forcing. The size of each ensemble (10 members) is determined by the availability of archived CESM-DPLE output necessary for RASM-WRF boundary conditions. Hence, output for 100 winters per each ensemble allows for statistical analysis of polynya occurrence in the past and near future.

3.2 Self-organizing maps

The self-organizing map (SOM), an artificial neural network based on a competitive learning algorithm (Kohonen, 2001), has been widely used to visualize input data vectors onto a low dimensional map of nodes and to objectively classify complex data sets in meteorology and oceanography (see Liu and Weisberg, 2011). For January-March 2018, the RASM/WRF pan-Arctic (>65 °N) 6-hourly surface wind fields (i.e., 10 m U and V components) are characterized using a SOM [4×4] map grid (i.e., 16 nodes or patterns). Time-dependent spatial features of near surface winds are identified and frequencies of occurrence of wind patterns favorable for a polynya are quantified using the SOM Toolbox 2.0 for MATLAB available at <http://www.cis.hut.fi/projects/somtoolbox>.

4 Results

4.1 2018 winter polynya – a case study

4.1.1 Near-surface air temperature around Greenland

Focusing on the most recent and the largest winter polynya event, daily near-surface air temperatures were examined from the weather stations around the Greenland coast for January-March 2018 (Fig. 3). A significant warming event (air temperatures rising above 0 °C) was observed over the northeastern Greenland region from mid-February to early March and captured well in the RASM simulation. This anomalous warming was most prominent along the northern Greenland coast (Figs. 3e and 3f), where the polynya was observed and simulated in RASM (Figs. 2a and 2d, respectively), but less pronounced over the mid-eastern Greenland coast (Figs. 3g and 3h). In contrast, no warming was measured in the southwestern stations for the same period (Figs. 3b-3d). This anomalous warming, with relative humidity rising above 90% (i.e., Station 04312; not shown), coincided with a strong reversal of the Arctic Oscillation (AO) index (https://www.cpc.ncep.noaa.gov/products/precip/CWlink/daily_ao_index/ao.shtml) from a positive to a strongly negative phase (Fig. 3a). Among the northeastern stations, the observed near-surface air temperature was positively correlated to the AO index (Fig. 3i). The maximum correlation coefficient (r) was time-lagged up to 11 days at the northern stations and only 3-4 days at the mid-eastern stations (Fig. 3j). Given that the anomalous warming started a few days earlier at the mid-eastern stations and

220 ceased a few days later at the northern stations, suggests the advection of warm air masses from the south. The prolonged warming at the northern stations could also be partly due to the release of oceanic heat from the polynya to the atmosphere.

Deleted: it may

225 On the other hand, near-surface air temperatures were inversely correlated with the AO index (Fig. 3i) in the southwestern Greenland region, with a shorter time-lag (one to two days) for a maximum correlation coefficient (Fig. 3j). We found that no anomalous warming was present in the southwestern Greenland region during February. In fact, near-surface air temperatures tended to gradually decrease from January through February, with the lowest temperatures observed between 22nd and 24th of February 2018 (Figs. 3b-3d). Thereafter, near-surface air temperatures rapidly increased at all southwestern stations, peaking in early March, corresponding to the strong negative AO phase, and the 230 second warming came approximately two to three weeks later.

235 Overall, the RASM hindcast simulation captured remarkably well the sudden increase of near-surface air temperatures in northeastern Greenland and its gradual decrease. RASM also reproduced well the cooling in February and then the warming over southwestern Greenland, in terms of its magnitude and spatio-temporal variability (Fig. 3). One discrepancy in the RASM simulation against the weather stations data was a positive bias of near-surface air temperatures at Stations 04221 (Fig. 3b) and 04330 (Fig. 3g), possibly linked to the relatively coarse horizontal resolution of the RASM atmospheric component (i.e., 50 km), which is insufficient for resolving strong temperature gradients across the ocean/land/ice sheet boundary or the fidelity of the near surface temperature distribution over the 240 Greenland Ice Sheet.

Deleted: , timing,

4.1.2 Sea ice dynamics

245 The sudden anomalous warming over northern Greenland with temperatures above 0 °C was an extreme phenomenon in 2018, considering that the long-term mean (2011-2020) of February surface air temperature is -27.8 °C at Station 04301 (Fig. 3e; Cape Morris Jesup). In addition, there were other years of anomalous warming (Figs. 4b and 4c), albeit less pronounced, when previous, smaller, polynya events occurred during February 2011 and 2017 (Figs. 2b and 2c, respectively). RASM's realistic representation of the polynya, as well as the magnitude and timing of anomalous warming, grants confidence to the examination of relative contribution of thermodynamic ice melt to the generation of the polynya, [although the center of the simulated polynya is not exactly collocated with the observed one](#). 250 Figure 5 shows the RASM thermal sea ice surface, lateral and bottom melting terms were all negligible (< 1 cm) over the study region when integrated for the whole month of February 2018. Hence, the dramatic rise of near-surface air temperatures by more than 25 °C above climatology and their persistence around the freezing point for several days (Fig. 3e), had no impact on sea ice melt, nor on the preconditioning and development of the polynya north of Greenland. In agreement with Moore et al. (2018), we corroborate that this polynya was driven by mechanical redistribution of sea ice outside 255 of the study region (see Fig. 4a): i.e., this was a latent heat polynya. Based on RASM results, we calculate that between 15 and 25 February 2018, 192 km³ of sea ice was dynamically transported outside the study region (Fig. 4d). During two weeks prior to the 2018 polynya event, a mean

Deleted: The

Deleted: although the center of the simulated polynya is not exactly located as the observed one,

thermodynamic ice growth over the region was 0.72 km³/day (Fig. 4d), which is comparable to the rate in a non-polynya year such as 2019 (not shown). The peak sea ice growth of 3.1 km³/day occurred on 26 February right after the maximum daily dynamic ice removal and the anomalous warming period. However, this large sea ice removal during the polynya formation period was not fully replenished in the region by the end of March, even with dynamic and thermodynamic processes adding 81 km³ and 42 km³, respectively, after 26 February. Overall, the RASM integrated thermodynamic ice growth in the study region during February 2018 was approximately 50% higher (31.2 km³; Fig. 4d) due to the rapid ice growth during the polynya opening, compared to the ice growth during a non-polynya year (i.e., 20.8 km³ in February 2019; not shown).

Deleted: due to

4.1.3 Atmospheric-sea ice coupling

As dynamic processes dominated the overall winter sea ice in the study region, we have analyzed the wind data from the weather station (Station 04301; Station Nord) and have found that the polynya development was associated with strong and persistent winds from the south-southeast (Fig. 4g). We further examined how the spatial near-surface wind fields evolved over the time period associated with the sea ice divergence. The SOM analysis extracted 16 patterns from the total of 360 synoptic wind fields: the 6-hourly RASM 10 m U- and V-wind components during January-March 2018. Figure 6 shows the four major wind patterns most frequently identified (more than 77% occurrence) from the pre-polynya period (5 February 2018) until closing of most of the open water areas (13 March 2018). The RASM hindcast simulation confirmed that this polynya event was predominantly associated with southerly to southeasterly winds blowing over the northern Greenland region (Fig. 6b and 6c), consistent with the ERA-Interim reanalysis of 10 m wind fields (Fig. S2). Prior to the polynya event in 2018, surface winds were mainly from the north or northwest over the region (Fig. 6a), which yielded little or no sea ice divergence (Fig. 6e). When the major wind pattern shifted to between southerly to southeasterly winds over the northern Greenland region on 15 February 2018 (Fig. 6b), sea ice started to deform and diverge significantly (Fig. 6f). Beginning on 20 February 2018, the southeasterly wind became even more prominent, with 19% stronger wind speed over the northern coast of Greenland (Fig. 6c), which increased the sea ice deformation rate further and led to the maximum polynya opening (Fig. 6g). The largest observed and modeled polynya areas were identified on 25 February 2018 (see Figs. 2a and 2d, respectively). Thereafter, within a week, the shift of wind patterns in late February (Fig. 6d) reversed the dynamic sea ice volume (SIV) tendency from net loss to gain (Fig. 4d) and subsequently reduced the deformation rate back to nearly zero by early March (Fig. 6h)

4.2 Winter polynyas during 2010s

Deleted: events before 2018

4.2.1 February 2011 and 2017

Upon examining satellite-derived SIC over the data record between 2010 and 2020 (Figs. 7 and S3), we found that there were two additional polynya events in the same month, albeit smaller, over the northern Greenland region observed in February 2017 and 2011 (see Figs. 2b and 2c, respectively). Here, we found those polynya events when the daily mean satellite-derived SIC fell down to or below 90% over

Deleted: entire

Deleted: 1979

Deleted: ,

Deleted: , and S4

Deleted: that a

Deleted: occurred

310 the region (see Figs. 7a, 7g, and 7h). As was the case of the 2018 polynya, both the previous events
 also coincided with anomalous warming, peaking on 12 February 2011 (Fig. 4b) and on 8 February
 2017 (Fig. 4c), as measured at Station 04301 (Cape Morris Jesup). Near-surface air temperature
 315 hindcast simulation (Figs. 2e and 2f) and thus we investigated their causality with respect to their
 relative SIV reductions due to dynamical processes and/or thermodynamic ice melt (Figs. 4e and 4f).
 The RASM results confirmed that those were latent heat polynyas dominated by the mechanical
 redistribution of sea ice out of the region. Their sizes were smaller compared to the 2018 polynya, with
 320 the one in 2017 even smaller than the one in 2011, which we attribute to somewhat different wind
 patterns such as its direction, magnitude and duration (Figs. 4i and 4h). Table 1 shows that, during the
 polynya periods based on the RASM simulation (defined here when ice volume tendency is less than –
 10 km³/day, for more than 3 days), i.e., 12-15 February 2011 and 8-10 February 2017, 55 km³ and 42
 325 km³ of sea ice was dynamically removed outside the study region, respectively, which is much less
 compared to the ice loss of 189 km³ during the 2018 event (16-25 February 2018). The size of a
 polynya was proportional to the pseudo-wind stress (i.e., wind speed squared) integrated over the
 polynya period (Table 1). Thus, more turbulent (latent plus sensible) heat was lost during the 2011
 winter polynya (daily mean of –96.7 W/m² and maximum of –125 W/m²) when its size was bigger
 (Table 2). In addition, referenced to the February ice growth during a non-polynya year (for example,
 330 20.8 km³ in February 2019; not shown), the RASM thermodynamic ice growth integrated over the
 month of February was elevated: 33% higher in 2011 (27.6 km³; Fig. 4f) in the study region. Note that
 the daily mean turbulent heat flux in the study region was –48 W/m² during the 2018 winter polynya
 event with the maximum daily heat loss up to –182 W/m². However, the total daily turbulent heat loss
 was much larger than any other year because of the size and duration of open water areas (Table 2).

Deleted: , which occurred for three consecutive days in 2011, one day in 2017, and 10 days in 2018
Deleted: respectively

Deleted: negative

Deleted: .

4.2.2 What is driving changes in polynya frequency

335 Given the above analysis, an outstanding question is why winter polynyas became more frequently
 observed during the 2010s within the past four decades. Since observational data around northern
 Greenland are incomplete, we expanded the analysis of surface wind fields (10 m U and V components)
 from the RASM hindcast simulation near Station 04312 (Fig. 8a; Station Nord) and Station 04301 (Fig.
 8b; Cape Morris Jesup). Our analysis revealed that northward wind (blowing from south), required for
 340 opening of a winter polynya along the coast of northern Greenland, has recently become more frequent,
 stronger (i.e., wind speed >10 m/s), and more persistent (i.e., blowing for at least 2 consecutive days or
 longer). In addition, the three years of winter polynya occurrence with the wind conditions satisfied the
 above criteria. Based on the RASM hindcast simulation, the grid mean wind conditions in February
 2009 were similar to the ones in February 2017 near Station 04312 (Fig. 8a), but a notable polynya was
 345 not detected, possibly owing to the influence by such wind conditions over a smaller area within the
 main polynya region. The observational data also indicate a polynya favorable wind condition in 2009:
 i.e., relatively warmer air (–12.6 °C) blowing from the south-southwest with the maximum wind speed
 of 19 m/s on February 7th, 2009 at Station 04312 (Table S1). But, the satellite-derived mean daily SIC
 only dropped down to 94% in February 2009 (Fig. S4o) because the wind was possibly weaker,

Formatted: Font:(Default) +Theme Headings (Times New Roman)
Formatted: Heading 2
Formatted: Font:(Default) +Theme Headings (Times New Roman)
Formatted: Font:(Default) +Theme Headings (Times New Roman)
Deleted: was
Deleted: only
Deleted: for the whole period of 1980-2020
Deleted: satisfying
Deleted: , all coincidentally were La Niña winters (see https://origin.cpc.ncep.noaa.gov/products/analysis_monitoring/ensos/tu/fi/ONI_v5.php)

compared to the wind condition in February 2017 (Fig. 4i). Note that the early February 2009 data at Station 04312 are missing and the entire February 2009 data at Station 04301 are completely unavailable. Although changes in wind (i.e., direction and intensity) play a role in this region, we cannot completely rule out that a thinning of sea ice may promote more frequent polynyas in recent years. Hence, we additionally simulated two large ensembles and evaluated model representation of polynyas in section 4.4.

- Deleted: it
- Deleted: more frequently
- Deleted: the

4.3 Winter polynyas between 1980s and 2000s

It is interesting to point out that no major winter polynya was found during the 1980s to 2000s (Figs. S3 and S4) except one major event in December 1986; the lowest daily mean SIC was 82% over the study region on 15 December 1986 (Fig. S4i). There were two instances indicating possible polynya events because the regional daily mean SIC was below 90%, i.e., December 1984-January 1985 (Fig. S4g) and December 2002-January 2003 (Fig. S3i). However, they were not considered polynyas in this study since sea ice dynamic volume tendency (DVT) was not below the threshold ($\leq -10 \text{ km}^3/\text{day}$) for at least three consecutive days (Fig. S5). Compared to the recent polynyas in the 2010s, the polynya in December 1986 is larger than ones in 2011 and 2017 but smaller than one in 2018. The satellite SIC shows that the polynya occurred in a similar location to the ones in the 2010s (Fig. 9a) and the RASM simulation also captured it well (Fig. 9b). The RASM simulation confirms that the December 1986 event was not associated with significant ice melting (Fig. 9c) although it was slightly elevated, compared to the 2018 event (Fig. 5). Analogous to the recent 2010s polynyas, the December 1986 polynya is linked to a strong southerly wind of almost the same strength as the 2018 winter polynya (Fig. 9d and Table 2), but its duration was shorter (the strong southerly wind only lasted 5 days; Fig. S6). Other features in common are that anomalous atmospheric warming occurred during the polynya and that the near-surface air temperature was positively correlated ($r=0.45, p<0.05$) to the AO index (Fig. 9e). The correlation was lagged by 11 days, similar to the event in February 2018. However, no SSW was reported during winter 1986-1987 (see Butler et al., 2017) and it was an El Niño winter in contrast to the polynyas of the 2010s. The RASM hindcast simulation suggests that, between 12 and 16 December 1986, 142 km^3 of sea ice was dynamically transported outside the study region (Fig. 9f), which is 25% less than the amount of ice removed in 2018, and this was 2.6 times more than the ice transported in February 2011 (Table 1). However, the mean turbulent heat flux was much less in December 1986 than in February 2011 even though it was a larger event in terms of polynya size and wind intensity. This is possibly due to the fact that sea ice was thicker in 1986; for example, the mean SIT was 4.4 m for 5 days before the polynya (Table 1). Due to large open water areas in December 1986, the integrated turbulent heat loss was much larger compared to the polynyas in February 2011 and 2017.

- Formatted: Font:Bold
- Formatted: Font:Bold
- Formatted: Font:Bold
- Deleted: from
- Deleted: as a
- Deleted: bigg
- Deleted: like
- Deleted: his
- Deleted: Similar
- Deleted: it
- Deleted: relatively
- Deleted: h
- Deleted: than
- Deleted: one in February 2018
- Deleted: that
- Deleted: , for example,
- Deleted: ceased by 20 December 1986
- Deleted: (
- Deleted: ones in

4.4 Polynyas in a large ensemble of initialized decadal prediction simulations

According to the merged CryoSat-2/SMOS data, the mean SIT over the region does not exhibit a negative trend in February. For example, in the first week of February, the mean SIT was 2.74 m in 2011, 3.16 m in 2017, and 2.73 m in 2018. On the other hand, the RASM hindcast simulation indicates

- Deleted: 3

425 a gradual thinning of sea ice in the region and a long-term trend of SIT was -0.26 m/decade for all
 months during 1985-2019 (not shown). In order to further investigate the potential role of regional sea
 ice thickness reduction versus internal variability in the occurrence of winter polynya events, north of
 Greenland, we performed and examined two RASM 10-member ensembles forced with atmospheric
 output from the CESM-DPLE simulations. They are initialized 30 years apart, i.e. in December 1985
 430 and 2015, respectively, to represent different SIT conditions over the study region, with the former
 corresponding to a thicker ice regime (Fig. S1a; mean SIT of 3.3 m) and the latter to a thinner ice
 regime (Fig. S1b; mean SIT of 2.3 m). Note that the PIOMASS SIT also corroborates that sea ice was
 1.1 m thicker north of Greenland in November 1985 (Fig. S1d; mean SIT of 3.3 m) than in November
 2015 (Fig. S1e; mean SIT of 2.2 m). Each ensemble member is integrated forward for 10 years, thus
 435 resulting in 100 winters (i.e., 10 ensemble members for each 10-year simulation) for statistical analysis
 of polynya occurrence in each of the two ensembles. Note that because of the setup of these
 experiments (i.e. using the CESM DPLE atmospheric output as boundary conditions for forcing the
 RASM-DPLE **fully-coupled** simulations), we only compared the probability of polynya occurrence
 during winter months (~~December~~-March) instead of focusing on accurate temporal representation of
 such events in a similar location.

Deleted: over

440 Taking as the baseline the winter polynya in February 2017 (Fig. 4e and Table 1), which was the
 smallest among the three observed events, we defined “a latent heat polynya” in the RASM-DPLE
 ensemble members when a daily winter sea ice loss due to dynamic processes was greater than 10
 km^3/day for at least three consecutive days over the study area (see Fig. 4a). Note that the ~~three~~ other
 445 observed polynya events experienced ~~4~~, ~~5~~ and 10 consecutive days of dynamic sea ice loss greater than
 $10 \text{ km}^3/\text{day}$ during February ~~2011~~ (Fig. 4f), ~~December 1986~~ (Fig. 9f) and ~~February 2018~~ (Fig. 4d),
 respectively (Table 1). Analogous to the RASM hindcast simulation, Tables 3 and 4 list all the polynya
 occurrences from the two ensemble runs. We found ~~25~~ polynyas in the 1985-initialized ensemble (Table
 3) and ~~16~~ polynyas in the 2015-initialized ensemble (Table 4), out of the 100 winters each. Note that
 450 some ensemble members simulate more than 1 polynya event while some had no polynyas at all. The
 majority of the polynya events (~~84%~~ in the 1985-initialized and ~~69%~~ in the 2015-initialized runs) were
 similar in size to the smaller polynyas that were simulated in February of 2011 ~~or 2017~~. The longer-
 lasting polynyas (with dynamic sea ice removal $\geq 10 \text{ km}^3/\text{day}$ for five consecutive days or longer) were
 455 also produced in the RASM-DPLE simulations: ~~four~~ incidents in the 1985-initialized and ~~five~~ incidents
 in the 2015-initialized runs.

Deleted: January

Deleted: two

Deleted: of

Deleted: 17

Deleted: 14

Deleted: 88

Deleted: 64

Deleted: and

Deleted: of

Deleted: or greater

Deleted: two

Deleted: four

Deleted: But none of the ensemble members produced the total integrated turbulent heat flux as large as the 2018 polynya (Table 2).

Deleted: largest

Deleted: 9a

Deleted: 9b

Deleted: 9c

Deleted: 9d

Deleted: 9e

Deleted: 9f

We finish this analysis of RASM-DPLE results by comparing the ~~longest~~ polynya detected in each
 ensemble (see Table 3 and 4), i.e., from the ensemble member #4 in January 1988 (Fig. ~~10a~~) and the
 ensemble member #2 in January 2024 (Fig. ~~10b~~). As in the case of observed events, these latent heat
 460 polynyas were created due to dynamic sea ice transport ($\text{DVT} \leq -10 \text{ km}^3/\text{day}$ for seven consecutive
 days), resulting in significant sea ice removal of -120 km^3 out of the region in 1988 (Fig. ~~10c~~; Table 3)
 and -136 km^3 in 2024 (Fig. ~~10d~~; Table 4). By cross-examining wind patterns over northern Greenland
 (i.e., near Cape Morris Jesup), our analysis confirmed that these large polynya openings were also
 465 associated with very strong and persistent southerly winds in the RASM-DPLE simulations (Figs. ~~10e~~
 and ~~10f~~). We found that daily DVT and meridional wind were significantly correlated, and the size of

490 polynyas was highly dependent on the pseudo-wind stress integrated over the polynya period (Table 3
and 4). However, given the wind patterns, their duration and the integrated sea ice removal (-189 km^3)
during the observed polynya in February 2018 (Table 1), this event stands out within all the RASM
results analyzed in this study. Compared to the 2018 event, the slight difference in the magnitudes
between the largest polynyas of each ensemble still does not imply much significance of SIT in their
generation and evolution. Overall, the more frequent winter polynyas, produced in a thicker sea ice
regime between the two 30-year apart ensembles, implies that changes in SIT are not significant
495 contributors (at least up to now) to the generation of such events for this region during wintertime.
Therefore, the findings support that polynyas becomes prevalent when southerly winds are more
persistent and stronger in northern Greenland.

- Deleted: a
- Deleted: similar
- Deleted: number of
- Deleted: gets
- Deleted: the

5 Discussion

500 The recent studies (i.e., Moore et al., 2018; Ludwig et al., 2019) used an ice-ocean model to study the
polynya in February 2018, which means that the model is prescribed with reanalysis or gridded products
on every grid cell. On the other hand, RASM is a fully-coupled high resolution regional model, which
allows us to further study interactions between ice, ocean, and atmosphere. When RASM is used for
dynamical downscaling, atmospheric forcing is prescribed only along the lateral boundaries and nudged
above approximately 500 hPa. Hence, surface atmospheric forcing is predicted every coupling time
505 step. Although the RASM near surface wind fields agree well with the reanalysis data, it is possible that
a slight discrepancy in wind direction or magnitude near the study region may have shifted the center of
the polynya more westward than was observed. Although the RASM hindcast simulation relies on the
CFSv2 atmospheric conditions, the winter SIT on February 25th, 2018 is very unrealistic in CFSv2 with
too thick ice in the central Arctic. This is a common bias in current climate models (i.e., Watts et al.,
510 2021), resulting from limitations in or lack of representation of key physical processes, partly due to
their coarse model resolution (e.g., Chassignet et al., 2020). Since the CFSv2 wind and temperature are
used only above approximately 500 hPa, the dynamic downscaling approach using the high-resolution
fully-coupled RASM allows resolving the fine-scale processes and provides valuable insight into the
mechanism of the generation and evolution of winter polynyas off the northern coast of Greenland. We
515 examined the occurrence of such events in RASM against the satellite observations in winter months
(December-March) over the past four decades (1979-2020). The results from the RASM hindcast
simulation suggest that the size of a winter latent heat polynya in this region is sensitive to the direction,
magnitude and duration of near-surface winds.

- Formatted: Font color: Text 1
- Deleted: ly
- Deleted: ed
- Deleted: s
- Deleted: are sub
- Deleted: 4
- Deleted: er
- Deleted: to
- Deleted: :
- Deleted: one of
- Deleted: es
- Deleted: 4
- Deleted: T

520 Subject to the limited sample sizes, we find that the generation of a winter polynya in this region
requires strong southerly winds (i.e., speeds $\geq 10 \text{ m/s}$ and lasting for more than 2 days largely over the
study region based on the RASM hindcast simulation). Table 1 and Figure 8 show that the stronger and
more persistent the southerly wind blows, the larger the winter polynya becomes regardless of ice
conditions in the study region, but it is not immediately clear what causes such changes in wind
525 patterns. At the same time, it is reported that the southerly winds might reduce sea ice export through
Fram Strait by a slowdown of sea ice drift (Wang et al., 2021). The RASM simulation captures such

- Deleted: greater than at least
- Deleted: ; however, there were no such extreme weather events before 2010
- Deleted: On the other hand

decline in 2018 and overall agrees with the observed interannual variability of ice export through the Fram Strait (Fig. S7; see also Smesdrud et al., 2017), indicating that atmospheric wind variability is well represented over the region. However, apart from the polynya region, sea ice coverage is overestimated especially north of Svalvard (Fig. 1b) where basal melting plays a role (Fig. 5c). This may suggest that ocean heat delivered via the West Spitsbergen Current is not strong enough to melt sea ice along its path into the Arctic. It can also be speculated that this region might have been affected by increasing extreme winter storm activities in recent years that are associated with anomalous warming events. However, no significant trends were found in January-February during 1979-2015 (Rinke et al., 2017).

- Deleted: to
- Deleted: the
- Deleted: in the
- Deleted: It

Considering the overall vulnerability and fate of some of the thickest sea ice in the Arctic under the recent warming climate, sea ice may become susceptible to modulation by the atmospheric forcing. Sensitivity experiments indicate that a polynya would occur under the same atmospheric conditions of February 2018 even in the thick ice conditions like the winter of 1979 (Moore et al., 2018). By taking advantage of an ensemble approach, the internal variability is better assessed with respect to the occurrence of such coastal polynyas during extreme events. The RASM-DPLE study shows that the frequency, size and duration of winter polynyas is unaffected by sea ice thinning, showing no apparent difference in polynya size or frequency of occurrence over time between the two ensembles initialized with two different sea ice regimes 30 years apart: 1980s vs 2010s. One interesting feature is that winter polynyas mostly (16 out of 25) occur in December-January in the 1985-initialized simulations whereas they are more prevalent (12 out of 16) in January-February in the 2015-initialized ensemble runs. In addition, the range of 5-day mean SIT before the occurrence of each polynya was 2.9-4.4 m (with a mean of 3.7 m and standard deviation of 0.36 m) in the 1985-initialized RASM-DPLE ensemble and 1.8-3.4 m (with a mean of 2.8 m and standard deviation of 0.36 m) in the 2015-initialized RASM-DPLE ensemble (Table 3 and 4, respectively). Although sea ice is thicker, the 1985-initialized ensemble simulations produce 56% more polynyas than the 2015-initialized ones. Hence, regardless of the SIT decline over the region, this study suggests that the primary necessary condition for a winter polynya occurrence is strong and persistent southerly winds. With the maximum wind speed exceeding 20 m/s and duration for 10 days, the 2018 winter polynya remains unique by any of these metrics.

- Deleted: The s
- Deleted: s
- Deleted: r
- Deleted: in
- Deleted: However, t
- Deleted: simulations indicate
- Deleted: not
- Deleted: such
- Deleted: characteristics
- Deleted: s
- Deleted: and increase in its
- Deleted: ies
- Deleted: 0
- Deleted: For example
- Deleted: the
- Deleted: 8
- Deleted: 40
- Deleted: 3
- Deleted: the
- Deleted: 35
- Deleted: s
- Deleted: occurring
- Deleted: reversal isthe observed polynya
- Deleted: D
- Deleted: sudden stratospheric warming (
- Deleted:)

When the polynyas occurred over the region, they coincided with a reversal (from the positive to negative) of the daily AO index. For instance, during the 2017-2018 winter, the AO shifted significantly from a positive phase to a strong negative phase between mid-February to early March (Fig. 3a), which was associated with weakening of the polar vortex and allowed warmer air into the Arctic. Alternatively, Kim et al. (2014) argued that due to sea ice loss especially in the Barents-Kara seas, the weakened stratospheric polar vortex preferentially induced a negative phase of the AO at the surface, resulting in warm air moving into the Arctic. This anomalous warming event occurred coincidentally after the SSW event observed in mid-February 2018 (Moore et al., 2018; Rao et al. 2018), which developed into a vortex split (Lü et al., 2020). Subsequently, the winter weather was severe with intense cold air across Europe in March 2018 (Overland et al., 2020). Although the exact cause of SSW variability is still under debate, SSWs are generally known to cause anomalous warming over Greenland and impact surface weather patterns down to mid-latitudes (Butler et al., 2017). The frequency of their occurrence

620 is enhanced by El Niño conditions (Polvani et al., 2017) and may also be enhanced during La Niña
winters depending on SSW definitions (Song and Son, 2018). For example, La Niña was in winter
2017-2018, while El Niño was in December 1986. In the recent winters with SSW (Butler et al., 2017;
625 Rao et al., 2018; Knight et al., 2021), no winter polynya events occurred except in February 2018. In
other words, the winter polynyas in 1986, 2011, and 2017 were not associated with SSW.

Deleted: true for
Deleted: ;
Deleted:
Deleted: f

630 The opening of polynyas in the region between the Wandel and Lincoln seas primarily depends on a
large-scale surface pressure pattern change, resulting in strengthening of intense southerly winds that
are short-lived and sporadic. Toward the end of each polynya event, the rate of thermodynamic ice
growth generally increased (Figs. 4d, 4e and 4f). The maximum rate of ice growth happened not when
635 sea ice removal was largest with strong southerly wind but when air temperature started to significantly
drop (Fig. 4). Upward surface turbulent heat fluxes continued even after DVT became positive (i.e.,
Table 2 and Fig. 4) when wind direction was reversed (i.e., Fig. 6d) until the open water area was
completely covered by ice. In general, the rate of new ice formation in winter is expected to be high
during an open water phase, such as leads and polynyas, due to an intense turbulent heat (sensible and
640 latent) loss to the atmosphere. In some polynya regions, turbulent heat loss is as high as 300 W/m² such
as along the Weddell Sea coast (see Morales Maqueda et al., 2004). However, the mean daily turbulent
heat loss in the study region was about 61 W/m² in 2018 (with the daily maximum of 124 W/m²), which
is similar to the model estimates; the average/maximum daily heat fluxes from the ocean to the
atmosphere are 40 and 124 W/m², respectively (Ludwig et al., 2019). Those values are much smaller
645 than that in the St. Lawrence Island polynya (412 W/m²; Pease, 1987) and Okhotsk Sea coast (471
W/m²; Alfultis and Martin, 1987), but comparable to the Northeast Water (NEW) polynya (31 W/m²;
Morales Maqueda et al., 2004). Nevertheless, new sea ice formation was somewhat slowed down in the
study region because the air-sea temperature difference was much smaller than in the other regions
listed above due to the anomalously warm air carried by southerly winds over northern Greenland.

Deleted: 48
Deleted: a
Deleted: is

650 RASM estimated a maximum polynya size of 13,000 km², where SIT is less than 10 cm, on that day
while Ludwig et al. (2019) calculated it as large as approximately 60,000 km² using satellite SIC
although there are large uncertainties between different satellite algorithms. For example, MODIS SIC
underestimates open water areas by about 50%, compared to the Advanced Microwave Scanning
Radiometer 2 (AMSR2) SIC (see Fig. 3 in Ludwig et al. (2019)). On the other hand, the RASM polynya
size is based on SIT and its maximum size could increase to 29,400 km² if a threshold of SIT less than
25 cm is applied. According to the RASM simulation between February 1st to March 31st, 2018, the
study region lost approximately 208 km³ of sea ice due to mechanical ice removal. Although sea ice was
655 added later by the dynamic replenishment (102 km³) and the thermodynamic ice growth (64 km³), the
deficit of SIV was about 42 km³, which is equivalent to the 36 cm reduction of mean SIT over the study
region. However, von Albedyll et al. (2021) showed that that the thermodynamic and dynamic
processes were almost equally contributed to the mean SIT change between February 25th and March
31st, 2018. Ludwig et al. (2019) calculated the ice volume thermodynamically produced (33 km³)
including the polynya period (February 14th to March 31st, 2018) using the freezing degree day
parameterization, but RASM produced slightly more sea ice for the same period (53 km³). Although an
660 ice cover was re-established with a mean thickness of 1.96 m at the end of March 2018 (von Albedyll et

Deleted: the

Deleted: Finally, unlike the events in 2011 and 2017, sea ice in the northern Greenland region was not fully restored even a month after the peak ice removal on February 25th, 2018 (Fig. 4d).

Deleted: The

Deleted: the

Deleted: Based on

Deleted:

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Deleted: The

Deleted: higher

Deleted: above

Formatted: Superscript

680 al., 2021), sea ice in the northern Greenland region was not fully restored even a month after the peak
ice removal on February 25th, 2018 unlike the events in 2011 and 2017 (Fig. 4d). The CryoSat-2/SMOS
data also confirmed the similar sea ice loss in the region; at the end of March 2018, the mean sea ice
north of Greenland was relatively thinner (2.45 m), compared to the beginning of February (2.73 m).
685 Therefore, one could hypothesize that the 2018 winter polynya event could have contributed to the
preconditioning of the polynya event in the following summer (Schweiger et al., 2021), which was
observed at a similar location. Hence the winter event might have been unprecedented by yet another
measure, as polynya events have never repeated within a year over the study region except during 2018.

Deleted: Henc

6 Summary

690 Following the previous studies of the 2018 winter polynya by Moore et al. (2018) and Ludwig et al.
(2019), this study demonstrates that an additional three observed winter open water events occurred
under a specific wind pattern, which removes sea ice mechanically out of the region, i.e., December
1986 as well as February 2011 and 2017. In all of the events, the wind direction was primarily from the
south and southwest (Figs. 4i and 4h). The size of polynya depended on the strength and persistence of
southerly winds, and the polynya closure was a result of the relaxation of these conditions. Although the
695 atmospheric conditions were associated with SSW in February 2018, other winter polynyas occurred in
non-SSW winters. Rather, they coincided with a reversal of the AO from a positive to a negative phase.
The polynya was larger in February 2011 compared to the one in 2017 because wind duration was
longer: 4 days of strong southerly wind (> 10 m/s) in 2011 compared to 2 days in 2017 (Fig. 8a). Hence,
31% more sea ice was removed in the former, although wind speed and wind stress were similar
700 between the two years (Table 1). The observed polynya in February 2018 was the largest one over the
satellite SIC record since 1979. It resulted from much stronger and more persistent southerly-
southeasterly winds (Fig. 4g) off the coast of northern Greenland, the direction which could also be
more favorable for polynya opening (Kawaguchi et al., 2010). Although southerly wind in December
1986 was as strong as in February 2018, its duration was shorter and thus not only the size of the
705 polynya, but also the turbulent heat flux, was relatively smaller. Sea ice was significantly thicker in the
1980s in north of Greenland, but more polynyas were not prevalent with thinner ice in the winters of
2015-2025 from the two RASM-DPLE ensembles. Given the more frequent rate of winter polynya
occurrence and their apparent lack of sensitivity to the initial sea ice thickness, we conclude that a
dominant cause of these winter polynyas stems from internal variability of atmospheric forcing rather
710 than from the forced response to a warming climate.

Deleted: study

Deleted: all

Deleted: February 2011 and 2017

Deleted: as

Deleted:

Deleted: and

Deleted: Although

Deleted: s

Deleted: the

Deleted: ern

Deleted: similar

Acknowledgement

This work was funded by NSF OPP IAA1603602 (NPS), DOE RGMA DE-SC0014117 (NPS), NSF
OPP-1603544 (CU Boulder), the Canada 150 Chair program (J. S.), and the Ministry of Science and
Higher Education in Poland (R. O.) in the frame of international project agreement 3808/FAO/2017/0.
715 The PNNL is operated for the U.S. DOE by Battelle Memorial Institute under contract DE-AC05-
76RL01830. In addition, computing resources were provided by the U.S. DOD High Performance

730 Computer Modernization Program (HPCMP). [The merging of CryoSat-2 und SMOS data was funded by the ESA project SMOS & CryoSat-2 Sea Ice Data Product Processing and Dissemination Service and data from February 2011 to 2018 were obtained from <https://www.meereisportal.de> \(grant: REKLIM-2013-04\). We thank the reviewers for providing helpful comments, which improved an earlier version of this publication.](#)

Competing interestsCode/data availability

735 All RASM simulations are archived in the HPCMP archive system and will be available upon publication according to the U.S. DOD data policy.

Competing interests

740 The authors declare no competing interests.

Author contributions

745 Y. L. and W. M. conceived the study and wrote the paper. A. C., S. K., R. O., and M. S. undertook the implementation and development of the RASM simulation. J. C., J. C. K., J. S., and H. W. contributed to the interpretation of results. All authors except S. K. and A. C. commented on the manuscript.

References

- Alfultis, M. A., & Martin, S. (1987). Satellite passive microwave studies of the Sea of Okhotsk ice cover and its relation to oceanic processes, 1978–1982. *Journal of Geophysical Research: Oceans*, 92(C12), 13013-13028.
- 750 Bintanja, R., & Van der Linden, E. C. (2013). The changing seasonal climate in the Arctic. *Scientific reports*, 3(1), 1-8.
- Butler, A. H., Sjöberg, J. P., Seidel, D. J., & Rosenlof, K. H. (2017). A sudden stratospheric warming compendium. *Earth Syst. Sci. Data*, 9, 63–76. <https://doi.org/10.5194/essd-9-63-2017>
- Cassano, J. J., DuVivier, A., Roberts, A., Hughes, M., Seefeldt, M., Brunke, M., et al. (2017). Development of the Regional Arctic System Model (RASIM): near-surface atmospheric climate sensitivity. *Journal of Climate*, 30(15), 5729-5753.
- 755 Cavaleri, D. J., Gloersen, P., & Campbell, W. J. (1984). Determination of sea ice parameters with the Nimbus 7 SMMR. *Journal of Geophysical Research: Atmospheres*, 89(D4), 5355-5369.
- Chassignet, E. P., Yeager, S. G., Fox-Kemper, B., Bozec, A., Castruccio, F., Danabasoglu, G., ... & Xu, X. (2020). Impact of horizontal resolution on global ocean–sea ice model simulations based on the experimental protocols of the Ocean Model Intercomparison Project phase 2 (OMIP-2). *Geoscientific Model Development*, 13(9), 4595-4637.
- 760 Craig, A. P., Vertenstein, M., & Jacob, R. (2012). A new flexible coupler for earth system modeling developed for CCSM4 and CESM1. *The International Journal of High Performance Computing Applications*, 26(1), 31-42.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., et al. (2011). The ERA-Interim reanalysis: Configuration and performance of the data assimilation system. *Quarterly Journal of the royal meteorological society*, 137(656), 553-597.
- 765 DuVivier, A. K., Cassano, J. J., Craig, A., Hamman, J., Maslowski, W., Nijssen, B., et al. (2016). Winter atmospheric buoyancy forcing and oceanic response during strong wind events around southeastern Greenland in the Regional Arctic System Model (RASIM) for 1990–2010. *Journal of Climate*, 29(3), 975-994.
- Graham, R. M., Cohen, L., Petty, A. A., Boisvert, L. N., Rinke, A., Hudson, S. R., et al. (2017). Increasing frequency and duration of Arctic winter warming events. *Geophysical Research Letters*, 44(13), 6974–6983. <https://doi.org/10.1002/2017GL073395>.
- 770 Hamman, J., Nijssen, B., Brunke, M., Cassano, J., Craig, A., DuVivier, A., et al. (2016). Land surface climate in the regional Arctic system model. *Journal of Climate*, 29(18), 6543-6562.
- Hamman, J., Nijssen, B., Roberts, A., Craig, A., Maslowski, W., & Osinski, R. (2017). The coastal streamflow flux in the Regional Arctic System Model. *Journal of Geophysical Research: Oceans*, 122(3), 1683-1701.
- 775 Itkin, P., Spreen, G., Cheng, B., Doble, M., Girard-Arduin, F., Haapala, J., et al. (2017). Thin ice and storms: Sea ice deformation from buoy arrays deployed during N-ICE2015. *Journal of Geophysical Research: Oceans*, 122(6), 4661-4674.
- Kawaguchi, Y., Nihashi, S., Mitsudera, H., & Ohshima, K. I. (2010). Formation mechanism of huge coastal polynyas and its application to Okhotsk northwestern polynya. *Journal of physical oceanography*, 40(11), 2451-2465.
- 780 Kim, B. M., Son, S. W., Min, S. K., Jeong, J. H., Kim, S. J., Zhang, X., ... & Yoon, J. H. (2014). Weakening of the stratospheric polar vortex by Arctic sea-ice loss. *Nature communications*, 5(1), 1-8.
- Knight, J., Scaife, A., Bett, P. E., Collier, T., Dunstone, N., Gordon, M., ... & Walker, B. (2021). Predictability of European Winters 2017/2018 and 2018/2019: Contrasting influences from the Tropics and stratosphere. *Atmospheric Science Letters*, 22(1), e1009.
- 785 Kohonen, T. (2001). *Self-Organizing Maps*, third ed. Springer-Verlag, New York.
- Kwok, R. (2018). Arctic sea ice thickness, volume, and multiyear ice coverage: losses and coupled variability (1958–2018). *Environmental Research Letters*, 13(10), 105005. <https://doi.org/10.1088/1748-9326/aae3ec>.
- 790 Kwok, R., Cunningham, G. F., Wensnahan, M., Rigor, I., Zwally, H. J., & Yi, D. (2009). Thinning and volume loss of the Arctic Ocean sea ice cover: 2003–2008. *Journal of Geophysical Research*, 114(C7), C07005. <https://doi.org/10.1029/2009JC005312>.
- Large, W., & Yeager, S. G. (2009). The global climatology of an interannually varying air–sea flux data set. *Climate dynamics*, 33(2-3), 341-364.

Deleted: Grosfeld, K.; Treffeisen, R.; Asseng, J.; Bartsch, A.; Bräuer, B.; Fritzsche, B.; Gerdes, R.; Hendricks, S.; Hiller, W.; Heygster, G.; Krumpen, T.; Lemke, P.; Melsheimer, C.; Nicolaus, M.; Ricker, R. and Weigelt, M. (2016), Online sea-ice knowledge and data platform <www.meereisportal.de>. *Polarforschung*, Bremerhaven, Alfred Wegener Institute for Polar and Marine Research & German Society of Polar Research, 85 (2), 143-155, doi:10.2312/polfor.2016.011. -

- Liu, Y., & Weisberg, R. H. (2011). A review of Self-Organizing Map applications in meteorology and oceanography. In *Self-Organizing Maps - Applications and Novel Algorithm Design*, Edited by J. I. Mwasiagi, InTech, Rijeka, Croatia, ISBN 978-953-307-546-4, pp.253-272.
- 805 Lü, Z., Li, F., Orsolini, Y. J., Gao, Y., & He, S. (2020). Understanding of European cold extremes, sudden stratospheric warming, and Siberian snow accumulation in the winter of 2017/18. *Journal of Climate*, 33(2), 527-545.
- Ludwig, V., Spreen, G., Haas, C., Istomina, L., Kauker, F., & Murashkin, D. (2019). The 2018 North Greenland polynya observed by a newly introduced merged optical and passive microwave sea-ice concentration dataset. *Cryosphere*, 13(7).
- 810 Maslowski, W., Kinney, J. C., Higgins, M., & Roberts, A. (2012). The future of Arctic sea ice. *Annual Review of Earth and Planetary Sciences*, 40.
- Maykut, G. A. (1982). Large-scale heat exchange and ice production in the central Arctic. *Journal of Geophysical Research*, 87(C10), 7971. <https://doi.org/10.1029/JC087iC10p07971>.
- 815 ~~Meier, W. N., Fetterer, F., Windnagel, A. K., & Stewart, J. S. (2021). NOAA/NSIDC Climate Data Record of Passive Microwave Sea Ice Concentration, Version 4. Boulder, Colorado USA. NSIDC: National Snow and Ice Data Center. doi: <https://doi.org/10.7265/efmz-2t65>.~~
- Meier, W. N., Hovelsrud, G. K., van Oort, B. E. H., Key, J. R., Kovacs, K. M., Michel, C., et al. (2014). Arctic sea ice in transformation: A review of recent observed changes and impacts on biology and human activity. *Reviews of Geophysics*, 52(3), 185–217. <https://doi.org/10.1002/2013RG000431>.
- 820 Moore, G. W. K., Schweiger, A., Zhang, J., & Steele, M. (2018). What Caused the Remarkable February 2018 North Greenland Polynya? *Geophysical Research Letters*. <https://doi.org/10.1029/2018GL080902>.
- Morales Maqueda, M. A., Willmott, A. J., & Biggs, N. R. T. (2004). Polynya Dynamics: a Review of Observations and Modeling. *Reviews of Geophysics*, 42(1), RG1004. <https://doi.org/10.1029/2002RG000116>.
- 825 Overland, J., Hall, R., Hanna, E., Karpechko, A., Vihma, T., Wang, M., & Zhang, X. (2020). The polar vortex and extreme weather: the Beast from the East in winter 2018. *Atmosphere*, 11(6), 664.
- Pease, C. H. (1987). The size of wind-driven coastal polynyas. *Journal of Geophysical Research: Oceans*, 92(C7), 7049-7059.
- Peings, Y., Labe, Z. M., & Magnusdóttir, G. (2021). Are 100 ensemble members enough to capture the remote atmospheric response to +2° C Arctic sea ice loss? *Journal of Climate*, 34(10), 3751-3769.
- 830 Polvani, L. M., Sun, L., Butler, A. H., Richter, J. H., & Deser, C. (2017). Distinguishing stratospheric sudden warmings from ENSO as key drivers of wintertime climate variability over the North Atlantic and Eurasia. *Journal of Climate*, 30(6), 1959-1969.
- Rampal, P., Weiss, J., Marsan, D., & Bourgoin, M. (2009). Arctic sea ice velocity field: General circulation and turbulent-like fluctuations. *Journal of Geophysical Research*, 114(C10), C10014. <https://doi.org/10.1029/2008JC005227>.
- 835 Rao, J., Ren, R., Chen, H., Yu, Y., & Zhou, Y. (2018). The stratospheric sudden warming event in February 2018 and its prediction by a climate system model. *Journal of Geophysical Research: Atmospheres*, 123(23), 13-332.
- ~~Ricker, R., Kauker, F., Schweiger, A., Hendricks, S., Zhang, J., & Paul, S. (2021). Evidence for an increasing role of ocean heat in Arctic winter sea ice growth. *Journal of Climate*, 34(13), 5215-5227.~~
- 840 ~~Ricker, R., Hendricks, S., Kaleschke, L., Tian-Kunze, X., King, J., & Haas, C. (2017). A weekly Arctic sea-ice thickness data record from merged CryoSat-2 and SMOS satellite data. *The Cryosphere*, 11(4), 1607-1623.~~
- Rinke, A., Maturilli, M., Graham, R. M., Matthes, H., Handorf, D., Cohen, L., et al. (2017). Extreme cyclone events in the Arctic: Wintertime variability and trends. *Environmental Research Letters*, 12(9), 094006.
- Roberts, A., Craig, A., Maslowski, W., Osinski, R., DuVivier, A., Hughes, M., et al. (2015). Simulating transient ice-ocean Ekman transport in the Regional Arctic System Model and Community Earth System Model. *Annals of glaciology*, 56(69), 211-228.
- 845 Saha, S., Moorthi, S., Wu, X., Wang, J., Nadiga, S., Tripp, P., et al. (2011). NCEP Climate Forecast System Version 2 (CFSv2) 6-hourly Products. Research Data Archive at the National Center for Atmospheric Research, Computational and Information Systems Laboratory. <https://doi.org/10.5065/D61CITXF>.
- 850 Schweiger, A. J., Steele, M., Zhang, J., Moore, G. W. K., & Laidre, K. L. (2021). Accelerated sea ice loss in the Wandel Sea points to a change in the Arctic's Last Ice Area. *Communications Earth & Environment*, 2(1), 1-11.

Deleted: Meier, W. N., Fetterer, F., Savoie, M., Mallory, S., Duerr, R., & Stroeve, J. (2017a). *NOAA/NSIDC Climate Data Record of Passive Microwave Sea Ice Concentration, Version 3*. Boulder, Colorado USA. NSIDC: National Snow and Ice Data Center. doi: <https://doi.org/10.7265/NS9P2ZTG>. Accessed 27 Sep 2019. - ... [1]

Deleted: Accessed 01 Jul 2020.

- 860 Schweiger, A., Lindsay, R., Zhang, J., Steele, M., Stern, H., & Kwok, R. (2011). Uncertainty in modeled Arctic sea ice volume. *Journal of Geophysical Research: Oceans*, 116(C8).
- Serreze, M. C., & Francis, J. A. (2006). The Arctic amplification debate. *Climatic change*, 76(3-4), 241-264.
- Song, K., & Son, S. W. (2018). Revisiting the ENSO–SSW relationship. *Journal of Climate*, 31(6), 2133-2143.
- Spreen, G., Kwok, R., & Menemenlis, D. (2011). Trends in Arctic sea ice drift and role of wind forcing: 1992-2009. *Geophysical Research Letters*, 38(19). <https://doi.org/10.1029/2011GL048970>.
- 865 Steele, M., Morley, R., & Ermold, W. (2001). PHC: A global ocean hydrography with a high-quality Arctic Ocean. *Journal of Climate*, 14(9), 2079-2087.
- Stroeve, J., & Notz, D. (2018). Changing state of Arctic sea ice across all seasons. *Environmental Research Letters*, 13(10), 103001.
- 870 Stroeve, J., Barrett, A., Serreze, M., & Schweiger, A. (2014). Using records from submarine, aircraft and satellites to evaluate climate model simulations of Arctic sea ice thickness. *The Cryosphere*, 8(5), 1839-1854.
- [von Albedyll, L., Haas, C., & Dierking, W. \(2021\). Linking sea ice deformation to ice thickness redistribution using high-resolution satellite and airborne observations. *The Cryosphere*, 15\(5\), 2167-2186.](#)
- Wang, Q., Ricker, R., & Mu, L. (2021). Arctic sea ice decline preconditions events of anomalously low sea ice volume export through Fram Strait in the early 21st century. *Journal of Geophysical Research: Oceans*, 126(2), e2020JC016607.
- 875 [Watts, M., Maslowski, W., Lee, Y. J., Kinney, J. C., & Osinski, R. \(2021\). A spatial evaluation of Arctic sea ice and regional limitations in CMIP6 historical simulations. *Journal of Climate*, 34\(15\), 6399-6420.](#)
- Weijer, W., Veneziani, M., Stössel, A., Hecht, M. W., Jeffery, N., Jonko, A., et al. (2017). Local Atmospheric Response to an Open-Ocean Polynya in a High-Resolution Climate Model. *Journal of Climate*, 30(5), 1629–1641.
- 880 <https://doi.org/10.1175/JCLI-D-16-0120.1>.
- Yeager, S.G., Danabasoglu, G., Rosenbloom, N.A., Strand, W., Bates, S.C., Meehl, G.A., Karspeck, A.R., Lindsay, K., Long, M.C., Teng, H. and Lovenduski, N.S., 2018. Predicting near-term changes in the Earth System: A large ensemble of initialized decadal prediction simulations using the Community Earth System Model. *Bulletin of the American Meteorological Society*, 99(9), pp.1867-1886.
- 885 Zhang, J., & Rothrock, D. A. (2003). Modeling global sea ice with a thickness and enthalpy distribution model in generalized curvilinear coordinates. *Monthly Weather Review*, 131(5), 845-861.

Formatted: Font:Italic

895 Table 1. Characteristics of the polynya events, when daily dynamic sea ice removal is greater than 10 km³/day (i.e., dynamic volume tendency (DVT) ≤ -10 km³/day) for more than three consecutive days, and the corresponding wind condition near a weather station, as well as sea ice thickness (SIT) over northern Greenland from the RASM hindcast simulation (1980-2020).

Year	Date when DVT ≤ -10 km ³ /day (No. of polynya days)	Five-day mean SIT (m) before polynya	Daily Mean DVT (km ³ /day)	Total sea ice removal (km ³)	North-South wind during polynya			Nearest weather station
					Time-integrated Pseudo-wind stress* (m ² /s ² ·day)	Mean speed (m/s)	Max. speed (m/s)	
1986	<u>12-16 Dec (5 days)</u>	<u>4.4</u>	<u>-28.4</u>	<u>-142</u>	<u>717</u>	<u>11.0</u>	<u>17.6</u>	<u>Cape Morris Jessup</u>
2011	12-15 Feb (4 days)	3.6	-13.8	-55	476	10.2	15.1	Station Nord
2017	8-10 Feb (3 days)	3.3	-14.0	-42	354	9.5	15.3	Station Nord
2018	16-25 Feb (10 days)	3.2	-18.9	-189	486	11.3	21.8	Cape Morris Jessup

*only for the northward wind component

Deleted: 1272

Deleted: 1

905 Table 2. Daily mean net turbulent (sensible and latent) heat flux (W/m^2) and its standard deviation (s.d.) during the polynya events over open water areas (km^2), where sea ice thickness (SIT) is less than 10 cm, from the RASM hindcast and DPLE simulations. The negative values indicate heat loss from the polynya to the atmosphere.

RASM Cases	Year	Date	Daily Turbulent Flux (W/m^2)			Daily Open Water Area (km^2)			Total integrated turbulent heat (W)	
			max	mean	s.d.	max	mean	s.d.		
Hindcast	1986	14-18 Dec	-48.8	-18.9	18.5	7,811	4,120	2,822	-3.56×10^{11}	
	2011	15-18 Feb	-116	-91.6	21.7	601	322	236	-1.25×10^{11}	
	2017	10 Feb	-1.59	-1.59	n/a	172	172	n/a	-2.74×10^8	
	2018	20 Feb-1 Mar	-124	-60.9	29.4	13,000	4,210	4,390	-2.03×10^{12}	
DPLE	1985-initialized ensemble #4	1988	17-21 Jan	-194	-103	29.8	429	309	130	-1.54×10^{11}
	2015-initialized ensemble #2	2024	20-24 Jan	-110	-74.1	23.8	6,090	2,540	2,340	-8.87×10^{11}

- Deleted: 125
- Deleted: 96
- Deleted: 7
- Deleted: 23
- Deleted: 2
- Formatted Table
- Deleted: 80
- Deleted: 0.289
- Deleted: 182
- Deleted: 48.2
- Deleted: 49.2
- Deleted: 196
- Deleted: 100
- Deleted: 67.3
- Deleted: 202
- Deleted: 62.6
- Deleted: 43.4

930 Table 3. The polynya events, defined when daily dynamic sea ice removal is greater than 10 km³/day (i.e., dynamic volume tendency (DVT) ≤ -10 km³/day) for more than three consecutive days, and the corresponding wind condition near Cape Morris Jessup as well as sea ice thickness (SIT) over northern Greenland from the RASM-DPLE simulation, initialized on December 1st, 1985. A correlation coefficient (*r*) is calculated between daily DVT and North-South wind for December-March and the bold indicates the largest polynya event.

DPLE ensemble member	Year	Month	No. of days when DVT ≤ -10 km ³ /day	Five-day mean SIT (m) before polynya	Daily mean DVT (km ³ /day)	Total Ice removal (km ³)	North-South wind during polynya			<i>r</i> (<i>p</i> <0.01) between daily DVT and North-South wind <u>Dec-Mar</u>	
							Time-integrated pseudo-wind stress* (m ² /s ² ·day)	Mean speed (m)	Max. speed (m)		
1	1986	Jan	3	4.1	-16.1	-48	307	9.7	13.6	-0.73	Deleted: January
1	1986	Mar	4	4.4	-12.6	-50	31	2.5	4.3	-0.73	Deleted: 75
1	1988	Mar	3	4.0	-11.5	-35	168	7.4	9.9	-0.76	Deleted: 75
1	1990	Mar	4	3.5	-15.6	-62	395	8.6	15.5	-0.72	Deleted: 74
<u>1</u>	<u>1991</u>	<u>Dec</u>	<u>3</u>	<u>3.8</u>	<u>-17.6</u>	<u>-53</u>	<u>244</u>	<u>8.6</u>	<u>12.2</u>	<u>-0.66</u>	Deleted: Jan
<u>2</u>	<u>1991</u>	<u>Dec</u>	<u>6</u>	<u>3.8</u>	<u>-22.3</u>	<u>-134</u>	<u>761</u>	<u>10.6</u>	<u>14.7</u>	<u>-0.81</u>	Deleted: 75
3	1987	Jan	4	3.8	-21.4	-85	475	10.1	19.0	-0.69	Deleted: 75
<u>4</u>	<u>1986</u>	<u>Dec</u>	<u>6</u>	<u>3.7</u>	<u>-16.9</u>	<u>-101</u>	<u>439</u>	<u>7.6</u>	<u>15.2</u>	<u>-0.78</u>	Deleted: 75
4	1988	Jan	7	4.2	-17.2	-120	438	7.6	12.0	-0.64	Deleted: 74
4	1989	Feb	5	4.0	-20.7	-104	651	11.2	15.7	-0.70	Deleted: 5
4	1991	Feb	3	3.2	-25.2	-76	550	12.6	17.8	-0.81	Deleted: 4
<u>4</u>	<u>1991</u>	<u>Dec</u>	<u>3</u>	<u>3.6</u>	<u>-13.5</u>	<u>-40</u>	<u>230</u>	<u>8.3</u>	<u>12.9</u>	<u>-0.78</u>	Deleted: 0
5	1986	Jan	4	3.8	-17.0	-68	260	7.3	13.7	-0.70	Deleted: 71
5	1987	Jan	4	3.5	-16.5	-66	121	4.9	8.5	-0.68	Deleted: 68
<u>5</u>	<u>1987</u>	<u>Dec</u>	<u>4</u>	<u>3.4</u>	<u>-13.9</u>	<u>-55</u>	<u>438</u>	<u>10.1</u>	<u>14.1</u>	<u>-0.61</u>	Deleted: 67
5	1988	Mar	3	3.8	-20.0	-60	121	4.8	11.2	-0.61	Deleted: 50
<u>6</u>	<u>1986</u>	<u>Dec</u>	<u>3</u>	<u>3.4</u>	<u>-12.6</u>	<u>-38</u>	<u>111</u>	<u>5.6</u>	<u>10.0</u>	<u>-0.74</u>	Deleted: 74
6	1994	Feb	4	2.9	-22.0	-88	700	13.0	16.7	-0.73	Deleted: 74
<u>7</u>	<u>1990</u>	<u>Dec</u>	<u>3</u>	<u>3.3</u>	<u>-20.8</u>	<u>-62</u>	<u>283</u>	<u>9.4</u>	<u>12.6</u>	<u>-0.73</u>	Deleted: 74
7	1994	Jan	3	3.8	-14.7	-44	255	8.5	12.3	-0.65	Deleted: 1
8	1992	Jan	3	4.3	-13.6	-41	277	9.2	15.0	-0.71	Deleted: 1
8	1995	Jan	3	3.3	-16.9	-51	172	6.9	13.0	-0.63	Deleted: 3
<u>9</u>	<u>1989</u>	<u>Dec</u>	<u>4</u>	<u>4.0</u>	<u>-13.2</u>	<u>-53</u>	<u>228</u>	<u>7.1</u>	<u>10.9</u>	<u>-0.71</u>	Deleted: 6
9	1995	Mar	3	3.8	-16.3	-49	322	10.2	11.6	-0.69	Deleted: 2
10	1987	Feb	3	3.6	-13.2	-40	145	6.6	10.1	-0.64	Deleted: 59

* only for the northward wind component

935

Table 4. Same as Table 3, but for the RASM–DPLE simulation, initialized on December 1st, 2015.

DPLE ensemble member	Year	Month	No. of days when DVT ≤ -10 km ³ /day	Five-day mean SIT (m) before polynya	Daily mean DVT (km ³ /day)	Total Ice removal (km ³)	North-South wind during polynya			r ($p < 0.01$) between daily DVT and North-South wind (Dec-Mar)	
							Time-integrated pseudo-wind stress* (m ² /s ² ·day)	Mean speed (m)	Max. speed (m)		
1	2016	Feb	5	2.7	-14.4	-72	165	5.0	10.1	-0.75	Deleted: Jan
1	2022	Feb	3	2.6	-15.1	-45	216	8.2	11.5	-0.62	Deleted: 63
1	2023	Jan	3	2.5	-14.4	-43	321	10.0	14.3	-0.78	Deleted: 82
1	2023	Jan	3	1.8	-15.4	-46	299	9.8	13.1	-0.78	Deleted: 82
2	2016	Jan	3	2.8	-16.4	-49	192	7.8	10.9	-0.76	Deleted: 77
2	2024	Jan	7	3.3	-19.4	-136	841	9.8	19.2	-0.78	Deleted: 79
4	2019	Mar	3	3.0	-15.4	-46	235	7.4	16.9	-0.63	Deleted: 65
5	2018	Jan	5	3.0	-20.8	-104	740	10.8	17.9	-0.77	Deleted: 8
5	2018	Mar	3	3.0	-15.4	-46	129	5.6	12.9	-0.77	Deleted: 8
5	2020	Feb	5	3.0	-15.5	-77	379	8.4	12.2	-0.75	
6	2016	Jan	4	2.8	-12.1	-48	379	9.6	12.3	-0.77	Deleted: 78
6	2018	Dec	3	3.0	-15.1	-45	358	10.4	16.2	-0.71	
6	2024	Jan	4	2.8	-20.0	-80	352	8.1	15.2	-0.72	
7	2023	Dec	4	3.4	-16.4	-66	305	8.2	14.8	-0.70	
8	2017	Feb	3	2.9	-14.5	-43	558	13.1	17.0	-0.79	Deleted: 8
8	2025	Jan	5	2.8	-18.5	-92	531	9.9	15.8	-0.76	Deleted: 74

960 *only for the northward wind component







Figure 1: (a) The Visible Infrared Imaging Radiometer Suite (VIIRS) Nighttime Imagery (Day/Night Band, Enhanced Near Constant Contrast) over the northern Greenland on 25 February 2018 (adapted from NASA Worldview at <https://worldview.earthdata.nasa.gov>) and sea ice thickness from (b) the RASM hindcast simulation as well as (c) CFS version 2 reanalysis over the Arctic on 25 February 2018. The rectangular box indicates the polynya area.

Deleted:
Deleted: 24

975

980



Deleted:

985 Figure 2: Mean sea ice thickness (SIT) over the northern Greenland region from the CryoSat-2/SMOS merged product on (a) 22-28 February 2018, (b) 8-14 February 2017, and (c) 13-19 February 2011 as well as daily SIT from the RASM hindcast simulation on (d) 25 February 2018, (e) 11 February 2017, and (f) 16 February 2016.





Figure 3. (a) The observed daily AO index and (b)-(h) the daily mean near-surface air temperature (T , °C; **blue**) during January-March 2018 from the selected weather stations around Greenland as shown in (i) and (j) as circle markers. The RASM daily 2-m air temperature (**light blue**) is from the nearest grid cell and the observed long-term daily mean (LTM, 2011-2020; **blue-dashed**) is only at Station 04301. The correlation coefficient (r) is statistically significant ($p < 0.05$) between the observed and the simulated air temperature is shown in the upper-left. The vertical **dashed** lines indicate the start and end dates of the 2018 winter polynya period. The spatial maps show (i) the maximum correlation coefficients and (j) their lagged time-scales (days) between the AO index and the RASM air temperature. Note that grid cells not statically significant ($p > 0.05$) are indicated as white. The color-scale of the markers quantifies the relationship between the AO index and the observed air temperature at each weather station.

Deleted:
 Deleted: black
 Deleted: black
 Deleted: (red-dashed)





Figure 4. The observed daily near-surface air temperature (T , °C; blue) from Station 04301 (Cape Morris Jesup as ●) in (a) and the RASM daily 2-m air temperature (light blue) from the nearest grid cell during January-March of (b) 2017 and (c) 2011; the observed long-term daily mean (LTM, 2011-2020; blue-dashed). The correlation coefficient (r) between the observed and the simulated air temperature is shown in the upper-left. Time rate of change of the RASM daily sea ice volume (km^3/day) due to thermodynamic (red) and dynamic (blue) tendency during January-March of (d) 2018, (e) 2017, and (f) 2011, spatially integrated for the black-shaded area in (a). Wind rose plots of 3-hourly wind data at Station 04312 (Station Nord as ▲) during February of (g) 2018, (h) 2017, and (i) 2011: wind speed (U , m/s) and direction. The missing data are indicted as white in each color bar.

010



Deleted:
 Deleted: black
 Deleted: black




Figure 5. Monthly integrated sea ice melt terms over the northern Greenland region during February 2018 from the RASM hindcast simulation: (a) surface top, (b) lateral, and (c) basal melt (cm).

Deleted:






Figure 6. Four major patterns with self-organizing maps (frequency of occurrence as %) of the near-surface 6-hourly wind fields from the RASM simulation during the time periods including before and after the 2018 polynya event: (a) 5-11 February (93%), (b) 15-20 February (83%), (c) 20-26 February (93%), and (d) 3-13 March (95%). The wind vectors were sub-sampled at every 3 grid-cells for a plotting purpose. RASM daily sea ice divergence (%/day) within each period above is shown on (e) 10 February, (f) 16 February, (g) 24 February, and (h) 4 March 2018.



025



Deleted:




030 Figure 7. (a)-(j) Satellite-derived (NASA team algorithm) daily mean sea ice concentration (SIC, μ ; black) for the northern Greenland region (see Fig. 4a) during January-March from 2011 to 2020. The grey shading depicts one standard deviation (σ ; gray) from the mean.



Deleted:

035 Figure 8. The northward wind component greater than 5 m/s from the RASM hindcast simulation during February of 1980-2020 at the nearest grid to (a) Station 04312 (Station Nord) and (b) Station 04301 (Cape Morris Jesup). Each square represents the six-hourly surface wind and its color indicates a magnitude of wind speed.



040 **Figure 9.** Mean sea ice concentration (NASA Team algorithm) on 15 December 1986 over the northern Greenland region from (a) the Nimbus-7 SMMR and (b) the RASM hindcast simulation. (c) Monthly integrated sea ice melt (top, lateral, and basal) for December 1986 and (d) near-surface (at 10 m) wind fields on 12 December 1986 from the RASM hindcast simulation. (e) RASM daily near-surface air temperature (light blue) from the nearest grid cell to Station 04301 (Cape Morris Jesup) and daily AO index (blue) in December 1986. The correlation coefficient (r) between the AO index and the simulated air temperature is shown in the upper-left. (f) Time rate of change of the RASM daily sea ice volume (km^3/day ; blue) due to dynamic tendency during December 1986, spatially integrated for the region shown in Fig. 4a. Daily mean net turbulent (sensible and latent) heat flux (W/m^2 ; red) is also shown over the same region where sea ice thickness is less than 10 cm from the RASM hindcast simulations. The negative values indicate loss from the region.





Figure 10. Sea ice thickness (SIT, m), sea ice dynamic volume tendency (DVT, km^3/day) for the northern Greenland region (see Fig. 4a), and north(positive)-south(negative) wind component (m/s) during winter months from the ensemble member #4 in the 1985-initialized run and the ensemble member #2 in the 2015-initialized run: SIT on (a) 22 January 1988 and (b) 14 January 2024, daily DVT in January-March of (c) 1988 and (d) 2024, and six-hourly north-south wind in January-March of (e) 1988 and (f) 2024.

Deleted:

Formatted: Font:(Default) +Theme Body (Times New Roman), 9 pt, Bold

Deleted: 9

Meier, W. N., Fetterer, F., Savoie, M., Mallory, S., Duerr, R., & Stroeve, J. (2017a). *NOAA/NSIDC Climate Data Record of Passive Microwave Sea Ice Concentration, Version 3*. Boulder, Colorado USA. NSIDC: National Snow and Ice Data Center. doi: <https://doi.org/10.7265/N59P2ZTG>. Accessed 27 Sep 2019.

Meier, W. N., Fetterer, F., & Windnagel, A. K. (2017b). *Near-Real-Time NOAA/NSIDC Climate Data Record of Passive Microwave Sea Ice Concentration, Version 1*. doi: <https://doi.org/10.7265/N5FF3QJ6>.