

Competing influences ~~Net effect of the ocean, atmosphere and solid earth on ice-sheet-atmosphere interactions reduces simulated transient Miocene Antarctic ice sheet variability~~

Lennert B. Stap¹, Constantijn J. Berends¹, Meike D.W. Scherrenberg¹, Roderik S.W. van de Wal^{1,2}, and Edward G.W. Gasson³

¹Institute for Marine and Atmospheric research Utrecht, Utrecht University, 3584 CC Utrecht, the Netherlands

²Faculty of Geosciences, Department of Physical Geography, Utrecht University, Utrecht, the Netherlands

³College of Life and Environmental Sciences, University of Exeter, Exeter, United Kingdom

Correspondence: L.B. Stap (L.B.Stap@uu.nl)

Abstract. Benthic $\delta^{18}\text{O}$ levels vary strongly during the warmer-than-modern early- and mid-Miocene (23 to 14 Myr ago), suggesting a dynamic Antarctic ice sheet (AIS). So far, however, realistic simulations of the Miocene AIS have been limited to equilibrium states under different CO_2 levels and orbital settings. Earlier transient simulations lacked ice-sheet-atmosphere interactions, and used a present-day rather than Miocene Antarctic bedrock topography. Here, we quantify the effect of ice-sheet-atmosphere interactions, running IMAU-ICE using climate forcing from Miocene simulations by the general circulation model GENESIS. Utilising a recently developed matrix interpolation method enables us to interpolate the climate forcing based on CO_2 levels (between 280 and 840 ppm) as well as [varying ice sheet configurations](#) (between no ice and a large [East Antarctic ice sheet](#)). We furthermore implement recent reconstructions of Miocene Antarctic bedrock topography. We find that the positive albedo-temperature feedback, partly compensated by ~~the negative ice volume-precipitation feedback a~~ [negative feedback between ice volume and precipitation](#), increases hysteresis in the relation between CO_2 and ice volume (V). Together, these ice-sheet-atmosphere interactions decrease the amplitude of ~~AIS variability caused by~~ [Miocene AIS variability in idealised transient simulations. Forced by quasi-orbital 40-kyr forcing \$\text{CO}_2\$ cycles, the ice volume variability reduces by 21% in transient simulations when ice-sheet-atmosphere interactions are included, compared to when forcing variability is only based on \$\text{CO}_2\$ changes.](#) Thereby, ~~they these interactions~~ also diminish the contribution of AIS variability to benthic $\delta^{18}\text{O}$ fluctuations. ~~Furthermore, we show that under equal atmospheric and oceanic forcing, the amplitude of 40-kyr transient AIS variability becomes 10% smaller~~ [Evolving bedrock topography](#) during the early- and mid-Miocene ~~, due to the evolving bedrock topography. Lastly, we quantify the influence of ice shelf formation around the Antarctic margins, by comparing simulations with Last Glacial Maximum (LGM) basal melt conditions, to ones in which ice shelf growth is prevented. Ice shelf formation increases hysteresis in the CO_2 - V relation, and amplifies~~ [reduces ice volume variability by 10%, under equal 40-kyr AIS variability by 19% using LGM basal melt rates, and by 5% in our reference setting cycles of atmosphere and ocean forcing.](#)

1 Introduction

The dynamics of the Antarctic ice sheet (AIS) during the early- and mid-Miocene (23 to 14 Myr ago) are a topical issue in (paleo)climate science. This issue is of great interest, because ~~the climatic conditions over the course~~ of this period the climatic conditions, such as the atmospheric CO₂ concentration (Kürschner et al., 2008; Foster et al., 2012; Badger et al., 2013; Greenop et al., 2014; Super et al., 2018; Steinhorsdottir et al., 2021b), ranged from those of the recent past to those predicted for the future (~200 to >800 ppm, see Steinhorsdottir et al. (2021a), for a review). Benthic $\delta^{18}\text{O}$ records show sizeable orbital-timescale fluctuations during this time, an indication of potential vigorous Antarctic ice volume variability (e.g. Zachos et al., 2008; Liebrand et al., 2011, 2017; Holbourn et al., 2013; Levy et al., 2019). The $\delta^{18}\text{O}$ signal is, however, also shaped by deep-sea temperature changes. How much of the signal is committed by each component, and thus how the AIS evolved during the Miocene, is currently still under debate. On the one hand, geological evidence supports strong AIS dynamism (Pekar and DeConto, 2006; Shevenell et al., 2008), with ice periodically advancing and retreating e.g. over Wilkes Land (Sangiorgi et al., 2018) and the Ross Sea sector (Hauptvogel and Passchier, 2012; Levy et al., 2016; Pérez et al., 2021). Some studies, on the other hand, advocate a more stable small AIS, based on ice-proximal sea-surface temperature records from dinoflagellate cyst assemblages (Bijl et al., 2018) and TEX₈₆ data (Hartman et al., 2018). In addition, a recent analysis of clumped isotope and Mg/Ca records has found high bottom-water temperatures, which combined with the benthic $\delta^{18}\text{O}$ record paradoxically suggests a Miocene AIS volume persistently larger than at present day (Modestou et al., 2020). Finally, using climate modelling and data comparisons, it has been suggested that the AIS varied mostly in spatial extent, rather than volume, which imprinted on Miocene $\delta^{18}\text{O}$ variability through a strong effect on deep-water temperatures (Bradshaw et al., 2021).

Ice-sheet models can be used to understand the physical processes driving Miocene AIS variability. Broadly stated, earlier modelling efforts have followed two approaches to this task. One approach is to use ice-sheet models bidirectionally coupled to climate models, with as much physical realism, in terms of processes included and resolution, as is computationally feasible for long-term simulations. Gasson et al. (2016) adopted this approach, using an isotope-enabled setup of a 3D ice-sheet model coupled to a regional climate model nested in a general circulation model (GCM). They showed the importance of including ice-sheet-climate feedbacks, ice-calving parameterisations, and changes in the ice sheet's oxygen isotopic composition, for simulating strong AIS-induced benthic $\delta^{18}\text{O}$ changes. Halberstadt et al. (2021) used the same model set-up to determine the influence of CO₂ changes and uplift of the Transantarctic Mountains on glacial variability during the Miocene. These studies were so far limited to steady-state simulations, in which the climate forcing is kept constant until the ice sheet has equilibrated. Transient ice sheet behaviour can be very different, as was e.g. demonstrated for the North American and Eurasian ice sheets during the Late Pleistocene (Abe-Ouchi et al., 2013). The second approach, which is the one taken in this study, is therefore to focus on transient AIS changes, applying gradually changing climatic conditions. Earlier studies employed 1D ice-sheet models in combination with parameterised temperatures (De Boer et al., 2010), or relatively simple climate models (Langebroek et al., 2009, 2010; Stap et al., 2017). Recently, Stap et al. (2019) showed that orbital-timescale CO₂ changes cause smaller transient AIS variability than equilibrium simulations suggest, using the 3D thermodynamical Parallel Ice Sheet

55 Model (PISM) forced by Miocene climates obtained from GCM (COSMOS) simulations (Stärz et al., 2017). Conceptual considerations indicate that the disequilibrium between ice volume and CO₂ also leads to out-of-phase changes of these quantities (Stap et al., 2020). However, the COSMOS climate simulations were performed using only one ice sheet configuration. Hence, the effect of surface whitening (darkening) by ice sheet advance (retreat) on temperature, the so-called albedo-temperature feedback, was not included. Neither was the ~~ice-volume-precipitation-feedback, which causes precipitation on the ice sheet to decrease with increasing ice volume~~
60 ~~ice-sheet desertification effect, i.e. the reduction of snow deposition on a growing ice sheet (e.g. Oerlemans, 2004).~~ Furthermore, the effect of mass balance changes due to surface height changes (surface-height-mass-balance feedback) could only be crudely captured using a scalar lapse-rate correction for temperature, which is known to be a simplification (Fortuin and Oerlemans, 1990). Moreover, a present-day Antarctic bedrock topography was used, while during the Miocene the topography was different, which also significantly affects the surface mass balance and ice flow (Stap et al.,
65 2016; Colleoni et al., 2018; Paxman et al., 2020).

In this study, we ~~examine the effect of~~ expand the scope of previous studies by performing transient simulations (cf. Gasson et al., 2016; Colleoni et al., 2018; Paxman et al., 2020; Halberstadt et al., 2021), now including ice-sheet-atmosphere
~~feedbacks, as well as bedrock topography, on transient Miocene AIS variability~~interaction
70 ~~(cf. De Boer et al., 2010; Stap et al., 2016, 2017).~~ We run the 3D thermodynamical ice-sheet model IMAU-ICE (De Boer et al., 2013; Berends et al., 2018), that is based on the commonly-used shallow ice and shallow shelf approximations and therefore comparable to PISM. Climate forcing is obtained from Miocene simulations by the GCM GENESIS, in which different CO₂ levels and ice sheet configurations ranging from zero ice to a substantial ~~ice volume~~ East Antarctic ice sheet are applied (Burls et al., 2021). Utilising a recently developed matrix interpolation routine (Berends et al., 2018), this set-up enables us to in-
75 terpolate the climate forcing based on land ice changes in addition to CO₂ changes. In this manner, the ~~albedo-temperature and ice-volume-precipitation feedbacks are represented, and because effects of ice sheet changes on local temperatures and precipitation are represented. Because~~ surface height changes are incorporated in the forcing, the surface-height-mass-balance feedback is captured more realistically as well. ~~Performing several experiments that include both equilibrium~~ We perform both equilibrium simulations, keeping the forcing CO₂ level constant, and idealised transient experiments, in which we vary the CO₂
80 level on long and short quasi-orbital timescales (40, 100 and transient simulations, we test the influence of these feedbacks under varying CO₂ levels. Furthermore changes. In this way, we quantify the ~~effect on transient Miocene AIS volume evolution of using bedrock topographies (Hochmuth et al., 2020a; Paxman et al., 2019)~~
influence of the ice-sheet-atmosphere feedbacks on the variability of the AIS during the Miocene. In addition, we investigate
85 the effect of bedrock topography changes over the Miocene, by using topographies pertaining to the early- (23-24 Myr ago) and mid-Miocene (14 Myr ago) , as well as the (Hochmuth et al., 2020a; Paxman et al., 2019). We also carry out simulations using Eocene (34 Myr ago) and present-day ~~present-day~~ bedrock topographies. Lastly, we ~~investigate where and under which conditions ice shelves form, and what their influence on the grounded ice volume is~~ test the sensitivity of our results to the parameterisation of basal melt underneath the floating ice shelves. This is achieved by performing ~~and comparing simulations~~

90 ~~, in which relatively mild, and severe basal melt rates underneath the ice shelves are applied~~ identical simulations as our reference experiment, but imposing relatively mild as well as harsh sub-shelf melt rates, easing and prohibiting ice shelf growth respectively. We quantify where and under which conditions ice shelves form and perish in our simulations, and how they influence the variability of the grounded ice volume.

2 Models and methods

95 For this study, we employ the ice-sheet model IMAU-ICE (v1.1.1-MIO) (De Boer et al., 2013; Berends et al., 2018) to simulate the Antarctic ice sheet, using atmospheric input data obtained from Miocene climate simulations by the GCM GENESIS (Burls et al., 2021).

2.1 Ice-sheet model

IMAU-ICE is a three-dimensional thermodynamical ice-sheet model, which uses a superposition of the shallow ice approxi-
 100 mation (SIA) and shallow shelf approximation (SSA) to simulate the dynamics of grounded and floating ice. In the interior of the ice sheet, the ice flow is dominated by the SIA, while towards the fast-flowing ice streams on the fringes the SSA gains importance, becoming the governing component on the ice shelf (Winkelmann et al., 2011). No additional grounding-line parameterisations are included. We use a ~~40 x 40 km~~ 40x40-km resolution grid covering the Antarctic continent. This resolution ensures the feasibility of performing a large ~~numbers~~ number of simulations, while still capturing the Antarctic ice flow in
 105 the amount of detail we deem appropriate for our purpose. Longitude and latitude coordinates are assigned to the grid using an inverse oblique stereographic projection (Reerink et al., 2010). For further settings, please see Appendix A.

2.1.1 Basal mass balance

Basal melt underneath the ice shelves (~~M~~ $M(x, y)$) is calculated using a combination of the parameterisation of Pollard and DeConto (2009) and a linear relation to ocean temperature change (Beckmann and Goosse, 2003; Martin et al., 2011):

$$110 \quad M(x, y) = (z_{deep}(x, y) - 1)[(1 - z_{expo}(x, y))S_m(x, y) + z_{expo}(x, y)M_{expo}] - (z_{deep}(x, y)M_{deep}), \quad (1)$$

with

$$S_m(x, y) = \rho_w c_p \gamma_T \frac{T_w - T_{freeze}}{H_{fus} \rho_i} \frac{T_w - T_{freeze}(x, y)}{H_{fus} \rho_i}. \quad (2)$$

~~Here, the weighing factors~~ $z_{deep} = \frac{h_w - 1800 \text{ m}}{200 \text{ m}}$ ~~Weighing factor~~ z_{deep} grows with water depths ($h_w(x, y)$) larger than 1800 m:

$$z_{deep}(x, y) = \frac{h_w(x, y) - 1800 \text{ m}}{200 \text{ m}}, \quad (3)$$

115 and $z_{expo} = \frac{\alpha_{sub} - 80^\circ}{30^\circ} e^{-0.4 * r_{open}}$ ~~depend on the water depth (h_w), $z_{expo}(x, y)$ depends on~~ the widest subtended angle to the open ocean (~~α_{sub}~~ $\alpha_{sub}(x, y)$) and the shortest linear distance to the open ocean (~~r_{open}~~ $r_{open}(x, y)$):

$$z_{expo}(x, y) = \frac{\alpha_{sub}(x, y) - 80^\circ}{30^\circ} e^{-0.4 * r_{open}(x, y)}. \quad (4)$$

Both factors are clipped at 0 and 1. The melt rate for the exposed shelves (M_{expo}) varies linearly between 3 m/yr for an applied CO₂ concentration of 280 ppm to 6 m/yr for CO₂ concentrations of 400 ppm and higher. The melt rate for the deep-water shelves (M_{deep}) similarly varies between 5 and 10 m/yr. Melt rate $S_m S_w(x, y)$ is dependent on the water density (ρ_w), specific heat (c_p), thermal exchange velocity (γ_T), water temperature (T_{water}), pressure-adjusted freezing point (~~T_{freeze}~~ $T_{freeze}(x, y)$), latent heat of fusion (H_{fus}) and shelf ice density (ρ_i). Because GENESIS has a slab-ocean component (Sect. 2.2), realistic water temperatures cannot be taken from the GCM results (Gasson et al., 2016; Halberstadt et al., 2021). Instead, we use an ad hoc global ocean temperature parametrisation, adapted from De Boer et al. (2013). Similar to melt rates M_{expo} and M_{deep} , T_w varies linearly as a function of CO₂ between -1.7°C (280 ppm) and 2°C (≥ 400 ppm).

Currently, calving is neglected. The only calving routine implemented in IMAU-ICE is thickness calving, i.e. calving floating ice with a thickness below a certain threshold. We exclude this in our simulations, because it could hinder the growth of new ice shelves (e.g. Ritz et al., 2001). In combination with the heavily parameterised sub-shelf melt calculation, this makes the mass balance of the ice shelves arguably a weak point of our modelling effort. ~~Therefore, in a separate sensitivity experiment We therefore perform sensitivity experiments~~ (Table 1) ~~, we to investigate how our results depend on ocean forcing. We~~ consider an extreme case using Last Glacial Maximum (LGM)-like melt parameter settings: $M_{expo} = 0$ m/yr, $M_{deep} = 2$ m/yr, and $T_w = -5^\circ\text{C}$ (De Boer et al., 2013). In another experiment, ice shelves are effectively inhibited from growing by applying a constant melt rate of 400 m/yr similar to the ABUM experiment of the ice-sheet model intercomparison project ABUMIP (Sun et al., 2020). By comparing the results of these extreme scenarios to our reference experiment, we can quantify where and under which conditions ice shelves form, and how they subsequently influence the variability of the grounded ice volume.

2.1.2 Surface mass balance

The surface mass balance (SMB) calculation uses precipitation (~~$P_{applied}$~~ $P_{applied}(x, y)$) and 2-m air temperature ~~$T_{applied}$~~ $T_{applied}(x, y)$), obtained from Miocene climate simulations by GENESIS (Sect. 2.2) utilising a matrix interpolation routine (Sect. 2.3, Appendix B). The GENESIS fields are remapped from their native T31 grid (Sect. 2.2) to the ice-sheet model grid using bilinear interpolation based on longitude and latitude. Monthly SMB is the sum of accumulation and refreezing, minus ablation (Berends et al., 2018). Refreezing is dependent on the available liquid water content, and the superimposed water content (Huybrechts and de Wolde, 1999; Janssens and Huybrechts, 2000). Accumulation (~~Acc~~ $Acc(x, y)$) is calculated as:

145 $Acc(x, y) = P_{applied}(x, y) * f_{snow}(x, y), \quad (5)$

with f_{snow} the temperature-dependent snow fraction (Ohmura, 1999). Ablation ($Abl(x, y)$) follows from an insolation-temperature melt parameterisation (Bintanja et al., 2002):

$$Abl(x, y) = c_1(T_{applied}(x, y) - 273.15 K) + c_2(Q_{TOA}(x, y) * (1 - \alpha(x, y))) + c_3, \quad (6)$$

where α is the internally-calculated surface albedo, and the parameters are set to $c_1 = 0.788 \text{ m yr}^{-1} \text{ K}^{-1}$, $c_2 = 0.004 \text{ m}^3 \text{ J}^{-1}$, and $c_3 = -0.50 \text{ m yr}^{-1}$. Incoming solar radiation ($Q_{TOA}(x, y)$) is derived as follows. First, a background albedo ($\alpha_{bg}(x, y)$) is determined: 0.1 for ocean-covered grid cells, 0.2 for soil, and 0.5 for ice. Thereafter, the effect of snow, with an albedo (α_{snow}) of 0.85, is added:

$$\alpha(x, y) = \alpha_{snow} - (\alpha_{snow} - \alpha_{bg}(x, y)) * \exp[-15 * z_{firn}(x, y)] - 0.015 * Abl_{yr-1}(x, y), \quad (7)$$

where $Abl_{yr-1}(x, y)$ is the ablation calculated for the previous year. The firn layer depth ($z_{firn}(x, y)$, clipped at 0 m and 10 m) is changed over time by addition of snowfall minus melt. The albedo $\alpha(x, y)$ is not allowed to be lower than the background albedo or higher than the albedo of snow. In Eq. 6, incoming solar radiation at the top of the atmosphere ($Q_{TOA}(x, y)$) is taken from Laskar et al. (2004), and kept constant at the present-day values, so as to isolate climate variability caused by CO_2 changes. This surface mass balance parameterisation has recently been shown to yield realistic results over the present-day Greenland ice sheet (Fettweis et al., 2020).

2.2 Climate model

GENESIS (version 3.0) (Thompson and Pollard, 1997; DeConto et al., 2012) has a spectral resolution of T31 ($\sim 3.75^\circ \times 3.75^\circ$). The atmospheric model is coupled to $2^\circ \times 2^\circ$ surface models including a non-dynamical 50-m slab diffusive mixed-layer ocean with dynamical sea ice, and dynamic vegetation (BIOME4). The applied paleogeography comes from Herold et al. (2008), with dedicated modifications to Antarctica based on simulations by DeConto and Pollard (2003). Different simulations were carried out as described in Burls et al. (2021); a subset is used in this study. The simulations used here have a distinct CO_2 level (280 ppm, 840 ppm), Antarctic ice sheet configuration (volumes of 0 km^3 (no ice) and $18.975 \times 10^6 \text{ km}^3$ (full ice sheet)), and orbital settings (eccentricity = 0.05, obliquity = 22.5° , longitude of precession = 90° (cold Antarctic summer), ecc. = 0, obl. = 23.5° , longitude of precession = N/A (medium, modern day settings), ecc. = 0.05, obl. = 24.5° , longitude of precession = 270° (warm Antarctic summer)).

For our reference simulations, we use the 2-m air temperature, precipitation, and surface height fields of simulations. Here, we use simulation 1fumebi (cold) and for the cold climate, and simulation 3nomebi (warm), bilinearly interpolated to the ice-sheet model grid for the warm climate (Fig. S1). Simulation 1fumebi has a CO_2 level of 280 ppm ($1 \times$ pre-industrial (PI) level), a full ice sheet, and medium Antarctic summers, while simulation 3nomebi has an AIS configuration with a full East Antarctic ice sheet ($18.975 \times 10^6 \text{ km}^3$ volume, area shown in Fig. S2). Simulation 3nomebi is forced by a CO_2 level of 840 ppm ($3 \times$ PI), no ice sheet, and medium Antarctic summers. The suffix 'bi' in the names of the GENESIS simulations indicates that with no ice cover. Both simulations use present-day orbital settings. The vegetation on non-glaciated land, which

has an influence on the regional climate, is determined by the dynamic vegetation model BIOME4~~was used, which is of no direct concern for the current study.~~

2.3 Matrix interpolation method

180 The applied climate forcing is interpolated between the cold (1fumebi) and warm (3nomebi) GENESIS fields using a matrix interpolation method (Berends et al., 2018), described in detail in Appendix B. Briefly, the applied temperature interpolation is based on a prescribed value for CO₂ and on the internally modelled ice sheets. The higher (lower) the CO₂ level, and the smaller (larger) ice coverage, the more the climate forcing is like the warm (cold) climate. The feedback of the ice sheets on the climate is calculated via the effect on absorbed insolation through changes in surface albedo. The precipitation
185 interpolation is determined solely from the ice volume. Because the cold climate is generally drier, this constitutes a negative ice-volume-precipitation feedback, representing the ice-sheet desertification effect in our simulations.

2.4 Simulations

We perform an experiment using our default settings (REF), as well as ~~12-11~~ sensitivity experiments with settings altered as indicated in Table 1. Each experiment consists of 15 simulations, 11 of which are steady-state, and 4 of which are transient
190 simulations (Table 2).

The steady-state simulations are run for 150 kyr with a constant forcing CO₂ level; enough time for the ice sheet to equilibrate with the forcing climate. Mind that the CO₂ level, rather than the climate itself, is kept fixed, which means the climate can still change due to land ice changes. First, six of these steady-state simulations are run, using CO₂ levels bracketing the
195 280 to 840 ppm range: 280, 392, 504, 616, 728, and 840 ppm. In our reference case, the initial conditions are obtained from recent reconstructions of the Antarctic bathymetry (Hochmuth et al., 2020a) and bedrock topography (Paxman et al., 2019) pertaining to 23 to 24 million years (Myr) ago (dataset: Hochmuth et al., 2020b). No ice is present at the start of these runs. ~~The CO₂ levels are set to different levels bracketing the 280 to 840 ppm range: 280, 392, 504, 616, 728, and 840 ppm.~~ These simulations constitute the lower, ascending branch of the hysteresis curve in the equilibrium CO₂-ice volume relation (blue
200 ~~lines-connected diamonds~~ in figures, e.g. Fig. 1b). Thereafter, the remaining five steady-state simulations are continued from results of the 280-ppm IMAU-ICE run, with CO₂ levels of 392, 504, 616, 728, and 840 ppm. These simulations constitute the upper, descending branch of the hysteresis curve in the equilibrium CO₂-ice volume relation (red ~~lines-connected diamonds~~ in figures, e.g. Fig. 1b).

205 Next, we perform the ~~transient simulations~~idealised transient simulations (Table 2). Three of these are ~~continued from the started from the final state of the equilibrium~~ 840-ppm IMAU-ICE run. The evolution of CO₂ over time has a triangular shape (e.g. pink line in Fig. 1a). The CO₂ is first linearly reduced from 840 to 280 ppm, and then increased back to 840 ppm~~.~~ In the three simulations, this triangular CO₂ evolution is implemented over the course of 100 ~~kyr~~ and 400 kyr, and five consecutive times over 40 kyr (200 kyr ~~total~~in total; e.g. pink line in Fig. 1c). The final transient simulation is continued from the 280-ppm

210 IMAU-ICE run. ~~In this simulation, and forced with an inverted triangular 40-kyr CO₂ is first gradually evolution. That means the CO₂ is first linearly increased from 280 to 840 ppm, and then reduced-decreased back to 280 ppm five consecutive times over 40 kyr. This is essentially the normal transient 40-kyr run mirrored. (e.g. purple line in Fig. 1c).~~

3 Results

215 3.1 Reference simulations

As a minimal check for ~~our model~~ model performance and parameter settings, we first perform steady-state ~~simulations-control simulations as in De Boer et al. (2015). These are executed~~ using ERA40 present-day (PD; 1957-2002) precipitation and temperature forcing (Uppala et al., 2005). Constant present-day-like melt parameter settings are used to calculate the basal mass balance: $M_{exp} = 3$ m/yr, $M_{deep} = 5$ m/yr, and $T_w = -1.7^\circ\text{C}$. Depending on whether the run is started from Bedmachine PD
220 ice and topography (Morlighem et al., 2020) or without ice and an isostatically rebounded topography, we simulate a total ice volume above floatation of 56.6 meter sea level equivalent (m.s.l.e.; 23.1×10^{15} m³) or 55.0 m.s.l.e. (22.4×10^{15} m³) respectively. This is very close to the 55.0 m.s.l.e. obtained from remapping the Bedmachine data to our grid. Like other SIA/SSA-based ice-sheet models (c.f. De Boer et al., 2015), IMAU-ICE generally simulates slightly thinner ice in the interior and thicker ice along the margins (Fig. ~~S2~~S3). The grounding line is simulated quite well, although slightly more advanced in the Filchner-Ronne,
225 and the Amery embayments. Due to lack of calving, the simulated ice shelves are extended much farther than the observations. However, since the increase in back pressure due to ~~the extra shelf area~~ these overestimated shelf areas is very small (Reese et al., 2018b), this is unlikely to affect the evolution of the grounded ice. In this study, we will therefore focus on grounded ice.

In the reference equilibrium simulations using Miocene forcing (experiment REF), the AIS consists only of disconnected
230 glaciers in high mountainous regions in case of a 840-ppm CO₂ level ~~-(Fig. 2a).~~ It grows into a full East AIS (EAIS) and small West AIS (WAIS) in case of 280-ppm forcing (Fig. 2b). In the latter case, the WAIS grounds on Marie Byrd Land and the Peninsula, while not growing into the Ross and Filchner-Ronne embayments. ~~The~~ Intermediate states for different forcing CO₂ levels are shown in Fig. S4.

235 We show the reference CO₂-equilibrium ice volume (above floatation) (V_{eq}) relation ~~decreases monotonically and shows considerable hysteresis (in Fig. 1b). Depending on the timescale of the imposed forcing variability, the transient simulations yield peak~~ (Pollard and DeConto, 2005). It decreases monotonically and exhibits considerable hysteresis, as displayed by the difference between the ascending (blue) and descending (red) branches. Transient simulations with different imposed timescales for the forcing variability are indicated by the dashed (400 kyr) and dotted (100 kyr) black lines. Peak ice volumes
240 of 45.48 45.5 m.s.l.e. (400 kyr) and 37.81 37.8 m.s.l.e. (100 kyr) are yielded. In agreement with Stap et al. (2020), this peak is only reached ~~after later in time than~~ the CO₂ minimum (Fig. 1a), ~~and relatively earlier.~~ The peak is attained relatively earlier in time (Fig. 1a), i.e. at a lower CO₂ level (Fig. 1b), in the 400-kyr simulation than in the shorter 100-kyr simulation. In case of

40-kyr forcing cycles, the equilibrium cycle is attained very quickly and does not depend on the initial conditions (Fig. 1c,d; black lines partly overlapping). In the equilibrium cycle, the ice volume varies between 3.6 m.s.l.e. at 728 ppm ~~to and~~ 30.7 m.s.l.e. at 532 ppm, a 27.1 m.s.l.e. difference. Since the maximum ice volume is reached longer after the minimum CO₂ level (252 ppm difference) than the minimum ice volume after the maximum CO₂ level (112 ppm difference), the growth phase lasts longer than the decay phase. That also means that ice sheet decay is on average 40% faster than growth, but both rates, in particular the growth rate, are certainly not constant. In the transient simulations, the AIS initially builds up slowly in the high mountainous regions at ~ 0.1 m.s.l.e./kyr, speeding up to a maximum of 3 m.s.l.e./kyr when the surface mass balance (SMB) becomes positive on a large part of the continent as separate ice domes start to merge ~~-(gradients in Fig. 1c)~~. When the transient ice volume is in between the ascending and descending branches of the equilibrium CO₂-V_{eq} relation, ~~the rate of i.e.~~ larger than the ascending (blue) volume but smaller than the descending (red) volume (Fig. 1d), the rate of transient ice volume change is relatively small (< 1 m.s.l.e./kyr).

In earlier research using IMAU-ICE, different values for ~~tuning the SMB~~ parameter c_3 in Eq. 6 have been used (e.g. De Boer et al., 2013, 2014; Berends et al., 2018). Our reference setting $C_3 = 0.50 \text{ m yr}^{-1}$ leads to satisfactory PD results, but using $C_3 = 0.25 \text{ m yr}^{-1}$ as De Boer et al. (2014) did, a similar PD ice volume of 55.5 m.s.l.e. is obtained (not shown). Using this setting (experiment SMB; Table 1) increases surface melt, with as a result onset of AIS growth and decay at lower CO₂ levels (Fig. ??a,b). ~~We choose our REF setting of $C_3 = 0.50 \text{ m yr}^{-1}$, such that variability in our simulated CO₂ domain is maximised.~~
~~A different tuning might lead to quantitatively different CO₂ levels, but not qualitatively. Furthermore, in the REF simulations we ignore the effect of orbital forcing variability completely. Performing experiment INSOL (Table 1), however, we make an assessment of the indirect effect of insolation changes (through temperature) on Miocene AIS variability. We now interpolate between the results of GENESIS simulations 1fucobi (cold Antarctic summer) and 3nowabi (warm Antarctic summer). Mind that the influence of insolation on SMB (through the second term on the r. h.s. of Eq. 6) remains constant at PD level. The indirect insolation effect changes in concert with CO₂; we therefore recalculate the weight given to S5). Our REF settings yield an ice sheet similar in extent to the results of Stap et al. (2019) and DeConto and Pollard (2003) at 280 ppm, which disappears almost completely at 840 ppm. This ensures that we can most effectively study large-scale variability of the East-Antarctic ice sheet. A different tuning would shift the CO₂ changes compared to ice extent changes in our interpolation method (Eq. B5) as $\epsilon_{CO_2} = 0.72$ in this case. In these simulations, the maximum equilibrium ice volume (at 280 ppm) is larger: 54.2 m.s.l.e. compared to 48.4 m.s.l.e. in case REF. This volume increase is established mostly in regions with low bedrock topography, through a thicker WAIS in Marie Byrd Land and Ellsworth Land, and EAIS in MacRobertson Land and Coats Land (not shown). Furthermore, hysteresis in the at which these ice sheet sizes are simulated, but does not (qualitatively) affect the shape of the CO₂-V_{eq} relation is diminished, in consonance with Langebroek et al. (2009), and transient ice volume variability is amplified (Fig. ??c,d). In this study, we perform idealised simulations, focussing on influences of the ocean, atmosphere and solid earth on transient Miocene AIS variability. When aiming to make realistic simulations, however, the INSOL simulations indicate that the matrix interpolation routine should be extended to include orbital forcing. ice volume relation.~~

3.2 Influence of ice-sheet-atmosphere interaction~~between ice sheets and atmosphere~~

Using a matrix interpolation routine enables us to interpolate the climate forcing for our ice-sheet model based on both CO₂ and ~~ice-volume~~ice-sheet size. Here, we make a comparison to results of our model set-up using a simpler ~~glacial~~-index routine, as e.g. used in Stap et al. (2019), in which the interpolation is solely based on the CO₂ level. In experiment ~~FEEDB-NOFEEDB~~ (Table 1), we implement $w_{tot} = \frac{CO_2 - CO_{2,cold}}{CO_{2,warm} - CO_{2,cold}}$ instead of Eqs. B7 (for temperature) and B11 (for precipitation) to interpolate between the cold ~~and warm climates~~.(1fumebi) and warm (3nomebi) climates.

Because the positive albedo-temperature feedback is not invoked in this case, hysteresis in the CO₂-V_{eq} relation is smaller (Fig. 3b, solid and dashed, red and blue lines). The temperature declines more quickly towards lower CO₂ values, causing the equilibrium simulations to yield larger ice volumes ~~-(Fig. 3b, blue lines)~~. At 280 ppm, V_{eq} in the NOFEEDB case is 55.8 m.s.l.e., compared to 48.4 m.s.l.e. in the difference in ice thickness with the REF simulation is again REF case. This difference in ice volume is mainly manifested in regions with low bedrock topography, through a thicker WAIS in Marie Byrd Land, and EAIS in MacRobertson Land and Coats Land (not shown). Because the AIS deglaciates at similar levels as in the REF case (Fig. 3b, red lines), the difference between the CO₂ levels of glaciation and deglaciation is smaller in the NOFEEDB case. That reduces the hysteresis in the CO₂-V_{eq} relation.

This also results in larger transient ice volume variability (Fig. 3a,b)~~-, black and orange lines~~. Peak ice volumes are increased to 54.3 m.s.l.e. (400 kyr forcing CO₂ cycle) and 47.4 m.s.l.e. (100 kyr). Forced by 40-kyr CO₂ cycles, the amplitude of variability is 34.1 m.s.l.e. in the equilibrium cycle ~~-, 26% larger than in the REF case. Stated the other way around, the in the NOFEEDB case. That means the~~ ice-sheet-atmosphere interactions, that are included in the REF case, decrease the amplitude of AIS variability by 21%.

As mentioned ~~before~~above, the difference between the REF and ~~FEEDB-NOFEEDB~~ results is mainly governed by the albedo-temperature feedback. However, in the REF case, precipitation decreases with increasing ice volume, via Eq. B10 combined with a ~~drier climate at lower CO₂~~cold climate that is generally drier. This constitutes a negative ice-volume-precipitation feedback. The effect of this feedback alone is shown by a comparison ~~to between~~ the results of experiment ~~FEEDB-onlyP~~ NOFEEDB and those of NOFEEDB-P (Table S1, Fig. 3c,d). In this experiment, the ~~glacial~~-index method is only used for precipitation, and temperature is interpolated in the normal ~~manner~~-sense (Eq. B7). The hysteresis in the CO₂-V_{eq} relation is now larger, and the amplitude of transient variability in the equilibrium cycle of the 40-kyr simulation is 19.3 m.s.l.e., ~~29.43%~~ smaller than in the ~~REF-NOFEEDB~~ case. In contrast, using the ~~glacial~~-index method only for temperature (experiment ~~FEEDB-onlyT~~NOFEEDB-T), the amplitude of 40-kyr transient variability becomes 6% larger (36.1 m.s.l.e.) than in the regular ~~FEEDB case. In conclusion, NOFEEDB case. The results of experiment NOFEEDB-T are much closer to those of experiment NOFEEDB. Therefore, of the albedo-temperature feedback is the strongest of these two ice sheet-atmosphere feedbacks, the~~ albedo-temperature feedback is dominant in our experiments.

3.3 ~~Influence~~ Effect of bedrock topography changes

Recent geological reconstructions have indicated that Antarctic bedrock topography has evolved substantially since the Eocene-Oligocene boundary (34 Myr ago), mostly due to thermal subsidence and glacial erosion (Paxman et al., 2019). In the REF simulations, we have used the bedrock topography reconstruction pertaining to the Oligocene-Miocene boundary (23-24 Myr ago). In experiment TOPO 14Ma (Table 1), we instead use the one pertaining to the mid-Miocene (14 Myr ago). A larger part of West Antarctica (Marie Byrd Land and Palmer Land) has subsided below sea level in this reconstruction ~~, which~~ (Fig. S6a,b). This leads to a smaller simulated WAIS at low CO₂ levels in ~~this the TOPO 14Ma~~ case (Fig. 4). The bedrock subsidence in Victoria Land and the Amery embayment has a smaller effect on equilibrium ice volume. The total equilibrium ice volume above floatation is 44.6 m.s.l.e. at 280 ppm, compared to 48.4 m.s.l.e. in the REF case (-8%). The amplitude of transient ice volume variability is affected as well, ~~as it even on relatively short orbital (obliquity) timescales. It~~ decreases to 24.4 m.s.l.e. (-10% w.r.t. REF) when 40-kyr forcing cycles are imposed (Fig. 5a,b).

The effect of long-term Antarctic landscape evolution on ice volume becomes more apparent when we compare simulations using bedrock conditions of the Eocene-Oligocene boundary (34 Myr ago; TOPO 34Ma) and relaxed PD topography (TOPO PD) ~~-(Fig. S6c,d)~~. In the latter case, less ice builds up in Victoria Land and Coats Land where the initial bedrock topography has subsided below sea level, compared to the TOPO 14Ma case (Fig. S3aS7a). The equilibrium ice volume at 280 ppm reduces to 42.8 m.s.l.e. (-12% w.r.t. REF), and the amplitude of transient 40-kyr ice volume variability to 23.3 m.s.l.e. (-14% w.r.t. REF; Fig. ??a,bS8a,b dotted lines). Conversely, it increases to 28.4 m.s.l.e. (+5% w.r.t. REF; Fig. S7b), and is centered around a generally larger ice volume, in experiment TOPO 34Ma (Fig. ??a,bS8a,b, long-dashed lines). This is also due to the fact that the EAIS is more stable, as shown by larger ice volumes already at high CO₂ levels in the CO₂-V_{eq} relation, particularly in the upper, descending branch. Hysteresis is indeed larger than in the REF case, because the positive surface-height-mass-balance feedback is stronger due to the build-up of thicker ice. At 280 ppm, the equilibrium ice volume is 57.4 m.s.l.e. in this experiment (+19% w.r.t. REF). A much thicker ice sheet is built up in Coats Land, the Amery embayment, and Marie Byrd Land (Fig. S3bS7b). Still in this case, the ice sheet does not significantly advance into the Ross and Filchner-Ronne embayments. This is different when we use the ~~older-earlier~~ geological reconstructions of Antarctic paleotopography at the Eocene-Oligocene boundary of Wilson et al. (2012) ~~-(Fig. S6e,f)~~. At 280 ppm, the ice then grounds in the Ross embayment (TOPO Wilson_mean, using the mean of the minimum and maximum reconstructions; Fig. ~~(Fig. S3e))~~S7c), or in both embayments (TOPO Wilson_max, using the maximum reconstruction; Fig. S3dS7d). Due to the higher CO₂ levels at which deglaciation takes place, a relatively large ice volume persists throughout 40-kyr transient simulations (Fig. ??eS8c,d). In contrast to the other experiments, the initial conditions of these transient simulations now affect the ice volume during the equilibrium cycle. The ice sheet remains generally larger when the equilibrium 280-ppm simulation, rather than the 840-ppm simulation, is used as starting point (not shown).

3.4 Influence of ice shelf formation Sensitivity to ocean forcing

We perform and compare two additional sensitivity experiments that are end-members with respect to basal melt rate settings, to assess the influence of ice shelf formation on AIS variability access the sensitivity of our results to the parameterisation of ocean forcing. In experiment BMB LGM (Table 1), we use Last Glacial Maximum (LGM)-like basal melt parameter settings in Eq. 1: $M_{expo} = 0$ m/yr, $M_{deep} = 2$ m/yr, and $T_w = -5^\circ\text{C}$ (De Boer et al., 2013). The mild basal melt rates in this case create an environment in which ice shelves can grow relatively easily. At the other extreme, in experiment BMB no_shelves, ice shelves are effectively inhibited from growing by applying a constant melt rate of 400 m/yr. The effect of ice shelf formation becomes notable at from 504 ppm in the equilibrium simulations (~~lower, ascending branch~~) (Fig. 6a,b, solid and dashed blue lines). Ice ~~shelf shelves~~ form along the margins at Coates Land and Oates Land (Fig. 7a). While in Coates Land, this leads to a modest thickening of the grounded ice, this happens to a much lesser degree in Oates Land. In the long 400-kyr transient simulation, these ice shelves form after around 110 kyr, at ~ 530 ppm (Fig. 6b). At 392 ppm in the equilibrium simulations, and after around 170 kyr at ~ 360 ppm in the 400-kyr transient simulation, ice shelves appear approximately simultaneously along almost the entire remainder of the (East and West) Antarctic margin (Fig. 7b). The effect on grounded ice is particularly noticeable in West-Antarctica, and Wilkes Land, George V Land and MacRobertson Land, where the ice sheet thickens by kilometers. In the deglaciation phase these ice shelves remain influential until 728 ppm in the equilibrium simulations (~~upper, descending branch~~ Fig. 6b, solid and dashed blue lines), and until around 360 kyr at ~ 728 ppm in the 400-kyr transient simulation ~~-, (Fig. 6a,b, green and orange lines)~~. In the transient simulations, the amplitude of variability is increased. The hysteresis in the CO_2 -V relation ~~-, and the amplitude of variability are increased~~ is increased as well, although only slightly. In the 40-kyr transient simulations, the ice volume variability amounts to 30.7 m.s.l.e. in case LGM BMB and 25.7 m.s.l.e. in case LGM no_shelves, a 5 m.s.l.e. (19%) difference (Fig. 6c,d).

4 Discussion

The CO_2 levels at which ~~East-Antaretic~~ East Antarctic glaciation is simulated, are known to be dependent on the climate model used to construct the ~~climatic~~ climate forcing (Gasson et al., 2014). This could (partly) explain why in our study EAIS inception occurs at higher CO_2 levels than in Stap et al. (2019). They used climate forcing from COSMOS which has a higher climate sensitivity (reported as 4.1 K per CO_2 doubling) than GENESIS (2.9 K). Note that in neither study greenhouse gasses other than CO_2 are varied, probably biasing the CO_2 concentrations high. ~~In contrast, our climate input comes from the same model as Halberstadt et al. (2021), who obtain relatively large EAIS volumes at CO_2 levels up to 1140 ppm. In absolute sense, this CO_2 difference can simply be caused by the value of~~ Another factor that has a large influence on the glaciation threshold is the ablation factor that we use in our surface mass balance calculation (C_3 in Eq. 6), as demonstrated by our SMB sensitivity experiment (Fig. ~~??~~ S5). Arguably the best way to further constrain this parameter in the future is by performing a comprehensive comparison of our mass balance calculation, and other methods and observations, as was done previously for the Greenland ice sheet (Fettweis et al., 2020). ~~Furthermore, they additionally use a regional climate model for downscaling~~

and refrain from using bias corrections for the temperature and precipitation forcing. Nevertheless, these factors likely do not explain the narrower CO₂ range between inception of the East- and West-Antarctic ice sheet in our simulations compared to Halberstadt et al. (2021). At the root of this discrepancy can be the different schemes we use to calculate precipitation and ablation, which may be more sensitive to temperature changes. Differences in ice dynamics, e.g. in the treatment of ice flow enhancement and basal sliding, might cause further disparity. Probably for the same reasons, we obtain larger transient AIS volume variability than Stap et al. (2019), when using a similar glacial-index method (experiment FEEDB). Conducting an intermodel comparison of early AIS dynamics would be advisable for a more comprehensive investigation of the different results yielded by ice sheet models. NOFEEDB). We nonetheless corroborate earlier findings that disequilibrium between the AIS and the forcing climate causes smaller transient ice volume variability than steady-state simulations suggest (Stap et al., 2019), as well as out-of-phase changes between these quantities (Stap et al., 2020).

Here, we include ice-sheet-atmosphere feedbacks in our reference case. We have shown that these feedbacks tend to increase hysteresis in the CO₂-equilibrium ice volume relation, and decrease the amplitude of AIS variability under equal 40-kyr forcing CO₂ cycles by 21%. This implies that correspondingly, the (Fig. 3). Also on longer timescales AIS variability is suppressed by the ice-sheet-atmosphere feedbacks. This is partly due to the smaller equilibrium ice volumes at low CO₂ levels, and partly to decreased growth rates of the transient ice volume relative to the change of equilibrium ice volume. In short, the ice-sheet-atmosphere feedbacks cause a slower build-up of the AIS to smaller peak ice volumes. Correspondingly, the contribution of AIS volume changes to benthic $\delta^{18}\text{O}$ fluctuations gets smaller, although the. This calls for a larger contribution of deep-sea temperatures to the amplitude of Miocene benthic $\delta^{18}\text{O}$ fluctuations. The actual strength of the impact will, however, depend on the isotopic composition of Antarctic snow as well (Gasson et al., 2016; Rohling et al., 2021).

Furthermore, we use recent reconstructions of Antarctic bedrock topography during the Miocene (Hochmuth et al., 2020a; Paxman et al., 2019), rather than the present day as in Stap et al. (2019), as a boundary condition in our AIS simulations. In agreement with Colleoni et al. (2018) and Paxman et al. (2020), we find that the WAIS becomes more vulnerable over the early- and mid-Miocene (Fig. 5). This is demonstrated by the lower equilibrium ice volumes at equal CO₂ levels in our experiment TOPO 14Ma, which has a bedrock topography pertaining to the mid-Miocene (14 Myr ago), compared to REF, in which the topography is from the Mioene-Oligocene Oligocene-Miocene boundary (23-24 Myr ago). We also corroborate the finding of Paxman et al. (2020), that from 34 to 24 million years ago bedrock changes primarily concern the EAIS(Fig. ??), while from 24 Myr ago onwards mostly the WAIS is affected. Here, we have additionally examined the effect of the evolving Antarctic landscape on transient Miocene AIS variability, which is 10% smaller when subjected to 40-kyr forcing CO₂ cycles in TOPO 14Ma than in REF, due to impeded WAIS growth. Hence, CO₂ minima must have dropped increasingly further over the course of the early- and mid-Miocene to obtain AIS variability with similar amplitude, and consequently a similar contribution of the AIS to Miocene benthic $\delta^{18}\text{O}$ variability.

410 We simulate the inception of ice shelves only at relatively low CO_2 levels. Ice shelves first appear along the coasts of Oates Land and Coates Land when the CO_2 concentration drops below ~ 530 ppm, but the effect of ice shelves on grounded ice volume only becomes significant below ~ 360 ppm CO_2 when the EAIS - with a volume of almost 30 m.s.l.e. - is already well-established. This transition from a land-based to a marine ice sheet at CO_2 levels around 400 ppm is in general agreement with other model results (Halberstadt et al., 2021), and geological data inferences (Naish et al., 2009; Levy et al., 2016, 2019).

415 Below ~ 360 ppm CO_2 , ice shelves fringe almost the entire continent in our simulations. This includes the Wilkes Subglacial Basin and the Ross Sea sector, where records from IODP Site U1356 and ANDRILL AND-2A show evidence of periodic ice retreat and readvance (Hauptvogel and Passchier, 2012; Levy et al., 2016; Pérez et al., 2021; Sangiorgi et al., 2018). In this study, we focus on the variability of a generally small AIS, which is located primarily on the East side of the continent, while the WAIS size remains relatively limited. In the Ross and Filchner-Ronne embayments, however, growth of extended ice shelves never occurs in our Miocene simulations, not even when the basal melt rates are set to LGM values. On the one hand, it is certainly possible that our basal melt parameterisation is not suited for the simulation of (large-scale) Antarctic ice shelf inception. On the other hand, the fact that the Ross and Filchner-Ronne ice shelf shelves do grow when we impose steady present-day climatic-climate forcing, points out that West-Antarctic ice shelves have to be sustained by a sufficiently large ice flow coming from the WAIS. That means that in addition to the oceanic conditions, suitable atmospheric conditions are

420 very important for ice shelf formation as well. Possibly Therefore, WAIS growth is therefore hampered by the air temperatures remaining too warm in our forcing climate simulations, because in the cold case (simulation 1fumebi) there is only a small WAIS present (Fig. S2) in the forcing climate simulation. In the future, the forcing steady-state climate simulations used in the matrix interpolation could-should preferably be constructed by a coupled set-up of the same ice-sheet and climate models that are deployed in the transient simulations. This would ensure congruence between the results of the equilibrium and transient

425 simulations, so that the transient ice sheet will not grow beyond its largest size in the forcing equilibrium results.

Although the AIS remains mostly grounded, the formation of floating ice shelves does have an impact on our result. This is demonstrated by comparing simulations in which LGM basal melt rates are applied, to simulations in which ice shelf growth is prevented (Figs. 6,7). The 40-kyr transient AIS variability is amplified by 19% in case of LGM basal melt rates - and 5% in the reference case - compared to the case without ice shelves. In our simulations, ice shelves first appear along the coasts of Oates Land and Coates Land when the CO_2 concentration drops below ~ 530 ppm ($T_w = 2^\circ\text{C}$), but the effect of ice shelves on grounded ice volume only becomes significant below ~ 360 ppm CO_2 ($T_w = 0.8^\circ\text{C}$) when the EAIS - with a volume of almost 30 m.s.l.e. - is already well-established. This transition, from a land-based ice sheet to one that is in contact with the ocean, at CO_2 levels around 400 ppm is in general agreement with other model results (Halberstadt et al., 2021), and geological data inferences (Naish et al., 2009; Levy et al., 2016, 2019). Below ~ 360 ppm CO_2 , ice shelves fringe almost the entire continent in our simulations. This includes the Wilkes Subglacial Basin and the Ross Sea sector, where records from IODP Site U1356 and ANDRILL AND-2A show evidence of periodic ice retreat and readvance (Hauptvogel and Passchier, 2012; Levy et al., 2016; Pérez et al., 2021; Sangiorgi et al., 2018). When we increase CO_2 levels from cold conditions in our simulations, the ice shelves keep having an effect on grounded AIS ice volume until relatively high

435

440

445 CO₂ levels of ~ 728 ppm ($T_w = 2^\circ\text{C}$) are reached. Only then does the AIS retreat into the Wilkes Subglacial Basin, suggesting that in our model set-up periodic AIS waxing and waning in this region has to be governed by substantial CO₂ fluctuations. Growth of ice shelves hence increases hysteresis in the CO₂-ice volume relation, in general agreement with the findings of Garbe et al. (2020). In line with this result, we find a different effect on final ice volume (above floatation) in steady-state PD simulations with a 400-m/yr melt rate, depending on whether the simulation is started with or without ice. Starting from an

450 isostatically rebounded PD topography without ice, we obtain a volume of 39.5 m.s.l.e., 28% smaller than the PD run with normal basal melt rates (Fig. S4aS9a,c). When the simulation is started from a PD topography including grounded and floating ice, the final volume is 48.4 m.s.l.e., which is only 14% smaller than the normal PD run (Fig. S4bS9b,d). Hence, in our model, the effect of ice shelves on grounded ice volume is smaller when existing ice shelves are melted away, than when ice shelves cannot form in the first place. Despite having a longer model runtime and different surface mass balance forcing, these

455 simulations are similar to the ABUMIP ABUM experiment (Sun et al., 2020). Our simulated 8.2 m.s.l.e. loss of ice volume above floatation after ice sheet removal under PD forcing conditions (starting with ice) is on the large side of the wide range of model responses (~ 1 to 11 m.s.l.e. loss) reported there. Showing complete collapse of the WAIS and strong retreat in the Amery embayment, but limited ice loss in the Recovery, Wilkes and Aurora subglacial basins, our spatial response is most similar to that of ABUMIP models PISM1, PSU3D1, and BISICLES. However, our simulated response, using a Coulomb

460 sliding law ($q = 0.3$), is weaker than that of ELMER/ICE and SICOPOLIS, that used a Weertman sliding law ($m = 3$), and that of a different version of IMAU-ICE that used the Schoof flux boundary condition (Schoof, 2007) at the grounding line. A stronger response of grounded ice volume to ocean warming and ice shelf break-up might lead to earlier retreat, i.e. at a lower CO₂ level, of the AIS in e.g. the Wilkes Subglacial Basin, hence reducing hysteresis in the CO₂-ice volume relation.

465 ~~Here, we have conducted idealised transient simulations, addressing the effects of the ocean, atmosphere and solid earth on~~
~~Miocene AIS evolution. Moving forward towards realistic transient simulations of AIS evolution, our results of experiment~~
~~INSOL indicate that orbital forcing variability will need to be accounted for in the model set-up (Fig. ??), for instance~~
~~by integrating it in the matrix interpolation method (e.g. Ladant et al., 2014; Tan et al., 2018). A further improvement to be~~
~~made is the implementation of a more advanced scheme to calculate the basal mass balance underneath the ice shelves;~~
~~several alternatives have been developed in recent years (e.g. Reese et al., 2018a; Lazeroms et al., 2019; Pelle et al., 2019).~~
~~This should be accompanied by ocean forcings supplied by sophisticated (3D) ocean models. As a final prospect, inclusion~~
~~of a coupled sediment model will ultimately open up the possibility of quantifying the transient effect of changing bedrock~~
~~topography on AIS dynamics over the Miocene (Pollard and DeConto, 2003, 2020). Our climate input comes from the same~~
~~model (GENESIS) as Halberstadt et al. (2021), who obtain relatively large EAIS volumes at CO₂ levels up to 1140 ppm.~~

475 Differences between their results and ours can be explained by the different ice-sheet models, and surface mass balance
schemes used. We use an insolation temperature melt (ITM) scheme (Bintanja et al., 2002), accounting for the direct effect
of insolation on surface melt, whereas they deploy a positive degree day (PDD) scheme. Additionally, they use a regional
climate model for downscaling the forcing, which allows for capturing orographic precipitation in more detail. We use a bias
correction to obtain a more detailed reference climate. This may, however, lead to overestimated temperatures, because we

480 use a present-day (ERA-40) base climate, which is slightly warmer than our PI reference (Stap et al., 2019). In model tests, we simulate higher AIS ice volumes at high CO₂ levels when we refrain from using this bias correction, although this is dependent on the ablation factor (c_3 in Eq. 6; not shown). All in all, even with the bias correction included, we need very large CO₂ variations on relatively short orbital timescales (40 kyr) to obtain substantial East Antarctic ice sheet variability. This could point to a relatively stable EAIS, and a consequent reduced contribution to benthic $\delta^{18}\text{O}$ fluctuations during the Miocene (Bijl et al., 2018; Hartman et al., 2018). Alternatively, the WAIS could play a more major role in establishing ice-sheet variability. As explained above, WAIS growth is hampered in our simulations by the climate forcing. However, recently geological evidence was found for a dynamic Miocene WAIS in the Ross Sea section (Marschalek et al., 2021), in agreement with the modelling results of e.g. Gasson et al. (2016) and Halberstadt et al. (2021). The generally low benthic $\delta^{18}\text{O}$ values during the Miocene in that case would require persistent high ocean temperatures, which were indeed recently found by Modestou et al. (2020). Further modelling efforts are still needed to reconcile these warm ocean temperatures with the existence of a large marine AIS.

5 Conclusions

Using the ice-sheet model IMAU-ICE, we have performed steady-state and transient simulations of the Antarctic ice sheet (AIS) during the early- and mid-Miocene (23 to 14 Myr ago). Climate forcing was obtained from Miocene simulations by GENESIS (Burls et al., 2021), with different CO₂ levels and ice sheet configurations, using a recently developed matrix interpolation routine (Berends et al., 2018). ~~We find competing influences of the atmosphere, ocean and solid earth on transient Miocene AIS variability.~~

Compared to a simpler ~~glacial~~ index method, as used e.g. in Stap et al. (2019), our matrix interpolation method incorporates two important ice-sheet-atmosphere feedbacks: the positive albedo-temperature feedback, and ~~the a~~ negative ice-volume-precipitation feedback. The former tends to increase hysteresis in the CO₂-~~V~~ ice volume above floatation (V) relation and decrease the amplitude of transient AIS variability, ~~while the latter~~. ~~The negative ice-volume-precipitation feedback, which represents the ice-sheet desertification effect,~~ has the opposite effect (Fig. 43). The albedo-temperature feedback is ~~the strongest of the two, as in~~ dominant in our simulations. In the 40-kyr transient reference simulation, the ice volume variability amplitude is 21% smaller than when ~~a glacial-an~~ index method is used. ~~Hence, the net effect of accounting for ice-sheet-atmosphere interactions in the interpolation of climate forcing reduces simulated transient Miocene Antarctic ice sheet variability.~~

Differences between the geological reconstructions of Antarctic bedrock topography at the Oligocene-Miocene boundary (23-24 Myr ago) and the mid-Miocene (14 Myr ago) are mainly located in ~~West-Antartica~~ West-Antarctica. The subsidence of land below sea level during the early- and mid-Miocene impedes WAIS build-up. Consequently, it diminishes hysteresis in the CO₂-~~ice volume above floatation (V)~~-V relation and reduces 40-kyr AIS variability by 10% in ~~strength~~ amplitude (Figs. 4, 5). Hence, ~~under equal atmospheric and oceanic forcing conditions, the amplitude of transient AIS variability becomes~~

~~progressively smaller during CO₂ minima must have dropped increasingly further over the course of the early- and mid-Miocene, due to glacial erosion, to obtain AIS variability with similar amplitude.~~

515

~~The importance of ice shelf formation is demonstrated by comparing simulations in which LGM basal melt rates are applied, to simulations in which ice shelf growth is prevented (Figs. 6, 7). Ice shelves start to form around CO₂ levels of ~530 ppm at the margins of Coates Land and Oates Land and at ~360 ppm approximately simultaneously around all other margins. They continue to affect grounded ice volume until the forcing. Here, we have conducted idealised transient simulations, addressing the effect of CO₂ concentration is increased to ~728 ppm in our 400-kyr transient simulation. The formation of ice shelves leads to a thickening of the hinterlying grounded ice, particularly of the WAIS, and consequently increases hysteresis in the CO₂-V relation. 40-kyr transient AIS variability is amplified by 19% in case of LGM basal melt rates – and 5% in the reference case – compared to variability on Miocene AIS evolution. Moving forward towards realistic transient simulations of AIS evolution, orbital forcing variability will need to be accounted for in the model set-up, for instance by integrating it in the matrix interpolation method (e.g. Ladant et al., 2014; Tan et al., 2018). A further improvement to be made is the ease without ice shelves, implementation of a more advanced scheme to calculate the basal mass balance underneath the ice shelves; several alternatives have been developed in recent years (e.g. Reese et al., 2018a; Lazeroms et al., 2019; Pelle et al., 2019). This should be accompanied by ocean forcing supplied by sophisticated (3D) ocean models. As a final prospect, inclusion of a coupled sediment model will ultimately open up the possibility of quantifying the transient effect of changing bedrock topography on AIS dynamics over the Miocene (Pollard and DeConto, 2003, 2020).~~

520
525
530

~~In our simulations, the ice-sheet-atmosphere feedbacks and the evolution of bedrock topography, that is due mostly to glacial erosion, hence tend to reduce the contribution of AIS changes to orbital timescale benthic $\delta^{18}\text{O}$ variability during the Miocene. Conversely, the formation of ice shelves eases the gaining of ice volume, and increases this contribution. Realistic simulations of Miocene AIS evolution are left to future research.~~

535 *Code availability.* The code for IMAU-ICE v1.1.1-MIO is available from <https://github.com/IMAU-paleo/IMAU-ICE/releases/tag/v1.1.1-MIO>.

Data availability. All relevant data related to this article will be made openly accessible from the PANGAEA database (<https://pangaea.de/>).

Appendix A: Ice-sheet model settings

In IMAU-ICE, Glenn’s flow law exponent is set to $n = 3$, and the SIA- and SSA-based velocities are enhanced by factors 5 and 1 respectively. The basal stress is calculated using a Coulomb power law model for pseudo-plastic till (exponent $q = 0.3$ and threshold velocity $u_{threshold} = 100 \text{ m/s}$) (Bueler and Brown, 2009). The till friction angle, which determines the

540

yield stress, varies linearly between 5° for bedrock elevations of -1000 m and lower, and 20° for those of 0 m and above. The geothermal heat flux is taken from Shapiro and Ritzwoller (2004). A basic elastic lithosphere-relaxing asthenosphere (ELRA) model (Lingle and Clark, 1985; Le Meur and Huybrechts, 1996) is used to calculate bedrock adjustment, using a mantle density of 3300 kg m⁻³, a lithospheric flexural rigidity of 1.0 x 10²⁵ kg m² s⁻², and a relaxation time of 3000 yr. The sea level is kept constant at present-day level (0 m) ~~because local Antarctic sea level changes are very uncertain during the Miocene as they are mainly due to Antarctic ice sheet changes. Local sea level changes can differ substantially from the global mean (e.g. Stocchi et al., 2013), and their calculation would require solving the sea-level equation which is not yet available in IMAU-ICE. We have performed three tests, applying a constant 7 m higher sea level (mimicking the absence of the Greenland ice sheet), as well as applying internally calculated global mean, and minus the global mean, sea level changes. These tests have shown negligible differences in comparison to our reference results (not shown).~~

Appendix B: Climate matrix forcing

Using a so-called matrix method (Pollard, 2010; Pollard et al., 2013), a time-continuous climate forcing for the ice-sheet model can be calculated. This is achieved by interpolating between the GENESIS time-slice climate simulations every 10 model years, based on the varying CO₂ levels and ice coverage. The matrix interpolation routine used in this study is described in detail in Berends et al. (2018), and was previously deployed to simulate the evolution of the North-American, Eurasian, Greenland and Antarctic ice sheets during the past 3.6 Myr (Berends et al., 2019, 2021).

B1 Surface air temperature

For the applied surface air temperature, first ~~two weighing factors~~ we calculate a scalar weighing factor for CO₂ (w_{CO_2}) and a spatially varying one for ice-sheet coverage (~~w_{ice}~~ are calculated $w_{ice}(x, y)$).

$$w_{CO_2} = \frac{CO_2 - CO_{2,cold}}{CO_{2,warm} - CO_{2,cold}}, \quad (B1)$$

where CO_2 is the applied CO₂ level, and $CO_{2,cold}$ and $CO_{2,warm}$ the low (280 ppm) and high (840 ppm) CO₂ levels applied in the climate forcing simulations. Weight factor ~~w_{ice}~~ $w_{ice}(x, y)$ is based on scaling between local absorbed insolation of the forcing climates, accounting for the effect of surface whitening (darkening) by ice sheet advance (retreat) on temperature. In order to ~~calculate w_{ice}~~ obtain $w_{ice}(x, y)$, first the absorbed insolation I_{abs} $I_{abs}(x, y)$ is calculated from the incoming insolation (Q_{TOA} $Q_{TOA}(x, y)$) and internally calculated surface albedo (~~α~~ $\alpha(x, y)$). This is done every 10-yr coupling time step for the modeled field ($I_{abs,mod}$), ~~and the fields interpolated between~~. For the cold (1fumebi) and warm (3nomebi) GCM snapshot fields ($I_{abs,cold}$, $I_{abs,warm}$) ~~this is done at the start of each simulation by running the SMB model (Sect. 2.1.2), without ice dynamics, for 10 yrs to determine $\alpha(x, y)$.~~

$$I_{abs}(x, y) = (1 - \alpha(x, y)) * Q_{TOA}(x, y). \quad (B2)$$

Incoming insolation $Q_{TOA}(x, y)$ (Laskar et al., 2004) is kept constant at the present-day values, so that differences between the cold, warm and modeled $I_{abs}(x, y)$ fields are solely determined by the albedo. Then, the weighing field for insolation $w_{ins}(x, y)$ is calculated:

$$w_{ins}(x, y) = \frac{I_{abs, mod} - I_{abs, cold}}{I_{abs, warm} - I_{abs, cold}} \frac{I_{abs, mod}(x, y) - I_{abs, cold}(x, y)}{I_{abs, warm}(x, y) - I_{abs, cold}(x, y)} \quad (B3)$$

Weighing factors w_{CO_2} and $w_{ins}(x, y)$ are clipped at -0.25 and 1.25, to avoid spurious extreme temperatures and precipitation. Finally, the weighing field for ice-sheet coverage is calculated from the smoothed (using a Gaussian-Shepard smoothing filter with a radius of 200 km) and average (over the ice-sheet model $griddomain$) w_{ins} , to account for local as well as regional effects:

$$w_{ice}(x, y) = \frac{1}{7} w_{ins, smoothed}(x, y) + \frac{6}{7} w_{ins, avg}. \quad (B4)$$

The viability of this partitioning $w_{ins, smoothed}(x, y)$ and $w_{ins, avg}$ was demonstrated by Berends et al. (2018), who simulated ice-sheet evolution over the last glacial cycle, showing that model results agree well with available data in terms of ice-sheet extent, sea-level contribution, ice-sheet surface temperature and contribution to benthic $\delta^{18}O$. For each grid cell in the domain the final weighing factor $w_{tot}(x, y)$ is calculated as:

$$w_{tot}(x, y) = \epsilon * w_{CO_2} + (1 - \epsilon) * w_{ice}(x, y). \quad (B5)$$

ϵ_{CO_2} is the ratio between the effects of CO_2 and ice-sheet coverage, predetermined (in our reference simulation) on the basis of. To determine ϵ for our simulations, we use GENESIS simulations 3fumebi (840 ppm CO_2 , full-large ice sheet) and 1nomebi (280 ppm CO_2 , no ice sheet). These climate simulations isolate the separate effects of CO_2 and the ice sheet respectively. The (spatially averaged) relative influence of CO_2 on temperature is estimated as the isolated effect of CO_2 divided by the combined influence of CO_2 and ice-sheet coverage.

$$\epsilon_{CO_2, ref} = \frac{\bar{T}_{3fumebi} - \bar{T}_{1fumebi}}{(\bar{T}_{3fumebi} - \bar{T}_{1fumebi}) + (\bar{T}_{1nomebi} - \bar{T}_{1fumebi})} \approx \frac{\bar{T}_{3fumebi} - \bar{T}_{1fumebi}}{\bar{T}_{3nomebi} - \bar{T}_{1fumebi}} = 0.62, \quad (B6)$$

where \bar{T} is the average temperature over the Antarctic domain. In the reference case $\epsilon_{CO_2} = 0.62$. The weighing factors are clipped at -0.25 and 1.25, to avoid spurious extreme temperatures and precipitation. Synergy between the effects of CO_2 and ice-sheet coverage is relatively small, so that $\epsilon \approx (\bar{T}_{3fumebi} - \bar{T}_{1fumebi}) / (\bar{T}_{3nomebi} - \bar{T}_{1fumebi})$.

Next, the reference temperature ($T_{ref}(x, y)$) and surface height ($h_{ref}(x, y)$) are obtained by interpolating between the cold ($T_{cold}, h_{cold}(x, y)$) and warm ($T_{warm}, h_{warm}(x, y)$) forcing climate fields:

$$T_{ref}(x, y) = w_{tot}(x, y) * T_{warm}(x, y) + (1 - w_{tot}(x, y)) * T_{cold}(x, y) \quad (B7)$$

and likewise

$$h_{ref}(x, y) = w_{tot}(x, y) * h_{warm}(x, y) + (1 - w_{tot}(x, y)) * h_{cold}(x, y). \quad (B8)$$

The applied 2-m air temperature field $T_{applied}(x, y)$ is obtained by first correcting for the height differences between the modeled ($h_{mod}(x, y)$) and reference ($h_{ref}(x, y)$) surface height using a uniform lapse rate $\lambda = 8$ K/km. Finally, a spatially-varying bias correction based on the difference between the modeled preindustrial temperature (T_{PI}) and the 1957-2002 ERA40 reanalysis data (Uppala et al., 2005) (T_{ERA40}) is applied:

$$T_{applied}(x, y) = T_{ref}(x, y) - \lambda * (h_{mod}(x, y) - h_{ref}(x, y)) + (T_{ERA40}(x, y) - T_{PI}(x, y)). \quad (B9)$$

B2 Precipitation

Precipitation is weighed based on the total ice volumes modeled (V_{mod}) and in the cold (V_{cold}) and warm (V_{warm}) GENESIS forcing climates (V_{cold} , V_{warm}):

$$w_{tot} = 1 - \frac{V_{mod} - V_{warm}}{V_{cold} - V_{warm}}. \quad (B10)$$

A reference precipitation field is obtained, by interpolating between the warm-cold (P_{warm}) and cold-warm (P_{cold}) states: forcing fields:

$$P_{ref}(x, y) = \exp \left[w_{tot} * \frac{\log[P_{warm}]}{P_{warm}} \log[P_{warm}(x, y)] + (1 - w_{tot}) * \frac{\log[P_{cold}]}{P_{cold}} \log[P_{cold}(x, y)] \right]. \quad (B11)$$

Similarly as for the temperature, a bias correction is applied:

$$P_{ref,corr}(x, y) = P_{ref}(x, y) * \frac{\frac{P_{ERA40} + 0.008}{P_{PI} + 0.008} \frac{P_{ERA40}(x, y) + 0.008}{P_{PI}(x, y) + 0.008}}{1}. \quad (B12)$$

The term 0.008 is added to avoid problems caused by division by very small numbers. And finally, the precipitation field is downscaled to the finer ice-sheet model grid, using a Clausius-Clapeyron relation (Lorius et al., 1985; Huybrechts, 2002; De Boer et al., 2013):

$$P_{applied}(x, y) = P_{ref,corr}(x, y) * \left(\frac{T_{I,ref}}{T_I} \frac{T_{I,ref}(x, y)}{T_I(x, y)} \right)^2 * \exp \left[22.47 * \left(\frac{273.16K}{T_{I,ref}} \frac{273.16K}{T_{I,ref}(x, y)} - \frac{273.16K}{T_I} \frac{273.16K}{T_I(x, y)} \right) \right] \quad (B13)$$

with

$$T_I(x, y) = 88.9 + 0.67 * T_{ref}(x, y), \quad (B14)$$

$$T_{I,ref}(x, y) = 88.9 + 0.67 * (T_{ref}(x, y) - \lambda * (h_{mod}(x, y) - h_{ref}(x, y))). \quad (B15)$$

Author contributions. LBS designed the research and performed the experiments, with technical assistance from CJB. LBS, CJB, MDWS
625 and RSWvdW analysed the results. EGWG provided the GENESIS results used as climate forcing in this study. LBS drafted the paper, with
input from all co-authors.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. L.B. Stap is funded by the Dutch Research Council (NWO), through VENI grant VI.Veni.202.031. C.J. Berends is
supported by PROTECT, which has received funding from the European Union's Horizon 2020 research and innovation programme under
630 grant agreement no. 869304. We thank our colleague Peter Bijl for commenting on an earlier draft of this article. We further thank the
researchers that reconstructed the Antarctic paleotopographies and paleobathymetries for making them publicly available, and in particular
Isabel Sauermilch for providing them to us. Last but not least, we would like to thank two anonymous reviewers, whose comments have
helped to improve the quality of this article.

References

- 635 Abe-Ouchi, A., Saito, F., Kawamura, K., Raymo, M. E., Okuno, J., Takahashi, K., and Blatter, H.: Insolation-driven 100,000-year glacial cycles and hysteresis of ice-sheet volume, *Nature*, 500, 190–193, <https://doi.org/10.1038/nature12374>, 2013.
- Badger, M. P. S., Lear, C. H., Pancost, R. D., Foster, G. L., Bailey, T. R., Leng, M. J., and Abels, H. A.: CO₂ drawdown following the middle Miocene expansion of the Antarctic Ice Sheet, *Paleoceanography*, 28, 42–53, <https://doi.org/10.1002/palo.20015>, 2013.
- Beckmann, A. and Goosse, H.: A parameterization of ice shelf–ocean interaction for climate models, *Ocean Modelling*, 5, 157–170, [https://doi.org/10.1016/S1463-5003\(02\)00019-7](https://doi.org/10.1016/S1463-5003(02)00019-7), 2003.
- 640 Berends, C. J., de Boer, B., and Van de Wal, R. S. W.: Application of HadCM3@Bristolv1.0 simulations of paleoclimate as forcing for an ice-sheet model, ANICE2.1: set-up and benchmark experiments, *Geoscientific Model Development*, 11, 4657–4675, <https://doi.org/10.5194/gmd-11-4657-2018>, 2018.
- Berends, C. J., de Boer, B., Dolan, A. M., Hill, D. J., and Van de Wal, R. S. W.: Modelling ice sheet evolution and atmospheric CO₂ during the Late Pliocene, *Climate of the Past*, 15, 1603–1619, <https://doi.org/10.5194/cp-15-1603-2019>, 2019.
- 645 Berends, C. J., de Boer, B., and Van de Wal, R. S. W.: Reconstructing the evolution of ice sheets, sea level, and atmospheric CO₂ during the past 3.6 million years, *Climate of the Past*, 17, 361–377, <https://doi.org/10.5194/cp-17-361-2021>, 2021.
- Bijl, P. K., Houben, A. J. P., Hartman, J. D., Pross, J., Salabarnada, A., Escutia, C., and Sangiorgi, F.: Paleoceanography and ice sheet variability offshore Wilkes Land, Antarctica–Part 2: Insights from Oligocene–Miocene dinoflagellate cyst assemblages, *Climate of the*
- 650 *Past*, 14, 1015–1033, <https://doi.org/10.5194/cp-14-1015-2018>, 2018.
- Bintanja, R., Van de Wal, R. S. W., and Oerlemans, J.: Global ice volume variations through the last glacial cycle simulated by a 3-D ice-dynamical model, *Quaternary International*, 95, 11–23, [https://doi.org/10.1016/S1040-6182\(02\)00023-X](https://doi.org/10.1016/S1040-6182(02)00023-X), 2002.
- Bradshaw, C. D., Langebroek, P. M., Lear, C. H., Lunt, D. J., Coxall, H. K., Sosdian, S. M., and de Boer, A. M.: Hydrological impact of Middle Miocene Antarctic ice-free areas coupled to deep ocean temperatures, *Nature Geoscience*, 14, 429–436, [https://doi.org/10.1038/s41561-](https://doi.org/10.1038/s41561-021-00745-wT)
- 655 [021-00745-wT](https://doi.org/10.1038/s41561-021-00745-wT), 2021.
- Bueler, E. and Brown, J.: Shallow shelf approximation as a “sliding law” in a thermomechanically coupled ice sheet model, *Journal of Geophysical Research: Earth Surface*, 114, <https://doi.org/10.1029/2008JF001179>, 2009.
- Burls, N. J., Bradshaw, C. D., De Boer, A. M., Herold, N., Huber, M., Pound, M., Donnadieu, Y., Farnsworth, A., Frigola, A., Gasson, E., et al.: Simulating Miocene warmth: insights from an opportunistic multi-model ensemble (MioMIP1), *Paleoceanography and Paleoclimatology*, p. e2020PA004054, <https://doi.org/10.1029/2020PA004054>, 2021.
- 660 Colleoni, F., De Santis, L., Montoli, E., Olivo, E., Sorlien, C. C., Bart, P. J., Gasson, E. G. W., Bergamasco, A., Sauli, C., Wardell, N., et al.: Past continental shelf evolution increased Antarctic ice sheet sensitivity to climatic conditions, *Scientific Reports*, 8, 1–12, <https://doi.org/10.1038/s41598-018-29718-7>, 2018.
- De Boer, B., Van de Wal, R. S. W., Bintanja, R., Lourens, L. J., and Tuentner, E.: Cenozoic global ice-volume and temperature simulations with 1-D ice-sheet models forced by benthic $\delta^{18}\text{O}$ records, *Annals of Glaciology*, 51, 23–33, 2010.
- 665 De Boer, B., Van de Wal, R. S. W., Lourens, L. J., Bintanja, R., and Reerink, T. J.: A continuous simulation of global ice volume over the past 1 million years with 3-D ice-sheet models, *Climate Dynamics*, 41, 1365–1384, <https://doi.org/10.1007/s00382-012-1562-2>, 2013.
- De Boer, B., Lourens, L. J., and Van De Wal, R. S. W.: Persistent 400,000-year variability of Antarctic ice volume and the carbon cycle is revealed throughout the Plio-Pleistocene, *Nature Communications*, 5, 1–8, <https://doi.org/10.1038/ncomms3999>, 2014.

- 670 De Boer, B., Dolan, A. M., Bernales, J., Gasson, E., Goelzer, H., Golledge, N. R., Sutter, J., Huybrechts, P., Lohmann, G., Rogozhina, I., Abe-Ouchi, A., Saito, F., and van de Wal, R. S. W.: Simulating the Antarctic ice sheet in the late-Pliocene warm period: PLISMIP-ANT, an ice-sheet model intercomparison project, *The Cryosphere*, 9, 881–903, <https://doi.org/10.5194/tc-9-881-2015>, 2015.
- DeConto, R. M. and Pollard, D.: Rapid Cenozoic glaciation of Antarctica induced by declining atmospheric CO₂, *Nature*, 421, 245–249, <https://doi.org/10.1038/nature01290>, 2003.
- 675 DeConto, R. M., Pollard, D., and Kowalewski, D.: Modeling Antarctic ice sheet and climate variations during Marine Isotope Stage 31, *Global and Planetary Change*, 88–89, 45–52, <https://doi.org/10.1016/j.gloplacha.2012.03.003>, 2012.
- Fettweis, X., Hofer, S., Krebs-Kanzow, U., Amory, C., Aoki, T., Berends, C. J., Born, A., Box, J. E., Delhasse, A., Fujita, K., Gierz, P., Goelzer, H., Hanna, E., Hashimoto, A., Huybrechts, P., Kapsch, M.-L., King, M. D., Kittel, C., Lang, C., Langen, P. L., Lenaerts, J. T. M., Liston, G. E., Lohmann, G., Mernild, S. H., Mikolajewicz, U., Modali, K., Mottram, R. H., Niwano, M., Noël, B., Ryan, J. C., Smith, A., Streffing, J., Tedesco, M., van de Berg, W. J., van den Broeke, M., van de Wal, R. S. W., van Kampenhout, L., Wilton, D., Wouters, B., Ziemann, F., and Zolles, T.: GrSMBMP: intercomparison of the modelled 1980–2012 surface mass balance over the Greenland Ice Sheet, *The Cryosphere*, 14, 3935–3958, <https://doi.org/10.5194/tc-14-3935-2020>, 2020.
- 680 Fortuin, J. P. F. and Oerlemans, J.: Parameterization of the annual surface temperature and mass balance of Antarctica, *Annals of Glaciology*, 14, 78–84, <https://doi.org/10.3189/S0260305500008302>, 1990.
- 685 Foster, G. L., Lear, C. H., and Rae, J. W. B.: The evolution of pCO₂, ice volume and climate during the middle Miocene, *Earth and Planetary Science Letters*, 341, 243–254, <https://doi.org/10.1016/j.epsl.2012.06.007>, 2012.
- Garbe, J., Albrecht, T., Levermann, A., Donges, J. F., and Winkelmann, R.: The hysteresis of the Antarctic ice sheet, *Nature*, 585, 538–544, <https://doi.org/10.1038/s41586-020-2727-5>, 2020.
- Gasson, E., Lunt, D. J., DeConto, R., Goldner, A., Heinemann, M., Huber, M., LeGrande, A. N., Pollard, D., Sagoo, N., Siddall, M., Winguth, A., and Valdes, P. J.: Uncertainties in the modelled CO₂ threshold for Antarctic glaciation, *Climate of the Past*, 10, 451–466, <https://doi.org/10.5194/cp-10-451-2014>, 2014.
- 690 Gasson, E., DeConto, R. M., Pollard, D., and Levy, R. H.: Dynamic Antarctic ice sheet during the early to mid-Miocene, *Proceedings of the National Academy of Sciences*, 113, 3459–3464, <https://doi.org/10.1073/pnas.1516130113>, 2016.
- Greenop, R., Foster, G. L., Wilson, P. A., and Lear, C. H.: Middle Miocene climate instability associated with high-amplitude CO₂ variability, *Paleoceanography*, 29, 845–853, <https://doi.org/10.1002/2014PA002653>, 2014.
- 695 Halberstadt, A. R. W., Chorley, H., Levy, R. H., Naish, T., DeConto, R. M., Gasson, E., and Kowalewski, D. E.: CO₂ and tectonic controls on Antarctic climate and ice-sheet evolution in the mid-Miocene, *Earth and Planetary Science Letters*, 564, 116908, <https://doi.org/10.1016/j.epsl.2021.116908>, 2021.
- Hartman, J. D., Sangiorgi, F., Salabarnada, A., Peterse, F., Houben, A. J. P., Schouten, S., Brinkhuis, H., Escutia, C., and Bijl, P. K.: Paleocene and ice sheet variability offshore Wilkes Land, Antarctica—Part 3: Insights from Oligocene–Miocene TEX₈₆-based sea surface temperature reconstructions, *Climate of the Past*, 14, 1275–1297, <https://doi.org/10.5194/cp-14-1275-2018>, 2018.
- 700 Hauptvogel, D. W. and Passchier, S.: Early–Middle Miocene (17–14 Ma) Antarctic ice dynamics reconstructed from the heavy mineral provenance in the AND-2A drill core, Ross Sea, Antarctica, *Global and Planetary Change*, 82, 38–50, <https://doi.org/10.1016/j.gloplacha.2011.11.003>, 2012.
- 705 Herold, N., Seton, M., Müller, R. D., You, Y., and Huber, M.: Middle Miocene tectonic boundary conditions for use in climate models, *Geochemistry, Geophysics, Geosystems*, 9, <https://doi.org/10.1029/2008GC002046>, 2008.

- Hochmuth, K., Gohl, K., Leitchenkov, G., Sauermilch, I., Whittaker, J. M., Uenzelmann-Neben, G., Davy, B., and De Santis, L.: The evolving paleobathymetry of the circum-Antarctic Southern Ocean since 34 Ma: A key to understanding past cryosphere-ocean developments, *Geochemistry, Geophysics, Geosystems*, 21, e2020GC009122, <https://doi.org/10.1029/2020GC009122>, 2020a.
- 710 Hochmuth, K., Paxman, G. J. G., Gohl, K., Jamieson, S. S. R., Leitchenkov, G. L., Bentley, M. J., Ferraccioli, F., Sauermilch, I., Whittaker, J., Uenzelmann-Neben, G., Davy, B., and DeSantis, L.: Combined palaeotopography and palaeobathymetry of the Antarctic continent and the Southern Ocean since 34 Ma, <https://doi.org/10.1594/PANGAEA.923109>, 2020b.
- Holbourn, A., Kuhnt, W., Clemens, S., Prell, W., and Andersen, N.: Middle to late Miocene stepwise climate cooling: Evidence from a high-resolution deep water isotope curve spanning 8 million years, *Paleoceanography*, 28, 688–699, <https://doi.org/10.1002/2013PA002538>,
715 2013.
- Huybrechts, P.: Sea-level changes at the LGM from ice-dynamic reconstructions of the Greenland and Antarctic ice sheets during the glacial cycles, *Quaternary Science Reviews*, 21, 203–231, [https://doi.org/10.1016/S0277-3791\(01\)00082-8](https://doi.org/10.1016/S0277-3791(01)00082-8), 2002.
- Huybrechts, P. and de Wolde, J.: The dynamic response of the Greenland and Antarctic ice sheets to multiple-century climatic warming, *Journal of Climate*, 12, 2169–2188, [https://doi.org/10.1175/1520-0442\(1999\)012<2169:TDROTG>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<2169:TDROTG>2.0.CO;2), 1999.
- 720 Janssens, I. and Huybrechts, P.: The treatment of meltwater retention in mass-balance parameterizations of the Greenland ice sheet, *Annals of Glaciology*, 31, 133–140, <https://doi.org/10.3189/172756400781819941>, 2000.
- Kürschner, W. M., Kvaček, Z., and Dilcher, D. L.: The impact of Miocene atmospheric carbon dioxide fluctuations on climate and the evolution of terrestrial ecosystems, *Proceedings of the National Academy of Sciences*, 105, 449–453, <https://doi.org/10.1073/pnas.0708588105>, 2008.
- 725 Ladant, J.-B., Donnadieu, Y., Lefebvre, V., and Dumas, C.: The respective role of atmospheric carbon dioxide and orbital parameters on ice sheet evolution at the Eocene-Oligocene transition, *Paleoceanography*, 29, 810–823, <https://doi.org/10.1002/2013PA002593>, 2014.
- Langebroek, P. M., Paul, A., and Schulz, M.: Antarctic ice-sheet response to atmospheric CO₂ and insolation in the Middle Miocene, *Climate of the Past*, 5, 633–646, <https://doi.org/10.5194/cp-5-633-2009>, 2009.
- Langebroek, P. M., Paul, A., and Schulz, M.: Simulating the sea level imprint on marine oxygen isotope records during the middle Miocene using an ice sheet–climate model, *Paleoceanography*, 25, <https://doi.org/10.1029/2008PA001704>, 2010.
730
- Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A. C. M., and Levrard, B.: A long-term numerical solution for the insolation quantities of the Earth, *Astronomy & Astrophysics*, 428, 261–285, <https://doi.org/10.1051/0004-6361:20041335>, 2004.
- Lazeroms, W. M. J., Jenkins, A., Rienstra, S. W., and Van de Wal, R. S. W.: An analytical derivation of ice-shelf basal melt based on the dynamics of meltwater plumes, *Journal of Physical Oceanography*, 49, 917–939, <https://doi.org/10.1175/JPO-D-18-0131.1>, 2019.
- 735 Le Meur, E. and Huybrechts, P.: A comparison of different ways of dealing with isostasy: examples from modelling the Antarctic ice sheet during the last glacial cycle, *Annals of Glaciology*, 23, 309–317, <https://doi.org/10.3189/S0260305500013586>, 1996.
- Levy, R., Harwood, D., Florindo, F., Sangiorgi, F., Tripathi, R., Von Eynatten, H., Gasson, E., Kuhn, G., Tripathi, A., DeConto, R., et al.: Antarctic ice sheet sensitivity to atmospheric CO₂ variations in the early to mid-Miocene, *Proceedings of the National Academy of Sciences*, 113, 3453–3458, <https://doi.org/10.1073/pnas.1516030113>, 2016.
- 740 Levy, R. H., Meyers, S. R., Naish, T. R., Gollledge, N. R., McKay, R. M., Crampton, J. S., DeConto, R. M., De Santis, L., Florindo, F., Gasson, E. G. W., et al.: Antarctic ice-sheet sensitivity to obliquity forcing enhanced through ocean connections, *Nature Geoscience*, 12, 132–137, <https://doi.org/10.1038/s41561-018-0284-4>, 2019.
- Liebrand, D., Lourens, L., Hodell, D. A., Boer, B. d., Van de Wal, R. S. W., and Pälike, H.: Antarctic ice sheet and oceanographic response to eccentricity forcing during the early Miocene, *Climate of the Past*, 7, 869–880, <https://doi.org/10.5194/cp-7-869-2011>, 2011.

- 745 Liebrand, D., de Bakker, A. T. M., Beddow, H. M., Wilson, P. A., Bohaty, S. M., Ruessink, G., Pälike, H., Batenburg, S. J., Hilgen, F. J., Hodell, D. A., et al.: Evolution of the early Antarctic ice ages, *Proceedings of the National Academy of Sciences*, 114, 3867–3872, <https://doi.org/10.1073/pnas.1615440114>, 2017.
- Lingle, C. S. and Clark, J. A.: A numerical model of interactions between a marine ice sheet and the solid earth: Application to a West Antarctic ice stream, *Journal of Geophysical Research: Oceans*, 90, 1100–1114, <https://doi.org/10.1029/JC090iC01p01100>, 1985.
- 750 Lorius, C., Jouzel, J., Ritz, C., Merlivat, L., Barkov, N. I., Korotkevich, Y. S., and Kotlyakov, V. M.: A 150,000-year climatic record from Antarctic ice, *Nature*, 316, 591–596, <https://doi.org/10.1038/316591a0>, 1985.
- Marschalek, J. W., Zurli, L., Talarico, F., van de Flierdt, T., Vermeesch, P., Carter, A., Beny, F., Bout-Roumazeilles, V., Sangiorgi, F., Hemming, S. R., et al.: A large West Antarctic Ice Sheet explains early Neogene sea-level amplitude, *Nature*, 600, 450–455, <https://doi.org/10.1038/s41586-021-04148-0>, 2021.
- 755 Martin, M. A., Winkelmann, R., Haseloff, M., Albrecht, T., Bueler, E., Khroulev, C., and Levermann, A.: The Potsdam Parallel Ice Sheet Model (PISM-PIK) – Part 2: Dynamic equilibrium simulation of the Antarctic ice sheet, *The Cryosphere*, 5, 727–740, <https://doi.org/10.5194/tc-5-727-2011>, 2011.
- Modestou, S. E., Leutert, T. J., Fernandez, A., Lear, C. H., and Meckler, A. N.: Warm middle Miocene Indian Ocean bottom water temperatures: Comparison of clumped isotope and Mg/Ca-based estimates, *Paleoceanography and Paleoclimatology*, 35, e2020PA003927, <https://doi.org/10.1029/2020PA003927>, 2020.
- 760 Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., Eisen, O., Ferraccioli, F., Forsberg, R., Fretwell, P., et al.: Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet, *Nature Geoscience*, 13, 132–137, <https://doi.org/10.1038/s41561-019-0510-8>, 2020.
- Naish, T., Powell, R., Levy, R., Wilson, G., Scherer, R., Talarico, F., Krissek, L., Niessen, F., Pompilio, M., Wilson, T., et al.: Obliquity-paced Pliocene West Antarctic ice sheet oscillations, *Nature*, 458, 322–328, <https://doi.org/10.1038/nature07867>, 2009.
- 765 Oerlemans, J.: Antarctic ice volume and deep-sea temperature during the last 50 Myr: a model study, *Annals of Glaciology*, 39, 13–19, <https://doi.org/10.3189/172756404781814708>, 2004.
- Ohmura, A.: Precipitation, accumulation and mass balance of the Greenland ice sheet, *Zeitschrift für Gletscherkunde und Glazialgeologie*, 35, 1–20, 1999.
- 770 Paxman, G. J. G., Jamieson, S. S. R., Hochmuth, K., Gohl, K., Bentley, M. J., Leitchenkov, G., and Ferraccioli, F.: Reconstructions of Antarctic topography since the Eocene–Oligocene boundary, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 535, 109–134, <https://doi.org/10.1016/j.palaeo.2019.109346>, 2019.
- Paxman, G. J. G., Gasson, E. G. W., Jamieson, S. S. R., Bentley, M. J., and Ferraccioli, F.: Long-Term Increase in Antarctic Ice Sheet Vulnerability Driven by Bed Topography Evolution, *Geophysical Research Letters*, 47, e2020GL090003, <https://doi.org/10.1029/2020GL090003>, 2020.
- 775 Pekar, S. F. and DeConto, R. M.: High-resolution ice-volume estimates for the early Miocene: Evidence for a dynamic ice sheet in Antarctica, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 231, 101–109, <https://doi.org/10.1016/j.palaeo.2005.07.027>, 2006.
- Pelle, T., Morlighem, M., and Bondzio, J. H.: Brief communication: PICOP, a new ocean melt parameterization under ice shelves combining PICO and a plume model, *The Cryosphere*, 13, 1043–1049, <https://doi.org/10.5194/tc-13-1043-2019>, 2019.
- 780 Pérez, L. F., De Santis, L., McKay, R. M., Larter, R. D., Ash, J., Bart, P. J., Böhm, G., Brancatelli, G., Browne, I., Colleoni, F., et al.: Early and middle Miocene ice sheet dynamics in the Ross Sea: Results from integrated core-log-seismic interpretation, *GSA Bulletin*, <https://doi.org/10.1130/B35814.1>, 2021.

- Pollard, D.: A retrospective look at coupled ice sheet–climate modeling, *Climatic Change*, 100, 173–194, <https://doi.org/10.1007/s10584-010-9830-9>, 2010.
- 785 Pollard, D. and DeConto, R. M.: Antarctic ice and sediment flux in the Oligocene simulated by a climate–ice sheet–sediment model, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 198, 53–67, [https://doi.org/10.1016/S0031-0182\(03\)00394-8](https://doi.org/10.1016/S0031-0182(03)00394-8), 2003.
- Pollard, D. and DeConto, R. M.: Hysteresis in Cenozoic Antarctic ice-sheet variations, *Global and Planetary Change*, 45, 9–21, <https://doi.org/10.1016/j.gloplacha.2004.09.011>, 2005.
- Pollard, D. and DeConto, R. M.: Modelling West Antarctic ice sheet growth and collapse through the past five million years, *Nature*, 458, 329–332, <https://doi.org/10.1038/nature07809>, 2009.
- 790 Pollard, D. and DeConto, R. M.: Continuous simulations over the last 40 million years with a coupled Antarctic ice sheet–sediment model, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 537, 109 374, <https://doi.org/10.1016/j.palaeo.2019.109374>, 2020.
- Pollard, D., Kump, L. R., and Zachos, J. C.: Interactions between carbon dioxide, climate, weathering, and the Antarctic ice sheet in the earliest Oligocene, *Global and Planetary Change*, 111, 258–267, <https://doi.org/10.1016/j.gloplacha.2013.09.012>, 2013.
- 795 Reerink, T. J., Kliphuis, M. A., and van de Wal, R. S. W.: Mapping technique of climate fields between GCM’s and ice models, *Geoscientific Model Development*, 3, 13–41, <https://doi.org/10.5194/gmd-3-13-2010>, 2010.
- Reese, R., Albrecht, T., Mengel, M., Asay-Davis, X., and Winkelmann, R.: Antarctic sub-shelf melt rates via PICO, *The Cryosphere*, 12, 1969–1985, <https://doi.org/10.5194/tc-12-1969-2018>, 2018a.
- Reese, R., Gudmundsson, G. H., Levermann, A., and Winkelmann, R.: The far reach of ice-shelf thinning in Antarctica, *Nature Climate Change*, 8, 53–57, <https://doi.org/10.1038/s41558-017-0020-x>, 2018b.
- 800 Ritz, C., Rommelaere, V., and Dumas, C.: Modeling the evolution of Antarctic ice sheet over the last 420,000 years: Implications for altitude changes in the Vostok region, *Journal of Geophysical Research: Atmospheres*, 106, 31 943–31 964, <https://doi.org/10.1029/2001JD900232>, 2001.
- Rohling, E. J., Yu, J., Heslop, D., Foster, G. L., Opdyke, B., and Roberts, A. P.: Sea level and deep-sea temperature reconstructions suggest quasi-stable states and critical transitions over the past 40 million years, *Science Advances*, 7, eabf5326, <https://doi.org/10.1126/sciadv.abf5326>, 2021.
- 805 Sangiorgi, F., Bijl, P. K., Passchier, S., Salzmann, U., Schouten, S., McKay, R., Cody, R. D., Pross, J., Van De Flierdt, T., Bohaty, S. M., et al.: Southern Ocean warming and Wilkes Land ice sheet retreat during the mid-Miocene, *Nature Communications*, 9, 1–11, <https://doi.org/10.1038/s41467-017-02609-7>, 2018.
- 810 Schoof, C.: Ice sheet grounding line dynamics: Steady states, stability, and hysteresis, *Journal of Geophysical Research: Earth Surface*, 112, <https://doi.org/10.1029/2006JF000664>, 2007.
- Shapiro, N. M. and Ritzwoller, M. H.: Inferring surface heat flux distributions guided by a global seismic model: particular application to Antarctica, *Earth and Planetary Science Letters*, 223, 213–224, <https://doi.org/10.1016/j.epsl.2004.04.011>, 2004.
- Shevenell, A. E., Kennett, J. P., and Lea, D. W.: Middle Miocene ice sheet dynamics, deep-sea temperatures, and carbon cycling: A Southern Ocean perspective, *Geochemistry, Geophysics, Geosystems*, 9, <https://doi.org/10.1029/2007GC001736>, 2008.
- 815 Stap, L. B., Van De Wal, R. S. W., De Boer, B., Bintanja, R., and Lourens, L. J.: The MMCO-EOT conundrum: Same benthic $\delta^{18}\text{O}$, different CO_2 , *Paleoceanography*, 31, 1270–1282, <https://doi.org/10.1002/2016PA002958>, 2016.
- Stap, L. B., Van de Wal, R. S. W., De Boer, B., Bintanja, R., and Lourens, L. J.: The influence of ice sheets on temperature during the past 38 million years inferred from a one-dimensional ice sheet–climate model, *Climate of the Past*, 13, 1243–1257, [https://doi.org/10.5194/cp-](https://doi.org/10.5194/cp-13-1243-2017)
- 820 13-1243-2017, 2017.

- Stap, L. B., Sutter, J., Knorr, G., Stärz, M., and Lohmann, G.: Transient variability of the Miocene Antarctic ice sheet smaller than equilibrium differences, *Geophysical Research Letters*, 46, 4288–4298, <https://doi.org/10.1029/2019GL082163>, 2019.
- Stap, L. B., Knorr, G., and Lohmann, G.: Anti-phased Miocene ice volume and CO₂ Changes by transient Antarctic ice sheet variability, *Paleoceanography and Paleoclimatology*, 35, e2020PA003 971, <https://doi.org/10.1029/2020PA003971>, 2020.
- 825 Stärz, M., Jokat, W., Knorr, G., and Lohmann, G.: Threshold in North Atlantic-Arctic Ocean circulation controlled by the subsidence of the Greenland-Scotland Ridge, *Nature Communications*, 8, 1–13, <https://doi.org/10.1038/ncomms15681>, 2017.
- Steinthorsdottir, M., Coxall, H., De Boer, A. M., Huber, M., Barbolini, N., Bradshaw, C. D., Burls, N. J., Feakins, S. J., Gasson, E., Henderiks, J., et al.: The Miocene: The future of the past, *Paleoceanography and Paleoclimatology*, 36, e2020PA004 037, <https://doi.org/10.1029/2020PA004037>, 2021a.
- 830 Steinthorsdottir, M., Jardine, P. E., and Rember, W. C.: Near-Future pCO₂ During the Hot Miocene Climatic Optimum, *Paleoceanography and Paleoclimatology*, 36, e2020PA003 900, <https://doi.org/10.1029/2020PA003900>, 2021b.
- Stocchi, P., Escutia, C., Houben, A. J. P., Vermeersen, B. L. A., Bijl, P. K., Brinkhuis, H., DeConto, R. M., Galeotti, S., Passchier, S., Pollard, D., et al.: Relative sea-level rise around East Antarctica during Oligocene glaciation, *Nature Geoscience*, 6, 380–384, <https://doi.org/10.1038/ngeo1783>, 2013.
- 835 Sun, S., Pattyn, F., Simon, E. G., Albrecht, T., Cornford, S., Calov, R., Dumas, C., Gillet-Chaulet, F., Goelzer, H., Golledge, N. R., et al.: Antarctic ice sheet response to sudden and sustained ice-shelf collapse (ABUMIP), *Journal of Glaciology*, 66, 891–904, <https://doi.org/10.1017/jog.2020.67>, 2020.
- Super, J. R., Thomas, E., Pagani, M., Huber, M., O’Brien, C., and Hull, P. M.: North Atlantic temperature and pCO₂ coupling in the early-middle Miocene, *Geology*, 46, 519–522, <https://doi.org/10.1130/G40228.1>, 2018.
- 840 Tan, N., Ladant, J.-B., Ramstein, G., Dumas, C., Bachem, P., and Jansen, E.: Dynamic Greenland ice sheet driven by pCO₂ variations across the Pliocene Pleistocene transition, *Nature Communications*, 9, 1–9, <https://doi.org/10.1038/s41467-018-07206-w>, 2018.
- Thompson, S. L. and Pollard, D.: Greenland and Antarctic mass balances for present and doubled atmospheric CO₂ from the GENESIS version-2 global climate model, *Journal of Climate*, 10, 871–900, [https://doi.org/10.1175/1520-0442\(1997\)010<0871:GAAMBF>2.0.CO;2](https://doi.org/10.1175/1520-0442(1997)010<0871:GAAMBF>2.0.CO;2), 1997.
- 845 Uppala, S. M., Kållberg, P. W., Simmons, A. J., Andrae, U., Bechtold, V. D. C., Fiorino, M., Gibson, J. K., Haseler, J., Hernandez, A., Kelly, G. A., et al.: The ERA-40 re-analysis, *Quarterly Journal of the Royal Meteorological Society*, 131, 2961–3012, <https://doi.org/10.1256/qj.04.176>, 2005.
- Wilson, D. S., Jamieson, S. S. R., Barrett, P. J., Leitchenkov, G., Gohl, K., and Larter, R. D.: Antarctic topography at the Eocene–Oligocene boundary, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 335, 24–34, <https://doi.org/10.1016/j.palaeo.2011.05.028>, 2012.
- 850 Winkelmann, R., Martin, M. A., Haseloff, M., Albrecht, T., Bueler, E., Khroulev, C., and Levermann, A.: The Potsdam Parallel Ice Sheet Model (PISM-PIK) – Part 1: Model description, *The Cryosphere*, 5, 715–726, <https://doi.org/10.5194/tc-5-715-2011>, 2011.
- Zachos, J. C., Dickens, G. R., and Zeebe, R. E.: An early Cenozoic perspective on greenhouse warming and carbon-cycle dynamics, *Nature*, 451, 279–283, <https://doi.org/10.1038/nature06588>, 2008.

Table 1. Name and description of the experiments carried out. Each experiment comprises 15 simulations, as explained in the main text and listed in Table 2.

Simulation name	Description
REF	Default settings (see Sect. 2)
SMB	Increased ablation: $c_3 = -0.25 \text{ m yr}^{-1}$
INSOL	Interpolation between forcing climates with low (Ifucobi) and high summer insolation (3nowabi)
FEEDB - NOFEEDB	Interpolation of climate forcing only based on CO_2
FEEDB-onlyT - NOFEEDB-T	Interpolation of temperature only based on CO_2
FEEDB-onlyP - NOFEEDB-P	Interpolation of precipitation only based on CO_2
TOPO 14Ma	Bedrock topography pertaining to 14 Myr ago (Hochmuth et al., 2020a; Paxman et al., 2019)
TOPO 34Ma	Bedrock topography pertaining to 34 Myr ago (Hochmuth et al., 2020a; Paxman et al., 2019)
TOPO PD	Present-day bedrock topography (Morlighem et al., 2020), isostatically rebounded after ice sheet removal
TOPO Wilson_max	Bedrock topography pertaining to Late Eocene (Wilson et al., 2012), max reconstruction
TOPO Wilson_mean	Bedrock topography pertaining to Late Eocene (Wilson et al., 2012), mean of min and max reconstructions
BMB LGM	LGM ocean conditions ($M_{expo} = 0 \text{ m/yr}$, $M_{deep} = 2 \text{ m/yr}$, and $T_w = -5^\circ\text{C}$)
BMB no_shelves	Constant sub-shelf melt rate $M = 400 \text{ m/yr}$

Table 2. Description of the 15 simulations spanning each experiment.

Type	Initial conditions	CO_2 levels	Runtime (model years)
Steady-state	Miocene topography, no ice	280, 392, 504, 616, 728, and 840 ppm	150 kyr
Steady-state	280-ppm steady-state IMAU-ICE run	392, 504, 616, 728, and 840 ppm	150 kyr
Transient	840-ppm steady-state IMAU-ICE run	linearly from 840 to 280 to 840 ppm	100, 400, and 200 (5x40) kyr
Transient	280-ppm steady-state IMAU-ICE run	linearly from 280 to 840 to 280 ppm	200 (5x40) kyr




Figure 1. (a) Evolution of forcing CO₂ levels (red/pink) and ice volume above floatation (in meters sea level equivalent; black) over time (relative to the length of the simulation), for the transient 100-kyr (dotted) and 400-kyr (long-dashed) REF simulations. (b) Relation between CO₂ and equilibrium ice volume (ascending branch, blue; descending branch, red), and transient ice volume (black as in (a)). (c) and (d) Same for the regular transient 40-kyr simulations starting from warm (CO₂ red/pink, ice volume solid black) and reversed cold (CO₂ blue/purple, ice volume dashed black) transient 40-kyr simulations conditions. Arrows in (b) and (d) indicate the progression direction of ice volume. This direction is similar (lower branch ascending in (d), upper branch descending) in further figures the transient ice volume evolutions of both simulations partly overlap.




Figure 2. (a) Evolution of forcing CO_2 levels (red) and Simulated equilibrated grounded ice volume (black) over relative time, for the transient 40-kyr thickness from experiment REF (solid) with 840-ppm, and SMB (dotted) simulations. (b) Relation between 280-ppm CO_2 and equilibrium ice volume (REF forcing, solid; SMB, dotted), and transient starting without ice volume. Only the equilibrium cycle (final 40 kyr) is shown. (c) and (d) Same for the REF (solid) and INSOL (dashed, orange) transient 40-kyr simulations. Cyan areas indicate ice shelf extent, black areas ice volume, and grey areas ice-free land.




Figure 3. (a) Evolution of forcing CO₂ levels (red/pink) and ice volume over relative time, for the transient 40-kyr REF (black) and FEEDB/NOFEEDB (orange) simulations. (b) Relation between CO₂ and equilibrium ice volume (REF, solid red and blue; FEEDB/NOFEEDB, dashed red and blue), and transient ice volume (as in (a)). Only the equilibrium cycle (final 40 kyr) is shown. (c) and (d) Same for the FEEDB-onlyP/NOFEEDB-P (dotted red, blue and orange) and FEEDB-onlyT/NOFEEDB-T (long-dashed red, blue and orange) equilibrium and transient 40-kyr simulations. The amplitudes of transient ice volume variability (in m.s.l.e.) in the equilibrium cycle are indicated in panels (a) and (c).




Figure 4. (a) Simulated equilibrated grounded ice thickness from experiment TOPO 14Ma with 280-ppm CO₂ forcing. Cyan areas indicate ice shelf extent, grey areas ice-free land. (b) Difference in ice thickness between TOPO 14Ma and REF with 280-ppm CO₂ forcing (grounding line and continental edge as in (a)).




Figure 5. (a) Evolution of forcing CO₂ levels (red line) and ice volume over relative time, for the transient 40-kyr REF (black) and TOPO 14Ma (orange) simulations. (b) Relation between CO₂ and equilibrium ice volume (REF, solid red and blue; TOPO 14Ma, dashed red and blue), and transient ice volume (as in (a)). Only the equilibrium cycle (final 40 kyr) is shown.




Figure 6. (a) Evolution of forcing CO₂ levels (red line) and ice volume over relative time, for the transient 400-kyr BMB no_shelves (solid orange) and BMB LGM (dashed green) simulations. (b) Relation between CO₂ and equilibrium ice volume (BMB no_shelves, solid red and blue; BMB LGM, dashed red and blue), and transient ice volume. (c) and (d) Same for the transient 40-kyr simulations. Only the equilibrium cycle (final 40 kyr) is shown.




Figure 7. (a) Simulated equilibrated grounded ice thickness from experiment BMB LGM with 504-ppm, and (b) 392-ppm CO₂ forcing, starting without ice. Cyan areas indicate ice shelf extent, grey areas ice-free land.