

Brief communication: Do 1.0°C, 1.5°C or 2.0°C matter for the future evolution of Alpine glaciers?

Loris Compagno^{1,2}, Sarah Eggs^{1,2}, Matthias Huss^{1,2,3}, Harry Zekollari^{4,5,1,2}, and Daniel Farinotti^{1,2}

¹Laboratory of Hydraulics, Hydrology and Glaciology (VAW), ETH Zurich, Zurich, Switzerland.

²Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Birmensdorf, Switzerland.

³Department of Geosciences, University of Fribourg, Fribourg, Switzerland.

⁴Department of Geoscience and Remote Sensing, Delft University of Technology, Netherlands.

⁵Laboratoire de Glaciologie, Université libre de Bruxelles, Belgium.

Correspondence: Loris Compagno <compagno@vaw.baug.ethz.ch>

Abstract. With the Paris Agreement, the urgency of limiting ongoing anthropogenic climate change has been recognized. More recent discussions have focused on the difference of limiting the increase in global average temperatures below 1.0, 1.5, or 2.0°C compared to pre-industrial levels. Here, we assess the impacts that such different scenarios would have on both the future evolution of glaciers in the European Alps and the water resources they provide. Our results show that even half-degree differences in global temperature targets have important implications for the changes predicted until 2100, and that – for the most optimistic scenarios – glaciers might start to partially recover owing to possibly decreasing temperatures after the end of the 21st century.

1 Introduction

Climate change is one of the largest challenges that society will face during this century. There is overwhelming consensus that limiting the increase of global average temperatures below a certain threshold is essential if adverse effects are to be avoided. Put forward in 2015 and adopted by 196 parties, the Paris Agreement concurred that such a threshold is “*well below 2°C above pre-industrial levels*”, and that efforts should be pursued to “*limit the temperature increase to 1.5°C*” (UN, 2015). Even under these ambitious climate targets, important environmental changes, such as changes in water availability, migration of species, or glacier loss, are expected to occur. The Special Report on Global Warming of 1.5°C (IPCC, 2018), for example, focused on the effects that would result if the 1.5°C target was to be met, and highlighted that even under such conditions, important and mostly negative changes would occur. Five years after the Agreement, there is compelling evidence that any reduction of anthropogenic climate change will pay off in the longer term, both from an ecological (IPBES, 2019) and economical (WEF, 2020) perspective. Discussions in preparation for the renewed pledges by signatory parties have thus focused on the differences between scenarios close to the original targets. For the upcoming 26th United Nations Climate Change Conference of the Parties, for example, emphasis is put onto separating the impacts occurring under scenarios of 1.0, 1.5, and 2.0°C of warming above pre-industrial levels. In this brief communication, we focus on the differing impacts that such scenarios would have on glaciers and related water resources in the European Alps. Across the world, glaciers are amongst the most prominent indicators

for climatic change, providing visual evidence for climatic changes occurring over decades. Previous work estimated that every kg of additionally emitted CO₂ would result in a long-term global glacier mass loss of ca. 16 kg (Marzeion et al., 2018) – with important implications for the corresponding landscapes, downstream ecosystems, and water supplies (IPCC, 2018, 2019) . Whilst future projections for the glacier evolution of the European Alps already exist under different representative concentrations pathways (RCPs) (e.g. Zekollari et al., 2019; Marzeion et al., 2020), targeted information on policy-relevant climate targets such as the difference between 1.5 and 2.0°C is not readily available. Here we analyse glacier evolution under low global warming scenarios by re-running the mass-balance ice-flow model GloGEMflow (Huss and Hock, 2015; Zekollari et al., 2019), with updated climate projections from the 6th phase of the Coupled Model Intercomparison Project (CMIP6). In particular, we focus on Shared Socioeconomic Pathways (SSPs) that would result in global mean temperature changes of +1.0, +1.5 and +2.0°C by 2100 compared to pre-industrial conditions (1850-1900), and provide projections for both future glacier geometries and glacier runoff until the end of the century. In an explorative analysis, we also consider projections until the year 2300, albeit acknowledging that large uncertainties exist when doing so.

35 2 Data and Methods

We simulate the evolution of all 3927 glaciers in the Alps independently based on (i) outlines provided by the Randolph Glacier Inventory version 6 (RGI 6.0) (RGI Consortium, 2017) and (ii) the consensus ice thickness estimate (Farinotti et al., 2019).

Glacier evolution is modelled with the combined mass-balance ice-flow model GloGEMflow (Huss and Hock, 2015; Zekollari et al., 2019), a model that showed robust performance in recent model intercomparison projects (e.g. Marzeion et al., 2020). Accumulation is computed based on precipitation and temperature, with a linear transition between solid and liquid precipitation occurring for temperatures between 0.5°C and 2.5°C. Ablation is calculated with a temperature-index model (Hock, 2003), and refreezing is determined based on heat conduction. The mass balance module runs at a monthly resolution, including a parametrization for day-to-day temperature variability. It is calibrated to reproduce glacier specific-geodetic volume changes (Zemp et al., 2019) spanning approximatively the period 1959–2013 and covering 57 % of the Alpine glacier area. For more details on the mass balance module and its calibration, refer to Huss and Hock (2015).

For all glaciers with a length >1 km in RGI 6.0 (i.e. 795 glaciers, accounting for 95 % of the Alpine glacier volume), we compute ice flow based on the shallow ice approximation. The computations follow Zekollari et al. (2019), including the iterative initialisation procedure used to generate a transient glacier geometry for the RGI inventory year (i.e. 2003 for the majority of Alpine glaciers). For all glaciers with length <1 km, instead, glacier evolution is modelled with an elevation-dependent parametrization that has shown to be in good agreement with higher-order ice-flow modeling results (Huss et al., 2010). For more information on GloGEMflow’s glacier evolution modelling, please refer to Zekollari et al. (2019).

For each glacier, water runoff is calculated at a monthly resolution by summing rain and melt (from ice, snow and firn) and by subtracting refreezing. Throughout the simulations, runoff is computed for the area comprised within the RGI 6.0 outlines, implying that, after glacier retreat, runoff contributions from rain and snow melt are still accounted for (in line with Huss and Hock, 2018).

For the past (1950-2020), GloGEMflow is forced with monthly 2-m air temperature and precipitation taken from the ensemble daily gridded observational dataset (E-OBS v.21.0) at 0.1° (11 km) horizontal resolution (Cornes et al., 2018). For the future (2020-2100), we rely on 128 CMIP6 global circulation model (GCM) members. We use direct GCM output, since CMIP6 GCM-results have not yet been downscaled by regional climate models, and since the spatial resolution of the climate data was recently shown to only marginally affect model results if suitable model calibration at the glacier-specific scale is performed (Compagno et al., 2021). Consistence between past and future climate is ensured by applying the de-biasing procedure proposed by Huss and Hock (2015). In a nutshell, the procedure uses a set of additive and multiplicative correction factors to adjust both the long-term mean and the long-term variability of the coarse-resolution GCMs (100 km) to the level of the high-resolution E-OBS data (see Huss and Hock, 2015, for more details).

To evaluate model performance, we compare modelled mass balances with observations provided by the World Glacier Monitoring Service (WGMS, 2020) for 72 glaciers in the European Alps. For glacier-wide annual mass balance, we find a bias of $0.07 \text{ m w.e. a}^{-1}$, a Root Mean Square Error (RMSE) of $1.07 \text{ m w.e. a}^{-1}$, and a squared correlation coefficient (r^2) of 0.29 (see Fig. S1). Observations aggregated to elevation bands yield a bias of $-0.01 \text{ m w.e. a}^{-1}$, a RMSE of $1.47 \text{ m w.e. a}^{-1}$, and a r^2 of 0.52. For glacier-wide winter balance, instead, a bias of $0.20 \text{ m w.e. a}^{-1}$, a RMSE of $0.74 \text{ m w.e. a}^{-1}$, and a r^2 of 0.26 is found (see Fig. S2). Albeit correlation coefficients are rather low, the small biases provide confidence in the regional results, which neither indicate over- nor underestimation. The latter is also confirmed by the comparison of the total glacier volume change in Switzerland (comprising almost 60% of the Alpine glacier volume at present) during the period 2000-2020: our modelled loss is of 24.9 km^3 , which is very close to the observation-based estimate of 23.6 km^3 (Grab et al., in press). Similarly, the modelled rate of glacier area loss ($1.4 \% \text{ a}^{-1}$ for the period 2003–2016) is in very good agreement with the rate of $1.3 \% \text{ a}^{-1}$ derived by Paul et al. (2020) using Landsat images.

For our main analysis, finally, we only select those GCM members that yield a 21st century global warming of $+1.0$, $+1.5$ and $+2.0^\circ\text{C}$ compared to pre-industrial. The selected members correspond to SSPs that assume low emissions, i.e. SSP119, SSP126 and SSP245, and are used to force GloGEMflow for the period 2020-2100. This selection is performed by computing the mean global 2-m air temperature change over both land and ocean between the period 2071-2100 and pre-industrial (1850-1900) levels, and by selecting members that are within a given range of temperature increase: $+1 \pm 0.25^\circ\text{C}$ (3 model members), $+1.5 \pm 0.25^\circ\text{C}$ (11 model members), and $+2 \pm 0.25^\circ\text{C}$ (14 model members). To determine the corresponding warming in the European Alps (see Fig. 1 and S3), we extract all GCM grid cells that are within latitudes of 42.5°N - 49.5°N , longitudes of 4°E - 16°E , and at elevations above 500 m.

3 Results and Discussion

The GCMs selected within the three temperature change targets (i.e. $+1.0$, $+1.5$, and $+2.0^\circ\text{C}$) show a global mean warming between 1850-1900 and 2071-2100 of $+1.00 \pm 0.11$, $+1.49 \pm 0.17$, and $+2.07 \pm 0.18^\circ\text{C}$. For the European Alps, this corresponds to a temperature change of $+0.98 \pm 0.56$, $+1.80 \pm 0.72$ and $+2.51 \pm 0.73^\circ\text{C}$, respectively (Fig. 1a). During the summer months (JJA), the temperature increase in the Alps for the two warmer climate targets is even stronger, and is of 0.96 ± 0.88 ,




Figure 1. Evolution of annual global and Alpine (a) average air temperature and (b) precipitation total compared to the pre-industrial (1850-1900) baseline. Series are smoothed with a 30-year running mean. The coloured bands show the standard deviation of the projection given by individual GCM members. n is the considered numbers of GCM members.

2.09 \pm 1.24, and 2.81 \pm 1.23°C, respectively. These differences agree well with previous assessments (Vautard et al., 2014), and are attributed to stronger warming over land than over ocean, to enhanced warming at high latitudes as compared to low latitudes, and to amplified warming at higher elevations (Vautard et al., 2014). For a global warming of +1.0, +1.5, and +2 °C, global-scale annual precipitation sums for 2071-2100 are projected to increase by 18 \pm 8, 29 \pm 9, and 42 \pm 10 mm a⁻¹, respectively, up from 1141 mm a⁻¹ in the pre-industrial baseline (Fig. 1b). For the Alps, the annual precipitation increase is anticipated to be of 37 \pm 94, 29 \pm 135, and 46 \pm 145 mm a⁻¹, respectively (the pre-industrial baseline is 1210 mm a⁻¹), with slightly higher precipitation changes over the winter season.

In all three considered warming scenarios, glaciers will be drastically affected. By 2100, glaciers in the European Alps are projected to lose 44 \pm 21 % (+1.0°C), 68 \pm 12 % (+1.5°C), and 81 \pm 8 % (+2.0°C) of their 2020 ice volume (see Fig. 2a). The evolution of the glacier area will be on a par (Fig. 2b). By 2100, Alpine glaciers are projected to lose 47 \pm 16% (+1.0°C), 65 \pm 10 % (+1.5.°C), and 77 \pm 8 % (+2.0°C) of their 2020 area (Fig. 2b). Note that, for the +1.0°C temperature target, the projected volume and area losses are only slightly higher than the committed loss, i.e. the loss obtained by assuming the climatic conditions to stabilize at the 1988-2017 levels. Indeed, Zekollari et al. (2019) estimated such committed loss to be of 37 \pm 6 % for glacier volume and 35 \pm 7 % for the glacier area.

Figure 3a-c shows the regional distribution of the modelled glacier evolution. By 2100, only 1,484 \pm 324 (+1.0°C), 1,033 \pm 253 (+1.5°C) and 724 \pm 201 (+2.0°C) of today's roughly 4,000 glaciers are anticipated to remain, i.e. to have a volume >0 by then. In the three scenarios, Grosser Aletschgletscher (Fig. 3d), the largest glacier of the Alps, might lose as much as 45 \pm 12 %, 64 \pm 11 %, and 77 \pm 6 % of its 2020 volume, respectively. Rhonegletscher (Fig. 3e), one of the most studied and visited glaciers, is projected to lose 55 \pm 24 %, 78 \pm 11 %, and 91 \pm 7 % of its volume in the same time span, whilst values for other glaciers are found in Table S1. Our results confirm that global warming can be significantly amplified at the regional scale, and that

glaciers in the European Alps sensitively respond to even small levels of global atmospheric warming, such as a global air
110 temperature change of $+1.0^{\circ}\text{C}$. Most importantly, however, our results show significant differences between the future glacier
evolution projected for the three warming scenarios, indicating that any effort to further limit warming will have important
effects on glaciers and, thus, the Alpine environment.

Future effects notably include the water availability from presently glacierized areas. Our results indicate that, on a yearly
basis, glacier runoff for the period 2080-2100 might decrease by as much as $25 \pm 6\%$ ($+1.0^{\circ}\text{C}$), $32 \pm 8\%$ ($+1.5^{\circ}\text{C}$), and
115 $36 \pm 10\%$ ($+2.0^{\circ}\text{C}$) compared to 2000-2020 levels (Fig. 2c). The decrease follows above-average glacier melt during the
last decades, and indicates that peak glacier discharge already occurred in the past for many glaciers in the European Alps
(consistent with the findings of e.g. Huss and Hock, 2018). Our model results also show a transition in the timing of the
maximum glacier discharge during the year: whilst it occurred in August historically (2000-2020), it is expected to occur in
July ($+1.0$ and $+1.5^{\circ}\text{C}$ scenarios) or even June-July ($+2.0^{\circ}\text{C}$) by 2080-2100 (Fig. 2d). Finally, our simulations project the
120 2080-2100 August runoff to be $36 \pm 15\%$ ($+1.0^{\circ}\text{C}$), $44 \pm 14\%$ ($+1.5^{\circ}\text{C}$), and $55 \pm 15\%$ ($+2.0^{\circ}\text{C}$) lower on average than it
was in 2000-2020. This shift in the magnitude and timing of glacier runoff (i) is similar to what projected in earlier studies
(e.g. Huss and Hock, 2018), (ii) is a direct consequence of both the reduced glacier volumes (which causes the annual runoff
to decrease) and the increased temperature (which causes snow and ice melt to occur earlier in the year), and (iii) will have
important consequences for water availability (Huss and Hock, 2018), particularly during dry summer months. For the densely
125 populated European Alps and the areas downstream, such changes will not only impact ecosystems but also a number of water
users, including agriculture or hydropower operators for instance (Beniston et al., 2018).

In line with the bulk of the literature, the above results all refer to the time horizon until 2100. To gain insights into glacier
evolution beyond this horizon, we run GloGEMflow with three GCM members that provide climate data until 2300 and that
project mean global temperature changes below $+2.0^{\circ}\text{C}$ for 2300 (see Fig. S3). Note that one of these GCM members was
130 already considered in the simulations until 2100, whilst the remaining two were not because they show a warming beyond
 $+2.0^{\circ}\text{C}$ for 2100 (Fig. S3). The three members all come from SSP126 but stem from three different GCMs (see Fig. S3).

These results show that, owing to slow lowering of air temperatures and enhanced precipitation implied by this particular
scenario after 2100, slow glacier recovery might happen (Fig. S3). While glacier volume losses of 80-85% are calculated for
2100, the experiment projects Alpine glaciers to re-gain a part of this volume, reaching a total volume between 28% and 53%
135 of the 2020 level by 2300. Although this result is only based on three GCM members and is thus very uncertain, it suggests
that considering projections beyond 2100 might change the current perception of a possibly irreversible glacier loss. From the
physical point of view, the result that glaciers might re-grow after a potential cooling global temperatures is not surprising.
Still, increasing the number of GCM members that consider such longer-term horizons and having different research groups
performing similar analyses would help verifying the robustness of this preliminary finding. We also stress that decisive climate
140 action would be required for steering global temperatures towards such an evolution (i.e. SSP126).




Figure 2. Modelled evolution of total glacier (a) volume, (b) area, (c) annual runoff, and (d) monthly runoff. The total refers to all glaciers of the European Alps. Time series in c are smoothed with a 30-year running mean. In all panels, the thick line represents the mean and the transparent band corresponds to one standard deviation of the results obtained by forcing GloGEMflow with the selected GCM members. The numbers of GCM members is given (n) in panel (a).

4 Conclusions

Whilst there is overwhelming consensus that decisive action has to be taken to limit unwanted consequences of ongoing climatic change (UN, 2015; IPCC, 2018; IPBES, 2019; IPCC, 2019; WEF, 2020), the debate around which temperature targets to pursue is all but settled. In this contribution, we quantified the impact that a global temperature change of +1.0°, +1.5° or
 145 +2.0°C compared to pre-industrial levels would have on glaciers of the European Alps. We show that this would correspond to an Alpine warming slightly above global levels and to a moderate increase in precipitation. Under such conditions, Alpine glaciers are projected to lose $44 \pm 21\%$, $68 \pm 12\%$, and $81 \pm 8\%$ of their 2020 ice volume by 2100, indicating that even in




Figure 3. Relative change in glacier volume between 2020 and 2100 for scenarios of (a) +1.0 °C, (b) +1.5 °C and (c) +2.0 °C of global temperature increase. Each dot represents one glacier, the dot's size being proportional to the glacier area as given by the Randolph Glacier Inventory. The labels "GA" and "Rh" identify "Grosser Aletschgletscher" and "Rhonegletscher", respectively. The modelled geometries of these two glaciers are shown in panels (d and e) at inventory date (dashed), for the year 2020 (grey), and for the year 2100 under the three selected scenarios (colours).

the most moderate scenario (+1.0 °C), about half of the present-day glacier volume will be lost by the end of the century. Preliminary results based on a strong mitigation scenario (SSP126) and running until 2300 indicate that slow recovery might happen after that, emphasizing the interest in considering projections that reach beyond the 21st century. The changes in glacier volume will strongly impact the water yield from presently glacierized surface, with 2080-2100 annual average runoff decreasing by $25 \pm 6\%$ (for a global warming of +1.0 °C), $32 \pm 8\%$ (+1.5 °C) and $36 \pm 10\%$ (+2.0 °C) compared to 2000-2020 levels. Changes in monthly runoff – with a runoff peak projected to occur 1 to 2 months earlier by the end of the century – will be even more pronounced, with August peak runoff reductions of $36 \pm 15\%$, $44 \pm 14\%$, and $55 \pm 15\%$, respectively. For glacierized catchments and the areas downstream, these changes in water availability can be expected to have important consequences ranging from the functioning of ecosystems to the generation of hydro-electricity. Even if our preliminary results based on a strong mitigation scenario indicate that slow glacier recovery might happen after the 21st century, we hope that the presented results will contribute in making the point that, when it comes to global climate, every half-degree counts.

Code availability. Code and data availability upon request

160 *Author contributions.* DF, LC, SE conceived the study. LC and SE performed the numerical modelling with support from MH, HZ and DF. The original code was developed by MH (mass balance component) and by HZ (ice flow component). SE retrieved, homogenized and analyzed all CMIP6 GCMs, with support of LC, MH and DF. LC and DF wrote the manuscript and produced the figures, with contributions from all other co-authors.

Competing interests. There are no competing interests.

165 *Acknowledgements.* We are grateful to the Swiss National Science Foundation, and for the funding provided for project Nr. 184634 in particular. HZ acknowledges the funding from a Marie Skłodowska-Curie Individual Fellowship (Grant 799904). We thank the RGI consortium for the global glacier inventory data. We acknowledge the ECA&D project for the E-OBS dataset, and CMIP for the GCM outputs. We also thank the WGMS for providing mass balance and length change measurements.

References

- 170 Beniston, M., Farinotti, D., Stoffel, M., Andreassen, L. M., Coppola, E., Eckert, N., Fantini, A., Giacona, F., Hauck, C., Huss, M., et al.: The European mountain cryosphere: a review of its current state, trends, and future challenges, *The Cryosphere*, 12, 759–794, <https://doi.org/10.5194/tc-12-759-2018>, 2018.
- Compagno, L., Zekollari, H., Huss, M., and Farinotti, D.: Limited impact of climate forcing products on future glacier evolution in Scandinavia and Iceland, *Journal of Glaciology*, pp. 1–17, <https://doi.org/10.1017/jog.2021.24>, 2021.
- 175 Cornes, R. C., van der Schrier, G., van den Besselaar, E. J. M., and Jones, P. D.: An ensemble version of the E-OBS temperature and precipitation data sets, *Journal of Geophysical Research: Atmospheres*, 123, 9391–9409, <https://doi.org/10.1029/2017JD028200>, 2018.
- Farinotti, D., Huss, M., Fürst, J. J., Landmann, J., Machguth, H., Maussion, F., and Pandit, A.: A consensus estimate for the ice thickness distribution of all glaciers on Earth, *Nature Geoscience*, 12, 168–173, <https://doi.org/10.1038/s41561-019-0300-3>, 2019.
- Grab, M., Mattea, E., Bauder, A., and Huss, M.: Ice thickness distribution and bedrock topography of all Swiss glaciers based on ground penetrating radar, *Journal of Glaciology*, 0, 0000–0000, in press.
- 180 Hock, R.: Temperature index melt modelling in mountain areas, *Journal of Hydrology*, 282, 104–115, [https://doi.org/10.1016/S0022-1694\(03\)00257-9](https://doi.org/10.1016/S0022-1694(03)00257-9), 2003.
- Huss, M. and Hock, R.: A new model for global glacier change and sea-level rise, *Frontiers in Earth Science*, 3, 54, <https://doi.org/10.3389/feart.2015.00054>, 2015.
- 185 Huss, M. and Hock, R.: Global-scale hydrological response to future glacier mass loss, *Nature Climate Change*, 8, 135–140, <https://doi.org/10.1038/s41558-017-0049-x>, 2018.
- Huss, M., Jouvet, G., Farinotti, D., and Bauder, A.: Future high-mountain hydrology: a new parameterization of glacier retreat, *Hydrology and Earth System Sciences*, 14, 815–829, <https://doi.org/10.5194/tc-12-759-2018>, 2010.
- IPBES: Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services, E.S. Brondizio, J. Settele, S. Díaz, and H.T. Ngo (eds.), IPBES secretariat, Bonn, Germany., <https://www.ipcc.ch/srocc/download/>, 2019.
- 190 IPCC: Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty, V. Masson-Delmotte, P. Zhai, H.-O. Pörtner, C. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J.B.R. Matthews, Y. Chen, X. Zhou, M.I. Gomis, E. Lonnoy, T. Maycock, M. Tignor, and T. Waterfield (eds.), Cambridge University Press, <https://www.ipcc.ch/sr15/download/>, 2018.
- 195 IPCC: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate, H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, N.M. Weyer (eds.), 2019.
- Marzeion, B., Kaser, G., Maussion, F., and Champollion, N.: Limited influence of climate change mitigation on short-term glacier mass loss, *Nature Climate Change*, 8, 305–308, <https://doi.org/10.1038/s41558-018-0093-1>, 2018.
- 200 Marzeion, B., Hock, R., Anderson, B., Bliss, A., Champollion, N., Fujita, K., Huss, M., Immerzeel, W. W., Kraaijenbrink, P., Malles, J., Maussion, F., Radić, V., Rounce, D. R., Sakai, A., Shannon, S., Wal, R., and Zekollari, H.: Partitioning the Uncertainty of Ensemble Projections of Global Glacier Mass Change, *Earth's Future*, 8, <https://doi.org/10.1029/2019EF001470>, <https://onlinelibrary.wiley.com/doi/10.1029/2019EF001470>, 2020.

- 205 Paul, F., Rastner, P., Azzoni, R. S., Diolaiuti, G., Fugazza, D., Le Bris, R., Nemeč, J., Rabatel, A., Ramusovic, M., Schwaizer, G., et al.: Glacier shrinkage in the Alps continues unabated as revealed by a new glacier inventory from Sentinel-2, *Earth System Science Data*, 12, 1805–1821, <https://doi.org/10.5194/essd-12-1805-2020>, 2020.
- RGI Consortium: Randolph Glacier Inventory 6.0, <https://doi.org/10.7265/N5-RGI-60>, 2017.
- UN: United nations Paris agreement, United nations framework convention on climate change, Rio de Janeiro, Brazil, and New York, United States, https://unfccc.int/sites/default/files/english_paris_agreement.pdf, 2015.
- 210 Vautard, R., Gobiet, A., Sobolowski, S., Kjellström, E., Stegehuis, A., Watkiss, P., Mendlik, T., Landgren, O., Nikulin, G., Teichmann, C., et al.: The European climate under a 2 C global warming, *Environmental Research Letters*, 9, 034006, <https://doi.org/10.1088/1748-9326/9/3/034006>, 2014.
- WEF: The Global Risks Report 2020, 15th Edition, Tech. rep., World economic forum, Cologne/Geneva, Switzerland, http://www3.weforum.org/docs/WEF_Global_Risk_Report_2020.pdf, 2020.
- 215 WGMS: Fluctuations of glaciers database, <https://doi.org/10.5904/wgms-fog-2020-08>, 2020.
- Zekollari, H., Huss, M., and Farinotti, D.: Modelling the future evolution of glaciers in the European Alps under the EURO-CORDEX RCM ensemble, *The Cryosphere*, 13, 1125–1146, <https://doi.org/10.5194/tc-13-1125-2019>, 2019.
- Zemp, M., Huss, M., Thibert, E., Eckert, N., McNabb, R., Huber, J., Barandun, M., Machguth, H., Nussbaumer, S. U., Gartner-Roer, I., Thomson, L., Paul, F., Maussion, F., Kutuzov, S., and Cogley, J. G.: Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016, *Nature*, 568, 382–386, <https://doi.org/10.1038/s41586-019-1071-0>, 2019.
- 220