

Rectification and validation of a daily satellite-derived Antarctic sea ice velocity product

Deleted: : impacts on ice kinematics

Tian R. Tian^{1,2}, Alexander D. Fraser^{2,1}, Noriaki Kimura³, Chen Zhao^{2,1}, Petra Heil^{4,2}

¹Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, TAS 7001, Australia

5 ²Australian Antarctic Program Partnership, Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, TAS 7001, Australia

³Atmosphere and Ocean Research Institute, The University of Tokyo, Kashiwa, Japan

⁴Australian Antarctic Division, 203 Channel Highway, Kingston, TAS 7050, Australia

Correspondence to: Tian R. Tian (tian.tian@utas.edu.au)

10 **Abstract.** Antarctic sea ice kinematic plays a crucial role in shaping the [Southern Ocean](#) climate and ecosystems. Satellite
passive microwave-derived sea ice motion data have been used widely for studying sea ice motion and deformation, and
provide daily, global coverage at a relatively low spatial-resolution (in the order of 60 [km](#) × 60 km). In the Arctic, several
validated data sets of satellite observations are available and used to study sea ice kinematics, but far fewer validation studies
exist for the Antarctic. Here, we compare the widely-used passive microwave-derived Antarctic sea ice motion product by
15 Kimura et al. (2013) with buoy-derived velocities, and interpret the effects of satellite observational configuration on the
representation of Antarctic sea ice kinematics. We identify two issues in the Kimura et al. (2013) product: (i) errors in two
large triangular areas within the eastern Weddell Sea and western Amundsen Sea relating to an error in the input satellite
data composite, and (ii) a more subtle error relating to invalid assumptions for the average sensing time of each pixel. Upon
rectification of these, performance of the daily composite sea ice motion product is found to be a function of latitude, relating
20 to the number of satellite swaths incorporated (more swaths further south as tracks converge), and the heterogeneity of the
underlying satellite signal (brightness temperature here). Daily sea ice motion vectors calculated using ascending- and
descending-only satellite tracks (with a true ~24 h time scale) are compared with the widely-used combined product
(ascending and descending tracks combined together, with an inherent ~39 h time scale). This comparison reveals that
kinematic parameters derived from the shorter time scale velocity datasets are higher in magnitude than the combined
25 dataset, indicating a high degree of sensitivity to observation time scale. We conclude that the new generation of “swath-to-
swath” (S2S) sea ice velocity datasets, encompassing a range of observational time scales, is necessary to advance future
research into sea ice kinematics.

1 Introduction

1

30 Antarctic sea ice plays a central role for Southern Ocean climate and ecosystems. Sea ice is a crucial component in the climate system due to its role in controlling the transfer of energy, gases and moisture between the polar ocean and atmosphere (Walsh, 1983), impact on climate variability through the ice-albedo feedback (Ebert and Curry, 1993), and on deep water formation as well as fresh water budget by modifying the upper ocean salinity and density gradient during ice growth and melt (Goosse et al., 1997; Yang and Neelin, 1993). [Because of its](#) relatively high albedo compared to open water, especially when covered by snow, [sea ice](#) reflects a large portion of the incoming shortwave radiation, thus strongly influencing the Earth's surface energy budget as well as the atmospheric thermo-circulation (Curry et al., 1995). Furthermore, sea ice production and melt modify the near-surface water, e.g., ice production in coastal polynyas may initiate the formation of dense shelf water, the precursor of Antarctic Bottom Water, one of the densest water masses in the global circulation (Rintoul et al., 2001). Sea ice also plays an important role in marine ecosystems and biogeochemical cycles, as a unique habitat for primary production (Arrigo and Thomas, 2004), and is affected by a network of complex chemical, biological and physical interactions (Dieckmann and Hellmer, 2010). Hence, understanding Antarctic sea ice physical processes is critical across many disciplines.

Sea ice kinematics describes sea ice motion and deformation. Accurate characterization of sea ice kinematics is crucial to investigate modifications of the polar sea ice cover and sea ice mass balance (Kwok, 2011). Sea ice motion is driven by external oceanic and atmospheric forcing as well as internal stresses (Kottmeier and Sellmann, 1996). It is also impacted by physical boundary conditions, e.g., coastline, icebergs, or ice tongues, as well as material properties, e.g., thickness, concentration, and strength (Heil et al., 2011). Sea ice deformation is defined as the spatial gradients in ice velocity driven by winds, currents, and internal stress (Kirwan, 1975; Heil et al., 1998; Marsan et al., 2004). Sea ice deformation may change the drag between ocean and air (Spreen et al., 2017), as well as impacting the distribution of ice thickness and strength (Marsan et al., 2004).

Sea ice kinematics has been studied extensively in the Arctic using satellite observations (e.g., Weiss and Marsan, 2004; Hutchings et al., 2005; Spreen et al., 2017; Hutter et al., 2018). However, satellite-based research of Antarctic sea ice kinematics has been much more limited (Emery et al., 1991; Giles et al., 2011; Hakkinen, 1995; Heil et al., 1998, 2006, 2009; Kimura, 2004; Kimura et al., 2013). In order to characterize Antarctic sea ice kinematics and its effects on physical ice properties accurately, more research on sea ice deformation processes is required (Hutter et al., 2018). With the development of remote sensing technology, satellite observations of Antarctic sea ice motion have been improved in recent years (Spreen et al., 2017; Lavergne et al., 2021). In particular, passive microwave (PM)-derived sea ice motion products give daily, Antarctic-wide coverage with a gridded data format, which may be suitable for characterizing large-scale sea ice kinematics and evaluating basin-scale sea ice simulations (Sumata et al., 2014). However, detailed exploration of this kind of composite satellite observations is required. The spatial resolution of PM-derived sea ice products (25 - 62.5 km resolution) is relatively low compared to that provided by synthetic aperture radar (SAR) sea ice motion products ([150 m resolution for ESA's](#)

[Sentinel1A/B](#)), and complete validation of PM-derived motion datasets has yet to be achieved, owing to coastal
65 contamination, numerous missing data and artifacts, plus limitations in coverage of buoy deployments necessary for
complete characterization and validation (Heil et al., 2006; Sumata et al., 2014; Szanyi et al., 2016).

The procedure most widely applied for producing daily sea ice motion vectors is the Maximum Cross-Correlation (MCC)
algorithm on satellite-derived daily composites of microwave brightness temperature (T_B) or backscatter coefficient (Ninnis
70 et al., 1986; Emery et al., 1991; Kimura et al., 2013). The MCC method determines the spatial difference between image
subsets in two consecutive T_B or backscatter coefficient composite maps by maximizing the cross-correlation coefficient
between two composite images of T_B or backscatter coefficient. Sea ice motion products derived from daily-averaged maps
of the satellite imagery using MCC algorithm have been referred to as “daily-map” (DM) products ((Lavergne et al., 2021).
DM products typically have a regular nominal time scale of approximately one or two days, hence potentially bias the
75 estimated sea ice kinematic parameters by not considering a variety of time scales (Lavergne et al., 2021). Polar orbiting
satellite swath convergence means a number of swaths are merged (typically by per-pixel mean calculation) to produce
composite maps of satellite signals (T_B or radar backscatter), which precipitates a reduction in accuracy of sea ice motion
vectors ((Lavergne et al., 2021). However, due to the advantages of a) regular large-scale coverage with a consistent grid, b)
relative ease of calculation, and c) relative ease of use and interpretation for the dataset end user, DM products have been
80 widely used for recent studies of sea ice kinematics (Szanyi et al., 2016; Hutter et al., 2018). Several PM-derived DM
products have been used for investigating sea ice motion in both Arctic and Antarctic, including the Ocean and Sea Ice
Satellite Application Facility (OSI SAF) sea ice drift product OSI-405-c (Lavergne, 2016), henceforth referred to as the OSI
SAF product; the NASA Polar Pathfinder Daily 25 km Equal-Area Scalable Earth Grid (EASE-Grid) Sea Ice Motion
Vectors product (Tschudi et al., 2019) from the National Snow and Ice Data Center (NSIDC), henceforth referred to as the
85 NSIDC product; and the daily sea ice motion product produced by Kimura et al. (2013), henceforth referred to as the
KIMURA product.

Daily input data of the KIMURA product include both the horizontal and vertical polarization channel T_B images for both
ascending (ASC) and descending (DES) orbital sections (i.e., four independent images) from the NASA’s Advanced
90 Microwave Scanning Radiometer - Earth Observing System (AMSR-E) and Advanced Microwave Scanning Radiometer 2
(AMSR2), on the sun-synchronous Global Change Observation Mission – Water 1 (GCOM-W1) platform. The final
KIMURA daily sea ice velocity product is then derived as the mean value of these four datasets. AMSR2, used by the
KIMURA product after 2012, passes over each [observed](#) point on the Earth surface at approximately the same local (solar)
time, and for any given pixel, the sensing times for ASC and DES passes differ. The local overpass time of the ASC pass in
95 the Antarctic sea ice zone (i.e., $\sim 60^\circ$ to 78° S) is around local 15:00 (Fig.1, red bar), and the DES pass is around local
midnight (Fig. 1, blue bar). At higher latitudes where several adjacent swaths observe the same pixel, the local times of
observation for these several swaths are centered around the nominal overpass time, but offset by several orbital periods (i.e.,

Deleted: motion

offset from the nominal local time by $\pm n \times 98$ minutes, where n can be 1, 2 or more depending on latitude). The daily
 100 combined KIMURA sea ice motion product (represented by the green bar in Fig. 1) is the mean of the ASC and DES data.
 As such, it has a true (nominal) time base of approximately 39 h. This exceeds that of the ASC and DES motion products
 (which have a nominal time base of 24 h). Previous buoy-based research demonstrated that shorter time scale observations
 can represent sea ice motion and kinematic parameters with higher magnitude than longer time scale observations (Hoerber
 and Gube-Lenhardt, 1987; Heil and Allison, 1999), but analogous research for satellite-based observational datasets is quite
 105 limited. We postulate that a shorter time scale sea ice motion dataset (i.e., derived from ASC or DES DM composites, with
 their inherent ~ 24 h time scale) will result in higher estimates of the kinematic parameter magnitude compared to a longer
 time scale dataset (i.e., a product combining ASC and DES data). Thus, to investigate the influence of the time scale on the
 accuracy of the PM-derived sea ice motion and derived kinematic parameter magnitude, individual ASC and DES KIMURA
 datasets are generated and investigated here.

110




Figure 1. Representation of the time scale of the combined (ASC and DES averaged together) KIMURA daily sea ice motion product, as well as the time scales of ASC and DES components.

115 Recently sea ice motion has been estimated from PM-derived T_B partial-overlap swath pairs from a wide range of temporal
 baselines (from one orbit of ~ 90 minutes, to 3 days). This approach is referred as “swath-to-swath” (S2S) (Lavergne et al.,
 2021). For S2S, the time base for derived motion is a function of the temporal separation of individual swath in each pair,
 hence differs between pairs (Lavergne et al., 2021). As such, the raw S2S-derived motion swath do not constitute a regular,
 full coverage time series. However, since S2S most accurately provides ice motion from individual swaths across a range of
 120 time bases, Lavergne et al. (2021) concluded that sea ice motion based on S2S is more accurate than that from DM.

We hypothesize that a greater number of satellite swaths used for averaged maps of satellite signals (e.g., T_B DMs) may
 negatively impact the performance of PM-derived sea ice motion vectors. Here, we investigate this hypothesis by performing
 comparisons against drifting buoy-derived motion in different latitude ranges with the expectation that DM-based KIMURA
 product performs better at lower latitudes, where swath overlap averaging is lower. We note that the OSI SAF sea ice motion
 125 product is derived by averaging over two days to increase the signal-to-noise ratio (SNR) of this operational product
 (Lavergne, 2016). This extended averaging interval arises, in part, due to OSI SAF integrating multiple sensors. Because of
 the extended time base of the OSI SAF product, direct comparison with the NSIDC and KIMURA product in a kinematic
 parameter context is not possible. Hence the OSI SAF dataset is not considered further in this study, although implications of

130 a longer temporal baseline are considered in the discussion. Version 3 of NSIDC dataset is known to display a high degree of autocorrelation in neighboring vectors (Szanyi et al., 2016) which would be detrimental to the derivation of kinematic parameters. Version 4 of the NSIDC product has reduced this autocorrelation to some extent (Tschudi et al., 2019), however this product still assimilates from several instruments, so retrieval of kinematic parameters will suffer from cross-instrument averaging. Furthermore, as the NSIDC product does not use AMSR2, spatial resolution is inherently lower than KIMURA product. For these reasons, the NSIDC dataset is not considered further here. [A comparison of the KIMURA, OSI SAF and NSIDC sea ice motion products against buoy observations is provided in Appendix A.](#) The KIMURA product is the only PM-derived dataset we consider here; it provides daily gridded sea ice motion vectors of the complete Southern Ocean at 60 km horizontal resolution.

140 The objectives of this study are to a) evaluate the KIMURA ice motion product separately for the ASC, DES and combined datasets by comparison with coincident drifting buoy-derived velocities, and b) investigate impacts of the observational time scale of the ice motion products on representation of Antarctic sea ice kinematic magnitude. This study will provide characterization of the satellite sensor configuration and observational time scale on the accuracy of derived sea ice motion, as well as the ability to represent Antarctic sea ice kinematic parameters.

145 2 Datasets

2.1 PM-derived sea ice motion

The KIMURA sea ice motion products (Kimura et al., 2013) used here were produced at 60 km×60 km resolution on a regular 145×145 pixel grid covering the entire Southern Ocean, with daily data from 2012-06-23 to 2020-06-01. These were calculated by applying the MCC method to 36 GHz, 10 km resolution AMSR2 T_B images (both vertical and horizontal polarization, ASC and DES) from the Arctic Data archive System (ADS) of the Japanese National Institute of Polar Research (NIPR). [The MCC method determines the spatial offset that maximizes the cross-correlation coefficient between two \$T_B\$ arrays \(at a resolution of 10 km per px\) in consecutive images \(from an ASC observation to the next ASC observation, or DES to DES\).](#) For this procedure, 6×6 grid cell (60 km×60 km) arrays are utilized. At first, ice motion vector quantized to an integer number of grid intervals (10 km) is computed. Then detections of false cross-correlation matches are automatically filtered out by the following procedure: (1) eliminate vectors with a correlation coefficient of less than 0.5 of the MCC coefficient; (2) retain the “reliable” vectors with more than 0.78 of the correlation coefficient, i.e. within the 95% confidence interval; (3) vectors with a correlation coefficient of between 0.5 and 0.78 of the MCC coefficient are eliminated if they differ from the “reliable” vectors (within a search radius of 100 km) by more than 23 cm/s (i.e. 2 pixels per day). Finally, filtered sea ice motion vectors, on a 10 km×10 km grid, are organized into a 60 km×60 km resolution dataset. In this procedure, at least six vectors are utilized for each resulting vector. If the number of the available vectors is less than six, that grid is regarded as missing data. The final KIMURA combined daily dataset is calculated as the mean value of four datasets

derived from the horizontal and vertical polarization channel images from ASC and DES satellite passes. By combining the ASC and DES passes in this fashion, the true characteristic time scale becomes approximate 39 hours, as described in Figure 1. In this work, we also produce and analyze daily sea ice motion vectors computed for the shorter time scale (true 24 h) ASC and DES swath composites.


2.2 In situ buoy-derived sea ice motion

Velocities derived from buoy positions are suitable for validating satellite-derived sea ice motion (Hoerber and Gubelhardt, 1987; Heil and Allison, 1999). Here we evaluate the three KIMURA datasets using three buoys deployed on ice floes in the Weddell and Ross seas (Table 1 and Fig. 2). These have been named the southern Weddell Sea buoy (Fig. 2, red), the central Weddell Sea buoy (Fig. 2, green) and the Ross Sea buoy (Fig. 2, orange). In the Weddell Sea, several drifting sea ice buoys were deployed in 2018 (Schröder, 2018). Only two of these are analyzed in this paper as the buoys were deployed in meso-scale arrays, hence at sub-grid scale to the KIMURA data. Others in this deployment were near the coast where the KIMURA ice-motion product suffers from coastal contamination. Raw positions were quality-controlled and any erroneous data removed. A weighted polynomial interpolator (Scargle, 1982; Heil et al., 2001) was applied to the quality-controlled position measurements to yield an equi-temporal hourly time series. In the Ross Sea, a total of eight position-only buoys were deployed as part of the *Polynyas, Ice Production, and seasonal Evolution in the Ross Sea (PIPERS)* cruise during austral autumn of 2017 (Tison et al., 2020). Four were excluded due to coastal proximity, and another one due to a short time series. Here we analyze only one of the three remaining buoys as they shared similar trajectories.

Table 1. Buoy deployment information

Buoy location	Buoy ID	Latitude Coverage	Active interval
Southern Weddell Sea	ACED-01	74.5° – 75.5° S	2018-04-04 – 2018-07-04
Central Weddell Sea	ACED-03	71.1° – 75.5° S	2018-04-08 – 2018-08-20
Ross Sea	sbd79220	65.0° – 73.4° S	2017-05-20 – 2017-11-26

Deleted: The Kimura et al. (2013) dataset combines both the daily ASC and DES sea ice motion datasets by taking their average. ...



185

Figure 2. Mean daily number of AMSR2 swaths for (a) ASC and (b) DES passes. Trajectories of three buoys use in this study are also shown.

3 Methods

3.1 Validation of KIMURA-derived ice motions datasets using buoy data

190 To evaluate the skill of the three KIMURA datasets to represent sea ice motion, we compare these with buoy-derived sea ice velocity, using three validation metrics: Pearson's correlation coefficient, regression slope and RMS deviation (RMSD). Since AMSR2 is in a sun-synchronous orbit, it passes over each point on the Earth at roughly the same local (solar) time. For daily motion comparison buoy time stamps, referenced in UTC time, are converted to local (solar) time, to investigate the start and end time for daily motion comparison. To calculate daily buoy velocity, a 24-hour interval of buoy positions is selected, to coinciding with the satellite overpass time. The choice of the start time for a 24 h interval is informed by calculating all validation metrics in hourly intervals from -48 h (local midnight of two days before) to +48 h (local midnight of two days later). The ASC, DES and combined start times which are found to maximize the correlation between KIMURA and buoy sea ice velocity were fixed for all analyses. Spatially, KIMURA pixels contributing to the daily velocity were selected by weighting the velocity in the pixels that the buoy passed through in the preceding 24 h.

200 We note that the number of AMSR2 swaths (Fig. 2, blue shading) increases rapidly poleward of 74° S for both ASC and
DES passes. This is particularly evident in the ASC case, where a strong discontinuity is observed at this latitude, owing to
the oblique, forward-looking viewing geometry of the instrument. In both ASC and DES cases, more satellite swaths will be
merged together at higher latitude (i.e., a higher degree of temporal "smearing" is expected at higher latitudes, with potential
detriment to derived motion products). By extension, fewer swaths will be merged at lower latitudes, (i.e., at northern
205 extremes, the DM product tends toward the S2S product, with reduced temporal smearing). Based on the results of Lavergne
et al (2021), we expect better performance of the KIMURA product at lower latitudes for this reason. To investigate the
effect of compositing more satellite swaths on the accuracy of the KIMURA product, buoy comparisons are separated based
on their latitude (Table 1). We separate the central Weddell Sea buoy to four 1° latitude segments and the Ross Sea buoy to
four 2° latitude segments to support this analysis (the southern Weddell Sea buoy has insufficient latitudinal range for this
210 analysis).

3.2 Calculation of sea ice divergence

Various differential kinematic parameters (DKPs) are frequently used to represent sea ice kinematics (Molinari and Kirwan,
1975; Hutchings et al., 2012): divergence (D), shear deformation rate (pure shear), normal deformation rate (stretching) and
vorticity (rotation). Here we use D to investigate a hypothesized increase in DKP magnitude when considering the shorter
215 (24 h) time scale ASC and DES datasets, compared to the combined dataset. To generate daily maps of D from the
KIMURA product, D at the center of four neighboring pixels was calculated as:

$$D = \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y}, \quad (1)$$

where u and v are daily sea ice velocities in the eastward and northward directions, respectively, while x and y represent
eastward and northward components. We evaluate the magnitude of D for the combined, ASC and DES datasets by
220 computing the monthly RMS deviation (hereafter RMSD, i.e., a measure of monthly variability). Variability of D over a
period of one month is used here to assess the relative magnitude of the DKPs. Furthermore, based on the latitude hypothesis
that more satellite swaths become merged causing more smoothing at higher latitudes, we also expect that D RMSD will
decrease from north to south.

4 Results

225 4.1 KIMURA data assessment

Here, we validate the three KIMURA sea ice motion datasets using daily velocity from buoys in the Ross and Weddell seas. First, we report some problems discovered in the KIMURA datasets.

4.1.1 Problem 1: Triangular regions with persistent low velocity

230 Initial inspection of the KIMURA product reveals the existence of two triangular regions with recurring low velocity (every second day). These are in the western Amundsen Sea for the ASC dataset (triangular region in Fig. 3a) and the Weddell Sea for the DES dataset (triangular region in Fig. 3c). These problem regions are also evident in the combined product, although their impact is masked (halved) due to the averaging of ASC and DES velocities. Upon investigation, this occurs at an acquisition time of around midnight UTC, where swath duplication has erroneously occurred in the T_B DM product (from NIPR, ADS). Such areas with replicated values in the source T_B images manifest as areas of near-zero velocity in the
235 KIMURA product. To rectify this problem, we produced a new version of the KIMURA product using 10 km resolution AMSR2 Level-3 36 GHz T_B images from the Japan Aerospace Exploration Agency (JAXA) ASC and DES T_B daily composites which do not exhibit erroneous swath duplication. A new combined product is produced as the mean of these two with the same 60 km horizontal grid resolution. Rectified KIMURA velocity maps (Fig. 3b and d) reveal that this problem
240 has been overcome. We note that the sea ice velocity in other regions changes slightly, which is due to small differences between the NIPR and JAXA T_B DM composites, but investigation is beyond the scope of this work.

Deleted: motion




Figure 3. Sea ice speed of (a) the original KIMURA DES dataset (produced from NIPR T_B composite), and (b) the rectified KIMURA DES dataset (produced from JAXA T_B composite) on 2017-09-15. (c) and (d): as for (a) and (b), but for ASC dataset on 2017-09-14. Arrows represent ice motion vectors, which are plotted as averaged 5×5 grid cells (300×300 km resolution) to enhance visibility. The triangular areas shown in red solid lines in (a) and (c) mark the regions of low velocity.

4.1.2 Problem 2: Incorrect assumption of 24 h time separation

The work described above revealed a second problem with the KIMURA product: an interval of exactly 24 h time separation between consecutive T_B composites was assumed, which is rarely correct. We analyze actual time separation between each

250 data pair collected on contiguous days. In order to characterize the variability of time separation, we calculate the standard deviation of time separation for both ASC and DES datasets. The result for September 2017 is displayed in Fig. 4 (all months similar, not shown here).

The mean time separation is close to 24 h throughout each month, however there is a region of high standard deviation around midnight UTC, which occurs in the Amundsen Sea for the ASC swaths (Fig. 4a) and the eastern Weddell Sea for the
255 DES swaths (Fig. 4b). This indicates that the actual time separation of pixels in subsequent daily composites can regularly be up to 34 h, or as low as 14 h in these two areas. Thus, the resulting velocity is regularly incorrect by up to 40%, and the combined KIMURA dataset exhibits errors for these locations. However, any error is compensated by the following day's velocity field, since the time intervals oscillate around 24 h. As such, it will not have a strong effect on long-term sea ice motion, but is highly detrimental when deriving sea ice kinematic parameters. To fix this problem, the actual time separation
P60 needs to be taken into account when producing daily sea ice velocity of KIMURA ASC, DES and combined datasets. The appropriate correction has been performed for all remaining analyses presented here.

Deleted: motion




Figure 4. Standard deviation of time separation for AMSR2 (a) ASC and (b) DES datasets in September 2017. The shaded background represents AMSR2-derived sea ice concentration on 2017-09-01 (Melsheimer and Spreen, 2019).

265

In summary, two problems have been discovered in the original KIMURA product, and have been rectified in all KIMURA data used in this study. Henceforth, the corrected KIMURA product is referred to as the KIMURA_{new} product. The official release of the rectified KIMURA sea ice motion is a work in progress.

270 4.2 Validation of KIMURA_{new} product

The KIMURA_{new} product is compared to buoy-derived sea ice motion. The buoy start and end times needed to maximize correlation coincide exactly with the local (solar) times of corresponding satellite passes, i.e., 15:00, 0:00 and 8:00 UTC for ASC, DES and combined data, respectively, giving confidence in our methodology. Validation metrics are given in Table 2.

Table 2. Comparison between KIMURA_{new} dataset and buoys

275

Validation parameter	Component of velocity	KIMURA _{new} datasets	Ross Sea buoy (Oct 2017)	Central Weddell Sea buoy (July 2018)	Southern Weddell Sea buoy (May 2018)
Correlation coefficient	Eastward	ASC	0.92	0.94	0.92
		DES	0.90	0.92	0.80
		Combined	0.92	0.94	0.87
	Northward	ASC	0.94	0.97	0.97
		DES	0.95	0.95	0.95
		Combined	0.96	0.95	0.96
Regression slope	Eastward	ASC	0.98	0.95	0.98
		DES	0.97	0.87	0.99
		Combined	1.06	0.97	1.17
	Northward	ASC	0.78	0.83	1.07
		DES	0.91	0.91	1.10
		Combined	0.92	0.93	1.10
RMSD (m day ⁻¹)	Eastward	ASC	4.9×10^3	2.1×10^3	2.5×10^3
		DES	5.6×10^3	2.5×10^3	3.7×10^3
		Combined	5.0×10^3	2.0×10^3	3.3×10^3
	Northward	ASC	5.8×10^3	2.6×10^3	2.3×10^3
		DES	5.0×10^3	2.7×10^3	3.2×10^3
		Combined	4.4×10^3	2.6×10^3	3.0×10^3





Figure 5. Scatterplot of comparisons between KIMURA_{new} dataset (ASC is red, DES is blue and combined is green) and buoy velocities in eastward (a, b and c) and northward (d, e and f) directions, Oct 2017 for the Ross Sea buoy, July 2018 for Central Weddell Sea buoy, and May 2018 for Southern Weddell Sea buoy. The black solid line represents the regression slope of 1.0.

The validation metrics (Table 2) demonstrate that the KIMURA_{new} dataset represents buoy velocities well. The mean value of correlation coefficient of comparison between three KIMURA_{new} datasets and three buoys is 0.93, and of regression slope is 0.97. The monthly average sea ice motion speed is around 15.8 km day^{-1} for Ross Sea in October 2017, 9.7 km day^{-1} for central Weddell Sea in July 2018 and 8.3 km day^{-1} for southern Weddell Sea in May 2018. We notice that comparison of KIMURA_{new} with the Ross Sea buoy (Table 2) exhibits relatively large RMSD (from 4.4 to 5.8 km day^{-1}), considerably larger than that of the two Weddell Sea buoys (from 2.0 to 3.7 km day^{-1}), as well as larger than the validation of DM-derived sea ice motion RMSD (from 2.3 to 2.9 km day^{-1}) from Lavergne et al. (2021). The comparisons between KIMURA_{new} and buoys (Fig. 5) confirm this for sea ice in the Weddell Sea. Here the regression slopes are closer to 1.0, and point lie closer to the regression lines than for the Ross Sea comparison. This difference in RMSD prompted further investigation into the performance difference between the Ross and Weddell seas. Since the MCC algorithm requires discernible structure in the

underlying T_B imagery to produce accurate vectors, we suggest this RMSD discrepancy is caused by a differing degree of heterogeneity of T_B between the Ross Sea and Weddell Sea ice during these buoy deployments. To investigate why the
295 RMSD of the Ross Sea result is higher than that of the Weddell Sea, we analyze the spatial standard deviation of 36 GHz 10 km vertically- and horizontally polarized T_B (i.e., the input data to the MCC algorithm) in limited areas around both buoy deployments.




300 **Figure 6.** ASC, horizontally-polarized daily T_B composite acquired on 2018-07-01. Buoy trajectories are indicated by black lines in the Weddell and Ross seas. Square black boxes display the locations of subsets centered around the central Weddell Sea and Ross Sea buoys analyzed in Table 3.

305

310 **Table 3.** Standard deviation of T_B for subregions in the Weddell Sea and Ross Sea shown in Fig. 6. Time periods are the corresponding time of buoys within these subregions. Satellite orbits and polarizations are ASC-horizontal (ASC, H), ASC-vertical (ASC, V), DES-horizontal (DES, H), and DES-vertical (DES, V).

Region of box	Latitude and longitude of center of box	Time	Orbit and polarization	Std. dev. (K)
Weddell Sea	73.5° S, 40.3° W	2018-07-01 ~ 2018-07-31 (31 days)	ASC, H	6.7
			ASC, V	5.4
		DES, H	6.8	
		DES, V	5.5	
Ross Sea	72.0° S, 179.0° E	2017-07-01 ~ 2017-07-31 (31 days)	ASC, H	5.0
			ASC, V	2.3
		DES, H	4.8	
		DES, V	2.1	

315 Firstly, we select subregions of 31×31 T_B pixels (i.e., 310×310 km) that cover the Ross and Weddell Sea buoy areas (Fig. 6). Then we compute the standard deviation within each subregion using all data while the buoys remain within the subregion, and for all four combinations of satellite orbital section and polarization. Standard deviation of T_B in the Ross Sea is lower than the Weddell Sea in all satellite orbital sections and polarizations, with a mean of 3.6 K for Ross Sea and 6.1 K for the Weddell Sea. This indicates that the spatial variability of T_B within the sea ice of the Ross Sea is lower than that of the Weddell Sea (for these cases). The relative homogeneity in the underlying T_B data in the western Ross Sea, a region dominated by new ice formed in the Ross Ice Shelf and Terra Nova Bay polynyas, reduces the ability of the MCC to retrieve accurate sea ice motion, due to less discernible structure in the underlying T_B imagery (in agreement with Meier et al., 320 (2000), who found that errors are maximized in regions of new ice).



325 **Figure 7.** Comparison between KIMURA_{new} dataset (ASC is red, DES is blue and combined is green) and buoy velocity using (a, d) correlation coefficient, (b, e) regression slope and (c, f) RMSD. Vertical dashed lines in the central Weddell Sea panels (lower row) mark the latitude of the AMSR2 swath discontinuity (74° S) shown in Fig. 2.

330 To investigate the hypothesis that DM-based daily sea ice motion vector performance is latitude-dependent, we calculate the three validation metrics in different latitude ranges along the buoy trajectories (Fig. 7). The Southern Weddell Sea buoy is excluded from this analysis due to insufficient latitudinal range. We find that the validation metrics for both remaining buoys generally improve as they move further north (Fig. 7). For the central Weddell Sea buoy comparison, eastward velocity results (Fig. 7 lower row solid lines) improve from south to north across the swath number discontinuity (74° S), confirming our hypothesis that regions with fewer satellite swaths may yield improved performance for DM-based sea ice motion retrieval. However, northward ice velocity results in the central Weddell Sea case (dashed lines) exhibit a slight RMSD (Fig. 335 7f) deterioration from south to north. We attribute this unexpected deterioration to anisotropic T_B heterogeneity, noting that the heterogeneity in the northward component is lower than that in the eastward component (shown in Table 4), giving higher confidence in the result of the eastward component, and underscoring the importance of heterogeneous imagery when using the MCC algorithm.

340

Table 4. Eastward and northward components of standard deviation of T_B images in subregion of the central Weddell Sea buoy from July 2018, corresponding to the time buoy resided within the black box (Fig. 6). Satellite orbits and polarizations are ASC-horizontal (ASC, H), ASC-vertical (ASC, V), DES-horizontal (DES, H), and DES-vertical (DES, V).

Direction	Orbit and polarization	Std. dev. (K)
Eastward	ASC, H	3.3
	ASC, V	2.8
	DES, H	3.3
	DES, V	2.7
Northward	ASC, H	2.7
	ASC, V	2.3
	DES, H	2.7
	DES, V	2.3

345 The validation of the Ross Sea buoy (Fig. 7, upper row) indicates that both the eastward and northward performance improve progressively from south to north. The entire Ross Sea buoy trajectory is located to the north of the swath number discontinuity (i.e., in the relatively low swath number regime, as indicated in Fig. 2), so we suppose that the gradual satellite swath decreases from south to north is not the key reason of increasing accuracy of the $KIMURA_{new}$ product here. Based on
350 our findings about the importance of underlying image heterogeneity, we expect that there may be a higher degree of T_B homogeneity to the south, whereas in the north, approaching the ice edge, lower and patchier sea ice concentration produces relatively high heterogeneity (giving many features for the MCC algorithm to track). In order to represent and analyze the image homogeneity for the southern and northern parts of the Ross Sea buoy trajectory, we examine the standard deviation of T_B images in two 20×20 pixel subsets to approximately match the 2° latitude bin used in computing the validation metrics
355 at the southern and northern ends of its trajectory, respectively (Fig. 8). We find that the standard deviation of T_B in the southern box is much lower than in the northern box, with a mean of 2.3 K for the southern box and 17.0 K for the northern box of the Ross Sea buoy (Table 5). This analysis indicates that T_B in the northern part of Ross Sea buoy trajectory exhibits more structure than in the southern part. Thus, we conclude that the performance of MCC-based sea ice motion algorithms is sensitive to structure in the underlying T_B imagery, with potential regional ramifications which are considered in the
360 discussion.




Figure 8. Map of ASC-horizontal T_B image on 2017-09-13. Black trajectory and square boxes display subsets at start and end of the Ross Sea buoy.

365

Table 5. Standard deviation of T_B images in two subregions along the Ross Sea buoy trajectory. Times correspond to the buoy within each box. Satellite orbits and polarizations are ASC-horizontal (ASC, H), ASC-vertical (ASC, V), DES-horizontal (DES, H), and DES-vertical (DES, V).

Latitude of center of box	Time Period	Orbit and polarization	Std. dev. (K)
73.3°S	2017-05-24 ~ 2017-06-23 (30 days)	ASC, H	3.3
		ASC, V	1.3
		DES, H	3.3
		DES, V	1.3
65.0°S	2017-10-25 ~ 2017-11-23 (30 days)	ASC, H	23.8
		ASC, V	11.0
		DES, H	23.2
		DES, V	10.5

370 The three buoys used in this study were all deployed during Antarctic winter, when the sea ice south of the marginal ice zone is relatively dry. A follow-up summertime comparison would provide important information regarding the performance of the KIMURA_{new} product during the sea ice melt season, where the microwave emissivity contrast is reduced, potentially giving a more homogeneous T_B field near the ice edge, and reducing the performance of this technique.

4.3 KIMURA_{new}-derived sea ice deformation

375 Accurate sea ice motion data are crucial for calculation of sea ice kinematic parameters. Here we investigate the effect of using the shorter time scale ASC and DES sea ice motion products to retrieve sea ice D , and compare it to those derived from the longer time scale combined product.

4.3.1 Sea ice divergence

380 Fig. 9 illustrates per-pixel D RMSD for the month of July 2017 derived from the KIMURA_{new} ASC, DES and combined datasets. The spatial mean circumpolar D RMSD (Fig. 9) for the combined dataset ($1.08 \times 10^{-6} \text{ s}^{-1}$) is lower in magnitude than that of both the ASC ($1.31 \times 10^{-6} \text{ s}^{-1}$) and DES datasets ($1.22 \times 10^{-6} \text{ s}^{-1}$), indicating that the ASC and DES datasets can represent a higher magnitude of differential sea ice motion than the combined dataset. The magnitude of derived sea ice kinematics is demonstrated to be sensitive to time scale, and we have shown that shorter time scale input data produces higher sea ice D values.

385 To investigate the impacts of swath number on representing sea ice D magnitude, we compare the July 2017 mean D RMSD calculated by KIMURA_{new} ASC, DES and combined datasets in subregions north and south of the swath number discontinuity (74° S for both ASC and DES, Fig. 2). The mean D RMSD at higher latitudes is about 9%, 14% and 16% lower than at lower latitudes (i.e., north of 74° S) for the ASC, DES and combined dataset, respectively. This indicates that DM-estimated sea ice D RMSD are lower at higher latitudes, potentially due to a greater degree of smoothing due to more merged swath at higher latitudes, i.e., an undesirable bias may be introduced as a consequence of the observation configuration. While we have not investigated physical sea ice properties, which may also give rise to the lower DKP magnitude south of the swath discontinuity, we note that in the major Southern Ocean basins sea ice might be thinner and mechanically weaker at higher southern latitudes than that further north, since much of it has formed recently in coastal polynyas (particularly in the south-western Ross and Weddell seas). As such, we would expect a greater magnitude of sea ice
395 D variability there. In this case, using an S2S dataset to derive sea ice motion may be required to remove observational bias.




Figure 9. Mean D RMSD for (a) ASC, (b) DES and (c) combined KIMURA_{new} datasets of July 2017. The red dashed ellipse highlights the high D RMSD filamentary structure in the DES map, described in the text.

Fig. 9 also indicates that there are several regions with distinct features when using ASC and DES sea ice motion to calculate D . However, the central northern Weddell Sea stands out, exhibiting a region of high D RMSD in the DES (Fig. 9b, red dashed circle), which is not mirrored in the ASC RMSD D map. Such filamentary structures exist in the Weddell Sea during most wintertime months (not shown), and occur in ASC and DES maps equally, but the location is non-stationary. Given the non-stationary nature of this signal, we suggest that it is not an artifact of the satellite or analysis procedure. Noting the sun-synchronous observation platform, we suppose that this filamentary structure might be a response to external (oceanic or atmospheric) forcing with a ~ 24 h period, such as tidal currents or near-surface wind stress, but further research is beyond the scope of this paper. That this structure is a) not present in the ASC map, and b) muted in the combined map underscores the importance of considering observational time scale and overpass time when interpreting maps of DKPs.

5 Discussion

In this study, we firstly analyze DM-based Antarctic sea ice motion retrieval on sea ice kinematics by separating PM-derived KIMURA_{new} sea ice motion product into ASC and DES. Results show that performance of KIMURA_{new} datasets is a function of latitude, and that the time scale of the composite dataset has crucial influence on retrieved sea ice kinematic magnitude. Here, we discuss the impacts of features of the satellite observational configuration on sea ice kinematics representation, and how to improve observations in the future.

5.1 Impact of features of satellite observational dataset on kinematics representation

415 We find a strong link between the heterogeneity of underlying satellite-derived maps (i.e., T_B in this study) and the performance of satellite-derived sea ice velocity dataset when using MCC algorithm. From the results shown in Table 4, the Weddell Sea T_B composite presents as a relatively heterogeneous T_B field, giving better performance of sea ice velocity vectors compared to the Ross Sea comparison (Sect. 4.2). There is a higher degree of homogeneity in the southern/central Ross Sea than in the northern Ross Sea, and validation metrics (Fig. 7a and b) consequently improve to the north (Sect. 4.2).
420 The KIMURA_{new} product is derived from both horizontal and vertical polarizations, and T_B heterogeneity for horizontal polarization is found here to be generally higher than that for vertical polarization T_B (Table 3, 4 and 5), likely because the difference between emissivity of the first-year sea ice and open ocean at 36 GHz is significantly larger for horizontal polarization than for vertical polarization (Mathew et al., 2009). Thus, we expect that sea ice velocity only derived by horizontal polarization, instead of using both polarizations, may be useful when more accurate characterization of kinematics
425 is desired.

Higher spatial resolution imagery may also be advantageous in terms of providing underlying imagery which is relatively heterogeneous (i.e., improving MCC algorithm performance). In addition to the results shown in Table 2 (showing a high RMSD in the Ross Sea buoy validation metrics), we also investigate the standard deviation of T_B in the Ross Sea for several other months in 2017 and 2018 (not shown here), and find that the relatively homogeneous 36 GHz T_B in the Ross Sea is likely to be a persistent feature of ice in this sector. Also, other parameters of satellite signals, such as [radar backscattering cross-section](#), may provide more independent structure for the MCC algorithm and may be more successful in a S2S framework than DM. For this reason, the inclusion of scatterometer data may give rise to the good performance of the OSI SAF sea ice drift product OSI-405 (Lavergne, 2016), albeit giving an (often undesirable) longer time scale [due to the inclusion of several instruments](#). Also, we suggest that all MCC-based sea ice motion products report cross-correlation
430 [signal-to-noise ratio](#), as an objective measure of vector confidence and [uniqueness of matching](#) (i.e., as a metric for heterogeneity within the MCC search window; Altena et al., 2021).
435

Here we suggest that DM-derived kinematic maps represent not only sea ice kinematics, but also features of the orbit, which is highly undesirable. However, the underlying sea ice motion datasets are still valuable for studies on bulk ice export/flux (e.g., Drucker et al., 2011), which favor low RMSD datasets assimilating data from many instruments, with temporal
440 baseline a secondary consideration. In this case, the satellite motion datasets based on DM techniques (including the KIMURA product) are still valuable (e.g., Drucker et al., 2011). When vector quality is paramount, inclusion of multiple instruments is important. Both S2S and DM products using scatterometer data in addition to radiometers (e.g., the OSI SAF product) would benefit from the inclusion of higher resolution of scatterometer data (e.g., Lindsley and Long, 2010).

Deleted: motion

Deleted: motion

Deleted: motion

A recent study by Lavergne et al. (2021) shows that an S2S sea ice motion dataset provides a diverse set of temporal baselines from which to more rigorously gage kinematic processes across a wide variety of time scales. However, interpretation of an S2S product, with its inherently irregular temporal grid and spatial coverage, is likely to be more difficult than DM-derived product. Even when using only a single instrument, DM-derived datasets with their inherently long, low diversity time scale cannot track sea ice deformation processes to the same degree as S2S products. Thus, looking to the future of sea ice kinematic studies, a dataset with the flexibility to provide a wide range of time scales is necessary.

5.2 Future development of ice kinematics representation

To fully and accurately represent sea ice kinematics, some avenues for targeted research pathways are suggested here. From recent research, S2S-derived sea ice trajectories obtained from AMSR2 imagery have been found to be more accurate than the DM-derived equivalents (Lavergne et al., 2021). However, S2S products give better coverage of sea ice motion dataset in the Arctic compared to the Antarctic due to Antarctic sea ice existing at lower latitudes. Thus, combining DM and S2S concepts together and producing a mixed dataset that takes advantage of both approaches is recommended for improvement in ice kinematic representation in the future while maintaining ease of use for end users.

More satellites and instruments are needed to acquire more frequent swaths and obtain a more comprehensive sea ice motion dataset, and to provide a wider diversity of overpass times to avoid the problem of aliasing shown in Fig. 9. For example, two satellites in a tandem configuration might produce a 12-hour time scale DM product, rather than the 24-hour time scale available from a single platform.

Recently, Lavergne et al. (2021) discussed the Copernicus Imaging Microwave Radiometer (CIMR) mission for both the Arctic and Antarctica. CIMR will be a conically-scanning sun-synchronous microwave radiometer mission by the European Space Agency, with a variety of attributes favorable for production of higher accuracy and resolution sea ice motion products, and is planned for launch in 2026 (Lavergne et al., 2021). However, sun-synchronous platforms are altitude-constrained (Brown, 1998), which inherently limit their spatial resolution for a given antenna aperture. In the case of CIMR, there is a need to support other mission objectives, such as sea surface temperature remote sensing, for which sun-synchronicity is important. Given that sun synchronicity is only important for lower latitude applications (indeed, there is no diurnal insolation cycle for much of the sea ice season in both hemispheres), we suggest that a future dedicated sea ice PM radiometer platform may dispense with this constraint, and gain spatial resolution from having a lower orbit.

6 Conclusions

This study presents the first comprehensive validation of the KIMURA sea ice motion products (Kimura et al., 2013) for the [Southern Ocean](#). Three daily satellite-based PM sea ice motion datasets are investigated: the commonly used combined

product (based on data from both the ASC and DES components of each orbit), as well as the pure ASC and DES components themselves. Our detailed investigation of the performance of these products reveals two problems with the KIMURA product: triangular regions with persistent low ice motion for the ASC dataset in the western Amundsen Sea and the DES dataset in the Weddell Sea, and an inaccurate time separation assumption resulting in incorrect daily velocity estimates of up to 40%. Both problems have been amended for the KIMURA_{new} datasets used in this study, and a corrected version of the KIMURA product for general distribution is under development.

After rectifying both of these problems, we analyze performance of the KIMURA_{new} ASC, DES and combined datasets using buoy-derived sea ice velocity in the Antarctic. Results indicate that the KIMURA_{new} dataset can represent buoy velocities well in general, with mean values of correlation coefficient and regression slope of 0.97 and 0.93, respectively. The validation metrics indicate that at lower latitudes, where fewer AMSR2 swaths are available to be merged to derive the daily T_B composite image than at higher latitudes, the performance of the KIMURA_{new} datasets is better than at higher latitudes. Our results support the assertion of Lavergne et al. (2021) that the S2S products can give more accurate sea ice motion vectors than the DM-based products. We also report that for the cases studied here, 36 GHz T_B in the Ross Sea is more homogeneous than in the Weddell Sea, and homogeneity of T_B appears to underlie the reduced performance of the KIMURA_{new} dataset in the Ross Sea. Lastly, the result of Antarctic sea ice *D* variability shows that ASC and DES datasets with their shorter inherent time scale (~24 h) can represent sea ice deformation processes with higher magnitude than the longer time scale (~39 h) combined dataset. We conclude by suggesting that S2S products with their wide diversity of time scales ((Lavergne et al., 2021) will likely underpin the future of sea ice kinematics research.

Appendix A: Validation of OSI SAF and NSIDC sea ice motion products

In addition to the KIMURA_{new} dataset validation presented in the main text, the OSI SAF and NSIDC sea ice motion products have also been validated using the same buoy-derived sea ice velocity dataset, and their validation results have been compared with KIMURA_{new} combined dataset. Because the time scale of OSI SAF product is 48 hours, the comparison between buoy and OSI SAF velocities is performed using a 48-hour time scale.

510 **Table A1.** Comparison of OSI SAF, NSIDC and KIMURA_{new} combined sea ice velocity against buoy-derived velocities

<u>Validation parameter</u>	<u>Component of velocity</u>	<u>Product name</u>	<u>Ross Sea buoy (Oct 2017)</u>	<u>Central Weddell Sea buoy (July 2018)</u>	<u>Southern Weddell Sea buoy (May 2018)</u>
<u>Correlation coefficient</u>	<u>Eastward</u>	<u>OSI SAF</u>	<u>0.92</u>	<u>0.99</u>	<u>0.94</u>
		<u>NSIDC</u>	<u>0.83</u>	<u>0.93</u>	<u>0.90</u>
		<u>KIMURA_{new}</u>	<u>0.92</u>	<u>0.94</u>	<u>0.87</u>
	<u>Northward</u>	<u>OSI SAF</u>	<u>0.97</u>	<u>0.98</u>	<u>0.97</u>
		<u>NSIDC</u>	<u>0.80</u>	<u>0.91</u>	<u>0.94</u>
		<u>KIMURA_{new}</u>	<u>0.96</u>	<u>0.95</u>	<u>0.96</u>
<u>Regression slope</u>	<u>Eastward</u>	<u>OSI SAF</u>	<u>1.24</u>	<u>0.98</u>	<u>1.04</u>
		<u>NSIDC</u>	<u>1.18</u>	<u>0.98</u>	<u>1.10</u>
		<u>KIMURA_{new}</u>	<u>1.06</u>	<u>0.97</u>	<u>1.17</u>
	<u>Northward</u>	<u>OSI SAF</u>	<u>0.96</u>	<u>1.12</u>	<u>1.10</u>
		<u>NSIDC</u>	<u>1.13</u>	<u>0.88</u>	<u>1.04</u>
		<u>KIMURA_{new}</u>	<u>0.92</u>	<u>0.93</u>	<u>1.10</u>
<u>RMSD (m day⁻¹)</u>	<u>Eastward</u>	<u>OSI SAF</u>	<u>4.5×10³</u>	<u>0.8×10³</u>	<u>1.6×10³</u>
		<u>NSIDC</u>	<u>7.4×10³</u>	<u>2.5×10³</u>	<u>3.0×10³</u>
		<u>KIMURA_{new}</u>	<u>5.0×10³</u>	<u>2.0×10³</u>	<u>3.3×10³</u>
	<u>Northward</u>	<u>OSI SAF</u>	<u>2.7×10³</u>	<u>1.5×10³</u>	<u>2.1×10³</u>
		<u>NSIDC</u>	<u>7.9×10³</u>	<u>3.5×10³</u>	<u>3.3×10³</u>
		<u>KIMURA_{new}</u>	<u>4.4×10³</u>	<u>2.6×10³</u>	<u>3.0×10³</u>




Figure A1. Scatterplot comparison of OSI SAF (blue), NSIDC (red) and KIMURA_{new} combined (green) sea ice velocity product against buoy velocities in eastward (a, b and c) and northward (d, e and f) directions, Oct 2017 for the Ross Sea buoy, July 2018 for Central Weddell Sea buoy, and May 2018 for Southern Weddell Sea buoy. The black solid line represents a perfect regression slope of 1.0 with no offset.

The validation metrics (Table A1) show that all three datasets can represent buoy velocities generally well. The OSI SAF product generally represents buoy velocities with higher accuracy than the NSIDC and KIMURA_{new} product, with a mean correlation coefficient of 0.96 and of regression slope of 1.07 (OSI SAF) compared to 0.89 and 1.05 (NSIDC), 0.93 and 0.97 (KIMURA_{new}). The RMSD between buoy and satellite-derived velocity in the Ross Sea is far higher in the Ross Sea than in the Weddell Sea for all products, indicating the importance of heterogeneity of the underlying remotely-sensed fields for all products. Scatterplots of comparisons between OSI SAF and NSIDC with buoys (Fig. A1) also indicate that performance in the Weddell Sea is better than in the Ross Sea.

Data availability. The KIMURA_{new} ASC, DES and combined sea ice motion dataset used in this study is publicly available and can be accessed using the following DOI: https://data.aad.gov.au/metadata/records/AAS_4506_sea_ice_motion_corrected_Tian_2021

530 *Author contributions.* TRT processed the sea ice motion data, carried out the analysis and wrote the manuscript. NK provided ASC and DES data, and contributed to dataset rectification. ADF and PH conceived the project and contributed to the analysis. All authors contributed to the discussion and provided input during the writing process.

Competing interests. The authors declare that they have no conflict of interest.

535 *Acknowledgements.* AMSR2 Level 3 T_B data and scanning time information were accessed through JAXA's G-Portal. Ross sea buoy data was accessed from Dr Ted Maksym. This project received grant funding from the Australian Government as part of the Antarctic Science Collaboration Initiative program, it contributes to Project 6 of the Australian Antarctic Program Partnership (Project ID ASC1000002). TRT was supported through an Australian Government Research Training Program (RTP) scholarship at the Institute for Marine and Antarctic Studies, University of Tasmania. The Antarctic Science Foundation (ASF) provided generous support and funding to TRT during the COVID-19 pandemic. PH is supported through the Australian Antarctic Science Project 4506, and the International Space
540 Science Institute (Bern, Switzerland) project #405. Computation and storage facilities were provided by The National eResearch Collaboration Tools and Resources project (NeCTAR). We appreciate Drs Damian Murphy and Andrew Klekociuk for discussions and Dr Misako Kachi for her help on JAXA scanning time information. [We thank both reviewers and the editor for their encouraging comments.](#)

References

- 545 Altena, B., Kääh, A. and Wouters, B.: Correlation dispersion as a measure to better estimate uncertainty of remotely sensed glacier displacements, *Cryosph. Discuss.*, (September), 1–23, doi:10.5194/tc-2021-202, 2021.
- Arrigo, K. R. and Thomas, D. N.: Large scale importance of sea ice biology in the Southern Ocean, *Antarct. Sci.*, 16(4), 471–486, doi:10.1017/S0954102004002263, 2004.
- Brown, C. D.: *Spacecraft Mission Design*, Second Edition., 1998.
- Curry, J. A., Schramm, J. L. and Ebert, E. E.: Sea ice-albedo climate feedback mechanism, *J. Clim.*, 8(2), 240–247,
550 doi:10.1175/1520-0442(1995)008<0240:SIACFM>2.0.CO;2, 1995.
- Dieckmann, G. S. and Hellmer, H. H.: The Importance of Sea Ice: An Overview, in *Sea Ice*, vol. 2, pp. 1–22, Wiley-Blackwell, Oxford, UK., 2010.
- Drucker, R., Martin, S. and Kwok, R.: Sea ice production and export from coastal polynyas in the Weddell and Ross Seas, *Geophys. Res. Lett.*, 38(17), 4–7, doi:10.1029/2011GL048668, 2011.
- 555 Ebert, E. E. and Curry, J. A.: An intermediate one-dimensional thermodynamic sea ice model for investigating ice-atmosphere interactions, *J. Geophys. Res.*, doi:10.1029/93jc00656, 1993.
- Emery, W. J., Fowler, C. W., Hawkins, J. and Preller, R. H.: Fram Strait satellite image-derived ice motions, *J. Geophys. Res.*, doi:10.1029/90JC02273, 1991.

- Giles, A. B., Massom, R. A., Heil, P. and Hyland, G.: Semi-automated feature-tracking of East Antarctic sea ice from
560 Envisat ASAR imagery, *Remote Sens. Environ.*, 115(9), 2267–2276, doi:10.1016/j.rse.2011.04.027, 2011.
- Goosse, H., Campin, J. M., Fichefet, T. and Deleersnijder, E.: Impact of sea-ice formation on the properties of
Antarctic bottom water, *Ann. Glaciol.*, doi:10.3189/s0260305500014154, 1997.
- Hakkinen, S.: Seasonal simulation of the Southern Ocean coupled ice-ocean system, *J. Geophysical Res.*,
doi:10.1029/95jc02441, 1995.
- 565 Heil, P. and Allison, I.: The pattern and variability of Antarctic sea-ice drift in the Indian Ocean and western Pacific
sectors, *J. Geophys. Res. Ocean.*, doi:10.1029/1999jc900076, 1999.
- Heil, P., Lytle, V. I. and Allison, I.: Enhanced thermodynamic ice growth by sea-ice deformation, *Ann. Glaciol.*,
doi:10.3189/1998aog27-1-433-437, 1998.
- Heil, P., Fowler, C. W., Maslanik, J. A., Emery, W. J. and Allison, I.: A comparison of East Antarctic sea-ice motion
570 derived using drifting buoys and remote sensing, *Ann. Glaciol.*, doi:10.3189/172756401781818374, 2001.
- Heil, P., Fowler, C. W. and Lake, S. E.: Antarctic Sea-ice velocity as derived from SSM/I imagery, *Ann. Glaciol.*, 44,
361–366, doi:10.3189/172756406781811682, 2006.
- Heil, P., Massom, R. A., Allison, I., Worby, A. P. and Lytle, V. I.: Role of off-shelf to on-shelf transitions for East
Antarctic sea ice dynamics during spring 2003, *J. Geophys. Res. Ocean.*, 114(C9), 2009.
- 575 Heil, P., Massom, R. A., Allison, I. and Worby, A. P.: Physical attributes of sea-ice kinematics during spring 2007 off
East Antarctica, *Deep. Res. Part II Top. Stud. Oceanogr.*, 58(9–10), 1158–1171, doi:10.1016/j.dsr2.2010.12.004, 2011.
- Hoeber, H. and Gube-Lenhardt, M.: The eastern Weddell Sea drifting buoy data set of the Winter Weddell Sea
Project (WWSP) 1986., *Berichte zur Polarforsch. Reports Polar Res.*, 37, 1987.
- Hutchings, J. K., Heil, P. and Hibler, W. D.: Modeling linear kinematic features in sea ice, *Mon. Weather Rev.*,
580 133(12), 3481–3497, doi:10.1175/MWR3045.1, 2005.
- Hutchings, J. K., Heil, P., Steer, A. and Hibler, W. D.: Subsynoptic scale spatial variability of sea ice deformation in
the western Weddell Sea during early summer, *J. Geophys. Res. Ocean.*, 117(1), doi:10.1029/2011JC006961, 2012.
- Hutter, N., Losch, M. and Menemenlis, D.: Scaling Properties of Arctic Sea Ice Deformation in a High-Resolution
Viscous-Plastic Sea Ice Model and in Satellite Observations, *J. Geophys. Res. Ocean.*, 123(1), 672–687,
585 doi:10.1002/2017JC013119, 2018.
- Kimura, N.: Sea ice motion in response to surface wind and ocean current in the Southern Ocean, *J. Meteorol. Soc.
Japan*, 82(4), 1223–1231, doi:10.2151/jmsj.2004.1223, 2004.
- Kimura, N. and Wakatsuchi, M.: Large-scale processes governing the seasonal variability of the Antarctic sea ice,
Tellus, Ser. A Dyn. Meteorol. Oceanogr., 63(4), 828–840, doi:10.1111/j.1600-0870.2011.00526.x, 2011.
- 590 Kimura, N., Nishimura, A., Tanaka, Y. and Yamaguchi, H.: Influence of winter sea-ice motion on summer ice cover
in the Arctic, *Polar Res.*, 32(1), 20193, 2013.
- Kirwan, A. D.: Oceanic Velocity Gradients, *J. Phys. Oceanogr.*, doi:10.1175/1520-

- 0485(1975)005<0729:ovg>2.0.co;2, 1975.
- 595 Kottmeier, C. and Sellmann, L.: Atmospheric and oceanic forcing of Weddell Sea ice motion, *J. Geophys. Res. C Ocean.*, 101(C9), 20809–20824, doi:10.1029/96JC01293, 1996.
- Kwok, R.: Satellite remote sensing of sea-ice thickness and kinematics: A review, *J. Glaciol.*, 56(200), 1129–1140, doi:10.3189/002214311796406167, 2011.
- Lavergne, T.: Low Resolution Sea Ice Drift Product User’s Manual, [online] Available from: https://osisaf.metsis.met.no/sites/osisaf.metsis.met.no/files/user_manuals/osisaf_cdop2_ss2_pum_sea-ice-drift-lr_v1p8.pdf, 600 2016.
- Lavergne, T. and Eastwood, S.: Low Resolution Sea Ice Drift Product User’s Manual, GBL LR SID–OSI 405, [online] Available from: http://osisaf.met.no/p/ice/lr_ice_drift.html#lavergne:2010:cmcc-jgr, 2010.
- Lavergne, T., Piñol Solé, M., Down, E. and Donlon, C.: Towards a swath-to-swath sea-ice drift product for the Copernicus Imaging Microwave Radiometer mission, *Cryosphere*, 15(8), doi:10.5194/tc-15-3681-2021, 2021.
- 605 Lindsley, R. D. and Long, D. G.: Adapting the SIR algorithm to ASCAT, in *International Geoscience and Remote Sensing Symposium (IGARSS)*, 2010.
- Marsan, D., Stern, H., Lindsay, R. and Weiss, J.: Scale dependence and localization of the deformation of Arctic sea ice, *Phys. Rev. Lett.*, 93(17), doi:10.1103/PhysRevLett.93.178501, 2004.
- Mathew, N., Heygster, G. and Melsheimer, C.: Surface emissivity of the Arctic sea ice at AMSR-E frequencies, *IEEE 610 Trans. Geosci. Remote Sens.*, 47(12), doi:10.1109/TGRS.2009.2023667, 2009.
- Meier, W. N., Maslanik, J. A. and Fowler, C. W.: Error analysis and assimilation of remotely sensed ice motion within an Arctic sea ice model, *J. Geophys. Res. Ocean.*, 105(C2), doi:10.1029/1999jc900268, 2000.
- Melsheimer, C. and Spreen, G.: AMSR2 ASI sea ice concentration data, Antarctic, version 5.4 (NetCDF) (July 2012 - December 2018), , doi:10.1594/PANGAEA.898400, 2019.
- 615 Molinari, R. and Kirwan, A. D.: Calculations of Differential Kinematic Properties from Lagrangian Observations in the Western Caribbean Sea, *J. Phys. Oceanogr.*, doi:10.1175/1520-0485(1975)005<0483:codkpf>2.0.co;2, 1975.
- Ninnis, R. M., Emery, W. J. and Collins, M. J.: Automated extraction of pack ice motion from Advanced Very High Resolution Radiometer imagery, *J. Geophys. Res.*, doi:10.1029/jc091ic09p10725, 1986.
- Rintoul, S. R., Hughes, C. W. and Olbers, D.: Chapter 4.6 The Antarctic Circumpolar Current system, *Int. Geophys.*, 620 77(C), 271–302, doi:10.1016/S0074-6142(01)80124-8, 2001.
- Scargle, J. D.: Studies in astronomical time series analysis. II - Statistical aspects of spectral analysis of unevenly spaced data, *Astrophys. J.*, doi:10.1086/160554, 1982.
- Schröder, M.: The Expedition PS111 of the Research Vessel POLARSTERN to the southern Weddell Sea in 2018, Bremerhaven, Germany., doi: 10.2312/BzPM_0718_2018, 2018.
- 625 Spreen, G., Kwok, R., Menemenlis, D. and Nguyen, A. T.: Sea-ice deformation in a coupled ocean–sea-ice model and in satellite remote sensing data, *Cryosph.*, 11(4), 1553, 2017.

- Sumata, H., Lavergne, T., Girard-Arduin, F., Kimura, N., Tschudi, M. A., Kauker, F., Karcher, M. and Gerdes, R.: An intercomparison of Arctic ice drift products to deduce uncertainty estimates, *J. Geophys. Res. Ocean.*, 119(8), 4887–4921, doi:10.1002/2013je009724, 2014.
- 630 Szanyi, S., Lukovich, J. V., Barber, D. G. and Haller, G.: Persistent artifacts in the NSIDC ice motion data set and their implications for analysis, *Geophys. Res. Lett.*, 43(20), 10,800-10,807, doi:10.1002/2016GL069799, 2016.
- Tison, J. L., Maksym, T., Fraser, A. D., Fraser, A. D., Corkill, M., Corkill, M., Kimura, N., Nosaka, Y., Nomura, D., Nomura, D., Nomura, D., Vancoppenolle, M., Ackley, S., Stammerjohn, S., Wauthy, S., Van Der Linden, F., Van Der Linden, F., Carnat, G., Sapart, C., De Jong, J., Fripiat, F. and Delille, B.: Physical and biological properties of early winter Antarctic sea ice in the Ross Sea, *Ann. Glaciol.*, doi:10.1017/aog.2020.43, 2020.
- 635 Tschudi, M. A., Meier, W. N. and Stewart, J. S.: An enhancement to sea ice motion and age products, *Cryosph. Discuss.*, 1–29, doi:10.5194/tc-2019-40, 2019.
- Walsh, J. E.: The role of sea ice in climatic variability: Theories and evidence, *Atmos. - Ocean*, 21(3), 229–242, doi:10.1080/07055900.1983.9649166, 1983.
- 640 Weiss, J. and Marsan, D.: Scale properties of sea ice deformation and fracturing, *Comptes Rendus Phys.*, 5(7), 735–751, doi:10.1016/j.crhy.2004.09.005, 2004.
- Yang, J. and Neelin, J. D.: Sea-ice interaction with the thermohaline circulation, *Geophys. Res. Lett.*, 20(3), 217–220, doi:10.1029/92GL02920, 1993.
- 645