

# Antarctic contribution to future sea level from ice shelf basal melt as constrained by ice discharge observations

Eveline C. van der Linden<sup>1</sup>, Dewi Le Bars<sup>1</sup>, Erwin Lambert<sup>1</sup>, and Sybren Drijfhout<sup>1,2</sup>

<sup>1</sup>Royal Netherlands Meteorological Institute, Utrechtseweg 297, 3731 GA, De Bilt, The Netherlands

<sup>2</sup>Institute for Marine and Atmospheric Research Utrecht, Department of Physics, Utrecht University, Princetonplein 5, 3584 CC, Utrecht, The Netherlands

**Correspondence:** Eveline C. van der Linden (linden@knmi.nl)

**Abstract.** Antarctic mass loss is the largest contributor to uncertainties in sea level projections on centennial timescales. In this study we aim to constrain future projections of the contribution of Antarctic dynamics by using ice discharge observations. The contribution of Antarctica's ice discharge is computed with ocean thermal forcing from 14 earth system models (ESMs) and linear response functions (RFs) from 16 ice sheet models for three shared socio-economic pathway (SSP) scenarios. New compared to previous studies, basal melt sensitivities to ocean temperature changes were calibrated on four decades of observed ice discharge changes rather than using observation-based basal melt sensitivities. Calibration improved historical performance, but did not reduce the uncertainty in the projections. The results show that even with calibration the acceleration during the observational period is underestimated for the Amundsen region, indicating missing physics. Also the relative contribution of the Amundsen region is underestimated. The Amundsen contribution and sea level acceleration are improved by choosing an Amundsen-specific calibration (rather than Antarctic-wide), quadratic basal melt parameterisation (rather than linear) and thermal forcing near the ice shelf base (rather than the deepest layer above the continental shelf). With these methodological choices we arrive at a median dynamic sea level contribution of 0.14 m in 2100 relative to 1995-2014 under SSP2-4.5, sitting in between projections of previous multi-model studies (ISMIP6 and LARMIP-2). Our results show that constraining the basal melt parameterisation on Amundsen ice discharge rather than applying the basal melt sensitivities used in ISMIP6 and LARMIP-2 leads to higher sea level contributions. We also show that uncertainties associated with ESMs and RFs affect the projected sea level contribution more than our methodological choices in the calibration and basal melt computation method. This suggests that constraining the basal melt relation in ISMIP6 and LARMIP-2 with ice discharge observations in the Amundsen region will lead to higher future estimates than those presented in IPCC AR6.

## 1 Introduction


Sea level rise poses an increasing threat to densely populated coasts and deltas worldwide (Hinkel et al., 2014). Even if the 1.5 degree target of the Paris Agreement is met, global mean sea level will rise several meters in the longer term (Clark et al., 2016; Fox-Kemper et al., 2021). At present, a global acceleration of sea level rise is visible in satellite measurements and the sea level is already rising more than twice as fast as the average rate over the twentieth century (Nerem et al., 2018; Dangendorf et al., 2019).

25 Mass loss from land ice (ice sheets and glaciers) is currently accelerating and is now (over the period 2006–2018) the largest contributor to the global mean sea level rise (Fox-Kemper et al., 2021). Antarctic ice sheet (AIS) mass loss has tripled over the last decade (Shepherd et al., 2018), which can be mainly attributed to increased ice discharge in the Amundsen Sea (Rignot et al., 2019). Models and geological data indicate that the AIS will cause most of the sea level rise over thousands of years (Bamber et al., 2019). Moreover, melt of Antarctic land ice is the largest contributor to uncertainties on centennial timescales (Palmer et al., 2020; van de Wal et al., 2019). The degree of acceleration of future sea level changes is mainly determined by dynamic processes on the AIS. The underlying processes are 1) increased melt from below by warmer ocean water (basal melt) and 2) increased calving (iceberg formation) triggered by basal melt and/or surface melt (Rignot and Jacobs, 2002; Pritchard et al., 2012; Liu et al., 2015; van den Broeke, 2005). It is important to gain a better understanding of the many uncertainties about the Antarctic contribution to sea level rise that exist and to reduce these uncertainties when possible to support adaptation planning (Haasnoot et al., 2020). Uncertainties associated with the Antarctic contribution to sea level rise appear to be increasing since more and more models and processes are included in the uncertainty assessments. Using similar methodologies to each other, the estimated Antarctic contribution in Levermann et al. (2020) shows increased uncertainty compared to its previous study (Levermann et al., 2014) and expert judgment assessments of Bamber et al. (2019) give higher uncertainties than before (Bamber and Aspinall, 2013). To address this issue, our study aims to gain more insight in the Antarctic contribution to, and uncertainties in, future sea level changes and provides directions for reducing these uncertainties.

Future projections of Antarctic mass loss are based on modelling studies, in which ice sheet models are used as a standalone unit and forcing is provided by earth system models (ESMs). Over the last decade, ice sheet modelling has advanced from single model studies to model intercomparison projects (MIPs). In these projects, earth system modelling and ice sheet modelling are combined to make projections of land ice. The Ice Sheet Model Intercomparison Project for CMIP6 (ISMIP6) (Nowicki et al., 2016) and Linear Antarctic Response Model Intercomparison Project (LARMIP-2) (Levermann et al., 2020) are currently used as one basis for projections of the Antarctic land ice evolution (Fox-Kemper et al., 2021). ISMIP6 (Seroussi et al., 2020) provides process-based projections of the sea level contribution of the AIS based on a variety of ice sheet models that are forced by atmosphere and ocean output from CMIP5 ESMs. ISMIP6 made a selection of six ESMs based on two main criteria. The first criterion is based on their performance in reproducing the mean state of the current climate (atmosphere and ocean) near Antarctica, but did not include trends. The second criterion ensures that the ESM selection includes a diversity of warming rates over the 21st century so that the uncertainty-range in projections is captured (Barthel et al., 2020; Nowicki et al., 2020). One risk of this selection process is that models with a relatively bad performance over the historical period in terms of trends could have been chosen. In ISMIP6 basal melt is calibrated on basal melt observations with two options for calibration: the mean AIS and Pine Island’s grounding line (Jourdain et al., 2020). LARMIP-2 focuses on ice sheet mass loss due to ice shelf basal melt (Levermann et al., 2014, 2020). In that study, the temperature melt-relation is parameterised with a linear dependency on thermal forcing. ISMIP6 and LARMIP-2 have thirteen ice sheet models in common and are primarily based on the CMIP5 ESMs and scenarios (RCPs) as forcing. Payne et al. (2021) demonstrate that the estimated AIS mass loss in ISMIP6 models with CMIP6 forcing is similar compared to using CMIP5 forcing. Edwards et al. (2021) estimated probability distributions for projections under the SSP scenarios based on CMIP6 ESMs, by using statistical emulation of the ISMIP6 ice sheet models.

60 Our study follows LARMIP-2 to account for the sensitivity of ice sheet models to climate change by using linear response functions (RF) of ice sheet models. The LARMIP-2 RFs were obtained by prescribing for five regions an immediate change in basal melt of the ice shelves and simulating the resulting ice loss with the ice sheet model. The changes in the volume above flotation of the ice sheet are then calculated to obtain the sea-level equivalent ice loss. In this way a relationship between basal melt and the related contribution to sea level is obtained for each region: the linear response function. Additionally, 65 a relationship between thermal forcing and basal melt is used to compute basal melt from ocean temperatures: the basal melt parameterisation. These relationships, together with a time-dependent warming derived from ESMs, then lead to a time-dependent mass loss of the ice sheet. This method was applied by Levermann et al. (2014, 2020) to a number of ice sheet models. In those studies, CMIP5 models were used to diagnose the relationship between global surface air temperature (GSAT) and ocean temperature changes around Antarctica, and GSAT was used as a driver of the method. The advantage of using 70 GSAT over ocean temperature changes as a driver is that also uncertainties in GSAT changes were included in the uncertainty estimate. Furthermore, GSAT is easier to derive, but it does not account for (future changes in) Southern Ocean dynamics. It could be expected that a regional metric has a better relation with forcing underneath ice shelves. Therefore, the current study improves this step by using subsurface ocean temperature as the driver (Lambert et al., 2021). In addition to the linear melt parameterisation as in the Levermann et al. (2020) study, a more advanced quadratic basal melt parameterisation is applied 75 since observation-based evidence suggests a nonlinear relationship between melting and ocean temperature (Jenkins et al., 2018).

The basal melt parameterisations are calibrated on the sea level contribution derived from observation-based changes in grounding line ice discharge (Rignot et al., 2019), rather than on basal melt as is done in ISMIP6. One advantage of using ice discharge measurements is that they capture the entire ice sheet through satellite measurements of ice height and velocity and 80 therefore are better constrained than basal melt estimates which are not measured for the full ice sheet and for the full time period that we use for calibration. Moreover, when using basal melt for calibration, basal melt observations are required long before the actual ice discharge acceleration takes place due to the delayed response of ice discharge to basal melt. The advantage of this new approach is that ice discharge acceleration during the historical period is directly derived from observations. Since basal melt has a delayed impact on ice discharge, using ice discharge observations for calibration constrains the basal melt 85 even before the observational period. As a calibration target, the mass loss estimates of Rignot et al. (2019) were chosen over Shepherd et al. (2018) for two reasons. The first reason is that Rignot et al. (2019) does not include surface mass balance processes which makes the data directly comparable with the linear response functions that only represent the contribution of Antarctic dynamics. The second reason is that the Rignot et al. (2019) record starts earlier which allows us to look into mass loss acceleration during a longer period. Two different calibration methods are applied: a regional calibration on the Amundsen 90 sector and one at the continental scale. By applying the same melt relation to the past and the future, we ensure that the physics is consistent with four decades of observed mass loss. Here, the assumption is that no new processes are taking place. Using different warming scenarios and RFs for a variety of models, we arrive at a new estimate of the future mass loss of Antarctica and the Amundsen sector that is constrained by observed ice discharge.



**Figure 1.** Flow diagram of procedure. Observational constraints are indicated in orange, main computations of the Levermann et al. method in green (including model experiments by the modelling groups), calibration methods in yellow, bias-adjustment in grey and (intermediate) output data in blue. The continuous lines represent direct pathways while the dashed lines refer to iterative processes or optional choices during calibration.

## 2 Methodology

95 In this study the contribution of changes in Antarctica’s ice discharge to sea level changes is computed with state-of-the-art  
 100 ESMS from Coupled Model Intercomparison Project Phase 6 (CMIP6; Eyring et al. 2016) and linear response functions from  
 the Linear Antarctic Response MIP (LARMIP-2; Levermann et al. 2020) ice sheet models. The basic procedure of this study  
 follows that of Levermann et al. (2020) with a number of modifications. First, we give a brief explanation of the procedure as  
 illustrated in Fig. 1.

100 All computations are performed for five ocean sectors around the Antarctic continent (Fig. 2). The regional mean subsurface  
 ocean temperatures are taken from each CMIP6 ESM and bias-adjusted with a global ocean reanalyses dataset (Sect. 2.1).

Then basal melt is computed from these bias-adjusted temperatures with a basal melt parameterisation and a first guess for the basal melt sensitivity (calibration parameter) which is derived from basal melt observations (Sect. 3.1). The resulting basal melt anomalies are fed into the linear response functions to compute the regional sea level contribution for each of the five sectors (Sect. 2.3). The sum of the five regions gives the summed Antarctic sea level contribution. The calibration starts either after the regional sea level computation (regional calibration) or after computing the summed Antarctic response (Antarctic-wide calibration) (Sect. 2.4). For each ESM-RF combination, the resulting sea level contribution is compared with observed grounding line ice discharge from Rignot et al. (2019). The basal melt sensitivity is used as the calibration parameter to improve the fit with observations. This is an iterative procedure. The calibration is performed on the Amundsen region (regional calibration) and for the sum of all regions (Antarctic-wide). Optionally, a model selection could be performed based on a comparison with observed ice discharge (Rignot et al., 2019). Details of each step are described in the subsections that follow.

## 2.1 Ocean forcing

Earth system models from CMIP6 are used as a basis for the computations, guaranteeing implementation of state-of-the-art models in the analysis and projections. The ocean forcing consists of annual mean simulated subsurface ocean temperatures which are obtained from ESM output instead of estimating them from scaling coefficients and GSAT as in LARMIP-2 (Lambert et al., 2021). The ocean temperatures are taken from the historical experiment (1850-2014) and the Shared Socioeconomic Pathway (SSP) scenarios SSP1-2.6, SSP2-4.5 and SSP5-8.5 (2015-2100). Only models that have data available at the Earth System Grid Federation (ESGF) data server for the historical experiment and all three SSP scenarios (at the time of study) are considered. In addition, models should provide data for the full period (1850-2100) without any data gaps since the computation of the delayed ice sheet response to basal melt requires a continuous time series. Table 1 summarises which models have been taken into account.

Ocean temperatures are averaged over five oceanic sectors: the East AIS (EAIS), Ross, Amundsen, Weddell and Peninsula sector (Fig. 2), and averaged vertically over a range of 100 m, centered around the depth of the ice shelf base (Table 2). In addition, temperatures in an ocean layer around the depth of the continental shelf near the ice shelf front (800-1000 m) were used to assess the impact of thermal forcing depth on the projections (Table 3)(Sect. 3.3.2). The deeper ocean layer is chosen as it approximately represents the deeper water masses on the continental shelf that have access to the cavities under ice shelves. Different from Levermann et al. (2020), the Peninsula sector is defined as a separate ocean sector rather than using the same ocean sector coordinates as the Amundsen sector.


The ocean temperature time series are corrected for model drift by removing the long term trend diagnosed by the linear trend in the pre-industrial control (piControl) experiment (Fig. 3). For models that did not provide suitable data for the pi-Control experiment, the model drift is not removed. Although the ocean temperature bias has no clear relation with projected temperature trends in ESMs (Little and Urban, 2016), it affects the magnitude of basal melt in the quadratic parameterisation. Therefore, before computing the basal melt the time mean ocean temperatures are bias-adjusted with global ocean reanalyses called the Global Reanalysis Ensemble Product (GREP). GREP can be obtained from the Copernicus Marine Server at 1 de-

**Table 1.** CMIP6 ESMs that have been evaluated. For each region the subsurface ocean temperature bias (in K) compared to the GREP reanalysis is indicated over the period 1993-2018, including years 2015-2018 for the SSP2-4.5 scenario. The ‘drift correction’ column indicates whether the piControl experiment was used for model drift correction. The bottom rows show the mean and standard deviation ( $\sigma$ ) of the ESM biases and the mean ocean temperature (in  $^{\circ}\text{C}$ ) and standard deviation of the GREP reanalysis product.

CMIP6 ESM	EAIS	Weddell	Amundsen	Ross	Peninsula	Drift correction
ACCESS-CM2	-0.33	-0.11	-1.05	-1.26	0.09	–
CAMS-CSM1-0	0.24	-0.05	0.22	-0.94	0.39	piControl
CAS-ESM2-0	1.43	0.79	0.20	-0.18	2.18	–
CMCC-ESM2	0.31	-0.23	0.51	-0.10	0.58	piControl
CanESM5	-0.55	-0.43	-0.07	-0.80	-0.21	piControl
EC-Earth3	0.06	-0.57	1.17	0.71	-0.33	–
EC-Earth3-Veg	-0.10	-0.58	0.84	0.44	-0.34	piControl
GFDL-ESM4	0.05	-0.38	0.45	-1.00	0.20	piControl
INM-CM4-8	-0.37	0.32	-0.66	-0.17	0.19	piControl
INM-CM5-0	-0.74	-0.24	-1.16	-1.11	-0.16	piControl
MIROC6	0.81	0.55	1.58	1.40	0.29	–
MPI-ESM1-2-LR	-0.31	0.03	0.08	-0.59	-0.41	piControl
MRI-ESM2-0	-0.12	-0.10	-0.12	-0.31	0.32	–
NorESM2-MM	-0.92	-0.45	-0.71	-0.84	-0.74	piControl
Bias Mean	-0.04	-0.10	0.09	-0.34	0.15	-
Bias $\sigma$	0.59	0.40	0.78	0.74	0.67	-
GREP Mean	0.53	-0.79	1.37	-0.18	-0.24	
GREP $\sigma$	0.23	0.21	0.24	0.53	0.21	

**Table 2.** Mean ice shelf depth (in m) for the five sectors in Fig. 2.

Sector	Depth (m)
EAIS	369
Weddell	420
Amundsen	305
Ross	312
Peninsula	420



**Figure 2.** Ocean sector definition.

green horizontal resolution over the period during which altimetry data observations are available (1993-2018). It is constructed by postprocessing of four reanalyses: GLORYS2V4 from Mercator Ocean (France), ORAS5 from ECMWF, FOAM/GloSea5 from Met Office (UK), and C-GLORS05 from CMCC (Italy). It should be noted, however, that the reanalysis data may also be biased due to a paucity of assimilated data and the absence of ice shelves in the physical ocean models.


140 Averaged over all CMIP6 ESMs the subsurface temperature is cold-biased for the EAIS, Weddell and Ross sectors over the 1993-2018 period. For the Amundsen and Peninsula sectors the mean simulated temperature is warm-biased (Table 1). For all regions, the sign of the bias differs between individual models. The ocean temperature time series of the individual models are corrected by the ensemble mean of the reanalysis products over the 1993-2018 time period over the entire historical and future period to obtain the bias-adjusted ocean temperatures (Fig. 3).

## 145 2.2 Basal melt parameterisation

CMIP6 ESMs do not represent ice shelf cavities and the related thermal and dynamical properties. Coastal ocean temperatures should therefore be translated into these cavities. This can be done by using a parameterisation that relates the far-field (coastal) ocean temperature to basal melt. Most of the simple basal melt parameterisations assume a relation with thermal forcing, i.e. the difference between the *in situ* temperature of sea water ( $T_o$ ) and the *in situ* freezing-melting point temperature ( $T_f$ ):

$$150 \quad TF = T_o - T_f. \quad (1)$$

Our main method employs a quadratic melt relation with thermal forcing (Table 3) as the quadratic relation was suggested to outperform a linear relation (Favier et al., 2019). However, we will also apply a linear relation so that we can compare our



**Figure 3.** Annual mean subsurface ocean temperature time series averaged over all ESMs (green), model drift- and bias-adjusted, and the GREP ensemble mean (orange). Both are smoothed by a five-year running average filter. The temperature is derived from a 100-m thick layer centered around the mean depth of the ice shelf base as specified in Table 2. The historical experiment (1850-2014) is combined with SSP2-4.5 (2015-2018) for this visualisation. Note that the tick distances of the vertical axis are the same for all regions, but the ranges are different.

**Table 3.** Overview of basal melt computation and calibration methods applied in this study. Two different depths were used for the thermal forcing: centered around the mean depth of the ice shelf base and the layer at 800-1000 m depth. Also, two different basal melt parameterisation methods were employed: linear and quadratic. Each parameterisation has been calibrated Antarctic wide and regionally on the Amundsen region. Finally, median basal melt sensitivities used in LARMIP-2 ( $11.5 \text{ m yr K}^{-1}$ ) and ISMIP6 AntMean method ( $2.6 \text{ m yr K}^{-2}$ ) have been applied in the linear and quadratic parameterisation, respectively.

Thermal forcing depth	Parameterisation relation	Basal melt sensitivity
Ice shelf base	Quadratic	Amundsen calibration
800-1000 m	Linear	Antarctic-wide calibration
		ISMIP6 AntMean Median
		LARMIP-2 Median

results with the linear relation used in Levermann et al. (2020). The linear relation is defined as:

$$m = \gamma_l \left( \frac{\rho_{sw} c_{po}}{\rho_i L_i} \right) TF, \quad (2)$$



**Table 4.** Physical constants.

parameter	symbol	value	unit
ice density	$\rho_i$	917	kg m <sup>-3</sup>
sea water density	$\rho_{sw}$	1028	kg m <sup>-3</sup>
specific heat capacity of ocean mixed layer	$c_{po}$	3947	J kg <sup>-1</sup> K <sup>-1</sup>
latent heat of fusion of ice	$L_i$	$3.34 \times 10^5$	J kg <sup>-1</sup>
heat exchange velocity	$\gamma$	calibrated	m s <sup>-1</sup>
liquidus slope	$\lambda_1$	-0.0575	°C PSU <sup>-1</sup>
liquidus intercept	$\lambda_2$	0.0832	°C
liquidus pressure coefficient	$\lambda_3$	$7.59 \times 10^{-4}$	°C m <sup>-1</sup>

155 where  $m$  is the basal melt and  $\gamma_l$  is the linear calibration parameter. It assumes a constant heat exchange, independent on the local stratification and circulation. The quadratic relation is defined as:

$$m = \gamma_q \left( \frac{\rho_{sw} c_{po}}{\rho_i L_i} \right)^2 TF |TF|. \quad (3)$$

160 where the quadratic calibration parameter is  $\gamma_q$ . The basal melt sensitivity is defined as  $\gamma_l \left( \frac{\rho_{sw} c_{po}}{\rho_i L_i} \right)$  for the linear relation and  $\gamma_q \left( \frac{\rho_{sw} c_{po}}{\rho_i L_i} \right)^2$  for the quadratic relation. The quadratic relation assumes that the heat exchange scales with the buoyancy-driven cavity circulation and that this scales linearly with the large-scale temperature gradient. The values of the physical constants  $\rho_{sw}$ ,  $c_{po}$ ,  $\rho_i$  and  $L_i$  are given in Table 4. The freezing-melting point temperature  $T_f$  is computed from the ocean salinity  $s_o$  and the thermal forcing depth  $z_b$ :

$$T_f = \lambda_1 s_o + \lambda_2 + \lambda_3 z_b. \quad (4)$$

Favier et al. take  $T_o$  and  $T_f$  either as local or as the product of local and the average over the entire ice draft of a given sector. 165 The thermal forcing depth is the depth of the ice shelf base or 800-1000 m (Table 3). In the current study, a purely nonlocal forcing is applied, similar to DeConto and Pollard (2016) and Levermann et al. (2020). This is because the linear response functions are derived from a homogeneous melt perturbation over the entire ice draft and therefore a single basal melt value is required per region for each time step. The values of  $T_o$  are computed as averages over the five (far-field) oceanic sectors, around the depth of the ice shelf base (see Table 2) or a deeper layer (800-1000 m depth). Since CMIP6 ESMs do not resolve 170 cavities, the far-field ocean temperature is taken. The underlying assumption is that the ocean temperature remains constant while it is advected into the cavity. The computation of  $T_f$  is based on a constant salinity value for each oceanic sector, which is computed from the far-field salinity climatology of the reanalysis data. The resulting values of  $T_f$  are approximately -1.6 °C in each sector.

Note that the melt is positive if the ocean temperature exceeds the freezing-melting point temperature and negative (i.e. water is refreezing) otherwise. In the current study, basal melt anomalies are used to compute the sea level contribution. The basal melt anomalies are defined as the difference in basal melt between time  $t$  and the baseline time period, 1850-1930. This period was chosen since it is long enough to reduce the impact of natural variability on the baseline but short enough so that it doesn't include the trends due to anthropogenic forcing.

### 2.3 Sea level contribution

Linear response functions (RFs) from LARMIP-2 will be used to compute the cumulative sea level contribution  $\Delta S$  (in meters) due to a change in basal melt  $\Delta m$  for each of the five sectors (Fig. 2):

$$\Delta S(t) = \int_0^t d\tau \Delta m(\tau) \cdot RF(t - \tau). \quad (5)$$

The sum of the five regional sea level contributions gives the total Antarctic sea level contribution.

LARMIP-2 provides RFs of 16 ice sheet models. Combined with the 14 ESMs (Table 1), this results in 224 ESM-RF combinations for the projections.


### 2.4 Calibration

For each ESM-RF pair, the basal melt parameterisation is calibrated on observed ice discharge from Rignot et al. (2019) (grey lines in Fig. 6). The basal melt parameterisation can be calibrated with the heat exchange velocity  $\gamma$ . It should be noted that  $\gamma_l$  and  $\gamma_q$  have a different order of magnitude in the linear and quadratic parameterisation, respectively, and are not directly comparable. The root-mean-square error (RMSE) between the observed and modelled cumulative changes in ice discharge for each year, weighted equally, over the period 1979-2017 for each ESM-RF pair is determined over a wide range of  $\gamma$  values for Eq. (2) and Eq. (3).

$$\text{RMSE} = \sqrt{\frac{\sum_{t=1}^T (\Delta S_{\text{simulated}}(t) - \Delta S_{\text{observed}}(t))^2}{T}} \quad (6)$$

The RMSE is computed over the full time series to constrain models on the cumulative sea level change as well as the acceleration. The  $\gamma$  value giving the lowest RMSE for each ESM-RF pair provides the calibrated basal melt sensitivity. Since the observational uncertainty is small compared to the intermodel spread (Fig. 6), it was not taken into account in the calibration. Note that this calibration step is a key difference with the Levermann et al. studies. Levermann et al. (2020) does not calibrate the basal melt parameterisation on ice discharge, but uses melt sensitivities derived from observations.

The calibration is applied regionally on the Amundsen region and Antarctic-wide (Table 3), resulting in two basal melt sensitivities for each ESM-RF pair for a given parameterisation. Figure 5 shows the basal melt sensitivities corresponding with the calibrated  $\gamma$  values for the linear and quadratic basal melt parameterisation and for the two calibration regions. For the Antarctic-wide calibration, the same  $\gamma$  value is applied to each region. The smallest RMSE between the summed discharge



**Figure 4.** Thermal forcing anomalies for SSP1-2.6, SSP2-4.5 and SSP5-8.5 including all evaluated CMIP6 ESMs (Table 1) from 1950 to 2100 relative to the baseline period 1850-1930. The shaded regions indicate the intermodel spread (17th to 83rd percentiles) in ocean subsurface temperature between the ESMs.

over all regions in observations and models determines the calibrated  $\gamma$  value. For the Amundsen calibration, the calibrated  $\gamma$  value is determined by the best fit between the modelled response and observations over only the Amundsen region. The resulting  $\gamma$  values are then applied to the other four regions to obtain the Antarctic summed response.

In addition, to assess the impact of our calibration method on the sea level projections, a single basal melt sensitivity (i.e. the calibration parameter  $\gamma$ ) derived from observed basal melt has been applied to all ESM-RF pairs. This calibration parameter is derived from the median basal melt sensitivity that was used in LARMIP-2 for the linear parameterisation and ISMIP6 AntMean method for the quadratic parameterisation (Table 3).

For all basal melt computation and calibration methods, the sea level contribution of the Amundsen region and the total AIS are analysed. The RMSE between observed and modelled ice discharge for these two regions was used to assess the impact of model selection on projections of the Antarctic dynamics contribution to sea level.

### 3 Results


#### 3.1 Basal melt computation and calibration

Basal melt is computed from subsurface thermal forcing anomalies from CMIP6 ESMs. The subsurface ocean temperature time series over the historical period are shown in Figure 3. Figure 4 shows the thermal forcing for part of the historical and future period (1950-2100). Over the 21st century, all regions show a median increase in thermal forcing but the magnitude varies between individual regions and becomes scenario dependent around year 2050.

The basal melt parameterisations are calibrated by fitting the sea level response of each ESM-RF pair on the changes in  
220 observed ice discharge over the full 1979-2017 period (Rignot et al., 2019). This exercise shows that the median basal melt  
sensitivity value resulting in the lowest RMSE differs between the Antarctic-wide calibration and calibration on the Amundsen  
region (Fig. 5). For the Amundsen region a higher median basal melt sensitivity than for the Antarctic-summed response  
improves the fit. The Antarctic-wide calibration includes regions with a small or negative past contribution to sea level, resulting  
in a lower basal melt sensitivity. The relatively high magnitude of the median basal melt sensitivity of the Amundsen region  
225 is consistent with the higher sensitivity to ocean warming as described in Dinniman et al. (2016). The contribution of ice  
discharge to sea level over the observational period is positive and (at least partly) attributable to ocean warming for both the  
Amundsen region and the total AIS (Pritchard et al., 2012). Therefore, for each ESM-RF pair the calibration parameter, and  
thus the basal melt sensitivity, should be positive for both Antarctica and the Amundsen region. If the best fit (lowest RMSE)  
is associated with a negative basal melt sensitivity, this means that the ESM-RF combination could not be calibrated. Between  
230 83% and 90% of all ESM-RF pairs could be calibrated, dependent on the parameterisation type and calibration region, as  
indicated on top of the boxes in Fig. 5. These percentages show that for the Antarctic-wide calibration region, the quadratic  
parameterisation has a higher percentage of positive values than the linear parameterisation. The boxplots only represent the  
ESM-RF pairs with positive basal melt sensitivities. These calibrated ESM-RF pairs are used in the hindcasts and projections  
of changes in ice discharge.

235 For the linear parameterisations, we compared our calibrated basal melt sensitivities to the values used in LARMIP-2 (Lev-  
ermann et al., 2020) (green shading in Fig. 5). This comparison shows that our Antarctic-wide calibration results in a median  
basal melt sensitivity just below the lower bound of the LARMIP-2 interval. Regional calibration on the Amundsen sector  
results in a median basal melt sensitivity above the LARMIP-2 range. Furthermore, the spread in our calibrated basal melt  
sensitivities is much larger than the spread in the observation-based range. For the Amundsen calibration, more than half of  
240 the ESM-RF pairs have a higher calibrated basal melt sensitivity than the observation-based LARMIP-2 range. These ESM-  
RF pairs will underestimate historical ice discharge when applying the lower, observation-based melt sensitivity. Vice versa,  
for the Antarctic-wide calibration, about half of the ESM-RF pairs have a lower calibrated sensitivity than the LARMIP-2  
range. These ESM-RF pairs will overestimate historical ice discharge when applying the higher melt sensitivity from within  
the LARMIP-2 range.

245 A similar comparison was made for the quadratic parameterisation, with the basal melt sensitivities applied in ISMIP6  
(Jourdain et al., 2020). Here, the median Antarctic-wide calibrated basal melt sensitivity sits at the lower end of the range of the  
Antarctic mean (AntMean) calibration option (blue shading in Fig. 5) applied in ISMIP6. The Amundsen calibration results in a  
median basal melt sensitivity at the top end of the AntMean range. In ISMIP6, also a calibration on Pine Island's grounding line  
basal melt was applied as an option (yellow shading in Fig. 5), which is the highest observed basal melt of the AIS. Only some  
250 calibrations of ESM-RF pairs outside the 95th percentile range resulted in  $\gamma$  values within the PIGL range. However, it should  
be remarked that the ISMIP6 PIGL calibration also includes negative ocean temperature corrections all around Antarctica that  
counter-balance the effects of the large  $\gamma$  values (Jourdain et al., 2020). Similar to the linear parameterisation, about half of the  
ESM-RF pairs has a calibrated melt sensitivity higher than the ISMIP6 AntMean range for the Amundsen calibration. These



**Figure 5.** Box-and-whisker plots of basal melt sensitivity values corresponding with the calibrated  $\gamma$  values of ESM-RF pairs. Only the sensitivities of calibrated  $\gamma$  values greater than zero are shown in the plot. The percentage of ESM-RF pairs with positive  $\gamma$  values is indicated by the green values on top of the boxes for each region. The horizontal orange line indicates the median value, boxes indicate the 25-75 percentile range and whiskers the 5-95 percentile range. Values beyond this range are not shown. The shaded regions indicate basal melt sensitivity ranges that are used in other studies. The green shading represents the basal melt sensitivity range of 7-16  $\text{m yr}^{-1} \text{K}^{-1}$  used in Levermann et al. (2020). The blue and yellow shading indicate the 5-95% range of the basal melt sensitivities corresponding with the  $\gamma$  values used for the nonlocal quadratic parameterisation in ISMIP6 (Jourdain et al., 2020) for both the Antarctic mean (AntMean) and Pine Island’s grounding line (PIGL) calibration option, respectively. For PIGL the 95% bound is 84  $\text{m yr}^{-1} \text{K}^{-2}$ , which is outside the scale of the vertical axis.

model pairs will underestimate historical ice discharge in the Amundsen region when applying the ISMIP6 AntMean basal melt sensitivity.

For the quadratic parameterisation, the sensitivity of the calibration parameter to the thermal forcing is tested. In this way, the impact of the uncertainty in the reanalysis dataset on the sea level projections is explored. This has been done by adding a positive temperature perturbation to the temperature time series near the ice shelf base of each ESM. The temperature perturbation is equal in size to one standard deviation between the reanalysis products (see the shaded orange regions in Fig 3). The resulting calibrated basal melt sensitivities are listed in Table 5 (Ice shelf base +  $1\sigma$ ). As expected, the higher ocean temperatures lead to stronger forcing in the quadratic parameterisation and therefore a lower basal melt sensitivity is required for the best fit with observations.

To summarise, a comparison of the calibrated basal melt sensitivity values in our study and equivalents in LARMIP-2 (Levermann et al., 2020) and the ISMIP6 AntMean method (Jourdain et al., 2020) suggests that calibration on past ice discharge

**Table 5.** Sensitivity of calibration parameter of the quadratic parameterisation to thermal forcing. Values indicate median basal melt sensitivity in  $\text{m yr}^{-1} \text{K}^{-2}$  for calibrated  $\gamma$  values based on three types of thermal forcing. The Antarctic-wide calibration (QA) and regional Amundsen calibration (QR) are shown. For comparison the median value of the AntMean calibration that is used in ISMIP6 (QM) is shown. The first thermal forcing type is the thermal forcing as shown in Fig. 3, which is based on the bias-adjusted ocean subsurface temperature timeseries of the ESMs near the ice shelf base. The second type is based on the same ocean temperature timeseries raised with one standard deviation ( $1\sigma$ ) that expresses the spread between the ocean reanalysis products (GREP  $\sigma$  in Table 1). The third type is the thermal forcing at 800-1000 m depth.

Thermal forcing	Antarctic-wide (QA) [ $\text{m yr}^{-1} \text{K}^{-2}$ ]	Amundsen (QR) [ $\text{m yr}^{-1} \text{K}^{-2}$ ]	ISMIP6 Antmean (QM) [ $\text{m yr}^{-1} \text{K}^{-2}$ ]
Ice shelf base	2.3	3.7	2.6
Ice shelf base + $1\sigma$	1.8	3.4	-
800-1000 m	1.2	5.5	-


265 rather than on basal melt observations results in relatively low basal melt sensitivities for the Antarctic-wide calibration. The Amundsen sector is more consistent with the high end of the basal melt sensitivity ranges applied in LARMIP-2 and the AntMean calibration option of ISMIP6. It should be noted that calibration on ice discharge leads by definition to a better fit with past ice discharge for individual ESM-RF pairs. Remarkably, the spread in the calibrated melt sensitivities is much higher than the observation-based ranges of LARMIP-2 and the ISMIP6 AntMean method. ESM-RF pairs with calibrated melt  
270 sensitivity values outside the observation-based ranges either underestimate or overestimate past ice discharge when using observation-based sensitivities.

### 3.2 Hindcasts of Antarctic and Amundsen sea level contribution

Hindcasts of the dynamic contribution of the Amundsen region and the total AIS to sea level rise are made to assess how well changes in ice discharge could be reproduced after calibration over the period 1979-2017. The calibration is performed  
275 by fitting the sea level on observations using a least squares fit of the sea level contribution for each year, weighted equally, over the hindcast period. The results of the linear and quadratic parameterisation are about equal when applied to the region of calibration (same RMSE; Table 6). However, the quadratic parameterisation performs better (lower RMSE) after calibration on an independent region than the linear parameterisation (i.e. when calibrated on the Amundsen region and applied to the total AIS or vice versa). Observations confirm that the quadratic relation can be better used when calibrating on (partly) independent  
280 regions (Jenkins et al., 2018). In the remainder of this article, therefore, our main results are based on the quadratic basal melt parameterisation. The linear parameterisation is used for making projections with the LARMIP2 median basal melt sensitivity (Sect. 3.3.1). The differences in the projections between the quadratic and linear parameterisation are further discussed in Sect. 3.3.2.

**Table 6.** RMSEs of the least squares fit of the median sea level contribution of each year, weighted equally, between calibrated results and ice discharge observations of Rignot et al. (2019). Results are shown for combinations of the two parameterisations, linear (L) and quadratic (Q) and two calibration methods, regional Amundsen (R) and Antarctic-wide (A), for two hindcast regions: Antarctic Ice Sheet (AIS) and the Amundsen region.

Basal melt method	RMSE AIS [mm]	RMSE Amundsen [mm]
Linear Amundsen	14.9	1.4
Quadratic Amundsen (QR)	7.2	1.4
Linear Antarctic-wide	1.7	2.7
Quadratic Antarctic-wide (QA)	1.6	2.4



**Figure 6.** Impact of calibration target region on sea level illustrated by hindcasts showing the sea level contribution over the period 1979-2017 based on all calibrated ESM-RF pairs for the total AIS (left panel) and the Amundsen region (right panel). The historical experiment is extended with SSP2-4.5 scenario for the years 2015-2017. The red lines indicate the median contribution based on the regional Amundsen calibration, whereas the blue lines indicate the median contribution for the Antarctic-wide calibration. Only the quadratic parameterisation with thermal forcing near the ice shelf base is shown. The observation-based changes in ice discharge from Rignot et al. (2019) are shown in grey. The shaded area indicates the associated likely range (17th to 83rd percentiles) for the modelled response and the observational error for the Rignot et al. (2019) data.

Figure 6 shows the hindcasts of all ESM-RF pairs using the calibrated basal melt sensitivities (Fig. 5). The two panels show the hindcasts for the total AIS and the Amundsen region, as specified in the titles. The total Antarctic sea level response is based on the summed contribution over the five sectors (Fig. 2). The colors represent two calibration methods, where red is the calibration on the Amundsen region and blue the Antarctic-wide calibration. The observed ice discharge values (Rignot et al., 2019) are shown in grey.

First, we evaluate the cumulative magnitude of the modelled sea level contributions over the period 1979-2017 (Table 7). The median Amundsen calibration overestimates the cumulative AIS contribution by about 30% whereas the median Antarctic-wide calibration underestimates the contribution by about 10%. For the Amundsen region, the cumulative contribution is underes-

**Table 7.** The median cumulative sea level contribution ( $\Delta S$ ) over the hindcast period 1979-2017 and the rate ( $dS/dt$ ) over the last decade (2008-2017) of the hindcast period for the two calibration methods (Amundsen and Antarctic-wide) and for the ice discharge observations of Rignot et al. (2019). Results are shown for the quadratic basal melt parameterisation with thermal forcing near the ice shelf base.


Source	AIS		Amundsen region	
	$\Delta S$ [mm]	$dS/dt$ [mm yr <sup>-1</sup> ]	$\Delta S$ [mm]	$dS/dt$ [mm yr <sup>-1</sup> ]
Ice discharge observations	13.1	0.58	9.7	0.48
Amundsen calibration (QR)	17.5	0.84	7.6	0.27
Antarctic-wide calibration (QA)	11.8	0.45	4.3	0.17
Amundsen calibration (QR) - top 10%	16.6	0.86	9.3	0.44
Antarctic-wide calibration (QA) - top 10%	13.3	0.60	5.3	0.24

295 estimated by the median response of the Amundsen calibration (ca. 20%) and strongly underestimated by the Antarctic-wide calibration (ca. 60%). Both calibration methods do not give an agreement in terms of the cumulative sea level contribution because of the choice to calibrate on the time series rather than on the cumulative sum. Even though the Antarctic-wide calibration is (by construction) closer to the observed Antarctic ice discharge than the Amundsen calibration, the strong underestimation of the Amundsen region still means that the response in other regions is overestimated. It should be kept in mind that the errors in the individual regions compensate each other, resulting in a summed Antarctic response that is close to observations.

300 Second, we evaluate the evolution of the sea level response over time. For the Antarctic-wide calibration, the median value overestimates changes in Antarctic discharge before 2001 and underestimates them thereafter. This means that the sea level acceleration over the full period cannot be captured with the Antarctic-wide calibration, making it likely that it will be underestimated in future projections as well. This is also visible in the ice discharge rate over the last decade of the hindcast (Table 7), which is lower than in observations. In a similar way, the Amundsen calibration overestimates the changes in Amundsen discharge before 2005 and underestimates them thereafter. So for the Amundsen region, even when using the Amundsen-specific calibration, the acceleration is not captured by the median response and the rate over the last decade of the hindcast is underestimated. It should be noted that not just the acceleration of the Amundsen contribution cannot be reproduced, but the relative dominance of Amundsen with respect to the total Antarctic contribution cannot be reproduced either (about 70% in observations, about 30-40% in our results).

310 Since the Amundsen region is the most important contributing region to the summed Antarctic response over the hindcasting period, we tested whether a selection of models could better capture past ice discharge in the Amundsen region. The top 10% calibrated models with the best fit to ice discharge observations (Fig. A1) were selected for both the Amundsen and Antarctic-wide calibration. The selection was based on the model performance in the calibration region. As a logical consequence, the top 10% ESM-RF pairs from the two calibration methods performs better on the cumulative sea level contribution in






**Figure 7.** Projections showing the calibrated sea level contribution over the period 2000-2100 based on SSP5-8.5, for the total AIS (left panel) and the Amundsen region (right panel). The red lines indicate the median contribution based on the regional Amundsen calibration, whereas the blue lines indicate the median contribution for the Antarctic-wide calibration. Results are shown for the quadratic parameterisation and thermal forcing near the ice shelf base. The shaded area indicates the associated likely ranges (17th to 83rd percentiles).

the calibration region (Table 7). Interestingly, the same selection of models also performs better in the region that was not used for the calibration. After Antarctic-wide selection the Amundsen sea level contribution in the hindcasts are closer to observations. Unfortunately, for the Antarctic-wide selection, the contributions of the other regions increase as well, which increases their error relative to observations. The Amundsen selection resulted in higher estimates than for the full model suite in the Amundsen region itself (by construction), but lower estimates (closer to observations) for the Antarctic summed response. As a result, the Amundsen contribution relative to the total AIS improves after model selection on the Amundsen region. Nevertheless, the mean response of the top 10% models could not reproduce the observed acceleration over the historical period in the Amundsen region. This means that despite its overestimation of the cumulative sum over the hindcast period for the AIS, the Amundsen calibration will presumably underestimate future projections of the sea level contribution for the Amundsen region.

### 3.3 Sea level contribution projections

In this section, projections of the sea level contribution due to basal melt for the AIS and the Amundsen region are presented. The projections comprise the 21st century. Computations start in the year 1850 so that the delayed contribution of ice discharge due to basal melt is included in the future sea level response. Figure 7 shows our main projections for the SSP5-8.5 scenario, based on the calibrated basal melt sensitivities for the quadratic parameterisation and thermal forcing near the ice shelf base. We assess two metrics: the cumulative magnitude and the rate of the sea level response. The cumulative sea level response is computed by taking the difference between the year 2100 and the average over the period 1995-2014. The sea level response rate at the end of the 21st century is indicative of differences in committed sea level rise beyond 2100. The sea level response rate is computed by a linear regression on the sea level response over the period 2081-2100.

First, we present the calibrated projections for the three SSP scenarios and explore the impact of calibration on projections of the sea level contribution. Second, the sensitivity of projections to methodological choices, such as the parameterisation relation (quadratic/linear), thermal forcing depth (ice shelf base/800-1000 m) and model selection (Earth system model/Ice sheet model) is explored.



**Figure 8.** Projected Antarctic sea level response for SSP1-2.6, SSP2-4.5 and SSP5-8.5. Top panels show the sea level contribution in 2100 compared to the period 1995-2014 and bottom panels the sea level rise rates over the period 2081-2100. The spread is determined by the calibrated ESM-RF pairs. The green numbers indicate the median values (corresponding with the green lines), whereas the boxes show the 25-75 percentiles and the whiskers the 5-95 percentiles. The left hand side shows projections using a single median basal melt sensitivity from the ISMIP6 AntMean method (QM) and from LARMIP-2 (LM). The basal melt computation methods on the right hand side are our main projections with calibrated basal melt sensitivities on ice discharge observations of the Amundsen region (QR) and the total AIS (QA). ESM-RF pairs that could not be calibrated are removed from all basal melt methods so that the same models are included in the comparison. If ESMs did not simulate year 2100, 2099 was used instead.

### 3.3.1 Impact of calibration on sea level projections

To understand how calibration of individual ESM-RF combinations on past ice discharge influences the results compared to using observation-based basal melt sensitivities, we also made projections in which a single basal melt sensitivity is applied in all ESM-RF combinations. This single value is the median basal melt sensitivity applied in LARMIP-2 ( $11.5 \text{ m yr K}^{-1}$ ) (Levermann et al., 2020) for the linear parameterisation (LM) and the median nonlocal basal melt sensitivity applied in ISMIP6 for the AntMean method ( $2.6 \text{ m yr K}^{-2}$ ) (Jourdain et al., 2020) for the quadratic basal melt parameterisation (QM). The resulting projections from these basal melt computation methods are included in Figures 8 and 9. In these figures, the green numbers


correspond with the median values of the projections. The median projected values are used to quantify the impact of the basal melt method on the sea level projections.

345 First, the sea level contribution of the total AIS is analysed. Figure 8 shows the projected sea level response for each SSP scenario and different basal melt computation methods. The computations methods include the median MIP basal melt sensitivities (QM, LM) and the calibrated sensitivities (QA and QR). The top panels represent the cumulative projections and the bottom panels the sea level response rate over the period 2081-2100. Not surprisingly, a higher emission scenario leads to a higher sea level contribution. Absolute differences between the basal melt computation methods within one SSP scenario  
350 become more explicit for the higher emission scenarios, but relative differences within one SSP scenario are comparable. To compare relative differences we use the ratio of the sea level projections between the highest and lowest basal melt method, which is QR/QA for the AIS sea level contribution. The ratio QR/QA (1.6) is only slightly larger than the ratio between the SSP5-8.5 and SSP1-2.6 scenario (1.4; averaged over all methods), indicating that the influence of the basal melt computation method on the sea level response is more or less similar to the impact of the emission scenarios. Since the highest sea level pro-  
355 jections result from the Amundsen calibration method and the lowest sea level projections from the Antarctic-wide calibration method, this means that this difference can be entirely attributed to the calibration region.

The projections of the AIS using the median basal melt sensitivities applied in ISMIP6 (QM) and LARMIP-2 (LM) fall in between the two calibrated projections. This is consistent with the median basal melt sensitivity of LARMIP-2 and ISMIP6, which is located above the median Antarctic-wide calibrated value and below the Amundsen-calibrated value, respectively  
360 (Fig. 5). Even though the spread between the basal melt methods is extended by using the calibration methods, using single basal melt sensitivities based on basal melt observations with different parameterisation types (linear/quadratic) also leads to a large spread in the projections ( $LM/QM = 1.3$ ; averaged over all SSPs). We remark that the top 10% best-performing models in reproducing ice discharge observations (Fig. A1), result in estimates that fall in between the Antarctic-wide (QA) and Amundsen calibration (QR) methods, reducing the spread (Fig. A2).

365 As a next step, the AIS sea level response rates are assessed at the end of the 21st century (2081-2100). These are important for sea level differences beyond 2100. The ratio QR/QA for the sea level response rates (1.6; averaged over all SSPs) shows that the influence of the calibration region on the response rate is smaller than the effect of the SSP scenarios ( $SSP5-8.5/SSP1-2.6 = 2.1$ ; averaged over all basal melt methods). The effect of the SSP scenarios is stronger for the quadratic parameterisations (QM, QA, QR) than for the linear one (LM). Consequently, the highest median response rate in SSP5-8.5 and SSP2-4.5 is using the QR basal melt method, whereas in SSP1-2.6 the response rate based on the median LARMIP-2 basal melt sensitivity  
370 (LM) is highest. This could be explained by the linear (rather than quadratic) relation with thermal forcing (see Sect. 3.3.2), which is independent on the absolute ocean temperature (which is linked to the SSP scenarios). It should also be noted that the Amundsen calibration is more skewed towards higher sea level response rates than the other basal melt methods. This is a result of the higher basal melt sensitivities that were required to fit the modelled historical Amundsen sea level contribution to  
375 ice discharge observations.

Second, the sea level projections of the Amundsen region are analysed (Fig. 9). For the Amundsen region, the highest projection is given by the Amundsen calibration, whereas the lowest projection is based on the median LARMIP-2 basal melt



**Figure 9.** Same as Fig. 8 but for the Amundsen region.

method. The ratio of the highest to lowest basal melt method ( $QR/LM = 1.9$ ) is larger than the ratio between the SSP5-8.5 and SSP1-2.6 scenario (1.3; averaged over all methods), indicating that the influence of the basal melt computation method on the sea level response is larger than the impact of the SSP scenarios. Also for the Amundsen sea level response rates, the impact of the basal melt method ( $QR/LM = 2.1$ ) is slightly larger than the impact of the SSP scenario ( $SSP5-8.5/SSP1-2.6 = 1.8$ ). This demonstrates that the rate is much more sensitive to the SSP scenario than the cumulative sum, indicating increasing differences between SSP scenarios beyond 2100.

The Amundsen calibration is considered to give the most realistic estimate for future projections of ice discharge in the Amundsen region. Considering the strong underestimation of past ice discharge rate in the Amundsen region using the Antarctic-wide calibration (Table 7), we expect that the future projections for the Amundsen region will be too low when using this method. The Amundsen projections using the median LARMIP-2 basal melt sensitivity (LM) are lower than for the Antarctic-wide calibration method and therefore are also expected to underestimate the sea level contribution of the Amundsen region. The projection based on the median ISMIP6 sensitivity (QM) is probably also too low, since even the hindcasts based on the Amundsen calibration slightly underestimated observed ice discharge in the Amundsen region (Table 7).

We conclude that for the AIS the cumulative sea level variations associated with basal melt computation methods are about equal to variations between different SSP scenarios. For the Antarctic sea level response rate, the SSP scenario is more impor-

**Table 8.** Projected dynamic contributions to sea level in meters from the AIS in 2100 relative to 1995-2014. The numbers for LARMIP-2, ISMIP6 and SMB are obtained from the IPCC AR6 report (Fox-Kemper et al., 2021). Note that for the ISMIP6 estimate surface mass balance contributions are removed as our study only accounts for changes in ice discharge. The columns show the 17th, 50th and 83rd percentiles of the distribution.

Scenario	Forcing/Source	17%	50%	83%
SSP5-8.5/RCP8.5	Antarctic-wide calibration (QA)	0.06	0.11	0.19
	Amundsen calibration (QR)	0.09	0.17	0.41
	Median ISMIP6 sensitivity (QM)	0.05	0.12	0.27
	Median LARMIP-2 sensitivity (LM)	0.08	0.15	0.32
	ISMIP6 AR6 (excl. SMB)	0.10	0.13	0.17
	LARMIP-2 AR6	0.10	0.20	0.39
SSP2-4.5/RCP4.5	Antarctic-wide calibration (QA)	0.05	0.09	0.16
	Amundsen calibration (QR)	0.07	0.14	0.34
	Median ISMIP6 sensitivity (QM)	0.04	0.10	0.22
	Median LARMIP-2 sensitivity (LM)	0.06	0.12	0.26
	ISMIP6 AR6 (excl. SMB)	0.07	0.12	0.16
	LARMIP-2 AR6	0.09	0.17	0.33
SSP1-2.6/RCP2.6	Antarctic-wide calibration (QA)	0.04	0.07	0.14
	Amundsen calibration (QR)	0.06	0.12	0.28
	Median ISMIP6 sensitivity (QM)	0.04	0.08	0.19
	Median LARMIP-2 sensitivity (LM)	0.06	0.11	0.23
	ISMIP6 AR6 (excl. SMB)	0.06	0.11	0.15
	LARMIP-2 AR6	0.08	0.15	0.29

395 tant than the basal melt method. In contrast, for the Amundsen region the basal melt method impacts the projections (cumulative sum and rate) more than the SSP scenarios. For the Amundsen region, we also conclude that the Amundsen calibration probably gives the most reliable projections since the Amundsen calibration already underestimated past ice discharge and its acceleration in the hindcasts, and the other methods give even lower estimates.

400 Furthermore, we compared our estimates with the emulated ISMIP6 and LARMIP-2 studies as presented in IPCC AR6 (Table 8). Despite the different method applied, the resulting projections of Antarctica’s sea level contribution are in line with previous multi-model studies (ISMIP6, LARMIP-2). The Amundsen calibration results in median estimates of 0.17 m for SSP5-8.5, 0.14 m for SSP2-4.5 and 0.12 m for SSP1-2.6, sitting in between the ISMIP6 and LARMIP-2 projections, as presented in IPCC AR6 (Fox-Kemper et al., 2021). It should be noted that this can only partly be attributed to the calibration on ice discharge observations, since our projections using the median ISMIP6 AntMean sensitivity (QM) and the median

LARMIP-2 sensitivity (LM) result in lower estimates than for ISMIP6 AR6 and LARMIP-2 AR6, respectively, which could be attributed to methodological differences other than the basal melt sensitivity. The differences with ISMIP6 and LARMIP-2 will be further discussion in Sect. 4.

### 3.3.2 Impact of methodological choices on projections


In this section we explore what the impact is of several methodological choices on the sea level response projections of the AIS and Amundsen region. These choices include the parameterisation relation (quadratic/linear), thermal forcing depth (ice shelf base/800-1000 m) and model selection (Earth system model/Ice sheet model). Additionally, we further motivate our choices to use the quadratic parameterisation with thermal forcing near the ice shelf base in our main projections (Fig. 7; QA and QR in Figs. 8 and 9).

First we assess the impact of the parameterisation type on the calibrated projections for the AIS and the Amundsen region (Fig. 10). To this end we applied two different parameterisations: a linear (Eq. 2) and a quadratic relation (Eq. 3) with thermal forcing. Both relations are calibrated on observed ice discharge (Fig. 5) using the Antarctic-wide and the Amundsen calibration. The results show that if the parameterisation is used to make projections for the same region as the region that is used for calibration, the cumulative sea level contribution is almost equal for both parameterisations. This means that calibration on past ice discharge strongly constrains the future response if applied to the region of projections.

On the other hand, if the calibration is performed in the Amundsen region and applied to make Antarctic projections, or vice versa, clear differences between the linear and quadratic relation appear. For the Amundsen calibration, the quadratic parameterisation results in lower projections for the Antarctic-wide contribution than the linear parameterisation. This can be expected, since the quadratic parameterisation is dependent on the absolute ocean temperature, whereas the linear parameterisation only uses temperature anomalies. By its definition the quadratic relation with thermal forcing implies that sectors that are melted by warmer waters are more sensitive than the colder sectors, even if the same basal melt sensitivity is applied. So if the Amundsen calibration is applied to colder ocean sectors than the Amundsen sector, this leads to less basal melt for a similar temperature increase, since the ocean temperatures are lower. In a similar way, Antarctic-wide calibration of the linear parameterisation leads to a lower basal melt sensitivity and thus lower projections for the Amundsen region than the quadratic parameterisation.

Favier et al. (2019) demonstrate that the quadratic parameterisation gives better results in representing ocean-induced melting under ice shelves than the linear forcing when compared with ocean-ice-sheet coupled simulations. Furthermore, Holland et al. (2008) show with an ocean model that total ice shelf basal melt increases quadratically as the ocean offshore of the ice front warms. Moreover, the quadratic relationship between thermal forcing and basal melt is confirmed by observations (Jenkins et al., 2018). These arguments are an important motivation to apply the quadratic parameterisation in our study.


Second, we assessed the impact of the thermal forcing depth on the calibrated projections (Fig. 11). For this experiment, thermal forcing and basal melt sensitivity are based on ocean temperature at two different depths: 100 m centered around the mean depth of the ice shelf base (similar to LARMIP-2) and an ocean layer around the depth of the continental shelf near the ice shelf front. The deeper ocean layer is chosen for comparison since the relevant water masses that drive the melting



**Figure 10.** Projections of Antarctic sea level contribution for SSP5-8.5 for all calibrated ESM-RF combinations for the AIS (left) and Amundsen region (right). Results are shown for thermal forcing near the ice shelf base. The bars show the median projections for the Antarctic-wide and regional Amundsen calibration using the quadratic (orange) and linear (blue) parameterisations. The spread indicates the 17th to 83rd percentiles.

close to the grounding line originate from the deepest depth of the bed near the ice shelf front, which we approximate as 800-1000 m. We only use the quadratic parameterisation, which is dependent on the absolute ocean temperature. Surprisingly, for the deeper layer, the Antarctic-wide calibration leads to a lower basal melt sensitivity, whereas the Amundsen calibration leads to a higher basal melt sensitivity than the corresponding basal melt sensitivities near the ice shelf base (Table 5). This can be explained by the differences in the water temperature and the warming rates of the two layers. For the Amundsen region, the ocean temperature in the deeper 800-1000 m layer warms slower than the ocean temperature near the ice shelf base (Fig. A3), although the temperature itself is comparable in magnitude. Therefore, a higher basal melt sensitivity is required to match ice discharge observations. In contrast, for all other regions, the ocean layer at 800-1000 m depth is warmer than the temperature near the depth of the ice shelf base, resulting in a higher ocean forcing. In the Weddell, Ross and the Peninsula regions, the temperature also warms faster in the deeper layer than in the layer at the depth of the ice shelf base, resulting also in stronger ocean forcing. Due to the stronger ocean forcing in the 800-1000 m depth layer, the calibrated basal melt sensitivity is lower for the Antarctic-wide calibration.

For the AIS projections, the lower Antarctic-wide basal melt sensitivity for 800-1000 m depth is largely compensated by a larger ocean forcing for the Antarctic-wide calibration. This results in a similar sea level contribution for the 800-1000 m-based projections compared to using the thermal forcing near the depth of the ice shelf base. However, the high Amundsen basal melt sensitivity for the 800-1000 m depth combined with the larger Antarctic-wide ocean forcing leads to higher estimates for the AIS projections. Projections for the Amundsen region are oppositely affected. The ocean forcing is smaller at 800-1000 m depth than near the ice shelf base, and combined with a lower basal melt sensitivity for the Antarctic-wide calibration this leads to much smaller projections. For the Amundsen region itself, the higher basal melt sensitivity only partly compensates for the smaller ocean forcing, resulting in a smaller sea level projection for the forcing at 800-1000 m compared to forcing near the ice shelf base. As a result, the fraction of Amundsen compared to the total Antarctic contribution is larger for the thermal forcing near the ice shelf base than for the 800-1000 m depth layer. Since this fraction was already smaller than in observations



**Figure 11.** Projections of the sea level contribution of the AIS (left) and Amundsen region (right) for SSP5-8.5 for all calibrated ESM-RF combinations using the *quadratic* parameterisation. The bars indicate the median sea level contribution in 2100 relative to 1995-2014. The thermal forcing and basal melt sensitivity are based on ocean temperature at two different depths: 100 m centered around the mean depth of the ice shelf base (blue) and 800-1000 m depth (orange). The black lines indicate the 17th to 83rd percentiles.

in the hindcast experiments using thermal forcing near the ice shelf base (Sect. 3.2), we argue that using thermal forcing near the ice shelf base leads to more realistic results than thermal forcing in the 800-1000 m depth layer.

We conclude that the depth of thermal forcing has a large influence on the resulting sea level contribution in future projections. Most straightforward, it influences the thermal forcing in the projections, which is depth-dependent, but also region-dependent. However, when calibration is applied, the thermal forcing depth also affects the strength of the basal melt sensitivity through its evolution over the historical period. The thermal forcing near the ice shelf base leads to a more realistic contribution of the Amundsen region compared to the total AIS, and is therefore applied throughout this study.

### 3.3.3 Modelling uncertainties associated with Earth System and Ice Sheet Models

In this section, we assess the role of CMIP6 ESMs and RFs of the LARMIP-2 ice sheet models in projection uncertainties for the AIS by comparing the sea level contributions for the Amundsen calibration, which is considered to perform better than the Antarctic-wide calibration for the Amundsen region (Sect. 3.2) and arguably also for the total AIS contribution (Sect. 4). These models cause the spread of the projections for a specific basal melt method (see the shaded regions in Fig. 7 and the error bars in Figs. 8-11). Fig. 12 shows the projected Antarctic sea level contribution for each individual CMIP6 ESM for the Amundsen calibration. Here, the spread for each ESM is determined by the linear response functions of the ice sheet models. Noticeably, the differences between the scenarios are small compared to the differences between individual ESMs, despite the bias adjustment with ocean reanalysis data. As a measure of ESM spread, we compute the standard deviation between the median values (bar heights). The intermodel standard deviation varies from 144 mm for SSP1-2.6 to 205 mm for SSP5-8.5.

The ESM with the strongest median sea level contribution (CAS-ESM2-0) also exhibits the largest warming over the 21st century for each individual ocean sector and has the second highest median calibrated basal melt sensitivity for the Amundsen region (not shown). Also, it has the fourth lowest ranking in reproducing historical ice discharge compared to the other ESMs. Remarkably, the five ESMs with the highest RMSE for the Amundsen region (when comparing their historical performance



480 to ice discharge observations) are amongst the six models with the highest cumulative sea level contribution for the AIS in the projections. This suggests that applying ESM selection based on the performance of ESMs in reproducing ice discharge observations in the Amundsen region would result in lower estimates of the Antarctic dynamics contribution to sea level projections. However, a potential selection of CMIP6 ESMs based on ice discharge can only be considered if the sensitivity of ice discharge to basal melt perturbations is well represented by the linear response functions (Sect. 4).


485 Fig. 12 also shows the projected Antarctic sea level contribution for the RF of each individual ice sheet model. Here, the spread in the error bars is determined by the CMIP6 ESMs. The RF spread is also greater than the scenario-induced spread. Similar as for the ESMs, we computed the intermodel standard deviation between ice sheet models as a measure of ice sheet model spread. The standard deviation between the median values varies from 46 mm for SSP1-2.6 to 62 mm for SSP5-8.5.

The RF of the ice sheet model giving the smallest median sea level contribution (GRIS LSC) has the second lowest calibrated  
490 basal melt sensitivity for the Amundsen region and could not be calibrated in combination with half of the ESMs. We remark that this RF also gave the smallest signal in LARMIP-2 (Levermann et al., 2020). The RF of the ice sheet model with the smallest calibrated basal melt sensitivity (PISM DMI) also could not be calibrated when combined with the forcing for 6 out of the 14 ESMs. Moreover, GRIS LSC and PISM DMI have the highest RMSE when compared with observed ice discharge. This suggests that RF selection based on reproducing historical ice discharge would result in higher future estimates of the sea  
495 level contribution.

We also compared the spread associated with the ESMs and RFs with the spread in the emission scenarios and basal melt methods. This was done by computing the standard deviation between the median estimates of the Amundsen calibration (QR) for the three SSP scenarios (28 mm for QR) and the standard deviation between the median estimates of the four basal melt methods for each SSP scenario (21 mm for SSP1-2.6 to 31 mm for SSP5-8.5). The spread between ESMs and RFs is thus  
500 larger than the spread between the three SSPs and four basal melt methods.

As a final assessment, the RMSE over the Amundsen region was used to rank the historical performance of individual combinations of ESM-RF pairs. The top 10% best-performing ESM-RF pairs have slightly lower estimates for the Antarctic contribution but similar estimates for the Amundsen contribution (Fig. A2). As a result the relative contribution of the Amundsen region increases compared to the total Antarctic dynamics contribution to sea level, as was also visible in the hindcasts of  
505 the top 10% models (Table 7).

To summarise, this assessment of individual models shows that modelling uncertainties of ESMs as well as ice sheet models are a greater source of uncertainties in Antarctic mass loss projections than the emission scenarios and the basal melt computation methods applied in this study. The uncertainties associated with the ocean temperature evolution from ESMs is even larger than those from ice sheet models, despite the bias adjustment that has been applied to the subsurface temperatures. We also  
510 find some relations between historical model performance and future projections, which point at model selection as a potential next step to better understand the future contribution of Antarctic dynamics to sea level changes.



**Figure 12.** Projected Antarctic sea level changes for SSP1-2.6 (blue), SSP2-4.5 (orange) and SSP5-8.5 (red) over the 21st century, defined as the difference between year 2100 and the period 1995-2014. The top panel shows the projections for each CMIP6 ESM, where the errorbars indicate the 17th to 83rd percentiles (computed from the associated RF timeseries). The bottom panel shows the projections for each RF, where the errorbars indicate the 17th to 83rd percentiles (computed from the associated ESMs). Basal melt is computed with the quadratic parameterisation which is calibrated on the Amundsen region (QR). Note the differences in the vertical scale.

#### 4 Discussion

In this study, projections of the sea level contribution of the AIS and the Amundsen region are presented that were calibrated on four decades of ice discharge observations. Calibration was applied on the basal melt parameterisation. The contribution of Antarctica's ice discharge to sea level changes is computed using ocean forcing from state-of-the-art ESMs from Coupled Model Intercomparison Project Phase 6 (CMIP6) applied to linear response functions from LARMIP-2 ice sheet models. The major strength of this method is that multiple climate and ice sheet models can be combined to assess their full range of modelling uncertainties. A drawback of the method is that non-linearities between thermal forcing and ice sheet mass loss, related to ice sheet instabilities and ocean dynamics, are not considered because we use the linear response functions framework.

Consistent with Levermann et al. (2020), the ocean sectors in our study are somewhat wider than the continental shelf. The advantage of a wider region is that it allows for more assimilated observations in the reanalysis product that is used for the bias adjustment of ocean temperature (the continental shelf region is only sparsely sampled). Furthermore, it should also be noted that we used basal melt anomalies and not absolute basal melt in the computation and calibration of the sea level contribution.

525 This is because that allows us to better represent observed melt but the downside is that anomalies are a second order effect that is harder to model and observe. We also remark that the linear response functions are derived from ice sheet model experiments with an homogeneous basal melt increase over each entire ice shelf. Therefore, apart from the five regions for which the linear response functions were derived, no spatial patterns and effects are taken into account.

The inability of our models to represent the observed acceleration (Fig. 6) could be explained by ice sheet/ocean feedbacks that are not represented in the models. Recent studies suggest a positive feedback between ice sheet melting and subsurface ocean warming (Bronselauer et al., 2018; Golledge et al., 2019; Sadai et al., 2020) that could explain this deficiency in the models. One reason to introduce the quadratic parameterisation was to account for the observed non-linear relation between ice melt and ocean forcing (Jenkins et al., 2018). However the feedback between surface freshening due to meltwater release, subsurface warming, and enhanced basal ice shelf melt is not represented by this parameterisation. It should also be noted that

530 our study does not address the impact of surface melt on calving nor marine ice cliff instability processes that would lead to higher projections.

In the current generation of ESMs (CMIP6) ice shelf cavities are not represented, leading to deficiencies in the representation of ocean currents and ice-ocean interactions (Mathiot et al., 2017). Including ice shelf cavities in ESMs would better resolve how the inflow/ambient temperature is affected by mixing with meltwater and ocean dynamical processes inside the cavity.

540 Also, the resolution of most CMIP6 ESMs is not high enough to resolve the ocean circulation on the continental shelf, including the Antarctic Slope Current (Thompson et al., 2018). This could lead to a mismatch between observed and simulated ocean warming in the coastal regions. Due to these ocean model deficiencies, temperature-melt relations are typically parameterised (Favier et al., 2019). We have chosen to use a simple quadratic scaling with far-field thermal forcing (Eq. 3), which could be calibrated on the heat exchange velocity  $\gamma$  and applied to all models. This parameterisation performs relatively well when

545 compared with ocean-ice sheet coupled simulations (Favier et al., 2019). The quadratic relation between ice shelf basal melt and thermal forcing is also confirmed by ocean model experiments (Holland et al., 2008) and observations (Jenkins et al., 2018).

Calibration of the  $\gamma$  value in the basal melt parameterisation results for 10-17% of the ESM-RF pairs in a value of zero, which means that, in some cases, the calibration method is invalid. However, we found that each ESM could lead to a successful

550 (positive  $\gamma$ ) calibration if combined with several RFs, so it is the ESM-RF combination which determines whether calibration is successful. Unsuccessful calibration occurs when the ESM produces large historical natural variability, and the lagged response in the RF translates this into a reduced mass loss over the specific period. In these cases, the ESM produces a weak signal-to-noise ratio in terms of historical warming (the observation period is too short). Overall, the calibration of each ESM-RF pair is dependent on the magnitude and phasing of natural variability (in ocean temperatures and observed mass loss). For the

555 calibrated ESM-RF combinations, the large number of pairs reduces the impact of natural variability on the resultant calibrated  
projections.

Calibrating the basal melt parameterisations on observed ice discharge is a way to get more correct historical sea level trends,  
which was not assessed in ISMIP6. Calibration of individual ESM-RF pairs increased the spread in basal melt sensitivities but  
decreased spread in the hindcast experiments of Antarctica's sea level contribution. Unfortunately, calibration of the basal melt  
560 relation on ice discharge did not reduce the spread in future projections of the ice dynamics contribution to sea level compared  
to using observation-based basal melt sensitivities. However, the ice sheet models used to derive the response functions could  
all be biased in the same direction, resulting in a too high or too low sensitivity to changes in basal melt. For example, if the ice  
sheet models are not sensitive enough to basal melt perturbations, calibration will result in high-biased melt rates to compensate  
the low-biased sensitivity. In this case, getting the correct historical ice discharge would not give so much confidence that the  
565 response to future warming is correct.

To compute projected sea level change, we have made the assumption that the calibrated gamma values are constant. There  
are, however, reasons to assume that basal melt sensitivities will change in the future. In the projections (Fig. 4), especially  
for SSP5-8.5, all coastal regions, especially the Weddell and Ross sectors, experience a warming signal which is not present  
in the historical period. As the open ocean outside the cavities warms, it could be expected that this warming will at a certain  
570 moment also be transported inside the cavities, and contribute there to basal melt and ice discharge. New calibration will then  
lead to larger Antarctic-wide basal melt sensitivities. This means that calibrated basal melt sensitivities that link open ocean  
subsurface temperatures outside cavities to basal melt underneath ice shelves could be time-evolving.

In this study, an Antarctic-wide and regional Amundsen calibration of the basal melt parameterisation have been applied.  
The relation between thermal forcing and basal melt is more difficult to derive for the full AIS. The reason is that it includes  
575 regions in which ocean warming has not been causally linked to changes in ice dynamics as the warming was too small or  
absent over the historical period. However, regions with small ice discharge during the calibration period are expected to melt  
as the climate warms. The calibration could therefore result in a basal melt sensitivity which is too low for future projections.  
Moreover, calibrating on the Antarctic-wide response strongly underestimates the historical mass loss in the Amundsen region,  
which accounts for more than 70% of the observed historical sea-level contribution. Therefore, the Antarctic-wide calibration  
580 gives information about a lower bound for the future projections: i.e. what would happen if the total AIS would keep the  
same basal melt sensitivity to ocean warming in the future. The Amundsen region is considered the best region for calibration  
since it has been shown that the Amundsen mass loss is dominated by ice discharge due to basal ice shelf melting (Pritchard  
et al., 2012). Previous studies have shown that ice dynamical changes were causally linked to ocean warming during the  
observational record (Rignot et al., 2019). It could be expected that when ocean temperatures increase and experience similar  
585 warming rates in other regions, the basal melt sensitivity will also increase in those regions. It should also be noted that  
the quadratic parameterisation does introduce some regional difference in basal melt sensitivity due to its dependence on  
the absolute temperature, resulting in a lower sensitivity in colder cavities. When the high basal melt sensitivities derived  
from the Amundsen calibration are applied to the other regions, the resulting basal melt will thus be smaller due to the colder  
temperatures. The nonlinear relation between melt and temperature change found in observations (Jenkins et al., 2018) suggests

590 that the quadratic relation based on the Amundsen region might be applicable to the cold-water sectors, although individual regions might still respond differently to similar forcing due to differences in ice and ocean dynamics and ice geometries. The Amundsen calibration is therefore considered more reliable for future projections of the total AIS than the Antarctic-wide calibration, even though it overestimates the total Antarctic contribution to sea level over the historical period.

The AIS projections using our methodology with median MIP sensitivities (LM, QM; Fig. 8) resulted in lower projections  
595 than in the original MIPs as presented in AR6 (Table 8). The differences between the AIS projections using our methodology with median MIP sensitivities and the original MIPs can be attributed to differences in thermal forcing and modelling of the ice sheet response. It could thus be expected that calibration of the basal melt parameterisation in ISMIP6 and LARMIP-2 on the Amundsen region will result in higher projections of the Antarctic sea level contribution than the projections presented in IPCC AR6. We used a different set of ESMs, which can lead to large differences in the modelled response (see Sect. 3.3.3).  
600 These large intermodel differences in ESMs point at model selection as a promising next step to reduce uncertainties in future projections of the contribution of ice dynamics to sea level changes. Since we only used temperature anomalies from ESMs as forcing, the selection criteria should not be based on the mean climate but on climate trends. Furthermore, LARMIP-2 uses global mean temperature as the driver of the method, whereas we use bias-adjusted ocean temperature from the ESMs. The methodological differences with ISMIP6 AR6 are even larger than for LARMIP-2 since ISMIP6 does not use the linear  
605 response functions framework but runs offline ice sheet models to account for the ice sheet response. Despite all these differences in methodology, we arrive at projections which are in line with previous multi-model assessments of the contribution of Antarctic mass loss to future sea level.

## 5 Conclusions

This study presents calibrated projections of the contribution of Antarctica's ice discharge to sea level in 2100 compared to  
610 present-day (1995-2014). Since there is still high uncertainty in the temperature-basal melt relation (Dinniman et al., 2016), we applied a new approach to constrain this relation (Fig. 1). This was done by calibrating the modelled response on ice discharge observations rather than observation-based estimates of basal melt. The new projections of the sea level contribution are therefore constrained by historical ice discharge observations of the Amundsen region and the total Antarctic ice sheet. Ocean thermal forcing is based on regional subsurface ocean temperature from 14 CMIP6 ESMs and 3 SSP scenarios and  
615 bias-adjusted with GREP ocean reanalysis data. The changes in ice discharge are calculated with 16 linear response functions (RF) based on ice sheet model experiments from LARMIP-2.

The results show that a large part of the calibrated basal melt sensitivities are higher than those derived from melt obser-  
vations, which is related to a wider spread in the calibrated basal melt sensitivities. The median basal melt sensitivities from calibration on ice discharge are for the Amundsen (Antarctic-wide) calibration higher (lower) than the median values applied  
620 in ISMIP6 (AntMean method) and LARMIP-2. However, even with calibration on past ice discharge, the acceleration of the sea level contribution during the observational period is underestimated for the Amundsen region, indicating missing physics. Also the relative contribution of the Amundsen region to the AIS sea level contribution is underestimated.

We find that the depth of thermal forcing has a large influence on the resulting sea level contribution in future projections. In our study we applied the same thermal forcing depth as in Levermann et al. (2020), which is the forcing near the ice shelf base. Using a thermal forcing depth near the ice shelf base rather than the deepest ocean layer above the continental shelf leads to a larger relative contribution of the Amundsen region to the total Antarctic sea level contribution, which is closer to observations.


For the Amundsen region, the basal melt method impacts the sea level contribution more than the SSP scenarios, whereas for the AIS the SSP scenarios dominate the sea level contribution over the basal melt method. However, differences related to the SSP scenarios and our methodological choices in the calibration and basal melt computation are small compared to the uncertainties associated with ESMs and RFs.

The calibration shows that the two main studies on which the IPCC AR6 Antarctic sea level contributions are based (ISMIP6 and LARMIP-2) use median basal melt sensitivities that are higher than the median Antarctic-wide calibrated values that we found, but lower than the median Amundsen calibrations. The Amundsen calibration performs better in simulating the sea level acceleration and the dominance of the Amundsen region over the historical period compared to Antarctic-wide calibration, and performs arguably better than the Antarctic-wide calibration when it comes to future projections (Sect. 4). The Amundsen calibration results in median estimates of 0.12 m for SSP1-2.6, 0.14 m for SSP2-4.5 and 0.17 m for SSP5-8.5, sitting in between the ISMIP6 and LARMIP-2 projections, as presented in IPCC AR6 (Fox-Kemper et al., 2021).


The basal melt calibration on Amundsen ice discharge leads to higher future sea level projections than projections using the median ISMIP6 AntMean and LARMIP-2 basal melt sensitivities. However, the LARMIP-2 and ISMIP6 projections as presented in IPCC AR6 are higher than our projections using their median basal melt sensitivity but otherwise applying the same procedure. This indicates that methodological differences between our study and LARMIP-2/ISMIP6 other than the basal melt sensitivity lead to higher projections (Fig. 1). If the Amundsen calibration would thus be combined with the methodological framework of ISMIP6 and LARMIP-2 as presented in IPCC AR6, our results suggest that the estimate of the Antarctic contribution to sea level would be higher than in the original studies.

## 6 Code and data availability

- Linear response functions from LARMIP-2 (Levermann et al., 2020): <https://github.com/ALevermann/Larmip2020/tree/master/RFunctions>
- Global ocean reanalyses: [https://resources.marine.copernicus.eu/product-detail/GLOBAL\\_REANALYSIS\\_PHY\\_001\\_026/INFORMATION](https://resources.marine.copernicus.eu/product-detail/GLOBAL_REANALYSIS_PHY_001_026/INFORMATION)
- Antarctic ice discharge (Rignot et al., 2019): [https://www.pnas.org/doi/suppl/10.1073/pnas.1812883116/suppl\\_file/pnas.1812883116.sd01.xlsx](https://www.pnas.org/doi/suppl/10.1073/pnas.1812883116/suppl_file/pnas.1812883116.sd01.xlsx)
- Other code available from reasonable request to the author.



**Figure A1.** Similar as Fig. 6, but for top 10% best-performing ESM-RF pairs.




**Figure A2.** Similar as Fig. 7, but for top 10% best-performing ESM-RF pairs.

*Author contributions.* EvdL, SD and DLB designed the study. DLB downloaded the CMIP6 data from the ESGF node and wrote the code to read it. EvdL performed the computations and prepared the manuscript with contributions from all co-authors.

655 *Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* We acknowledge the editor and two anonymous reviewers for their constructive comments which improved the quality of the paper. This publication was supported by the Knowledge Programme Sea Level Rise which received funding from the Dutch Ministry of Infrastructure and Water Management. This publication was supported by the project RECEIPT (REmote Climate Effects and their Impact on European sustainability, Policy and Trade) which received funding from the European Union's Horizon 2020 Research and Innovation  
 660 Programme under Grant agreement no. 820712. This publication was supported by PROTECT. This project has received funding from the European Union's Horizon 2020 research and innovation program under Grant agreement no. 869304, PROTECT contribution number 30.



**Figure A3.** Annual mean subsurface ocean temperature time series of the CMIP6 multi-model mean, model drift- and bias-adjusted, for temperatures centered around the mean depth of the ice shelf base (solid lines) and temperatures between 800-1000 m depth (dashed lines).

## References

- Bamber, J. L. and Aspinall, W. P.: An expert judgement assessment of future sea level rise from the ice sheets, *Nature Climate Change*, 3, 424–427, <https://doi.org/10.1038/nclimate1778>, 2013.
- 665 Bamber, J. L., Oppenheimer, M., Kopp, R. E., Aspinall, W. P., and Cooke, R. M.: Ice sheet contributions to future sea-level rise from structured expert judgment, *Proceedings of the National Academy of Sciences of the United States of America*, 166, 11 195–11 200, <https://doi.org/10.1073/pnas.1817205116>, 2019.
- Barthel, A., Agosta, C., Little, C. M., Hattermann, T., Jourdain, N. N., Goelzer, H., Nowicki, S., Seroussi, H., Straneo, F., and Bracegirdle, T. T.: CMIP5 model selection for ISMIP6 ice sheet model forcing: Greenland and Antarctica, *Cryosphere*, 14, 855–879, <https://doi.org/10.5194/tc-14-855-2020>, 2020.
- 670 Bronselaer, B., Winton, M., Griffies, S. M., Hurlin, W. J., Rodgers, K. B., Sergienko, O. V., Stouffer, R. J., and Russell, J. L.: Change in future climate due to Antarctic meltwater, *Nature*, 564, 53–58, <https://doi.org/10.1038/s41586-018-0712-z>, 2018.
- Clark, P. U., Shakun, J. D., Marcott, S. A., Mix, A. C., Eby, M., Kulp, S., Levermann, A., Milne, G. A., Pfister, P. L., Santer, B. D., Schrag, D. P., Solomon, S., Stocker, T. F., Strauss, B. H., Weaver, A. J., Winkelmann, R., Archer, D., Bard, E., Goldner, A., Lambeck, K.,   
 675 Pierrehumbert, R. T., and Plattner, G. K.: Consequences of twenty-first-century policy for multi-millennial climate and sea-level change, *Nature Climate Change*, 6, 360–369, <https://doi.org/10.1038/nclimate2923>, 2016.
- Dangendorf, S., Hay, C., Calafat, F. M., Marcos, M., Piecuch, C. G., Berk, K., and Jensen, J.: Persistent acceleration in global sea-level rise since the 1960s, *Nature Climate Change*, 9, 705–710, <https://doi.org/10.1038/s41558-019-0531-8>, <http://dx.doi.org/10.1038/s41558-019-0531-8http://www.nature.com/articles/s41558-019-0531-8>, 2019.
- 680 DeConto, R. M. and Pollard, D.: Contribution of Antarctica to past and future sea-level rise, *Nature*, 531, 591–597, <https://doi.org/10.1038/nature17145>, <http://dx.doi.org/10.1038/nature17145>, 2016.



- Dinniman, M. S., Asay-Davis, X. S., Galton-Fenzi, B. K., Holland, P. R., Jenkins, A., and Timmermann, R.: Modeling ice shelf/ocean interaction in Antarctica: A review, *Oceanography*, 29, 144–153, <https://doi.org/10.5670/oceanog.2016.106>, 2016.
- 685 Edwards, T. L., Nowicki, S., Marzeion, B., Hock, R., Goelzer, H., Seroussi, H., Jourdain, N. C., Slater, D. A., Turner, F. E., Smith, C. J., McKenna, C. M., Simon, E., Abe-Ouchi, A., Gregory, J. M., Larour, E., Lipscomb, W. H., Payne, A. J., Shepherd, A., Agosta, C., Alexander, P., Albrecht, T., Anderson, B., Asay-Davis, X., Aschwanden, A., Barthel, A., Bliss, A., Calov, R., Chambers, C., Champollion, N., Choi, Y., Cullather, R., Cuzzone, J., Dumas, C., Felikson, D., Fettweis, X., Fujita, K., Galton-Fenzi, B. K., Gladstone, R., Golledge, N. R., Greve, R., Hattermann, T., Hoffman, M. J., Humbert, A., Huss, M., Huybrechts, P., Immerzeel, W., Kleiner, T., Kraaijenbrink, P., Le clec'h, S., Lee, V., Leguy, G. R., Little, C. M., Lowry, D. P., Malles, J. H., Martin, D. F., Maussion, F., Morlighem, M., O'Neill, J. F., 690 Nias, I., Pattyn, F., Pelle, T., Price, S. F., Quiquet, A., Radić, V., Reese, R., Rounce, D. R., Rückamp, M., Sakai, A., Shafer, C., Schlegel, N. J., Shannon, S., Smith, R. S., Straneo, F., Sun, S., Tarasov, L., Trusel, L. D., Van Breedam, J., van de Wal, R., van den Broeke, M., Winkelmann, R., Zekollari, H., Zhao, C., Zhang, T., and Zwinger, T.: Projected land ice contributions to twenty-first-century sea level rise, *Nature*, 593, 74–82, <https://doi.org/10.1038/s41586-021-03302-y>, 2021.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model 695 Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Favier, L., Jourdain, N. C., Jenkins, A., Merino, N., Durand, G., Gagliardini, O., Gillet-Chaulet, F., and Mathiot, P.: Assessment of Sub-Shelf Melting Parameterisations Using the Ocean-Ice Sheet Coupled Model NEMO(v3.6)-Elmer/Ice(v8.3), *Geoscientific Model Development Discussions*, pp. 1–40, <https://doi.org/10.5194/gmd-2019-26>, 2019.
- 700 Fox-Kemper, B., H. T. Hewitt, C. Xiao, G. Aðalgeirsdóttir, S. S. Drijfhout, T. L. Edwards, N. R. Golledge, M. Hemer, R. E. Kopp, G. Krinner, A. Mix, D. Notz, S. Nowicki, I. S. Nurhati, L. Ruiz, Sallée, J.-B., Slangen, A. B. A., and Yu, Y.: Ocean, Cryosphere and Sea Level Change, Tech. rep., 2021.
- Golledge, N. R., Keller, E. D., Gomez, N., Naughten, K. A., Bernales, J., Trusel, L. D., and Edwards, T. L.: Global environmental consequences of twenty-first-century ice-sheet melt, *Nature*, 566, 65–72, <https://doi.org/10.1038/s41586-019-0889-9>, <http://dx.doi.org/10.1038/s41586-019-0889-9>, 2019. 705
- Haasnoot, M., Kwadijk, J., Van Alphen, J., Le Bars, D., Van Den Hurk, B., Diermanse, F., Van Der Spek, A., Oude Essink, G., Delsman, J., and Mens, M.: Adaptation to uncertain sea-level rise; how uncertainty in Antarctic mass-loss impacts the coastal adaptation strategy of the Netherlands, *Environmental Research Letters*, 15, <https://doi.org/10.1088/1748-9326/ab666c>, 2020.
- Hinkel, J., Lincke, D., Vafeidis, A. T., Perrette, M., Nicholls, R. J., Tol, R. S., Marzeion, B., Fettweis, X., Ionescu, C., and Levermann, A.: 710 Coastal flood damage and adaptation costs under 21st century sea-level rise, *Proceedings of the National Academy of Sciences of the United States of America*, 111, 3292–3297, <https://doi.org/10.1073/pnas.1222469111>, 2014.
- Holland, P. R., Jenkins, A., and Holland, D. M.: The response of Ice shelf basal melting to variations in ocean temperature, *Journal of Climate*, 21, 2558–2572, <https://doi.org/10.1175/2007JCLI1909.1>, 2008.
- Jenkins, A., Shoosmith, D., Dutrieux, P., Jacobs, S., Kim, T. W., Lee, S. H., Ha, H. K., and Stammerjohn, S.: West Antarctic Ice Sheet retreat 715 in the Amundsen Sea driven by decadal oceanic variability, *Nature Geoscience*, 11, 733–738, <https://doi.org/10.1038/s41561-018-0207-4>, <http://dx.doi.org/10.1038/s41561-018-0207-4>, 2018.
- Jourdain, N. C., Asay-Davis, X., Hattermann, T., Straneo, F., Seroussi, H., Little, C. M., and Nowicki, S.: A protocol for calculating basal melt rates in the ISMIP6 Antarctic ice sheet projections, *The Cryosphere*, 14, 3111–3134, <https://doi.org/10.5194/tc-14-3111-2020>, <https://tc.copernicus.org/articles/14/3111/2020/>, 2020.

- 720 Lambert, E., Le Bars, D., Goelzer, H., and van de Wal, R. S.: Correlations Between Sea-Level Components Are Driven by Regional Climate Change, *Earth's Future*, 9, 1–17, <https://doi.org/10.1029/2020EF001825>, 2021.
- Levermann, A., Winkelmann, R., Nowicki, S., Fastook, J. L., Frieler, K., Greve, R., Hellmer, H. H., Martin, M. A., Meinshausen, M., Mengel, M., Payne, A. J., Pollard, D., Sato, T., Timmermann, R., Wang, W. L., and Bindschadler, R. A.: Projecting Antarctic ice discharge using response functions from SeaRISE ice-sheet models, *Earth System Dynamics*, 5, 271–293, <https://doi.org/10.5194/esd-5-271-2014>, 2014.
- 725 Levermann, A., Winkelmann, R., Albrecht, T., Goelzer, H., Golledge, N. R., Greve, R., Huybrechts, P., Jordan, J., Leguy, G., Martin, D., Morlighem, M., Pattyn, F., Pollard, D., Quiquet, A., Rodehacke, C., Seroussi, H., Sutter, J., Zhang, T., Van Breedam, J., Calov, R., Deconto, R., Dumas, C., Garbe, J., Hilmar Gudmundsson, G., Hoffman, M. J., Humbert, A., Kleiner, T., Lipscomb, W. H., Meinshausen, M., Ng, E., Nowicki, S. M., Perego, M., Price, S. F., Saito, F., Schlegel, N. J., Sun, S., and Van De Wal, R. S.: Projecting Antarctica's contribution to future sea level rise from basal ice shelf melt using linear response functions of 16 ice sheet models (LARMIP-2), *Earth System Dynamics*, 11, 35–76, <https://doi.org/10.5194/esd-11-35-2020>, 2020.
- 730 Little, C. M. and Urban, N. M.: CMIP5 temperature biases and 21st century warming around the Antarctic coast, *Annals of Glaciology*, 57, 69–78, <https://doi.org/10.1017/aog.2016.25>, 2016.
- Liu, Y., Moore, J. C., Cheng, X., Gladstone, R. M., Bassis, J. N., Liu, H., Wen, J., and Hui, F.: Ocean-driven thinning enhances iceberg calving and retreat of Antarctic ice shelves, *Proceedings of the National Academy of Sciences of the United States of America*, 112, 3263–3268, <https://doi.org/10.1073/pnas.1415137112>, 2015.
- 735 Mathiot, P., Jenkins, A., Harris, C., and Madec, G.: Explicit representation and parametrised impacts of under ice shelf seas in the  $z_*$ -coordinate ocean model NEMO 3.6, *Geoscientific Model Development*, 10, 2849–2874, <https://doi.org/10.5194/gmd-10-2849-2017>, 2017.
- Nerem, R. S., Beckley, B. D., Fasullo, J. T., Hamlington, B. D., Masters, D., and Mitchum, G. T.: Climate-change-driven accelerated sea-level rise detected in the altimeter era, *Proceedings of the National Academy of Sciences of the United States of America*, 115, 2022–2025, <https://doi.org/10.1073/pnas.1717312115>, 2018.
- 740 Nowicki, S., Goelzer, H., Seroussi, H., Payne, A. J., Lipscomb, W. H., Abe-Ouchi, A., Agosta, C., Alexander, P., Asay-Davis, X. S., Barthel, A., Bracegirdle, T. J., Cullather, R., Felikson, D., Fettweis, X., Gregory, J. M., Hattermann, T., Jourdain, N. C., Kuipers Munneke, P., Larour, E., Little, C. M., Morlighem, M., Nias, I., Shepherd, A., Simon, E., Slater, D., Smith, R. S., Straneo, F., Trusel, L. D., Van Den Broeke, M. R., and Van De Wal, R.: Experimental protocol for sea level projections from ISMIP6 stand-alone ice sheet models, *Cryosphere*, 14, 2331–2368, <https://doi.org/10.5194/tc-14-2331-2020>, 2020.
- 745 Nowicki, S. M., Payne, A., Larour, E., Seroussi, H., Goelzer, H., Lipscomb, W., Gregory, J., Abe-Ouchi, A., and Shepherd, A.: Ice Sheet Model Intercomparison Project (ISMIP6) contribution to CMIP6, *Geoscientific Model Development*, 9, 4521–4545, <https://doi.org/10.5194/gmd-9-4521-2016>, 2016.
- Palmer, M. D., Gregory, J. M., Bagge, M., Calvert, D., Hagedoorn, J. M., Howard, T., Klemann, V., Lowe, J. A., Roberts, C. D., Slangen, A. B., and Spada, G.: Exploring the Drivers of Global and Local Sea-Level Change Over the 21st Century and Beyond, *Earth's Future*, 8, <https://doi.org/10.1029/2019EF001413>, 2020.
- 750 Payne, A. J., Nowicki, S., Abe-Ouchi, A., Agosta, C., Alexander, P., Albrecht, T., Asay-Davis, X., Aschwanden, A., Barthel, A., Bracegirdle, T. J., Calov, R., Chambers, C., Choi, Y., Cullather, R., Cuzzone, J., Dumas, C., Edwards, T. L., Felikson, D., Fettweis, X., Galton-Fenzi, B. K., Goelzer, H., Gladstone, R., Golledge, N. R., Gregory, J. M., Greve, R., Hattermann, T., Hoffman, M. J., Humbert, A., Huybrechts, P., Jourdain, N. C., Kleiner, T., Munneke, P. K., Larour, E., Le clec'h, S., Lee, V., Leguy, G., Lipscomb, W. H., Little, C. M., Lowry, D. P., Morlighem, M., Nias, I., Pattyn, F., Pelle, T., Price, S. F., Quiquet, A., Reese, R., Rückamp, M., Schlegel, N. J., Seroussi, H., Shepherd, A., Simon, E., Slater, D., Smith, R. S., Straneo, F., Sun, S., Tarasov, L., Trusel, L. D., Van Breedam, J., van de

- Wal, R., van den Broeke, M., Winkelmann, R., Zhao, C., Zhang, T., and Zwinger, T.: Future Sea Level Change Under Coupled Model Intercomparison Project Phase 5 and Phase 6 Scenarios From the Greenland and Antarctic Ice Sheets, *Geophysical Research Letters*, 48, 1–8, <https://doi.org/10.1029/2020GL091741>, 2021.
- 760 Pritchard, H. D., Ligtenberg, S. R., Fricker, H. A., Vaughan, D. G., Van Den Broeke, M. R., and Padman, L.: Antarctic ice-sheet loss driven by basal melting of ice shelves, *Nature*, 484, 502–505, <https://doi.org/10.1038/nature10968>, 2012.
- Rignot, E. and Jacobs, S. S.: Rapid bottom melting widespread near antarctic ice sheet grounding lines, *Science*, 296, 2020–2023, <https://doi.org/10.1126/science.1070942>, 2002.
- 765 Rignot, E., Mouginot, J., Scheuchl, B., Van Den Broeke, M., Van Wessem, M. J., and Morlighem, M.: Four decades of Antarctic ice sheet mass balance from 1979–2017, *Proceedings of the National Academy of Sciences of the United States of America*, 116, 1095–1103, <https://doi.org/10.1073/pnas.1812883116>, 2019.
- Sadai, S., Condrón, A., DeConto, R., and Pollard, D.: Future climate response to Antarctic Ice Sheet melt caused by anthropogenic warming, *Science Advances*, 6, 1–9, <https://doi.org/10.1126/sciadv.aaz1169>, 2020.
- 770 Seroussi, H., Nowicki, S., Payne, A. J., Goelzer, H., Lipscomb, W. H., Abe-Ouchi, A., Agosta, C., Albrecht, T., Asay-Davis, X., Barthel, A., Calov, R., Cullather, R., Dumas, C., Galton-Fenzi, B. K., Gladstone, R., Golledge, N. R., Gregory, J. M., Greve, R., Hattermann, T., Hoffman, M. J., Humbert, A., Huybrechts, P., Jourdain, N. C., Kleiner, T., Larour, E., Leguy, G. R., Lowry, D. P., Little, C. M., Morlighem, M., Pattyn, F., Pelle, T., Price, S. F., Quiquet, A., Reese, R., Schlegel, N. J., Shepherd, A., Simon, E., Smith, R. S., Straneo, F., Sun, S., Trusel, L. D., Breedam, J. V., Van De Wal, R. S., Winkelmann, R., Zhao, C., Zhang, T., and Zwinger, T.: ISMIP6 Antarctica: A multi-model ensemble of the Antarctic ice sheet evolution over the 21st century, *Cryosphere*, 14, 3033–3070, <https://doi.org/10.5194/tc-14-3033-2020>, 2020.
- 775 Shepherd, A., Ivins, E., Rignot, E., Smith, B., van den Broeke, M., Velicogna, I., Whitehouse, P., Briggs, K., Joughin, I., Krinner, G., Nowicki, S., Payne, T., Scambos, T., Schlegel, N., A. G., Agosta, C., Ahlstrøm, A., Babonis, G., Barletta, V., Blazquez, A., Bonin, J., Csatho, B., Cullather, R., Felikson, D., Fettweis, X., Forsberg, R., Gallee, H., Gardner, A., Gilbert, L., Groh, A., Gunter, B., Hanna, E., Harig, C., Helm, V., Horvath, A., Horwath, M., Khan, S., Kjeldsen, K. K., Konrad, H., Langen, P., Lecavalier, B., Loomis, B., Luthcke, S., McMillan, M., Melini, D., Mernild, S., Mohajerani, Y., Moore, P., Mouginot, J., Moyano, G., Muir, A., Nagler, T., Nield, G., Nilsson, J., Noel, B., Otosaka, I., Pattle, M. E., Peltier, W. R., Pie, N., Rietbroek, R., Rott, H., Louise Sandberg-Sørensen, Ingo Sasgen, H. S., Scheuchl, B., Schrama, E., Schröder, L., Seo, K.-W., Simonsen, S., Slater, T., Spada, G., Sutterley, T., Talpe, M., Tarasov, L., van de Berg, W. J., van der Wal, W., van Wessem, M., Vishwakarma, B. D., Wiese, D., and Wouters, B.: Mass balance of the Antarctic Ice Sheet from 1992 to 2017, *Nature*, 558, 219–222, <https://doi.org/10.1038/s41586-018-0179-y>, <http://www.nature.com/articles/s41586-018-0179-y>, 2018.
- 780 Thompson, A. F., Stewart, A. L., Spence, P., and Heywood, K. J.: The Antarctic Slope Current in a Changing Climate, *Reviews of Geophysics*, 56, 741–770, <https://doi.org/10.1029/2018RG000624>, 2018.
- van de Wal, R. S., Zhang, X., Minobe, S., Jevrejeva, S., Riva, R. E., Little, C., Richter, K., and Palmer, M. D.: Uncertainties in Long-Term Twenty-First Century Process-Based Coastal Sea-Level Projections, *Surveys in Geophysics*, 40, 1655–1671, <https://doi.org/10.1007/s10712-019-09575-3>, <https://doi.org/10.1007/s10712-019-09575-3>, 2019.
- 790 van den Broeke, M.: Strong surface melting preceded collapse of Antarctic Peninsula ice shelf, *Geophysical Research Letters*, 32, 1–4, <https://doi.org/10.1029/2005GL023247>, 2005.