Indication of high basal melting at <u>the</u> EastGRIP drill site on the Northeast Greenland Ice Stream

Ole Zeising^{1,2} and Angelika Humbert^{1,2}

¹Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, Bremerhaven, Germany ²University of Bremen, Department of Geosciences, Germany

Correspondence: Ole Zeising (ole.zeising@awi.de)

Abstract. The accelerated ice flow of ice streams that reach far into the interior of the ice sheet, is associate sheets, is associated with lubrication of the ice sheet base by basal melt water. However, the amount of basal melting under the large ice streams – such as the Northeast Greenland Ice Stream (NEGIS) – are is largely unknown. In-situ measurements of basal melt rates are important from various perspectives as they indicate the heat budget, the hydrological regime and the role relative importance of

- 5 sliding in glacier motion. The few previous estimates of basal melt rates in the NEGIS region were $0.1 \,\mathrm{m\,a^{-1}}$ and more, based on radiostratigraphy methods. These finding findings raised the question of the heat source, since even an increased geothermal heat flux could not deliver the necessary amount of heat. Here, we present basal melt rates at the recent deep drill site East-GRIP, located in the center of NEGIS. Within two subsequent years, we found basal melt rates of $(0.16 - 0.22) \pm 0.01 \,\mathrm{m\,a^{-1}}$, $0.19 \pm 0.04 \,\mathrm{m\,a^{-1}}$ that are based on analysis of repeated phase-sensitive radar measurements. In order to quantify the contri-
- 10 bution of processes that eause a heat flux into the icecontribute to melting, we carried out an assessment of the energy sources balance at the interface and found the subglacial water system to play a key role in facilitating such high melt rates.

1 Introduction

Ice sheet models are used to quantify the contribution of the Greenland Ice Sheet (GrIS) to future sea-level rise under different climatic scenarios. The In these simulations, the distinctive extent of Greenland's largest ice stream – the Northeast Greenland

15 Ice Stream (NEGIS, Fig. 1) – can only be represented well if an reproduced well if a higher-order approximation is considered for the momentum balance and initial states are based on inversion (Goelzer et al., 2018) or involve subglacial hydrological models (Smith-Johnsen et al., 2020a). Primarily, this is due to the model's ability inability to accurately represent lubrication and thus the subsequent sliding at the ice stream base that occurs.

The NEGIS is the only large ice stream in Greenland, extending from a distance of 100 km from the ice divide over a length of about 700 km towards the coast (Fahnestock et al., 1993, 2001b; Joughin et al., 2001). It drains about 12 % of Greenland's ice through three major outlet glaciers Nioghalvfjerdsbrae, Zachariæ Isstrøm and Storstrømmen Glacier (Rignot and Mouginot, 2012). Loss of the floating tongue of Zachariæ Isstrøm has already led to acceleration and ice flow acceleration and increased mass loss (Mouginot et al., 2015). Consequently, it is expected and projected that NEGIS will contribute significantly to sea level sea-level rise in the future (Khan et al., 2014), highlighting the importance to understand its lubrication the general ice

25 flow dynamics and its driving mechanisms.

One hypothesis for the genesis of NEGIS is locally increased basal melt rates melting at the onset area that enable basal sliding as basal melt water enables and enhances basal sliding (Fahnestock et al., 2001a; Christianson et al., 2014; Franke et al., 2021) and forms a subglacial hydrological system (Fahnestock et al., 2001a; Christianson et al., 2014; Franke et al., 2021). The coupling with basal sliding is facilitated via the water pressure, so that the sliding velocity rises with increasing water pressure (e.g., Beyer et al., 2018).

30 . However, little is known about the amount of subglacial water system below the 2000 - 3000 m below the up to $\sim 3300 \text{ m}$ thick ice sheet.

First estimates of basal melt rates by Fahnestock et al. (2001a) and later by Keisling et al. (2014) and MacGregor et al. (2016) are based on the interpretation of chronology in radiostratigraphy. Both-All three studies found melt rates of $0.1 \,\mathrm{m\,a^{-1}}$ and more – which is extremely large for inland ice. However, these estimates may be prone to limited validity given the

35 assumptions about the flow regime and constant accumulation rate. The cause for such intensive melt was attributed to a high geothermal heat flux which possibly originates from a remnant of the passing of the passage of Greenland over the Icelandic hot spot (Fahnestock et al., 2001a; Rogozhina et al., 2016; Martos et al., 2018; Alley et al., 2019).

In order to <u>conductdirectly observe</u>, among other things, <u>direct observations of</u> flow regimes and basal conditions of ice streams, an ice core is drilled in the course of the East Greenland Ice-Core Project (EastGRIP) near the onset of the NEGIS.

- 40 Here, surface velocities reach about $57 \,\mathrm{ma}^{-1}$ (Hvidberg et al., 2020) and the NEGIS widens (Fig. 1). Smith-Johnsen et al. (2020a) forced an ice model with a locally increased geothermal heat fluxes heat flux below the EastGRIP drill siteand. They found that a heat flux of $0.97 \,\mathrm{Wm}^{-2}$ (corresponding to a basal melting melt rate of $0.1 \,\mathrm{ma}^{-1}$ (Fahnestock et al., 2001a)) is necessary to reasonably reproduce the velocities of NEGIS. By utilizing a coupled subglacial hydrology and ice sheet model, Smith-Johnsen et al. (2020b) demonstrated the large impact of an uncertainty in geothermal heat flux on the flow of NEGIS
- 45 arising from the subglacial hydrological system, hence from basal melting and water pressure, as well as from friction.

However, measurements with an adequate accuracy are still required to narrow down the basal melt rates further. Here, we present the first estimates of basal melting melt rates from repeated in-situ phase-sensitive radar measurements from the EastGRIP drill site and consider the contribution of different heat sources at the ice base.

2 Data and Methods

50 2.1 Data

We deployed an autonomous phase-sensitive radio-echo sounder (ApRES; Brennan et al., 2014; Nicholls et al., 2015) within a near surface trench at the EastGRIP drill site (Fig. 1). The radar performed a measurement once a day during winter from 08/2017 - 04/2018 and 08/2018 - 05/2019. Within a single measurement, the ApRES transmits a sequence of 100 chirps with a duration of 1-s-s in which the frequency of the transmitted electromagnetic wave is increased from 200 to 400 MHz MHz.



Figure 1. Surface ice flow velocity map of the Greenland Ice Sheet (Joughin et al., 2018). The box in the overview map (upper left corner) marks the boundaries of the main figure showing northeast Greenland and the Northeast Greenland Ice Stream (NEGIS), which drains into the three major outlet glaciers, namely Nioghalvfjerdsbrae (79°N Glacier, 79NG), Zachariæ Isstrøm (ZI) and Storstrømmen Glacier (SG). The location of the EastGRIP drill site is denoted by the black triangle.

55 2.2 Basal melting The Ice thickness evolution

The method we use to derive a basal melt rate is based on the ice thickness evolution equation that is valid in both, the Eulerian and Lagrangian reference system

$$\frac{\partial H}{\partial t} = -\operatorname{div} \boldsymbol{Q} + a_s - a_b \tag{1}$$

with the ice thickness H, the time t, the volume flux Q, the surface mass balance a_s and the basal melt rate a_b (positive for melting) (e.g., Greve and Blatter, 2009). Equation (1) states that a temporal change in ice thickness is caused by a changing volume flux arising from deformation and accumulation or ablation at the ice surface and base. It is worth to note that a basal melt rate larger than the accumulation rate only leads to thinning of the glacier, if the volume flux cannot supply sufficient amount of ice to balance this out. The volume flux Q is defined as the vertically integrated horizontal velocities $v_x, v_y(x, y, z, t)$

65
$$\boldsymbol{Q} = \begin{pmatrix} Q_x \\ Q_y \end{pmatrix} = \begin{pmatrix} \int_0^H v_x \, \mathrm{d}z \\ \int_0^H v_y \, \mathrm{d}z \end{pmatrix}.$$
(2)

(Greve and Blatter, 2009) and represents the ice thickness change due to deformation and sliding, thus stretching or compression in the horizontal direction. This may, for example, be due to changes in basal velocities or ice creeping across a bedrock undulation. Using the continuity equation for incompressible materials, div v = 0, and Leibniz's integral rule we can rewrite div Q as

70 div
$$\mathbf{Q} = \frac{\partial}{\partial x} \int_{0}^{H} v_x dz + \frac{\partial}{\partial y} \int_{0}^{H} v_y dz = \int_{0}^{H} \frac{\partial v_x}{\partial x} + \frac{\partial v_y}{\partial y} dz = \int_{0}^{H} \frac{\partial v_z}{\partial z} dz = \int_{0}^{H} \dot{\varepsilon}_{zz} dz$$
 (3)

with $\dot{\varepsilon}_{zz}$ the vertical strain rate $\dot{\varepsilon}_{zz} = \partial v_z / \partial z$.

The recorded ApRES time series allows for a precise estimation of vertical displacements of internal layers and of changes in ice thickness ΔH from the vertical displacement of the basal reflector relative to the instrument and of internal layers from consecutive measurements. However, basal melt rates can not be derived directly, so other quantities that influence the

- 75 measured ice thickness *H* need to be derived first. Here and in the following, the term applying the ice thickness evolution equation (Eq. (1)) to the ApRES measurements requires some modifications. Since the ApRES is located within a trench below the surface, the 'measured ice thickness' *H* is defined as the range between the ApRES and the ice base, which was determined using the depth of the basal return. The total ice thickness – the range from the surface to the ice base – is about $\frac{7 \text{ to } 8 \text{ m}}{7 \text{ to}}$ 8 m thicker and includes the upper firn and snow layers.
- 80 A change in the measured ice thickness Thus, ΔH within the time interval Δt is caused is independent on the surface mass balance, $a_s = 0 \text{ ma}^{-1}$, but influenced by firn densification ΔH_f , by vertical strain $\Delta H_{\varepsilon_{zz}}$ and by basal melting ΔH_b (Nicholls et al., 2015):

$$\frac{\Delta H}{\Delta t} = \frac{\Delta H_f}{\Delta t} + \frac{\Delta H_{\varepsilon_{zz}}}{\Delta t} + \frac{\Delta H_b}{\Delta t}.$$

Thus, the basal melt rate a_b is defined as

$$a_b = \frac{\Delta H}{\Delta a}$$

In contrast to the total ice thickness, the change in the measured ice thickness is not affected by accumulation and only by the densification occurring below the radar instrument that significantly affects the vertical displacement in the upper ~ 100 m. As this is not considered in Eq. (1), we add the term $\Delta H_f / \Delta t$ to correct for the densification process below the ApRES.

The thickness change due to vertical strain is the integral over the entire vertical column of the vertical strain component ε_{zz}

90

$$\Delta H_{\varepsilon_{zz}} = \int_{0}^{H} \varepsilon_{zz}(z) \, \mathrm{d}z.$$

Vertical strain is defined as the Equation (3) states that the divergence of the volume flux in Eq. (1) can be expressed by the depth integrated vertical strain rate. However, we derive vertical displacements u_z from ApRES measurements instead of

vertical gradient of the vertical displacement component $u_z(z)$ velocities v_z . Thus, we can calculate strain $\varepsilon_{zz} = \partial u_z / \partial z$ for a 95 time period of Δt . Therefore, Eq. (3) needs to be reformulated as

$$\int_{0}^{H} \dot{\varepsilon}_{zz} \, \mathrm{d}z = \frac{1}{\Delta t} \int_{0}^{H} \varepsilon_{zz} \, \mathrm{d}z = \frac{\Delta H_{\varepsilon}}{\Delta t} \tag{4}$$

with the change in ice thickness due to vertical strain ΔH_{ε} . Finally, the modified ice thickness evolution equation can be written as

$$\varepsilon_{zz}(z)\frac{\Delta H}{\Delta t} = \frac{\partial u_z(z)}{\partial z}\frac{\Delta H_f}{\Delta t} + \frac{\Delta H_\varepsilon}{\Delta t} - a_b .$$
(5)

- 100 All three quantities ΔH , ΔH_f and $\frac{\Delta H_{\varepsilon_{zz}}}{\Delta H_{\varepsilon}}$, which are needed to derive a_b , are described by vertical displacements and hence by the measurement itself in a consistent manner.
 - ΔH is derived from the vertical displacement of the basal return. As densification processes are limited in depth, ΔH_f can be estimated based on the vertical displacement of the layers within the ice.
 - In order to estimate $\Delta H_{\varepsilon_{zz}}$, ε_{zz} and thus ΔH_{ε} , a regression analysis of the vertical displacements needs to be calculated
 - and the displacement extrapolated to the surface (the vertical displacement of layers above the noise-level depth limit *h* (depth at which the noise-level location of the ApRESmeasurement prevents an unambiguous estimation) needs to be analyzed and extrapolated) and to the ice base. The vertical gradient of the vertical displacement is the vertical strain.
 - ΔH_f is the intercept of the regression function at the surface.

Negative values of all three quantities contribute to the thinning of the ice thickness.

110 2.3 ApRES processing

105

We use an ApRES time series to achieve a reliable estimation of the annual mean basal melt rate. In order to derive amplitudeand phase-depth profiles for the ApRES signal processing, we followed the processing described by Brennan et al. (2014), Nicholls et al. (2015) and Stewart et al. (2019). For the conversion from travel time to depth, we used a vertical propagation velocity of $\frac{168,194 \text{kms}^{-1}}{168,914 \text{kms}^{-1}}$ according to relative permittivity of $\frac{\epsilon_r = 3.15}{\epsilon_r} = 3.15$ (Fujita et al., 2000). To

- 115 derive vertical displacements of layers within the ice as well as for the and of the basal return from the ApRES time series, we used a modification of the process described by Vaňková et al. (2020) modified the processing of Vaňková et al. (2020) to reduce measurement errors (details below). Both methods are based on estimated phase differences derived phase differences estimated from cross-correlation of individual depth segments. In contrast to Vaňková et al. (2020), we compare the first measurement with each repeated measurement instead of pairwise time-consecutively measurements to reduce measurement
- 120 errors. Here, the ApRES time series is used to achieve a reliable estimation of the annual mean basal melt rate.

FirstFirstly, we divided the depth profile into 6 m wide range segments with a 3 m overlap from a depth of 20 m below the antennas to 20 m above the ice base and a wider segment of 10 m (-9 to +1 m) around the basal return, characterized by

a strong increase in amplitude. Each In order to derive vertical displacements, each depth segment of the first measurement

 (t_1) was cross-correlated with the same segment of each repeated measurements measurement (t_i) . This is in contrast to

- 125 Vaňková et al. (2020), who derived displacements from pairwise time-consecutive measurements $(t_{i-1} t_i)$. The lag of the minimum mean phase difference obtained from the cross-correlation gives the cumulative displacement at the given depth. The range of expected lag was limited by the estimation to the previous measurement $(t_1 t_{i-1})$. This results in a time series of displacements for each segment individually. The vertical displacement of the basal segment is the change in the measured ice thickness ΔH . Segments whose time series contain outliers or whose shift deviates significantly from their neighboring
- 130 segments were discarded.

135

Next, we estimate the vertical strain ε_{zz}^{obs} as described in Equation (??) and quantify ΔH_f as well as $\Delta H_{\varepsilon_{zz}}$. ΔH_{ε} . To avoid influences of firn densification on the determination of ε_{zz}^{obs} , we excluded all segments above a depth of 250m (~9% of all segments). In addition, segments below the noise-level depth limit (depth at which the noise-level of the ApRES measurement prevents an unambiguous estimation) of $h \approx 1450$ m were excluded (~45% of all segments). Furthermore, outliers were filtered out (~7%). We found a linear fit $u_z(z)$ to match the curve of of

$$u_z(z) = \varepsilon_{zz}^{\text{obs}} \cdot z + \Delta H_f, \quad 250 \,\text{m} \le z \le h \tag{6}$$

that best matches the cumulative vertical displacements of the remaining ~ 400 segments within the ice (below a depth of 250 m to exclude layers affected by firm densification) best

$$u_z(z) = \varepsilon_{zz}^{\text{obs}} \cdot z + \Delta H_f, \quad 250 \,\text{m} \le z \le h$$

140 . The gradient of this fit is ε_{zz}^{obs} and the offset at the surface is ΔH_f . However, ε_{zz} for $z \ge h$ is unknown. Here, we used two scenarios in order estimate a range $\Delta H_{\varepsilon_{zz}}$ and provide a comparison to another often considered scenario (to estimate ΔH_{ε} (Fig. 2, Appendix Fig. A1). First, we assumed that ε_{zz} is constant with depth:

$$\varepsilon_{zz}^{\text{const}}(z) = \varepsilon_{zz}^{\text{obs}}, \quad 0 \le z \le H$$
(7)

As a second scenario, we used a vertical strain distribution (ε^{sim}_{zz}) obtained from an ice sheet model based on inverse surface
flow velocities (Rückamp et al., 2020). Here, ε^{sim}_{zz} increases with depth and reaches values of roughly twice ε^{obs}_{zz} at the base. For comparison, we also used a Dansgaard–Johnsen distribution of the vertical strain (Dansgaard and Johnsen, 1969), despite no-slip at the base is likely unrealistic (MacGregor et al., 2016). This assumes a constant ε^{di}_{zz} from the surface to the depth at which the uniform horizontal velocity changes to shear flow. From here, the vertical strain-rate decreases linearly to zero. As the depth of increased shear flow at EastGRIP is yet unknown, we assume that shear flow is occurring from the noise-level
depth limit (h) on, in order to underestimate ΔH_{εzz} and thus to overestimate the basal melt rate:

 $\varepsilon_{zz}^{\rm dj}(z) = \begin{cases} \varepsilon_{zz}^{\rm obs} & 0 \le z \le h \\ \varepsilon_{zz}^{\rm obs} \left(1 - \frac{h-z}{h-H}\right) & h < z \le H \end{cases}$

In order to be less dependent on a single measurement, we averaged the annual mean values of e_{zz}^{obs} , ΔH_f and ΔH from compute for each of the last 65 records (days (records; roughly 25% of the measurements) of a year an annual melt rate and compute from these 65 melt rate estimates a mean annual value by averaging. Finally, $\Delta H_{\varepsilon_{zz}} \Delta H_{\varepsilon}$ was derived from Equation Eq. (4) for the two vertical strain distributions ($\Delta H_{\epsilon_{zz}}^{\text{const}}, \Delta H_{\epsilon_{zz}}^{\text{sim}}, \Delta H_{\epsilon_{zz}}^{\text{const}}, \Delta H_{\epsilon_{zz}}^{\text{sim}}$), and the basal melt rate $a_{\rm b}$ from Equation Eq. (5). Given errors are based on the standard deviation of the estimates based on the considered 65 measurements and a 1%

uncertainty in the signal propagation speed in ice (Fujita et al., 2000). For visualization, we calculated the cumulative vertical displacement referenced to the ice bed.base (Fig. 2).

3 Results

155

- The analysis of the 2017/18 ApRES time series revealed a measured ice thickness (distance between radar and ice base) 160 of roughly 2668 m at the EastGRIP drill site with an annual mean change of $-0.471 \,\mathrm{m\,a^{-1}}$ (Tab. Table 1, Fig. 2). The firm densification – the intercept (see Fig. 2) of the linear fit at z = 0 m (the elevation of the ApRES) – occurring below the radar is 0.074 ma^{-1} . We derived a vertical strain of $-0.068 \times 10^{-3} \varepsilon_{zz} = -0.068 \times 10^{-3}$ from reliable estimates of vertical displacements feasible to a depth of 1450m. The dynamic thinning of the ice derived from the two scenarios ranges
- from $-0.181 \,\mathrm{m\,a^{-1}} \left(\Delta H_{\epsilon_{ex}}^{\mathrm{dj}} \Delta H_{\epsilon_{ex}}^{\mathrm{const}} \right)$ to $-0.194 \,\mathrm{m\,a^{-1}} \left(\Delta H_{\epsilon_{ex}}^{\mathrm{sim}} \right)$. The firn densification the intercept of the linear fit at the 165 surface (see Fig. 2) – occurring below the radar is $0.074 \,\mathrm{m\,a^{-1}}$. ΔH^{sim}_{coc}). This results in a basal melt rate between 0.203 and 1, Appendix Fig. A2). The annual mean change in measured ice thickness is 27 mm (or 6%) less-lower and the firn densification has increased by $15 \,\mathrm{mm}$ (or $20\,\%$). Thus, the larger compared to the values derived in 2017/18. The resulting basal melt rate varies from 0.160 to 0.173 ma⁻¹ and are 27% of 0.167 ± 0.018 ma⁻¹ is $\sim 20\%$ lower than the year before. Finally, we
- 170

derive an averaged melt rate over both years of $0.19 \pm 0.04 \,\mathrm{m\,a^{-1}}$.

Table 1. Results for measured ice thickness change (ΔH), firm densification (ΔH_f), vertical strain (ε_{zz}), dynamic ice thickness change obtain obtained from a constant vertical strain ($\Delta H_{\epsilon}^{const} \Delta H_{\epsilon}^{const}$) and based on a simulation ($\Delta H_{\epsilon}^{sim} \Delta H_{\epsilon}^{sim}$), find densification ($\Delta H_{\epsilon}^{const}$) and basal melting $(a_{\rm b})$ for both time series projected to 365 days.

Year	ΔH (m)	$\Delta H_{f}(\mathbf{m})$	$\varepsilon_{zz} \ (\times 10^{-3})$	$\Delta H_{\epsilon}^{\text{const}}(\mathbf{m}) \Delta H_{\epsilon}^{\text{sim}} \Delta H_{\epsilon}^{\text{const}}(\mathbf{m})$	$\frac{\Delta H_{f}}{\Delta H_{f}} \stackrel{\text{sim}}{\sim} (\text{m})$	
2017/18	-0.471 ± 0.008	-0.074 ± 0.001	-0.068 ± 0.001	-0.181 ± 0.001	-0.194 ± 0.001	-0.074 ± 0.00
2018/19	-0.444 ± 0.006	-0.089 ± 0.002	-0.068 ± 0.002	-0.182 ± 0.005	-0.195 ± 0.005	-0.089 ± 0.00

Discussion 4

We used estimated vertical displacements from the upper half of the ice column to estimate the dynamic thinning, since noise prevents an unambiguous estimation of the vertical strain for the lower half. To cover a range of variations in the dynamic thin-



Figure 2. Derived vertical displacements u_z of the surface $(\Delta H; blue dot)$ and selected internal layers referenced to the ice bed-base from 2017/18 ApRES time series. Derived displacements used for melt rate estimations are marked by red dots and of layers within the firn by gray dots. The estimated displacements between a depth of 250 m and h are used to calculate a linear fit (solid black line) which, the gradient of which is the vertical strain. Extrapolations to the bottom are shown by the dashed lines. The offset at the ice bed-base is caused by basal melting and the difference between the surface intercept of the linear fit and the derived surface displacement is the firn compaction.

- 175 ning, we used three_two different scenarios for vertical strain distribution. As the Dansgaard-Johnsen distribution describes a decrease in vertical velocity to zero at the ice base (representing a no-slip boundary condition), we do not consider this further. The vertical strain distribution obtained from the inverse ice sheet model is expected to give a more realistic representation. The The resulting dynamic thinning of the simulated vertical strain and the constant strain differs only slightly. However, we cannot exclude that larger strain values are reached at the base, which would lead to an overestimation of the basal melt rates. In case of
- 180 a non-existing melt rate, the dynamic thinning of the lower half of the ice column would be in average more than four times as large as the one of the upper half. However, a strong increase is not found in higher-order ice sheet simulations (Rückamp et al., 2020). A frequently used strain distribution (e.g., Fahnestock et al., 2001a; Keisling et al., 2014; MacGregor et al., 2016) that takes into account deviating strain within a shear zone is the Dansgaard–Johnsen distribution model (Dansgaard and Johnsen, 1969). As this model assumes a linearly decreasing strain in the shear zone that reaches zero at the ice base, the resulting basal melt

- 185 rate at EastGRIP would be even larger. However, the Dansgaard–Johnsen model represents a no-slip boundary condition at the ice base. As this is an unrealistic assumption in an ice stream, we did not consider the Dansgaard–Johnsen model further. The derived vertical strain is based on more than 300 vertical displacements estimated between the firn-ice transition and about 1450 m. In contrast, the estimation of the displacement of the basal return is based on the phase shift of only one segment around the basal return, slightly above the noise levelnoise-level. This makes the determination more prone to errors. Instead
- 190 of comparing the first measurement (t_1) with all repeated measurements (t_i) , the pairwise comparison of time-consecutively time-consecutive measurements $(t_{i-1} \text{ and } t_i)$, as it is shown by Vaňková et al. (2020), leads to a lower thinning rate of ΔH in 2017/18 than in 2018/19 $(-0.441 \pm 0.004 \text{ m a}^{-1} \text{ in } 2017/18, -0.467 \pm 0.009 \text{ m a}^{-1} \text{ in } 2018/19)$. Thus, the variability found is not necessarily a variability of the ice sheet system but can rather be influenced by the methodology.

A variation in the selected depth limit of densification, to exclude segments affected by densification, causes slight changes in vertical strain and thus in basal melt rate in the order of millimeters. However, we observed an increased densification rate within the considered 65 records. The increased densification can possibly be a result of increased load from the camp at the surface.

Our derived basal melt rates of 0.16 to 0.22 ma^{-1} are slightly rate of $0.19 \pm 0.04 \text{ ma}^{-1}$ is above previous estimates from airborne radar measurements. Fahnestock et al. (2001a) and MacGregor et al. (2016) found melt rates in the order of 0.1 ma^{-1}

200 in the vicinity of the EastGRIP drill sidesite, but larger melt rates of $> 0.15 \,\mathrm{m\,a^{-1}}$ further upstream in the onset region of NEGIS. Both studies used a constant vertical strain over depth where basal melting occurs. Smith-Johnsen et al. (2020a) found that basal melt rates of $0.1 \,\mathrm{m\,a^{-1}}$, derived from a heat flux of $0.97 \,\mathrm{W\,m^{-2}}$, are needed at the location of the EastGRIP drill site to reproduce the NEGIS in an ice sheet model.

205 4.1 Considerations of the energy balance at the ice base

210

In order to constrain the heat flux required to sustain the basal melt rates a_b derived in this study, we consider the energy balance at the ice base. As for any surface across which a physical quantity may not be continuous, a jump condition is formulated. In typical continuum mechanical formulation, the jump ($[[\psi]]$) of a quantity ψ is defined as $[[\psi]] = \psi^+ - \psi^-$, meaning the difference in the quantity ψ across the interface (Greve and Blatter, 2009). The jump condition of the energy at the ice base reads as

$$[[\boldsymbol{q}\cdot\boldsymbol{n}]] - [[\boldsymbol{v}\cdot\mathbf{t}\cdot\boldsymbol{n}]] + [[\boldsymbol{\rho}\left(\boldsymbol{u}+\frac{1}{2}\boldsymbol{v}^{2}\right)(\boldsymbol{v}-\boldsymbol{w})\cdot\boldsymbol{n}][\boldsymbol{\rho}^{\mathrm{i}}\left(\boldsymbol{u}+\frac{1}{2}\boldsymbol{v}^{2}\right)(\boldsymbol{v}-\boldsymbol{w})\cdot\boldsymbol{n}]] = [[\boldsymbol{q}\cdot\boldsymbol{n}]] - [[\boldsymbol{v}\cdot\mathbf{t}\cdot\boldsymbol{n}]] + \rho_{\underline{i}}^{\mathrm{i}}a_{\mathrm{b}}[[\boldsymbol{u}]] = 0$$
(8)

with the heat flux qq, the velocity v, the velocity of the singular surface ww, the normal vector n pointing outwards of the ice body, the Cauchy stress t, the ice density ρⁱ and the internal energy u (Greve and Blatter, 2009). The jump of the heat flux [[q · n]] becomes (q_{geo} + q_{sw}) · n - κ(T) grad T, with q_{geo} (q^{geo} + q^{sw}) · n - κ(T) grad T, with q^{geo} the geothermal heat flux
and q_{sw} q^{sw} the heat flux from subglacial water with a temperature above pressure melting point, T temperature and κ thermal

conductivity. For the jump in work of surface forces we find

$$\llbracket \boldsymbol{v} \cdot \mathbf{t} \cdot \boldsymbol{n} \rrbracket = \boldsymbol{v}^{\mathrm{sw}} \cdot \mathbf{t}^{\mathrm{sw}} \cdot \boldsymbol{n} - \boldsymbol{v}^{\mathrm{i}}_{\mathrm{b}} \cdot \mathbf{t}^{\mathrm{i}} \cdot \boldsymbol{n}$$
⁽⁹⁾

with t^{sw} the Cauchy stress of the subglacial water side of the singular surface, v_{b}^{i} the ice velocity and t^{i} the stress field of the ice at the base.

We split the traction vector of the subglacial water in a normal and tangential component, with the water pressure $p_w p^{sw}$ and the stress in the normal directionand use for the tangential component an empirical relation. Following the same approach as at an ice shelf base (Greve and Blatter, 2009), we employ an empirical relation

$$\mathbf{t}^{\mathrm{sw}} \cdot \mathbf{n} = -p_{\underline{\mathbf{w}}} \sum_{\sim}^{\mathrm{sw}} \mathbf{n} + C^{\mathrm{i/sw}} \rho_{\underline{\mathbf{w}}} \sum_{\sim}^{\mathrm{sw}} |\mathbf{v}^{\mathrm{sw}}|^2 \mathbf{e}_{\mathrm{t}}$$
(10)

with $e_t = v^{sw}/|v^{sw}|$ and $e_t \perp n$. The roughness drag coefficient at the underside of the ice is $C^{i/sw}$, similar as a Manning roughness is taken into account in subglacial conduits. So that the part of the subglacial water becomes

$$\boldsymbol{v}^{\mathrm{sw}} \cdot \mathbf{t}^{\mathrm{sw}} \cdot \boldsymbol{n} = -p_{\underline{\mathbf{w}}} \mathop{\sim}_{\sim}^{\mathrm{sw}} \boldsymbol{v}^{\mathrm{sw}} \cdot \boldsymbol{n} + \boldsymbol{v}^{\mathrm{sw}} \cdot C^{\mathrm{i}/\mathrm{sw}} \rho_{\underline{\mathbf{w}}} \mathop{\sim}_{\sim}^{\mathrm{sw}} |\boldsymbol{v}^{\mathrm{sw}}|^2 \boldsymbol{e}_{\mathrm{t}} = -p_{\underline{\mathbf{w}}} \mathop{\sim}_{\sim}^{\mathrm{sw}} \boldsymbol{v}_{\perp}^{\mathrm{sw}} + C^{\mathrm{i}/\mathrm{sw}} \rho_{\underline{\mathbf{w}}} \mathop{\sim}_{\sim}^{\mathrm{sw}} |\boldsymbol{v}_{\parallel}^{\mathrm{sw}}|^3$$
(11)

with v_{\perp}^{sw} , v_{\parallel}^{sw} the normal and tangential velocity of the subglacial water, respectively. This formulation is quite similar to the treatment of the jump condition at an ice shelf base. For the traction vector at the ice base, we follow the same procedure and find

$$230 \quad \mathbf{t}^{\mathrm{i}} \cdot \boldsymbol{n} = -N\boldsymbol{n} + \tau_{\mathrm{b}}\boldsymbol{e}_{\mathrm{t}} \tag{12}$$

with N the normal component and $\tau_{\rm b}$ the component in the tangential plane. For $v^{\rm i}_{\rm b} \cdot {f t}^{\rm i} \cdot {m n}$ we find

$$\boldsymbol{v}_{b}^{i} \cdot \boldsymbol{t}^{i} \cdot \boldsymbol{n} = -N \boldsymbol{v}_{b}^{i} \cdot \boldsymbol{n} + \tau_{b} \boldsymbol{v}_{b}^{i} \cdot \boldsymbol{e}_{t}.$$
(13)

With the jump of the internal energy [[u]] = L, we are can reformulate Eq. (8) to

$$q_{\underline{\underline{goo}}}^{\perp} + q_{\underline{\underline{sw}}}^{\perp} = \rho_{\underline{\underline{i}}}^{i} a_{\mathrm{b}} L + \kappa(T) \operatorname{grad} T - \underline{\underline{+}} p_{\underline{\underline{w}}}^{sw} v_{\perp}^{sw} + \underline{-} C^{i/sw} \rho_{\underline{\underline{w}}}^{sw} | v_{\parallel}^{sw} |^{3} + \underline{-} N v_{\mathrm{b}}^{i} \cdot n - \underline{\underline{+}} \tau_{\mathrm{b}} v_{\mathrm{b}}^{i} \cdot e_{\mathrm{t}}.$$

$$(14)$$

- 235 The tangential components $C^{i/sw}\rho_w |v_{\parallel}^{sw}|^3 C^{i/sw}\rho_w^{sw} |v_{\perp}^{sw}|^3$ and $\tau_b v_b^i \cdot e_t$ are frictional heating and are dominating the contribution of heat arising from work of surface forces. They need to be seen as two end members of the system: either the ice is only in contact with a vertically extensive thick subglacial hydrological system, then $C^{i/sw}\rho_w |v_{\parallel}^{sw}|^3 C^{i/sw}\rho_w^{sw}|v_{\parallel}^{sw}|^3$ is at place, or the subglacial hydrological system is permanently in contact with a lubricated base, then the second term $\tau_b v_b^i \cdot e_t$ is governing. The components are visualized in Fig. 3.
- Next, we aim at constraining the individual terms for which we use the following material parameters: $\rho_1 = 910 \text{ kgm}^{-3} \rho_1^i = 910 \text{ kgm}^{-3}$, the latent heat of fusion, $L = 335 \text{ kJ kg}^{-1}$, and the thermal conductivity for ice at $\frac{273.15 \text{ K} \kappa (273.15 \text{ K})}{2.07 \text{ Wm}^{-1} \text{ K}^{-1}}$ the pressure melting point of $270.81 \text{ K} \kappa (270.81 \text{ K}) = 2.10 \text{ Wm}^{-1} \text{ K}^{-1}$ (Greve and Blatter, 2009).

We consider three scenarios: (i) there is only temperate ice that is melting, (ii) heat is required to warm the ice to the pressure melting point and (iii) friction at the base is contributing significantly to melt ratebasal melting. (i) For temperate ice and no

245 heat arising from work of surface forces, we find a melt rates of at least $\frac{0.16 \text{ ma}^{-1} \ 0.19 \text{ ma}^{-1}}{0.19 \text{ ma}^{-1}}$ to correspond to a heat flux of $\frac{1.56 \text{ Wm}^{-2}}{1.84 \text{ Wm}^{-2}}$. (ii) Considering grad *T* to be between 10^{-1} and 10^{-3} ma^{-1} raises less than 10^{-1} Km^{-1} , this increases the required heat flux into the ice by 0.207 to 0.0207 Wm^{-2} from scenario (i) by up to 0.21 Wm^{-2} , as this additional heat is required to warm the ice to the pressure melting point.

(iii) Heat arising from work of the surface forces may, however, reduce the required heat flux into the ice to melt this amountof ice. To this end, we need to estimate the magnitude of the components of the stress tensors.

We assume that the normal stress component N is hydrostatic and bridging stresses to be negligible. With a mean density of ice of 910 kg m^{-3} we find $N = 23.8 \text{ MPa} p^i = 23.8 \text{ MPa}$. The normal velocity is of the order of the basal melt rate $v_b^{\perp} \approx 0.2 \text{ ma}^{-1} v_b^{\perp} \approx -0.2 \text{ ma}^{-1}$ by assuming the velocity of the interface (w) to be zero. The normal component of the ice side is then in the order of 0.15 W m^{-2} . For the tangential components of the ice side, we consider the shear stress at the base to be $\tau_b \approx 1$ to 100 kPa. This compares to basal shear stress found by Rückamp et al. (2020) of $\frac{50 \text{ kPa} 50 \text{ kPa}}{50 \text{ kPa}}$. To constrain the sliding velocity, we assume it to be maximum the surface velocity of 57 ma^{-1} and minimum half of the surface velocity. This leads to a tangential component on the ice side to be up to $\frac{150 \text{ mWm}^{-2}}{(\text{Appendix } 0.15 \text{ W m}^{-2})}$ (Fig. 4).

Next, we constrain the normal component of the subglacial water $p_w v_{\perp}^{sw} p_{\perp}^{sw} v_{\perp}^{sw}$. A water pressure of 10 to 23 MPa is consistent with subglacial hydrological modelling (Beyer et al., 2018; Smith-Johnsen et al., 2020a). Assuming the normal velocity

- to be at most as large as the basal melt rate, we find the range of this term to be between 51 to 116 mW m⁻² (Appendix 0.05 to $0.12 W m^{-2}$ (Fig. 4). The tangential component $C^{i/sw} \rho_w |v_u^{sw}|^3 - C^{i/sw} \rho_w |v_u^{sw}|^3$ needs an assumption on the roughness $C^{i/sw}$, for which we consider a range from the roughness of the ice shelf base to maximum a a maximum roughness ten times larger than this 10^{-3} to 10^{-2} . The motivation for this is that ice shelf roughness is governed by convection cells at the interface, whereas in the inland ice, the interaction with the bedrock may lead to a larger roughness. As nothing is known about the shape of the subglacial conduit, the range of velocity cannot be constrained well. We consider a similar speed like the speed similar
- to the one of the ocean $0.1 \,\mathrm{m\,s^{-1}}$, but as surface rivers easily reach $1.0 \,\mathrm{m\,s^{-1}}$, we take this as an upper limit (Fig. 4).

To summarize Thus, the contribution of friction to the energy available for basal melting may account for at least $\sim 200 \,\mathrm{mW \,m^{-2}} \sim 0.20 \,\mathrm{W}$ with the potential to be by far larger based on the assumptions we made.

- In particular To summarize, the jump in the tangential component (friction) has the potential to govern the heat budget, depending on flow speeds in subglacial water and roughness of the ice base, as can be seen in Fig. 4. However, assuming the geothermal heat flux to be in the order of $\mathcal{O}(q_{\text{geo}}) \approx 0.25 \,\mathrm{W \,m^{-2}}$ demonstrates $\mathcal{O}(q^{\text{geo}}) \approx 0.25 \,\mathrm{W \,m^{-2}}$ makes evident that the key player in facilitating such high melt rates is the subglacial water system, that may supply the ice base with an additional heat flux.
- 275 We have focused our consideration onto the interface between a subglacial water layer and the ice, as this drives the basal melt rate. However, observations of Christianson et al. (2014) highlight the existence of a wet till layer beneath the ice stream. Depending on the thickness of the water layer, the velocity and pressure of the water and the porosity of the till layer, complex interaction between the till and water may arise, too. Kutscher et al. (2019) present high resolution simulations of a comparable system that highlight the importance of studying this interface as well. To date, it is unclear which vertical extent of the water



Figure 3. Sketch of the energy balance at the ice base. Thermal components are colored in orange, mechanical components are in blue-purple color. For detail explanation see main text.



Figure 4. Magnitude of terms in the energy jump condition at the ice base. (a) Contribution of work of surface forces in normal direction from water pressure (left panellines) and tangential direction ice overburden pressure (middle and right panelsdots). The (b) Contribution of work of surface forces in tangential direction from friction on the subglacial waterand ice side are shown. (c) Contribution of work of surface forces in tangential direction from friction on the middle and right panel, respectively ice side.

280 layer is required to decouple the interaction of a water-till interface from the ice-water interface and thus the influence on the basal melt rate.

Implications of large basal melting is predominantly its effect on sliding via the effective normal pressure. Considering sliding to be the dominant part of ice stream motion, large basal melt rates are also plausible in the respect that the subglacial hydrological system needs to be sustained over time and hence creep closure of cavities or conduits needs to be balanced by melting.

285

Future measurements at EastGRIP after successful completion will shed more light onto the sliding speed and may also provide more information on the characteristics of the subglacial hydrological system. This will enable the community to put our melt rate estimates into further context.

5 Conclusions

- We estimated annual mean basal melt rates at the EastGRIP drill site from time series of high-precise high-precision phasesensitive radar measurements. We derived the change in the measured ice thickness, thinning from firn densification occurring below the instrument and the vertical strain in the upper 1450 m of the roughly 2668 m thick ice. Three Two different scenarios for vertical strain distribution were used to to quantify a plausible range of dynamic thinning. Thus, we derived a range of melt rates from 0.16 to 0.22 m a^{-1} an averaged melt rate of $0.19 \pm 0.04 \text{ m a}^{-1}$. We are aware that these melt rates require
- an extremely large amount of heat that we suggest to arise from the subglacial water system and the geothermal heat flux. However, these melt rates are based on measurements with a modern ice penetrating radar whose penetration depth is limited due to transmitting power. Thus, no assumptions on past accumulation rates or other uncertainties in age reconstruction are involved. Our major uncertainty is the vertical strain in the lower part of the ice stream. This could be overcome if a more powerful radar with a similar vertical resolution could be operated autonomously over several months which we want to accumulate herewith.

Code availability. Own developed MATLAB routines for ApRES processing are available from the corresponding author on request.

Data availability. Raw data of the ApRES measurements (https://doi.org/10.1594/PANGAEA.931018) are submitted to the World Data Center PANGAEA.



Figure A1. Observed vertical strain distribution (solid <u>black</u> line) and different extensions to the ice base (dashed lines) for three_two scenarios based on a constant (dashed black line), a Dansgaard–Johnsen (Dansgaard and Johnsen, 1969) and a simulated (Rückamp et al., 2020) (Rückamp et al., 2020, dashed orange line) distribution.



Figure A2. Derived vertical displacements u_z of the surface which equals ΔH (blue dot) and selected internal layers referenced to the ice bed base from 2018/19 ApRES time series. Derived displacements used for melt rate estimations are marked by red dots and of layers within the firm by gray dots. The estimated displacements between a depth of 250 m and h are used to calculate a linear fit (solid black line)which, the gradient of which is the vertical strain. Extrapolations to the bottom are shown by the dashed lines. The offset at the ice bed base is caused by basal melting and the difference between the surface intercept of the linear fit and the derived surface displacement is the firm compaction.

305 *Author contributions.* O.Z., A.H. conducted part of the field study. O.Z. processed the ApRES data and estimated the resulting basal melt rates. A.H. has designed the study and conducted the analysis of the energy jump condition. O.Z. and A.H. discussed the basal melt rate distribution and time series. O.Z. and A.H. wrote the manuscript jointly.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. Data has been acquired at the EGRIP EastGRIP camp that kindly hosted this activity as an associate project. EGRIP

- 310 EastGRIP is directed and organized by the Centre for Ice and Climate at the Niels Bohr Institute, University of Copenhagen. It is supported by funding agencies and institutions in Denmark (A. P. Møller Foundation, University of Copenhagen), USA (US National Science Foundation, Office of Polar Programs), Germany (Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research), Japan (National Institute of Polar Research and Arctic Challenge for Sustainability), Norway (University of Bergen and Bergen Research Trond Mohn Foundation), Switzerland (Swiss National Science Foundation), France (French Polar Institute Paul-Emile Victor, Institute for Geosciences and Environ-
- 315 mental research), <u>Canada (University of Manitoba</u>) and China (Chinese Academy of Sciences and Beijing Normal University). We like to thank Johanna Kerch, Sepp Kipfstuhl and Daniel Steinhage for their support in the deployment and the dismount of the measuring device. We are grateful for discussions with Thomas Kleiner and Martin Rückamp (AWI) and Ralf Greve (ITLS, Japan) on vertical strain-rates and our results.

References

335

- 320 Alley, R. B., Pollard, D., Parizek, B. R., Anandakrishnan, S., Pourpoint, M., Stevens, N. T., MacGregor, J. A., Christianson, K., Muto, A., and Holschuh, N.: Possible Role for Tectonics in the Evolving Stability of the Greenland Ice Sheet, Journal of Geophysical Research: Earth Surface, 124, 97–115, https://doi.org/10.1029/2018JF004714, 2019.
 - Beyer, S., Kleiner, T., Aizinger, V., Rückamp, M., and Humbert, A.: A confined–unconfined aquifer model for subglacial hydrology and its application to the Northeast Greenland Ice Stream, The Cryosphere, 12, 3931–3947, https://doi.org/10.5194/tc-12-3931-2018, 2018.
- 325 Brennan, P. V., Lok, L. B., Nicholls, K., and Corr, H.: Phase-sensitive FMCW radar system for high-precision Antarctic ice shelf profile monitoring, IET Radar, Sonar & Navigation, 8, 776–786, https://doi.org/10.1049/iet-rsn.2013.0053, 2014.
 - Christianson, K., Peters, L. E., Alley, R. B., Anandakrishnan, S., Jacobel, R. W., Riverman, K. L., Muto, A., and Keisling, B. A.: Dilatant till facilitates ice-stream flow in northeast Greenland, Earth and Planetary Science Letters, 401, 57–69, https://doi.org/10.1016/j.epsl.2014.05.060, 2014.
- 330 Dansgaard, W. and Johnsen, S.: A flow model and a time scale for the ice core from Camp Century, Greenland, Journal of Glaciology, 8, 215–223, https://doi.org/10.3189/S0022143000031208, 1969.
 - Fahnestock, M., Bindschadler, R., Kwok, R., and Jezek, K.: Greenland Ice Sheet Surface Properties and Ice Dynamics from ERS-1 SAR Imagery, Science, 262, 1530–1534, https://doi.org/10.1126/science.262.5139.1530, 1993.
 - Fahnestock, M. A., Abdalati, W., Joughin, I., Brozena, J., and Gogineni, P.: High geothermal heat flow, basal melt, and the origin of rapid ice flow in central Greenland, Science, 294, 2338–2342, https://doi.org/10.1126/science.1065370, 2001a.
- Fahnestock, M. A., Joughin, I., Scambos, T. A., Kwok, R., Krabill, W. B., and Gogineni, S.: Ice-stream-related patterns of ice flow in the interior of northeast Greenland, Journal of Geophysical Research: Atmospheres, 106, 34 035–34 045, https://doi.org/10.1029/2001JD900194, 2001b.
 - Franke, S., Jansen, D., Beyer, S., Neckel, N., Binder, T., Paden, J., and Eisen, O.: Complex Basal Conditions and Their Influence on
- 340 Ice Flow at the Onset of the Northeast Greenland Ice Stream, Journal of Geophysical Research: Earth Surface, 126, e2020JF005689, https://doi.org/10.1029/2020JF005689, 2021.
 - Fujita, S., Matsuoka, T., Ishida, T., Matsuoka, K., and Mae, S.: A summary of the complex dielectric permittivity of ice in the megahertz range and its applications for radar sounding of polar ice sheets, in: Physics of ice core records, pp. 185–212, Hokkaido University Press, 2000.
- 345 Goelzer, H., Nowicki, S., Edwards, T., Beckley, M., Abe-Ouchi, A., Aschwanden, A., Calov, R., Gagliardini, O., Gillet-Chaulet, F., Golledge, N. R., et al.: Design and results of the ice sheet model initialisation initMIP-Greenland: An ISMIP6 intercomparison, The Cryosphere, 12, 1433–1460, https://doi.org/10.5194/tc-12-1433-2018, 2018.
 - Greve, R. and Blatter, H.: Dynamics of Ice Sheets and Glaciers, Springer Science & Business Media, https://doi.org/10.1007/978-3-642-03415-2, 2009.
- 350 Hvidberg, C. S., Grinsted, A., Dahl-Jensen, D., Khan, S. A., Kusk, A., Andersen, J. K., Neckel, N., Solgaard, A., Karlsson, N. B., Kjær, H. A., et al.: Surface velocity of the Northeast Greenland Ice Stream (NEGIS): assessment of interior velocities derived from satellite data by GPS, The Cryosphere, 14, 3487–3502, https://doi.org/10.5194/tc-14-3487-2020, 2020.
 - Joughin, I., Fahnestock, M., MacAyeal, D., Bamber, J. L., and Gogineni, P.: Observation and analysis of ice flow in the largest Greenland ice stream, Journal of Geophysical Research: Atmospheres, 106, 34 021–34 034, https://doi.org/10.1029/2001JD900087, 2001.

- 355 Joughin, I., Smith, B. E., and Howat, I. M.: A complete map of Greenland ice velocity derived from satellite data collected over 20 years, Journal of Glaciology, 64, https://doi.org/10.1017/jog.2017.73, 2018.
 - Keisling, B. A., Christianson, K., Alley, R. B., Peters, L. E., Christian, J. E., Anandakrishnan, S., Riverman, K. L., Muto, A., and Jacobel, R. W.: Basal conditions and ice dynamics inferred from radar-derived internal stratigraphy of the northeast Greenland ice stream, Annals of Glaciology, 55, 127–137, https://doi.org/10.3189/2014AoG67A090, 2014.
- 360 Khan, S. A., Kjær, K. H., Bevis, M., Bamber, J. L., Wahr, J., Kjeldsen, K. K., Bjørk, A. A., Korsgaard, N. J., Stearns, L. A., Van Den Broeke, M. R., et al.: Sustained mass loss of the northeast Greenland ice sheet triggered by regional warming, Nature Climate Change, 4, 292–299, https://doi.org/10.1038/nclimate2161, 2014.
 - Kutscher, K., Geier, M., and Krafczyk, M.: Multiscale simulation of turbulent flow interacting with porous media based on a massively parallel implementation of the cumulant lattice Boltzmann method, Computers & Fluids, 193, 103733, https://doi.org/10.1016/j.compfluid.2018.02.009, 2019.
 - MacGregor, J. A., Fahnestock, M. A., Catania, G. A., Aschwanden, A., Clow, G. D., Colgan, W. T., Gogineni, S. P., Morlighem, M., Nowicki, S. M., Paden, J. D., et al.: A synthesis of the basal thermal state of the Greenland Ice Sheet, Journal of Geophysical Research: Earth Surface, 121, 1328–1350, https://doi.org/10.1002/2015JF003803, 2016.
 - Martos, Y. M., Jordan, T. A., Catalán, M., Jordan, T. M., Bamber, J. L., and Vaughan, D. G.: Geothermal heat flux reveals the Iceland hotspot
- track underneath Greenland, Geophysical research letters, 45, 8214–8222, https://doi.org/10.1029/2018GL078289, 2018.

365

375

380

- Mouginot, J., Rignot, E., Scheuchl, B., Fenty, I., Khazendar, A., Morlighem, M., Buzzi, A., and Paden, J.: Fast retreat of Zachariæ Isstrøm, northeast Greenland, Science, 350, 1357–1361, https://doi.org/10.1126/science.aac7111, 2015.
 - Nicholls, K. W., Corr, H. F., Stewart, C. L., Lok, L. B., Brennan, P. V., and Vaughan, D. G.: A ground-based radar for measuring vertical strain rates and time-varying basal melt rates in ice sheets and shelves, Journal of Glaciology, 61, 1079–1087, https://doi.org/10.3189/2015JoG15J073, 2015.
- Rignot, E. and Mouginot, J.: Ice flow in Greenland for the international polar year 2008–2009, Geophysical Research Letters, 39, https://doi.org/10.1029/2012GL051634, 2012.
 - Rogozhina, I., Petrunin, A. G., Vaughan, A. P., Steinberger, B., Johnson, J. V., Kaban, M. K., Calov, R., Rickers, F., Thomas, M., and Koulakov, I.: Melting at the base of the Greenland ice sheet explained by Iceland hotspot history, Nature Geoscience, 9, 366–369, https://doi.org/10.1038/ngeo2689, 2016.
- Rückamp, M., Goelzer, H., and Humbert, A.: Sensitivity of Greenland ice sheet projections to spatial resolution in higher-order simulations: the Alfred Wegener Institute (AWI) contribution to ISMIP6 Greenland using the Ice-sheet and Sea-level System Model (ISSM), The Cryosphere, 14, 3309–3327, https://doi.org/10.5194/tc-14-3309-2020, 2020.
- Smith-Johnsen, S., de Fleurian, B., Schlegel, N., Seroussi, H., and Nisancioglu, K.: Exceptionally high heat flux needed to sustain the
 Northeast Greenland Ice Stream, The Cryosphere, 14, 841–854, https://doi.org/10.5194/tc-14-841-2020, 2020a.
 - Smith-Johnsen, S., Schlegel, N.-J., de Fleurian, B., and Nisancioglu, K. H.: Sensitivity of the Northeast Greenland Ice Stream to Geothermal Heat, Journal of Geophysical Research: Earth Surface, 125, e2019JF005 252, https://doi.org/doi.org/10.1029/2019JF005252, 2020b.
 - Stewart, C. L., Christoffersen, P., Nicholls, K. W., Williams, M. J., and Dowdeswell, J. A.: Basal melting of Ross Ice Shelf from solar heat absorption in an ice-front polynya, Nature Geoscience, 12, 435–440, https://doi.org/10.1038/s41561-019-0356-0, 2019.
- 390 Vaňková, I., Nicholls, K. W., Corr, H. F., Makinson, K., and Brennan, P. V.: Observations of tidal melt and vertical strain at the Filchner-Ronne Ice Shelf, Antarctica, Journal of Geophysical Research: Earth Surface, 125, e2019JF005280, https://doi.org/10.1029/2019JF005280, 2020.