Quantifying the effects of background concentrations of crude oil pollution on sea ice albedo

Benjamin Heikki Redmond Roche¹ and Martin D. King¹

¹Department of Earth Sciences, Royal Holloway, University of London, Egham, Surrey, TW20 0EX, UK

5 Correspondence to: Professor Martin King (m.king@rhul.ac.uk)

Abstract. Sea ice albedo plays an important role in modulating the climate of the Earth and is affected by low background concentrations of oil droplets within the ice matrix that absorb solar radiation. In this study the albedo response of three different types of sea ice (melting, first-year, and multi-year sea ice) are calculated at increasing mass ratios $(0-1000 \text{ ng g}^{-1})$ of crude oil by using a coupled atmosphere-sea

- 10 ice radiative-transfer model (TUV-snow) over the optical wavelengths 400–700 nm. The different types of quasi-infinite thickness sea ice exhibit different albedo responses to oil pollution, with a 1000 ng g⁻¹ mass ratio of oil causing a decrease to 70.9% in multi-year sea ice, 47.2% in first-year sea ice, and 22.1% in melting sea ice relative to the unpolluted albedo at a wavelength of 400 nm. The thickness of the sea ice is also an important factor, with realistic thickness sea ices exhibiting similar results, albeit with a
- 15 weaker albedo response for multi-year sea ice to 75.3%, first-year sea ice to 66.3%, and melting sea ice to 35.9%. The type of oil also significantly affects the response of sea ice albedo, with a relatively opaque and heavy crude oil (*Romashkino* oil) causing a significantly larger decrease in sea ice albedo than a relatively transparent light crude oil (*Petrobaltic* oil). The size of the oil droplets polluting the oil also plays a minor role in the albedo response, with weathered submicron droplets (0.05–0.5 μm radius) of
- 20 Romashkino oil being the most absorbing across the optical wavelengths considered. Therefore, the work presented here demonstrates that low background concentrations of small submicron to micron-sized oil droplets have a significant effect on sea ice albedo. All three types of sea ice are affected, however first-year sea ice and particularly melting sea ice are very sensitive to oil pollution; thus, the Arctic may become more vulnerable to oil pollution as the ice becomes progressively thinner and younger in response to a
- 25 changing climate.

1 Introduction

Arctic sea ice has significantly declined from its 1981–2010 spring and summer averages, both in extent and thickness of sea ice cover (Fetterer et al., 2017). The summer sea ice minimum has decreased 13.1% per decade from the 1981–2010 average, with an average extent of 6.85 million km² in 1979–1992 compared to an average extent of 4.44 million km² from 2007–2020 (Thoman et al., 2020). Perennial sea ice cover decline is between 12.2% and 13.5% for first-year sea ice, and 15.6% and 17.5% for multiyear sea ice per decade, respectively (Comiso, 2012; Tschudi et al., 2019). Consequently, it is now very likely that an ice-free Arctic Ocean, a so called 'Blue Ocean Event', will be realised by the mid-century unless there is a rapid reduction in greenhouse gas emissions (Notz and Stroeve, 2018). In response to the 'blue'

- 35 Arctic Ocean, there has been a significant interest in developing northern shipping routes which can decrease journey lengths from Europe to Asia by up to 40% (Ho, 2010; Eguíluz et al., 2016; Kikkas and Romashkina, 2018). Additionally, 13% of total global undiscovered oil reserves are estimated to be in the Arctic Ocean and their exploitation is of great geopolitical importance for the Arctic (Bird et al., 2008; Krivorotov and Finger, 2019; Czarny, 2019). Therefore, well established concerns prevail for the 40 associated impacts that blowouts, pollution from offshore drilling, production, and transportation of oil
- may have in the Arctic (Koivurova and Vanderzwaag, 2007; WWF-Canada, 2011; Gulas et al., 2017). Sea ice albedo plays a key role in modulating the climate of the Earth. The high latitude, radiative balance, is primarily controlled by shortwave solar radiation which significantly effects both sea ice and snow cover in the region (e.g., Perovich et al., 1998; Flanner et al., 2007). Several different physical
- 45 properties affect sea ice albedo, the most significant of which are: density, grain size, grain shape, brine content, thickness, snow cover, and light-absorbing impurities (e.g., Perovich et al., 1998; Perovich, 2003; Marks and King, 2014; Hancke et al., 2018). The wavelength integrated and spectral albedos for different types of sea ice have previously been considered in literature (e.g., Grenfell and Maykut, 1977; Grenfell and Perovich, 1984; Pervoich et al., 1986; Buckley and Trodahl, 1987; Grenfell, 1991; Perovich, 1996;
- 50 Hanesiak et al., 2001); utilising these optical properties this study focuses on three types of sea ice: melting, first-year, and multi-year sea ice. The optical properties of these different types of sea ice are described in Marks and King (2014) and Lamare et al (2016). The albedo of sea ice is wavelength dependent with maximum albedo values occurring at 390 nm in pure ice, where absorption is at a minimum (Warren at al., 2006). The absorption of light absorbing impurities are also wavelength
- 55 dependent and affect where the maximum albedo occurs depending on their absorption spectra and the amount of the impurities that are contained within the ice. Different light absorbing impurities in sea ice such as volcanic dust, mineral dust, black carbon and soot have previously been examined for their effect on albedo (e.g., Warren and Wiscombe, 1980; Warren, 1984; Light et al., 1998; Doherty et al., 2010; Marks and King, 2013, 2014; Lamare et al., 2016; Marks et al., 2017) and were found to have significant
- 60 effects on sea ice albedo, even at very low concentrations. Indeed, Glaeser and Vance (1971) released oil on top of ice and found that oil absorbs 30% more heat from the sun than normal ice. However, aside from limited and extreme field studies by NORCOR (1975), and Gavrilo and Tarashkevich (1992), the effects that oil pollution has upon sea ice albedo have not previously been considered in literature, so this is explored here in a modelling study.
- 65 As oil is released into sea water it is influenced by several weathering processes: evaporation, dispersion, wave action, sedimentation, photo-oxidation and bioremineralisation (e.g. Daling et al., 1990; Resby and Wang, 2004; Dillipiane et al., 2021). The physicochemical properties of oil (e.g., water-in-oil emulsion viscosity, density, pour point) are constrained by oil composition, wave energy, and ice conditions, which largely determine the fate of oil in cold marine environments (Brandvik and Faksness,
- 70 2009; Brandvik et al., 2010; Singsaas et al., 2020). In low energy environments where nonbreaking waves occur, oil spilled at the surface will spread into a slick owing to gravity, viscosity, and surface tension. In high energy environments where plunging, spilling and breaking waves occur, oil droplets are entrained into the water and continually resurface, resulting in the droplet size decreasing (e.g. Delvigne and Sweeney, 1988; Wang et al., 2005; Z. Li et al., 2008a, 2008b; C. Li et al., 2017; Wilkinson et al., 2017).
- 75 In these high energy conditions, oil can spread several square kilometres in several hours and several hundred square kilometres within several days (Berenshtein et al., 2020). There have been extensive

modelling studies into both large-scale submarine blowouts (e.g., Johansen, 2003; Zheng et al., 2003; Lima Neto et al., 2008; Socolofsky et al., 2008; Fraga et al., 2016; Dissanayake et al., 2018) as well as surface oil slicks (e.g., Spaulding et al., 1992; Reed and Rye, 1995; Daling et al., 1997; Papadimitrakis et

- 80 al., 2005, 2011; Gamzaev, 2009), with the consensus being that knowledge of oil droplet size distribution is fundamental to accurately model ocean oil spills (Nissanka and Yapa, 2018). There have also been several lab (Hesketh et al., 1991; Masutani and Adams, 2000; Tang and Masutani, 2003; Brandvik et al., 2013, 2014; Wang et al., 2018) and field experiments (Johansen et al., 2003; Brandvik et al., 2010) which have generally focused on very high levels of pollution within a relatively short time frame from the
- 85 release of oil. However, away from the initial spill site the size of oil droplets dispersed by breaking waves is typically a log-normal distribution of smaller spherical particles (Otremba, 2007; Z. Li et al., 2011; Johansen et al., 2013; Haule et al., 2015; Haule and Freda, 2016).

Three major Joint Industry Programs into the behaviour and fate of oil in sea ice have taken place – the first was a four year SINTEF led project looking at 'Oil Spill Contingency for Arctic and Ice-covered

- 90 Waters' (Sørstrøm et al., 2010). The second was a three year International Association of Oil and Gas Producers led project looking at 'Arctic Oil Spill Response Technology' (Dickins, 2017). The third was a three year collaboration between SINTEF and the Research Council of Norway looking at the 'Fate, Behaviour and Response to Oil Drifting into Scattered Ice and Ice Edge in the Marginal Ice Zone' (Singsaas et al., 2020). Smaller studies have considered the hyperspectral features of oil-polluted sea ice
- 95 (Praks et al., 2004; Ivanov et al., 2005; Liu et al., 2016, 2018; Chao et al., 2017); it is possible to calculate the albedo from the bidirectional reflection distribution function (BDRF), however these are not comparable to this study as background mass ratios of oil are examined here. In another study, Gavrilo and Tarashkevich (1992) found that crude oil can migrate vertically through multiyear ice if it is pumped either into or below the ice. They also found that the presence of oil decreased both the albedo and
- 100 mechanical strength of the multiyear sea ice but did not measure the direct relative change in ice albedo in order to parameterise the effects. A similar study by NORCOR (1975) also concluded that the presence of oil released onto the ice may accelerate melting by 1 to 3 weeks but did not parameterise the effects either. Whilst it is now recognised that oil can be trapped in ice in a variety of ways (e.g. Dickins and Buist, 1981; Buist and Dickins, 1983; Buist et al., 1983; Drozdowski et al., 2011) what is missing from 105 the literature is an understanding of how oil exists at low concentrations over a large spatiotemporal range
- in cold marine environments and what their climatic significance may be. Studies from the Gulf of Thailand and South China Sea since the 1970's indicate that mass ratios

of oil are highly variable and dependent on location and season but can be up to 75 ng g⁻¹ (75 ppb), even far offshore (Law and Mahmood, 1986; Wongnapapan et al., 1999). In regions of intensive shipping and

- 110 marine transportation, particularly near to offshore oil fields, concentrations of oil ranging from several ppb to ppm are common (Haule and Freda, 2016), and oil concentrations of ship effluent discharge are only limited to a value of 15 ppm (i.e., 15,000 ng g⁻¹) (IMO, MARPOL Annex I). Mega oil spills are also capable of transporting significant quantities of oil vast distances in sea water. Following the Deepwater Horizon blowout oil mass ratios exceeding 100 ng g⁻¹ were transported via currents over 1000 km from
- 115 the spill site more than 60 days after the event occurred (Berenshtein et al., 2020). Therefore, it is important to consider a wide range of oil concentrations to replicate a variety of scenarios.

Oil droplets frequently weather more slowly and are more durable in cold environments than warm environments (Venkatesh et al., 1990; Singsaas et al., 2020). Smaller droplets (<70 µm diameter) are

affected by physical processes induced by atmospheric forcing (e.g., shear-induced turbulence, sea-

- 120 surface waves, Langmuir circulations, Ekman transport, and thermal convection) in the euphotic zone, resulting in the continuous mixing of the top of the water column and the formation of the ocean mixed layer (Edson et al., 2007; Sullivan and McWilliams, 2010; Özgökmen et al., 2012; D'Asaro, 2014; Xiao and Yang, 2020). The droplets remain in suspension for extended periods of time (i.e., weeks to months) in the presence of adequate mixing energy, with their buoyancy acting as a constant resistant force to the
- 125 mixing (Ryerson et al., 2012; Yang et al., 2014; Zhao et al., 2016; Uttieri et al., 2019; Xiao and Yang, 2020). The following papers (e.g., Lunel, 1993; Reed et al., 2009a, 2009b; Bandara and Yapa, 2011; Brandvik et al., 2013, 2021; Johansen et al., 2013; Zhao et al., 2014a, 2014b, 2016, 2017a, 2017b; North et al., 2015; Zeinstra-Helfrich et al., 2015a, 2015b, 2016; Nissanka and Yapa, 2016, 2017, 2018; Z. Li et al., 2017a, 2017b; Wang et al., 2018) demonstrate the typical sizes of oil droplets in sea water (3–500 μm,
- 130 locally up to 12,000 μm) and also focus on relatively large concentrations of crude oil representing the immediate area around a spill site, whereas the study presented here focuses on relatively low background concentrations of weathered crude oil over a large area. The focus of these studies is on droplet sizes between microns to millimetres which demonstrate a logarithmic dependence between number density and droplet size that increases as droplet size decreases (Lunel, 1993; Brandvik et al., 2013, 2021;
- 135 Johansen et al., 2013; Zhao et al., 2017b; Nissanka and Yapa, 2017; C. Li et al., 2017). These larger micron to millimetre sized droplets are characteristic of oil released in the subsurface, such as during a blowout or similar specific acute event of small geographical extent, which have not undergone significant weathering or transport (e.g., Johansen et al., 2013; Brandvik et al., 2013). The number density of smaller sized droplets is potentially not reported owing to limitations of measuring equipment or techniques, and
- 140 as pointed out by Otremba (2007), they are measuring the larger radius tail of a log-normal distribution of a number density against droplet radius. It is likely that the most numerous droplets affecting wide areas of polar sea ice are these smaller sized weathered oil droplets. The smallest droplets may not be considered in these studies owing to four factors: (i) the proximity and timing of measurements to droplet release; (ii) equipment limitation in droplet size-range detection (e.g., LISST-100X below 2.5 μm)
- 145 coupled with the effect of background noise at the smallest scale (e.g., Brandvik et al., 2013); (iii) droplets smaller than 0.2 µm being considered as dissolved (e.g., North et al., 2015); (iv) the difficulty and high uncertainty of direct measurements in situ (e.g., Gonçalves et al., 2016). In addition, Nissanka and Yapa (2017) suggest that many of the equilibrium models utilised in the literature (e.g., Delvigne and Sweeney, 1988; Spaulding et al., 1992; Delvigne and Hulsen, 1994; Reed and Rye, 1995; Tkalich and Chan, 2002;
- 150 Reed et al., 2009a, 2009b; Johansen et al., 2013, 2015) omit to consider the temporal modification of oil droplet size distributions owing to droplet break up and coalescence.

The study presented here utilises a coupled atmospheric and sea ice radiative-transfer model to quantify how oil pollution may affect the albedo of three different types of sea ice (melting sea ice, first-year sea ice, and multi-year sea ice). The principle aim of this study is evaluating the impacts increasing

155 amounts of two crude oils that band an envelope of crude oils optically have on sea ice albedo. Previous studies have demonstrated that the size and absorption cross section of light absorbing impurities and the type of sea ice affect how sensitive a sea ice is to increasing mass densities of pollutant (e.g., Marks and King, 2013, 2014; Lamare et al., 2016; Marks et al., 2017). Therefore, the effect that oil has on the three different types of sea ice and at different oil droplet sizes are also evaluated.

160 2 Methods

The methodology is separated into three sections: a description of the radiative-transfer model, a description of the optical properties of the pollutants used in the model, and the different model runs that were examined in this study.

2.1 Radiative-transfer model

- 165 The Tropospheric Ultraviolet and Visible Radiation model was originally written by Madronich and Flocke (1998) and later adapted into a coupled atmosphere-snow/sea ice model (TUV-snow) by Lee-Taylor and Madronich (2002). The model is capable of calculating radiative-transfer parameters from the top of the atmosphere into the sea ice and snow layers found at the ocean-atmosphere interface, and has been used in several studies (e.g., King et al., 2005; Marks and King, 2013, 2014; Lamare et al., 2016;
- 170 Marks et al., 2017). The model utilises an eight-stream discrete-ordinates 'DISORT' algorithm (Stamnes et al., 1988) with a pseudo-spherical correction, and is comprehensively described by Lee-Taylor and Madronich (2002). In this study the albedo response of three different types of sea ice (multi-year sea ice, first-year sea ice and, melting sea ice) to increasing amounts of two different crude oils are calculated using the TUV-snow model. The sea ice layers are assumed to be horizontally homogeneous, weakly
- absorbing and highly scattering. The optical properties of the sea ice layers are described by an asymmetry factor, g, a wavelength independent scattering cross-section, σ_{scatt} , a wavelength dependent absorption cross-section, σ_{abs} , and the sea ice density (Lee-Taylor and Madronich, 2002; France et al., 2011; Reay et al., 2012; Marks and king, 2013, 2014; Lamare et al., 2016; Marks et al., 2017). The optical and physical parameters of the three sea ices emanate from several field studies (Grenfell and Maykut, 1977; Perovich,
- 180 1990, 1996; Timco and Frederking, 1996; Gerland et al., 1999; Simpson et al., 2002; Fisher et al., 2005; King et al., 2005; France et al., 2011; Marks and King, 2013) and are summarised in Table 1. The density of sea ice has been fixed in this study to be comparable to previous work (Lamare et al., 2016), however sea ice density can range between extremes of 700–950 kg m⁻³ (Grenfell and Maykut, 1977; Perovich, 1990, 1996; Timco and Frederking, 1996; Gerland et al., 1999).
- 185

Table 1. TUV-snow model sea ice input parameters derived from literature (Grenfell and Maykut, 1977; Perovich, 1990, 1996; Timco and Frederking,l 1996; Gerland et al, 1999) and previously used in other studies (e.g., King et al, 2005; France, 2008; France et al, 2011; Marks and King, 2013, 2014; Lamare et al, 2016).

190

Type of sea ice	Density	Scattering cross	Asymmetry	Quasi-infinite ice	Realistic ice
	(kg m ⁻³)	section (m ² kg ⁻¹)	parameter (g)	thickness (cm)	thickness (cm)
Melting	800	0.03	0.98	1500	250
First-year	800	0.15	0.98	1500	80
Multi-year	800	0.75	0.98	1500	250

The atmosphere and sea ice in the model are separated into 201 layers comprising a total thickness of 90 km and are shown in Table 2. Marks (2017) found it important to have a large number

of layers at the interface so 201 layers offers a good compromise between computational time and

- 195 precision. The optical effects are considered both in vacuo, by using a quasi-infinite thickness sea ice, and at realistic thicknesses of sea ice. The sea ice is set to be optically thick at a quasi-infinite thickness of 15 m (Lamare et al., 2016) therefore making albedo changes at the ice-atmosphere interface independent of the layer under the sea ice and allowing for a fair comparison of the effects explored here. The realistic thickness sea ices are also considered to allow for a more realistic effect on the
- 200 different sea ice albedos to be observed. The realistic multi-year and melting sea ice are assigned a thickness of 2.5 m and the first-year sea ice is assigned a thickness of 0.8 m (Bourke and Garrett, 1987; Laxon et al., 2003; Weeks, 2010).

Table 2. TUV-snow model sea ice input parameters derived from literature and detailed in Marks and205King (2014).

Ice thickness (m)	Number of layers	Layer structure
0.8	141	1 cm increments 0–0.1 m; 10 cm increments 0.1–0.2 m; 2 cm increments 0.2–0.6 m; 1 cm increments 0.6–0.7 m; 1 mm increments 0.7–0.8 m
2.5	141	1 cm increments 0–0.1 m; 10 cm increments 0.1–2.2 m; 2 cm increments 2.2–2.4 m; 1 mm increments 2.4–2.5 m
15	141	1 cm increments 0–0.1 m; 10 cm increments 0.1–1 m 1 m increments 1–14 m; 10 cm increments 14–14.9 m; 1 mm increments 14.9–15 m
Atmosphere (90 km thick)	60	(10 cm increments 0.8–1 m; 4.5 m increments 1–10 m; 10 m increments 10–100) (1.875 m increments 2.5–10 m; 10 m increments 10–100 m;) 1 m increments 15–20 m; 10 m increments 20–100 m; 100 m increments 100–1000 m; 1 km increments 1–10 km; 2 km increments 10–42 km; 4 km increments 42–90 km

The 'flat plate' irradiance reflectance of the surface layer of the sea ice is calculated in the model and is referred to as albedo in the work described here, represented by Eq. (1):

210

$A = \frac{E_u}{E_u}$	(1)
$E_{\rm d}$	(-)

Where E_u is the upwelling irradiance and E_d is the downwelling irradiance. The oil droplets in this study are considered to have a log-normal size distribution that is representative of weathered oil according to experimental measurements by Otremba (2007). The log-normal distribution is centred on a droplet radius size of 0.05 µm and spans a range of 0.05–5 µm, in line with Haule and Freda (2016). Furthermore, a wide range of oil mass ratios are considered (0–1000 ng g⁻¹) so that the data is applicable for a variety of different scenarios. The wavelength independent albedo of the layer under the sea ice is fixed to 0.1, representative of a dark ocean layer (Payne, 1972). The sea ice is illuminated isotropically
to allow ready comparison with previous work (e.g., Marks and King, 2013, 14; Lamare et al., 2016; Marks et al., 2017) and remove a solar zenith angle dependency. As shown elsewhere (e.g., Marks and King, 2013), the isotropic illumination allows any solar zenith angle dependence to be disregarded for simplicity. The atmosphere is assumed to be aerosol free with a typical ozone column of 300 Dobson units. The distance between the Earth and the Sun is set to 1 astronomical unit. The model produced a
wavelength dependent surface albedo at a 1 nm interval over a spectrum covering 400–700 nm, corresponding to the wavelength range of the available oil optical properties and typical optical range

- measured in the field. The wavelength range 400–700 nm was considered in this study as the absorption coefficient of oil decreases with increasing wavelength whereas the absorption coefficient of ice increases with wavelength, thus at wavelengths longer than 700 nm the absorption of solar photons is dominated
 by ice.
- 230 by ice.

2.2 Optical properties of crude oil

The model utilises the absorption spectra for ice as defined in Warren and Brandt (2008). The under-ice surface, snow cover, average Arctic black carbon (5.5–19.2 ng g⁻¹), and average aerosol mass ratios ($\leq 193.2 \text{ ng g}^{-1}$) are not considered for the quasi-infinite sea ice in order to independently assess the 235 effect that oil pollution has on the albedo of the different types of sea ice (Zdanowicz et al., 1998; Grenfell et al., 2002 Jiao et al., 2014). However, a very low background mass ratio of black carbon ($\ln g g^{-1}$) is included, in line with previous studies (e.g., Marks and King, 2013; 2014), and representative of the most pristine snowpack away from the stations at both Dome C, Antarctica and Summit, Greenland (Zatko et al., 2013). For the realistic thickness sea ices, a typically low Arctic background black carbon mass ratio of 5.5 ng g⁻¹ is selected (Grenfell et al., 2002). However, snow is not considered to allow for the effect of 240 oil pollution on sea ice albedo to remain independent. The droplets are evenly distributed throughout the ice to be representative of seasonal oil movement in the ice (NORCOR, 1975; Martin, 1979; Drozdowski et al., 2011). The TUV-snow model is capable of considering layers of pollutant in the ice (e.g., Marks and King, 2013; Lamare et al., 2016; Marks et al., 2017); however, until there is a greater knowledge of 245 the thickness, type and location of these layers for background mass ratios of oil, it is beyond the scope of the present work. The mass ratio of oil in the ice is small and therefore the light scattering properties of the ice matrix dominate, thus only the absorption of light by the oil droplets is considered. The total absorption in sea ice is represented by Eq. (2):

250
$$\sigma_{\text{total}} = \sigma_{ice}(\lambda) + \chi_{oil}\sigma_{oil}(\lambda)$$

(2)

Where $\sigma_{ice}(\lambda)$ is the wavelength-dependent absorption cross section of sea ice, χ_{oil} is the mass ratio of crude oil in the sea ice, and σ_{oil} is the wavelength dependent mas absorption cross-section by crude oil.

255

The optical properties of several types of crude oil, lubricating oil and petroleum are explored by Otremba (2000), who parameterised the real and imaginary parts of their refractive index. Otremba (2000) established that, for the oils capable of forming a surface film, *Romashkino* crude oil has the largest imaginary refractive index and absorption coefficient values whereas *Petrobaltic* crude oil has the

- smallest values, with other oils bracketed by these parameters. *Petrobaltic* is described as a relatively transparent light crude oil, whereas *Romashkino* is described as a heavy, relatively opaque crude oil (Otremba, 2005; 2007; De Carolis et al., 2014). Whilst both crude oils have a variety of uses, including as marine engine fuels, *Romashkino* can be considered a typical marine engine Heavy Fuel Oil (De Carolis et al., 2014; IMO, MARPOL Annex 1). In the model *Petrobaltic* and *Romashkino* are assigned characteristic light and heavy crude oil densities of 0.77 g cm⁻³ and 0.94 g cm⁻³, respectively (Hollebone,
- 265 2015). Both mechanically and chemically dispersed small droplets tend to be spherical in nature, so they are considered to be spherical in the model (Johansen et al., 2013). A log-normal distribution of number density versus radius with a peak radius of 0.05 μm is utilised, corresponding to the weathered oil experimentally measured in Otremba (2007) and used in other studies (e.g., Haule et al., 2015; Haule and Freda, 2016). The log-normal distribution is represented by Eq. (3):
- 270

$$\frac{dN}{dln(r)} = \frac{No}{\sqrt{2}\pi} \frac{1}{ln(s)} \exp\left(\frac{-(ln(r) - ln(r_m))^2}{2ln^2(s)}\right)$$
(3)

Where *r* is the droplet radius, r_m is the median radius of the size distribution, *s* is the geometric standard deviation, N_0 is the total particle density, and *N* is the particle density (Seinfeld and Pandis, 2006). To quantify the size effect a range of droplet sizes spanning 0.05–5 µm (Haule and Freda, 2016) are considered, and the geometric standard deviation of the mean is taken from Otremba (2007). Owing to the size of the droplets and the range of the spectrum considered (400–700 nm), Mie calculations (Bohren and Huffman, 1983) are required to record the efficiency of light absorption for particles of different sizes within the Mie regime, in a sea ice or brine matrix. There are two similar modern determinations of the refractive index of ice (Warren and Brandt, 2008; Picard et al., 2016), to stay consistent with previous work (e.g., Marks and King, 2013; 2014) the values of Warren and Brandt (2008) are used in this paper. The real refractive index of the brine was 1.3402 at a wavelength of 600 nm (Quan

and Fry, 1995) and set to a temperature of 0° C and a salinity of 35‰, the smallest temperature and largest salinity values available.

Mie calculations of spherical oil droplets are weighted by log-normal size distributions using a modified version of the 'BHMIE' computational code (Bohren and Huffman, 1983). The calculations utilise the real (0° C) and imaginary refractive index values of *Romashkino* and *Petrobaltic* oil from Otremba (2000) and are applied in the model in agreement with literature (Otremba, 2007; Haule and Freda, 2016). The Mie calculations compute the mass absorption cross sections of the oil from the absorption efficiencies. The mass absorption coefficients of the oils are used in the TUV-snow computational code (Lee-Taylor and Madronich, 2002) to calculate the effect that increasing mass ratios

of Romashkino and Petrobaltic oil have on the albedo of the different types of sea ice.

2.3 Calculation of albedo on sea ice with different loadings of crude oil

Albedos of quasi-infinite first-year sea ice doped with different mass ratios of both *Petrobaltic* and *Romashkino* from 0–1000 ng g⁻¹ (0, 5, 10, 25, 100, 150, 200, 250, 300, 350, 400, 450, 500, and 1000 ng g⁻¹) are calculated as a function of wavelength. The study presented here is then repeated for quasiinfinite multi-year, first-year, and melting sea ice. The effect of oil droplet size (in the range 0.05–5.0

um) are then considered for a single wavelength (400 nm) for the oil with largest absorption cross section (Romashkino) and for quasi-infinite sea ice with the largest penetration depth to light (melting sea ice)

300 (Marks and King, 2014). These conditions produce the largest variation in sea ice albedo with changing droplet size. The mass absorption coefficient of droplets (in the range 0.005-50 µm) are also considered for the wavelengths 400-700 nm. The albedo of quasi-infinite melting sea ice polluted with increasing amounts of black carbon (1-100 ng g⁻¹) and Romashkino oil are also considered to investigate how sea ice albedo responds to increasing mass ratios of oil as a function of background pollution of sea ice. The albedos of realistic thickness melting, first-year, and multi-year sea ice are also calculated at increasing

305

mass ratios of both *Petrobaltic* and *Romashkino* from 0 to 1000ng g⁻¹.

3 Results

The results are separated into five sections: how the type of oil and increasing mass ratios of oil effect quasi-infinite sea ice albedo, the sensitivity of each type of quasi-infinite sea ice to increasing oil, the effect of droplet size distribution on quasi-infinite sea ice albedo, the impact that increasing 310 background mass ratios of black carbon have on the further reduction of quasi-infinite sea ice albedo by oil, and the effect oil has on each type of realistic thickness sea ice.

3.1 The effect of increasing mass ratios of different types of oil on the albedo of quasi-infinite sea ice

- 315 Figure 1 illustrates the surface albedo of quasi-infinite first-year sea ice between 400–700 nm as a function of *Petrobaltic* and *Romashkino* oil mass ratios that are evenly distributed throughout the ice and increase from 0–1000 ng g⁻¹. As the mass ratios of both oils increases sea ice albedo decreases. The Mie calculation demonstrates the oil droplets are most absorbing at a wavelength of 400 nm where ice is the least absorbing (Warren and Brandt, 2008). The albedo at 400 nm with no oil pollutant is 0.87, with an addition of 10 ng g⁻¹ of *Romashkino* resulting in a decrease in albedo to 97.4% of the unpolluted albedo 320 and 10 ng g⁻¹ of *Petrobaltic* to 99.8%. Therefore, the *Romashkino* is more absorbing per unit mass than the *Petrobaltic*, in line with Otremba (2000), and the albedo decreases dramatically with increasing mass ratios of oil. For instance, at a mass ratio of 100 ng g⁻¹ Romashkino and Petrobaltic cause an albedo decrease to 83.8% and 98.3% of the unpolluted albedo, respectively, and at 1000 ng g⁻¹, Romashkino and *Petrobaltic* cause a decrease to 47.2% and 88.2% of the unpolluted albedo, respectively. The albedo 325 response to increasing mass ratios of oil is non-linear (Fig. 2) and is exemplified by the doubling of *Romashkino* mass ratios from 100 to 200 ng g⁻¹ causing a decrease in albedo to 89.8% of the 100 ng g⁻¹
- value, whilst doubling of the mass ratios from 500 to 1000 ng g⁻¹ causes a decrease in albedo to 78.3% of the 500 ng g⁻¹ value. For *Petrobaltic* there is also a non-linear response as mass ratios increase but it is 330 significantly less pronounced (98.5% and 94.8% of the unpolluted albedo, respectively).

The change in albedo is wavelength dependent, and the Mie calculations indicate the effect of oil significantly decreases as wavelength increases and the ice becomes more absorbing. At a wavelength of 550 nm, 1000 ng g⁻¹ of *Romashkino* causes albedo to decrease to 81.6% of the unpolluted albedo, and at 700 nm, this decrease is to 95.5% of the unpolluted albedo. The effect *Petrobaltic* has on albedo above 500 nm is negligible, even at larger mass ratio. Therefore, at shorter wavelengths (400–550 nm) the albedo 335

is more responsive to the oil whereas at longer wavelengths (550–700 nm), the albedo is more responsive to the ice.



340 Fig. 1. Wavelength dependent albedo of first-year sea ice at increasing mass ratios of *Romashkino* (solid) and *Petrobaltic* (dotted) oils (0–1000 ng g⁻¹) with a median droplet radius of 0.05 μm. The first-year sea ice is quasi-infinite (15 m thick), and the properties are described in Table 1. These data demonstrate sea ice albedo is significantly more responsive to *Romashkino* oil than *Petrobaltic* oil.

3.2 The sensitivity of each type of quasi-infinite sea ice to increasing oil

- 345 Figure 2 illustrates the surface albedo response of quasi-infinite multi-year, first-year, and melting sea ice as mass ratios of *Romashkino* and *Petrobaltic* oils increase from 0–1000 ng g⁻¹. The three types of sea ice have different unpolluted albedos: melting sea ice 0.72, first-year sea ice 0.87, and multi-year sea ice 0.94 at a wavelength of 400 nm respectively, owing to their different scattering cross sections (Perovich, 1996; Marks and King, 2014). The three types of sea ice also respond differently to increasing mass ratios of oil, analogous to previous studies exploring the response of sea ice to mineral dust (Lamare et al., 2016). As the mass ratio of *Romashkino* oil increases in multi-year sea ice the albedo decreases to 98.8% at 10 ng g⁻¹, 92.4% at 100 ng g⁻¹, and 70.9% at 1000 ng g⁻¹ of the unpolluted albedo at a wavelength of 400 nm. In first-year sea ice, the albedo decreases to 97.4% at 10 ng g⁻¹, 83.8% at 100 ng g⁻¹, and 47.2% at 1000 ng g⁻¹ of the unpolluted albedo at a wavelength of 400 nm. In melting sea ice, the albedo at a wavelength of 400 nm.
- decreases to 94.9% at 10 ng g⁻¹, 68.8% at 100 ng g⁻¹, and 22.1% at 1000 ng g⁻¹ of the unpolluted albedo

at a wavelength of 400 nm. The same response occurs with *Petrobaltic* oil pollution, although the effect is much weaker, with multi-year sea ice albedo decreasing to 94.5% at 1000 ng g^{-1} ; first-year sea ice albedo decreasing to 88.2% at 1000 ng g^{-1} ; and melting sea ice albedo decreasing to 77% at 1000 ng g^{-1} of the unpolluted albedo at a wavelength of 400 nm. Therefore, the most responsive type of sea ice to oil pollution is melting sea ice whilst multi-year sea ice is the least responsive.

360

365

At longer wavelengths, the three types of sea ice are less responsive to increasing mass ratios of either *Romashkino* or *Petrobaltic* as the absorption coefficient of the ice increases (Warren, 2019). At a wavelength of 500 nm increasing mass ratios of *Romashkino* oil significantly effect albedo, however at wavelengths longer than 500 nm the effect of oil pollution is negligible for the three types of sea ice. Evidently the albedos of the three types of sea ice are very wavelength dependent and the response in albedo to increasing mass ratios of oil significantly decreases as ice absorption increases at longer wavelengths (Warren and Brandt, 2008; Warren, 2019).

Both the type of oil and the type of ice are significant in changing the albedo. There is a large difference in how responsive each type of sea ice is to increasing mass ratios of oil and melting sea ice is

- 370 significantly more affected by oil pollution than multi-year sea ice. There is also a significant difference in the effect that different oils have on sea ice albedo. At a wavelength of 400 nm and a relatively low increase in mass ratios of oil (10–100 ng g⁻¹), *Romashkino* oil causes a relative decrease in albedo to 72.5% in melting sea ice, 86% in first-year sea ice, and 93.4% in multi-year sea ice. At the same wavelength and increasing mass ratios of *Petrobaltic* oil, there is a relative decrease in albedo to 99.3%
- 375 in multi-year sea ice, 98.5% in first-year sea ice, and 97% in melting sea ice. Therefore, as mass ratios of oil increase, it is the type of oil that has a larger effect on how responsive sea ice albedo is, rather than what the type of ice is.



Fig. 2. Albedos of multi-year, first-year, and melting sea ice with increasing mass ratios of *Romashkino* (A, B, and C) and *Petrobaltic* (D, E, and F) oils (0–1000 ng g⁻¹) with a median droplet radius of 0.05 μm. The selected wavelengths are 400 nm, 500 nm, 600 nm, and 700 nm. The three types of sea ice are quasi-infinite, and their properties are described in Table 1. These data demonstrate a large difference in how responsive each type of sea ice is to increasing mass ratios of *Romashkino* or *Petrobaltic* oil.

3.3 The effect of oil droplet size distribution on the albedo of quasi-infinite sea ice



385

Fig 3. The albedo response of melting sea ice as mass ratios of *Romashkino* oil increase $(0-1000 \text{ ng g}^{-1})$ at a wavelength of 400 nm. The different coloured lines represent different lognormally distributed droplet sizes with median radii of 0.05–5.0 µm. The melting sea ice is quasi-infinite, and the properties are described in Table 1. These data demonstrate the sea ice albedo to be most responsive to the smallest droplet sizes (0.05-0.5 µm).

390

Figure 3 illustrates the surface albedo response of melting sea ice at a wavelength of 400 nm as mass ratios of *Romashkino* oil increase from 0–1000 ng g⁻¹ for different sized spherical droplets. Melting sea ice was selected as it is liable to have the least conservative effect owing to it having the largest penetration depth to light of the three types of sea ice. The different droplet sizes examined are characteristic of oils that have undergone different extents of weathering and are common in coastal areas, with radii of 0.05 µm, 0.25 µm, 0.5 µm, 1.5 µm, 2.5 µm, 3.5 µm, and 5 µm, in line with Haule and Freda (2016) and utilising the geometric standard deviation that were experimentally measured in Otremba (2007). The albedo is most responsive to the smallest droplets sizes (0.05–0.5 µm) and progressively less responsive as the droplets increase in size. The difference in albedo response to the smallest droplets is small, with a decrease in albedo at a mass ratio of 1000 ng g⁻¹ to 21.7% of the unpolluted value for a 0.25 µm droplet, 22.1% for a 0.05 µm droplet, and 23% for a 0.5 µm droplet. Figure 4 shows the mass absorption coefficient of a larger droplet size range (0.005–50 µm) of both *Romashkino* and *Petrobaltic* oils, which are assigned a geometric standard deviation of e¹ and are displayed at a wavelength of 400,

- 405 500, 600, and 700 nm. The mass absorption coefficient of oil droplets is a function of droplet size in the Mie regime and can be seen to peak for *Romashkino* oil at a wavelength of 400 nm between 0.05 to 0.5 μm in Fig. 4. The relationship shown in Fig. 4 is an interplay between the value of the absorption efficiency from the Mie calculation and the increasing mass of a larger droplet, resulting in a decreasing mass absorption coefficient as the droplet size increases. As the wavelength increases the critical radius
- 410 (i.e., the radius with maximum mass absorption coefficient) for *Romashkino* also changes from 0.1 μm at 400 nm to 0.5 μm at 700 nm. Similarly, for *Petrobaltic* the critical radius changes with increasing wavelength from 1 μm at 400 nm to 5 μm at 700 nm. For *Romashkino* there is only a minor decrease in mass absorption coefficient for droplets smaller than the critical radius, with a 0.005 μm droplet decreasing to 87.9% of the value at 400 nm, and 80.2% at 700 nm. However, there is a significant decrease
- 415 in mass absorption coefficient for droplets larger than the critical radius, with a 50.0 µm droplet decreasing to 5.4% of the value at 400 nm, and 27.7% of the value at 700 nm. For *Petrobaltic* there is a comparable minor decrease in mass absorption coefficient for droplets smaller than the critical radius, with a 0.005 µm droplet decreasing to 78.6% of the value at 400 nm, and 46.2% at 700 nm. However, droplets larger than the critical radius do not experience the same reduction that is seen in the *Romashkino*,
- 420 and the value only decreases to 75.5% at 400 nm, and 90.8% at 700 nm for the radii considered, presumably owing to the significantly weaker mass absorption coefficient. Figure 4 demonstrates the need for the size distribution of oil to be known for accurate reproductions of oil spills to be possible.





3.4 The effect of increasing background black carbon on the albedo of quasi-infinite sea ice

- Figure 5 illustrates the surface albedo of melting, first-year, and multi-year sea ice as a function of increasing mass ratios of *Romashkino* oil from 0–1000 ng g⁻¹ and background black carbon levels from 1–100 ng g⁻¹. The upper row of Fig. 5 (A, C, and E) indicate albedo versus increasing mass ratios of *Romashkino* oil, whereas the lower row (B, D, and F) is a metric for the sensitivity of an ice to oil in the presence of black carbon, where sensitivity is the rate of change in albedo with increasing mass ratios of *Romashkino* oil (i.e., ΔA/ΔM).
- Black carbon has a very strong absorption cross section over shortwave wavelengths which is approximately ~10 m² g⁻¹ between 400–700 nm (Marks and King, 2013), although global mean values range between 4.3–15 m² g⁻¹ (Bond et al., 2013). If black carbon is present the three types of sea ice have a decreased sensitivity to increasing mass ratios of oil pollution, as the largest decrease in albedo occurs in the cleanest ices with the smallest black carbon mass ratios. At a 1 ng g⁻¹ mass ratio of black carbon,
- 440 the albedos of the three sea ices are very responsive to low mass ratios of oil of less than 100 ng g⁻¹. Melting sea ice is the most responsive sea ice to oil and has the largest sensitivity at 1ng g⁻¹ of black carbon, whereas multi-year sea ice is the least responsive and has the lowest sensitivity comparatively. As the mass ratio of black carbon increases, the sensitivity of the three sea ices to increasing mass ratios of oil rapidly decreases. Even at typical Arctic values of black carbon, which range between 5.5–19.2 ng
- g⁻¹, the sensitivity is significantly decreased (Grenfell et al., 2002; Jiao et al., 2014). The 100 ng g⁻¹ black carbon mass ratios are extreme values (Jiao et al., 2014) and significantly decrease the albedo of all three ices without any oil pollution present: multi-year sea ice albedo decreases to 59% of its 1 ng g⁻¹ black carbon value, the first-year sea ice albedo decreases to 32.4%, and the melting sea ice decreases to 11.8%. At such high mass ratios of black carbon, the responsive of the sea ice is significantly decreased as mass
- 450 ratios of oil increase. Therefore, oil pollution will have a significantly greater effect on the albedo of the cleanest ice that is not already polluted by large mass ratios of black carbon.

Fig. 5. Albedos of multi-year (A), first-year (C) and melting (E) sea ice with increasing mass ratios of *Romashkino* oil (0–1000 ng g⁻¹) with a peak droplet radius of 0.05 μm. The gradients (^{ΔA}/_{Δm}) of plots A, C, and E are shown for multi-year (B), first-year (D), and melting (F) sea ice, respectively. The selected wavelength is 400 nm and the background levels of black carbon increase: 1 ng g⁻¹ (black line), 10 ng g⁻¹ (red line), 20 ng g⁻¹ (blue line), and 100 ng g⁻¹ (green line). The three types of sea ice are quasi-infinite, and their properties are described in Table 1. These data demonstrate that increasing mass ratios of black carbon rapidly decrease the albedo response of each type of sea ice to *Romashkino* oil.

460 **3.5** The effect of oil, type of sea ice, and mass ratio on realistic thickness sea ice albedo

465

Calculations in section 3.2 are replicated with realistic thickness melting, first-year, and multiyear sea ice as a function of increasing mass ratios of *Romashkino* and *Petrobaltic* oils from 0–1000 ng g^{-1} and are illustrated in Fig. 6. Typical Arctic thicknesses of 2.5 m for multi-year and melting sea ice (Bourke and Garrett, 1987; Laxon et al., 2003; Weeks, 2010), and 0.8 m for first-year sea ice (Weeks, 2010) are used with a realistic background Arctic black carbon mass ratio of 5.5 ng g^{-1} (Grenfell et al., 2002). The realistic thickness sea ices have different unpolluted albedos from the quasi-infinite sea ice (melting sea ice 0.41, first-year sea ice 0.58, and multi-year sea ice 0.87 at a wavelength of 400 nm respectively) and are comparable with previous literature (Marks and King, 2014; Lamare et al., 2016). The albedo response of each sea ice is decreased from the quasi-infinite sea ice albedo response owing to

- 470 the presence of the dark ocean layer below the ice, with a 1000 ng g⁻¹ mass ratio of *Romashkino* oil resulting in the albedo decreasing to 75.3% for the multi-year sea ice; 66.3% for the first-year sea ice; and 35.9% for the melting sea ice relative to the unpolluted albedo at a wavelength of 400 nm. The albedo response to increasing mass ratios of *Romashkino* oil at longer wavelengths is weaker and essentially insignificant at wavelengths longer than 500 nm. Similarly, the albedo response to *Petrobaltic* oil at a
- 475 1000 ng g⁻¹ mass ratio is negligible, with albedo decreasing to 96.8% for the multi-year sea ice; 97% for the first-year sea ice; and 90.9% for the melting sea ice relative to the unpolluted albedo at a wavelength of 400 nm. Evidently the thickness and type of sea ice, as well as the type of oil, are important components in the effect that increasing mass ratios of oil have on sea ice albedo. As the ice becomes thinner photons are likely to penetrate the ice and be absorbed by the underlying dark ocean layer. In the quasi-infinite
- 480 sea ice photons were only weakly absorbed by the ice and efficiently by the oil. The first-year sea ice is relatively less responsive to increasing mass ratios of oil than either the melting sea ice or multi-year sea ice owing to its 0.8 m thickness compared to a 2.5 m thickness. This decreased response in albedo is indicated by the first-year sea ice albedo decreasing by 0.2 as the *Romashkino* oil mass ratio increases from 0 to 1000 ng g⁻¹, whereas the multi-year sea ice albedo decreases by 0.22. The dramatic difference
- 485 in albedo response to *Romashkino* oil and *Petrobaltic* oil indicates that the type of oil has a greater effect on sea ice albedo than the type or thickness of sea ice.

Fig. 6. Albedos of multi-year, first-year, and melting sea ice with increasing mass ratios of *Romashkino* (A, B, and C) and *Petrobaltic* (D, E, and F) oils (0–1000 ng g⁻¹) with a median droplet radius of 0.05 μm.
The selected wavelengths are 400 nm, 500 nm, 600 nm, and 700 nm. The melting sea ice and multi-year sea ice are 2.5 m thick; the first-year sea ice is 0.8 m thick, and their properties are described in Table 1. These data demonstrate that the type of oil rather than the type of sea ice may have the largest effect on the albedo of realistic thickness sea ice.

495 4 Discussion

500

The discussion is separated into seven sections: the Mie calculations, the effect of different droplet size distributions, the response of sea ice albedo to *Petrobaltic* or *Romashkino* oil, the effect of *Romashkino* and *Petrobaltic oil* on different types of sea ice, the albedo response of quasi-infinite sea ice and realistic thickness sea ice to oil pollution, potential implications of the study, and potential limitations of the study.

4.1 Mie calculations

Several studies have explored the effects of aerosol pollution on snow and sea ice (e.g., Warren and Wiscombe, 1980; Warren, 1984; Clarke and Noone, 1985; Warren and Clarke, 1990; Light et al., 1998; Grenfell et al., 2002; Aoki et al., 2003; Lee-Taylor and Madronich, 2002; Jacobson, 2004; Flanner et al., 2007; Zender et al., 2009; Doherty et al., 2010; Dumont et al., 2010; Zatko et al., 2013; Marks and King, 2013; 2014; Lamare et al., 2016; Marks et al., 2017) but there has been limited research into the effect of oil pollution other than two basic field studies by NORCOR (1975) and Gavrilo and Tarashkevich (1992). To the authors knowledge, this is the first time the effect that small mass ratios of dispersed submicron weathered oil droplets have on the wavelength dependent albedo of sea ice are
considered. Whilst studies have looked at how oil pollution affects the inherent optical properties (i.e., absorption, scattering, backscattering, and attenuation coefficients) of liquid seawater (e.g., Otremba, 2007; Haule et al., 2015; Haule and Freda, 2016), they have not examined what the effect is on sea ice and its albedo. In this study, background levels of oil pollution are considered over a wide area and not within close proximity to any single large and acute event (e.g., Macondo Oil Spill or Exxon Valdez).

- 515 Oil is known to entrain itself into sea ice in several ways: (1) oil beneath the sea ice is frozen into the sea ice and the ice sheet continues to grow underneath the sea ice, with the oil being fully encapsulated in the ice matrix within 18–72 hours (NORCOR, 1975; Dickins and Buist, 1981; Buist and Dickins, 1983; Buist et al., 1983; Wilkinson et al., 2015); (2) Oil will drift with the sea ice in broken ice fields when concentrations of sea ice are greater than 60–70% and is incorporated into the sea ice during freeze up 520 (Venkatesh et al., 1990); (3) the oil can become trapped in small cavities under the sea ice and may or
- may not be encapsulated, depending on the time of year (Drozdowski et al., 2011). Once encapsulated in the sea ice the oil can migrate to the surface owing to its buoyancy, however this rate is heavily dependent upon the amount of brine drainage within the sea ice and therefore the time of year (Drozdowski et al., 2011). Martin (1979) found that during the winter oil formed thin lenses in the sea ice, which then
- 525 distributed themselves throughout the brine channel feeder systems in the spring (February to March) as these expanded, and eventually formed horizontal layers in the upper part of the sea ice (NORCOR, 1975). Martin (1979) also found that in summer oil at the ice surface leads to melt-pond formation owing to the absorption of solar energy; once on the surface, oil will be reintroduced to the ocean in a weathered or emulsified form by melting through the ice or flowing off the sides. These studies have focused on
- 530 macroscopic quantities of oil whereas the work presented here focuses on microscopic sized background concentrations of oil which require more work. The mass ratio of 1000 ng g⁻¹ of oil in ice is possibly a large value for diffuse background pollution but is included to provide a significant upper limit to the effect of background oil pollution on sea ice. Therefore, it is unclear whether the droplets are in the matrix

or brine channels of the sea ice, so in the work presented here they are assumed to be evenly distributed

- 535 throughout the ice to replicate the different stages of the seasonal variation. The droplets are also considered to be in a pure sea ice medium (Warren and Brandt, 2008) as opposed to a brine medium (Quan and Fry, 1995) for optical calculations. The refractive index of the medium in which the oil droplet is contained may affect the value of its mass absorption coefficient established from a Mie calculation. However, as indicated by Fig. 7, the difference in mass absorption coefficient for the oil droplets between
- 540 a sea ice medium (refractive index of 1.3091 at a wavelength of 600 nm) and brine (refractive index of 1.3402 at a wavelength of 600 nm) medium are inconsequential owing to their similar refractive indices (Quan and Fry, 1995; Warren and Brandt, 2008). Thus, the position of the oil droplet within the sea ice may not be important.
- The shape of the oil droplets may also be an important factor that influences the Mie calculations. 545 Whilst the shape of oil droplets in sea ice or brine is not known, immiscible fluids are known to adopt spherical morphologies to reduce their surface relative to volume and both mechanically and chemically dispersed droplets are spherical in nature (Johansen et al., 2013). Brine channels typically have a width of 0.1 mm (Lieb-Lappen et al., 2017) which is far larger than the droplet sizes considered here, so it is unlikely that droplet morphology is affected when being transported through the brine channels
- 550 (NORCOR, 1975; Martin, 1979). Therefore, the oil droplets are assumed to be spherical in the model as spheres are computationally facile for Mie calculations and the shape must be known for the calculation.

4.2 Droplet size distribution

Tens of microns to millimetre scale droplets are present in seawater immediately after the release of oil. from either the subsurface or at the surface (e.g., Spaulding et al., 1992; Reed and Rye, 1995; Daling et al., 1997; Johansen, 2003; Zheng et al., 2003; Lima Neto et al., 2008; Socolofsky et al., 2008; 560 Gamzaev, 2009; Papadimitrakis et al., 2005, 2011; Fraga et al., 2016; Dissanayke et al., 2018). Fingas and Hollebone (2014) summarise much of the literature on the behaviour of oil in ice-infested waters. however all the research discussed involves releasing large quantities of oil either directly under or onto the ice in acute events, whereas this study focuses on background quantities of oil. After release the 565 droplets can travel several hundred square kilometres within several days (Berenshtein et al., 2020) and are exposed to weathering processes that continually decrease the droplet size (e.g., Daling et al., 1990; Resby and Wang, 2004; Dillipiane et al., 2021). The dispersed droplet distribution sizes can be described by a lognormal distribution of submicron droplets with an experimentally measured peak for weathered oil centred around a median radius of 0.05 μm (Otremba, 2007; Z. Li et al., 2011; Johansen et al., 2013; Haule et al., 2015; Haule and Freda, 2016). The focus of this study is on the small mass ratios (i.e., 1000 570 ng g⁻¹ or less) of submicron weathered oil droplets that are incorporated into the sea ice in polluted polar seas. It is likely that small droplets in this range are the most numerous and will affect large areas of polar sea ice, in addition to the fact that droplets that have been treated with dispersants will also be smaller (C. Li et al., 2017). Therefore, lognormal distributions of droplets with medians in the range 0.005-50 um 575 are considered for completeness (Fig. 4). The droplets considered in Fig. 4 were assigned a geometric standard deviation size e^1 as it is a mid-range value from Haule and Freda (2016) and to keep the

distribution width equivalent for each droplet size. The geometric standard deviation used for all other figures is consistent with measurements from Otremba (2007) and used in Haule and Freda (2016). With reference to Fig. 3, it is apparent that sea ice albedo is weakly sensitive to the droplet size of

- 580 oil in the range explored. Weathered submicron droplets of absorbing oils such as *Romashkino* have a greater effect on albedo than larger micron-scale droplets, with Fig. 4 indicating the critical radius changes marginally from 0.1 µm at a wavelength of 400 nm to 0.5 µm at a wavelength of 700 nm. These weathered submicron droplets are roughly equivalent in size to the wavelength of light (~0.5 µm) and therefore Mie light scattering must be considered (Petty, 2006). The *Petrobaltic* oil mass absorption coefficient also
- 585 decreases as the droplet size increases, albeit more slowly than the *Romashkino* oil owing to its very low values. This weak dependence of sea ice albedo to droplet size may have implications for remote sensing and droplet size estimates may be important to calculate the effect of oil pollution on sea ice albedo for future events.

4.3 Effect of different oils in quasi-infinite sea ice

590 Both the real and imaginary refractive indices of *Romashkino* and *Petrobaltic* oil are described as the respective largest and smallest values that oils capable of creating surface films possess, according to Otremba (2000). Therefore, *Romashkino* have a significantly larger mass absorption cross section than *Petrobaltic*; roughly 13-fold greater for a 0.5 μm droplet at a wavelength of 400 nm. The resulting differences in effect on sea ice albedo are therefore very significant, with 1000 ng g⁻¹ of *Romashkino*

595 decreasing albedo of first-year ice from 0.87 to 0.41 (a decrease to 47% relative to the unpolluted value)

at 400 nm, whereas *Petrobaltic* decreases the albedo from 0.87 to 0.77 (a decrease to 88.2% relative to the unpolluted value). As the ice becomes more absorbing at longer wavelengths (Warren and Brandt, 2008) this difference is decreased: at 550 nm 1000 ng g⁻¹ of *Romashkino* causes a decrease in albedo to 81.6% and *Petrobaltic* to 99.3%. At 700 nm the effect is very low for both oils. Thus, even at small mass ratios, heavy, and opaque oils such as *Romashkino* (0.94 g cm³) will have a significant effect on sea ice albedo whilst relatively transparent and light oils, such as *Petrobaltic* (0.77 g cm³), may have an

insignificant effect on albedo (Otremba, 2005; 2007; Hollebone, 2015). Therefore, both *Romashkino* and *Petrobaltic* can be regarded as the upper and lower respective brackets of the effect that oil pollution can have on sea ice albedo. This may be of note as the most commonly used fuel for commercial vessels is
Heavy Fuel Oil which is defined as having a density of 0.9 g cm⁻³ (IMO, MARPOL Annex I) and is therefore analogous to *Romashkino* oil.

4.4 Response of melting, first-year, and multi-year sea ice to oil

600

The three different sea ices considered here are all modelled at a quasi-infinite thickness of 15 m, in line with similar studies (e.g., Marks and King, 2014; Lamare et al., 2016) to ensure that the albedo was independent of the layer under the sea ice and any effects that a non-optically thick sea ice may have, whilst the properties of sea ice type, mass ratio and droplet size are considered. Although such large thicknesses are not typical in nature and this causes the unpolluted albedo of all three sea ices to be exaggerated from their natural values, it is useful to understand how the Romashkino and Petrobaltic fundamentally affect albedo. The multi-year sea ice is the least affected by either type of oil, with a decrease to 70.9% and 94.5% of the unpolluted albedo at a mass ratio of 1000 ng g⁻¹ of *Romashkino* oil and *Petrobaltic* oil, respectively. The multi-year sea albedo ice is only affected by mass ratios of *Romashkino* oil exceeding 100 ng g⁻¹. However, multi-year sea ice is susceptible to oil pollution as Comfort and Purves (1982) confirmed that oil does eventually migrate to the surface, despite a thickness of several metres. First-year sea ice is more sensitive to oil pollution and decreased to 47.2% and 88.2%

- 620 of the unpolluted albedo at a mass ratio of 1000 ng g⁻¹ of *Romashkino* oil and *Petrobaltic* oil, respectively. The first-year sea ice albedo is affected by mass ratios of *Romashkino* exceeding 50 ng g⁻¹. The most sensitive sea ice to oil pollution by a considerable margin is the melting sea ice which decreased to 22.1% and 77% of the unpolluted albedo at a mass ratio of 1000 ng g⁻¹ of *Romashkino* oil and *Petrobaltic* oil, respectively. The melting sea ice albedo is affected by mass ratios of *Romashkino* exceeding 25 ng g⁻¹.
- 625 Melting sea ice is the least reflective type of sea ice with a scattering cross section of 0.03 compared to 0.15 (first-year sea ice) and 0.75 (multi-year sea ice) (Perovich, 1996; Marks and King, 2014), resulting in it becoming substantially more absorbing as pollutants are added. Whilst these findings are unsurprising, they are concerning as multi-year Arctic sea ice has declined dramatically from 64% of the total sea ice in March 1985 to 30% in March 2020, with 4+ year old sea ice decreasing from 33% to
- 630 4.4% respectively (Perovich et al., 2020). As these data show, this decline in perennial types of sea ice may render the Arctic more vulnerable to increased oil pollution in the region, particularly as it is opened to both shipping and oil extraction.

4.5 The effect of depth

The work described here has focused on quasi-infinite sea ice throughout to ensure that it is 635 optically thick and therefore independent of the layer under the sea ice. Section 4.5 explores the response of realistic sea ice thicknesses and black carbon mass ratios to offer a realistic comparison with the quasiinfinite thicknesses. In comparing the quasi-infinite sea ice with the realistic sea ice (i.e., Fig. 2 and Fig. 6) it is apparent that the melting sea ice is less responsive to increasing mass ratios of Romashkino oil and essentially not affected by *Petrobaltic* oil. The photons are being continuously scattered in the quasi-640 infinite sea ice and this increases the likelihood of the photons being absorbed by oil droplets, whereas in the realistic ice the photons have a greater chance of passing through the ice and being absorbed by the dark layer under the sea ice. This is also apparent in the first-year sea ice which is less affected by increasing mass ratios of both oils than the multi-year sea ice, despite having a smaller scattering cross section, because it is thinner. Even though the realistic types of sea ice are less responsive to increasing 645 mass ratios of oil, their albedos are still significantly decreased by the *Romashkino* oil. Therefore, it appears that the type of oil has the biggest effect on how responsive sea ice is to increasing mass ratios of oil owing to the large variability in oil optical properties. This differs from a similar study (Lamare et al. 2016) which concluded that the optical properties of sea ice played a more important role on the response of sea ice albedo than the type of pollutant (e.g., windblown aerosols).

650 4.6 Implications of study

Arctic multi-year and first-year sea ice are declining at 17.5% and 13.5% per decade respectively and melting sea ice is becoming more prevalent earlier in the year (Comiso, 2012; Tschudi et al., 2019). First-year and particularly melting sea ice may be more responsive to oil pollution than multi-year sea ice, so these trends indicate that sea ice albedo in the Arctic may become more vulnerable to background
levels of oil pollution as the relative amount of these types of ice dominate in the Arctic. This increasing vulnerability will be exacerbated by potential future extraction of the vast hydrocarbon reserves in the Arctic Ocean (Bird et al., 2008; Krivorotov and Finger, 2019; Czarny, 2019) and the development of northern sea routes (Ho, 2010; Eguíluz et al., 2016; Kikkas and Romashkina, 2018), resulting in vast quantities of Heavy Fuel Oil akin to *Romashkino* oil passing through the region and potentially exposing it to increased pollution (De Carolis et al., 2014; IMO, MARPOL Annex 1). Furthermore, the albedo of clean sea ice is more responsive to oil pollution than sea ice polluted with higher mass ratios of aerosols such as black carbon, so future oil pollution may imperil vast areas of pristine Arctic sea ice even at low background mass ratios.

4.7 Uncertainties

There are a few potential uncertainties to this study stemming primarily from the fact that it is a modelling study and that, to the authors knowledge, this is the first instance that the effect of background oil pollution on sea ice albedo has been considered. A limitation with the Mie calculations is the uncertainty of whether the oil droplets are spherical in the ice. A careful study of the particle morphology of oil droplets in ice is required to improve the veracity of future modelling studies. However, it is expected that the results presented here would be much better than an order of magnitude level. A second

potential limitation of this study is that it has not considered the effect of a snowpack on top of the sea ice. Sea ice is frequently covered with snow and sea ice tends to only be free of snow where it is removed by katabatic winds or during the melting season (Weeks, 2010). Whereas similar studies (e.g., Marks and King, 2013, 2014; Lamare et al., 2016) consider layers of snow on top of the ice, they differ in that the

- 675 pollutants explored are atmospherically deposited directly on top of both the ice and snow. For the oil considered here to be incorporated into the ice it must be frozen into the ice directly from the seawater below (NORCOR, 1975; Martin, 1979; Dickins and Buist, 1981; Buist and Dickins, 1983; Buist et al., 1983; Wilkinson et al., 2015; Drozdowski et al., 2011). However, Owens et al. (2005) found that if snow comes into contact with oil, for instance if the snow-ice forms from the surface flooding of an ice floe, it
- 680 can initially absorb up to 70% of the oil thus decreasing the albedo of the snow. Furthermore, a layer of oil underneath the snowpack may expedite melting from below as the oil may be up to 6 °C warmer than the air temperature (Chen, 1972). The amount of snow cover on sea ice varies significantly across the Arctic (Zhou et al., 2021) so some areas may be more vulnerable to oil pollution earlier in the spring. Nevertheless, even a thin snow cover (less than 1 cm) can increase the albedo of sea ice and can be
- 685 considered optically thick when it exceeds 10 cm (France et al., 2011; Marks and King, 2013). Therefore, similar to the work by Light et al (1998), the findings of this study may only be valid during the ablation season when snow cover has melted or been removed by wind.

Throughout this study the typical mid-range values for sea ice have been selected and are reported in Table 1. To assess any uncertainties that may arise from the variability of scattering cross sections reported in literature (Grenfell and Maykut, 1977; Perovich, 1990, 1996; Timco and Frederking, 1996; Gerland et al., 1999; Simpson et al., 2002; Fisher et al., 2005; King et al., 2005; France et al., 2011; Marks and King, 2013) Fig. 8 is additionally presented and indicates the response of realistic types of sea ice with both lower and higher scattering cross sections. There is a natural variability in sea ice density between 700–950 kg m⁻³ (Grenfell and Maykut, 1977; Perovich, 1990, 1996; Timco and Frederking,

- 695 1996; Gerland et al, 1999) which can be propagated to a variation in scattering cross section of approximately a factor of 0.88 and 1.19 of the original values for the lowest and highest reported densities, respectively. These ranges are much smaller than the natural variation in the scattering of sea ice, as shown in Fig. 8. The variation due to density of ice in this study is not considered important.
- This manuscript is a sensitivity study that considers a wide variety of optical and physical parameters for the oil pollution of sea ice. Ultimately, the results indicate that background concentrations of oil pollution may have an important effect on sea ice albedo. It is anticipated that this study will act as a foundation for more complex studies to follow that, coupled with field and lab-based experiments, may explore the effects such as the movement of oil within the sea ice column in greater detail.

Fig. 8. Wavelength dependent albedo of multi-year (A), first-year (B), and melting sea ice (C). Shown here is the variability in albedo of each type of sea ice based on scattering cross sections described in literature (Grenfell and Maykut, 1977; Perovich, 1990, 1996; Timco and Frederking, 1996; Gerland et al., 1999; Simpson et al., 2002; Fisher et al., 2005; King et al., 2005; France et al., 2011; Marks and King, 2013), ranging from 0.5–1 m² kg⁻¹ for multi-year sea ice; 0.1–0.2 m² kg⁻¹ for first-year sea ice; and 0.01–0.05 m² kg⁻¹ for melting sea ice. Shown in red are the data with no oil pollution present and shown in blue are the data with 1000 ng g⁻¹ of *Romashkino* oil present. The thicker lines show the values of typical ice that are used in this study. The melting sea ice and multi-year sea ice are 2.5 m thick; the first-year sea ice is 0.8 m thick, and the background concentration of black carbon is set to 5.5 ng g⁻¹.

715 **5 Conclusion**

Background mass ratios of crude oil (≤1000 ng g⁻¹) can have a significant effect on sea ice albedo. The albedo response is dependent on the type of oil, with the relatively light absorbing *Romashkino* oil having the largest effect owing to its large mass absorption coefficient. The albedo response is also dependent on the type of sea ice, with a 1000 ng g⁻¹ mass ratio of *Romashkino* oil resulting in quasiinfinite thickness (15 m thick) multi-year sea ice decreasing to 70.9%, first-year sea ice decreasing to 47.2%, and melting sea ice decreasing to 22.1% relative to the unpolluted albedo at a wavelength of 400 nm. Sea ice thickness also plays an important role on the albedo response with realistic thickness multi-year sea ice decreasing to 35.9% of the unpolluted value. The size of the oil droplets has a weak effect on the albedo response,

- 725 with weathered submicron *Romashkino* oil droplets (0.05–0.5 μm) being the most absorbing in the visible spectrum (400–700 nm). This work demonstrates that the type of oil, type of ice, thickness of ice, and droplet size are important components in the response of sea ice albedo to background mass ratios of crude oil pollution. The type of oil is the most important factor in how responsive sea ice is to increasing mass ratios of oil and is more important than the type of sea ice being polluted. By collecting these data
- 730 in the event of a future spill, it may be possible to model what the effect of background oil pollution may be. This may become increasingly important as multi-year sea ice continues to decrease, first-year sea ice becomes more dominant and there is earlier melting of sea ice in the year.

Data availability. All data have been published using Zenodo and can be accessed at: DOI.10.5281/zenodo.6583629.

735 *Acknowledgements*. The work contained in this paper contains work conducted during a PhD study undertaken as part of the Centre for Doctoral Training (CDT) in Geoscience and the Low Carbon Energy Transition. It is sponsored by Royal Holloway, University of London via their GeoNetZero CDT Studentship whose support is gratefully acknowledged.

Edited by: Yevgeny Aksenov

740 References

- Aoki, T., Hachikubo, A., and Hori, M.: Effects of snow physical parameters on shortwave broadband albedos. J. Geophys. Res., 108, 4916, https://doi.org/10.1029/2003JD003506, 2003.
- Bandara, U. C. and Yapa, P. D.: Bubble Sizes, Breakup, and Coalescence in Deepwater Gas/Oil Plumes, J. Hydraul. Eng., 137, 729–738, https://doi.org/10.1061/(ASCE)HY.1943-7900.0000380, 2011.
- 745 Berenshtein, I., Paris, C. B., Perlin, N., Alloy, M. M., Joye, S. B., and Murawski, S.: Invisible oil beyond the Deepwater Horizon satellite footprint, Sci. Adv., 6, eaaw8863, https://doi.org/10.1126/sciadv.aaw8863, 2020.
 - Bird, K. J., Charpentier, R. R., Gautier, D. L., Houseknecht, D. W., Klett, T. R., Pitmna, J. K., Moore, T. E., Schenk, C. J., Tennyson, M. E., and Wandrey, C. J.: Circum-Arctic resource appraisal; estimates
- 750 of undiscovered oil and gas north of the Arctic Circle, U.S. Geol. Surv., 2008-3049, https://doi.org/10.3133/fs20083049, 2008.
 - Bohren, C. F. and Huffman, D. R. (Eds.): Absorption and Scattering of Light by Small Particles, John Wiley & Sons, Weinheim, Germany, 1983.
 - Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M.
- G., Ghan, S., Kaercher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: a scientific assessment, J. Geophys. Res.: Atmos., 118, 5380–5552, https://doi.org/10.1002/jgrd.50171, 2013.
 - Bourke, R. H. and Garrett, R. P.: Sea ice thickness distribution in the Arctic Ocean, Cold Reg. Sci. Technol., 13, 259–280, https://doi.org/10.1016/0165-232X(87)90007-3, 1987.
 - Brandvik, P. J. and Faksness, L.-G.: Weathering processes in Arctic oil spills: Meso-scale experiments with different ice conditions, Cold Reg. Sci. Tech., 55, 160–166, https://doi.org/10.1016/j.coldregions.2008.06.006, 2009.
- Brandvik, P. J., Daling, P. S., and Myrhaug, L. J.: Mapping weathering properties as a function of ice conditions: a combined approach using a flume basin verified by large-scale field experiments, in: Proceedings of the 33rd Arctic and Marine Oilspill Program (AMOP) Technical Seminar on Environmental Contamination and Response, Environment Canada, Ottawa, Ontario, Canada, 7–9 June 2010, 701–723, 2010.
 - 26

- Brandvik, P. J., Johansen, Ø., Leirvik, F., Farooq, U., and Daling, P. S.: Droplet breakup in subsurface oil releases – Part 1: Experimental study of droplet breakup and effectiveness of dispersant injection, Mar. Pollut. Bull., 73, 319–326, https://doi.org/10.1016/j.marpolbul.2013.05.020, 2013.
- Brandvik, P.J., Johansen, Ø., Farooq, U., Glen, A., and Leirvik, F.: Subsurface oil releases Experimental
- study of droplet distributions and different dispersant injection techniques Version 2, Technical Report SINTEF, Trondheim, Norway, A26122, 99pp., 2014.
 - Brandvik, P. J., Davies. E., Leirvik, F., Johansen, Ø., and Belore, R.: Large-scale basin testing to simulate realistic oil droplet distributions from subsea release of oil and the effect of subsea dispersant injection, Mar. Pollut. Bull., 163, 111934, https://doi.org/10.1016/j.marpolbul.2020.111934, 2021.
- 780 Buckley, R. G. and Trodahl, H. J.: Thermally driven changes in the optical properties of sea ice, Cold Reg. Sci. Technol., 14, 201–204, 1987.
 - Buist, I. A. and Dickins, D. F.: Fate and Behavior of Water-in-Oil Emulsions in Ice, Prepared by Dome Petroleum Ltd. For the Canadian Offshore Oil Spill Research Association (COOSRA), Calgary, Report No. CS 11., 1983.
- 785 Buist, I. A., Potter, S. G., and Dickins, D. F.: Fate and Behavior of Water-in-Oil-Emulsions in Ice, in: Proceedings of the Sixth Arctic Marine Oilspill Program Technical Seminar, Environment Canada, Ottawa., 1983.

- Chao, J., Liu, C., Li, Y., Lin, X., and Yan, Y.: Characteristics of the sea ice reflectance spectrum polluted by oil spills based on field experiments in the Bohai Sea, Acta. Oceanol. Sin., 36, 73–79, https://doi.org/10.1007/s13131-017-0995-1, 2017.
- Chen, E. C.: Arctic Winter Oil Spill Test, Technical Bulletin No. 68, Environment Canada, Ottawa, ON. 20 pp., 1972.
 - Clarke, A. D. and Noone, K. J.: Soot in the Arctic snowpack: A cause for perturbations in radiative transfer, Atmos. Environ., 19, 2045–2053, https://doi.org/10.1016/0004-6981(85)90113-1, 1985.
- Comfort, G. and W. Purves.: The Behaviour of Crude Oil Spilled Under Multi-Year Ice. EPS 4-EC-82 4, Environment Canada, Ottawa, ON., 76 pp., 1982.
 - Comiso, J. C.: Large Decadal Decline of the Arctic Multiyear Ice Cover, J. Clim., 25, 1176–1193, https://doi.org/10.1175/JCLI-D-11-00113.1, 2012.
- Czarny, R. S.: The Arctic: Numerous Interests and Multiple Players, Toruńskie Studia 800 Międzynarodowe, 12, 125-142, 2019.
 - D'Asaro, E.A.: Turbulence in the upper-ocean mixed layer, Annu. Rev. Mar. Sci., 6, 101–115, https://doi.org/10.1146/annurev-marine-010213-135138, 2014.
- Daling, P. S., Brandvik, P. J., Mackay, D., and Johansen, Ø.: Characterization of crude oils for environmental purposes, Oil Chem. Poll., 7, 199–224, https://doi.org/10.1016/S0269-8579(05)80027-9, 1990.
 - Daling, P. S., Aamo, O. M., Lewis, A., Strøm-Kristiansen, T.: Sintef/iku oil-weathering model: predicting oils' properties at sea, in: International Oil Spill Conference Proceedings (1997), Fort Lauderdale, Florida, USA, 2–10 April 1997, 297–307, 1997.
 - De Carolis, G., Adamo, M., and Pasquariello, G.: On the estimation of thickness of marine oil slicks from
- 810 sun-glittered, near-infrared MERIS and MODIS Imagery: The Lebanon Oil Spill Case Study, IEEE Trans. Geosci. Rem. Sci., 52, 559–573, https://doi.org/10.1109/TGRS.2013.2242476, 2014.

Delvigne, G. A. L. and Sweeney, C. E.: Natural dispersion of oil, Oil Chem. Pollut., 4, 281–310, https://doi.org/10.1016/S0269-8579(88)80003-0, 1988.

- Delvigne, G. A. L. and Hulsen, L. J. M.; Simplified Laboratory Measurements of Oil Dispersion
 Coefficient Application in Computations of Natural Oil Dispersion, in: Proceedings of the 17th
 Arctic Marine Oil Spill Program (AMOP) Technical Seminar, Vancouver, Canada, 8-10 June, 173-187, 1994.
 - Dickins, D.: Arctic Oil Spill Response Technology Joint Industry Programme Synthesis Report, International Association of Oil and Gas Producers, DF Dickins Associates, LLC, London, UK, 2017.

- Dickins, D. and Buist, I.: Oil and Gas Under Sea Ice, Prepared by Dome Petroleum Ltd. for the Canadian Offshore Oil Spill Research Association (COOSRA), Report CV-1, Volumes I and II., 1981.
- Dilliplaine, K., Oggier, M., Collins, R. E., Eicken, H., Gradinger, R., and Bluhm, B. A.: Crude oil exposure reduces ice algal growth in a sea-ice mesocosm experiment, Polar Biol., 44, 525–537, https://doi.org/10.1007/s00300-021-02818-3, 2021.
 - Dissanayake, A. L., Gros, J., and Socolofsky, S. A.: Integral models for bubble, droplet, and multiphase plume dynamics in stratification and crossflow, Environ. Fluid Mech., 18, 1167–1202, https://doi.org/10.1007/s10652-018-9591-y, 2018.
- Doherty, S. J., Warren, S. G., Grenfell, T. C., Clarke, A. D., and Brandt, R. E.: Light-absorbing impurities
 in Arctic snow, Atmos. Chem. Phys., 10, 11647–11680, doi:10.5194/acp-10-11647-2010, 2010.
 - Drozdowski, D., Nudds, S., Hannah, C. G., Niu, H., Peterson, I., and Perrie, W.: Review of oil spill trajectory modelling in the presence of ice, Canadian Technical Report of Hydrographic and Ocean Sciences, Fisheries and Ocean, Canada, 274 pp., 2011.
- Dumont, M., Brissaud, O., Picard, G., Schmitt, B., Gallet, J., and Arnaud, Y.: High-accuracy
 measurements of snow bidirectional reflectance distribution function at visible and NIR
 wavelengths-comparison with modelling result, Atmos. Chem. Phys., 10, 2507–2520, https://doi.org/10.5194/acp-10-2507-2010, 2010.
 - Edson, J., Crawford, T., Crescenti, J., Farrar, T., Frew, N., Gerbi, G., Helmis, C., Hristov, T., Khelif, D., Jessup, A., Jonsson, H., Li, M., Mahrt, L., McGillis, W., Plueddemann, A., Shen, L., Skyllingstad,
- E., Stanton, T., Sullivan, P., Sun, J., Trowbridge, J., Vickers, D., Wang, S., Wang, Q., Weller, R., Wilkin, J., Williams, III, A. J., Yue, D. K. P., Zappa, C.: The coupled boundary layers and air-sea transfer experiment in low winds, Bull. Amer. Meteor. Soc., 88, 341–356, https://doi.org/10.1175/BAMS-88-3-341, 2007.
- Eguíluz, V., Fernández-Gracia, J., Irigoien, X., and Duarte, C. M.: A quantitative assessment of Arctic shipping in 2010–2014, Sci. Rep., 6, 30682, https://doi.org/10.1038/srep30682, 2016.
- Fetterer, F., Knowles, K., Meier, W., Savoie, M., and Windnagel, A. K.: Sea Ice Index, Version 3, National Snow and Ice Data Center (NSIDC), Boulder, Colorado USA, https://doi.org/10.7265/N5K072F8, 2017.
- Fingas, M. F. and Hollebone, B. P.: Oil behaviour in ice-infested waters, in: International Oil Spill Conference Proceedings 2014, Savannah, Georgia, USA, 5–8 May, 1239-1250, 2014.
 - Fisher, F., King, M., and Lee-Taylor, J.: Extinction of UV-visible radiation in wet midlatitude (maritime) snow: implications for increased NOx emission, J. Geophys. Res., 110, D21301, https://doi.org/10.1029/2005JD005963, 2005.

Flanner, M. G., Zender, C. S., Randerson, J. T., and Rasch, P. J.: Present-day climate forcing and response from black carbon in snow, J. Geophys. Res.-Atmos., 112, D11202,

https://doi:10.1029/2006JD008003, 2007.

855

865

- Fraga, B., Stoesser, T., Lai, C. C. K., Socolofsky, S. A.: A LES-based Eulerian–Lagrangian approach to predict the dynamics of bubble plumes, Ocean Model., 97, 27–36, https://doi.org/10.1016/j.ocemod.2015.11.005, 2016.
- 860 France, J. L.: Chemical oxidation in snowpacks, Ph. D. Thesis, Royal Holloway, University of London, 2008.
 - France, J. L., King, M. D., Frey, M. M., Erbland, J., Picard, G., Preunkert, S., MacArthur, A., and Savarino, J.: Snow optical properties at Dome C (Concordia), Antarctica; implications for snow emissions and snow chemistry of reactive nitrogen, Atmos. Chem. Phys., 11, 9787–9801, https://doi.org/10.5194/acp-11-9787-2011, 2011.
 - Gamzaev, K. M.: Modeling the spread of an oil slick on the sea surface, J. Appl. Mech. Tech. Phys., 50, 466–469, https://doi.org/10.1007/s10808-009-0062-6, 2009.
 - Gavrilo, V. P. and Tarashkevich, V. N.: Study of oil intrusion influence on physical properties of sea ice, in: Proceedings of the Second International Offshore and Polar Engineering Conference, San Francisco, USA, 14-19 June, 666-669, 1992.
 - Gerland, S., Winther, J., Ørbæk, J., Liston, G., Øritsland, N., Blanco, A., and Ivanov, B.: Physical and optical properties of snow covering Arctic tundra on Svalbard, Hydrol. Process., 13, 2331–2343, https://doi.org/10.1002/(SICI)1099-1085(199910)13:14/15<2331::AID-HYP855>3.0.CO;2-W, 1999.
- 875 Glaeser, J. L. and Vance, G.: A study of the behaviour of oil spills in the arctic, Report Number 714/08/A/001,002, United States Coast Guard, Washington, DC, USA, 53 pp., 1971
 - Gonçalves, R. C., Iskandarani, M., Srinivasan, A., Thacker, W. C., Chassignet, E., and Knio, O. M.: A framework to quantify uncertainty in simulations of oil transport in the ocean, J. Geophys. Res.: Oceans, 121, 2058–2077, https://doi:10.1002/2015JC011311, 2016.
- 880 Grenfell, T. C.: A radiative transfer model for sea ice with vertical structure variations, J. Geophys. Res.: Oceans., 96, 16991–17001, https://doi.org/10.1029/91JC01595, 1991.
 - Grenfell, T. C. and Maykut, G. A.: The optical properties of ice and snow in the Arctic Basin, J. Glaciol., 18, 445–463, https://doi.org/10.3189/S0022143000021122, 1977.
- Grenfell, T. C. and Perovich, D. K.: Spectral albedos of sea ice and incident solar irradiance in the southern Beaufort Sea, J. Geophys. Res.: Oceans, 89, 3573–3580, https://doi.org/10.1029/JC089iC03p03573, 1984.
 - Grenfell, T. C., Light, B., and Sturm. M.: Spatial distribution and radiative effects of soot in the snow and sea ice during the SHEBA experiment, J. Geophys. Res.: Oceans, 107, SHE 7-1-SHE7-7, https://doi.org/10.1029/2000JC000414, 2002.
- 890 Gulas, S., Downton, M., D'Souza, K., Hayden, K., and Walker, T. R.: Declining Arctic Ocean oil and gas developments: Opportunities to improve governance and environmental pollution control, Mar. Pol., 75, 53–61, https://doi.org/10.1016/j.marpol.2016.10.014, 2017.
 - Hancke, K., Lund-Hansen, L. C., Lamare, M. L., Højlund Pedersen, S., King, M. D., Andersen, P., and Sorrell, B. K.: Extreme Low Light Requirement for Algae Growth Underneath Sea Ice: A Case

- 895 Study From Station Nord, NE Greenland, J. Geophys. Res.: Oceans, 123, 985–1000, https://doi.org/10.1002/2017JC013263, 2018.
 - Hanesiak, J. M., Barber, D. G., De Abreu, R. A., and Yackel, J. J.: Local and regional albedo observations of arctic first-year sea ice during melt ponding, J. Geophys. Res.: Oceans, 106, 1005– 1016, https://doi.org/10.1029/1999JC000068, 2001.
- 900 Haule, K., Darecki, M., and Toczek, H.: Light penetration in seawater polluted by dispersed oil: results of radiative transfer modelling, J. Eur. Opt. Soc. -Rapid, 10, 15052, https://dx.doi.org/10.2971/jeos.2015.15052, 2015.

920

- Haule, K. and Freda, W.: The effect of dispersed Petrobaltic oil droplet size on photosynthetically active radiation in marine environment, 23, 6506–6516, https://doi.org/ 10.1007/s11356-015-5886-4, 2016.
- Hesketh, R. P., Etchells, A. W., and Russell, T. W. F.: Experimental observations of bubble breakage in turbulent flow. Ind. Eng. Chem. Res. 30, 835–841, https://doi.org/10.1021/ie00053a005, 1991.
- Ho, J.: The implications of Arctic sea ice decline on shipping, Marine Policy, 34, 713-715, https://doi.org/10.1016/j.marpol.2009.10.009, 2010.
- 910 Hollebone, B.: The oil properties data appendix, in: Handbook of oil spill science and technology, Fingas, M., Ed. John Wiley and Sons, New York, 577–681, 2015.
 - IMO, MARPOL Annex I—Prevention of Pollution by Oil: http://www.imo.org/OurWork/Envi ronment/PollutionPrevention/OilPollution/Pages/Default.aspx, last access: 29 April 2021.
- Ivanov, B., Bezgreshnov, A., Kubyshkin, N., and Kursheva, A.: Spreading of oil products in sea ice and their influence on the radiation properties of the snow-ice cover, in: Proceedings of the International Conference on Port and Ocean Engineering under Arctic Conditions, POAC, Potsdam, New York, USA, 26–30 June 2005, 853–862, 2005.
 - Jacobson, M. Z.: Climate response of fossil fuel and biofuel soot, accounting for soot's feedback to snow and sea ice albedo and emissivity, J. Geophys. Res.-Atmos., 109, D21201, https://doi:10.1029/2004JD004945, 2004.
 - Jiao, C., Flanner, M. G., Balkanski, Y., Bauer, S. E., Bellouin, N., Berntsen, T. K., Bian, H., Carslaw, K. S., Chin, M., De Luca, N., Diehl, T., Ghan, S. J., Iversen, T., Kirkevåg, A., Koch, D., Liu, X., Mann, G. W., Penner, J. E., Pitari, G., Schulz, M., Seland, Ø., Skeie, R. B., Steenrod, S. D., Stier, P., Takemura, T., Tsigaridis, K., van Noije, T., Yun, Y., and Zhang, K.: An AeroCom assessment of
- 925 black carbon in Arctic snow and sea ice, Atmos. Chem. Phys., 14, 2399–2417, https://doi.org/10.5194/acp-14-2399-2014, 2014.
 - Johansen, Ø.: Development and verification of deep-water blowout models, Mar. Pollut. Bull., 47, 360–368. https://doi.org/10.1016/S0025-326X(03)00202-9, 2003.

Johansen, Ø., Rye, H., Cooper, C.: Deep spill – field study of a simulated oil and gas blow-out in deep

- 930 water. Spill Sci. Technol. Bull., 8, 433–443, https://doi.org/10.1016/S1353-2561(02)00123-8, 2003.
 - Johansen, Ø., Brandvik, P. J., and Farooq, U.: Droplet breakup in subsea oil releases Part 2: Predictions of droplet size distributions with and without injection of chemical dispersants, Mar. Pollut. Bull., 73, 327–335, https://doi.org/10.1016/j.marpolbul.2013.04.012, 2013.
- 935 Johansen, Ø., Reed, M., and Bodsberg, N. R.: Natural dispersion revisited, Mar. Pollut. Bull., 93, 20 m 26, https://doi.org/10.1016/j.marpolbul.2015.02.026, 2015.

- Kikkas, K. and Romashkina, E.: Potential Opportunities for the Arctic Transport Space, IOP Conf. Ser.: Earth Environ. Sci., 180, 012016, https://doi.org/10.1088/1755-1315/180/1/012016, 2018.
- King, M., France, J., Fisher, F., and Beine, H.: Measurement and modelling of UV radiation penetration
 and photolysis rates of nitrate and hydrogen peroxide in Antarctic sea ice: an estimate of the production rate of hydroxyl radicals in first-year sea ice, J. Photoch. Photobio. A, 176, 39–49, https://doi.org/10.1016/j.jphotochem.2005.08.032, 2005.
 - Koivurova, T. and Vanderzwaag, D. L.: The arctic council at 10 years: retrospect and prospects, University of British Columbia Law Review, 40 121–194, https://ssrn.com/abstract=1860308, 2007.

955

- Krivorotov A., Finger M.: State-Owned Enterprises in the Arctic, In: The GlobalArctic Handbook, Finger, M. and Heininen, L, Ed. Spring, Cham, Denmark, 45–62, https://doi.org/10.1007/978-3-319-91995-9_4, 2019.
- Kukulka, T., Law, K.L., Proskurowski, G.: Evidence for the influence of surface heat fluxes on turbulent
- 950 mixing of microplastic marine debris, J. Phys. Oceanogr., 46, 809–815, https://doi.org/10.1175/JPO-D-15-0242.1, 2016.
 - Lamare, M. L., Lee-Taylor, J., and King, M. D.: The impact of atmospheric mineral aerosol deposition on the albedo of snow & sea ice: are snow and sea ice optical properties more important than mineral aerosol optical properties?, Atmos. Chem. Phys., 16, 843–860, https://doi.org/10.5194/acp-16-843-2016, 2016.
 - Law, A. T. and Mahmood, Z. B.: Distribution of petroleum hydrocarbon in the south-western portion of the South China Sea, Ekspedisi Matahari '86, Serdang, Faculty of Fisheries and Marine Science, Universiti Putra, Malaysia, 1986.
- Laxon, S., Peacock, N., and Smith, D.: High interannual variability of sea ice thickness in the Arctic region, Nature, 425, 947–950, https://doi.org/10.1038/nature02050, 2003.
 - Lee-Taylor, J. and Madronich, S.: Calculation of actinic fluxes with a coupled atmosphere–snow radiative transfer model, J. Geophys. Res.-Atmos., 107, https://doi.org/10.1029/2002JD002084, 2002.
 - Li, C., Miller, J., Wang, J., Koley, S. S., and Katz, J.: Size Distribution and Dispersion of Droplets Generated by Impingement of Breaking Waves on Oil Slicks, J. Geophys. Res.: Oceans, 122, 7938– 7957, https://doi.org/10.1002/2017JC013193, 2017.
 - Li, Z., Lee, K., King, T., Boufadel, M. C., and Venosa, A. D.: Oil droplet size distribution as a function of energy dissipation rate in an experimental wave tank, in: Proceedings of the International Oil Spill Conference 2008, Savannah, Georgia, USA, 4–8 March 2008, 621–626, 2008a.
- Li, Z., Lee, K., King, T., Boufadel, M. C., and Venosa, A.D.: Assessment of chemical dispersant effectiveness in a wave tank under regular non-breaking and breaking wave conditions, Mar. Pollut. Bull., 56, 903–912, https://doi.org/10.1016/j. marpolbul.2008.01.031, 2008b.
 - Li, Z., Lee, K., King, T., Niu, H., Boufade, I.M. C., and Venosa, A. D.: Application of entropy analysis of in situ droplet-size spectra in evaluation of oil chemical dispersion efficacy, Mar. Pollut. Bull., 62, 2129–2136, https://doi:10.1016/j.marpolbul.2011.07.012, 2011.
- 975 Li, Z., Spaulding, M. L., French-McCay, D., Crowley, D., and Payne, J. R.: Development of a unified oil droplet size distribution model with application to surface breaking waves and subsea blowout releases considering dispersant effects, Mar. Pollut. Bull., 114, 247–257, https://doi.org/10.1016/j.marpolbul.2016.09.008, 2017a.

- Li, Z., Spaulding, M. L. and French-McCay, D.: An algorithm for modeling entrainment and naturally and chemically dispersed oil droplet size distribution under surface breaking wave conditions, Mar. Pollut. Bull., 119, 145–152, https://doi.org/10.1016/j.marpolbul.2017.03.048, 2017b.
 - Lieb-Lappen, R. M., Golden, E. J., and Obbard, R. W.: Metrics for interpreting the microstructure of sea ice using X-ray micro-computed tomography, Cold. Reg. Sci. Technol., 138, 24–35, https://doi.org/10.1016/j.coldregions.2017.03.001, 2017
- 985 Light, B., Eicken, H., Maykut, G. A., and Grenfell, T. C.: The effect of included participates on the spectral albedo of sea ice, J. Geophys. Res.: Oceans, 103, 27739– 27752, https://doi.org/10.1029/98JC02587, 1998.
 - Lima Neto, I. E., Zhu, D. Z., and Rajaratnam, N.: Bubbly jets in stagnant water, Int. J. Multiphase Flow, 34, 1130–1141. https://doi.org/10.1016/j.ijmultiphaseflow.2008.06.005, 2008.
- 990 Liu, B., Li, Y., Zhang, Q., and Han, L.: Assessing sensitivity of hyperspectral sensor to detect oil with sea ice, J. Spectrosc, 2016, 6584314, https://doi.org/10.1155/2016/6584314, 2016.
 - Liu, B., Li, Y., Liu, C., Xie, F., and Muller, J. -P.: Hyperspectral Features of Oil-Polluted Sea Ice and the Response to the Contamination Area Fraction, Sensors, 18, 234, https://doi.org/10.3390/s18010234, 2018.
- 995 Lunel, T.: Dispersion: Oil droplet size measurements at sea, in: Proceedings of the Sixteenth Arctic and Marine Oil Spill Program Technical Seminar, Environment Canada, Calgary, Canada, 7–9 June 1993, 1023–1055, 1993.
 - Madronich, S. and Flocke, S.: The role of solar radiation in atmospheric chemistry, in: Handbook of Environmental Chemistry, edited by: Boule, P., Springer-Verlag, New York, USA, 1–26, 1998.
- 000 Marks, A. A. and King, M. D.: The effects of additional black carbon on the albedo of Arctic sea ice: variation with sea ice type and snow cover, The Cryosphere, 7, 1193–1204, doi:10.5194/tc-7-1193-2013, 2013.

- Marks, A. A. and King, M. D.: The effect of snow/sea ice type on the response of albedo and light penetration depth (*e*-folding depth) to increasing black carbon, The Cryosphere, 8, 1625–1638, https://doi.org/10.5194/tc-8-1625-2014, 2014.
- Marks, A. A., Lamare, M. L., and King, M. D.: Optical properties of sea ice doped with black carbon an experimental and radiative-transfer modelling comparison, The Cryosphere, 11, 2867–2881, https://doi.org/10.5194/tc-11-2867-2017, 2017.
 - Martin, S.: A field study of brine drainage and oil entrainment in first-year sea ice, J. Glaciol., 22, 473–502, https://doi.org/10.3189/S0022143000014477, 1979.
- Masutani, M. S. and Adams, E. E.: Experimental study of multiphase plumes with application to deep ocean oil spills, University of Hawaii Final Report submitted to the US Department of the Interior Minerals Management Service, Contract No. 1435-01-98-CT-30964, 88pp., 2000.
- Nissanka, I. D. and Yapa, P. D.: Calculation of oil droplet size distribution in an underwater oil well
 blowout, J. Hydraul. Res., 54, 307–320, https://doi.org/10.1080/00221686.2016.1144656, 2016.
- Nissanka, I. D. and Yapa, P. D.: Oil slicks on water surface: Breakup, coalescence, and droplet formation under breaking waves, Mar. Pollut. Bull., 114, 480–493. https://doi.org/10.1016/j.marpolbul.2016.10.006, 2017.
- Nissanka, I. D. and Yapa, P. D.: Calculation of oil droplet size distribution in ocean oil spills: A review,
 Mar. Pollut. Bull., 135, 723–734, https://doi.org/10.1016/j.marpolbul.2018.07.048, 2018.

- NORCOR Engineering and Research Ltd.: The Interaction of Crude Oil with Arctic Sea Ice, Beaufort Sea Technical Report, No. 27, Beaufort Sea Project, Department of the Environment, Victoria, BC, Canada, 201 pp., 1975.
- North, E. W., Adams, E. E., Thessen, A. E., Schlag, Z., He, R., Socolofsky, S. A., Masutani, S. M., and
 Peckham, S. D.: The influence of droplet size and biodegradation on the transport of subsurface oil droplets during the Deepwater Horizon spill: a model sensitivity study, Environ. Res. Lett., 10, 024016, https://doi.org/10.1088/1748-9326/10/2/024016, 2015.
 - Notz, D. and Stroeve, J.: The trajectory towards a seasonally ice-free Arctic Ocean, 4, 407–416, https://doi.org/10.1007/s40641-018-0113-2, 2018.
- 030 Otremba, Z.: The impact on the reflectance in VIS of a type of crude oil film floating on the water surface, Opt. Express, 7, 123–134, https://doi.org/10.1364/OE.7.000129, 2000.
 - Otremba, Z.: Influence of oil dispersed in seawater on the bi-directional reflectance distribution function (BRDF), Opt. Appl., 35, 99–109, 2005.
- Otremba, Z.: Oil droplets as light absorbents in seawater, J. Opt. Soc. Am., 15, 8592–8597, https://doi.org/10.1364/OE.15.008592, 2007.
 - Owens, E. H., Dickins, D. F., and Sergy, G. A.: 2005. The behavior and documentation of oil spilled on snow- and ice-covered shorelines, in: Proceedings of the International Conference on Oil Spills, Miami Beach, Florida, USA, 15–19 May 2005, 513–519, 2005.
- Özgökmen, T. M., Poje, A. C., Fischer, P. F., Childs, H., Krishnan, H., Garth, C., Haza, A. C., and Ryan,
 E.: On multi-scale dispersion under the influence of surface mixed layer instabilities and deep flows,
 Ocean Model., 56, 16–30, https://doi.org/10.1016/j.ocemod.2012.07.004, 2012.
 - Papadimitrakis, J., Psaltaki, M., Christolis, M., and Markatos, N.: Three-dimensional oil spill modelling for coastal waters, J. Marine Env. Eng., 7, 249–260, 2005.
- Papadimitrakis, I., Psaltaki, M., and Markatos, K.: 3-D oil spill modelling. Natural dispersion and the spreading of oil-water emulsions in the water column, Glob. Nest J., 13, 325–338, https://doi.org/10.30955/gnj.000726, 2011.
 - Payne, R.: Albedo of the sea surface, J. Atmos. Sci., 29, 959–970, https://doi.org/10.1175/1520-0469(1972)029<0959:AOTSS>2.0.CO;2, 1972.
 - Perovich, D.: Theoretical estimates of light reflection and transmission by spatially complex and
- 050 temporally varying sea ice covers, J. Geophys. Res.: Oceans, 95, 9557–9567, https://doi.org/10.1029/JC095iC06p09557, 1990.
 - Perovich, D. K.: The Optical Properties of Sea Ice, US Army Corps of Engineers: Cold Regions Research and Engineering Laboratory, US Army Cold Regions Research and Engineering Laboratory, Virginia, USA, 96, 1996.
- 055 Perovich, D. K.: Complex yet translucent: the optical properties of sea ice, Phys. B: Condensed Matter, 338, 107–114, https://doi.org/10.1016/S0921-4526(03)00470-8, 2003.
 - Perovich, D. K., Maykut, G. A., and Grenfell, T. C.: Optical properties of ice and snow in the Polar Oceans. I: Observations, 1986 Technical Symposium Southeast, 7 August 1986, Orlando, USA, 0637, 232–241, 1986.
- 060 Perovich, D. K., Roesler, C. S., and Pegau, W. S.: Variability in Arctic sea ice optical properties, J. Geophys. Res.: Oceans, 103, 1193–1208, https://doi.org/10.1029/97JC01614, 1998.

- Perovich, D., Meier, W., Tschudi, M., Hendricks, S., Petty, A. A., Divine, D., Farrell, S., Gerland, S., Haas, C., Kaleschke, L., Pavlova, O., Ricker, R., Tian-Kunze, X., Webster, M., and Wood, K.: Arctic Report Card 2020: Sea Ice, available at: https://doi.org/10.25923/n170-9h57 (last access: 19 November 2021), 2020.
- Petty, G. (2nd Eds): A first course in atmospheric radiation, Sundog Publishing, Madison, Wisconsin, USA, 2006.

- Picard, G., Libois, Q., and Arnaud, L.: Refinement of the ice absorption spectrum in the visible using radiance profile measurements in Antarctic snow, The Cryosphere, 10, 2655–2672, https://doi.org/10.5194/tc-10-2655-2016, 2016.
- Praks, J., Eskelinen, M., Pulliainen, J., Pyhalahti, T., and Hallikainen, M.: Detection of oil pollution on sea ice with airborne and spaceborne spectrometer, 2004 IEEE International Geoscience and Remote Sensing Symposium, 20–24 September 2004, Anchorage, Alaska, USA, 276, https://doi.org/10.1109/IGARSS.2004.1369014, 2004.
- 075 Quan, X. and Fry, E.: Empirical equation for the index of refraction of seawater, OSA Appl. Opt., 34, 3477-3480, 1995.
 - Reay, H. J., France, J. L., and King, M. D.: Decreased albedo, efolding depth and photolytic OH radical and NO2 production with increasing black carbon content in Arctic snow, J. Geophys. Res.-Atmos., 117, D00R20, https://doi.org/10.1029/2011JD016630, 2012.
- 080 Reed, M. and Rye, H.: A three-dimensional oil and chemical spill model for environmental impact assessment, in: Proceedings of the 1995 International Oil Spill Conference, Long Beach, California, USA, February–March 1995, 61–66, https://doi.org/10.7901/2169-3358-1995-1-61, 1995.
- Reed, M., Johansen, Ø., and Leirvik, F.: Natural dispersion of heavy oil products and weathered crude oils, Interspill Conference 2009, Marseille, France, 12–14 May 2009, http://refhub.elsevier.com/S0025-326X(16)30823-2/rf0270, 2009a.
 - Reed, M., Johansen, Ø., Leirvik, F., and Brørs, B.: Numerical algorithm to compute the effects of breaking wave on surface oil spilled at Sea, SINTEF Technical Report, Trondheim, Norway, 75 pp., 2009b.
 - Resby, J. L. M. and Wang, U. M.: Update of weathering properties of Oseberg Blend, SINTEF Technical Report, Trondheim, Norway, 7 pp., 2004.
- 090 Ryerson, T. B., Camilli, R., Kessler, J. D., Kujawinski, E. B., Reddy, C. M., Valentine, D. L., Atlas, E., Blake, D. R., de Gouw, J., Meinardi, S., Parrish, D. D., Peischl, J., Seewald, J. S., and Warneke, C.: Chemical data quantify Deepwater Horizon hydrocarbon flow rate and environmental distribution, in: Proc. Natl. Acad. Sci., 109, 20246–20253, https://doi.org/10.1073/pnas.1110564109, 2012.
- Seinfeld, J. H. and Pandis, S. N. (2nd Eds.): Atmospheric chemistry and physics: from air pollution to climate change, John Wiley & Sons, New York, USA, 2006.
- Simpson, W., King, M., Beine, H., Honrath, R., and Zhou, X.: Radiation-transfer modeling of snow-pack photochemical processes during ALERT 2000, Atmos. Environ., 36, 2663–2670, https://doi.org/10.1016/S1352-2310(02)00124-3, 2002.
- Singsaas, I., Leirvik, F., Daling, P. S., Guénette, C., and Sørheim, K. R.: Fate and behaviour of weathered
 oil drifting into sea ice, using a novel wave and current flume, Mar. Pollut. Bull., 159, 111485, https://doi.org/10.1016/j.marpolbul.2020.111485, 2020.

Socolofsky, S. A., Bhaumik, T., and Seol, D.-G.: Double-plume integral models for near-field mixing in multiphase plumes, J. Hydraul. Eng., 134, 772–783, https://doi.org/10.1061/(ASCE)0733-9429(2008)134:6(772), 2008.

- 105 Sørstrøm, S. E., Brandvik, P. J., Buist, I., Daling, P. S., Dickins, D., Faksness, L. G., Potter, S., Fritt-Rasmussen, J., and Singsaas, I.: Joint industry program on oil spill contingency for Arctic and icecovered waters: Summary Report, SINTEF, Trondheim, Norway, A14181, 40 pp., 2010.
 - Spaulding, M. L., Howlett, E., Anderson, E., and Jayko, K.: OILMAP: A Global Approach to Spill Modeling, in: Proceedings of the Fifteenth Arctic Marine Oil Spill Program Technical Seminar, Environment Canada, Edmonton, Canada, 9–11 June 1992, 15–21, 1992.
- Stamnes, K., Tsay, S.-C., Wiscombe, W., and Jayaweera, K.: Numerically stable algorithm for discreteordinate-method radiative transfer in multiple scattering and emitting layered media, Appl.Optics, 27, 2502–2509, https://doi.org/10.1364/AO.27.002502, 1988.

110

- Sullivan, P. P., and McWilliams, J.C.: Dynamics of winds and currents coupled to surface waves, Annu. Rev. Fluid Mech. 42, 19–42, https://doi.org/10.1146/annurev-fluid-121108-145541, 2010
- Tang, L., Masutani, and S. M.: Laminar to turbulent flow liquid-liquid jet instability and breakup, In: Proceedings of the Thirteenth International Offshore and Polar Engineering Conference, Honolulu, Hawaii, USA, 25–30 May, 317–324, 2003.
- Thoman, R. L., Richter-Menge, J., and Druckenmiller, M. L.: Arctic Report Card 2020, Eds, 1–141 https://doi.org/10.25923/mn5p-t549, 2020.
 - Timco, G. W. and Frederking, R. M. W.: A review of sea ice density, Cold. Reg. Sci. Technol., 24, 1–6, https://doi.org/10.1016/0165-232X(95)00007-X, 1996.
 - Tkalich, P. and Chan, E. S.: Vertical mixing of oil droplets by breaking waves, Mar. Pollut. Bull., 44, 1219–1229, https://doi.org/10.1016/S0025-326X(02)00178-9, 2002.
- 125 Tschudi, M., Meier, W. N., and Stewart, J. S.: 2019. Quicklook Arctic Weekly EASE-Grid Sea Ice Age, Version 1, Boulder, CA, USA, NASA National Snow and Ice Data Center Distributed Active Archive Center, https://doi.org/10.5067/2XXGZY3DUGNQ, 2019.

Uttieri, M., Nihongi, A., Hinow, P., Motschman, J., Jiang, H., Alcaraz, M., and Strickler, J. R.: Copepod manipulation of oil droplet size distribution,Sci. Rep., 9, 1–8, https://doi.org/10.1038/s41598-018-37020-9, 2019.

- Venkatesh, S., El-Tahan, H., Comfort, G., and Abdelnour, R.: Modelling the behaviour of oil spills in iceinfested waters, Atmos.-Ocean, 28, 303–329, https://doi.org/10.1080/07055900.1990.9649380, 1990.
- Wang, S. D., Shen, Y. M., and Zheng, Y. H.: Two-dimensional numerical simulation for transport and 135 fate of oil spill in seas, Ocean Eng., 32, 1556–1571, https://doi.org/10.1016/j.oceaneng.2004.12.010, 2005.
 - Wang, B., Socolofsky, S. A., Lai, C. C. K., Adams, E. E., and Boufadel, M. C.: Behaviour and dynamics of bubble breakup in gas pipeline leaks and accidental subsea oil well blowouts, Mar. Pollut. Bull., 131, 72–86, https://doi.org/10.1016/j.marpolbul.2018. 03.053, 2018.
- 140 Warren, S. G.: Impurities in snow: effects on albedo and snow melt (Review), Ann. Glaciol., 5, 177–179, https://doi.org/10.3189/1984AoG5-1-177-179, 1984.
 - Warren, S. G.: Optical properties of ice and snow, Phil. Trans. R. Soc. A., 377, 20180161, http://dx.doi.org/10.1098/rsta.2018.0161, 2019.

Warren, S. G. and Wiscombe, W. J.: A model for the spectral albedo of snow. II: snow containing

- 145 atmospheric aerosols, J. Atmos. Sci., 37, 2734–2745, https://doi.org/10.1175/1520-0469(1980)037<2734:AMFTSA>2.0.CO;2, 1980.
 - Warren, S. G. and Clarke, A. D.: Soot in the atmosphere and snow surface of Antarctica, J. Geophys. Res.-Atmos., 95, 1811–1816, https://doi.org/10.1029/JD095iD02p01811, 1990.
- Warren, S. G., Brandt, R. E., and Grenfell, T. C.: Visible and near-ultraviolet absorption spectrum of ice
 from transmission of solar radiation into snow, Appl. Opt., 45, 5320, https://doi.org/10.1364/AO.45.005320, 2006.

Warren, S. and Brandt, R.: Optical constants of ice from the ultraviolet to the microwave: a revised compilation, J. Geophys. Res., 113, 1206–1225, https://doi.org/10.1029/2007JD009744, 2008.
Weeks, W.: On Sea Ice, University of Alaska Press, Fairbanks, Alaska, USA, 2010.

155 Wilkinson, J. P., Boyd, T., Hagen, B., Maksym, T., Pegau, S., Roman, C., Singh, H., and Zabilansky, L.: Detection and quantification of oil under sea ice: The view from below, Cold Reg. Sci. Technol., 109, 9-17, https://doi.org/10.1016/j.coldregions.2014.08.004, 2015.

Wilkinson, J. P., Beegle-Krause, C., Evers, K.-U., Hughes, N., Lewis, A., Reed, M., and Wadhams, P.: Oil spill response capabilities and technologies for ice-covered Arctic marine waters: a review of

160 recent developments and established practices, Ambio, 46, 423–441, https://doi.org/10.1007/s13280-017-0958-y, 2017.

Wongnapapan, P., Wattayakorn, G., and Snidvongs, A.: Petroleum hydrocarbon in seawater and some sediments of the South China Sea, Area I: Gulf of Thailand and East Coast of Peninsular Malaysia, in: Proceedings of the 1st Technical Seminar on Marine Fishery Resources Survey in the South China Sea, 105–110, 1999.

WWF-Canada Arctic Offshore Drilling Review, NEB File: OF-EP-Gen-AODR 01 Additional ResponsetoCFI#1and#2:https://awsassets.wwf.ca/downloads/wwf_canada___submission_to_neb_review_april_2011.pdf,last access: 1 March 2021, 2011.

165

- 170 Xiao, S., and Yang, D.: Effect of oil plumes on upper-ocean radiative transfer A numerical study, Ocean. Model., 145, 101522, https://doi.org/10.1016/j.ocemod.2019.101522, 2020.
 - Yang, D., Chamecki, M., and Meneveau, C.: Inhibition of oil plume dilution in Langmuir ocean circulation, Geophys. Res. Lett., 41, 1632–1638, https://doi.org/10.1002/2014GL059284, 2014.
- Zatko, M., Grenfell, T., Alexander, B., Doherty, S., Thomas, J., and Yang, X.: The influence of snow grain size and impurities on the vertical profiles of actinic flux and associated NOx emissions on the Antarctic and Greenland ice sheets, Atmos. Chem. and Phys., 13, 3547–3567, https://doi:10.5194/acp-13-3547-2013, 2013.
 - Zdanowicz, C. M., Zielinski, G. A., andWake, C. P.: Characteristics of modern atmospheric dust deposition in snow on the Penny Ice Cap, Baffin Island, Arctic Canada, Tellus B, 50, 506–520, https://doi.org/10.3402/tellusb.v50i5.16234, 1998.
 - Zeinstra-Helfrich, M., Koops, W., Dijkstra, K., and Murk, A. J.: Quantification of the effect of oil layer thickness on entrainment of surface oil, Mar. Pollut. Bull., 96, 401–409, https://doi.org/10.1016/j.marpolbul.2015.04.015, 2015a.

Zeinstra-Helfrich, M., Koops, W., and Murk, A. J.: The NET effect of dispersants — a critical review of

- 185 testing and modelling of surface oil dispersion, Mar. Pollut. Bull., 100, 102–111, https://doi.org/10.1016/j.marpolbul.2015.09.022, 2015b.
 - Zeinstra-Helfrich, M., Koops, W., and Murk, A. J.: How oil properties and layer thickness determine the entrainment of spilled surface oil, Mar. Pollut. Bull., 110, 184–193, https://doi.org/10.1016/j.marpolbul.2016.06.063, 2016.
- 190 Zender, C. S., Gallet, J.-C., Dominé, F., Picard, G., and Flanner, M. G.: Albedo reduction by black carbon in snow: Measurements and implications, Geophys. Res. Lett., http://dust.ess.uci.edu/ppr/ppr_ZGD09.pdf, 2009
 - Zhao, L., Boufadel, M. C., Socolofsky, S. A., Adams, E., King, T., and Lee, K.: Evolution of droplets in subsea oil and gas blowouts: Development and validation of the numerical model VDROP-J, Mar. Pollut. Bull., 83, 58–69, https://doi.org/10.1016/j.marpolbul.2014.04.020, 2014a.
- Zhao, L., Torlapati, J., Boufadel, M. C., King, T., Robinson, B., and Lee, K.: VDROP: A comprehensive model for droplet formation of oils and gases in liquids - Incorporation of the interfacial tension and droplet viscosity, Chem. Eng. J., 253, 93–106, https://doi.org/10.1016/j.cej.2014.04.082, 2014b.

195

- 200 Zhao, L., Shaffer, F., Robinson, B., King, T., D'Ambrose, C., Pan, Z., Gao, F., Miller, R. S., Conmy, R. N., and Boufadel, M. C.: Underwater oil jet: hydrodynamics and droplet size distribution, Chem. Eng. J., 299, 292–303, https://doi.org/10.1016/j.cej.2016. 04.061, 2016.
 - Zhao, L., Boufadel, M. C., King, T., Robinson, B., Gao, F., Socolofsky, S.A., and Lee, K.: Droplet and bubble formation of combined oil and gas releases in subsea blowouts, Mar. Pollut. Bull., 120, 203– 216, https://doi.org/10.1016/j.marpolbul.2017.05.010, 2017a.
 - Zhao, L., Gao, F., Boufadel, M. C., King, T., Robinson, B., Lee, K., and Conmy, R.: Oil jet with dispersant: macro-scale hydrodynamics and tip streaming, AICHE J., 63, 5222–5234, https://doi.org/10.1002/aic.15864, 2017b.
- Zheng, L., Yapa, P.D., and Chen, F.: A model for simulating deepwater oil and gas blowouts part I: 210 theory and model formulation, J. Hydraul. Res., 41, 339–351. https://doi.org/10.1080/00221680309499980, 2003.
- Zhou, L., Stroeve, J., Xu, S., Petty, A., Tilling, R., Winstrup, M., Rostosky, P., Lawrence, I. R., Liston, G. E., Ridout, A., Tsamados, M., and Nandan, V.: Inter-comparison of snow depth over Arctic sea ice from reanalysis reconstructions and satellite retrieval, The Cryosphere, 15, 345–367, https://doi.org/10.5194/tc-15-345-2021, 2021.