Impact of freshwater runoff from the southwest Greenland Ice Sheet on fjord productivity since the late 19th century

Mimmi Oksman¹, Anna Bang Kvorning¹, Signe Hillerup Larsen¹, Kristian Kjellerup Kjeldsen¹, Kenneth David Mankoff¹, William Colgan¹, Thorbjørn Joest Andersen², Niels Nørgaard-Pedersen³, Marit-Solveig Seidenkrantz⁴, Naja Mikkelsen¹, Sofia Ribeiro¹

¹Department of Glaciology and Climate, Geological Survey of Denmark and Greenland, Øster Voldgade 10, 1350 Copenhagen, Denmark.

²Department of Geosciences and Natural Resource Management, University of Copenhagen, Øster Voldgade 10, 1350 Copenhagen, Denmark.

³Department of Marine Geology, Geological Survey of Denmark and Greenland, C.F. Møllers Allé 8, 8000 Aarhus C, Denmark.

⁴Paleoceanography and Paleoclimate Group, Arctic Research Centre, and iClimate centre, Department of Geoscience, Aarhus University, Høegh-Guldbergs Gade 2, 8000 Aarhus C, Denmark.

Correspondence to: Mimmi Oksman (mio@geus.dk)

Abstract. Climate warming and the resulting acceleration of freshwater discharge from the Greenland Ice Sheet are impacting Arctic marine coastal ecosystems, with implications for their biological productivity. To accurately project the future of coastal ecosystems, and place recent trends into perspective, paleo-records are essential. Here, we showpresent runoff estimates from the late 19th century to present-day runoff estimates for a large sub-Arctic fjord system (Nuup Kangerlua, southwest Greenland) influenced by both marine- and land-terminating glaciers. We followed a multiproxy approach to reconstruct spatial and temporal trends in primary production from four sediment core records, including diatom fluxes and assemblage composition changes, biogeochemical and sedimentological proxies (total organic carbon,

- nitrogen, C/N-ratio, biogenic silica, δ¹³C, δ¹⁵N, and grain size distribution). We show that an abrupt increase in freshwater runoff in the mid-1990²s iwas reflected by a 3-fold increase in biogenic silica fluxes in the glacier-proximal area of the fjord. In addition to increased productivity, freshwater runoff modulates the diatom assemblages and drives the dynamics and magnitude of the diatom spring bloom. Our records indicate that marine productivity is higher today than it has been at any point since the late 19th century and suggest that increased mass loss of the Greenland Ice Sheet mayis-likely to continue promoting high productivity levels at sites proximal to marine-terminating glaciers. We highlight the importance
 - of paleo-records in offering a unique temporal perspective on ice-ocean-ecosystem responses to climate forcing beyond existing remote sensing or monitoring time-series.

1 Introduction

35

40

5

10

15

Arctic coastal ecosystems are experiencing profound and rapid changes that are predicted to intensify in the near future due to ongoing climate changes (Lannuzel et al., 2020). Arctic ecosystems are impacted by sea-ice thinning and retreat (Arrigo and van Dijken, 2015; Stroeve and Notz, 2018) and by increasing freshwater discharge from melting glaciers and terrestrial runoff (Hopwood et al., 2020). During the last two decades, freshwater runoff from the Greenland Ice Sheet (GrIS) has rapidly increased and doubled in volume, amounting to ca. 1 000 Gt year⁻¹ today (Bamber et al., 2018; Mankoff et al., 2020); and, under a warming climate, freshwater discharge will continue is predicted to increase in the future (Fettweis et al., 2013). Glacial freshwater discharge changes the physical and chemical properties of the water column

and nutrient (e.g., nitrate, phosphate and silica) dynamics, which in turn impacts primary production, the process by which primary producers convert solar energy into organic compounds, the length of the productive season, species composition and the distribution of biological communities (e.g., Hawkings et al., 2015; Hopwood et al., 2020). Recent studies have observed increased phytoplankton blooms and primary production rates as well as a more prolonged productive season in the high Arctic (Renaut et al., 2018, Lewis et al., 2020). However, these studies cover only the last few decades whereas

45

65

Greenland fjord ecosystems, at the interface between the ocean and the GrIS, maintain a high biological productivity which is essential to sustain important socio-economic activities such as fisheries and harvesting of indigenous resources (mainly marine mammals). Higher-trophic-level organisms, such as mammals and birds, are typically not directly

long-term records, providing the natural baseline for the recent changes, are scarce.

- 50 impacted by the freshwater increase yet changes in their food sources have cascading effects with severe implications to these organisms (e.g., Berthelsen, 2014). Long-term records of primary production changes, as well as knowledge on its spatial variability, are needed to estimate the fate of these unique ecosystems under a changing climate. However, our current knowledge of freshwater impacts on marine productivity is largely based on modern observations (Meire et al., 2017; Hopwood et al., 2018, 2020) and on monitoring data spanning only a few decades (Greenland Ecosystem
- 55 Monitoring program; GEM). These studies have shown that the influence of glaciers on marine productivity depends on multiple factors such as glacier type (i.e., marine- vs. land-terminating), fjord geometry, and on the resources (nutrients and light availability) that are limiting phytoplankton growth (Hopwood et al., 2020). In fjords with marine-terminating glaciers, sub-glacial discharge plays a key role in sustaining high productivity levels, as the upwelling meltwater plume brings crucial nutrients from the ambient deep water to the surface (Meire et al., 2017; Hopwood et al., 2018; Kanna et

60 al., 2018). However, in the future, this nutrient source can shut down as marine-terminating glaciers retreat onto land or if glacier grounding line depths changes significantly.

The GrIS is presently losing mass (between 10s and ~500 Gt yr⁻¹) with a long-term trend of increasing mass loss;, and this shrinking is most pronounced on the western side of the ice sheet (Mankoff et al., 2021). Nuup Kangerlua (Godthåbsfjord), situated in southwest Greenland, (Fig. 1) is one of the largest fjord systems in the world. The fjord is ca. 190 km long, 4 – 8 km wide, up to 625 m deep, and covers an area of ca. 2 013 km². The fjord has several sills; the main one is located at the mouth of the fjord at ca. 170 m water depth, with smaller sills in the side branches of the fjord

- (Mortensen et al., 2011). The fjord hydrography is influenced by ocean forcing (governed largely by the West Greenland Current, WGC) and both tidal and wind mixing (Juul-Pedersen et al., 2015), and the combination of atmosphere-ocean-ice forcing makes the fjord particularly sensitive to climatic changes. The main fjord branch, Nuup Kangerlua, is heavily
 influenced by glacial discharge and <u>actsfunctions</u> as a gateway for iceberg outflow, while the three side branches Kapisillit Kangerluat, Qoornup Sullua and Uummannap Sullua, do not house large outlet glaciers and are not directly influenced
- by glacial discharge (Fig. 1). The average yearly freshwater dischargerunoff between 2015 and 2019 was ca. 20.9 Gt year⁻¹ and solid ice discharge ca. 8.1 Gt year⁻¹ (Mankoff et al., 2020). The fjord system is heavily impacted by meltwater, especially during summer, which lowers the surface salinities (Meire et al., 2017). However, icebergs also have a
- 75 significant role in the freshwater flux as they can produce up to <u>ca. one quarter22.%</u> of the total meltwater (Van As et al., 2014). The fjord receives freshwater and sediment discharge from three marine-terminating glaciers (Kangiata Nunaata Sermia, KNS; Akullersuup Sermia, AS; and Narsap Sermia, NS) and three land-terminating glaciers (Qamanaarsuup Sermia; QS, Kangilinnguata Sermia; KS and Saqqap Sermia; SS) of which SS drains into Lake Tasersuaq, while the others drain into the fjord (Fig. 1). <u>The turbid meltwater reduces light penetration in the inner fjord close to the outlet</u>

point of Lake Tasersuaq creating a large spatial contrast across the fjord (Fig. 1; Meire et al., 2017). The inner part of the fjord is annually covered by sea ice from November to May, whereas the outer part stays ice free for most of the year. A large interannual environmental gradient exists inside the fjord as the inner part receives ice, freshwater, and sediments especially in the spring and late summer, creating salinity, temperature, and productivity gradients (Mortensen et al., 2013; Krawczyk et al. 2015a, 2018; Meire et al., 2016a, 2017). The turbid meltwater reduces light penetration in the inner fjord close to the outlet point of Lake Tasersuaq creating a large spatial contrast across the fjord (Fig. 1; Meire et al., 2017). The inner part of the fjord is annually covered by sea ice from November to May, whereas the outer part stays ice free for most of the year.



90 Figure 1. (a) Locations of studied sediment cores (with site numbers) SA13-ST6-36R (site 6), SA12-ST8-1 (site 8), DANA08-ST17 (site 17), and DANA08-ST20 (site 20), and the main freshwater sources feeding into the fjord; Kangiata Nunaata Sermia (KNS),

Akullersuup Sermia (AS), Narssap Sermia (NS), Kangilinnguata Sermia (KS), Qamanaarsuup Sermia (QS), Saqqap Sermia (SS) and Lake Tasersuaq (LT) in the Nuup Kangerlua fjord system. 1) Main fjord branch Nuup Kangerlua, 2) Qoornup Sullua, 3) Kapisillit Kangerluat, and 4) Uummannap Sullua. Yellow circle indicates the oceanographic station for sea-surface temperature measurements (Ribergaard et al., 2014). Redlines mark the division into inner, outer, and southern fjord used for the freshwater runoff estimates. (b) True color Sentinel-2 imagery of the fjord representing summer season conditions, from August 9th, 2018, obtained from http://carthexplorer.usgs.gov (last access: 30 Nov 2021).

95 '

100

105

110

Present-day total annual primary production in Nuup Kangerlua is generally high (84.6 - 139.1 g C m⁻² yr⁻¹) and it is maintained by diverse pelagic and benthic protist communities (Juul-Pedersen et al., 2015; Krawczyk et al., 2015a, Krawczyk et al., 2015b, 2018). Two phytoplankton blooms, of a similar magnitude, occur each year with the first bloom taking place during the spring in April/May and the second during the summer in July/August (Juul-Pedersen et al., 2015). The spring bloom plays a major role in Nuup Kangerlua accounting for 50-60 % of the annual production, and it is highly relevant for fueling the marine food web by starting the annual productive season (Hodal et al., 2012; Meire et al., 2016a). During recent decades, phytoplankton spring blooms in the Arctic have become markedly longer and they occur earlier due to sea-ice retreat (Kahru et al., 2011). However, in Arctic fjords, increased freshwater discharge has been suggested to be one of the main factors enhancing the phytoplankton spring bloom (Meire et al., 2016a). In Nuup Kangerlua, the timing, intensity, and location of the modern spring bloom is also controlled by a combination of sea-ice cover, upwelling of nutrient-rich waters, and wind forcing (Juul-Pedersen et al., 2015; Meire et al., 2015, 2016a). The spring bloom is typically dominated by pennate diatoms (that form silicified frustules) and thus the input of dissolved silica (DSi) from the GrIS is one of the critical components driving the primary production. Meltwater runoff is suggested to be tightly linked to DSi input to the fjord, withand surface runoff contributinges- 79 %, of the total silica input to Nuup Kangerlua, whereas subglacial discharge accounts only for 12 %, and solid ice discharge 9 % of the total silica input to Nuup Kangerlua (Meire et al., 2016b). Together, these three mechanisms sustain a high silica concentration and primary

production 30 – 80 km down fjord from downstream the marine terminating glaciers, with the highest production at ca.
 ~50 km off the glacier termini (Meire et al., 2016b, 2017).
 Diatoms are one of the principal contributors to marine primary production with in polar and sub-polar regions. The

silicified frustules of diatoms sink to the seabed and, over time, become archived in sedimentary records (Hasle and Syvertsen, 1996). While diatoms are widely distributed and abundant as a group, many species have narrow ecological preferences in terms of hydrographic conditions (e.g., salinity, temperature, and light availability) and this featurewhich makes them an excellent tool for reconstructing past environmental and productivity changes (Hasle and Syvertsen, 1996; von Quillfeldt, 2000, 2004; Krawczyk et al., 2018; Oksman et al., 2019; Weckström et al., 2020). In Nuup Kangerlua, diatom assemblages have been shown to respond mainly to physical conditions in the water column (temperature and salinity) rather than to biogeochemical properties (e.g., nutrients), and especially glacial discharge <u>appears to favors</u>

125 diatoms mostly when compared tomore than other protists groups (Krawczyk et al., 2015a, 2018). Recent studies on primary production around Greenland have focused either on process-level understanding, characterizing dynamics during the productive season, or on the annual cycle of species succession (e.g., Arendt et al., 2016; Meire et al., 2016a; Krawczyk et al. 2015a, 2015b, 2018; Holding et al., 2019; Luostarinen et al., 2020). Although some marine and coastal studies have focused on longer timescales (e.g., Limoges et al., 2020; Saini et al., 2020), records of decadal to centennial fjord productivity and particularly its response to increased freshwater discharge are poorly

4

Formatted: Superscript

Field Code Changed

documented. As underlined by the IPCC-SROCC 2019 report, long-term variations in fjord primary productivity remain poorly constrained, and thus we lack a solid knowledge of past productivity responses to cryosphere changes that can be used as a base for future projections (Bindoff et al., 2019).

- The aim of this study is to assess the impacts of freshwater runoff from the GrIS on primary productivity in the Nuup
 Kangerlua fjord system since the late 19th century (from ca. 1890 to present_day). We chose Nuup Kangerlua as a suitable glaciated fjord system due to its high productivity, magnitude of modern glacial discharge, and because the fjord has been monitored since 2008 and is a key site for the GEM. We extracted basin-wide freshwater runoff estimates between 1900 and 2019 from model hindcasts and reconstructed trends in primary production from four dated sediment records following a multiproxy approach including diatom fluxes and assemblage composition, biogenic silica (BSi) content, composition and origin of organic matter (TOC, C/N-ratio, carbon and nitrogen isotopes; δ¹³C and δ¹⁵N), and grain-size
- analysis. Alongside with the freshwater runoff estimates, we interpret spatial and temporal trends in our proxy data against historical records of regional air and sea_surface temperature, and changes in the front position of glaciers feeding into the fjord. Our combined records of freshwater runoff and primary production span more than a century and provide a novel insight into historical and recent changes and their spatial variability.

145 2 Material and methods

2.1 Sediment core material

A total of four sediment core records were included in this study, retrieved during three campaigns that took place between 2008 and 2013 (Fig. 1). During the campaigns, coring sites were carefully selected based on shallow seismic records and multibeam bathymetry data from previous research cruises (Harff et al., 2016; Jensen and Rasch, 2009; Mikkelsen et al., 150 2012, Seidenkrantz et al., 2014). The studied sediment cores were chosen based on their location and temporal resolution. Core SA12-ST8-1 (503 m water depth, 153 cm long, 64°16,416'N 51°19,777'W), was retrieved from the side branch Qoornup Sullua in May 2012 (Mikkelsen et al., 2012) and core SA13-ST6-36R (398 m water-depth, 96 cm-long, 64°29,076'N 50°42,553'W) was retrieved from the side branch Kapisillit Kangerluat in 2013 (Seidenkrantz et al. 2013). Both cores were collected on board the Greenlandic research vessel (RV) Sanna using a Rumorhr-lot-corer. Cores . 155 DANA08-17 (575m water-depth, 40 cm-long, 64°37,277'N 50°56,938'W) and DANA08-20 (496 m water-depth, 45 cmlong, 64°41,927'N 50° 21,566'W) were recovered with a multicorer from the main branch of Nuup Kangerlua in 2008, on board the Danish RV Dana. This fjord branch is heavily influenced by glacial discharge, as ice floes and icebergs travel through this channel to exit the fjord (Fig. 1). All sediment cores were stored in cold (4 - 7 °C) and dark conditions until opened. Sediment core sub-samples were frozen at -20 °C and freeze-dried before further processing. Sediment core 160 records are hereafter referred to by their site numbers as site 8 (SA12-ST8-1), 6 (SA13-ST6-36R), 17 (DANA08-17), and 20 (DANA08-20) and presented in the figures following an outer-to-inner fjord gradient (8, 6, 17, and 20) (Fig. 1).

2.2 Geochronology

165

The activity of ²¹⁰Pb, ²²⁶Ra and ¹³⁷Cs was analyzed from all studied cores by gamma-spectrometry at the Gamma Dating Center at the University of Copenhagen. From selected depth intervals, ca. 2 gram of freeze-dried sediment was measured per sample using Canberra ultralow-background Ge-detectors. ²¹⁰Pb was measured via its gamma-peak at 46.5 keV, ²²⁶Ra via the granddaughter ²¹⁴Pb (peaks at 295 and 352 keV) and ¹³⁷Cs via its peak at 661 keV. Chronologies for each site were obtained using a modified constant rate of supply (CRS) model (Andersen, 2017), where the activity below the lowermost sample was calculated based on a regression of activity versus accumulated mass depth for the upper part of the core. The obtained chronologies were validated by ¹³⁷Cs data.

170 2.3 Grain size analysis

175

The grain-size distribution of the sediment records was analyzed on 28, 19, 20, and 23 samples from sites 8, 6, 17, and 20 (respectively) using two complimentary methods: by-wet-sieving and laser diffraction. For wet-sieving, a minimum of 2 grams of freeze-dried sediment per sample, in orderwas used to obtain the percentages of clay and silt grains (<63 μ m), fine sand (63 – 150 μ m), and coarse sand (>150 μ m). Further analysis, using The laser diffraction analyses with a Malvern Mastersizer E/2000, wereas performed on a Malvern Mastersizer E/2000, done to obtain the distribution of clay (<2 μ m) and silt (2 – 63 μ m)-grains. Before laser diffraction measurements, samples of ca. 0.2 g of sediment were treated with a sodium pyrophosphate decahydrate-solution and ultrasonicated once for 2 minutes to separate aggregates.

2.4 Biogeochemical analysis

The total organic carbon (TOC) content of the sediments was measured at the Geological Survey of Denmark and Greenland using a LECO CS-200 instrument on sub-samples (1 g of freezedried sedimentweight). TOC was measured oin a total of 72 samples: 19 from site 8, 10 from site 6, 20 from site 17, and 23 from site 20. The inorganic carbon fraction was removed by treating the samples with HCl before measuring their TOC content. Sources of the organic material (terrestrial versus marine) were further investigated from 13 (site 8) and 10 (site 6) samples by measuring the isotopic composition of bulk organic carbon (δ¹³C) and nitrogen (δ¹⁵N). The isotopic composition analyses were done at the IRMS

- lab, Department of Geosciences and Natural Resources Management, University of Copenhagen. Ca. 20 30 mg per sample were homogenized before their carbon and nitrogen content and isotopic ratios were measured by Dumas combustion (1700 °C) on an elemental analyzer (CE 1110, Thermo Electron, Milan, Italy or Flash 2000, Thermo Scientific, Bremen, Germany). The EA iwas coupled in continuous flow mode to a Finnigan MAT Delta PLUS or Thermo Delta V Advantage isotope ratio mass spectrometer (Thermo Scientific, Bremen, Germany).
- 190 Biogenic silica was measured at the Geological Survey of Denmark and Greenland according to the 5h Na₂CO₃ extraction method (DeMaster, 1991) ion a total of 100 samples is 40, 17, 20, and 23 samples from sites 8, 6, 17, and 20, respectively. Samples of 30 ± 1 mg of freeze-dried sediment were leached using 40 ml of 1 % sodium carbonate (Na₂CO₃) in an 85 °C water bath for 5 hours. Subsamples (1 ml) were taken after 3, 4, and 5 hours of extraction and measured with a spectrophotometer Jenway 7305 using the blue ammonium-molybdate method (Mullin and Riley, 1955). Final biogenic
- 195 silica concentrations were calculated <u>and corrected for mineral dissolution</u> using the intercept of a linear regression line <u>through 3-, 4-, and 5-hour</u> sub-<u>samples</u> under the assumption that all biogenic silica dissolved during the first two hours, whereas minerogenic silica dissolves slowly at a constant rate throughout the extraction (Barão et al. 2015).

2.5 Diatoms

200

Diatom microscopy slides were prepared according to Battarbee et al., (2001) at the Geological Survey of Denmark and Greenland from 48, 21, and 20 samples from sites 8, 6, and 17, respectively. Ca. 0,1 grams of sediment per sample was treated with 10 % hydrochloric acid (HCl) and 30 % hydrogen peroxide (H₂O₂) in <u>an</u> 80 °C water bath for 4 hours to remove carbonates and organic material. Afterwards, samples were rinsed with distilled water and spiked with 0.1 ml of

210

205

a microsphere solution of 6.13×10^6 spheres/ml. Slides were then mounted using Naphrax® with a refractive index of 1.73. Diatom slides were analyzed with an Olympus BX51_light microscope using phase contrast optics at 1000X magnification, and a minimum of 300 valves were counted (excluding *Chaetoceros*). Diatom concentrations (valves g⁻¹ dry sediment) were calculated using the sum of microspheres counted per total diatom sum and the number of microspheres added in the diatom sample. The non-linear ordination method of correspondence analysis (CA) was utilized using the program PAST version 4.02 (Hammer et al., 2001) to detect ecological shifts and the major patterns of variation in the diatom assemblages. CA was run on relative abundance data for all taxa representing >1 %, and the resulting sample eigenvalues give a measure of the variance accounted for by the corresponding eigenvectors. To test correlation between CA axis scores, diatom taxa, grain size, geochemical parameters, and environmental variables (runoff, air and sea surface temperatures) a Pearson product-moment correlation coefficient and the associated level of statistical significance (p < 0.05) were used on the 5-year running mean average for each time series.

2.6 Freshwater runoff 1900-2019

- 215 We used regional climate model data to obtain runoff estimates for the period 1900-2019. For the period 1979-2019, we used the Greenland liquid water runoff dataset (Mankoff et al., 2020). This dataset is based on MAR v3.11, which is driven by the most recent reanalysis data ERA5 (Delahasse et al., 2020). For the period prior to 1979, we used MAR v3.5 (Fettweis et al., 2017) that is based on 20CRv2c reanalysis (Compo et al., 2011). The runoff data from the Greenland liquid water discharge dataset is considered to be more accurate than the MAR v3.5 dataset and thus, the latter data series
- 220 was calibrated prior to merging of the two datasets. Modelled runoff from both ice and tundra surfaces were extracted for the Nuup Kangerlua basin partitioned into three regions (Fig. 1); inner, outer, and southern part of the fjord in order to investigate spatial differences between the four sediment core sites.

2.7 Glacier front position

To assess temporal changes in the front positions of the outlet glaciers feeding into Nuup Kangerlua, we use historical aerial and satellite imagery. The historic imagery consists of oblique and vertical images from the Department of Data-supply and Efficiency, supplemented with Corona-imagery from the 1960s and satellite data from different Landsat missions retrieved from http://earthexplorer.usgs.gov (last access: 30 Nov 2021). All imagery was georeferenced relative to a 2 m ortho-photo mosaic (Korsgaard et al., 2016), which in this area of Greenland is based on aerial stereophotogrammetric imagery from summer 1985. The georeferencing is based on the method outlined by Bjørk et al., 2012).

Field Code Changed

3. Results

3.1 Geochronology

- The relatively high sedimentation rates in the fjord since the late 19th century allowed for obtaining fjord records with a sub-decadal resolution for all the cores (average sample resolution between 2 and 6 years).
- For site 8, the ²¹⁰Pb-based chronology indicates that the upper 55 cm of the record cover the time period between ca. 1890 and 2012 (Fig. 2), resulting in an average sample resolution of 2 years. Sedimentation rates are relatively high at site 8 and vary between 0.33 and 0.67 cm yr⁻¹, mean 0.45 ± 0.1 cm yr⁻¹ (Fig. 3).

For site 6, the topmost 21 cm of the sediment core used in this study represent the period from ca. 1890 to 2013, thus giving an average sample resolution of ca. 6 years (Fig. 2). Sedimentation rates are thus lower at site 6 than site 8 and vary between 0.13 and 0.25 cm yr⁻¹, mean 0.18 ± 0.03 cm yr⁻¹ (Fig. 3).

The cores from sites 17 and 20 cover shorter time intervals, from 1930 to 2008 and from 1956 to 2008, respectively (Fig. 2). Sedimentation rates are higher in these glacier-proximal sites and vary between 0.24 and 1.18 cm yr⁻¹ at site 17 (mean 0.57 ± 0.2 cm yr⁻¹) and between 0.46 and 2.33 cm yr⁻¹ (mean 1.10 ± 0.6 cm yr⁻¹) at site 20 (Fig. 3). The sediment cores from sites 17 and 20 were analyzed at 4-year and 2-year-sample resolutions, respectively.

245



Figure 2. Age-depth models, and ²¹⁰Pb and ¹³⁷Cs activities for the sediment cores SA12-ST8-1 (8), SA13-ST6-36R (6), DANA08-17 (17), and DANA08-20 (20).

3.2 Grain-size distribution analysis

- All the sediment cores consisted mostly of fine (<63 µm)-clay and silt grains-(Fig. 3). Thise <63 µm fraction of the sediment varied from 90.7 to 98.2 % (mean 96.6 ± 1.4 %) at site 8, from 28.2 to 93.0 % at site 6 (mean 76.7 ± 14.7 %), from 87.3 to 99.0 % (mean 94.2 ± 3.2 %) at site 17, and from 88.5 to 99.6 % (mean 96.7 ± 2.8 %) at site 20 (Fig. 3). More detailed distribution between silt (2 63 µm) and clay (<2 µm) grains is presented in Figure 3. The fraction of coarse material >63 µm is widely used as a proxy for ice rafted debris (IRD) carried by icebergs that generally melt relatively close to their source. The fine sand fraction (63 150 µm) varied between 1.2 and 3.7 % (mean 2.2 ± 0.7 %) at site 8, between 3.8 and 25.5 % (mean 7.5 ± 4.6 %) at site 6, between 0.6 and 4.8 % (mean 2.1 ± 1.1 %) at site 17, and between 0.2 and 10.8 % (mean 1.9 ± 2.4 %) at site 20, whereas the portion of coarse sand fraction (>150 µm) ranged between 0.2 and 5.8 % (mean 1.2 ± 1.0 %) at site 8, between 4.2 and 22.2 % (mean 8.4 ± 4.4 %) at site 6, between 0.5 and 10.9 % (mean 3.7 ± 2.4 %) at site 17, and between 0.1 and 8.2 % (mean 1.4 ± 1.7 %) at site 20 (Fig. 3). The variations in the grain sizes between the two methods are not in full agreement when it comes to the larger fraction. This might be caused
 - by the different sample volumes used, making the wet-sieving method more sensitive for estimating the fraction of larger grain sizes.

3.3 Biogeochemical analysis

Total organic carbon (TOC) content in sediments provides an integrated signal of production, deposition, and preservation
of organic matter over time, whereas the stable carbon and nitrogen isotopes and C/N-ratio can be used to infer the origin of organic matter (i.e., sourced from the terrestrial, marine, or sea ice environments). TOC contents were highest in the most glacial-distal area (sites 8 and 6) and substantially lower at the glacial-proximal sites (17 and 20) (Table 1 and Fig.
3). Sediment C/N-ratios were calculated from the sediment cores from sites 8 and 6 and show relatively low values, thus suggesting a predominantly marine origin (Table 1 and Fig. 3).

270

								1	
Site	TOC <u>(wt %)</u>	TN <u>(%)</u>	C/N-ratio	δ ¹³ C <u>(‰)</u>	δ ¹⁵ N <u>(‰)</u>	BSi <u>(mg/g⁻¹)</u>	BSi flux (mg cm ² yr ⁴)		Formatted Table
8	1.25 – 1.68 wt%	0.28 – 0.33 %	4.6 - 8	-23.321.4-‰	3.9 – 4.7 -‰	21.1 - 35.4-mg/g ⁻¹	7.1 – 19.2-mg cm ² yr ⁻¹		Formatted: Norwegian (Bokmål)
	(mean 1.4 ± 0.1	(mean 0.3 \pm	(mean 6.3 \pm	(mean -22.4 ±	(mean 4.3 ±	(mean 27.5 ± 3.7	(mean 12.1 ± 2.9-mg		
	₩t%)	0.02-%)	1.0)	0.6 -‰)	0.3 -‰)	mg/g ⁻¹)	em ² ·yr ⁻¹)		
6	0.72 – 1.08 wt%	0.10 - 0.14 %	6.6 - 8.2	-23.622.7-%	4.9 - 6.2-%	12.8 - 34.5 mg/g ⁻¹	2.1 – 5.8 mg cm² yr ⁻¹	ĺ	
	$(mean \ 0.7 \pm 0.08$	(mean 0.1 \pm	(mean 7.5 \pm	(mean -23.1 ±	(mean 5.9 ±	(mean 22.2 ± 6.0	$(\text{mean } 3.9 \pm 1.1 \text{-mg-cm}^2)$		
	wt%)	0.01-%)	0.4)	0.3-%)	0.3-‰)	mg/g⁻¹)	yr ⁻¹)		
17	0.09 – 0.30 wt%					5.3 - 28.9-mg/g ⁻¹	2.2 – 18.0 mg cm² yr⁻¹	ĺ	
	(mean 0.2 ± 0.06					(mean 12.6 ± 6.6	$(\text{mean } 7.5 \pm 5.1 \text{-} \text{mg - cm}^2)$		
	wt%)					mg/g⁻¹)	yr ⁻¹)		
20	0.007 - 0.07 - wt%					5.3 – 14.3-mg/g ⁻¹	$3.0 - 25.6 \text{ mg cm}^2 \text{yr}^4$	İ	
	$(mean\ 0.03\pm 0.02$					(mean 8.0 ± 2.4	$(mean 9.2 \pm 6.3 \text{-mg-cm}^2$		
	wt%)					mg/g - ¹)	yr - ¹)		
							1		

The stable nitrogen isotope composition in sediments can be used to infer nitrate (NO₃⁻) supply in the euphotic zone₂⁻ aAs phytoplankton consumes lighter nitrate the sediment becomes enriched in the heavier (δ^{15} N) nitrate indicating of increased productivity. The total nitrogen and isotopic composition of the sediments were not measured in the sediment cores from sites 17 and 20 due to the very low organic matter content (Table 1 and Fig. 3). Biogenic silica <u>concentrations wereas</u> highe<u>rst</u> in the glacial-distal sites than <u>in the</u> glacier-proximal sites (Table 1). <u>However, when accounting for differences</u> <u>in yet changes in</u> sedimentation rates<u>t</u> were <u>considered and</u>-BSi fluxes were higher in the glacial-proximal sites (Table 1 and Fig. 3).



Figure 3. Sediment grain-size distribution (wet-sieving and <u>laser diffractionMalvern</u>), sedimentation rate (cm/yr), diatom flux (valves x 10⁶ g yr⁻¹), BSi content (mg g⁻¹ dry mass); green line and BSi flux (mg cm² yr⁻¹); black line, total organic carbon (wt%), total nitrogen (%), C/N-ratio, δ^{13} C (%PDB) and δ^{15} N (%AIR) in sediment cores SA12-ST8-1 (8), SA13-ST6-36R (6), DANA08-17 (17)₂ and DANA08-20 (20).

3.4 Diatoms

Diatom assemblages were both rich and diverse at all sites expect for site 20, which was virtually barren of diatoms and is discussed in Section 4.1. At sites 8, 6, and 17, a total of 172, 138, and 93 different taxa were identified to species level and 22, 21, and 10 taxa to genus level, respectively. Total diatom concentrations (including resting spores and vegetative forms of *Chaetoceros* spp.) in the sediment samples varied from 1.2 to 12.2 x 10⁶ valves/g at site 8, from 2.3 to 35.8 x 10⁶ valves/g at site 6, and from 2.8 to 30.4 x 10⁶ valves/g at site 17 (Fig. 4). Annual diatom fluxes ranged between 0.5 and 5.1 x10⁶ valves cm⁻¹yr⁻¹ (mean 2.1 ± 1.0 x10⁶ valves cm⁻¹yr⁻¹) at site 8, 0.3 to 7.2 x10⁶ valves cm⁻¹yr⁻¹ (mean 1.6 ± 1.6 x10⁶ valves cm⁻¹yr⁻¹) at site 6 and between 0.8 and 15.8 x10⁶ valves cm⁻¹yr⁻¹ (mean 7.8 ± 4.3 x10⁶ valves cm⁻¹yr⁻¹) at site 17 (Fig. 3).

295

285

Diatom assemblage composition was relatively uniform between the three studied sites and diatom taxa (excluding *Chaetoceros* species) were generally dominated by *Detonula confervacea* (resting spore), *Thalassiosira antarctica* var. *borealis* (resting spore), *Fragilariopsis oceanica*, *F. cylindrus* and *T. nordenskioeldii* (Fig. 4). Diatoms were divided into 6 groups according to their ecological preferences; cold-water species, early spring bloomers, benthic species, sea-ice associated species and freshwater species (Table 2).

300

Table 2. Diatom groups, species in each group, ecological preferences, and references.

Group	Species	Ecology	Reference	
Cold-water species	Detonula confervacea, Thalassiosira antarctica var borealis (r.s. and veg), Bacteriosira bathyomphala, Thalassiosira anguste-lineata, Thalassiosira hyalina, and Shionodiscus trifultus	Species preferring relatively cold waters, blooming in the late summer.	Krawzcyk et al., 2015a; Oksman et al., 2019; Luostarinen et al., 2020	
Early spring bloomers	Fragilariopsis oceanica, Fragilariopsis cylindrus, Fragilariopsis reginae-jahniae, and Pauliella taenitae	First species to bloom in the spring during and immediately after sea_ice break-up and melt.	Quillfeldt 2000, 2004; Oksman et al., 2019; Weckström et al., 2020	
Sea-ice associated	Actinocyclus curvatulus, Coscinodiscus oculus-iridis, Fossula arctica, Porosira glacialis, Thalassiosira bulbosa, Rhisozolenia spp., Pleurosigma stuxbergii	Species associated with seasonal and high sea_ice concentrations or produce sea_ice <u>related</u> biomarkers (HBI III).	Oksman et al., 2019; Weckström et al., 2020 Belt et al., 2017	
Freshwater species	Eunotia spp., Cymbella spp., Cyclotella spp., Tabellaria spp., and Pinnularia spp.	Species living in freshwater environments.	Cremer, 1998; Foged, 1973	

Chaetoceros spp. are generally abundant in the sediment due to their good preservation and are likely to be overrepresented in the assemblage and thus were left out of the diatom assemblage percentage calculations discussed here | 310

315

(Fig. 4). However, to have a general ideaestimate of the concentrations of *Chaetoceros* spp. and Cchrysophyte cysts, we calculated flux rates (Fig. 4). *Chaetoceros* spp. (including vegetative forms and resting spores) were abundant in all the cores containing diatoms, making average flux rates of 0.4, 0.4, and 0.8 x 10⁶ valves cm⁻¹yr⁻¹ at sites 8, 6, and 17 respectively (Fig. 4). Additionally, Cchrysophyte cysts were counted, as this groupenus is associated with low-salinity waters during the summer/autumn bloom (Hasle and Heimdal, 1998; Krawczyk et al., 2015a) and can thus provide

For the diatom record from site 8, the first CA axis explains 36 % of the variation in the dataset (Fig. 4). The axis scores switch from mainly positive to mostly negative values around the late 1940's. At site 6, the first CA axis accounts for 49.7 % of the variation in the dataset and the CA axis scores change from negative to positive at ca. 1960's. At site 17, the first CA axis explains 29.5 % of the variation in the dataset. Scores for the first CA axis remain mainly negative until

ca. 1985 when they shift to positive values.

additional information on past salinity changes.

Formatted: Font: Not Italic



Figure 4. Total freshwater runoff (Gt/yr), Ddiatom valve concentration (10⁶ valves/g dry sediment), CA axis 1 scores (dashed lines indicates "0"), relative abundance (%) of the cold-water species, early spring bloomers, benthic species, sea-ice associated species, freshwater species, relative percentage of the most abundant <u>taxaspecies</u> in each core, *Chaetoceros* spp., and Cghrysophyte cyst fluxes (x10⁶ valves cm⁻¹yr⁻¹) in sediment cores SA12-ST8-1 (8), SA13-ST6-36R (6), and DANA08-17 (17). DANA08-20 (20) was devoid of diatom valves. Diatom species are scaled on different x-axes for better visualization of trends in the percentages of less abundant species.

325 3.5 Freshwater runoff

Total freshwater runoff estimates extracted from MAR were further dividedpartitioned into estimates representing inner fjord runoff, outer fjord runoff and southern fjord runoff (Fig. 5). Runoff estimates for the southern part of the fjord system (see Fig. 1 for division of the fjord regions) are excluded since they represent part of the fjord that is not investigated in this study. The estimate for the total freshwater runoff varies between 13.0 and 49.0 Gt year⁻¹ (mean 23.1 Gt/year⁻¹), for the inner fjord runoff between 9.1 and 41.1 Gt year⁻¹ (mean 18.0 Gt year⁻¹), and between 0.7 and 2.2 Gt year⁻¹ (mean 1.3 Gt year⁻¹) form the outer fjord (Fig 5). Glacial runoff makes up the largest proportion of the total runoff in Nuup Kangerlua, whereas the runoff from tundra and rainfall bothinto the fjord have a smaller contribution (Fig. 5).

Tundra 33 %



320



335

340

345

Figure 5. (a) Inner fjord (blue), outer fjord (yellow), and southern fjord (green) freshwater runoff estimates (Gt/yr) for 1900 – 2019.
 For definition of the different regions of the fjord, see red lines on Fig. 1. (b) Pie chart overshows the distribution of the average runoff from the ice (turquoise blue), tundra (green) and fjord rainfall (pink) between 2010 – 2019.

Correlation between tThe relationship between the different runoff estimates and historical climate records wasere tested, showing a positive correlation between the total and inner fjord runoff and the historical air temperature record (r = 0.61 and 0.63, respectively) as well as the with the Fyllas Bankesea-surface temperature (SST) record (r = 0.47 and 0.49) from Fyllas Banke oceanographic station shown in Fig. 1. In contrast, the outer fjord runoff estimates did not have a statistically significant correlation to neither air nor sea-surface temperatures. All statistically significant correlations between the studied proxies and the total, inner and outer fjord freshwater runoff, and air_ and sea_surface temperatures are presented in Table 3.

Table 3. Statistically significant correlations (p < 0.05) between studied-proxyles records (diatoms, geochemistry, grain size) and environmental variables (total, inner, and outer freshwater runoff, historical air temperatures and sea-surface temperatures 'SSTs') in sediment cores SA12-ST8-1 (8), SA13-ST6-36R (6), DANA08-17 (17) and DANA08-20 (20). Correlation coefficient (r) values are presented in brackets for each correlation. Positive correlations are in bold.

Formatted: Font: Italic

Site	Total runoff	Inner runoff	Outer runoff	Air temperature	SST
8	Chrysophyte cysts (-0.57)	Freshwater species (-0.37)	Diatom abundance (-0.44)	Early spring bloomers (-0.50)	Chrysophyte cysts (-0.39)
	Cocconeis costata (-0.44)	Cocconeis costata (-0.41)	Fragilariopsis oceanica (-0.39)	Fragilariopsis cylindrus (-0.52)	
	Clay; <2 µm (0.42)	Chrysophyte cysts (-0.56)	Clay; < 2 μm (0.38)	Chrysophyte cysts (-0.50)	
	Silt; 2-63 µm (-0.43)	Clay; <2 μm (0.40)	Silt; 2-62 µm (-0.43)		
	Fine sand; 63-150 µm (-0.44)	Silt; 2-63 µm (-0.41)	Fine sand; 63-150 µm (-0.51)		
	Fine grains; <63 µm (0.42)	Fine sand; 63-150 µm (-0.39)	Fine grains; <63 µm (0.44)		
		Fine grains; <63 µm (0.40)			
	BSi (0.57)	BSi (0.54)	TOC (0.52)	TOC (0.60)	
	BSi flux (0.61)	BSi flux (0.61)	TN (0.56)	TN (0.45)	
6	TOC (0.80)	TOC (0.80)	C/N ratio (0.40)	Benthic species (-0.39)	
	TN (0.64)	TN (0.63)	IRD; >150 µm (-0.49)	Chaetoceros spp. (-0.42)	
0	C/N ratio (0.44)	C/N ratio (0.42)		IRD; >150 µm (-0.43)	
	Chaetoceros spp. (-0.47)	Benthic species (-0.38)			
	IRD; >150 µm (-0.49)	Chaetoceros spp. (-0.48)			
		IRD; >150 µm (-0.46)			
	BSi (0.67)	BSi (0.64)	BSi (0.48)	BSi flux (0.30)	Early spring bloomers (-0.50)
	BSi flux (0.50)	BSi flux (0.48)	Diatom abundance (0.56)	Early spring group (-0.73)	Cold-water species (0.54)
	Early spring bloomers (-0.56)	Early spring bloomers (-0.58)	Cold-water species (0.50)	Cold-water species (0.60)	Freshwater species (-0.45)
	Sea ice associated (0.47)	Sea ice associated (0.46)	Chrysophyte cysts (-0.53)	Freshwater species (-0.55)	
17	Cold-water species (0.63)	Cold-water species (0.64)	Th. antarctica v borealis (0.59)	Th. antarctica v borealis (0.50)	
17	Freshwater species (-0.54)	Freshwater species (-0.57)		Fragilariopsis cylindrus (-0.68)	
	Chrysophyte cysts (-0.59)	Chrysophyte cysts (-0.57)			
	Th. antarctica v borealis (0.64)	Th. antarctica v borealis (0.65)			
	Thalassiosira nordenskioeldii (-0.49)	Thalassiosira nordenskioeldii (-0.46)			
	Fragilariopsis cylindrus (-0.52)	Fragilariopsis cylindrus (-0.53)			
20	BSi (0.76)	BSi (0.73)	BSi (0.66)	Clay; <2 μm (-0.50)	
	BSi flux (0.84)	BSi flux (0.81)	BSi flux (0.71)	IRD; >63 µm (0.53)	
	Sedimentation rates (0.73)	Sedimentation rates (0.71)	Sedimentation rates (0.62)		
	Fine grains; <63 µm (-0.50)	Clay; <2 μm (-0.68)	Fine sand; 63-150 µm (0.62)		
	Clay; <2 µm (-0.69)	IRD; >63 µm (0.80)	Fine grains; <63 µm (-0.70)		
	IRD; >63 µm (0.82)		IRD; >63 µm (0.63)		

4. Discussion

B55

350

In Arctic fjords, marine primary production is a crucial ecosystem function that sustains important ecosystem services such as fisheries and indigenous livelihoods (Berthelsen, 2014; Lydersen et al., 2014; Meire et al., 2017; Laufer-Meiser et al., 2021). Here, we present sub-decadal records of temporal and spatial changes on marine primary production (diatom fluxes, BSi and TOC) and diatom species composition in Nuup Kangerlua since the late 19th century and discuss these changes in relation to annual freshwater runoff estimates, historical air- and sea-surface temperatures, and glacier front dynamics.

4.1 Records of marine primary productivityon in Nuup Kangerlua since the late 19th century

360

Our primary productivityon indicators (diatom fluxes, BSi and TOC) from Nuup Kangerlua display large spatial variability between the studied sites (Fig. 3). In addition to variations in productivity between the sites, different proxies from the same core show minor dissimilarities (Fig. 3) as they all do not reflect the same productivity signal. In general, TOC and BSi values, representing organic carbon from primary and secondary producers and amount of biologically

produced silica (Table 1), point to a relatively high productivity in Nuup Kangerlua throughout the 20th century, when
 compared to modern values from other coastal Arctic sites where TOC in the surface samples remains <1 wt% and BSi
 <10 mg g⁻¹ (e.g., Ribeiro et al., 2017; Limoges et al., 2018a; Kumar et al., 2016, Detlef et al., 2021). Similarly, high values have been observed on the northwest Greenland shelf during the Holocene when climate conditions were favorable for high primary production (Limoges et al., 2020; Saini et al., 2020; Ribeiro et al., 2021). The modern fjord hydrography in Nuup Kangerlua has been shown to maintain productivity also in the outer fjord, as biomass and nutrients are partly advected from the inner fjord by currents, wind and tidal mixing which together with the warm, nutrient-rich waters

- S70 advected from the fine fjord by currents, which and that mixing which together with the warn, nutrent-field waters carried by the WGC, can increase the seasonal productivity (Juul-Pedersen et al., 2015). Also, the year-round ice-free conditions in the outer fjord may prolong the productive season, whereas glacial calving and sea-ice cover in the inner fjord shorten the productive window. Today, spatial trends in primary production in Nuup Kangerlua are mainly controlled by sea-surface temperature and
- salinity gradients along with nutrient availability, so that the highest productivity rates have been reported from the inner fjord area (Krawczyk et al., 2015b, 2018; Meire et al., 2016b, 2017). In contrast, our sediment records with the highest organic carbon content were retrieved from the outer fjord (site 8), where average TOC and BSi levels are up to 46- and 3-times higher, respectively, than in the inner fjord area (Fig. 3). While sedimentation rates are more than 2-times higher in the inner fjord, mean BSi fluxes are still slightly higher at the outer fjord site than at the two inner fjord sites 17 and 20
 (Table 2). Such an inner-outer fjord gradient of increasing productivity can be found elsewhere along the Arctic coasts
- (Arendt et al., 2016; Kumar et al., 2016; Ribeiro et al., 2017; Limoges et al., 2018a). Whereas productivity (<u>diatom</u> <u>productivity</u>, <u>BSi and TOC</u>) at the outer fjord <u>site (8)</u> has remained relatively stable (and high) since the late 19th century, this site records frequent low-amplitude fluctuations <u>in biogenic silica content and diatom production</u> occurring at a sub-decadal scale (<u>Fig. 3</u>).
- 385 The diatom productivity (absolute concentrations and fluxes) is highest at site 17, and this part of the Nuup Kangerlua fjord is characterized by very high rates of annual primary production (up to 120 g C m² year⁻¹) according to data from the Greenland Ecosystem Monitoring Programme (Juul-Pedersen et al., 2015; Meire et al., 2015). Diatoms alone account for about 95 % of the biomass here (Arendt et al., 2011; Meire et al., 2016a). The high productivity in this part of the fjord has been suggested to be maintained by the high dissolved silica (DSi) input from surface runoff combined with 390 upwelling of nutrient-rich deep waters (Meire et al., 2016b, 2017). The high diatommarine productionvity at site 17 can be explained by its ideal location receiving freshwater discharge from both surface runoff and several marine-terminating glaciers (Fig. 1). Subglacial discharge plays a key role in sustaining high productivity in this part of the fjord. Although the most glacier-proximal site (20) has a relatively high BSi flux (mean BSi flux of 9.2 mg cm² yr⁻¹-on avg.), no diatom valves were found apart from the a few in the uppermost parteentimeters of the sediment core. In general, biogenic silica 395 contents at site 20 are significantly lower until the late 1990s, suggesting low primary production here, compared to the other sites. The major increase in BSi and appearance of diatom valve fragments in the upper part of the core indicate an increase in productivity, although remaining significantly lower than at the other sites. The absence of diatoms and low

BSi values indicate, in turn, very low productivity levels. The near absence and overall poor preservation of diatom valves in the top of thisis record may also be explained by dissolution of the diatom valves in the water column or water-sediment

interface due to adverse physicochemical conditions (Ryves et al., 2006).

400

Formatted: Superscript

4.2 Sources of sedimentary organic matter in glacier-distal sites

The stable isotope carbon-13 (δ^{13} C) and the carbon to nitrogen (C/N) ratio can be used to trace whether the organic matter in the sediment originates from marine or terrestrial sources. Autochthonous organic carbon is produced by the primary producers using the CO2 dissolved in the seawater, whereas allochthonous organic carbon sources can be of terrestrial or 405 marine (e.g., sea ice) origin (Bianchi et al., 2020). At sites 8 and 6, the organic matter in the sediment can be traced to be of marine origin, as the measured δ^{13} C values (between -20 ‰ and -22 ‰) are typical for protein-rich marine phytoplankton, whereas the terrestrial organic carbon would generally be more depleted in $\delta^{13}C$ (Meyers, 1994). In our record from sites 6 and 8, we find no evidence of freshwater runoff increasing the amount of terrestrial carbon in the sediment, as the variations in δ^{13} C values are minimal and show no increase in pace with the freshwater runoff dataset 410 (Fig. 3). Despite the C/N-ratio from site 6 having a positive correlation with freshwater runoff, the very low values (average 6.3 at site 8 and average 7.4 at site 6) further support autochthonous organic matter sources, as C/N-ratios above 20 would indicate terrestrial carbon sources (Meyers, 1994). However, in marine sediments, bound inorganic nitrogen can constitute a large portion of the total nitrogen concentration and thus the marine organic matter might be more depleted in δ^{13} C than terrestrial organic matter (Kumar et al. 2016). <u>However, we found no clay-bound nitrogen when C/N-ratios</u> 415 were tested for land derived inorganic nitrogen according to Schubert and Calvert (2001). The C/N-ratios measured in this study were not corrected for the land-derived inorganic nitrogen and thus might be underestimated. These analyses $(\delta^{13}C \text{ and } C/N \text{-ratio})$ were performed only for the glacier-distal records (8 and 6) and not for the glacier-proximal records (17 and 20) due to extremely low carbon contents in these sediments (Fig. 3). Although the sedimentary organic carbon in our records is mainly from marine sources, we found freshwater diatoms in sites 8, 6, and 17 (Fig. 4) that originate 420 from terrestrial freshwater sources e.g., lakes and ponds, and enter the fjord via runoff (Jensen, 2003). Overall, the relative abundance of freshwater diatoms is extremely low (<2 %), but it represents a clear fingerprint of terrestrial origin as these species generally have very low to zero tolerance to saline waters and are not found in marine environments (Cremer, 1998; Foged, 1973) and thus support assumption that part of the organic carbon could originate from terrestrial sources. However, freshwater species were most abundant at site 6 (Fig. 4) where the C/N data suggest that the organic matter was 425 found to originate from marine sources (Fig. 3).

4.3 Freshwater discharge positively impacts primary production in the inner fjord

Our records proxy data display a substantial increase in <u>BSi content and fluxes marine primary production</u> at the glacier-proximal sites (17 and 20) from ca. 1995 onwards, as sedimentary BSi contents increase rapidly in pace with the inner fjord freshwater runoff (Fig. 6). In modern monitoring studies the inner fjord is found to displayhave high productivity, as upwelling of the sub-glacial freshwater discharge and the subsequent meltwater plume <u>bring_lead to upwelling</u> of <u>consential</u> nutrients from the <u>nutrient-rich</u> deep-water into the surface water layer for the primary producers (Meire et al., 2016b, 2017; Hopwood et al., 2018, 2020). Also, the input of dissolved silica (DSi) and its implications for increasing diatom productivity are is-highest in theis inner part of the fjord and s(Meire et al. 2016b). The modern monitoring data from Nuup Kangerlua reaches back to 2008 (Greenland Ecosystem Monitoring), however, our long-term records suggest that the high productivity levels, prevailing at the glacier-proximal area today, initiated already in the 1990's. The strong positive correlation between marine productivity (<u>BSi flux</u>) and freshwater runoff (Table 3) shows that increased productivity in this <u>part of the fjord is tightly associated to freshwater discharge</u>. This positive correlation suggests that

in the future, increased freshwater discharge is likely to have a positive socio-economic impact, as high productivity is reflected in the higher trophic levels such as fish and marine mammals (Boye et al., 2010). For example, halibut fisheries, which are economically important for Greenland (Berthelsen, 2014), have been shown to have a positive correlation with meltwater input (Meire et al., 2017).

AlthoughDespite most of the freshwater originatesing from glacial discharge (Fig. 5), the increasing relative abundance

440

of freshwater diatoms recorded from the 1990's (Fig. 6) indicates flushing of freshwater deposits (such as lake Tasersuaq) into the fjord. Together with the BSi flux, Ecoarse sediment grains (>63 µm) at site 20 display a strong positive correlation 445 with freshwater runoff (Table 3) suggesting that freshwater input to this site originates from solid ice discharge. The coarse sediment grain size in the sediment-generally represents the IRD deposited by icebergs which typically release most of the coarse grains close to the source as they melt. The melting of icebergs can cause water column vertical mixing in a similar way as subglacial discharge, and thus provide more nutrients to the surface waters (Kanna et al., 2018). Also, sedimentation rates are increasing at sites 17 and 20 along with freshwater runoff (Table 3), suggesting that a large amount 450 of sediment is being delivered to the fjord with freshwater discharge. Sub-glacial discharge from marine-terminating

- glaciers typically causes sediment suspension plumes (Fig. 1) which by upwelling, bringsadd nutrients from nutrient-rich deep waters to the surface for the primary producers (Meire et al., 2017; Hopwood et al., 2018, 2020). In the inner fjord, Agn increase in freshwater discharge accompanied by increased sediment fluxes and upwelling, would represent a shift from clear, stratified waters to turbid, mixed waters after the 1990's. In fact, our diatom record from site 17 shows that 455 species such as Thalassiosira nordenskioeldii and Fragilariopsis cylindrus, as well as Echrysophyte cysts, that prefer
- low-salinities and stratified waters (Hasle and Heimdal, 1998; von Quillfeldt, 2004; Krawczyk et al., 2015a, 2015b), decrease after the 1990's (Fig. 4), while the number of benthic species, associated to water turbidity and to low light conditions (Glud et al., 2002; Karsten et al., 2011) increases (Fig. 4). Interestingly, site 17 shows evidence of the suspension plume typically caused by the marine terminating glaciers, whereas the record from site 20 suggests increased

460 iceberg calving.

465

In contrast to the inner fjord, the record from the glacial distalouter fjord site (site 8) shows no correlation between productivity (diatom fluxes, BSi or TOC) and freshwater runoff throughout the entire studied period, and here, freshwater runoff is limited (Fig. 5). The diatom CA axis suggest an important change in the assemblage around the 1940-s (Fig. 6), from an assemblage dominated by early-spring bloomers to an assemblage dominated by benthic species and late-summer bloomers, e.g., Thalassiosira antarctica v. borealis r.s. (Fig. 4). This suggests an increase in water turbidity, and/or more nutrient -limitationed conditions (Glud et al., 2002; Karsten et al., 2011). However, the timing of this shift in the assemblage composition is not associated with any significant changes in the freshwater runoff and may therefore be

caused by changes in e.g., wind strength and consequently mixing of the water column.

Formatted: Font: Not Italic



475

Figure 6. (a) Three3-year average of annual summer (May to, June, July, and August) air temperature measured at Nuuk airport (black line) from Ribergaard₇ (2014) and (b) June sear-surface temperatures; SSTs (0 - 40 m) from Fylla Bank (blue line) from Ribergaard₇ (2014). (c) Inner fjord freshwater runoff estimates (Gt/yr). (d) BSi flux (mg cm² yr⁻¹) at site 6 (green line), site 17 (purple line) and site 20 (orange line). (e) BSi flux (mg cm² yr⁻¹) at site 8. (f) Relative percentage of freshwater diatoms at site 6. (g) Relative percentages of freshwater diatoms at site 17. (h) Diatom CA axis 1 scores at site 6 (green line) and site 17 (purple line), dashed lines indicate "0". (i) Diatom CA axis1 scores at site 8, dashed line indicates "0". (j) Relative percentage of early spring bloom species at site 6. (k) Relative percentage of early spring bloom species at site 17. (I) Changes in KNS glacier front position (m) in a relation to observed position in 2020. Shaded area marks the period from 1995 to present.

4.4 Freshwater discharge influences spring and summer bloom dynamics

480

Based on modern observations, the annual marine primary production in the-Nuup Kangerlua fjord-is sustained by two seasonal blooms of similar magnitude occurring during the early spring and the late summer (Juul-Pedersen et al., 2015). Our long-term perspective indicates that the dynamics and magnitude of these blooms have varied according to freshwater discharge (Fig. 6). We find that freshwater runoff leads to a stronger spring bloom signal at site 6, while promoting latesummer bloomers at site 17 (Table 3 and Fig. 4 and 6), and vice versa during periods of lower runoff. In terms of species 485 responses, the diatom records show that species blooming in early spring during the ice melt (Fragilariopsis spp.) or right after ice break-up in late spring (Thalassiosira nordenskioeldii) (Cremer, 1998; Jensen, 2003; Oksman et al., 2019) dominate the assemblage at site 6 during periods of higher freshwater discharge while these species at site 17 have a negative correlation with freshwater (Table 3). Yet, the species blooming during the late summer (Detonula confervacea and Thalassiosira antarctica var. borealis r.s.) as well as Chaetoceros spp., which typically follow nutrient exhaustion 490 and represent the last stage of the productive season (Krawczyk et al., 2015a), are more common at site 17 during periods of increased freshwater runoff (Fig. 4).

Modern observations have shown that today spring and summer bloom dynamics are mainly controlled by seasonal changes in salinity and temperature as well as fjord hydrography, and tidal and wind mixing (Juul-Pedersen et al., 2015; Krawczyk et al., 2015a, 2018; Meire et al., 2016a). Other environmental factors, in particular sea ice, might better explain

495 the differences observed at the two sites, as sea ice can also influence the timing and magnitude of the spring bloom (Meire et al., 2016a). At the glacier proximal site (17), a relatively high number of sea ice species (which have a positive correlation with freshwater runoff) and low number of early spring bloomers suggest sea-ice presence continuing late into the spring and limiting the spring production season (Fig. 4). However, the substantially high percentage (up to 79 % in the 1950's) of late summer bloomers alongside with elevated BSi and diatom flux (Fig. 3) are indicative of a productive 500 summer season.

The Sspring bloom ismakes up a significant part of the annual primary production in Nuup Kangerlua, (Juul-Pedersen et al., 2015) and increased spring production in the Arctic is a typical feature of the ongoing climatic warming, which has been evident in the Arctic Ocean for the recent decades (Renaut et al., 2018). The length of the productive season has implications for CO₂ capture as the fjords can function as carbon sinks (Meire et al., 2015; Rysgaard et al., 2012). In the

future, increasing freshwater discharge is likely to extend the productive season and contribute to spatial differences indespite_the timing of the blooms occurring in different parts of the fjord system. The spring bloom is likely to occur earlier in the side branch of the fjord, whereas the magnitude of the late summer bloom increases along the main ice flow pathway.

Formatted: Subscript

22

4.5 Freshwater impact reachinges into the glacial-distal side branch

510 Until now, the impacts of freshwater runoff on primary production have mainly been studied in the main branch of the Nuup Kangerlua fjord system (Juul-Pedersen et al., 2015; Meire et al., 2016a, 2016b, 2017; Krawczyk et al., 2018), whereas productivity changes in the side branches are less known. Today, the upwelling of nutrients by subglacial processes in Nuup Kangerlua is suggested to affect primary production downstream, to a distance of 10-100 km along the pathway of the advected bloom (Meire et al., 2016a; Hopwood et al., 2018). Here, we show that freshwater runoff 515 impacts primary production (BSi flux and TOC) in the glacier-distal site at the side branch of the fjord system (site 6). Here, both BSi flux and TOC strongly correlate with the freshwater runoff, despite this site being in the sheltered side branch ca. 110 km from the Kangiata Nunaata Sermia (KNS) termini and not directly on the main flow path of glacier influence (Fig. 1). The high number of early spring bloomers at site 6 indicates an earlier seasonal ice breakup and strong inflow of freshwater into the side branch as these species thrive in cold and fresh surface waters. Diatom production iwas 520 high (up to 36 x 106 valves / g) and a flux of 7.2 valves cm⁻¹yr⁻¹) during the 1930's and the 1940's, while freshwater runoff was elevated, and sea-surface temperatures weare relatively high (Figs. 3 and 6).- The high productivity is further supported inby the diatom species composition data, as by the Rhisozolenia genusera as it reaches the highest abundances of the record around 1945 (Fig. 4). This genus has been linked to a nutrient-rich and stratified water column, conditions that are generally related to meltwater pulses (Kemp et al., 2006).

525 4.6 Long-term variability in productivity links to atmospheric forcing

Teleconnections in the North Atlantic region, such as the North Atlantic Oscillation (NAO) and the Atlantic Multidecadal Oscillation (AMO), have been shown to drive marine primary production (Martinez et al., 2009). These decadal-scale ocean-atmosphere oscillations modulate sea-surface conditions through fluctuations in temperature, precipitation, and wind, and thus, impact ecosystem responses (Martinez et al., 2009; Racault et al., 2017). In west Greenland, the locations of glacier termini are driven primarily by atmospheric temperatures reflecting NAO phases, and atmospheric forcing has been defined as the primary driver of KNS ice margin changes (Lea et al., 2014; Bjørk et al., 2018). NAO phases modulate the climate so that during a negative NAO phase West Greenland experiences warmer climate, and increased precipitation, whereas during the positive phase the opposite occurs (e.g., Hurrell, 1995; Bjørk et al., 2018).

In Nuup Kangerlua, most of the glacier termini responded to the warm atmospheric temperatures during the negative
NAO phase between the 1930's and 1970's by retreat and mass loss (Fig. 7), with KNS having the largest retreat of ca. 8 km (Fig. 6). Our record shows increased marine productivity at site 6 during this negative NAO phase (Fig. 6), and here the TOC has a strong positive correlation with regional air temperatures and freshwater runoff records. The Đdiatom record from site 6 shows freshening of surface waters as *Chaetoceros* spp. and Cehrysophyte cyst fluxes are highest, together with higher relative percentage of early-spring bloomers (Fig. 4). Together, this suggests that fjord productivity responds to decadal ocean-atmosphere oscillations. After the 1970's, the NAO phase has remained mainly positive but interestingly since the 1990's the accelerated climate warming has over-ruled the NAO modulated climate signal (Seidenkrantz et al., 2009), and both temperatures and freshwater discharge have increased substantially despite the ongoing positive NAO phase (Fig. 6). During the recent decades, asynchronous behavior of the KNS and NS has primarily been attributed to subglacial melt (Motyka et al., 2017), however, both KNS and NS have retreated (Fig. 7) resulting in increased freshwater discharge and productivity in the inner fjord sites. The accelerated warming of the recent decade

23

may lead to substantial retreat of KNS in the future (Aschwanden et al., 2019). This can have a negative impact on productivity if the grounding depth changes significantly (reducing the fertilizing effect of nutrient upwelling) or if the ice margin retreats on to land. The retreat of marine-terminating glaciers is likely to have a strong impact on productivity, due to the key role that subglacial discharge plays in sustaining high fjord productivity both on interannual (Meire et al., 2017; Hopwood et al., 2018) and multi-decadal timescales as shown here.



Figure 7. Glacier terminus changes of Saqqap Sermia (SS), Kangilinnguata Sermia (KS), Narssap Sermia (NS), Qamanaarsuup Sermia (QS), Kangiata Nunaata Sermia (KNS), Akullersuup Sermia (AS), and Lake Tasersuaq (LT) in a relation to their 2020 position.

555 5. Conclusions

560

Long-term records of marine productivity are needed to define the natural baseline against which to assess recent cryosphere and coastal ecosystem changes. More accurate baselines ultimately provide more robust projections of future changes. Here, we have combined sub-decadal records of primary productivity changes with estimates of freshwater runoff since the late 19th century to determine the impacts of freshwater on fjord productivity for Nuup Kangerlua, southern West Greenland.

- We find that in the glacier-proximal area, **BSibiogenic silica** fluxes are tightly associated with freshwater runoff, and that freshwater imposes positive impacts on marine productivity. Our record shows non-uniform responses to freshwater inside the fjord system which highlights the importance of marine-terminating glaciers to marine productivity. However, we find that the impacts of freshwater discharge are wider than previously known, as freshwater has a positive impact on the marine productivity in the glacier-distal side branches of the fiord system outside of away from the main discharge
- the marine productivity in the glacier-distal side branches of the fjord system outside of away from the main discharge flow path-flow. Further, our diatom assemblage data shows that freshwater modulates the location and magnitude of the spring and late summer blooms, and these blooms are not synchronous inside the fjord. Freshening of the inner fjord prolongs the late spring sea-ice conditions leading to more extensive spring bloom in the side branch while summer bloom become larger in the inner fjord. Overall, however, wWe conclude that productivity is higher today than at any other time
- 570 since the late 19th century. A substantial increase in productivity initiated in the 1990-s in the inner fjord is coincident with localdue to the acceleration in the elimate warming and the following cryosphere changes (retreat ofin the glacier terminusfront line and sub-glacial freshwater input to the fjord) that are associated with acceleration in climate warming. Our diatom assemblage data shows that freshwater modulates the location and magnitude of the spring and late summer blooms, and that these blooms are not synchronous inside the fjord. Freshening of the inner fjord prolongs the late spring sea ice conditions leading to more extensive spring bloom in the side branch while summer blooms becomes larger in the
- inner fjord.

In the future, freshwater runoff is expected to increase and will likely result in higher marine productivity in Nuup Kangerlua, with positive influence on Greenlandic fisheries and other harvestable marine resources. It is likely that, in a warmer_near-term future with more ice-sheet melt, the fjord system with marine-terminating glaciers that have not

580

retreated beyond their grounding line, may become a greater CO₂ sink, as increasing freshwater runoff is related toincreasing marine production and a prolonged annual productive season due to increased nutrient availability. Ultimately,the future productivity of this system will be tightly linked to the fate of its marine-terminating glaciers.

References

Andersen, T.J.: Some Practical Considerations Regarding the Application of 210Pb and 137Cs Dating to Estuarine

585

Sediments, in: Applications of Paleoenvironmental Techniques in Estuarine Studies, Springer, Developments in

Paleoenvironmental Research (DPER), vol. 20, pp. 121-140, <u>https://doi.org/10.1007/978-94-024-0990-1_6</u>, 2017.

Formatted: Subscript

Field Code Changed

Arendt, K.E., Dutz, J., Jonasdottir, S.H., Jung-Madsen, S., Mortensen, J., Moller, E.F., and Nielsen, T.G.: Effects of suspended sediments on copepods feeding in a glacial influenced sub-Arctic fjord, J. Plankton Res., 33 (10), 1526-1537, doi:10.1093/plankt/fbr054, 2011.

590 Arendt, K.E., Agersted, M.D., Sejr, M.K., and Juul-Pedersen, T.: Glacial meltwater influences on plankton community structure and the importance of top-down control (of primary production) in a NE Greenland fjord, Estuar., Coast. Shelf Sci., 183, 123-135, 2016.

Arrigo, K.R. and van Dijken, G.I.: Continued increases in Arctic Ocean primary production, Prog. Oceanogr., 136, 60-70, 2015.

595 van As, D., Andersen, M.L., Petersen, D., Fettweis, X., van Angelen, J.H., Lenaerts, J.T.M., van den Broeke, M., Lea, J.M., Bøggild, C.E., Alhstrøm, A.P., and Steffen, K.: Increasing meltwater discharge from the Nuuk region of the Greenland ice sheet and implications for mass balance (1960-2021), J. Glaciol., 60, 314-322, doi: 10.3189/2014JoG13J065, 2014.

Aschwanden, A., Fahnestock, M.A., Truffer, M., Brinkerhoff, D.J., Hock, R., Khroulev, C., Mottram, R., and Khan, S.A.: Contribution of the Greenland Ice Sheet to sea level over the next millennium, Sci. Adv., 5, doi: 10.1126/sciadv.aav9396,

2019. Bamber, J.L., Tedstone, A.J., King, M.D., Howat, I.M., Enderlin, E.M., van den Broeke, M.R., and Noel, B.: Land Ice Freshwater Budget of the Arctic and North Atlantic Oceans: 1. Data, Methods, and Results, J. Geophys. Res. Oceans,

123, 1827-1837, https://doi.org/10.1002/2017jc013605, 2018.

605 Barão, L., Vandevenne, F., Clymans, W., Frings, P., Ragueneau, O., Meire, P., Conley, D.J., and Struyf, E.: Alkalineextractable silicon from land to ocean: A challenge for biogenic silicon determination, Limnol. Oceanogr. Meth., 13(7), 329-344, <u>https://doi.org/10.1002/lom3.10028</u>, 2015.

Battarbee, R.W., Jones, V.J., Flower, R.J., Cameron, N.G., Bennion, H., Carvalho, L., and Juggins, S.: Diatoms, in: Tracking Environmental Change Using Lake Sediments Volume 3, Terrestial, Algal, and Siliceous Indicators, edited by:
Smol, J.P., Birks, H.J.B., and Last, W.M., Kluwer Academic Publishers, Dordrecht, Netherlands, 155-202, 2001.

Belt, S.T., Brown, T.A., Smik, L., Tatarek, A., Wiktor, J., Stowasser, G., Assmy, P., Allen, C.S., and Husum, K.: Identification of C25 highly branched isoprenoid (HBI) alkenes in diatoms of the genus Rhizosolenia in polar and subpolar marine phytoplankton, Org. Geochem., 110, 65-72, 2017.

Berthelsen, T.: Coastal fisheries in Greenland, KNAPK report, Nuuk, 2014.

615 Bianchi, T.S., Arndt, S., Austin, W.E.N., Benn, D.I., Bertrand, S., Cui, X., Faust, J.C., Koxiorowska-Makuch, K., Moy, C.M., Savage, C., Smeaton, C., Smith, R.W., and Syvitski, J.: Fjord as Aquatic Critical Zones (ACZs), Earth Sci. Rev., 203, 103145, 2020.

Bindoff, N.L., Cheung, W.W.L., Kairo, J.G., Arístegui, J., Guinder, V.A., Hallberg, R., Hilmi, N., Jiao, N., Karim, M.S., Levin, L., O'Donoghue, S., Purca Cuicapusa, S.R., Rinkevich, B., Suga, T., Tagliabue, A., and Williamson, P.: Changing

- 620 Ocean, Marine Ecosystems, and Dependent Communities, 2019, in: IPCC Special Report on the Ocean and Cryosphere in a changing Climate, edited by: Pörtner, H.-O., Roberts, D.C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., Mintenbeck, K., Alegría, A., Nicolai, M., Okem, A., Petzold, J., Rama, B., Weyer, N.M., 2019. Bjørk, A.A., Kjær, K.H., Korsgaard, N.J., Khan, S.A., Kjeldsen, K.K., Andresen, C.S., Box, J.E., Larsen, N.K., and Funder, S.: An aerial view of 80 years of climate-related glacier fluctuations in southeast Greenland, Nat. Geosci., 5(6),
- 625 427-432, <u>https://doi.org/10.1038/ngeo1481</u>, 2012.

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

	Boye, T.K., Simon, M., and Madsen, P.T.: Habitat use of humpback whales in Godthåbsfjord, West Greenland, with	
	implications for commercial exploitation, J. Mar. Biolog. Assoc. of the United Kingdom, 90, 1529-1539, 2010.	
	Compo, G.P., Whitaker, J.S., Sardeshmukh, P.D., Matsui, N., Allan, R.J., Yin, X., Gleason, B.E., Vose, R.S., Rutledge,	
620	G., Bessemoulin, P., Bronnimann, S., Brunet, M., Crouthamel, K.I., Grant, A.N., Groisman, P.Y., Jones, P.D., Kruk,	
630	M.C., Kruger, A.C., Marshall, G.J., Maugeri, M., Mok, H.Y., Nordli, Ø., Ross, I.F., Irigo, R.M., Wang, X.L., Woodruff,	
	S.D., and Worley, S.J.: The Twentieth Century Reanalysis Project, Q.J.R. Meteorol. Soc., 137, 1-28,	C
	https://doi.org/10.1002/qj.//6, 2011.	-1
	Cremer, H.: The diatom flora of the Laptev Sea (Arctic Ocean), Bibliotheca diatomologica, 40, Berlin and Stuttgart, 1998.	
	Delhasse, A., Kittel, C., Amory, C., Hofer, S., van As, D., S. Fausto, R., and Fettweis, X.: Brief communication:	
635	Evaluation of the near-surface climate in ERA5 over the Greenland Ice Sheet, Cryosphere, 14, 957-965,	C
	https://doi.org/10.5194/tc-14-957-2020, 2020.	-1
	DeMaster, D.J.: Measuring biogenic silica in marine sediments and suspended matter, in: Marine Particles: Analysis and	
	Characterization, Geophysical Monograph 63, American Geophysical Union, edited by: Hurd, D.C. and Spencer, D.W.,	
6.40	Washington, D.C., 363-367, 1991.	
640	Detlet, H., Reilly, B., Jennings, A., Jensen, M.M., O Regan, M., Glasius, M., Olsen, J., Jakobsson, M., and Pearce, C.:	
	Holocene sea-ice dynamics in Petermann Fjord in relation to ice tongue stability and Nares Strait ice arch formation,	
	Cryosphere, 15, 4557-4380, https://doi.org/10.5194/tc-15-4557-2021, 2021.	
	Fettweis, X., Franco, B., Tedesco, M., van Angelen, J.H., Lenaerts, J.T.M., van den Broeke, M.R., and Gall, H.: The	
C 4 F	Cryosphere estimating the Greenland Ice Sheet surface mass balance contribution to ruture sea level rise using the regional	
645	atmospheric climate model MAR, Cryosphere, 7, 469-489, 2013.	
	Peterseting of the 1000 2015 Growland ice cheet surface mass helence using the regional elimete MAP model	
	Creacebare 11, 1015 1022, unusur the arrigenbare pet/11/1015/2017/ 2017	C
1	Enged N : Diatoms from southwest Greenland Meddelelser om Granland: 194 pr. 5. Kabenhavn, C.A. Reitsel, 1973	
650	Glud R N Kühl M Wenzhöfer F, and Rysgaard S: Benthic diatoms of a high Arctic fiord (Young Sound NE	- (
	Greenland): importance for ecosystem primary production, Mar. Ecol. Prog. Ser., 238, 15-29, 2002.	
	Hammer, Ø., Harper, D.A.T., and Ryan, P.D.: PAST: Paleontological Statistics Software Package for Education and Data	
	Analysis, Paleontol. Electron., 4(1):9pp, 2001.	
	Harff, J., Perner, K., Moros, M.: Deglacial history, coastal development, and environmental change in West Greenland	
655	during the Holocene: Results of the R/V "Maria S, Merian" expedition MSM05/03, 15th June to 4th July 2007. Meereswiss.	
	Ber., Warnemünde, 99, doi:10.12754/msr-2016-0099, 2026.	
	Hasle, G.R. and Heimdal, B.R.: The net phytoplankton from Kongsfjorden, Svalbard, July 1988, with general remarks on	
	species composition of arctic phytoplankton, Polar Res., 17, 31–52, 1998.	
	Hasle, G.R. and Syvertsen, E.E.: Marine diatoms, in: Identifying Marine Phytoplankton, edited by: Tomas, C.R.,	
660	Academic Press, California, 5-385, 1996.	
	Hawkings, J.R., Wadham, J.L., Tranter, M., Lawson, E., Sole, A., Cowton, T., Tedstone, A.J., Bartholomew, I., Nienow,	
	P., Chandler, D., and Telling, J.: The effect of warming climate on nutrient and solute export from the Greenland Ice	
	Sheet, Geochem. Perspect. Lett., 1, 94-104, doi:10.7185/geochemlet.1510, 2015.	

Field Code Changed

Field Code Changed

Field Code Changed

Formatted: English (United States)

Hodal, H., Falk-Petersen, S., Hop, H., Kristiansen, S., and Reigstad, M.: Spring bloom dynamics in Kongsfjorden,
Svalbard: Nutrients, phytoplankton, protozoans and primary production, Polar Biol., 35, 191-203, doi:10.1007/s00300-011-1053-7, 2012.

Holding, J.M., Markager, S., Juul-Pedersen, T., Paulsen, M.L., Møller, E.F., Meire, L., and Sejr, M.K.: Seasonal and spatial patterns of primary production in a high-latitude fjord affected by Greenland Ice Sheet run-off, Biogeosciences, 16, 3777-3792, 2019.

670 Hopwood, M.J., Carroll, D., Browning, T.J., Meire, L., Mortensen, J., Krisch, S., and Achterberg, E.P.: Non-linear response of summertime marine productivity to increased meltwater discharge around Greenland, Nat. Commun., 9:3256, doi: 10.1038/s41467-018-05488-8, 2018.

Hopwood, M.J., Carroll, D., Dunse, T., Hodson, A., Holding, J.M., Iriarte, J.L., Ribeiro, S., Achterberg, E.P., Cantoni, C., Carlson, D.F., Chierici, M., Clarke, J.S., Cozzi, S., Fransson, A., Juul-Pedersen, T., Winding, H.S., and Meire, L.:

 675 Review article: How does glacier discharge affect marine biogeochemistry and primary production in the Arctic?, Cryosphere, 14, 1347-1383, 2020.
 Hurrell, J.W.: Decadal trends in the North Atlantic Oscillation: regional temperatures and precipitation, Science, 269,

676-679, 1995.

Jensen, K.G.: Holocene hydrographic changes in Greenland coastal waters, Ph.D. thesis, The Geological Survey of Denmark and Greenland, Copenhagen, Denmark, 2003.

Jensen, L.M. and Rasch, M.: Nuuk Ecological Research Operations, 2nd Annual Report, National Environmental Research Institute, Aarhus University, Denmark, 80 pp., 2008.

Juul-Pedersen, T., Arendt, K.E., Mortensen, J., Blicher, M.E., Søgaard, D.H., and Rysgaard, S.: Seasonal and interannual phytoplankton production in a sub-Arctic tidewater outlet glacier fjord, SW Greenland, Mar. Ecol. Prog. Ser., 524, 27-38, 2015.

Kanna, N., Sugiyama, S., Ohashi, Y., Sakakibara, D., Fukamachi, Y., and Nomura, D.: Upwelling of macronutrients and dissolved inorganic carbon by a subglacial freshwater driven plume in Bowdoin Fjord, northwestern Greenland, J. Geophys. Res. Biogeosci., 123, 1666-1682. https://doi.org/10.1029/2017JG004248, 2018.

Kahru, M., Brotas, V., Manzano-Sarabia, M., and Mitchell, B.G.: Are phytoplankton blooms occurring earlier in the Arctic?, Glob. Chang. Biol., 17, 1733-1739, doi:10.1111/j.1365-2486.2010.02312.x, 2011.

Karsten, U., Schlie, C., Woelfel, J., and Becker, B.: Benthic diatoms in Arctic Seas – Ecological functions and adaptations, Polarforschung, 81 (2), 77-84, 2011.

Kemp, A.E.S., Pearce, R.B., Grigorov, I., Rance, J., Lange, C.B., Quilty, P., and Salter, I.: Production of giant marine diatoms and their export at oceanic frontal zones: Implications for Si and C flux from stratified oceans, Global
Biogeochem. Cycles, 20, GB4S04, doi:10.1029/2006GB002698, 2006.

Korsgaard, N.J., Nuth, C., Khan, S.A., Kjeldsen, K.K., Bjørk, A.A., Schomacker, A., and Kjær, K.H.: Digital elevation model and orthophotographs of Greenland based on aerial photographs from 1978-1987, Sci. Data, 3, https://doi.org/10.1038/sdata.2016.32, 2016.

Krawczyk, D.W., Witkowski, A., Juul-Pedersen, T., Arendt, K.E., Mortensen, J., and Rysgaard, S.: Microplankton
 succession in a SW Greenland tidewater glacial fjord influenced by coastal inflows and run-off from the Greenland Ice
 Sheet, Polar Biol., 38, 1515-1533, 2015a.

Field Code Changed

Field Code Changed

29

680

685

Krawczyk, D., Arendt, K.E., Juul-Pedersen, T., Sejr, M.K., Blicher, M.E., and Jakobsen, H.H.: Spatial and temporal distribution of planktonic protists in the East Greenland fjord and offshore waters, Mar. Ecol. Prog. Ser., 538, 99-116, 2015b.

705 Krawczyk, D., Meire, L., Lopes, C., Juul-Pedersen, T., Mortensen, J., Li, C.L., and Krogh, T.: Seasonal succession, distribution, and diversity of planktonic protists in relation to hydrography of the Godthåbsfjord system (SW Greenland), Polar Biol., 2033-2052, https://doi.org/10.1007/s00300-018-2343-0, 2018. Kumar, V., Tiwari, M., Nagoji, S., and Tripathi, S.: Evidence of anomalously low $\delta^{13}C$ of marine organic matter in an arctic fjord, Sci. Rep., 6:36192, doi:10.1038/srep36192, 2016. 710 Lannuzel, D., Tedesco, L., van Leeuwe, M., Campbell, K., Flores, H., Delille, B., Miller, L., Stefels, J., Assmy, P., Bowman, J., Brown, K., Castellani, G., Chierici, M., Crabeck, O., Damm, E., Else, B., Fransson, A., Fripiat, F., Geilfus, N.-X., Jacques, C., Jones, E., Kaartokallio, H., Kotovich, M., Meiners, K., Moreau, S., Nomura, D., Peeken, I., Rintala, J.-M., Steiner, N., Tison, J.-L., Vancoppenolle, M., Van der Linden, F., Vichi, M., and Wongpan, P.: The future of Arctic sea-ice biogeochemistry and ice-associated ecosystems, Nature Climate Change, 10, 983-992, 715 https://doi.org/10.1038/s41558-020-00940-4, 2020. **Field Code Changed** Laufer-Meiser, K., Michaud, A.B., Maisch, M., Byrne, J.M., Kappler, A., Patterson, M.O., Røy, H., and Jørgensen, B.B.: Potentially bioavailable iron produced through benthic cycling in glaciated Arctic fjords of Svalbard, Nat. Commun., 12, 1349, 2021. Lea, J.M., Mair, D.W.F., Nick, F.M., Rea, B.R., van As, D., Morlighem, M., Nienow, P.W., and Weidick, A.: Fluctuations 720 of a Greenlandic tidewater glacier driven by changes in atmospheric forcing: observations and modelling of Kangiata Nunaata Sermia, 1859-present, Cryosphere, 8, 2031-2045. doi:10.5194/tc-8-2031-2014, 2014. Lewis, K.M., van Dijken, G.L., and Arrigo, K.R.: Changes in phytoplankton concentration now drive increased Arctic Ocean primary production, Science, 369, 198-202, 2020. Limoges, A., Ribeiro, S., Weckström, K., Heikkilä, M., Zamelczyk, K., Andersen, T.J., Tallberg, P., Massé, G., Rysgaard, 725 S., Nørgaard-Pedersen, N., and Seidenkrantz, M.-S.: Linking the modern distribution of biogenic proxies in High Arctic Greenland shelf sediments to sea ice, primary production, and Arctic-Atlantic inflow, J. Geophys. Res. Biogeosci., 123, 760-786. https://doi.org/10.1002/2017JG003840, 2018a. Field Code Changed

Limoges, A., Massé, G., Weckström, K., Poulin, M., Ellegaard, M., Heikkilä, M., Geilfus, N.-X., Sejr, M.K., Rysgaard, S., and Ribeiro, S.: Spring succession and vertical export of diatoms and IP25 in a seasonally ice-covered high arctic 730 fjord, Front. Earth Sci., 6:226, doi:10.3389/feart.2018.00226, 2018.

- Limoges, A., Weckström, K., Ribeiro, S., Georgiadis, E., Hansen, K.E., Martinez, P., Seidenkrantz, M.-S., Giraudeau, J., Crosta, J., and Massé, G.: Learning from the past: Impact of the Arctic Oscillation on sea ice and marine productivity off northwest Greenland over the last 9,000 years, Glob. Change Biol., 26, 6767-6786, https://doi.org/10.1111/gcb.15334, 2020.
- 735 Luostarinen, T., Ribeiro, S., Weckström, K., Sejr, M., Meire, L., Tallberg, P., and Heikkilä, M.: An annual cycle of diatom succession in two contrasting Greenlandic fjords: from simple sea-ice indicators to varied seasonal strategists, Mar. Micropaleontol., 158: 101873, 2020.

Lydersen, C., Assmy, P., Falk-Petersen, S., Kohler, J., Kovacs, K.M., Reigstad, M., Steen, H., Strøm, H., Sundfjord, A., Varpe, Ø., Walczowski, W., Weslawski, J.M., and Zajaczkowski, M.: The importance of tidewater glaciers for marine mammals and seabirds in Svalbard, Norway, J. Mar. Syst., 129, 452-471, 2014.

Field Code Changed

Field Code Changed

30

Mankoff, K.D., Noël, B., Fettweis, X., Ahlstrøm, A.P., Colgan, W., Kondo, K., Langley, K., Sugiyama, S., van As, D., and Fausto, R.S.: Greenland liquid water discharge from 1958 through 2019, Earth Syst. Sci. Data, 12, 2811-2841, https://doi.org/10.5194/essd-12-2811-2020, 2020.

Field Code Changed Mankoff, K.D., Fettweis, X., Langen, P.L., Stendel, M., Kjeldsen, K.K., Karlsson, N.B., Noël, B., van den Broeke, M.R., 745 Colgan, W., Simonsen, S.B., Box, J.E., Solgaard, A., Ahlstrøm, A.P., Andersen, S.B., and Fausto, R.S.: Greenland ice sheet mass balance from 1840 through next week, Earth Syst. Sci. Data, https://doi.org/10.5194/essd-2021-131, 2021. **Field Code Changed** Martinez, E., Antoine, D., D'Ortenzio, F., and Gentili, B.: Climate-driven basin-scale decadal oscillations of oceanic phytoplankton, Science, 326, doi: 10.1126/science.1177012, 2009. Meire, L., Søgaard, D.H., Mortensen, J., Meysman, F.J.R., Soetaert, K., Arendt, K.E., Juul-Pedersen, T., Blicher, M.E., 750 and Rysgaard, S.: Glacial meltwater and primary production are drivers of strong CO2 uptake in fjord and coastal waters adjacent to the Greenland Ice Sheet, Biogeosciences, 12, 2347-2363, doi: 10.5194/bg-12-2347-2015, 2015. Meire, L., Mortensen, J., Rysgaard, S., Bendtsen, J., Boone, W., Meire, P., and Meysman, J.R.: Spring bloom dynamics in a subarctic fjord influenced by tidewater outlet glaciers (Godthåbsfjord, SW Greenland), J. Geophys. Res. Biogeosci., 121, 1581-1592. doi:10.1002/2015JG003240, 2016a. 755 Meire, L., Meire, P., Struyf, E., Krawczyk, D.W., Arendt, K.E., Yde, J.C., Juul-Pedersen, T., Hopwood, M., Rysgaard, S., and Meysman, F.J.R.: High export of silica from the Greenland Ice Sheet, Geophys. Res. Lett., 43, 9173-9182, doi:10.1002/2016GL070191, 2016b. Meire, L., Mortensen, J., Meire, P., Juul-Pedersen T., Sejr, M.K., Rysgaard, S., Nygaard, R., Huybrechts, P., and Meysman, F.J.R.: Marine-terminating glaciers sustain high productivity in Greenland fjords, Glob. Change Biol., 23, 760 5344-5357 2017 Meyers, P.A.: Preservation of elemental and isotopic source identification of sedimentary organic matter, Chem. Geol., 114 (3-4), 289-302, https://doi.org/10.1016/0009-2541(94)90059-0, 1994. Field Code Changed Mikkelsen, N., Lennert, A., Ribeiro, S., Nørgaard-Pedersen, N.: Marine geological sampling in the Godthåbsfjord region. Geological cruise report - R/V Sanna 10th-18th May 2012, Danmarks og Grønlands geologiske undersøgelse rapport 765 2012/89, 2012 Mortensen, J., Lennert, K., Bendtsen, J., and Rysgaard, S.: Heat sources for glacial melt in a sub-Arctic fjord (Godthåbsfjord) in contact with the Greenland Ice Sheet, J. Geophys. Res., 116, C01013, doi:10.1029/2010JC006528, 2011. Mortensen, J., Bendtsen, J., Motyka, R.J., Lennert, K., Truffer, M., Fahnestock, M., and Rysgaard, S.: On the seasonal 770 freshwater stratification in the proximity of fast-flowing tidewater glaciers in a sub-Arctic sill fjord, J. Geophys. Res. Oceans, 118, 1382-1395, doi:10.1002/jgrc.20134, 2013. Motyka, R.J., Cassotto, R., Trueffer, M., Kjeldsen, K.K., van As, D., Korsgaard, N.J., Fahnestock, M., Howat, I., Langen, P.L., Mortensen, J., Lennert, K., and Rysgaard, S.: Asynchronous behavious of outlet glaciers feeding Godthåbsfjord (Nuup Kangerlua) and the triggering of Narsap Sermia's retreat in SW Greenland, J. Glaciol., 63 (238), 288-308, 2017. 775 Mullin, J. and Riley, J.: The colorimetric determination of silicate with special reference to sea and natural waters, Anal. Chim. Acta, 12, 162-176, https://doi.org/10.1016/S0003-2670(00)87825-3, 1955. **Field Code Changed** Oksman, M., Juggins, S., Miettinen, A., Witkowski, A., and Weckström, K.: The biogeography and ecology of common diatom species in the northern North Atlantic, and their implications for paleoceanographic reconstructions, Mar. Micropaleontol., 148, 1-28, https://doi.org/10.1016/j.marmicro.2019.02.002, 2019.

Field Code Changed

780	von Quillfeldt, C.H. Common diatom species in Arctic spring blooms: Their distribution and Abundance, Bot. Mar., 43, 499-516, 2000.		
	von Quillfeldt, C.H.: The diatom Fragilariopsis cylindrus and its potential as an indicator species for cold water rather than for sea ice, Vie Milieu 54, 137-143, 2004.		
	Racault, MF., Sathyendranath, S., Brewin, R.J.W., Raitsos, D.E., Jackson, T., and Platt, T.: Impact of El Niño variability		
785	on oceanic phytoplankton, Front. Mar. Sci., 4:133, doi:10.3389/fmars.2017.00133, 2017.		
	Renaut, S., Devred, E., and Babin, M.: Northward expansion and intensification of phytoplankton growth during the early		
	ice-free season in Arctic, Geophys. Res. Lett., 45, 10,590-10,598, https://doi.org/10.1029/2018GLO78995, 2018.	 Field Code Changed	_
	Ribeiro, S., Limoges, A., Massé, Johansen, K.L., Colgan, W., Weckström, K., Jackson, R., Georgiadis, E., Mikkelsen,		
	N., Kuijpers, A., Olsen, J., Olsen, S.M., Nissen, M., Andersen, T.J., Strunk, A., Wetterich, S., Syväranta, J., Hendersson,		
790	A.C.G., Mackay, H., Taipale, S., Jeppesen, E., Larsen, N.K., Crosta, X., Giraudeau, J., Wengrat, S., Nuttall, S., Grønnow,		
	B., Mosbech, A., and Davidson, T.A.: Vulnerability of the North Water ecosystem to climate change, Nat. commun., 12,		
	4475, https://doi.org/10.1038/s41467-021-24742-0, 2021.	 Field Code Changed	_
	Ribeiro, S., Sejr, M.K., Limoges, A., Heikkilä, M., Andersen, T.J., Tallberg, P., Weckström, K., Husum, K., Forwick,		
	M., Dalsgaard, T., Massé, G., Seidenkrantz, MS., and Rysgaard, S.: Sea ice and primary production proxies in surface		
795	sediments from a High Arctic Greenland fjord: Spatial distribution and implications for paleoenvironmental studies,		
	Ambio, 46 (S1), 106-118, https://doi.org/10.1007/s13280-016-0894-2, 2017.	Field Code Changed	
	Ribergaard, M.H., Oceanographic investigations off West Greenland 2013, NAFO Scientific Council Documents, 14/001,		
	2014.		
	Rysgaard, S., Mortensen, J., Juul-Pedersen, T., Sørensen, L.L., Lennert, K., Søgaard, D.H., Arendt, K.E., Blicher, M.E.,		
800	Sejr, M.K., and Bendtsen, J.: High air-sea CO ₂ uptake rates in nearshore and shelf areas of Southern Greenland: Temporal		
	and spatial variability, Mar. Chem., 128-129, 26-33, 2012.		
	Ryves, D.B., Battarbee, R.W., Juggins, S., Fritz, S.C., and Anderson, N.J.: Physical and chemical predictors of diatom		
	dissolution in freshwater and saline lake sediments in North America and West Greenland, Limnol. Oceanogr. 51(3),		
	1355-1368, 2006.		
805	Saini, J., Stein, R., Fahl, K., Weiser, J., Hebbeln, D., Hillaire-Marcel, C., and de Vernal, A.: Holocene variability in sea		
	ice and primary productivity in the northeastern Baffin Bay, Arktos, 6, 55-73, https://doi.org/10.1007/s41063-020-00075-	 Field Code Changed	
1	y, 2020.	_	
	Schubert, C.J. and Calvert, S.E.: Nitrogen and carbon isotopic composition of marine and terrestrial organic matter in		
	Arctic Ocean sediments: implications for nutrient utilization and organic matter composition, Deep Sea Research, 48,		
810	789-810, https://doi.org/10.1016/S0967-0637(00)00069-8, 2001.		
I	Seidenkrantz, MS., Kuijpers, A., and Schmith, T.: Comparing past and present climate - a tool to distinguish between		
	natural and human-induced climate change. In: Beyond Kyoto: Addressing the challenges of climate change, IOP		
	Conference Series: Earth and Environmental Sciences (8), 012012, doi 10.1088/1755-13115/8/1012012, 2009.		
	Seidenkrantz, MS., Røy, H., Lomstein, B.A., Meire, L., and the shipboard and on-shore parties: Godthåbsfjord system		
815	and the West Greenland shelf with 'R/V Sanna', 11-16, Cruise report, Aarhus University, Denmark, 2014.		
	Stroeve, J. and Notz, D.: Changing state of Arctic sea ice across all seasons, Environ. Res. Lett., 13:103001, doi:		

10.1088/1748-9326/aade56, 2018.

Weckström, K., Roche, B., R., Miettinen, A., Krawczyk, D., Limoges, A., Juggins, S., Ribeiro, S., and Heikkilä, M.: Improving the plaeoceanographic proxy tool kit – On the biogeogeography and ecology of the sea ice-associated species Fragilariopsis oceanica, Fragilariopsis reginae-jahniae and Fossula arctica in the northern Atlantic, Mar. Micropaleontol.,

820

157, 101860, https://doi.org/10.1016/j.marmicro.2020.101860, 2020.

Author contribution

825 M.O. planned the study together with S.R. S.R. designed the project (*Greenshift*) and provided sediment core material together with N.N.P., M.-S.S., and N.M. M.O. conducted diatom, grain size, and BSi analysis, and A.B.K. assisted with BSi and grain-size analyses. T.J.A. carried out ²¹⁰Pb measurements. S.H.L. and W.T.C. provided freshwater runoff calculations for 1900-2010 together with K.D.M. who provided freshwater runoff data. K.K.K. provided data for ice margin position. M.O. wrote the manuscript with input from all the co-authors.

830

Acknowledgement

This study received financial support from Geocenter Danmark (project *Greenshift* grant 2018-5 to Sofia Ribeiro). The 2013 cruise with RV *Sanna* was funded by the Arctic Research Centre of Aarhus University, Denmark. SRI received additional funding from the Independent Research Fund Denmark (grant no. 9064-00039B). MSS was funded by the Danish Council for Independent Research (grant no. 7014-00113B (G-Ice) and 0135-00165B (GreenShelf) with additional funding from the European Union's Horizon research and innovation program under Grant Agreement no. 869383 (ECOTIP). We thank Ida Sofie Mikaelsdóttir Olsson, Annette Ryge and Charlotte Olsen for help with biogenic silica analysis and diatom slide preparation. Finally, we thank Thomas Juul-Pedersen and Ann Lennert for accommodating and supporting our sediment coring activities during the Greenland Ecosystem Monitoring cruise with RV *Sanna* in 2012.

835

Field Code Changed