

First evidence of microplastics in Antarctic snow

Alex R. Aves^{1,2}, Laura E. Revell¹, Sally Gaw¹, Helena Ruffell¹, Alex Schuddeboom¹, Ngaire E. Wotherspoon¹, Michelle LaRue², and Adrian J. McDonald^{1,2}

¹School of Physical and Chemical Sciences, University of Canterbury, Christchurch, New Zealand

²Gateway Antarctica, School of Earth and Environment, University of Canterbury, Christchurch, New Zealand

Correspondence: Alex R. Aves (alexandra.aves@pg.canterbury.ac.nz)

Abstract. In recent years, airborne microplastics have been identified in a range of remote environments. However, data throughout the Southern Hemisphere, in particular Antarctica, are largely absent to date. We collected snow samples from 19 sites across the Ross Island region of Antarctica. Suspected microplastic particles were isolated and their composition confirmed using micro-Fourier transform infrared spectroscopy (μ FTIR). We identified microplastics in all Antarctic snow samples at an average concentration of 29 particles L^{-1} , with fibres the most common morphotype and polyethylene terephthalate (PET) the most common polymer. To investigate sources, backward air mass trajectories were run from the time of sampling. These indicate potential long-range transportation of up to 6000 kilometers, assuming a residence time of 6.5 days. Local sources were also identified as potential inputs into the environment, as the polymers identified were consistent with those used in clothing and equipment from nearby research stations. This study adds to the growing body of literature regarding microplastics as a ubiquitous airborne pollutant, and establishes their presence in Antarctica.

1 Introduction

Over the last century plastics have become one of the most ubiquitous synthetic materials in the world, due to their versatility and durability. Despite their longevity, plastics degrade over time to produce microplastics (plastic particles <5 mm in diameter), and when present in the environment have the potential to cause significant ecological damage (MacLeod et al., 2021). Microplastics have negative effects on marine organisms (Wright et al., 2013), and act as vectors for persistent organic pollutants and other toxic substances (Mato et al., 2001; Rios et al., 2007), which are harmful to marine environments and organisms (Hermabessiere et al., 2017). Microplastics have been recognised as widespread pollutants in the marine environment (Ryan, 2015) and are known to be damaging to terrestrial ecosystems, (de Souza Machado et al., 2018), while their small size and relatively low density also allows them to become airborne and transported over large distances (Evangelidou et al., 2020).

Airborne microplastics have been identified in atmospheric fallout in a range of urban (Dris et al., 2015, 2016; Cai et al., 2017; Klein and Fischer, 2019; Knobloch et al., 2021) and remote regions worldwide (Allen et al., 2019; Bergmann et al., 2019; Brahney et al., 2020). It is now understood that microplastics transition between marine environments, terrestrial environments and the atmosphere via the plastic cycle (Horton and Dixon, 2018; Brahney et al., 2021). This allows microplastics to reach locations far from anthropogenic sources, such as the Arctic (Bergmann et al., 2019), the Tibetan Plateau (Zhang et al., 2021), European alpine regions (Allen et al., 2019; Bergmann et al., 2019; Materić et al., 2020, 2021) and conservation areas across

the continental United States (Brahney et al., 2020). Deposited microplastics may accelerate melting of the cryosphere when present on snow and ice in alpine or polar regions (Evangelidou et al., 2020). Microplastics may further influence climate by acting as cloud ice nuclei in the atmosphere (Ganguly and Ariya, 2019), and through their minor contribution to global radiative forcing (Revell et al., 2021).

30 ~~Antarctica was~~ With a few exceptions, such as lead pollution in the late 19th century (McConnell et al., 2014), Antarctica
was generally thought to be largely untouched by humans until the early 20th century due to its inaccessibility, extreme environmental conditions and barriers such as the Antarctic Circumpolar Current (Tin et al., 2014; Gordon, 1971). While the human footprint has increased over the last century, Antarctica ~~has been set aside as~~ is still a place of peace and science and is thought of as the last remaining true wilderness on earth (Tin et al., 2016). Due to this, Antarctica can act as an indicator of physical,
35 chemical, and biological effects caused from anthropogenic stresses (Huiskes et al., 2006). Research on microplastics in the Antarctic has focused on the marine environment, where particles have been detected in deep sea sediments in the Weddell Sea (Van Cauwenberghe et al., 2013), marine sediments from the western Antarctic Peninsula (Reed et al., 2018) and the Ross Sea (Munari et al., 2017), south of the Polar Front (Cózar et al., 2014) and in the surface waters of the Southern Ocean and Antarctic Peninsula (Absher et al., 2019; Cincinelli et al., 2017; Isobe et al., 2017; Suaria et al., 2020; Waller et al., 2017; Lacerda et al.,
40 2019). Microplastics were recently identified for the first time in a freshwater Antarctic Specially Protected Area (ASPA) on Livingston Island, which is used for long-term ecological monitoring due to its pristine nature and use as a reference for inland water research (González-Pleiter et al., 2020).

To date there is little information available regarding the presence of airborne microplastics in Antarctica. We collected freshly fallen snow samples from the Ross Island region of Antarctica in late 2019, and analyzed them to quantify the presence
45 and abundance of microplastics. Samples were collected close to two scientific research stations (Scott Base and McMurdo Station), and from 13 field sites up to 20 km from the research stations. We identified polymer composition using μ FTIR spectroscopy and analyzed air mass back-trajectories to identify the potential origins of sampled air masses. Further, we catalogued the composition of field equipment to understand local polymer sources.

2 Materials and methods

50 2.1 Field collection

Snow samples were collected in 500 mL stainless-steel bottles. Nineteen samples were collected in total with six from locations near research stations and thirteen from remote locations with minimal human disturbance (Fig. 1). Samples were collected using a stainless-steel scoop and stainless-steel funnel to fill each 500 mL bottle with snow from the top 2 cm of the surface. Samples were stored in a -25 °C freezer at Scott Base and were kept chilled with dry ice during transit to New Zealand. Upon
55 arrival in New Zealand, samples were stored at -20 °C.




Figure 1. Site locations for snow sample collections across the Ross Island region of Antarctica (S1–S19). Sample sites are marked in black, and locations of scientific research stations shown by blue triangles. Some site markers correspond to two sampling sites due to the scale of the map. Map data sourced from Matsuoka et al. (2021).

2.2 Laboratory analysis of microplastics

Snow samples were thawed in the sealed sample bottles at room temperature for 24–48 hours prior to analysis. Thawed samples were filtered through a glass apparatus attached to a vacuum using a cellulose nitrate membrane filter (Whatman nitrocellulose membrane, 50 mm diameter, 0.45 μm pore size). The volume of liquid (melted snow) was recorded at this step, before rinsing, to establish the volume of each individual sample bottle (Table. A2). Approximately 10–20 mL of 70% ethanol was used to rinse the filters, and a further 10–30 mL of 96% ethanol was added to soak the filter for 10 minutes, to prevent bacterial and viral growth for further biosecurity measures. The sides of the glassware were thoroughly rinsed with ultra-pure water ($<18\text{ M}\Omega$) to dislodge any microplastics adhered to the walls of the filtering equipment, and samples were dried under vacuum.

The dried cellulose nitrate filter was transferred to a 250 mL glass beaker for a wet peroxide oxidation (WPO) digestion to remove organic material present on the filter. An Iron (Fe(II)) sulfate solution (0.05 M) was prepared by adding 7.5 g of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ ($= 278.02\text{ g mol}^{-1}$) to 500 mL of ultra-pure water and 3 mL of concentrated sulfuric acid (Fenton's reagent). This study followed similar methods to those outlined by the National Oceanic and Atmospheric Administration (NOAA) (Masura et al., 2015) for WPO digestion, with 20 mL of 30% H_2O_2 , 100 mL ultra-pure water and 5 mL of the Fe(II) solution added to the beaker with the filter. The solution was then left, with the beaker opening covered in aluminium foil, for 20

minutes before being stirred on a hot plate at 45 °C for 2–3 hours with a magnetic stir bar. The magnetic stir bar had a white polytetrafluoroethylene (PTFE) coating, which was not identified in any field or blank samples.

The digested cellulose nitrate filter was rinsed with ultra-pure water to remove any attached material. The digested filtrate was then filtered under vacuum onto a Whatman glass fibre GF/C filter (47 mm diameter, 1.2 µm pore size) and all glassware was thoroughly rinsed with ultra-pure water. All reagents used were previously unopened and analytical grade, with blanks also undergoing identical analysis to control for contamination. The GF/C filter was then vacuum dried before being removed with tweezers into a closed petri dish and labelled accordingly for storage until analysis.

2.3 Visual characterisation of microplastics and µFTIR analysis

Filter papers were initially screened using a Leica MZ125 stereomicroscope with 10× magnification for visual identification. Suspected plastic particles were identified according to characteristics related to the morphotype, shape, colour and the physical characteristics of each particle (Bridson et al., 2020). ~~Microplastics~~ Suspected microplastics were characterized into four main morphotypes – fibres, films, fragments and beads. Colours were recorded for each suspected particle. Analysis was performed by the same individual to ensure consistency in identification and counts. Each filter was visually analyzed three times. Due to limitations of visual identification techniques (Knobloch et al., 2021), dark colours were difficult to differentiate. Therefore throughout this study, ‘blue’ includes blue, black and navy. It is recognised that due to human error, inability to transfer some particles due to their small size and brittleness, and the translucent and transparent nature of some microplastics, that there are limitations to this method which are hard to avoid. This may lead to the underestimation of microplastics in this study. The smallest particle identified in this study was 44 µm (non-plastic), meaning particles less than this size were not accounted for due to analysis limitations.

All suspected microplastics were chemically identified by micro-Fourier transform infrared spectroscopy (µFTIR). This procedure followed pre-existing methods for characterizing microplastics (Primpke et al., 2018). A Hyperion 2000 µFTIR microscope (Bruker Optics), attached to a Vertex 70 (Bruker Optics) spectrometer, was used to analyse particles plated on a calcium fluoride (CaF₂) disk (25 mm diameter), with a liquid nitrogen cooled Mercury-Cadmium-Telluride (MCT) detector. Each particle was manually transferred onto the disk using tweezers and a drop of 96% ethanol to aid in transfer. An optical overview image was recorded before infra-red measurements were performed at 15× magnification. Scans were run in transmission mode (10 scans, 4 cm⁻¹ resolution, spectral range of 4000–1000 cm⁻¹) using OPUS 7.8 software. All spectra were ~~saved and run against a Wiley spectral library (Databases: HIX, OSX, HDX, PPX, KLX, FMX, YX, DAX, HUX, QPX, WSAHX).~~ baseline corrected and CO₂ spectral range excluded, saved and compared against the following Wiley spectral libraries (Databases: Industrial chemicals, pure organic compounds; organosilicons; polymers, Hummel defined basic; Sadtler acrylates and Methacrylates; Sadtler fibers and textile chemicals; Sadtler fibers by microscope; Sadtler inorganics; Sadtler polymers and monomers (comprehensive); Sadtler polymers, Hummel; Sadtler standards (organic and polymeric compounds subset); Sigma-Aldrich library of FT-IR spectra). Particles returning a hit quality index (HQI) of >70% against the library reference spectra were accepted as microplastics. Those <70% that exhibited plastic characteristics from visual screening and possessed similar µFTIR spectra were analyzed further by the authors using Wiley peak picking

105 tools to identify characteristic peaks of plastic polymer types ~~-(Kroon et al., 2018).~~These particles matching characteristic spectral peaks were included in results. Due to the environmental degradation sampled particles have been subjected to, and the limitations of spectral libraries due to the use of high standard polymers, visual inspection of spectra is an essential step in identification of environmental microplastic analysis (De Frond et al., 2021; Shim et al., 2017).

2.4 Quality control

110 2.4.1 Field sampling

Each sampling site was selected ensuring the presence of fresh snow with no visible contamination or movement in the collection area. Non-plastic sampling equipment was used to avoid contamination and all equipment was rinsed in nearby snow prior to each sample collection. Leather gloves were worn by the sample collectors and the sample site selected upwind from any human movement to minimise contamination by the samplers. The lid of each sampling bottle was held without the inside
115 being touched during sample collection to avoid human contamination. Bottles were stored upright and kept in a cool box in snow during transportation.

2.4.2 Controls and blanks

Two field controls were collected during sample collection: one alongside a remote field site sample and one alongside a research station sample. The two field control bottles were left open during snow sampling and filled with ultra-pure water when returned to the laboratory. They subsequently underwent the same laboratory methods as all other samples. Two method
120 controls were prepared in stainless-steel bottles identical to the sampling bottles used for collection and were filled with ultra-pure water. These were stored in the laboratory freezer in New Zealand during sampling and underwent identical laboratory procedures as the samples to identify potential sources of contamination from the sampling bottles. For blank corrections we followed the methodology highlighted in Brander et al. (2020) and used in Vandermeersch et al. (2015), whereby particles matching the identical characteristics of those found in the blanks (laboratory and field blanks) were omitted from further analysis, and total daily laboratory blank findings were subtracted from the corresponding samples (Table. S1). Daily laboratory blanks were analyzed using the same procedure as all samples with 500 mL of ultra-pure water. The laboratory blanks were analyzed to compare results for each individual date and accounted for in the data for the samples filtered on the corresponding days. Particles found in field samples with identical characteristics to those found
130 in blanks were discarded and excluded from the results.

2.4.3 Laboratory analysis

Method recovery tests consisted of two 500 mL samples of ultra-pure water spiked with 5 polyethylene (PE) beads and 7 polymethyl methacrylate (PMMA) fibres sized between 500–2000 µm. The spiked samples were analyzed identically to the field samples to measure recovery rates of the spiked samples. ~~Daily laboratory blanks were analyzed using the same procedure as all samples with 500 mL of ultra-pure water.~~
135 Filtration and laboratory procedures were performed in a Labrocure fume hood

cabinet to limit contamination by airborne microplastics. Glassware was cleaned inside the fume hood three times with ultra-pure water and once with acetone. Aluminium foil was used to cover glassware openings to minimise contamination by airborne microplastics. ~~The laboratory blanks were analyzed to compare results for each individual date and accounted for in the data for the samples filtered on the corresponding days.~~ Non-synthetic (wool and cotton) clothing was worn during the laboratory and analysis process. ~~Particles found in field samples with identical characteristics to those found in blanks were discarded and excluded from the results.~~ Surfaces were sprayed and wiped down with 70% ethanol prior to microscopy and analysis work, with filters remaining covered except when manual extraction of particles occurred.

2.5 Clothing composition

National Antarctic programs provide essential clothing and field gear for staff and scientists. Some of the gear provided is mandatory whenever undertaking fieldwork or travelling outside of bases. The composition of field gear provided by the New Zealand National Antarctic program (including base layers, mid layers, outer layers, shoes, boot liners, gloves, bags, hats and accessories) was catalogued to determine potential local sources of synthetic particles into the Ross Island region (Table A1).

2.6 Trajectory analysis

Because samples were collected during, or shortly after a single snowfall event (Fig. 2), air mass trajectories were produced to understand potential source regions. Lagrangian air parcel trajectories were derived using the Hybrid Single Particle Lagrangian Integrated Trajectory Model (HYSPLIT) (Stein et al., 2015). HYSPLIT was run using meteorological data from the National Oceanic and Atmospheric Administration Global Forecast System (GFS), with a horizontal grid resolution of $0.25^\circ \times 0.25^\circ$ and a temporal resolution of 3 hours. To achieve this temporal resolution the GFS model is initialized from observational data every 6 hours and then the three hour forecast is included in the GFS output to fill the gaps in time (Stein et al., 2015).

The ensemble configuration of HYSPLIT was used to generate 27-member back-trajectory ensembles, which represent the uncertainty in the trajectories. For each ensemble trajectory, the meteorological data was offset in the x and/or y directions by one grid point, and/or 0.1 sigma units in the z direction, with 27 ensemble members covering all possible combinations of these offsets. Back trajectories were run from starting points at each of the sampling sites at the corresponding time of sampling and with starting heights of 500, 1000 and 2000 m. Recent published results show that the residence time for microplastic particles in the atmosphere may vary between 1 hour and 156 hours (Brahney et al., 2021). As such, the back trajectories were run for 156 hours to show the full range of possible sources.

3 Results

3.1 Blanks and sample extractions

Recovery rates for spiked samples were 100%. Across the sample controls an average of ~~one particle was found in daily laboratory blanks (n=11), three particles in field blanks (n=2) and two particles in method controls (n=2).~~ In all blanks, fibres




Figure 2. Snowfall over the measurement period from the ERA5 reanalysis. These values are averaged over the sample site area (77.75–78°S, 166.5–167.25°E). Orange circles indicate the time when samples were taken with nearby text indicating the sample number.

were the most common morphotypes. Several suspected microplastic fragments present in the controls and samples resembled that of the sampling vessel. 1.5 ± 0.89 particles were found in daily laboratory blanks ($n=11$), 3 ± 0 in field blanks ($n=2$) and 2 ± 1 particles in method controls ($n=2$). All fragments identified in field and laboratory blanks matched the colour and coating of the sampling bottles. No fragments were found in the daily blanks, meaning fibres were the predominant morphotype (Table. S1). Spectroscopic analysis confirmed that the outside bottle coating and fragments found in the blanks were polymethyl methacrylate (PMMA). Suspected microplastic particles in field samples of an identical colour and morphotype to those detected in the field and laboratory blanks were not analyzed further and discounted from the results. All reported PMMA still included in results was a different morphotype and colour to the corresponding sampling bottles, and therefore some PMMA is still shown in results. The lining of the lids of the sampling bottles was confirmed as clear silicone, which was not found in any of the samples or blanks. Particles with identical characteristics of those found in the field and laboratory blanks were excluded from further study and daily blank contamination was subtracted from the results of corresponding samples.

3.2 Concentration of microplastics in snow samples

Microplastics were found in all Antarctic snow samples with a total of 109 particles confirmed as microplastics using μ FTIR spectroscopy across the 19 field samples (Fig. 1). Microplastics were present at an average concentration of 29.4 ± 4.7 particles L^{-1} of melted snow (mean \pm one standard error) across all sites (Fig. 3). The average microplastic concentration was




Figure 3. Concentrations of microplastics (MP) particles L^{-1} at each sampling site in the Ross Island region. Insets provide an overview of the location of sampling sites in Antarctica. Map data sourced from Matsuoka et al. (2021).

22.5 ± 4.0 particles L^{-1} and 47.2 ± 8.4 particles L^{-1} at remote sites and base sites, respectively (Fig. 3, Supplementary Fig. 1). The highest concentration found was 82 particles L^{-1} at site S16 (Scott Base) and the lowest concentration found was 4 particles L^{-1} at site S2 (Erebus Glacier Tongue).

185 ~~Concentrations of microplastics (MP) particles L^{-1} at each sampling site in the Ross Island region. Insets provide an overview of the location of sampling sites in Antarctica. Map data sourced from Matsuoka et al. (2021).~~

3.3 Characterization of microplastics

~~Polymer types identified across all samples. (a) Number of each polymer type found across all Antarctic study sites. (b) Number of microplastic fragments and fibres identified in each colour category (films ($n=1$) excluded).~~

190 ~~Size distribution of microplastics across all samples categorized by morphotype (length).~~

A total of 13 different polymer types were identified across the snow samples when compared against a spectral reference library (Supplementary Fig. 2). Polyethylene terephthalate (PET) was the most frequently detected polymer type, found in 79% of the samples, comprising 41% of total polymers identified (Fig. 4a). Copolymers (CP, those containing two or more different monomers) equated to 17% of total polymers identified (Fig. 4a and Table A2). Polymer types comprising less than 10% of the total polymers identified included polymethyl methacrylate (PMMA, 9%), polyvinyl chloride (PVC, 9%), polyamide (PA, 6%), polyethylene (PE, 4%), alkyd (ALK, 4%), cellulose nitrate (CN, 4%) and other (6%). Polymer types present in the ‘other’

195

category with a detection frequency less than 4% include polytetrafluoroethylene (PTFE), polyvinylidene, polypropylene, silicone and polymethyl anhydride. The cellulose nitrate particles detected were not found in laboratory control samples and did not match the morphotype and colour of the filters used in the digestion stage, so were included in results.

200 Particles confirmed spectroscopically as microplastics were classified as fibres, fragments, and films (Supplementary Fig. 3). No beads were detected. Fibres were the most abundant morphotype (61%, Fig. ??4) and the majority of microplastics identified were blue (55%) and pink (23%, Fig. 4b). Blue was the most common colour of both fibres and fragments. The size range of microplastics detected varied between 50–3510 μm with an overall average size of 606 μm (Fig. ??4). A large proportion of microplastics were <1000 μm (81%), with 28% in the 0–200 μm size range (Fig. ??4). All fragments were
205 $\leq 1000 \mu\text{m}$ while fibres were present in all size groups (Fig. ??4). The mean size for fragments was 200 μm while the mean size for fibres was 850 μm , which skews the total averages (Fig. ??4).

3.4 Snowfall and back trajectories

All of the samples from remote sites (S1–S13) were collected over three days (30 November – 2 December 2019). A large snowfall event occurred one day prior to sampling on 30 November 2019 (Fig. 2), and no snowfall was reported over the
210 subsequent two days, therefore all remote samples originated from the same snowfall event. Because only the top 2 cm of snow was collected, the microplastics we found were likely deposited in the recent snowfall event.

Figure 5 shows Lagrangian back-trajectories beginning at the time and location of six of the different sampling sites. The six events shown were chosen to cover the largest spatial distances and time periods within the data set, with many of the excluded sites showing near-identical results to included sites. This figure only shows the results for trajectories that start with
215 an altitude of 2000 m, but trajectories initialised at 500 m and 1000 m show very similar results (Fig. A1 and A2). Distances travelled by air masses over 6 hours, 24 hours and 156 hours are shown in Table A3 to represent a range of potential airborne microplastic residence times. The residence time is the length of time that a particle can remain in the atmosphere which was estimated by Brahney et al. (2021) to be as long as 156 hours for microplastics prior to deposition at the sampling site. We acknowledge that microplastics could be suspended in air for longer than the time periods used, although
220 unlikely, and that they could have been derived from further afield or have been deposited and resuspended from land or the sea en route.

The trajectories presented in Fig. 5 indicate that the snowfall event was likely linked to a Ross Air Stream event, where strong near-surface winds flow from the southeast parallel to the Trans-Antarctic Mountains (Parish et al., 2006). The Ross Air Stream is primarily fed by air from the Siple Coast region in West Antarctica, caused by a complex mixture of katabatic winds
225 and barrier flows along the Trans-Antarctic mountains created by cyclones (Seefeldt and Cassano, 2012). Pressure gradients induced by cyclones to the north of the ice shelf force air to flow towards the Trans-Antarctic mountains, but the resulting wind lacks the kinetic energy to pass over the mountains causing a barrier flow. These conditions occur approximately 24% of the time, though are more common in the Austral winter, and bring warmer temperatures and strong winds to the Ross Island region (Coggins et al., 2014) along with clouds connected to snowfall (Jolly et al., 2018). The splitting of the Ross Air Stream
230 around Ross Island also impacts the formation of the Ross Sea polynya and McMurdo Sound polynya (Dale et al., 2017; Brett

et al., 2020). These open water regions provide a possible secondary local source of microplastics; in particular, the McMurdo Sound polynya is very close to site S2. The back trajectories shown in Fig. A1 for Sites 3, 7 and 13 could mean that the open water in the Ross Sea polynya is a relevant source.

In accordance with the climatology of the region, all of the trajectories show that the immediate source of the airflow is from the south following the Trans-Antarctic mountains. This means that for the majority of these samples, short term local transport is the most likely source of microplastics as the sampling sites are mostly north of the local bases (Scott Base and McMurdo Base). It is also possible that local small scale transport processes that are not captured by HYSPLIT could play a key role in transport at this scale. For the trajectories that clearly show a Ross Air Stream event (sites 2, 3, 7 and 13), the most likely distant sources are the Ross and Amundsen Seas as there are no manned stations or other likely sources along the trajectory path. The transport processes in this region can be very rapid with the trajectories from these sites on average covering a distance of 143 km in the first 6 hours and 469 km in 24 hours. With a 156 hour residence time, transport over thousands of kilometers is possible.

For site 19 there is a different set of meteorological conditions causing a divergence from the results from the other samples. The corresponding trajectories show a wide range of possible sources including the Antarctic Peninsula and Weddell Sea. While these conditions are relatively rare compared with the kind of Ross Air Stream flow examined earlier, they still provide a possible source of microplastics. These different conditions expand the set of possible sources to include more distant sources including many other Antarctic research stations. Given that an assumed maximum residence time of 156 hours leads to trajectories covering a total distance of over ~ 4800 km in transit (Table A3), transport from these stations appears possible but unlikely.

4 Discussion

4.1 Microplastic concentrations

Our work provides the first evidence of microplastics in Antarctic snow, and critically, the average concentration of microplastics found in this study are higher (29.4 ± 4.7 particles L^{-1}) than in the surrounding Ross Sea (1.7×10^{-4} particles L^{-1}) and those reported in East Antarctic sea ice (11.7 particles L^{-1}) (Kelly et al., 2020; Cincinelli et al., 2017). This dataset is limited by the low sample volumes due to the permitting restrictions for Antarctic samples. Therefore, we recommend this study to be replicated to further understand these preliminary findings. Larger volumes of snow (>10 L) or replicates from the same study sites would be beneficial for future research. Our reported concentrations of microplastics in surface snow near two Antarctic stations (47.2 ± 8.4 particles L^{-1} ; highest concentration 82.1 particles L^{-1}) identified similar concentrations to those present in supraglacial debris from an Italian glacier (74 particles kg^{-1} dry weight) (Ambrosini et al., 2019). For comparison, concentrations of microplastics in Arctic sea ice have been shown to range from 8 to 41 particles L^{-1} (Geilfus et al., 2019), while Arctic snow sampling identified higher concentrations of microplastics up to 14.4×10^3 particles L^{-1} (Bergmann et al., 2019), which may be attributed to the proximity to more populated regions.

4.2 Microplastic characteristics

The most frequent polymer type we detected was PET, which was found in 79% of all samples. Approximately 60% of all PET produced are used for synthetic fibres and 30% for plastic bottles (Ji, 2013). Our results differ to surrounding marine studies which found PE and PP were the most common polymer types in seawater samples from the Ross Sea region (Cincinelli et al., 2017). Comparatively, varnish (including acrylates) and rubber were the most common in Arctic snow collected using similar sampling techniques, with PMMA being the third most common polymer type in this study (Bergmann et al., 2019). Air samples collected over the West Pacific Ocean similarly found PET was present in the highest abundance (57%) (Liu et al., 2019b).

Fibres were the most abundant morphotype (60%) followed by fragments (39%) and films (1%, Fig. 224). These findings are consistent with previous studies whereby fibres were also the dominant morphotype of airborne microplastics due to their relatively low density and physical characteristics (Liu et al., 2019a; Dris et al., 2016; Bullard et al., 2021). Analysis of seawater samples in the surrounding Ross Sea identified fragments as the predominant morphotype (72%) (Cincinelli et al., 2017), in contrast to our results. Fibres were the most predominant morphotype in sediment from Terra Nova Bay (Munari et al., 2017), which is consistent with the findings of this research. This likely indicates that the distribution of microplastics around the Ross Sea region is heterogeneous depending on the sample type. Further research is required to understand the microplastic footprint in the region. Similar to previous studies, we identified darker colours (blue, black and navy) as the most common (Ambrosini et al., 2019; Liu et al., 2019a, b). Dark-coloured microplastics are likely efficient at absorbing solar radiation compared to lighter colours and are of particular concern in the cryosphere as they may accelerate melting (Evangelou et al., 2020).

As the size distribution of identified microplastics is skewed towards smaller particles (Fig. 224), it is likely that particles smaller than the smallest particle observed (50 μm) are present, but not able to be detected due to the magnification limit of the stereomicroscope (20 μm) and difficulties in handling particles $<50 \mu\text{m}$. The abundance of microplastics has previously been shown to increase with decreasing size (Isobe et al., 2017; Levermore et al., 2020), which corresponds to the findings of this study (Fig. 224). The size distribution from our study was comparable to those measured in the remote Pyrenees (Allen et al., 2019). We identified that only 10% of the microplastics were $<100 \mu\text{m}$ while findings in the Arctic reported 98% of particles were $<100 \mu\text{m}$ (Bergmann et al., 2019). While fibres were present across all size ranges found, the only fragments found were smaller than 1000 μm , consistent with previous studies (Revell et al. (2021) and references therein).

4.3 Origin of microplastics

Microplastics in Antarctica may originate from both local sources and long-range transport. Direct sources of microplastics to the Antarctic environment may include fragmentation of plastic equipment from research stations, clothing worn by base staff and researchers, and mismanaged waste. Microplastics may also enter the Antarctic environment via long range transport by ocean currents (Fraser et al., 2018), ocean to atmosphere exchange (Allen et al., 2020) and both short and long-range atmospheric transportation (Evangelou et al., 2020; Brahney et al., 2021).

Antarctic research stations on Ross Island, Scott Base (NZ) and McMurdo Station (US), ~~are within~~ have the closest proximity to the sampling sites, up to 20 km away (Fig. 1), with Zucchelli Station (Italian) the next closest at 350 km away, which is only operational over summer. Direct transport from Zucchelli station is highly unlikely given that it is north from our sampling sites and the winds primarily come from the south. McMurdo Station has a maximum capacity of ~~1,300 people which is met~~ 1,200 people in the summer months, and decreases to approximately ~~300–150~~ people over winter, whereas Scott Base has a capacity of 86 people over summer and typically hosts ~~12–11~~ staff over winter (COMNAP, 2017). The concentration of microplastics measured at the base sites (S14-19; 47 ± 8 particles L^{-1}) was higher on average than the remote sites (S1-13; 22 ± 4 particles L^{-1}) suggesting that microplastics originated from local sources.

Plastic products in use at research stations (including building materials, marker flags, safety equipment and tyre rubber) may fragment with environmental exposure and be a potential local source of microplastics into the environment. General wear and weathering from clothing and outdoor equipment used in the field may introduce plastics into more remote regions away from populated bases. In addition, enhanced ultra-violet fluxes due to the Antarctic ozone hole may accelerate the fragmentation of larger plastic products into microplastics (Williamson et al., 2019).

PET was the most common polymer found in snow samples making up 41% of total microplastics. The most common polymer found in catalogued gear was also PET which was present in 48% of the garments in varying percentages from 68-100% followed by PA which was present in 30% of the garments (Table A1). Fibres were also the most common morphotype in snow samples (Fig. ~~??~~4) suggesting an origin from textile garments (Acharya et al., 2021). The most common colours used in the clothing are black, orange, navy and grey (Table A2). These colours are consistent with those found in snow samples, with 55% of the particles being of a dark colour (black, blue, navy; Fig. 4b). Throughout some regions of Antarctica, polyamide flags are used to identify safe routes for travel and are visible in harsh weather conditions. Weathering of wayfinding flags may release microplastics into the environment, and due to the high volume used each year could have the potential to impact the surrounding environment. The flags are predominantly red, green, blue and black, with weathering processes resulting in a dulling of the flag colour to a pale pink or blue (Fig. 6). In this study, blue was the most common colour detected (55%) followed by pink (23%, Fig. 4b). Replacing the polyamide fabric used for wayfinding flags with an alternative non-synthetic material could reduce the local impact of microplastics on the environment caused by research stations in Antarctica.

Wastewater treatment plants (WWTPs) have been identified as a source of entry of microplastics to the environment worldwide (Browne et al., 2011). An excess of 700,000 synthetic fibres are released from an average 6 kg load of washing acrylic fibres (Napper and Thompson, 2016) and with tumble dryers being present these may also contribute to the presence of microplastics (Tao et al., 2022). The WWTP at Zucchelli Station has been hypothesised as a potential source of microplastic fibres into the Ross Sea while identifying microplastics in nearby benthic sediment (Cincinelli et al., 2017), however no WWTPs in Antarctica have been investigated for their contribution of microplastics to the environment to date. The Protocol on Environmental Protection to the Antarctic Treaty (Annex III and Annex IV) entered into force in 1998. The Protocol prohibits plastic waste from being released by ocean vessels within the Antarctic Treaty area and untreated sewage within 12

nautical miles of land or ice shelves. However, no annex for the release of effluent from research stations was established.

330 WWTPs are not mandatory at stations in Antarctica, leaving many research stations with insufficient wastewater treatment facilities (Gröndahl et al., 2009; Stark et al., 2016). The findings of the Antarctic Treaty Consultative Meeting (ATCM) in 2019 recommended that all governments with Antarctic interests eliminate products containing microplastic beads, work towards reducing microplastic release from wastewater systems and support monitoring of plastic pollution regarding human activity.

335 Future work should focus on quantifying the contribution of tumble dryers and wastewater discharge to the abundance of microplastics in Antarctica, as well as the most effective wastewater treatment process(es) for microplastic removal, which could ultimately be used at these bases.

4.3.2 Long-range transport of microplastics

The transportation of particulates, such as dust, across the Southern Ocean and into Antarctica from other continents has been explored in previous literature, identifying mid-latitude circumpolar westerly winds dominating atmospheric transport to these

340 regions, with Patagonia and New Zealand the most likely sources (Neff and Bertler, 2015). Transportation of microplastics into the Southern Ocean via deep water processes can also introduce synthetic pollutants from afar (Mountford and Maqueda, 2020). Atmospheric long-range transport has been identified as a contamination source of persistent organic pollutants into Antarctica (Kallenborn et al., 2013) as well as dust originating from Australia, Patagonia and the Northern Hemisphere (Li et al., 2008).

345 Short-range transport of microplastics from the bases to ~~the sampling sites~~ sampling sites close by (e.g. S14-S19) is more likely than long-range transport, given the sites' proximity to research bases and the climatology of the area. Yet sites further away may have more influence from long-range transportation, showing the potential influence of both short range and long range inputs. HYSPLIT trajectory modelling indicates that micro-plastics may have travelled from the Amundsen or Ross Seas to reach the remote sample sites and could possibly have come as far as the Weddell Sea (based on hypothesised

350 particle residence time in Brahney et al. (2021)). There are no significant anthropogenic sources identified along the trajectories of the majority of the air masses, such as other research stations, with the Ross Ice Shelf extending for approximately 800 km westward from the sampling sites.

Microplastic particles may have originated from local anthropogenic sources and been transported around Antarctica via a cycle of entrainment and deposition. Alternatively, microplastics may have originated from surface waters surrounding Antarc-

355 tica via co-emission with sea spray (Allen et al., 2020) and atmospheric transport. Given the air mass trajectories pass through the Ross Sea, Amundsen Sea and potentially the Weddell Sea, these are all possible sources with known presences of plastics and microplastics at these locations (Cincinelli et al., 2017; Munari et al., 2017; Van Cauwenberghe et al., 2013; Barnes et al., 2010). Recent atmospheric transport modelling indicates that Antarctica is a net importer of microplastics, with the flux of microplastics from mismanaged plastic waste in the ocean transferring to the atmosphere at the Antarctic coast likely exceeding

360 anthropogenic sources of microplastics on the continent (Brahney et al., 2021).

4.4 Implications and outlook

The implications of microplastics reaching remote regions such as Antarctica are vast. Antarctic organisms have adapted to extreme environmental conditions over many millions of years (Peck, 2018) and the rapid environmental changes due to anthropogenic influence is threatening the unique ecosystems present in the polar regions (Aronson et al., 2011; Convey and
365 Peck, 2019; Lee et al., 2017). Organisms' exposure to microplastics can lead to limited growth, negative effects on reproduction and impaired biological functions (Foley et al., 2018), increasing pressure on ecosystems. Epiplastic communities of bacteria, microalgae and invertebrates can form on environmental plastic particles, which may contribute to the movement and introduction of invasive species into the Antarctic region (Lacerda et al., 2019, 2020).

Negative effects of plastic pollution in Antarctic waters have been reported since 1990 when anthropogenic products, mainly
370 from fishing vessels, were found to be entangling fur seals (Croxall et al., 1990). The ingestion of microplastics by zooplankton, an ecologically important organism in marine ecosystems, can disrupt usual biological processes and negatively impact upon function and health (Cole et al., 2013). The ingestion of microplastics by Antarctic krill may cause a multitude of negative effects on the entire Antarctic food chain as they are a keystone species which a large number of organisms rely on to survive in the Southern Ocean (Hill et al., 2006). Plastic ingestion by the common Antarctic collembolan (*Cryptopygus antarcticus*)
375 has also been identified, indicating the presence of plastic in the Antarctic terrestrial food chain (Bergami et al., 2020).

The consumption of microplastics by higher predators in the Antarctic ecosystem has also been noted, with gentoo (*Pygoscelis papua*), Adélie (*Pygoscelis adeliae*), chinstrap (*Pygoscelis antarcticus*) and king (*Aptenodytes patagonicus*) penguins showing the presence of microplastics in their diet (Fragão et al., 2021; Le Guen et al., 2020). Atmospheric transport of microplastics increases availability to terrestrial organisms, with the potential for ingestion and consequently, negative health
380 effects. Endemic Antarctic penguin species could be exposed to microplastics during their breeding season which is spent on Antarctic fast ice with their colonies. The compounding effects of anthropogenic changes are putting Emperor penguin species at risk with current models predicting a population decline of 81% by 2100 (Jenouvrier et al., 2019).

Microplastics are an emerging contaminant for which appropriate controls and regulations have not been widely put in place (Waller et al., 2017). Members of the Antarctic Treaty committed to reducing plastic pollution in Antarctica and the
385 Southern Ocean at the Antarctic Treaty Consultative Meeting in 2019, urging Antarctic activities to: reduce their use of plastic care products containing microplastics, reduce release of microplastics from WWTP and support greater monitoring of plastic pollution in the region. As microplastics continue to pose a growing threat to the Antarctic ecosystem more rapid approaches are required throughout the Antarctic Treaty System to minimise the widespread impacts. Further research is required to determine the inputs of microplastics into the Antarctic environment, as well as developing a greater understanding of the role
390 of long-range transport for microplastic distribution.

The findings of this study highlight the global reach of plastic pollution and identifies the need for urgency in creating successful policy to reduce its extent and effects, both globally and locally. While studies in the Antarctic region currently focus on marine microplastic pollution, future research and policy needs to take a holistic approach to incorporate airborne and terrestrial impacts. The Protocol on Environmental Protection to the Antarctic Treaty (1991) aims to promote the protection

395 of the Antarctic environment and its place in the world as a natural reserve devoted to peace and science. Growing rates of pollution across the world make this a much greater challenge, and huge transdisciplinary efforts are required to ensure this can continue to be achieved.

5 Conclusions

This study confirms the presence of microplastics in Antarctic snow. Microplastics were identified on the Ross Ice Shelf and near Scott Base and McMurdo Station at an average concentration of 29 particles L⁻¹. Fibres were the most common morphotype identified, and PET was the most common polymer found. Outdoor clothing and other equipment used at the nearby research stations was catalogued to understand local inputs of microplastics, while also assessing the potential for long-range transport through back-trajectory modelling. Back-trajectories indicated that microplastics may have travelled a distance of up to 6000 km, depending on the residence time. Given that air masses passed over the bases prior to sampling, and that the polymers we identified are consistent with those catalogued, it is likely that ~~the majority of identified microplastics originated from local inputs from surrounding research stations~~ local inputs were a contributor to the microplastic pollution identified. Our results highlight the importance for further monitoring – both in Antarctica and in surrounding waters – to develop our understanding of the microplastic footprint in Antarctica and the threat they may pose to the Antarctic environment.




Figure 4. Polymer types identified across all samples. (a) Number of each polymer type found across all Antarctic study sites (PET: polyethylene terephthalate, CP: copolymers, PMMA: polymethyl methacrylate, PVC: polyvinyl chloride, PA: polyamide, other: (polytetrafluoroethylene (PTFE), polyvinylidene, polypropylene, silicone and polymethyl anhydride.), PE: polyethylene, ALK: alkyds, CN: cellulose nitrate); (b) Number of microplastic fragments and fibres identified in each colour category (films ($n=1$) excluded); (c) Size distribution of microplastics across all samples categorized by morphotype (length).




Figure 5. Back-trajectory ensembles generated by HYSPLIT for six of the collected samples. Each of these trajectories is run for 156 hours with the ensemble members generated by minor perturbations as described in section 2.6. Orange points indicate the locations of active Antarctic research stations that operate all year. All of these trajectories are started at 2000 m altitude but show very similar results to those started at 500 m and 1000 m. Note that each simulation is run to correspond to the time of sampling so is host to different meteorological conditions than other simulations.



Figure 6. Marker flags used in Antarctica at McMurdo Station. Before being placed outside (left) and after they were retrieved (right). Photo credit: Evan Thompson.




Figure A1. As for Fig. 5, but trajectories were initialised with a starting height of 500 m.




Figure A2. As for Fig. 5, but trajectories were initialised with a starting height of 1000 m.

Garment	Material composition	Notes
Base layers (tops and bottoms)	100% merino wool	Elasticated waste band and stitching likely to be nylon
Fleece shirt (full zip)	100% polyester	Colour: black
Fleece pants	100% polyester	Colour: black
Salopettes	100% nylon	Colour: navy and black
Softshell jacket	Outside: 100% nylon. Inside: 100% polyester.	Colour: orange and black
Primaloft jacket	Outer: 100% nylon Liner: 100% nylon. Filling: 100% polyester fibre.	Colour: orange and black
ECW jacket	Outer shell: 85% polyester, 15% cotton. Lining: 100% nylon. Filling: 80% duck down, 20% small feather. Fur trim: 90% acrylic, 10% polyester.	
Carhartt jacket	Shell: 100% cotton. Lining: 100% nylon. Interlining: 100% polyester.	Colour: navy
Windproof cap	Shell: 100% nylon. Lining: 100% polyester.	Colour: navy and grey
Beanie	100% polyester	Colour: black and orange
Balaclava	100% polyester	Colour: black and orange
Neck gaiter	100% polyester	Colour: grey
Goggles	Foam lined, plastic body	
ECW mitts	Outer: 94% nylon, 6% spandex. Outer back: 100% polyester. Palm: 100% leather. Shell liner: 100% polyester. Removable liner: 68% polyester, 32% olefin.	
Kinco gloves	Leather	

Fleece gloves	Shell: 96% polyester, 4% spandex. Palm: 50% nylon, 50% PU
Wool gloves/mitts	100% wool
Merino gloves	100% merino
Polypro gloves	Polypropylene
ECW boots	Synthetic and leather uppers
Sorel boots	Natural rubber and leather
Boot liners	100% synthetic
Socks	100% wool

Table A1: Catalogued material used in New Zealand Antarctic program gear.

Site	Morphotype	Colour	Size (µm)	Polymer	Volume (mL)
S1	fibre	blue	630	PET	207
	fragment	blue	226	acrylic copolymer	
	fragment	pink	326	acrylic melamine copolymer	
	fibre	blue	318	acrylic epoxy resin copolymer	
S2	fibre	blue	541	polyamide	250
S3	fibre	blue	1330	polyamide	233
	fibre	blue	885	PET	
	fibre	clear	256	PMMA	
	fibre	blue	1034	PET	
	fibre	blue	500	PET	
	fibre	blue	700	PET	
	fibre	red	139	PA	
	fibre	clear	1519	PMMA	
S4	fibre	blue	273	polyvinyl acetate copolymer	192
	fibre	clear	457	PET	
	fragment	blue	166	PE	
	fragment	blue	168	silicone	
	fragment	pink	130	alkyds	
	fragment	blue	90	polyethylene	
	fragment	pink	221	polyethylene	
S5	fibre	blue	1224	PET	216
	fibre	blue	883	PET	
	fibre	blue	419	PET	
	fibre	blue	329	PET	
	fragment	pink	313	PET	
	fragment	pink	144	PMMA	
S6	fragment	pink	405	polypropylene	208
	fibre	pink	252	PMMA	

	fibre	blue	255	polyamide	
	fibre	clear	544	PMMA	
S7	fibre	blue	519	polyvinylidene	184
	fragment	pink	225	PET	
	fibre	blue	751	PET	
	fragment	pink	189	polyethylene	
	fibre	clear	550	polyester copolymer	
	fibre	blue	905	PET	
	fibre	pink	3510	polyamide	
	fibre	clear	1534	PET	
	fragment	blue	89	cellulose nitrate	
	fragment	pink	215	acrylate copolymer	
S8	fibre	blue	2453	PET	221
	fragment	blue	744	acrylic copolymer	
	fragment	blue	981	acrylic copolymer	
	fibre	blue	660	acrylic copolymer	
	fibre	pink	258	acrylic copolymer	
S9	fragment	white	519	cellulose nitrate	203
S10	fibre	pink	185	polyamide	191
	fragment	blue	77	PMMA	
	fragment	blue	63	PMMA	
S11	fibre	clear	396	PET	235
	fragment	purple	85	PET	
S12	fibre	clear	889	PET	226
	fibre	clear	498	PTFE	
S13	fibre	blue	617	polyamide copolymer	166
	fibre	blue	1388	cellulose nitrate	
	fibre	blue	730	polyamide	
	fragment	pink	314	PET	
	fibre	pink	468	PET	
	fragment	pink	258	acrylic copolymer	

S14	film	blue	1497	PET	219
	fragment	green	199	alkyds	
	fibre	blue	1092	PET	
	fibre	blue	1453	PET	
	fibre	blue	1647	alkyds	
	fragment	pink	135	PMMA	
	fragment	blue	142	acrylic copolymer	
	fibre	pink	2003	PET	
	fibre	blue	1131	cellulose nitrate	
	fibre	clear	1841	PET	
S15	fibre	clear	960	PET	217
	fibre	blue	2005	PMMA	
	fibre	pink	983	PET	
S16	fibre	pink	726	PET	146
	fibre	blue	1322	methyl vinyl ether	
	fibre	pink	415	PET	
	fragment	blue	208	PVC	
	fragment	green	180	alkyds	
	fibre	blue	722	PET	
	fibre	blue	200	styrene copolymer	
	fibre	clear	300	PET	
	fibre	clear	1379	PET	
	fibre	clear	1462	PET	
	fibre	blue	421	poly(methacrylic an- hydride)	
	fragment	green	346	acrylic copolymer	
S17	fibre	blue	400	PET	156
	fragment	blue	50	PET	
	fragment	blue	68	PMMA	
	fragment	grey	336	PTFE	
	fragment	grey	63	PFTE	
	fragment	green	121	acrylic copolymer	
S18	fragment	blue	110	PVC	160

	fibre	blue	1458	PET	
	fibre	clear	484	PET	
	fibre	blue	1327	PET	
	fragment	blue	123	PVC	
	fibre	pink	467	PET	
	fibre	blue	465	PET	
	fibre	blue	404	PET	
	fragment	blue	128	PVC	
S19	fibre	blue	462	PET	182
	fibre	blue	901	PVC	
	fibre	pink	283	PET	
	fragment	pink	171	PET	
	fragment	pink	118	PET	
	fragment	blue	164	acrylic copolymer	
	fragment	blue	113	PVC	
	fragment	blue	116	PVC	
	fragment	blue	121	PVC	
	fragment	blue	165	PVC	

Table A2: Characteristics of microplastics identified [and sample volume](#). As discussed in the text, ‘blue’ includes blue, black and navy. ‘Size’ indicates the width for fragments and length for fibres.

Site number	6-hour distance (km)	24-hour distance (km)	156-hour distance (km)
S1	202	849	5737
S2	173	752	5972
S3	77	328	5006
S4	77	328	4985
S5	87	302	4660
S6	305	681	4004
S7	346	1003	4172
S8	285	1010	4019
S9	281	1060	4243
S10	285	1009	4127
S11	305	1243	5316
S12	43	321	4540
S13	45	321	4520
S14	52	323	4780
S15	60	320	4723
S16	65	315	5026
S17	65	315	4777
S18	68	304	4892
S19	68	304	4882

Table A3. Mean distance travelled by air masses for trajectory ensembles at each of the sites, at 6 hours, 24 hours and 156 hours prior to sampling. Trajectories were initialised at a starting height of 2000 m; starting heights of 1000 m and 500 m yielded similar results.

410 *Author contributions.* Conceptualization: LER and SG. Methodology: LER, HR and SG. Field sampling: ARA, ML, AJM. Investigation: ARA, NEW, AS. Visualization: ARA, NEW, ML, AS. Supervision: LER, SG. Writing – original draft: ARA, LER. Writing – review and editing: ARA, LER, SG, HR, AS, NEW, ML, AJM.

Data availability. **The microplastic data generated in this study has been provided in the appendix of this manuscript, including microplastic counts, sample volume, particle size, shape and polymer type. Relevant data to evaluate the conclusions of this paper**
415 **are present in either the main paper, the appendix, or the supporting information provided.**

Competing interests. The authors declare no competing interests.

Acknowledgements. The authors thank Nicole Lauren-Manuera, Alex Nicholls, Paula Brooksby, Justin Harrison, Antarctica New Zealand, staff and students from Gateway Antarctica and the Postgraduate Certificate of Antarctic Studies 2019 group. LER and SG were supported by the Royal Society of New Zealand Marsden Fund (contract number MFP-UOC1903). ARA was supported by Gateway Antarctica's Ministry of Foreign Affairs and Trade Scholarship in Antarctic and Southern Ocean Studies. We acknowledge mana whenua, Ngāi Tūāhuriri, on whose lands our analysis and writing took place.

References

- Absher, T. M., Ferreira, S. L., Kern, Y., Ferreira, A. L., Christo, S. W., and Ando, R. A.: Incidence and identification of microfibers in ocean waters in Admiralty Bay, Antarctica, *Environmental Science and Pollution Research*, 26, 292–298, 2019.
- 425 Acharya, S., Rumi, S. S., Hu, Y., and Abidi, N.: Microfibers from synthetic textiles as a major source of microplastics in the environment: A review, *Textile Research Journal*, p. 0040517521991244, 2021.
- Allen, S., Allen, D., Phoenix, V. R., Roux, G. L., Jiménez, P. D., Simonneau, A., Binet, S., and Galop, D.: Atmospheric transport and deposition of microplastics in a remote mountain catchment, *Nature Geoscience*, 12, 339–344, <https://doi.org/10.1038/s41561-019-0335-5>, 2019.
- 430 Allen, S., Allen, D., Moss, K., Le Roux, G., Phoenix, V. R., and Sonke, J. E.: Examination of the ocean as a source for atmospheric microplastics, *PloS one*, 15, e0232746, 2020.
- Ambrosini, R., Azzoni, R. S., Pittino, F., Diolaiuti, G., Franzetti, A., and Parolini, M.: First evidence of microplastic contamination in the supraglacial debris of an alpine glacier, *Environmental Pollution*, 253, 297–301, 2019.
- Aronson, R. B., Thatje, S., McClintock, J. B., and Hughes, K. A.: Anthropogenic impacts on marine ecosystems in Antarctica, *Annals of the*
435 *New York Academy of Sciences*, 1223, 82–107, 2011.
- Barnes, D. K., Walters, A., and Gonçalves, L.: Macroplastics at sea around Antarctica, *Marine environmental research*, 70, 250–252, 2010.
- Bergami, E., Rota, E., Caruso, T., Birarda, G., Vaccari, L., and Corsi, I.: Plastics everywhere: first evidence of polystyrene fragments inside the common Antarctic collembolan *Cryptopygus antarcticus*, *Biology letters*, 16, 20200093, 2020.
- Bergmann, M., Mützel, S., Pimpke, S., Tekman, M. B., Trachsel, J., and Gerdt, G.: White and wonderful? Microplastics prevail in snow
440 from the Alps to the Arctic, *Science Advances*, 5, eaax1157, <https://doi.org/10.1126/sciadv.aax1157>, 2019.
- Brahney, J., Hallerud, M., Heim, E., Hahnenberger, M., and Sukumaran, S.: Plastic rain in protected areas of the United States, *Science*, 368, 1257–1260, <https://doi.org/10.1126/science.aaz5819>, 2020.
- Brahney, J., Mahowald, N., Prank, M., Cornwell, G., Klimont, Z., Matsui, H., and Prather, K. A.: Constraining the atmospheric limb of the plastic cycle, *Proceedings of the National Academy of Sciences*, 118, e2020719118, <https://doi.org/10.1073/pnas.2020719118>, 2021.
- 445 Brander, S. M., Renick, V. C., Foley, M. M., Steele, C., Woo, M., Lusher, A., Carr, S., Helm, P., Box, C., Cherniak, S., et al.: Sampling and quality assurance and quality control: a guide for scientists investigating the occurrence of microplastics across matrices, *Applied Spectroscopy*, 74, 1099–1125, 2020.
- Brett, G., Irvin, A., Rack, W., Haas, C., Langhorne, P., and Leonard, G.: Variability in the distribution of fast ice and the sub-ice platelet layer near McMurdo Ice Shelf, *Journal of Geophysical Research: Oceans*, 125, e2019JC015678, 2020.
- 450 Bridson, J. H., Patel, M., Lewis, A., Gaw, S., and Parker, K.: Microplastic contamination in Auckland (New Zealand) beach sediments, *Marine Pollution Bulletin*, 151, 110867, 2020.
- Browne, M. A., Crump, P., Niven, S. J., Teuten, E., Tonkin, A., Galloway, T., and Thompson, R.: Accumulation of microplastic on shorelines worldwide: sources and sinks, *Environmental science & technology*, 45, 9175–9179, 2011.
- Bullard, J. E., Ockelford, A., O'Brien, P., and Neuman, C. M.: Preferential transport of microplastics by wind, *Atmospheric Environment*,
455 245, 118038, 2021.
- Cai, L., Wang, J., Peng, J., Tan, Z., Zhan, Z., Tan, X., and Chen, Q.: Characteristic of microplastics in the atmospheric fallout from Dongguan city, China: preliminary research and first evidence, *Environmental Science and Pollution Research*, 24, 24928–24935, <https://doi.org/10.1007/s11356-017-0116-x>, 2017.

Cincinelli, A., Scopetani, C., Chelazzi, D., Lombardini, E., Martellini, T., Katsoyiannis, A., Fossi, M. C., and Corsolini, S.: Microplastic
460 in the surface waters of the Ross Sea (Antarctica): occurrence, distribution and characterization by FTIR, *Chemosphere*, 175, 391–400,
2017.

Coggins, J. H., McDonald, A. J., and Jolly, B.: Synoptic climatology of the Ross Ice Shelf and Ross Sea region of Antarctica: k-means
clustering and validation, *International journal of climatology*, 34, 2330–2348, 2014.

Cole, M., Lindeque, P., Fileman, E., Halsband, C., Goodhead, R., Moger, J., and Galloway, T. S.: Microplastic ingestion by zooplankton,
465 *Environmental science & technology*, 47, 6646–6655, 2013.

COMNAP: Antarctic Station Catalogue, COMNAP, [https://static1.squarespace.com/static/61073506e9b0073c7eaaf464/t/
611497cc1ece1b43f0eeca8a/1628739608968/COMNAP_Antarctic_Station_Catalogue.pdf](https://static1.squarespace.com/static/61073506e9b0073c7eaaf464/t/611497cc1ece1b43f0eeca8a/1628739608968/COMNAP_Antarctic_Station_Catalogue.pdf), 2017.

Convey, P. and Peck, L. S.: Antarctic environmental change and biological responses, *Science Advances*, 5, eaaz0888, 2019.

Cózar, A., Echevarría, F., González-Gordillo, J. I., Irigoien, X., Úbeda, B., Hernández-León, S., Palma, Á. T., Navarro, S., García-de Lomas,
470 J., Ruiz, A., Fernández-de Puelles, M. L., and Duarte, C. M.: Plastic debris in the open ocean, *Proceedings of the National Academy of
Sciences*, 111, 10 239–10 244, 2014.

Croxall, J., Rodwell, S., and Boyd, I.: Entanglement in man-made debris of Antarctic fur seals at Bird Island, South Georgia, *Marine Mammal
Science*, 6, 221–233, 1990.

Dale, E. R., McDonald, A. J., Coggins, J. H., and Rack, W.: Atmospheric forcing of sea ice anomalies in the Ross Sea polynya region, *The
475 Cryosphere*, 11, 267–280, 2017.

De Frond, H., Rubinovitz, R., and Rochman, C. M.: μ ATR-FTIR Spectral Libraries of Plastic Particles (FLOPP and FLOPP-e) for the
Analysis of Microplastics, *Analytical Chemistry*, 93, 15 878–15 885, 2021.

de Souza Machado, A. A., Kloas, W., Zarfl, C., Hempel, S., and Rillig, M. C.: Microplastics as an emerging threat to terrestrial ecosystems,
Global change biology, 24, 1405–1416, 2018.

480 Dris, R., Gasperi, J., Rocher, V., Saad, M., Renault, N., and Tassin, B.: Microplastic contamination in an urban area: a case study in Greater
Paris, *Environmental Chemistry*, 12, 592–599, <https://doi.org/10.1071/EN14167>, 2015.

Dris, R., Gasperi, J., Saad, M., Mirande, C., and Tassin, B.: Synthetic fibers in atmospheric fallout: a source of microplastics in the environ-
ment?, *Marine pollution bulletin*, 104, 290–293, 2016.

Evangelidou, N., Grythe, H., Klimont, Z., Heyes, C., Eckhardt, S., Lopez-Aparicio, S., and Stohl, A.: Atmospheric transport is a major pathway
485 of microplastics to remote regions, *Nature communications*, 11, 1–11, 2020.

Foley, C. J., Feiner, Z. S., Malinich, T. D., and Höök, T. O.: A meta-analysis of the effects of exposure to microplastics on fish and aquatic
invertebrates, *Science of the Total Environment*, 631, 550–559, 2018.

Fragão, J., Bessa, F., Otero, V., Barbosa, A., Sobral, P., Waluda, C. M., Guímaro, H. R., and Xavier, J. C.: Microplastics and other anthro-
pogenic particles in Antarctica: using penguins as biological samplers, *Science of The Total Environment*, p. 147698, 2021.

490 Fraser, C. I., Morrison, A. K., Hogg, A. M., Macaya, E. C., van Sebille, E., Ryan, P. G., Padovan, A., Jack, C., Valdivia, N., and Waters,
J. M.: Antarctica’s ecological isolation will be broken by storm-driven dispersal and warming, *Nature climate change*, 8, 704–708, 2018.

Ganguly, M. and Ariya, P. A.: Ice Nucleation of Model Nanoplastics and Microplastics: A Novel Synthetic Protocol and the Influence of
Particle Capping at Diverse Atmospheric Environments, *ACS Earth and Space Chemistry*, 3, 1729–1739, 2019.

Geilfus, N.-X., Munson, K., Sousa, J., Germanov, Y., Bhugaloo, S., Babb, D., and Wang, F.: Distribution and impacts of microplastic
495 incorporation within sea ice, *Marine pollution bulletin*, 145, 463–473, 2019.

- González-Pleiter, M., Edo, C., Velázquez, D., Casero-Chamorro, M. C., Leganés, F., Quesada, A., Fernández-Piñas, F., and Rosal, R.: First detection of microplastics in the freshwater of an Antarctic Specially Protected Area, *Marine Pollution Bulletin*, 161, 111–113, 2020.
- Gordon, A. L.: Antarctic polar front zone, *Antarctic Oceanology I*, *Antarctic Res. Ser.*, 15, 205–221, 1971.
- Gründahl, F., Sidenmark, J., and Thomsen, A.: Survey of waste water disposal practices at Antarctic research stations, *Polar Research*, 28, 298–306, 2009.
- Hermabessiere, L., Dehaut, A., Paul-Pont, I., Lacroix, C., Jezequel, R., Soudant, P., and Duflos, G.: Occurrence and effects of plastic additives on marine environments and organisms: A review, *Chemosphere*, 182, 781–793, 2017.
- Hill, S., Murphy, E., Reid, K., Trathan, P., and Constable, A.: Modelling Southern Ocean ecosystems: krill, the food-web, and the impacts of harvesting, *Biological Reviews*, 81, 581–608, 2006.
- Horton, A. A. and Dixon, S. J.: Microplastics: An introduction to environmental transport processes, *Wiley Interdisciplinary Reviews: Water*, 5, e1268, 2018.
- Huiskes, A., Convey, P., and Bergstrom, D.: Trends in Antarctic terrestrial and limnetic ecosystems: Antarctica as a global indicator, pp. 1–13, Springer, 2006.
- Isobe, A., Uchiyama-Matsumoto, K., Uchida, K., and Tokai, T.: Microplastics in the Southern Ocean, *Marine pollution bulletin*, 114, 623–626, 2017.
- Jenouvrier, S., Holland, M., Iles, D., Labrousse, S., Landrum, L., Garnier, J., Caswell, H., Weimerskirch, H., LaRue, M., and Ji, R.: The Paris Agreement objectives will likely halt future declines of emperor penguins, *Global change biology*, 2019.
- Ji, L. N.: Study on preparation process and properties of polyethylene terephthalate (PET), in: *Applied Mechanics and Materials*, vol. 312, pp. 406–410, Trans Tech Publ, 2013.
- Jolly, B., Kuma, P., McDonald, A., and Parsons, S.: An analysis of the cloud environment over the Ross Sea and Ross Ice Shelf using CloudSat/CALIPSO satellite observations: the importance of synoptic forcing, *Atmospheric Chemistry and Physics*, 18, 9723–9739, 2018.
- Kallenborn, R., Breivik, K., Eckhardt, S., Lunder, C. R., Manø, S., Schlabach, M., and Stohl, A.: Long-term monitoring of persistent organic pollutants (POPs) at the Norwegian Troll station in Dronning Maud Land, Antarctica, *Atmospheric Chemistry and Physics*, 13, 6983–6992, 2013.
- Kelly, A., Lannuzel, D., Rodemann, T., Meiners, K., and Auman, H.: Microplastic contamination in east Antarctic sea ice, *Marine Pollution Bulletin*, 154, 111–130, 2020.
- Klein, M. and Fischer, E. K.: Microplastic abundance in atmospheric deposition within the Metropolitan area of Hamburg, Germany, *Science of The Total Environment*, 685, 96–103, <https://doi.org/https://doi.org/10.1016/j.scitotenv.2019.05.405>, 2019.
- Knobloch, E., Ruffell, H., Aves, A., Pantos, O., Gaw, S., and Revell, L. E.: Comparison of Deposition Sampling Methods to Collect Airborne Microplastics in Christchurch, New Zealand, *Water, Air, & Soil Pollution*, 232, 133, <https://doi.org/10.1007/s11270-021-05080-9>, 2021.
- Kroon, F., Motti, C., Talbot, S., Sobral, P., and Puotinen, M.: A workflow for improving estimates of microplastic contamination in marine waters: A case study from North-Western Australia, *Environmental Pollution*, 238, 26–38, 2018.
- Lacerda, A. L. d. F., Rodrigues, L. d. S., Van Sebille, E., Rodrigues, F. L., Ribeiro, L., Secchi, E. R., Kessler, F., and Proietti, M. C.: Plastics in sea surface waters around the Antarctic Peninsula, *Scientific reports*, 9, 1–12, 2019.
- Lacerda, A. L. d. F., Proietti, M. C., Secchi, E. R., and Taylor, J. D.: Diverse groups of fungi are associated with plastics in the surface waters of the Western South Atlantic and the Antarctic Peninsula, *Molecular Ecology*, 2020.

- Le Guen, C., Suaria, G., Sherley, R. B., Ryan, P. G., Aliani, S., Boehme, L., and Brierley, A. S.: Microplastic study reveals the presence of natural and synthetic fibres in the diet of King Penguins (*Aptenodytes patagonicus*) foraging from South Georgia, *Environment international*, 134, 105 303, 2020.
- Lee, J. R., Raymond, B., Bracegirdle, T. J., Chades, I., Fuller, R. A., Shaw, J. D., and Terauds, A.: Climate change drives expansion of Antarctic ice-free habitat, *Nature*, 547, 49–54, 2017.
- Levermore, J. M., Smith, T. E., Kelly, F. J., and Wright, S. L.: Detection of microplastics in ambient particulate matter using Raman spectral imaging and chemometric analysis, *Analytical Chemistry*, 92, 8732–8740, 2020.
- Li, F., Ginoux, P., and Ramaswamy, V.: Distribution, transport, and deposition of mineral dust in the Southern Ocean and Antarctica: Contribution of major sources, *Journal of Geophysical Research: Atmospheres*, 113, 2008.
- Liu, K., Wang, X., Fang, T., Xu, P., Zhu, L., and Li, D.: Source and potential risk assessment of suspended atmospheric microplastics in Shanghai, *Science of the Total Environment*, 675, 462–471, 2019a.
- Liu, K., Wu, T., Wang, X., Song, Z., Zong, C., Wei, N., and Li, D.: Consistent transport of terrestrial microplastics to the ocean through atmosphere, *Environmental science & technology*, 53, 10 612–10 619, 2019b.
- MacLeod, M., Arp, H. P. H., Tekman, M. B., and Jahnke, A.: The global threat from plastic pollution, *Science*, 373, 61–65, 2021.
- Masura, J., Baker, J. E., Foster, G. D., Arthur, C., and Herring, C.: Laboratory methods for the analysis of microplastics in the marine environment: recommendations for quantifying synthetic particles in waters and sediments, 2015.
- Materić, D., Kasper-Giebl, A., Kau, D., Anten, M., Greilinger, M., Ludewig, E., van Sebille, E., Röckmann, T., and Holzinger, R.: Micro- and Nanoplastics in Alpine Snow: A New Method for Chemical Identification and (Semi)Quantification in the Nanogram Range, *Environmental Science & Technology*, 54, 2353–2359, <https://doi.org/10.1021/acs.est.9b07540>, 2020.
- Materić, D., Ludewig, E., Brunner, D., Röckmann, T., and Holzinger, R.: Nanoplastics transport to the remote, high-altitude Alps, *Environmental Pollution*, 288, 117 697, <https://doi.org/10.1016/j.envpol.2021.117697>, 2021.
- Mato, Y., Isobe, T., Takada, H., Kanehiro, H., Ohtake, C., and Kaminuma, T.: Plastic resin pellets as a transport medium for toxic chemicals in the marine environment, *Environmental science & technology*, 35, 318–324, 2001.
- Matsuoka, K., Skoglund, A., Roth, G., de Pomereu, J., Griffiths, H., Headland, R., Herried, B., Katsumata, K., Le Brocq, A., Licht, K., Morgan, F., Neff, P. D., Ritz, C., Scheinert, M., Tamura, T., Van de Putte, A., van den Broeke, M., von Deschwandén, A., Deschamps-Berger, C., Van Liefferinge, B., Tronstad, S., and Melvær, Y.: Quantarctica, an integrated mapping environment for Antarctica, the Southern Ocean, and sub-Antarctic islands, *Environmental Modelling & Software*, 140, 105 015, <https://doi.org/10.1016/j.envsoft.2021.105015>, 2021.
- McConnell, J. R., Maselli, O. J., Sigl, M., Vallelonga, P., Neumann, T., Anshütz, H., Bales, R. C., Curran, M. A., Das, S. B., Edwards, R., et al.: Antarctic-wide array of high-resolution ice core records reveals pervasive lead pollution began in 1889 and persists today, *Scientific Reports*, 4, 1–5, 2014.
- Mountford, A. S. and Maqueda, M. M.: Modelling the accumulation and transport of microplastics by sea ice, *Journal of Geophysical Research: Oceans*, p. e2020JC016826, 2020.
- Munari, C., Infantini, V., Scoconi, M., Rastelli, E., Corinaldesi, C., and Mistri, M.: Microplastics in the sediments of Terra Nova Bay (Ross Sea, Antarctica), *Marine pollution bulletin*, 122, 161–165, 2017.
- Napper, I. E. and Thompson, R. C.: Release of synthetic microplastic plastic fibres from domestic washing machines: Effects of fabric type and washing conditions, *Marine pollution bulletin*, 112, 39–45, 2016.

- 570 Neff, P. D. and Bertler, N. A.: Trajectory modeling of modern dust transport to the Southern Ocean and Antarctica, *Journal of Geophysical Research: Atmospheres*, 120, 9303–9322, 2015.
- Parish, T. R., Cassano, J. J., and Seefeldt, M. W.: Characteristics of the Ross Ice Shelf air stream as depicted in Antarctic Mesoscale Prediction System simulations, *Journal of Geophysical Research: Atmospheres*, 111, 2006.
- Peck, L. S.: Antarctic marine biodiversity: adaptations, environments and responses to change, *Oceanography and Marine Biology*, 2018.
- 575 Primpke, S., Wirth, M., Lorenz, C., and Gerdts, G.: Reference database design for the automated analysis of microplastic samples based on Fourier transform infrared (FTIR) spectroscopy, *Analytical and bioanalytical chemistry*, 410, 5131–5141, 2018.
- Reed, S., Clark, M., Thompson, R., and Hughes, K. A.: Microplastics in marine sediments near Rothera research station, Antarctica, *Marine pollution bulletin*, 133, 460–463, 2018.
- Revell, L. E., Kuma, P., Le Ru, E. C., Somerville, W. R. C., and Gaw, S.: Direct radiative effects of airborne microplastics, *Nature*, 598, 462–467, <https://doi.org/10.1038/s41586-021-03864-x>, 2021.
- 580 Rios, L. M., Moore, C., and Jones, P. R.: Persistent organic pollutants carried by synthetic polymers in the ocean environment, *Marine Pollution Bulletin*, 54, 1230–1237, 2007.
- Ryan, P. G.: A brief history of marine litter research, pp. 1–25, Springer, Cham, 2015.
- Seefeldt, M. W. and Cassano, J. J.: A description of the Ross Ice Shelf air stream (RAS) through the use of self-organizing maps (SOMs), *Journal of Geophysical Research: Atmospheres*, 117, 2012.
- 585 Shim, W. J., Hong, S. H., and Eo, S. E.: Identification methods in microplastic analysis: a review, *Analytical methods*, 9, 1384–1391, 2017.
- Stark, J. S., Corbett, P. A., Dunshea, G., Johnstone, G., King, C., Mondon, J. A., Power, M. L., Samuel, A., Snape, I., and Riddle, M.: The environmental impact of sewage and wastewater outfalls in Antarctica: An example from Davis station, East Antarctica, *Water research*, 105, 602–614, 2016.
- 590 Stein, A., Draxler, R. R., Rolph, G. D., Stunder, B. J., Cohen, M., and Ngan, F.: NOAA’s HYSPLIT atmospheric transport and dispersion modeling system, *Bulletin of the American Meteorological Society*, 96, 2059–2077, 2015.
- Suaria, G., Perold, V., Lee, J. R., Lebouard, F., Aliani, S., and Ryan, P. G.: Floating macro-and microplastics around the Southern Ocean: Results from the Antarctic Circumnavigation Expedition, *Environment International*, 136, 105494, 2020.
- Tao, D., Zhang, K., Xu, S., Lin, H., Liu, Y., Kang, J., Yim, T., Giesy, J. P., and Leung, K. M.: Microfibers Released into the Air from a Household Tumble Dryer, *Environmental Science & Technology Letters*, 2022.
- 595 Tin, T., Lamers, M., Liggett, D., Maher, P. T., and Hughes, K. A.: Setting the scene: human activities, environmental impacts and governance arrangements in Antarctica, pp. 1–24, Springer, 2014.
- Tin, T., Summerson, R., and Yang, H. R.: Wilderness or pure land: tourists’ perceptions of Antarctica, *The Polar Journal*, 6, 307–327, 2016.
- Van Cauwenberghe, L., Vanreusel, A., Mees, J., and Janssen, C. R.: Microplastic pollution in deep-sea sediments, *Environmental pollution*, 182, 495–499, 2013.
- 600 Vandermeersch, G., Van Cauwenberghe, L., Janssen, C. R., Marques, A., Granby, K., Fait, G., Kotterman, M. J., Diogène, J., Bekaert, K., Robbens, J., et al.: A critical view on microplastic quantification in aquatic organisms, *Environmental Research*, 143, 46–55, 2015.
- Waller, C. L., Griffiths, H. J., Waluda, C. M., Thorpe, S. E., Loaiza, I., Moreno, B., Pachterres, C. O., and Hughes, K. A.: Microplastics in the Antarctic marine system: an emerging area of research, *Science of the Total Environment*, 598, 220–227, 2017.
- 605 Williamson, C. E., Neale, P. J., Hylander, S., Rose, K. C., Figueroa, F. L., Robinson, S. A., Häder, D.-P., Wängberg, S., and Worrest, R. C.: The interactive effects of stratospheric ozone depletion, UV radiation, and climate change on aquatic ecosystems, *Photochemical & Photobiological Sciences*, 18, 717–746, 2019.

- Wright, S. L., Thompson, R. C., and Galloway, T. S.: The physical impacts of microplastics on marine organisms: a review, *Environmental pollution*, 178, 483–492, 2013.
- 610 Zhang, Y., Gao, T., Kang, S., Allen, S., Luo, X., and Allen, D.: Microplastics in glaciers of the Tibetan Plateau: Evidence for the long-range transport of microplastics, *Science of The Total Environment*, 758, 143 634, <https://doi.org/10.1016/j.scitotenv.2020.143634>, 2021.