

Controls on Greenland moulin geometry and evolution from the Moulin Shape model

Lauren C. Andrews¹, Kristin Poinar^{2,3}, Celia Trunz⁴

¹Global Modeling and Assimilation Office, NASA Goddard Space Flight Center, Greenbelt, MD, 20771, USA

²Department of Geology, University at Buffalo, Buffalo, NY, 14260, USA

³Research and Education in eNergy, Environment and Water (RENEW) Program, University at Buffalo, Buffalo, NY, 14260, USA

⁴Geosciences Department, University of Arkansas, Fayetteville, AR, 72701, USA

Correspondence: Lauren C. Andrews (lauren.c.andrews@nasa.gov)

Abstract. Nearly all meltwater from glaciers and ice sheets is routed englacially through moulins, ~~which collectively comprise approximately 10–14% of the efficient englacial–subglacial hydrologic system.~~ Therefore, the geometry and evolution of moulins has the potential to influence subglacial water pressure variations, ice motion, and the runoff hydrograph delivered to the ocean. We develop the *Moulin Shape* (MouSh) model, a time-evolving model of moulin geometry. MouSh models ice deformation around a moulin using both viscous and elastic rheologies and models melting within the moulin through heat dissipation from turbulent water flow, both above and below the water line. We force MouSh with idealized and realistic surface melt inputs. Our results show that variations in surface melt change the geometry of a moulin by approximately ~~30~~20% daily and by over 100% seasonally. These size variations cause observable differences in moulin water storage capacity, moulin water levels, and subglacial channel size compared to a static, cylindrical moulin. Our results suggest that moulins are ~~significant~~important storage reservoirs for meltwater, with storage capacity and water levels varying over multiple timescales. Representing moulin geometry within subglacial hydrologic models would therefore improve ~~their accuracy~~the representation of subglacial pressures, especially over seasonal periods or in regions where overburden pressures are high.

1 Introduction

~~Surface-sourced meltwater delivered to the glacier bed influences the evolution of the subglacial hydrologic system and associated subglacial pressures (e.g., Iken and Bindshadler, 1986; Müller and Iken, 1973). The efficiency of the subglacial system, in turn, changes the flow patterns of the overlying ice on daily, seasonal, and multi-annual timescales (e.g., Hoffman et al., 2011; Iken and Bindshadler, 1986; Moon et al., 2014; Tedstone et al., 2015; Williams et al., 2020). Thus, glacial hydrology is a crucial factor in short-term calculations of mass loss on glaciers and ice sheets (Bell, 2008; Flowers, 2018). On the Greenland Ice Sheet, surface meltwater can take multiple paths, depending on its spatial origin. In the accumulation zone,~~

meltwater may percolate through snow and firn, remaining liquid (Forster et al., 2014) or refreezing (MacFerrin et al., 2019). In the ablation zone, meltwater runs over bare ice, coalesces into supraglacial streams, and pools into supraglacial lakes (e.g., Smith et al., 2015). These surficial water features—rivers, streams, lakes, aquifers, etc.—direct meltwater into englacial features that can deliver the water to the bed of the ice sheet (Andrews et al., 2014; Das et al., 2008; Miège et al., 2016; Poinar et al., 2017; Smith et al., 2015). Englacial features include moulins, which are near-vertical shafts with large surface catchments (~1–5 km² per moulin, Banwell et al., 2016; Colgan and Steffen, 2009; Yang and Smith, 2016), and crevasses, which are linear features with limited local catchments (~0.05 km² per crevasse, Poinar et al., 2017). Together, moulins and crevasses constitute the englacial hydrologic system.

Water fluxes through the englacial system, and therefore to the subglacial system, are non-uniform in space and time. Variations in subglacial water fluxes can modify the form of the subglacial hydrologic system and influence ice motion on diurnal to multi-year timescales (Hoffman et al., 2011; Tedstone et al., 2015). Quantifying these temporal variations in water fluxes to the glacier bed requires understanding the time evolution of the supraglacial and englacial water systems that deliver it. Ongoing research is making great strides in characterizing the supraglacial water network (Germain and Moorman, 2019; Smith et al., 2015; Yang et al., 2016). For instance, field observations from Greenland indicate that much of the supraglacial water network terminates into crevasses and moulins (Colgan et al., 2011a; McGrath et al., 2011; Poinar et al., 2015; Smith et al., 2015) and that these englacial features are important modulators of surface melt inputs to the ice sheet bed (Andrews et al., 2014; Cowton et al., 2013).

Our knowledge of moulin sizes, scales, and time evolution has largely been informed by exploration and mapping of the top tens to hundred meters of a few moulins (Benn et al., 2017; Covington et al., 2020; Gulley et al., 2009; Holmlund, 1988; Moreau, 2009). These sparse field data indicate that moulin shapes deviate greatly from simple cylinders. Furthermore, deployments of tethered sensors into Greenland moulins have encountered irregularities, including apparent ledges, plunge pools, and constrictions, below the depths of human exploration (Andrews et al., 2014; Covington et al., 2020; Cowton et al., 2013). These direct near-surface and indirect deep observations suggest that moulin geometry evolves a high degree of complexity at all depths.

State-of-the-art subglacial hydrology models are forced by meltwater inputs that enter the system through crevasses or moulins. These models generally represent the geometry of moulins in a simplified and time independent manner, for instance as a static vertical cylinder (e.g., Hewitt, 2013; Hoffman et al., 2016; Werder et al., 2013) or cone (Clarke, 1996; Flowers and Clarke, 2002; Werder et al., 2010). The basis for the cylindrical simplification arises from the assumption that depth-dependent variations in moulin size are small relative to the vertical scale of the moulin. The basis for time independence is the assumption that the moulin capacity is, again, small relative to that of the subglacial system. However, neither of these assumptions have been tested. Here, we explore the extent to which time evolution of moulin geometry affects the rate of subglacial meltwater input and subglacial pressure in channelized regions of the bed.

We present the Moulin Shape (MouSh) model, a new, physically based numeric model that evolves moulin geometry over diurnal and seasonal periods (Fig. 1). Surface-sourced meltwater delivered to the glacier bed drives the evolution of the

subglacial hydrologic system and associated subglacial pressures (e.g., Iken and Bindschadler, 1986; Müller and Iken, 1973). The efficiency of the subglacial system, in turn, changes the flow patterns of the overlying ice on daily, seasonal, and multi-annual timescales (e.g., Hoffman et al., 2011; Iken and Bindschadler, 1986; Moon et al., 2014; Tedstone et al., 2015; Williams et al., 2020). Thus, glacial hydrology is a crucial factor in short-term changes to glacier and ice sheet dynamics (Bell, 2008; Flowers, 2018).

On the Greenland Ice Sheet, surface meltwater can take multiple paths, depending on its origin location. In the accumulation zone, meltwater may percolate through snow and firn, remaining liquid (Forster et al., 2014) or refreezing (MacFerrin et al., 2019). In the ablation zone, meltwater runs over bare ice, coalesces into supraglacial streams, and pools into supraglacial lakes (e.g., Smith et al., 2015). These surficial water features – rivers, streams, lakes, aquifers, etc. – direct meltwater into englacial features that can deliver the water to the bed of the ice sheet (Andrews et al., 2014; Das et al., 2008; Miège et al., 2016; Poinar et al., 2017; Smith et al., 2015). Englacial features include moulins, which are near-vertical shafts with large surface catchments (~1–5 km² per moulin, Banwell et al., 2016; Colgan and Steffen, 2009; Yang and Smith, 2016), and crevasses, which are linear features with limited local catchments (~0.05 km² per crevasse, Poinar et al., 2017). Together, moulins and crevasses constitute a substantial fraction of the englacial hydrologic system.

Water fluxes through the englacial system, and therefore to the subglacial system, are non-uniform in space and time. Quantifying these temporal variations in water fluxes to the glacier bed requires understanding the time evolution of the supraglacial and englacial water systems that deliver it. Ongoing research is making great strides in characterizing the supraglacial water network (Germain and Moorman, 2019; Smith et al., 2017; Yang et al., 2016). For instance, field observations from Greenland indicate that much of the supraglacial water network terminates into crevasses and moulins (McGrath et al., 2011; Smith et al., 2015) and that moulins are important modulators of surface melt inputs to the ice sheet bed (Andrews et al., 2014; Cowton et al., 2013; Mejia et al., 2021).

Our knowledge of moulin sizes, scales, and time evolution has largely been informed by exploration and mapping of the top tens to hundred meters of a few moulins (Benn et al., 2017; Covington et al., 2020; Gulley et al., 2009; Holmlund, 1988; Moreau, 2009). These sparse field data indicate that moulin shapes deviate greatly from simple cylinders. Furthermore, deployments of tethered sensors into Greenland moulins have encountered irregularities including apparent ledges and plunge pools (Andrews et al., 2014; Covington et al., 2020; Cowton et al., 2013), and seismic (Rösli et al., 2016) and radar (Catania et al., 2008) studies suggest constrictions below the depths of human exploration. These direct near-surface and indirect deep observations suggest that moulin geometry evolves a high degree of complexity at all depths.

State-of-the-art subglacial hydrology models are forced by meltwater inputs that enter the system through crevasses or moulins. These models generally represent the geometry of moulins in a simplified and time-independent manner, for instance as a static vertical cylinder (e.g., Hewitt, 2013; Hoffman et al., 2016; Werder et al., 2013) or cone (Clarke, 1996; Flowers and Clarke, 2002; Werder et al., 2010). The basis for the cylindrical simplification arises from the assumption that depth-dependent variations in moulin size are small relative to the vertical scale of the moulin. The basis for time independence is the assumption that the moulin capacity is, again, small relative to that of the subglacial system. However, neither of these

assumptions have been tested. Here, we explore the extent to which time evolution of moulin geometry affects the rate of subglacial meltwater input and subglacial pressure in channelized regions of the bed.

We present the Moulin Shape (MouSh) model, a new, physically based numeric model that evolves moulin geometry over diurnal and seasonal periods. The MouSh model can be coupled to subglacial hydrology models to more completely characterize the time evolution of the englacial and subglacial hydrologic systems, which are intimately linked.

2 Methods

2.1 The role of moulins in Greenland meltwater transport

Modeling of the subglacial hydrologic system has revealed the importance of moulins in the initiation of subglacial channels under both alpine glaciers (Hoffman and Price, 2014) and ice sheets (Werder et al., 2013). Thus, moulins comprise an important component of the overall glacial hydrologic system but are usually omitted or represented simply in subglacial hydrology models.

We estimate the relative path length of moulins within the overall glacial hydrologic system by combining available maps of moulin locations (Andrews, 2015; Hoffman et al., 2018; Smith et al., 2015) and maps of subglacial hydraulic potential (Fig. 2a & 3a-b), following Shreve (1972):

$$\Phi = k\rho_i g(s - b) + \rho_w g b \quad (1)$$

Here, Φ is the subglacial potential; ρ_i and ρ_w are ice and water density, respectively; g is gravitational acceleration; and s and b are the surface and bed elevations, respectively, derived from available datasets (Howat et al., 2014; Morlighem et al., 2017, 2018). k is a spatially uniform parameter that represents the fraction of flotation; that is, the ratio of water pressure and ice overburden pressure, where $k=1$ indicates that water pressure is equal to ice overburden pressure at the bed, and $k=0$ indicates that water is at atmospheric pressure (e.g., Rippin et al., 2003; Shreve, 1972). In reality, k will not be uniform in space or time, nor indicative of whether there are subglacial channels; however, studies suggest that using k in flow routing improves prediction in regions with large subglacial water fluxes that would likely organize into efficient or channelized flow (Chu et al., 2016; Everett et al., 2016; Fried et al., 2015; Palmer et al., 2011). We use $k=1$ to approximate the likely channelized subglacial pathways.

We calculate theoretical flow accumulation via the hydraulic potential gradient in two separate areas of the Greenland Ice Sheet: the Pákitsoq and Russell Glacier regions (Fig. 3). Flow accumulation maps do not directly indicate whether flow is channelized or unchannelized, especially at far inland locations, where efficient subglacial drainage may still occur in the absence of channels (Meierbachtol et al., 2013). Thus, we define a flow accumulation area threshold of $\sim 1 \text{ km}^2$, which is similar to but generally smaller than the size of individual supraglacial drainage basins. As subglacial channels are generally initiated at moulin locations (Werder et al., 2013), this ensures that we do not underestimate potential subglacial channel

lengths. Furthermore, we only interpret subglacial accumulation pathways seaward of the 1,300-meter surface elevation contour as channelized. This threshold is supported by GPS analysis in the Pâkitsoq region (Andrews et al., 2018) and dye tracing in the Russell Glacier region (Chandler et al., 2013). Using this information and the distance between each moulin location and the nearest subglacial channel node, we estimate the total length of subglacial channelization in these two regions. We next extract the ice thickness at each moulin from the BedMachine dataset (Fig. 3e, Morlighem et al., 2018), following Andrews (2015) and Yang and Smith (2016). The total subglacial channel length, including the distance between moulins and the nearest flow pathway plus the sum of moulin depths, are used to estimate the percentage of the total subsurface (englacial-subglacial) path length that moulins comprise.

2.2 Physical moulin Moulin physical model

We develop the Moulin Shape (MouSh) model, a numeric model of moulin evolution that considers ice deformation and ice melt associated with the dissipation of energy from turbulently flowing meltwater (Fig. 1). We include here a detailed description of the model framework and each module that influences the time-evolving geometry of the modeled moulin- (Fig. 2a).

2.2.1 Moulin geometry coordinate system

We discretize our model in the vertical (z) and radial (r_1 and r_2) directions, treating the moulin as a stack of egg-shaped (semi-circular, semi-elliptical) holes in the ice that both change in size and move laterally relative to each other. We calculate moulin geometry (elliptical radii r_1 and r_2) and water level (h_w) with a 5-minute timestep dt . Model calculations are performed in cylindrical coordinates, where $\Pi(z)$ is the perimeter of the semi-circular, semi-elliptical moulin, using Ramanujan's approximation:

$$\Pi \approx \pi r_1 + \frac{1}{2} \pi [3(r_1 + r_2) - \sqrt{(3r_1 + r_2)(r_1 + 3r_2)}] \quad (21)$$

Here, r_1 and r_2 are the major and minor radii, respectively, for each node in the vertical direction. The minor radius r_1 is also the radius of the half-circle.

We calculate the cross-sectional area of the semi-circular, semi-elliptical moulin as follows:

$$A_m = \frac{\pi r_1}{2} (r_1 + r_2) \quad (2)$$

The plan-view orientation of the radii and the coordinate system, as detailed on a remotely sensed moulin, are indicated in Fig. 2-2b-d. The elliptical shape was chosen to reflect the observation that supraglacial meltwater flows into a moulin along a single side above the water line. This asymmetry leads to a nonuniform, noncircular geometry above the water level, which can affect the total amount and evolution of water storage at high water levels. This choice is in line with observations of a

Greenland moulin becoming more elliptical over time (Rössli et al., 2016). For simplicity, MouSh contains an option to set the moulin cross-sectional geometry to a circle, rather than an egg (see Supplement S2).

2- Each module is also dependent on the depth varying hydrostatic and cryostatic pressures. We subtract the cryostatic pressure, P_i , from the hydrostatic pressure, P_w , to get the total depth-dependent effective pressure N at all levels z within the moulin:

$$P_i = \rho_i g (H_i - b) \quad (3a)$$

$$P_w = \rho_w g (h_w - b) \quad (3b)$$

$$N = P_w - P_i \quad (3c)$$

where H_i is the ice thickness; h_w is the height of the water above the bed; z is the vertical coordinate; ρ_i and ρ_w are ice and water density, respectively; and g is gravitational acceleration (Table 1). In this formulation, positive pressures causes outward expansion of the moulin walls (radial growth), and negative pressures reduces the size of the moulin (radial closure). We use a flat bed at sea level for all model runs presented here, so bed elevation $b = 0$.

2.2 Ice deformation modules

We represent the deformation of the ice with the simplest possible combination of elastic and viscous components: a Maxwell rheology, where elastic and viscous deformation occur independently, without interaction (Turcotte and Schubert, 2002). The Maxwell timescale is equal to $(E \times A \times \tau^2)^{-1}$, or roughly 10–100 hours for typical Greenland ice. We approximate this as 1 day (Turcotte and Schubert, 2002). The Maxwell model comprises an elastic element (a spring) and a viscous element (a dashpot) in series and is standard in geophysical modeling. The response timescale in our Maxwell model is equal to $(E \times A \times \tau^2)^{-1}$ where E is Young’s modulus, A is the viscous flow law parameter, and τ is stress (Turcotte and Schubert, 2002). The Maxwell timescale is thus roughly 10–100 hours for typical Greenland ice. On timescales shorter than the Maxwell timescale, ice deformation is primarily elastic. On longer timescales, viscous deformation dominates.

2.2.2.1 Elastic deformation

Field measurements indicate that, nearly universally during the melt season, the water level in a moulin varies at a sub-hourly timescale (Andrews et al., 2014; Covington et al., 2020; Cowton et al., 2013; Iken, 1972). This variability is shorter than, but comparable to, the 1 day Maxwell timescale for ice; therefore, we must assume that elastic deformation plays a role in the response of the ice to variations in moulin water level.

The stress and deformational patterns around a borehole have been well studied in the rock mechanics literature (Amadei, 1983; Goodman, 1989; Priest, 1993). We Elastic deformation is described in Sect. 2.2.1. We represent total viscous deformation in two modes: (1) radial opening and closure of the moulin, which changes the size of the moulin (Sect. 2.2.2), and (2) vertical shear of the moulin, which changes the shape but not the size of the moulin (Sect. 2.2.3).

2.2.1 Elastic deformation

Field measurements indicate that, nearly universally during the melt season, the water level in a moulin varies at a sub-hourly timescale (Andrews et al., 2014; Covington et al., 2020; Cowton et al., 2013; Iken, 1972). This variability is shorter than, but comparable to, the Maxwell timescale for ice (10–100 hours; see Sect. 2.2); therefore, we must assume that elastic deformation plays a role in the response of the ice to variations in moulin water level.

Weertman (1971, 1973, 1996) applied dislocation fracture mechanics principles to vertical glaciological features: water-filled crevasses. These equations have applied to supraglacial lake drainages (Krawczynski et al., 2009) and slow ice hydrofracture (Poinar et al., 2017). However, these problems are Cartesian (linear), not cylindrical, so their solutions are not readily adaptable to a moulin. The stress and deformational patterns around cylindrical boreholes have been well studied in the rock mechanics literature (Amadei, 1983; Goodman, 1989; Priest, 1993). We therefore base our description of the stress field surrounding the moulin on that of a fluid-filled borehole in a porous rock medium, described by Aadnøy (1987) and based on the Kirsch equations, which describe stresses surrounding a circular hole in a rigid plate (Kirsch, 1898). We assume plane strain and approximate our moulin as a stack of such plates with analogous holes (Goodman, 1989). A subtle difference is that our moulin shape is not circular, but egg-shaped: half circular, half elliptical.

At each vertical level z in the moulin, we apply Hooke's Law to the stress field to calculate the strain, in horizontal cross-section, at all points on the moulin wall and in the surrounding ice, for both radii r_1 and r_2 . We then integrate these strains from an infinite distance (cylindrical coordinate $r = \infty$) to the moulin wall ($r_1, r_2 = r_m$). The result is a full derivation, based on the expected stress states in a borehole described by Aadnøy (1987), is in Supplement S1. We express the total radial elastic displacement, Δr_e , of a segment deformation r_e of a moulin with radius r_m segment as:

$$\Delta r_e = \frac{r_m}{E} [(1 + \nu)(PN - \frac{1}{2}(\sigma_x + \sigma_y) + \frac{1}{4}(\sigma_x - \sigma_y)(1 - 3\nu - 4\nu^2) + \frac{1}{4}\tau_{xy}(2 - 3\nu - 8\nu^2))] \quad (4)$$

Here, E is Young's modulus for uniaxial deformation, ν is Poisson's ratio, and σ_x , σ_y , and τ_{xy} are the background deviatoric and shear stresses that describe the regional setting of the moulin (typically compressive and of order 100 kPa). We apply this equation to both moulin radii, the semi-circular radius r_1 and the semi-elliptical radius r_2 , separately. The outward pressure in the moulin P is the difference between the hydrostatic and cryostatic pressures; Poinar and Andrews, 2021). The model is designed to accept user-defined deviatoric and shear stresses; we choose values $\sigma_x = 0$ kPa, $\sigma_y = 50$ kPa, and $\tau_{xy} = -50$ kPa. However, because the deviatoric and shear stresses are poorly constrained, we also test the simplification $\sigma_x = \sigma_y = \tau_{xy} = 0$ (Supplement S2), which simplifies the elastic deformation r_e :

$$P = \rho_w g(h_w - z) = \rho_i g(H_i - z) \quad (5)$$

$$\text{where } h_w r_e = \frac{r_m}{E} (1 + \nu) N \quad (5)$$

Unlike viscous deformation and melting, elastic deformation is the height instantaneous. However, we take advantage of the water above observation that elastic deformation is driven by changes in the cryostatic and hydrostatic pressures. Therefore, we express Eq. 4 and Eq 5 as an elastic ‘deformation rate’ for non-zero (Eq 6a) and zero (Eq 6b) stresses:

$$dr_e = \frac{1}{E} \left(r_m (1 + \nu) \frac{dN}{dt} + [(1 + \nu) \left(N - \frac{1}{2} (\sigma_x + \sigma_y) \right) + \frac{1}{4} (\sigma_x - \sigma_y) (1 - 3\nu - 4\nu^2) + \frac{1}{4} \tau_{xy} (2 - 3\nu - 8\nu^2)] \frac{dr_m}{dt} \right) dt \quad (6)$$

$$dr_e = \frac{(1+\nu)}{E} \left(r_m \frac{dN}{dt} + N \frac{dr_m}{dt} \right) dt \quad (7)$$

Equations 6 and 7 assume that both effective pressure and moulin radius vary smoothly over the bed, H_i is the ice thickness, and Δt time interval in question, which is generally true for small timesteps (5-minutes in our model). We apply Eq. 6 or 7 to both moulin radii, the semi-circular radius r_1 and the vertical coordinate semi-elliptical major radius r_2 , separately. When water is above the flotation level, elastic deformation opens the moulin elastically opens at all depths, below the water line. When the water level is below flotation, which is the typical case, elastic deformation closes the moulin at all depths. The values of the surface stresses σ_x , σ_y , and τ_{xy} determine the sign of the deformation above the water line.

Because we use a flat bed for all model runs presented, we define h_w as ‘moulin water level’, which only includes the hydrostatic water head, as opposed to ‘moulin hydraulic head’, which would include both the hydrostatic and elevation heads.

2.2.2.2 Viscous deformation

Over times longer than the ~ 1 day Maxwell timescale, viscous deformation is the dominant deformation process. We represent total viscous deformation in two modes: (1) radial opening and closure of the moulin, which changes the size of the moulin (Section 2.2.2.2.1), and (2) vertical shear of the moulin, which changes the shape but not the size of the moulin (Section 2.2.2.2.2).

2.2.2.2.1 Radial opening and closure

Moulins close when they lose their water source at the end of a melt season (Catania and Neumann, 2010). Similarly, boreholes close if they are not filled with drilling fluid with a density similar to ice (Alley, 1992). Our modeled moulin is intermediate to these edge cases, since it typically contains water. When the moulin is filled with water to the flotation level, it will stay open at its base and viscously close at and below the water level. When water is above flotation, the moulin will viscously open at all depths. When the water level is below flotation, which is the typical case, viscous deformation shrinks the moulin at all depths.

We sum the glacio-static stress, σ_i , and the hydrostatic stress, σ_w , to get the total depth dependent stress, σ_z , at all levels z within the moulin:

$$\sigma_i = -\rho_i g (H_i - z) \quad (6a)$$

$$\sigma_w = -\rho_w g (h_w - z) \quad (6b)$$

$$\sigma_z = \sigma_i + \sigma_w \quad (6c)$$

In this formulation, positive stress causes outward expansion of the moulin walls (radial growth), and negative stress reduces the size of the moulin (radial closure).

Moulins close when they lose their water source at the end of a melt season (Catania and Neumann, 2010). Similarly, boreholes close if they are not filled with drilling fluid with a density similar to ice (Alley, 1992). Our modeled moulin is intermediate to these edge cases because it typically contains water. When the moulin is filled with water to the flotation level, it will stay open at its base and viscously close at and below the water level. When the water level is above flotation, the moulin will viscously open in regions where hydrostatic pressure exceeds the cryostatic pressure. When the water level is below flotation, which is the typical case, viscous deformation shrinks the moulin at all depths.

We calculate strain rate ($\dot{\epsilon}$) from the total depth-dependent ~~stresseffective pressure~~ N (Eq. 6e3c) using Glen's Flow Law:

$$\dot{\epsilon} = \frac{E A(T, P) \cdot \left(\frac{1}{3} \sigma_z\right)^3}{(F_* A(T_i, P_i) \cdot \left(\frac{1}{3} N\right)^n)} \quad (8)$$

where F_* is ~~an~~ the flow law enhancement factor, and $A(T, P_i, P)$ is the flow law parameter. For the flow law parameter, we use the standard relationship from Cuffey and Paterson (2010, Equation 3.35); (2010, Eq. 3.35), which is a function of ice temperature T_i and ice pressure P_i .

~~We follow borehole studies by Naruse et al (1988) and Paterson (1977) to write strain, ϵ , in the radial direction as~~
We follow borehole studies by Naruse et al (1988) and Paterson (1977) to write strain, ϵ , in the radial direction as

$$\epsilon = \ln\left(\frac{r_f}{r_0}\right) \quad (89)$$

where a moulin with initial radius r_0 and final radius r_f underwent radial strain of ϵ .

We use the time derivative of Eq. (89) to calculate the change in moulin radius in the $(i+1)^{th}$ timestep as a function of the radius in the i^{th} timestep, separated by time Δt due to viscous deformation:

$$r_{i+1} = r_i \exp(\dot{\epsilon} \Delta t) \quad (9)$$

with strain rate given by Eq. (7). This is the same relationship used by Catania and Neumann (2010):

$$dr_v = r_m \exp(\dot{\epsilon} dt) - r_m \quad (10)$$

with strain rate given by Eq. (8). This is the same relationship used by Catania and Neumann (2010).

2.2.2.2.23 Shear deformation

We use Glen's Flow Law to calculate the change in shape of the moulin due to regional-scale ice flow. This deforms the entire moulin in bulk, shearing it in the vertical and shifting it laterally downstream, without changing its radii. Basal sliding is not currently included in the model. To represent deformation, we discretize the moulin as a stack of plates with elliptical (or circular) holes with a thickness dz and represent deformational ice flow as displacement between these plates.

We calculate $u_d(z)$, the rate of deformational ice flow u_d in the downstream direction, from ice temperature T and pressure PN , surface slope α , a constant enhancement factor E, F_{*2} and ice thickness H_i , using Glen's Flow Law (Cuffey and Paterson, 2010): (Cuffey and Paterson, 2010):

$$u_d = \left[2 E (\rho_i g \alpha)^n \int_{z=b}^{z=s} A(T) (H - z)^n dz \right]^{1/n} \quad (10)$$

We obtain ice deformation rates of $\sim 20 \text{ m yr}^{-1}$, which is typical of the ablation zone in western Greenland (Ryser et al., 2014).

$$2 \cdot u_d = 2 F_* (\rho_i g \alpha)^n \cdot \int_b^{H_i} A(T_i, P_i) (H_i - z)^n dz \quad (11)$$

We obtain ice deformation rates of $\sim 20 \text{ m yr}^{-1}$, which is typical of the ablation zone in western Greenland (Ryser et al., 2014).

2.3 Phase change modules

The second mode that changes the geometry of the moulin is ice ablation from or accretion to the moulin walls. During the melt season, the flow of water into and through the moulin generates turbulence, which as it dissipates acts to melt back the moulin walls, expanding the size of the moulin. There is also a small component of melting due to temperature differences between the water and surrounding ice. Outside the melt season, conduction of latent heat into the surrounding ice causes stagnant water to freeze back onto the moulin walls, contracting the size of the moulin.

2.2.3.1 Refreezing

We calculate freeze-on. Refreezing occurs in cold ice when water flow within the moulin is laminar (Reynold's number $Re < 1$), which occurs only outside absent or slow enough that the melt season.

The MouSh model conserves energy via a balance rate of sensible heat within conduction into the surrounding ice sheet and the latent heat of drops the water in the moulin. The model does not track sensible energy fluxes associated with water flow through the moulin. Equivalently, we assume that all water in the moulin is at the melting point, and that all sensible heat changes within the ice in the moulin walls are compensated by latent heat transfer via the refreezing of moulin water onto the moulin wall. We calculate the wintertime freeze on thickness, δ , at each timestep and at each depth z below the moulin water level as follows:

$$\delta(z) = \frac{k_i}{\rho_i L_f} \frac{\partial T}{\partial x} \Delta t \quad (11)$$

Here, k is the thermal conductivity of ice, L_f is the latent heat of freezing, and $\frac{\partial T}{\partial x}$ is the lateral temperature gradient in the ice at the moulin walls. This temperature gradient varies in space and time and entirely dictates the refreezing rate. We calculate the temperature of the ice sheet, $T(x, z)$, at each depth z using the one-dimensional, diffusion-only heat equation with Dirichlet boundary to the freezing point. These conditions for far field temperature, which we define at 30 meters from the moulin, and 0°C at the moulin wall. We treat the latent heat deposition at the moulin wall as a source term (Poinar et al., 2016). We treat each depth independently, thus assuming zero vertical diffusion or advection. We calculate the change in moulin water volume from freezing, V_{frz} , by summing the refrozen ice thickness, δ , around the perimeter of the moulin at all depths z ,

and converting ice volume to water volume; occur primarily outside the melt season. When these conditions are met, we apply a radial freezing term, which is parameterized economically, following Alley (2005):

$$dr_f = 2 \frac{T_i - T_{pmp}}{L_f} \sqrt{\frac{k_i C_p}{\pi \rho_i}} (\sqrt{t_t} - \sqrt{t_t - dt}) \quad (12)$$

Here, $T_i - T_{pmp}$ is the depth-varying difference between the far-field temperature (prescribed as from borehole temperature observations) and the moulin water temperature, which is taken as the pressure melting temperature T_{pmp} . C_p is the specific heat capacity of ice. The refreezing rates thus evolve exclusively based on the elapsed time since the cessation of turbulent flow, t_t .

We calculate the change in moulin water volume from freezing, V_{frz} , by summing the refrozen ice thickness in a timestep, dr_f , around the perimeter of the moulin at all depths z , and converting ice volume to water volume:

$$V_{frz} = \frac{\rho_i}{\rho_w} \int_0^{h_w} \Pi(z) \delta(z) dz \int_b^{h_w} \Pi(z) r_f(z) dz \quad (12)$$

We also include an option to model refreezing rates more economically, following Alley (2005):

$$\delta(z) = 2 \frac{\Delta T}{L_f} \sqrt{\frac{k C_p}{\pi \rho_i}} (\sqrt{t} - \sqrt{t - \Delta t}) \quad (13)$$

where ΔT is the depth varying difference between the far field temperature and 0°C , and C_p is the specific heat capacity of ice.

2.2.3.2 Moulin wall melting

During the melt season, turbulent energy dissipation from water flowing through the moulin melts back the moulin walls. We calculate the thickness of ice melted, $m(z)$, and add this new volume of meltwater to the water already in moulin, similarly to Eq. (12) for V_{frz} :

$$V_{wallmelt} = \frac{\rho_i}{\rho_w} \int_0^{h_w} \Pi(z) m(z) dz \quad (14)$$

The dissipation of turbulent energy and the associated melting of the surrounding ice will increase the local moulin radius. We parameterize turbulence in two separate spatial domains: (1) within the water column of the moulin (Section 2.2.3.2.1), where r_1 and r_2 are evolved uniformly, and (2) above the water level along the side of the moulin, as supraglacial input falls to the water level (Section 2.2.3.2.2). The second parameterization, while simplifying complex hydraulics and melting patterns, is necessary to offset the steady viscoelastic closure of the moulin above the water line, where only r_2 is evolved.

The parameterizations of turbulently driven melting we use in both regimes rely on two simplifications. First, the volume of water moving through each vertical model node is constant within each time step. This ensures that water mass is conserved and that all model elements below the water line are water filled; however, this eliminates the potential long-term storage of meltwater within plunge pools caused by non-uniform incision into the ice. Second, all energy generated from turbulent dissipation is instantaneously applied to melting the surrounding ice. This neglects any heat transport within the

water, which is a common approximation in subglacial models (e.g., Hewitt, 2013; Schoof, 2010; Werder et al., 2013), and allows us to make the simplifying assumption that meltwater entering the moulin is at 0°C and at the pressure melting temperature at all points below the water line. As part of our melt parameterization, we also include the effect of the temperature difference between the water and the surrounding ice (Jarosech and Gudmundsson, 2012) because, unlike for subglacial channels, we cannot assume that the surrounding ice is at the pressure melting temperature. Though we model instantaneous heat exchange, we note recent findings that the appropriate heat transfer coefficient can be difficult to determine (Sommers and Rajaram, 2020).

2.2.3.2.1 Melt belowThe parameterizations of turbulently driven melting we use in both regimes rely on three simplifications. First, the volume of water moving through each vertical model node is constant within each time step. This ensures that water mass is conserved and that all model elements below the water line are water filled; however, this eliminates the potential long-term storage of meltwater within plunge pools caused by non-uniform incision into the ice. Second, all energy generated from turbulent dissipation is instantaneously applied to melting the surrounding ice. This neglects any heat transport within the water, which is a common approximation in subglacial models (e.g., Hewitt, 2013; Schoof, 2010; Werder et al., 2013). Third, we also make the simplifying assumption that meltwater entering the moulin is at 0°C and at the pressure melting temperature T_{mp} at all points below the water line, though we do not model the impact of this temperature change on melting because moulin water temperatures are unknown.

Submerged zone: Below the water line

Below the waterline, the vertical velocity of the water is dictated by the hydraulic gradient within the system and the local cross-sectional area of the moulin. Under such conditions, head loss – the departure of the hydraulic head from that calculated by Bernoulli’s equation – reflects the energy dissipated as heat. We parameterize head loss using the Darcy–Weisbach equation, which relates water velocity (u_w) to changes in the hydraulic gradient (dh/dL (head loss per unit length along flow), via the hydraulic diameter (D_h), radius R_h and a dimensionless friction factor (f_R) and gravitational acceleration (g). Because water velocity is constrained by mass balance within the system, we calculate the head loss (dh/dL), or turbulent energy dissipated into melting the moulin walls, as follows:

$$\frac{dh}{dL} = \frac{u_w^2 f_R}{2D_h g} \quad (15)$$

The differential element dL represents the path length over which the water experiences head loss: $dL = \sqrt{dx^2 + dz^2}$ for horizontal distance dx and vertical drop dz . The friction factor (f_R) is a unitless model parameter that controls the rate of head loss within the system. Its value thus directly affects the amount of internal melting. Most subglacial models fix the Darcy–Weisbach friction factor, with values ranging from 0.01 to 0.5 (e.g., Colgan et al., 2011b; Schoof, 2010; Spring and Hutter, 1981) or use equivalent values of Manning’s n (e.g., Hewitt, 2013; Hoffman and Price, 2014). Such constant values, however, are somewhat at odds with field observations that indicate highly variable subglacial channel roughness over a range of time

and spatial scales (Gulley et al., 2014; Mankoff et al., 2017). Alternatively, other models parameterize channel roughness using a geometry-dependent friction factor (e.g., Boulton et al., 2007; Clarke, 2003; Flowers, 2008). All current parameterizations for time-varying roughness, however, were developed outside moulines and thus may not accurately represent conditions there, where observations of extensive ice scalloping, for instance, make moulines distinct from other conduits (Gulley et al., 2014). Thus, MouSh has options for fixed or variable friction factors. The overall effect on moulin geometry is modest.

For time and geometry-dependent parameterization of roughness, we choose the Bathurst parameterization (Bathurst, 1985):

$$f_k = \left(-1.987 \log \left(\frac{k_s}{5.15 R_h} \right) \right)^{-2} \quad (16)$$

Here, k_s is the surface roughness height and R_h is the channel hydraulic radius,

which is equal to the cross-sectional area divided by the wetted perimeter at heights at or below the water line. $\frac{dh_w}{dl} = \frac{u_w^2 f}{8 R_h g}$

$$(14)$$

The differential element dl represents the path length over which the water experiences head loss: $\partial l = \sqrt{\partial x^2 + \partial z^2}$ for horizontal distance dx and vertical drop dz . The friction factor f is a unitless model parameter that controls the rate of head loss within the system. Its value thus directly affects the amount of melting. Most subglacial models fix the Darcy-Weisbach friction factor, with values ranging from 0.01 to 0.5 (e.g., Colgan et al., 2011b; Schoof, 2010; Spring and Hutter, 1981) or use equivalent values of Manning's n (e.g., Hewitt, 2013; Hoffman and Price, 2014). Alternatively, other models parameterize channel roughness using a geometry-dependent friction factor (e.g., Boulton et al., 2007; Clarke, 2003; Flowers, 2008). Thus, MouSh has options for fixed or variable f .

The friction factor within the submerged zone is indicated by f_m and in the open channel zone by f_{oc} . To explore the impact of the chosen friction factor, we complete a sensitivity study (Sect. 2.3 and 3.2) where we vary the friction factor in water filled sections, f_m , over an expected range, centered on $f_m = 0.1$. We use a constant $f_m = 0.1$ for all other model runs presented.

Because we approximate the moulin as a half-circular, half-elliptical cylinder with perimeter Π , the hydraulic radius R_h of a water filled node is:

$$R_h = \frac{A_m}{\Pi} \quad (17)16.$$

for moulin perimeter Π . The Bathhurst parameterization produces a range of friction factors over approximately two orders of magnitude for typical Greenland moulines. This and other roughness parameterizations cannot adequately account for changing form roughness, including sinuosity and large-scale ice scalloping. To explore this, we complete a sensitivity study (Sect. 2.3 & 3.2) where we fix the friction factor over the expected range, centered on $f_k = 0.1$. We use a constant $f_k = 0.1$ for all other model runs presented.

~~We calculate the head loss used to determine the amount of~~ To calculate moulin wall melting ~~using, we use~~ a simple energy balance equation: following previous work (e.g., Gulley et al., 2014; Jarosch and Gudmundsson, 2012; Nossokoff, 2013):

$$\rho_i C_w \Delta T \frac{dS}{dt} + \rho_i (T_{pmp} - T_i) \frac{dA_m}{dt} + \rho_i L_f \frac{dS}{dt} \frac{dA_m}{dt} = \rho_w g Q \frac{dh_L}{dz} \quad (18)$$

$$Q \left(\rho_w g \frac{dh_w}{dl} \right) \quad (17)$$

The first term represents sensible heat, the small amount of energy needed to warm the surrounding ice to the pressure melting point. Otherwise, Eq. (18) follows previous work based in temperate ice conditions (e.g., Gulley et al., 2014; Jarosch and Gudmundsson, 2012; Nossokoff, 2013). Equation (18) can be rearranged and modified for our elliptical system such that the change in moulin radius due to melting is:

$\Delta r_t = \left[\frac{\rho_w g Q}{\pi \rho_i (C_w \Delta T + L_f)} \frac{dh_L}{dz} \right]$ The first term represents the energy needed to warm the surrounding ice to the pressure melting temperature of water T_{pmp} . Equation (17) can be rearranged and combined with equation (14) to provide the area of ice melted:

$$dA_t = Q_{out} \left(\rho_w g \frac{u_{wf}^2}{4Rhg} \right) \left(\rho_i C_w (T_{pmp} - T_i) + \rho_i L_f \right)^{-1} dt \quad (19)$$

Here, dh_L/dL is the head loss over the model node (Eq. 15), π is the wetted perimeter of a water filled moulin node; Q Where Q_{out} is the discharge from the moulin-subglacial system as dictated by the subglacial model component (Section Sect. 2.2.4.2); and $\Delta T = T_i - T_{pmp}$ is the temperature difference between the water (prescribed to be at the pressure melting point) and the surrounding ice, which can be assigned a number of different profiles from Greenland as described in Table 1. we can vary from site to site around Greenland as described in Table 1. Note that Eq. 18 determines the area of ice that is removed through melting. For each time step, we reframe Eq. 18 into radial melting within an egg-shaped moulin using information about the previous geometry and the assumption that melting occurs uniformly around the perimeter:

2.2.3.2.2 Melt above the water line

$$dr_t = 2dA_t / \left[\pi(5r_1 + 3r_2 - \sqrt{(3r_1 + r_2)(r_1 + 3r_2)}) \right] \quad (19).$$

Equation 19 is simplified when considering a circular geometry ($r_1 = r_2$).

Unsubmerged zone: Above the water line, a range variety of complex processes drive melting. A first-principles approach is would be to quantify melting due to the potential energy loss of falling water, following the work on terrestrial waterfalls (e.g., Scheingross and Lamb, 2017). (e.g., Scheingross and Lamb, 2017). However, nearly all waterfall-based parameterizations rely on abrasion between waterborne sediment and the substrate as the primary mechanism of erosion. Instead, we implement a simple parameterization for open-channel flow with the understanding that the complexities of thermal erosion are not completely captured. In our model, open-channel melting occurs only on the up-glacier wall of the moulin and follows two ad-hoc rules based on the slope between the vertical nodes: (1) open-channel turbulent melting is applied if the slope of the upstream moulin wall allows water to flow over it; and (2) a small, prescribed amount of melting is applied when the upstream

wall slope is vertical or overhung, because while water cannot flow directly along the ice, spray and other processes likely drive some amount of melting. These cases are respectively (1) the open-channel zone and (2) the falling water zone.

Open-channel zone:—In the open-channel zone, we use a similar approach as for melting below the water line. However, the hydraulic radius R_h is adjusted to reflect the observation that water runs down only one wall of the moulin, and a higher friction factor is used to represent complex geometries (e.g., Covington et al., 2020). Due to the presence of a discontinuity between open-channel and water-filled regions (at the water line), we parameterize the wetted perimeter hydraulic radius of open channel flow as follows:

$$R_{hopen} = \pi r_1 \quad (20)$$

To drive $0.5r_2$. We also use a higher open channel friction factor f_{oc} of 0.8 to parameterize observed extensive scalloping (e.g., Gully et al., 2014; Covington et al., 2020). We apply melting in the open-channel region, we use Eq. (17) with a hydraulic radius of only a portion the elliptical side of the moulin, defined by r_2 derived using Eq. 18. Note that the hydraulic radius prescribed for open-channel flow is likely larger than the small region over which water is flowing in the natural system (Fig. 2d), minimizing 2a,d). Further, the amount of resulting melt ΔA_{oc} is applied only to the model major radius to calculate dr_{oc} .

Falling water zone:—In the falling water zone, there is very limited interaction between the moulin walls and the water. For simplicity, we assume that a small fraction, f_f , of the potential energy lost as water falls reaches deposited into the moulin walls, perhaps impacting it as the kinetic energy of spray, and is used to melt the surrounding moulin. The change in radius due to this process is as follows:

$$\Delta r_f dr_{mf} = f_p \frac{(\rho_w/\rho_i)gQ_{in}}{L_f \Pi} dt \quad (21)$$

We set f_p , the fraction of potential energy applied to melting the surrounding walls, to 0.1% for the all model runs presented here.

We add the volume of ice melted to the water already in the moulin, similarly to Eq. 12 for V_{fz} . We calculate the change in moulin water volume from melting by summing the melted ice thickness, r_{mf} , around the perimeter of the moulin at all depths z , and converting ice volume to water volume:

$$V_{wallmelt} = \frac{\rho_i}{\rho_w} 2 \cdot dt \int_b^H \Pi(z) r_{mf}(z) + A_{oc}(z) + A_t(z) dz \quad (21)$$

2.4 Water flux into and out of the moulin

We enforce (Mass conservation of water mass within the moulin system as follows:)

$$\Delta V_{water} = (Q_{in} - Q_{out}) \Delta t + V_{wallmelt} - V_{fz} \quad (22)$$

The change in the volume of water stored in the moulin, ΔV_{water} , Water balance within a time step is proportional to the volumetric recharge from a supraglacial stream (Q_{in} , Sect. 2.4.1), discharge into and out of the moulin (Q_{in} through a subglacial channel (Q_{out} , Q_{base} ; Sect. 2.4.2) and Q_{out} , respectively) and the change in volume of water added due to melting (V_{wallmelt}) or removed due to refreezing (V_{frz}). The latter two terms are described, such that the volume of water in the system (V) is:

$$\frac{dV}{dt} = Q_{in} - Q_{out} + Q_{base} + \frac{(dV_{\text{wallmelt}} - dV_{\text{frz}})}{dt} \quad (22).$$

The integral term varies in space and time, with high melt rates above in Eq. (14) the water line during the melt season (when $Q_{in} > 0$), and (12), respectively. Discharge into moderate melt rates at and out of the system are described below the water line during and after the melt season, when there is water flow through the moulin ($Q_{out} > 0$) and refreezing below the water line throughout the winter (when $Q_{in} = Q_{out} = 0$). The MouSh model can also accept an additional prescribed base flow Q_{base} directly to the subglacial module. We design base flow as a loose approximation of additional subglacial water inputs from varied upstream sources, including other moulins on the same subglacial channel, regional basal melt, and the addition and removal of meltwater from subglacial storage. Base flow is generally required to maintain realistic moulin water levels. In the moulin runs forced by realistic Q_{in} , we represent subglacial flow from ~5 surrounding moulins by prescribing base flow as five times the running 5-day mean of Q_{in} . In other model runs, we do not include base flow. The addition of base flow is designed to represent the widespread seasonal evolution of surface melt; its inclusion maintains a slightly larger subglacial channel than would otherwise occur, which reduces otherwise unrealistically large daily swings in modeled moulin water level.

2.2.4.1 Meltwater runoff from the ice sheet surface

We force the MouSh model with time-varying water inputs from the supraglacial environment, Q_{in} . We use two different Q_{in} scenarios: a simple diurnal cosine with maximum and minimum discharges ranging between ~1 and 5 m^3s^{-1} , in rough agreement with observations near the margins (Eq. 23, Chandler et al., 2013; McGrath et al., 2011; Smith et al., 2017); and realistic supraglacial discharge over a melt season, determined by using in-situ surface melting data and internally drained catchment size and geometry (Yang and Smith, 2016).

We use the following cosine curve to represent our simplest form of supraglacial discharge into the moulin during sensitivity studies:

$$Q_{in} = \cos(\pi(t - 19.5)/12) + 3 \quad (23)$$

Here, t is time in days 2.4.1 Meltwater runoff from the ice-sheet surface

We force the MouSh model with time-varying water inputs from the supraglacial environment, Q_{in} . We use two different Q_{in} scenarios: a simple diurnal cosine with maximum and minimum discharges ranging between ~1 and 5 m^3s^{-1} , in rough agreement with observations near the margins (Eq. 23, Chandler et al., 2013; McGrath et al., 2011; Smith et al., 2017); and

realistic supraglacial discharge over a melt season, determined by using in-situ surface melting data and internally drained catchment size and geometry (Yang and Smith, 2016).

We use the following cosine curve to represent our simplest form of supraglacial discharge into the moulin during sensitivity studies:

$$Q_{in} = \cos(\pi(t - 19.5)/12) + 3 \quad (23)$$

Here, t is time in hours and Q_{in} is in $\text{m}^3\text{m}^{-3}\text{s}^{-1}$. This function has its daily peak at 19:30 hours and a daily minimum at 07:30.

~~Initial surface runoff values for the 2019 melt season were modified using a synthetic unit hydrograph derived for the ablation zone and parameters appropriate for western Greenland (Table 2, Smith et al., 2017). The parameters for the unit hydrograph were determined during the middle of the melt season and therefore may inaccurately represent routing delays at the beginning and end of the melt season.~~

~~The MouSh model can also accept base flow directly to the subglacial module. We design base flow as a loose approximation of additional subglacial water flow from varied upstream sources, including other moulins on the same subglacial channel, regional basal melt, and the addition and removal of meltwater from subglacial storage. The latter reflects the englacial void ratio used in many subglacial models. Base flow is generally required to maintain realistic moulin water levels. In the moulin runs forced by realistic Q_{in} , we represent subglacial flow from 5 surrounding moulins by prescribing base flow as five times the running 5-day mean of Q_{in} . This application mimics the seasonal evolution of surface melt and maintains a slightly larger subglacial channel than would otherwise occur, which reduces otherwise unrealistically large daily swings in modeled moulin water level.~~

~~2. To examine a set of realistic moulins, we select three supraglacial basins from Yang and Smith (2016) and derive their size, distance from terminus from information provided therein (Basin 1-3; Table 2). We derive surface runoff from MERRA-2 reanalysis (Gelaro et al., 2017; Smith et al., 2017). Surface runoff values for the 2019 melt season were modified using a synthetic unit hydrograph derived for the ablation zone and parameters appropriate for western Greenland (Table 2, Smith et al., 2017). The use of a unit hydrograph parameterizes the time and magnitude adjustments expected from meltwater routing over the ice surface. The parameters for the unit hydrograph were determined during the middle of the melt season and therefore may inaccurately represent routing delays at the beginning and end of the melt season. Further details on supraglacial and internal catchment characteristics are included in Sect. 2.5.2.~~

2.4.2 Water flow ~~to~~from the subglacial system

~~We couple the moulin model and a single evolving subglacial channel controlled by melt opening and creep closure (Covington et al., 2020; Schoof, 2010) using a reservoir constriction model (Covington et al., 2012) that simulate flows between the two elements.~~

~~The time rate of change of water level (h) is developed in Covington et al. (2012):~~

$\frac{dh}{dt} = \frac{1}{A_m(h)} (Q_{in} + Q_{base} - Q_{out})$ We couple the moulin model and a single evolving subglacial channel controlled by melt opening and creep closure (Covington et al., 2020; Schoof, 2010) using a reservoir-constriction model (Covington et al., 2012) that simulate flows between the moulin and subglacial channel. Following Covington et al. (2020), the rate of change of moulin water level h_w is

$$\frac{dh_w}{dt} = \frac{1}{A_m(h_w)} \frac{\partial V}{\partial t} \quad (24)$$

where $A_m(h)$ is the moulin cross-sectional area at the water level h . The water level (Eq. 3), volume is related to Q_{in} is the influx of surface meltwater, Q_{base} is the base flow at each time step (Sect. 2.2.4.1), and Q_{out} , where Q_{out} is the meltwater output from the subglacial channel, defined as follows:

$$Q_{out} = c_3 S^{5/4} \Psi / \sqrt{|\Psi|} \quad (25)$$

Here, S is the subglacial channel cross-sectional area. The hydraulic gradient $\Psi = -\rho_w \rho_i g \frac{d(h_w + b)}{dL}$ assumes zero bed slope and $\frac{d(h_w - b)}{dL}$ is a linear gradient in the moulin water level $(h_w - b)/L$ to the outlet at a horizontal distance L , where the pressure head is zero. In our calculations, the bed elevation b is zero. Finally, c_3 is a flux parameter:

$$c_3 = \frac{2^{5/4}}{\pi^{1/4}} \sqrt{\frac{\pi}{(\pi+2)\rho_w f}} \sqrt{\frac{\pi}{(\pi+2)\rho_w f}} \quad (26)$$

Equation (26) for c_3 follows Covington et al. (2020), who corrected a small error from the original Schoof (2010) (2020), who corrected a small error from the original Schoof (2010) formulation.

We use an equation from Schoof (2010) for the time rate of change in subglacial channel cross-section area S , with the first part describing the turbulent melting of the subglacial channel walls, and the second term describing closure due to the pressure of the overlying ice:

$$\frac{dS}{dt} = c_1 \frac{Q_{out}}{L_f} - c_2 N^n S^{-n} \quad (27)$$

We use an equation from Schoof (2010) for the time rate of change in subglacial channel cross-section area S , with the first part describing the turbulent melting of the subglacial channel walls, and the second term describing closure due to the pressure of the overlying ice:

$$dS = (c_1 Q_{out} \Psi - c_2 N^n S^{-n}) dt \quad (27)$$

Here, the constant $c_1 = \frac{1}{\rho_i L_f}$ with ρ_i the ice density and L_f the latent heat fusion of ice, the constant $c_2 = 2A n^{-n} (T_i, N) n^{-n}$ with the Glen's flow law parameters $A \approx 10^{-24}$ for the subglacial component defined as $A = 6 \cdot 10^{-24} \text{Pa}^{-3} \text{s}^{-1}$ and $n = 3$. The effective pressure $N = P_i - \rho_w g h_w$.

Replacing Q_{out} , Ψ , and N in Eq. (27) yields

$$\frac{dS}{dt} = c_1 c_3 S^{5/4} \left(\frac{\rho_w g h}{L} \right)^{3/2} \left(\frac{\rho_w g h_w}{L} \right)^{3/2} - c_2 (P_i - \rho_w g h_w)^n (P_i - \rho_w g h_w)^n S^{-n} dt \quad (28)$$

Equations (24) and (28) are numerically solved simultaneously, as in Schoof (2010) and Covington et al. (2021). The parameters used in this module are included in Table 1 and are the same as those used in the englacial component of MouSh, with the exception of apart from the flow law parameter A . In the englacial system, A is calculated from local temperature within the ice column, which can be as cold as -23°C in our study area (Iken et al., 1993), western Greenland (Iken et al., 1993). This contrasts with the temperature at the ice-bed interface, which must be at the melting point; thus, the subglacial component of MouSh uses a higher fixed A value.

2.2.4.3 Water volume contributions from melting and freezing

Water balance within the moulin and the subglacial channel is dictated by recharge from a supraglacial stream (Q_{in}), discharge through a subglacial channel (Q_{out}), and any change in volume due to melting or refreezing (related to $m(z)$, the radial melt or refreezing rate, in m s^{-1}), such that the volume of water in the moulin (V_m) is

$$\frac{dV_m}{dt} = Q_{in} - Q_{out} + \int_0^H \Pi(z) m(z) dz \quad (29)$$

The integral term varies in space and time, with high melt rates above the water line during the melt season (when $Q_{in} > 0$), and moderate melt rates at and below the water line during and after the melt season, when there is water flow through the moulin ($Q_{out} > 0$) and refreezing below the water line throughout the winter (when $Q_{in} = Q_{out} = 0$).

2.3 In its current configuration, the subglacial module provides a single set of outputs representative of conditions at the moulin. This is primarily because this study focuses on the evolution of a moulin and is not representative of a channel running from a moulin to the terminus in a natural system. A more complex subglacial model would more accurately resolve the spatial changes in subglacial channel geometry and flow.

2.5 Suites of model experiments

To examine the sensitivity of the MouSh model to uncertain parameters, ice and meltwater characteristics, and model choices, and difference from previous moulin parameterizations, we run four suites of experiments. While these experiments do not cover the complete range of possibilities, they were designed to address primary uncertainties in the MouSh model and examine how moulin geometry might vary spatially and temporally.

2.5.1 Quasi-equilibrium and the impact of diurnal supraglacial variability

Under steadily varying conditions such as a repeating diurnal variation, the modeled moulin reaches a quasi-equilibrium state independent of initial conditions with melting opposing viscous and elastic deformation and the only change being driven by shear deformation. We examine the quasi-equilibrium state and the impact of supraglacial variability on this state. Supraglacial runoff Q_m is highly variable with seasonal, event, and diurnal variability in surface melting modified by supraglacial drainage basin characteristics as it is routed to a moulin. Moulin water level and shape respond to these patterns of variability. To examine the impact of Q_m magnitude (mean) and Q_m amplitude (variability), we perform a series of model runs that vary the

magnitude of a cosine curve between 1 and 20 m^3s^{-1} with a fixed amplitude of 0.5 m^3s^{-1} and a series of runs that vary the amplitude of a cosine curve between 0 and 2 m^3s^{-1} with a fixed magnitude of 5.0 m^3s^{-1} . The amplitude as one half the diurnal range. These runs use Basin 1 ice conditions (Table 2; Sect. 2.5.4) with no base flow prescribed. Further details can be found in Supplement S2.1.

2.5.2 Sensitivity to uncertain parameters

We explored the sensitivity of our results to the values of seven parameters, shown in Fig. 43, with the prescribed ranges shown in Table 1. We studied the effect on the water level; the moulin radius at the equilibrium water level; the volume and water storage of the moulin; and the cross-sectional area of the subglacial channel at the end of a ten-day model run. These values reach equilibrium, with daily oscillations superimposed, after 3–5 days. We also tested the dependence of our results on the initial moulin radius, R_{i0} , which we varied across an order of magnitude from 0.565 to 5.0 meters.

We varied the value of a uniform deformation enhancement factor E over an order of magnitude ($E = 1$ to 9), which affects viscous flow of the ice surrounding the moulin. ~~We also tested the effect of ice temperature, independent of the enhancement factor. We used five different temperature profiles: cold ice temperatures (mean -15°C , range -23.1°C to the pressure melting point) measured in the center of Jakobshavn Isbræ (Iken et al., 1991); moderate ice temperatures (mean -7°C , range -13.5°C to the pressure melting point) measured at the GULL site in Pâkitsoq (Lüthi et al., 2015; Ryser et al., 2014); warmer ice temperatures (mean -5°C , range -9.3°C to the pressure melting point) measured at the FOXX site in Pâkitsoq (Lüthi et al., 2015; Ryser et al., 2014); a hypothetical linear profile from -5°C at the surface to 0°C at the bed; and, finally, a fully temperate ice column. These different ice temperature scenarios affected the creep closure rates of ice through the temperature-dependent softness parameter A by approximately a factor of 6 from the coldest profile (Iken et al., 1993) While the range of enhancement factors tested exceeds that likely to be observed in the field, the variation of an order of magnitude was chosen to match the range of other rheological parameters. We also tested the effect of ice temperature, independent of the enhancement factor. We used five different temperature profiles: cold ice temperatures (mean $\sim -15^\circ\text{C}$, range -23.1°C to the pressure melting point) measured in the center of Jakobshavn Isbræ (Iken et al., 1991); moderate ice temperatures (mean $\sim -7^\circ\text{C}$, range -13.5°C to the pressure melting point) measured at the GULL site in Pâkitsoq (Lüthi et al., 2015; Ryser et al., 2014); warmer ice temperatures (mean $\sim -5^\circ\text{C}$, range -9.3°C to the pressure melting point) measured at the FOXX site in Pâkitsoq (Lüthi et al., 2015; Ryser et al., 2014); a hypothetical linear profile from -5°C at the surface to 0°C at the bed; and, finally, a fully temperate ice column. These different ice temperature scenarios affected the creep closure rates of ice through the temperature-dependent softness parameter A by approximately a factor of 6 from the coldest profile (Iken et al., 1993)~~ compared to the fully temperate column.

We also examined moulin sensitivity to elastic deformation by varying Young's modulus (YE) of the ice column between 1–9 GPa (e.g., Vaughan, 1995). ~~We also tested (Vaughan, 1995) and the~~ sensitivity to the values of friction factors for the moulin walls. ~~These factors control melt rates associated with the dissipation of turbulent energy.~~ MouSh has two friction

factors: f_r (~~moulin friction; f_m~~ (below the water line) and f_c (~~open-channel friction;~~ above the water line). We varied these friction factors across two orders of magnitude. We did not vary the subglacial channel friction factor. Finally, we varied values for basal ice softness over two orders of magnitude and independently examined moulins over a range of ice thicknesses (670–1570 m) and corresponding distance from the terminus (~20–110 km), which in combination results in variations in hydraulic gradient.

2.45.3 Sensitivity to local conditions

We examined moulins over a range of ice thicknesses, and corresponding ~~distanced~~distances from the terminus, and appropriate Q_{in} forcings for three different representative locations on the ice sheet (Table 2). We designed this suite of model experiments to provide general guidance on Each moulin is associated with a supraglacial basin derived by Yang and Smith (2016). The moulins were selected based on ice thicknesses that broadly represent the range of variability in moulin ice thicknesses within the ablation zone of the western Greenland Ice Sheet and supraglacial drainage basin sizes and geometries that were visually similar to nearby drainage basins and approximately representative of the mean supraglacial drainage basin area for the given ice thicknesses (553m, 741m, and 1315m). To derive broadly representative Q_{in} values for each basin, we integrate 3-hourly modeled surface melting from a downscaled version of MERRA-2 (Gelaro et al., 2017) over the course of surface area of each moulin surface drainage basin. We then use synthetic unit hydrograph parameters derived for a supraglacial basin from western Greenland during the middle of the 2015 melt season (Smith et al., 2017) to estimate supraglacial discharge into each moulin.

The supraglacial discharge curves for each moulin are only meant to capture the seasonal change in discharge rates and diurnal variability and occasional increases in runoff due to surface melt events during the 2019 melt season and over a range of supraglacial catchment sizes. As part of this analysis, we. The primary goal of this exercise is to examine season-long and daily differences in model outputs and, the variation in each model component (viscous, elastic and phase change), and their the relative importance of each component in driving moulin geometry change at different representative locations of the western Greenland Ice Sheet.

2.5.4 Comparison to a cylindrical moulin

Subglacial models generally use a time-invariant vertical cylinder to represent moulins. To investigate and quantify the efficacy of our time-evolving moulin shape model, we drove MouSh and a static cylinder with the same meltwater inputs. We use the time-mean radius at the water level as the radius of the static cylinder; this is 1.4 m for Basin 1 and 1.3 m for Basin 2. We compared the resulting moulin water level, moulin capacity, subglacial cross-sectional area and meltwater input difference (due to melt generated within the model itself) across these runs. We compared the moulin water level values directly (variable water level — cylindrical water level — variable water level) and compared other metrics by percentage difference ($2 \cdot (\text{variable} - \text{cylindrical} - \text{variable}) / (\text{variable} + \text{cylindrical})$).

3 Results

3.1 Moulins as part of the channelized englacial-subglacial system

The capacity for water storage in the englacial system, relative to the subglacial system, depends on both the spatial density of moulins and the volumes of individual moulins. Below 1300 m elevation, moulin densities are ~ 0.27 per km^2 in Pâkitsoq (270 moulins) and ~ 0.16 per km^2 (704 moulins) in the Russell region (Fig. 2a, 3a-b). The total length of subglacial flow pathways are 765 km and 4,679 km for Pâkitsoq and Russell regions, respectively. The distribution of ice thicknesses at each moulin location are shown in Fig. 3c; the cumulative englacial path lengths are ~ 110 km in Pâkitsoq and ~ 446 km in the Russell region. Thus, we find that moulins comprise between 10% (Russell) and 14% (Pâkitsoq) of the path length that water takes from its entry to the englacial-subglacial system to its exit at the terminus or calving front. These lengths are not insignificant and suggest that moulin geometry and evolution may be important to subglacial processes.

2.5.5 Sensitivity to model choices

As part of MouSh development, we made several decisions about how to represent moulin geometry, water inputs, and the associated subglacial system that can directly impact the shape and water level of a modeled moulin. These decisions include (1) representing moulin cross-sectional area as a semi-elliptical, semi-circular “egg” instead of as a circle (Sect. 2.1 and 2.3.2); (2) the inclusion of estimated surface stresses in the representation of elastic deformation (Sect. 2.2.1); (3) the use of a parabolic ice sheet profile to determine the surface slope and distance to terminus for a given ice thickness (Cuffey and Paterson, 2010); (4) the use of prescribed base flow into the subglacial component of the model (Sect. 2.4); and (5) the use of a time-evolving subglacial channel. The first two choices pertain to the complexity of the model, with our choices being more complex; simplification may be beneficial in some circumstances. Choices 4 and 5 reflect the need for a subglacial hydrologic model and would be eliminated if MouSh was configured to function with either specific observational data or with a more comprehensive subglacial model. We also test the impact of the magnitude and diurnal variability of Q_{in} on the timescale for the moulin to reach quasi-equilibrium.

To explore the impact of our model choices for decisions 1-4, we perform a series of experimental comparisons against the seasonal run for Basin 1. This allows us to capture the effect of our choices during periods of increasing and decreasing Q_{in} . We change only the parameter of interest to isolate the effect on moulin water level and moulin capacity, the two variables that most directly affect water flow within the subglacial system.

To examine the effect of an evolving versus a fixed-radius subglacial channel, we complete a series of runs with the same ice thickness and distance from terminus as Basin 1 but use a simpler Q_{in} , the cosinusoidally varying function described in Sect. 2.4.1. Further description of these runs is included in Supplement S2.2.

3 Results

3.1 Quasi-equilibrium and dependence on Q_{in}

Under uniform supraglacial inputs, the moulin water level, radius, and water capacity reach equilibrium within 10 days (red line, Fig. S2). However, supraglacial inputs are rarely, if ever, uniform, so under constantly varying conditions, the moulin will reach a ‘quasi-equilibrium’ state. This is a mean state (geometry, water level, deformation rates) with superimposed variability on the timescale of variations in Q_{in} alone. Therefore, if the forcing is diurnal, the moulin will exhibit diurnal variability from a mean state. The quasi-equilibrium state is dependent on model characteristics and parameters (Sect. 3.2; Supplement S2.2).

The magnitude and amplitude of Q_{in} alter the moulin water level and major radius at the mean water level (a proxy for moulin geometry) in predictable ways (Fig. S2 and Fig. S3). Increasing the diurnal amplitude of Q_{in} increases the diurnal variability and mean moulin water (Fig. S2b, Fig. S4). This occurs due to the disparate timescales of ice deformation versus melting. The daily increase in Q_{in} raises the water level quickly because the subglacial channel is slow to expand by melting. Conversely, the nightly fall in Q_{in} is muted by a fast contraction of the subglacial channel. This behavior drives the daily peak in moulin water level higher above the mean than daily minimum water level falls below it (Fig. S2b). The “extra” time spent with higher water levels reduces the visco-elastic closure of the moulin while also increasing turbulent melting, resulting in a larger moulin, as indicated by the moulin radius at the mean water level (Fig. S2c). Higher diurnal amplitudes in Q_{in} magnify this effect.

As the Q_{in} magnitude increases, both the mean water level and its diurnal variability decrease (Fig. S3a-b). This occurs because the moulin becomes larger in response to increasing Q_{in} and subsequent increases in subglacial discharge. As the moulin and subglacial channel widen, they can readily accommodate the fluctuations in Q_{in} with more limited variations in moulin water level. This accommodation is evident in the moulin radius at the mean water level (Fig. S3c). Higher Q_{in} magnitude drives a linear increase in melt rates within the moulin alongside nonlinear increases in visco-elastic deformation, causing an overall nonlinear increase in mean moulin water level (Fig. S4). However, when moulin water levels exceed flotation, the moulin grows due to both visco-elastic deformation and melting, resulting in a larger than expected moulin (red line, Fig. S3c).

3.2 Sensitivity of MouSh to parameter values and deformational processes

A range of ice characteristics affect the time evolution of moulin geometry. These include the initial moulin size, temperature and viscosity of the ice column, viscosity of basal ice, friction factors, and ice thickness. ~~These~~Some of these factors are ~~either~~ highly spatially variable (e.g., ice thickness) ~~or~~and others are poorly known (e.g., basal ice viscosity). We quantify the effect of these factors on moulin water level and moulin volume, moulin geometry, and subglacial channel cross-sectional area over both multi-day and diurnal timescales by performing multiple independent sensitivity studies (~~Section~~Sect. 2.3).

We find that moulins reach sizes near equilibrium within 3–5 days, regardless of their initial radii, but that the equilibrium aspect ratio is sensitive to the initial radius (Fig. 4a–b). Moulins initialized with large radii (5 m) reach equilibrium shapes that are more elliptical than moulins initialized with more realistic radii (0.5 m). In particular, the equilibrium major radius is 36% larger and the minor radius is 27% larger for 5 m initial conditions compared to 0.5 m (Fig. 4a), which manifests as a 14% increase in moulin capacity (Fig. 4b). The initial condition also affects the magnitude of diurnal variations in major radius (97%), minor radius (26%), and moulin water storage (13%), with larger initialized moulins varying less (Fig. 5a). Regardless of initial condition, the major radius undergoes lower magnitude diurnal fluctuations than the minor radius. The initial moulin radius does not significantly affect equilibrium subglacial channel size (0.04%), moulin water level (0.02%), or moulin water storage (1.5%). Diurnal variations in subglacial channel size (0.3%), water level (0.4%), and moulin volume (5.4%) were all also insensitive to initial conditions (Fig. 5b).

We find that moulins reach a quasi-equilibrium within 10 days, where the mean moulin water level and the moulin radius at this location are constant. This quasi-equilibrium is independent of the initial moulin radius (Fig. 3a–b, Fig 4a–b), apart from locations above the water line (Supplement S2.2; Fig. S2.4g) where surface deviatoric and shear stresses impact moulin shape.

Three major parameters affect the degree of viscous and elastic deformation in the moulin: the ice flow enhancement factor E , the ice temperature profile $T(z)$, and Young's Young's modulus YE . We tested a span of reasonable values representative of Greenland ice (Table 1) and found a limited effect on moulin geometry. Equilibrium moulin water level, subglacial channel area, and their diurnal variabilities remain constant (<0.1% change) over the tested range of these parameters (Fig. 3d,f,h & 4d,f,h & 5d,f,h). Moulin capacity and water storage show moderate sensitivity (~20% in equilibrium value and ~40% in diurnal range) across the range of E and T scenarios tested; a decrease in moulin capacity and water storage pair with an increase in the diurnal variability for these variables. For instance, varying E across an order of magnitude grew the equilibrium major radius by 23% and shrank the equilibrium minor radius by 44%, with a net effect that moulins had 23% less volume and 20% less water storage capacity in softer ice ($E = 9$) compared to harder ice ($E = 1$) (Fig. 4e3c–d). Similarly, the different ice temperature profiles we tested caused variations of 11% in moulin major radius, 18% in moulin minor radius, and 24% in moulin capacity and moulin water storage, with warmer ice hosting smaller moulins (Fig. 4e3e–f).

We varied Young's Young's modulus, YE , across one order of magnitude. With the highest Young's Young's modulus, moulin major radius increased by 50% compared to the lowest, minor radius decreased by 15%, moulin water volume increased by 38%, and moulin capacity increased by 56% (Fig. 4g3g–h). The equilibrium water level decreased insignificantly (<0.1%) and the subglacial channel area increased insignificantly (<0.1%) across this range of YE . These effects are comparable to those of E, E^* , which we also varied over one order of magnitude, and T , which changed the englacial flow-law parameter A by approximately a factor of 6.

In contrast to the above parameters, we find that moulin geometry is strongly sensitive to the choice of basal ice softness (prescribed A_{basal}) and the friction factors used within the moulin (f_m and f_{oc}). Melting due to the dissipation of turbulent energy is partially controlled by the friction factors chosen for the moulin walls. The friction factor above the water line (f_{oc} ,

“open channel”) does not significantly affect moulin water level (<0.1% change for f_{oc} variations over two orders of magnitude), moulin volume (4%), moulin water storage (2%), or subglacial channel area (<0.1%) over either long or diurnal timescales (Fig. 4m3m–n and 5m4m–n). However, like the deformational parameters, the open channel friction factor does affect moulin radii, with the major radius growing by 36% as the open channel friction factor increases over two orders of magnitude, and the minor radius decreasing by 27%. This dampens the diurnal variability in both radii: (Fig. 4m).

Increasing the friction factor below the water line (f_m) had similar effects to changing f_{oc} . Increasing f_m by two orders of magnitude increased the cross-sectional area of the moulin by 106%, via a 15% increase in the major radius and a 95% increase in the minor radius. The water volume increased by 116% and the storage capacity increased by 100% (Fig. 4k3k–l) while the equilibrium water level and the subglacial channel area changed by <0.1%. Increasing the underwater friction factor f_m also increased the diurnal variability of the moulin capacity and water storage (Fig. 5k4k–l) by increasing the diurnal differential melt rate.

The two parameters which have the largest impact on moulin water level are the basal ice softness (A_{basal}) and the moulin location on the ice sheet, described jointly by the ice thickness (H) and distance from the terminus (L). This sensitivity indicates an interplay among these parameters, the subglacial hydraulic gradient, and moulin water level. We varied basal ice softness A by two orders of magnitude. Softer basal ice increased the size and storage capacity of the moulin: the major radius by 21%, the minor radius by 25%, the total capacity by 88%, and the stored water volume by 112% (Fig. 4i3i–j). These changes also increased the equilibrium water level by 57% and the subglacial channel area by 24%, unlike tests on englacial parameters (E , T , and Y), which did not affect the water level and/or subglacial channel area. These changes occur because softer basal ice increases the rate of subglacial creep closure, which reduces subglacial channel cross-sectional area, which reduces water throughflow in the moulin and increases water level, which in turn reduces the amount of viscous and elastic radial closure in the moulin. Increasing the basal ice softness A_{basal} to approximately $10^{-23} \text{ Pa}^{-3}\text{s}^{-1}$ increases the diurnal variability in the sizes of the subglacial channel and moulin (Fig. 5i4i–j); however, increasing A above this value causes moulin water levels to rise high enough that diurnal fluctuations are truncated by the ice thickness. Results at these values are therefore not shown on Fig. 4. The precise value of A where this transition occurs depends on other traits of the moulin, including ice thickness, resulting in an observed decrease in diurnal range that would not be present in thicker ice (Fig. 4j).

We co-varied ice thickness and distance from terminus using a parabolic approximation for a perfectly plastic ice surface profile (Cuffey and Paterson, 2010). Variations in ice thickness from 670 to 1570 m (80%) increase equilibrium subglacial conditions by 20% and increase equilibrium water levels by 107% (Fig. 4e3e–p). Increasing ice thickness and distance from the terminus increases the moulin major and minor radii by 4%, increases moulin volume by 97%, and increases moulin water storage by 114% (Fig. 4p). We also find significant increases in diurnal variability in subglacial channel size (28%), water level (105%), moulin radii (major radius 85% and minor radius 22%), moulin volume (130%), and moulin water storage (140%) in thicker ice farther from the terminus (Fig. 5e4e–p).

Overall, we find that MouSh-modeled moulins are primarily sensitive to the friction factors for water flow through the moulin, basal ice softness, and location on the ice sheet (ice thickness and distance from the terminus). The results are less

sensitive to englacial material factors that govern elastic and viscous deformation. The observed sensitivity to the ice thickness and distance from terminus signals that moulin geometry can vary spatially. The sensitivity to friction factors and basal ice softness indicates that the values of these poorly constrained parameters should be carefully chosen and kept in mind when interpreting model output.

3.2.21 Contributions to moulin shape

Figure 65 illustrates the role of each process that changes moulin radius under equilibrium conditions: phase change, viscous deformation, and elastic deformation. We use standard values for all parameters (Table 1) and we vary ice thickness and distance from terminus. We find that moulin shape is quite similar across different ice thicknesses, while mean water level, moulin capacity (Fig. 6a5a–e) and the diurnal range in moulin radius (Fig. 6g5g) increase with ice thickness. We also analyze temporal variations in each process (Fig. 6g5g). The times of maximum melt and maximum viscous closure are slightly offset, with peak melting occurring during the most rapid decline in viscous deformation (Fig. 6g5g). This offset aligns with the rising limb of the input hydrograph, when the moulin is small and increases in Q_{in} raise water level and, in turn, elevate englacial melt rates and reduce viscous deformation.

Melt rates both above and below the water line contribute to moulin growth (Fig. 6f5f–g). Melt above the water level occurs due to stream or waterfall erosive processes, which in MouSh occur only within a fraction of the total circumference (Fig. 2d2a,d), which manifests as growth of the major radius. The actual rate of melting, however, is also dictated by the area over which water flow occurs, which under our parameterization is related to the cross-sectional area of the moulin at any given depth (Fig. 6f5f).

Elastic deformation, like viscous deformation, closes the moulin except when the water level is above flotation. Elastic deformation rates are generally smaller than viscous rates, except between ~100–300 meters above the bed, where viscous deformation is minimized by cold ice temperatures (Lüthi et al., 2015; Ryser et al., 2014); (Lüthi et al., 2015; Ryser et al., 2014). Diurnally, elastic deformation varies with a similar pattern to viscous deformation, though over less range.

3.3 Moulin shape in different environments

We modeled the seasonal growth and collapse of moulins in a range of environments across the Greenland Ice Sheet using realistic melt forcings derived for the 2019 melt season (Section Sect. 2.4.1 and Sect. 2.5.3). These model runs varied with respect to ice thickness, moulin distance from the terminus, base flow, and the magnitude, diurnal range, and seasonal evolution of supraglacial inputs (Table 2; Fig. 7a6a). Overall, we find that moulin setting affects the scale of diurnal and seasonal variability in the size and water capacity of moulins as well as the evolution of subglacial channels (Fig. 76 and 87).

The sizes of all three modeled moulins reach equilibrium with the melt forcing within 10 days of the onset of the melt seasons (Fig. 7b6b–c). As the water flux increases over the next few weeks, each moulin grows in response to increasing supraglacial inputs, both diurnally and with a long-term trend (Fig. 7e), though this growth is more significant in thicker ice

(Fig. 8m-p6c and Fig. 7). The subglacial channel grows with a similar pattern, but interestingly, the setting and fluxes of Basin 1 and Basin 2 result in very similar subglacial channel cross-sectional areas (Fig. 7d). However, moulin water level shows diurnal variations superimposed on a relatively constant base value throughout the melt season (Fig. 7b) despite different water levels (Fig. 6d).

Though the three moulins all evolve in a similar fashion, there are differences in moulin water capacity, equilibrium water level (Fig. 7b), overall moulin geometry (Fig. 8j), and the magnitude of englacial deformation (Fig. 9g). Diurnal variation in moulin capacity is slightly larger in thicker ice, due to higher rates of deformation and greater melt rates within both the moulin (Fig. 9a). While the relative fraction of phase change (melting) to deformation is generally near 1, in the thickest ice, the absolute value of total deformation is generally higher than that of the phase change term (8a). This indicates that viscous and elastic opening play a role in maintaining moulin geometry when occurs because daily water levels level fluctuations are greater in thicker ice due to the non-linearity of ice creep in conjunction with the linearity of melt-driven growth (Sect. 3.1). Furthermore, in thick ice, visco-elastic deformation plays a relatively larger role in moulin evolution (dark purple line in Fig. 8b), for the same reason. The only exception is during periods of low diurnal variability in Q_m , which occurred around Day 30 of the 2019 melt season (Fig. 6a). During this period, the minimum daily supraglacial inputs are quite high (Fig. 9b), otherwise, the 7a); moulin would continually close water levels stayed near flotation for a few days (Fig. 7b), keeping effective pressure near zero and retarding deformation, and slightly increasing melt rates (Fig. 8b). In this case, viscous deformation hovers around zero (though causing moulin opening), resulting in a high ratio of elastic to viscous deformation and a high ratio of phase change to viscous deformation (purple line in Fig. 9b). There is an associated growth in moulin capacity (Fig. 7c). Ultimately, this is a response to multiple days where melt inputs do not exhibit substantial diurnal variability.

The ratio of elastic to viscous deformation generally ranges from ~0.4 to ~0.87, depending on ice thickness (Fig. 9b8b). Elastic deformation rates in the moulin depend on a linear function of ice thickness, while viscous rates are related to ice thickness cubed. Thus, at lower elevations, the elastic contribution is maximized (~0.8 of viscous deformation), while at high elevations, significant increases in viscous closure lowers the relative contribution of elastic deformation (~0.4 of viscous deformation). This increase in viscous closure in thick ice also minimizes subglacial channel size in thick inland ice (Fig. 7d6d), despite closure rates being retarded by daily periods of above overburden water levels persistently at or near the ice surface pressures.

Each moulin has a different equilibrium daily mean capacity (Fig. 7c). This, in addition to differences in supraglacial inputs, ensures that daily moulin water level variations are substantially different across moulins. Basin 1 exhibits the largest variation in daily moulin water level, followed by Basin 2 (Fig. 40a9a). Basin 3 shows the lowest daily change; however, this is due at least in part to the fact that water overtops the moulin nearly daily (Fig. 7b6b and 8m7m-n). Changing water levels drive changes in moulin and subglacial capacity. Over the melt season, daily change in moulin capacity can be as low as 5%

during lulls in diurnal melt variability (Basin 3) or as high as ~~3231%~~ following a recovery from a low melt day (Basin 1; Fig. ~~40b9b~~). However, in general all moulin display a similar daily change in capacity of ~~~10–25%, with no clear pattern as to which elevation experiences greater short-term changes in moulin capacity, 20%~~.

The subglacial system undergoes diurnal variations in channel size between 1 and 21% (Fig. ~~40e9c~~). These changes are similar in magnitude to daily capacity changes within the moulin but exhibit more variability across ice thicknesses and are related to the daily changes in moulin water level (Fig. ~~40a9a~~). This suggests that the time evolution of moulin geometry dampens the diurnal pressure fluctuations that drive subglacial channel growth and collapse. Evidence for this can be seen in the temporal pattern of moulin water level and subglacial channel cross-sectional area (Fig. ~~40a and 40e9a,c~~). To test this idea, we compared results from static and time-evolving moulin (Section ~~Sect.~~ 3.4).

3.4 Comparison to cylindrical moulin

~~To examine the role moulin evolution plays in modifying the subglacial hydrologic system, we compared moulin water levels, moulin capacity, and subglacial channel size between model runs with a fully evolving moulin and runs with a static cylindrical moulin. We performed these tests with realistic melt inputs based on the 2019 melt season (Section 2.4), at moulin with low and moderate ice thicknesses (553 m – Basin 1 and 741 m – Basin 2). We defined the radius of the fixed cylinder as 1.4 m and 1.3 m for Basin 1 and 2, respectively. This results in fixed cross-sectional areas ($\sim 6 \text{ m}^2$ and $\sim 5 \text{ m}^2$) within the range of the spatially invariant moulin cross-sectional areas $\sim 2\text{--}10 \text{ m}^2$ often prescribed in subglacial models (e.g., Andrews et al., 2014; Banwell et al., 2013; Bartholomew et al., 2012; Cowton et al., 2016; Meierbachtol et al., 2013; Werder et al., 2013). Inter-comparison of these runs allows us to examine the role moulin geometry has on subglacial pressures (Covington et al., 2020; Trunz et al., in review).~~

To examine the role moulin evolution plays in modifying the subglacial hydrologic system, we compared moulin water levels, moulin capacity, and subglacial channel size between model runs with a fully evolving moulin and runs with a static cylindrical moulin. We performed these tests with realistic melt inputs based on the 2019 melt season (Sect. 2.4.1, 2.5.3), at moulin with low and moderate ice thicknesses (553 m – Basin 1 and 741 m – Basin 2). We defined the radius of the fixed cylinder as 1.4 m and 1.3 m for Basin 1 and 2, respectively. This results in fixed moulin cross-sectional areas ($\sim 6 \text{ m}^2$ and $\sim 5 \text{ m}^2$) that are within the range of the spatially invariant moulin cross-sectional areas $\sim 2\text{--}10 \text{ m}^2$ often prescribed in subglacial models (e.g., Andrews et al., 2014; Banwell et al., 2013; Bartholomew et al., 2012; Cowton et al., 2016; Meierbachtol et al., 2013; Werder et al., 2013). Inter-comparison of these runs allows us to examine the role moulin geometry has on subglacial pressures (Covington et al., 2020; Trunz, 2021).

Comparison of water level, moulin capacity, moulin water storage, and subglacial cross-sectional area between fixed and evolving moulin show differences on both the diurnal and seasonal times scales (Fig. ~~4410~~). Moulin water levels (~~fixed-variable—fixed~~) can be substantial (Fig. ~~44a10a–b~~), with short term differences driven by variable melt conditions reaching a maximum of ~~-12997~~ m (Basin 1) and ~~-177145~~ m (Basin 2), but values can also be negative, indicating that the realistic run

moulin has higher water levels, up to 46 m for Basin 1 and 25 m for Basin 2. The long-term daily average differences are -286 m and -4935 m for Basin 1 and Basin 2, respectively. These differences are driven by a combination of differences in moulin capacity and subglacial channel size (Fig. ~~11b-e~~10c-f) and are despite a total increase in the meltwater input relative to a fixed into an evolving moulin, due to melt generated from turbulent dissipation (~~Fig. 11d~~), less than 2%. These results indicate that diurnal variability is an important component not effectively represented with a cylindrical moulin.

Generally, the evolving moulin is larger, (Fig. 10c-d), stores more water, generates more water through internal melting, and maintains a larger subglacial channel (Fig. ~~11b-d~~10e,f), which all contribute to the observed difference in water levels. Midway through the melt season, the evolving moulin exhibits capacities and storage volumes 10–50% only slightly larger than those of the fixed cylinder, but these capacity differences are exacerbated during higher Q_m values (Fig. ~~12b~~10c-d). As meltwater inputs taper at the end of the melt season (day ~97 in Fig. 11b100), the capacity and storage volume in the evolving moulin falls below that of the fixed cylinder, whose volume does not adjust in response to the forcings: (Fig. 10c-d). This seasonal evolution is consistent between the two ice thicknesses tested.

The capacity differences between the variable and fixed moulin contribute directly to dampening the supraglacial input signal and dampening of moulin water levels. This contributes directly to, in turn, drives an increase in subglacial channel size (Fig. ~~11d~~10e-f), both diurnally and over the season. The seasonal difference between the variable and fixed moulin forcing is relatively constant, though punctuated by dips associated with reduced moulin water level differences (Fig. ~~11a~~10a-b).

3.4.5 Impact of model choices on moulin geometry

Chosen parameterizations have the potential to impact the representation of moulin water level and capacity (Supplement S2). Overall, we find that a circular geometry has limited impact on moulin water level with the circular geometry having water levels that are less than 3 m higher than the egg-shaped geometry, though in nearly all instances the difference is less than 0.5m (Fig. S5a); however, the impact on capacity is slightly larger (the circular moulin is up to 47% smaller) and displays a seasonal trend as the egg-shaped moulin elongates along its elliptical axis (Fig. S5b). Altering the deviatoric and shear stresses used in the calculation of elastic deformation results in minimal changes, primarily above the water line. Moulin water levels are typically within 0.25 m of the control run (Fig. S5c). Prescribing the surface stresses to be zero results in a maximum increase in moulin capacity of less than 10% (Fig. S5d).

In contrast to the previous choices, the distance from the terminus (L) and the prescribed base flow (Q_{base}) can have a substantial impact on moulin water level and capacity (Fig. S5e-h). Distance from the terminus is defined by the position of a given moulin on the ice sheet, and as such is not a choice or parameter per se; and base flow is used here to mitigate the use of a simplistic subglacial hydrology model. Finally, we examine the impact of fixing the subglacial channel cross-sectional area S . Experimental results using a fixed S and a seasonally evolving melt curve resulted in extremely low and extremely high water levels resulting in complete moulin collapse or runaway growth, respectively. Therefore, we explore the impact of fixing S using a constant mean Q_m with an overlaid diurnal variability (Supplement Sect. S2.2.6). When the fixed S is smaller than a

variable S , moulin water levels are higher and exhibit less diurnal variability while moulin capacity is larger (Fig. S6). Further details are included in the Supplement S2.

4 Discussion

4.1 Formation mechanisms Timescales of moulins

~~Moulins can form through multiple processes, including cutmoulin formation and closure of supraglacial streams (Gulley et al., 2009) and vertical hydrofracture through cold ice (Das et al., 2008). The formation mechanism dictates the initial geometry, which then evolves rapidly in response to a range of ice and melt processes to reach equilibrium geometry.~~ evolution

~~On the Greenland Ice Sheet, moulin locations are generally disassociated with crevasse fields (Colgan et al., 2011; Phillips et al., 2011; Smith et al., 2015) and, instead, occupy regions of compressional stresses, including supraglacial lake basins (Catania et al., 2008; Poinar and Andrews, 2020; Stevens et al., 2015). Episodic local water inputs to the bed can instigate transient stresses that exceed the fracture toughness of ice, forming crevasses in otherwise compressional regimes (Christoffersen et al., 2018; Hoffman et al., 2018; Poinar and Andrews, 2020). Depending on the meltwater influx, crevasses that form moulins may propagate to the bed in less than a day (Stevens et al., 2015) or over multiple years (Poinar, 2015). Regardless of formation timescale, all known moulins on the Greenland Ice Sheet are thought to be vertical or near vertical.~~

We consider the formation timescales of moulins in the context of the shape evolution of a mature moulin. Using MouSh, we find that in the absence of external forcing, such as time-variable Q_{in} , the size of a moulin reaches its equilibrium value in ~ 1 –10 days. depending on ice and supraglacial input conditions (Fig. 5g, Fig. S2 and Fig. S3). This relaxation time is comparable to the Maxwell time for ice (10–100 hours), as expected for a linear visco-elastic system. Our relaxation time also compares well to the equilibration timescale defined by Covington et al. ~~(2020)~~(2020) for their modeled moulin – subglacial conduit system, which Trunz ~~et al. (in review)~~(2021) found to be 1–20 days. The most realistically sized moulins in Trunz ~~et al. (in review)~~(2021) had relaxation times closer to 1 day. Their modeled ~~channel~~ system was governed solely by melt and viscous deformation and lacked elastic deformation; this may explain their modestly longer relaxation time compared to ours.

~~If the process of moulin formation occurs on a timescale shorter than the 1–5 day relaxation time, the formation process likely will not influence the overall form of the englacial system at equilibrium. This time range includes hydrofracture during rapid lake drainage (~ 2 hours) and slow lake drainage (< 6 days, e.g., Selmes et al., 2011), and likely also the reactivation of existing moulins in ensuing melt seasons, which, based on the timing difference between surface melt onset and ice acceleration, occurs on a multi-day timescale (Andrews et al., 2018; Hoffman et al., 2011; Zwally et al., 2002). On the other hand, moulin formation by cut and closure occurs over years to decades (Gulley et al., 2009), well above the MouSh relaxation time and the Maxwell time for ice. The interdependence of formation and evolution of these moulins gives us less confidence in applying our model to moulins with cut and closure origins. Those moulins primarily occur in temperate near-surface ice within polythermal glaciers (Gulley et al., 2009) and have not been reported on the Greenland Ice Sheet.~~

4.2 Comparison of modeled and observed moulin geometries

Field observations suggest that moulin geometry evolves a high degree of complexity. Observations include anecdotes of difficulty deploying sensors to the bottom of a moulin, which suggests the presence of kinks, ledges, knickpoints, and other twists (Andrews et al., 2014; Covington et al., 2020; Cowton et al., 2013). Complex geometry revealed during mapping moulins above the water line further suggests that moulins are not simply vertical cylindrical shafts (Covington et al., 2020; Moreau, 2009).

The MouSh model suggests that the energy transfer from turbulent meltwater entering the moulin to the surrounding ice drives highly spatially variable melt rates above the water line. We incorporated the open-channel melt module to allow a large opening to emerge above the water line (Fig. 6a–e and 8). If the process of moulin formation occurs on a timescale shorter than the 1–10-day relaxation time, the formation process likely will not influence the overall form of the englacial system at equilibrium. This time range includes hydrofracture during rapid lake drainage (~2 hours) and slow lake drainage (<~6 days, e.g., Selmes et al., 2011), and likely also the reactivation of existing moulins in ensuing melt seasons, which, based on the timing difference between surface melt onset and ice acceleration, occurs over multiple days (Andrews et al., 2018; Hoffman et al., 2011). On the other hand, moulin formation by cut-and-closure occurs over years to decades (Gulley et al., 2009), well above the MouSh relaxation time and the Maxwell time for ice and are more likely to create subvertical englacial channels. The interdependence of formation and evolution of these moulins gives us less confidence in applying our model to moulins with cut-and-closure origins. Those moulins primarily occur in temperate near-surface ice within polythermal glaciers (Gulley et al., 2009) and have not been reported on the Greenland Ice Sheet.

4.2 Comparison of modeled and observed moulin geometries

Field observations suggest that moulin geometry evolves a high degree of complexity. Observations include anecdotes of difficulty deploying sensors to the bottom of a moulin, which suggests the presence of kinks, ledges, knickpoints, and other twists (Andrews et al., 2014; Covington et al., 2020; Cowton et al., 2013). Complex geometry revealed during mapping moulins above the water line further suggests that moulins are not simply vertical cylindrical shafts (Covington et al., 2020; Moreau, 2009).

The MouSh model suggests that the energy transfer from turbulent meltwater entering the moulin to the surrounding ice drives highly spatially variable melt rates above the water line. We incorporated the open-channel melt module to allow a large opening to emerge above the water line (Fig. 5a–e and 7). When we run MouSh without the open-channel module (Sect. 2.3.2.4.3), the surface expression of the moulin is much smaller than observed in remote sensing images and in some cases, the moulin will pinch closed at the ice-sheet surface. The open channel module also permits the development of an egg-shaped geometry, which is supported by seismic observations and a resonance model of a moulin, which suggest that the moulin increased in ellipticity over time (Röösli et al., 2016).

The value of the open-channel friction factor and the size of the spatial footprint over which melting occurs directly affects the size of the upper, air-filled chamber of the moulin, which differs from when treated as circular (Fig. S5b). MouSh consistently predicts ledges at the top and bottom of a consistent diurnal range in water level. Thus, we infer that energetic subaerial water flow drives formation of moulin complexity above the water line, and diurnal fluctuations around a steady multi-day water level drive ledge formation through underwatera differential in melting and visco-elastic deformation: above and below the water line. Energetic water flow is commonly observed at stream-fed moulins near the peak of the melt season (Pitcher and Smith, 2019)(Pitcher and Smith, 2019) or during and immediately following rapid lake drainage (Chudley et al., 2019);(Chudley et al., 2019). This suggests that complex moulin geometries form during periods of relatively consistent water supply. Conversely, multi-day rises in water level, driven by either the surface water supply (Q_{in}) or the basal water supply (baseflow), can erase geometric complexities such as ledges, as seen in MouSh results during a melt event (Fig. 87).

Above the water line, explored moulins in Greenland show highly variable shapes from moulin to moulin (e.g., Covington et al., 2020). Some moulins, for example the FOXX moulin, are nearly cylindrical within the explored depth (~100 m), with radii comparable to what we model (~2 meters). Others, like the Phobos moulin, open some tens of meters below the surface to large caverns with radii approaching 10 meters, a similar morphology to karst caves with narrow entrance shafts. MouSh can produce large openings above the water line if we use a suitably large open channel friction parameter, although we lack a narrow entrance shaft. These differences are due to the inability of model parameterizations to represent complex geometries such as scalloping, plunge pools and knickpoint migration (Gulley et al., 2014; Mankoff et al., 2017). Indeed, instead of modeling processes above the water line as turbulent open flow, they could be modeled using geomorphic parameterizations to model waterfall migration, perhaps resulting in the clearer development of steps and plunge pools. However, these parameterizations are generally based on contact abrasion and debris cover (e.g., Scheingross and Lamb, 2017), making the translation to thermal erosion difficult.

Below the water line, MouSh results indicate that a cylinder is a reasonable representation for newly formed moulins in Greenland. However, there are two caveats. First, moulin cross sectional area, and thus water storage capacity, can vary substantially over the course of a day or season (Fig. 11). Second, in instances where moulins are reactivated over multiple melt seasons (Chu, 2014; Smith et al., 2017), there can be substantial deformation, as suggested by cable breakage in boreholes (Ryser et al., 2014; Wright et al., 2016).

Above the water line, explored moulins in Greenland show highly variable shapes from moulin to moulin (e.g., Covington et al., 2020). Some moulins, for example the FOXX moulin, are nearly cylindrical within the explored depth (~100 m), with radii comparable to what we model (~2 meters). Others, like the Phobos moulin, open some tens of meters below the surface to large caverns with radii approaching 10 meters, a similar morphology to karst caves with narrow entrance shafts (Covington et al., 2020). MouSh can produce large openings above the water line if we use a suitably large open channel friction parameter, although we lack a narrow entrance shaft and substantial vertical variability. These differences are due to the inability of model parameterizations to represent complex geometries such as scalloping, plunge pools and knickpoint

migration (Gulley et al., 2014; Mankoff et al., 2017). Indeed, instead of modeling processes above the water line as turbulent open flow, they could, in the future, be modeled using geomorphic parameterizations to model waterfall migration, perhaps resulting in the clearer development of steps and plunge pools. This would require development and inclusion of a supraglacial channel model as well.

Below the water line, MouSh results indicate that a cylinder is a reasonable representation for newly formed moulins in Greenland. However, there are two caveats. First, moulin cross-sectional area, and thus water storage capacity, can vary substantially over the course of a day or season (Fig. 9) and features such as englacial crevasses and reservoirs may be present (e.g., McQuillan and Karlstrom, 2021). Second, in instances where moulins are reactivated over multiple melt seasons (Chu, 2014; Smith et al., 2017), there may be substantial deformation, as suggested by cable breakage in boreholes (Ryser et al., 2014; Wright et al., 2016).

Observations show a wide range of moulin volumes above the water line, and moulin volumes predicted by MouSh are sensitive to the value of the open-channel friction factor. Given the flexibility of model results, we should continue to rely on field exploration to measure moulin size and geometry above the water line and make efforts to constrain the parameters that affect sub-seasonal growth and collapse. MouSh results below the water line are less sensitive to uncertain parameter values, so we rate direct observations of underwater exploration of Greenland moulins at a lower priority geometry would be less relevant for model validation than subaerial observations. Overall, results from the MouSh model demonstrate that moulin geometry evolves substantially over diurnal to seasonal timescales and varies with ice conditions.

4.3 Diurnal water level oscillations and moulin size

Moulin geometry can directly alter the relationship between meltwater inputs and moulin water level changes – the primary driver of subglacial channel evolution (Andrews et al., 2014; Cowton et al., 2013). Field measurements of moulin water levels indicate diurnal oscillations of 3–12% (Covington et al., 2020), ~25% (Andrews et al., 2014) and >20% (Cowton et al., 2013) of overburden pressure. These diurnal fluctuations are larger than those observed in boreholes, which are generally thought to sample inefficient components of the subglacial hydrologic system (Andrews et al., 2014; Meierbachtol et al., 2013; Wright et al., 2016). (Andrews et al., 2014; Cowton et al., 2013). Field measurements of moulin water levels indicate diurnal oscillations of 3–12% (Covington et al., 2020), ~25% (Andrews et al., 2014), and >20% (Cowton et al., 2013) of overburden pressure with mean water levels of ~70% of overburden. These diurnal fluctuations are larger than those observed in boreholes, which are generally, though not always, thought to sample inefficient components of the subglacial hydrologic system (Andrews et al., 2014; Meierbachtol et al., 2013; Wright et al., 2016).

Our model results agree well with observations of moulin water level: diurnal fluctuations of 15–25% of overburden pressure, with larger oscillations in thicker ice. To explain larger-than-expected daily oscillations (~10%) in thinner ice, Covington et al. (2020)(2020) incorporated moulin cross-sectional area as a free parameter into their model. Matching field measurements of water level required a modeled moulin radius of ~5 m (~75 m² cross-sectional area) at ice thickness 500 m

and a much larger moulin (radius ~ 20 m and cross-sectional area ~ 1500 m²) at ice thickness 700 m (Covington et al., 2020);(Covington et al., 2020). For comparison, MouSh predicts average radii of ~ 1.3 to 1.4 m (~ 5 m² cross-sectional area) at these ice thicknesses using parameters described in Table 2, including substantially larger meltwater inputs compared to Covington et al. (2020);(2020). The drastic differences in moulin size despite similar variations in diurnal water level variability between our study and Covington et al. (2020);(2020) cannot easily be attributed to a single factor, but may be explained by our limited ability to model processes above the water line, our inclusion of ~~baseflow~~;base flow (Fig. S5g–f), substantial differences in meltwater input; (e.g., Figs S2 and S3), fluctuations in moulin capacity, or that their measured water levels were not from the same moulin they mapped englacially. Nevertheless, we observe substantial differences in water level between fixed and variable geometry moulins that are dependent on supraglacial inputs and at both elevations ice conditions (Fig. H10). Water levels are less variable and generally lower in the evolving moulins compared to the fixed cylindrical moulin. ~~We also note that for both ice thicknesses, the fixed moulin frequently overtops while the evolving moulin does not (Fig. 11a) for the same melt, ice, and subglacial conditions.~~ Thus, to match observed moulin water level fluctuations without evolving the moulin geometry, a fixed cross-sectional area substantially larger than the associated subglacial channel may be necessary, as reported in Covington et al. (2020).

4.3 Magnitude of viscous moulin deformation

Viscous and elastic deformation drive moulin closure. The role of elastic deformation in the glacial hydrologic system is discussed below (Sect. 4.4); viscous deformation is the primary mechanism of moulins, boreholes and subglacial channels (e.g., Catania and Neuman, 2010; Paterson, 1977, Shreve, 1972), with viscous deformation dependent on local effective pressure, ice characteristics, and the geometry of the feature of interest (Flowers, 2015). Viscous deformation within our moulin varies in response to meltwater inputs (Fig. 5g and Fig. 8a) with the highest deformation rates occurring at the water line (Fig. 5f) because at the water line, inward cryostatic pressure is least offset by outward hydrostatic pressure (see Eq. 3).

During our realistic runs, viscous deformation can exceed 0.5 m d⁻¹ for short periods at the highest elevation (Fig. 8a). These deformation rates are substantially larger than measured borehole deformation rates for the primary reasons that boreholes are often at or above flotation due to high subglacial water pressures (e.g., Ryser et al., 2014) or because creep measurements are recorded in much smaller boreholes in colder ice (e.g., Paterson, 1997).

A previous moulin modeling effort focused on understanding moulin closure rates (Catania and Neumann, 2010). Their results indicate that an air-filled moulin will close within a single day at the bed. However, in this instance there is no opposing hydrostatic pressure. While our modeled closure rates are similar to those calculated by Catania and Neuman (2010) near the surface, the moulins modeled here always contain water even at the end of the melt season (Fig. 6b). This continued retention of meltwater is in line with borehole observations that subglacial pressures tend to be highest outside the melt season (Downs et al., 2018) and preclude the presence of completely air-filled moulins in areas where viscous deformation rapidly shuts down the hydrologic system as supraglacial inputs fall.

4.4 The role of elastic deformation in ice sheet hydrology

Our model results indicate that the equilibrium moulin geometry is dictated by a balance of visco-elastic deformation and turbulence-driven melting (Fig. 5 and Fig. 8). In both the sensitivity study and realistic model runs, visco-elastic deformation generally closes the moulin, while melting of the surrounding ice consistently opens the moulin. The exception is when moulin water levels exceed flotation, in which case all three mechanisms open the moulin. In all model runs, we find that elastic and viscous deformation are of the same order of magnitude, and that the elastic mode can be between 40% and 80% of the viscous deformation (Fig. 5g and Fig. 8). The importance of elastic deformation holds even in the bottom few hundred meters of the ice column, where stress conditions are similar to those in subglacial models (Fig. 5f). However, the relative importance of viscous and elastic deformation in closing the moulin is also dependent on the values of Young's modulus and viscous enhancement factor (Fig. 5 and Fig. 8). Despite extensive study of these parameters, their values are difficult to constrain. Currently, the space of viscous and elastic parameter values could conceivably allow either elastic or viscous deformation to dominate the closure of a moulin. This underscores the importance of including both modes in the MouSh model.

Current subglacial hydrology models represent subglacial channel development (opening) by turbulent energy dissipation and destruction (closing) by viscous deformation alone. Some more recent work involving elastically responding storage elements or elastic flexure of the ice sheet has occurred (Clarke, 1996; Dow et al., 2015), and there have been efforts to use elastic deformation or fluid compressibility to improve numeric stability of channel equations (Clarke, 2003; Spring and Hutter, 1981, 1982). Interestingly, Clarke (2003) chose to use fluid compressibility due to model integration times. Yet, elastic deformation has generally been omitted from current models of subglacial channelization, even when modeling rapid changes in meltwater inputs (< 1 day; e.g., Hewitt, 2013; Hoffman et al., 2016; Werder et al., 2013). This choice is likely because the role of elastic deformation was considered negligible over timescales of subglacial evolution (e.g., days to weeks). However, the importance of elastic deformation in diurnally closing moulins, particularly in thinner ice (Fig. 8b), suggests that its exclusion from subglacial channel models could result in the underestimation of channel closure rates when water levels are below flotation.

This leads us to ask why elastic deformation is absent from subglacial models, particularly because its importance relative to viscous deformation is difficult to constrain given the current range of observed Young's modulus (Vaughan, 1995). Hypothetical subglacial channel models that included elastic deformation alongside viscous deformation would show less temporal asymmetry, particularly in thinner ice, where channel closure may be strongly dictated by elastic deformation. Elastic-incorporating models would also likely predict larger diurnal variations in channel size and moulin water level. This in turn would incite stronger local pressure gradients at the bed, increasing connectivity between the channel and the surrounding distributed system.

4.5 Moulin geometry and the englacial void ratio

Subglacial hydrology models use an englacial void ratio parameter to represent bulk storage and release of meltwater in the englacial system (see Flowers and Clarke (2002) for the best description). The englacial void ratio allows subglacial models to resolve observed diurnal fluctuations in water pressure and, if coupled to a dynamical ice model, corresponding diurnal variations in ice flow. This parameter represents bulk behavior and is usually set constant over the model domain, yet it must be tuned by comparing to local observations (e.g., Bartholomaeus et al., 2011; Hoffman et al., 2016; Werder et al., 2013).

Subglacial hydrology models use an englacial void ratio parameter to represent bulk storage and release of meltwater in the englacial system (see Flowers and Clarke (2002) for the best description). Because the englacial void ratio acts as short term, pressure dependent, storage for subglacial models, it can improve the representation of diurnal water pressure fluctuations in subglacial models (Flowers and Clarke, 2002) and, if coupled to a dynamical ice model, corresponding diurnal variations in ice flow. This parameter represents bulk behavior and is usually set constant over the model domain, yet it must be tuned by comparing to local observations (e.g., Bartholomaeus et al., 2011; Hoffman et al., 2016; Werder et al., 2013). The inclusion of moulins, potentially in addition to time varying representation of englacial fractures (Gajek et al., 2021), that evolve in response to meltwater inputs and subglacial pressures could reduce subglacial model dependence on this highly parameterized englacial storage, particularly in light of observations of time varying englacial features (Church et al., 2020) and meltwater content (e.g., Vankova et al, 2018).

Recent work suggests that fluctuations in water level are controlled by the size of the moulin near the water level (~~Trunz et al., in review~~);(Trunz, 2021): moulins with larger cross-sectional areas have lower diurnal variability in water level, if given the same melt input. Furthermore, our results suggest that the amount of water stored in a moulin is highly dependent on local conditions, such as water pressure on daily to seasonal timescales, and ice thickness (Fig. 6c and Fig. 7). Thus, we explore the possibility that detailed model-based information on moulin sizes and shapes could inform the englacial void ratio used in subglacial hydrology models. This would allow time dependence and finer spatial variation, including in the vertical dimension as well as horizontal, than is currently possible with a bulk parameter. Periods of increased supraglacial inputs can require a sizable increase in englacial void ratio for subglacial models to accurately predict moulin water level (Hoffman et al., 2016). ~~During these times, MouSh predicts rapid growth in moulin capacity (Fig. 7 and 8).~~(Hoffman et al., 2016). During these times, MouSh predicts rapid growth in moulin capacity (Fig. 7 and Fig. 8). This correspondence suggests plausible close ties between moulin size and the englacial void ratio: moulin size modifies englacial storage spatially and temporally.

MouSh can be used to infer both moulin size and shape, which would effectively change the englacial void ratio in all three spatial dimensions and time. The shape of the moulin imposes new temporal variability on water level and subglacial channel size: moulins with large near-surface chambers that funnel down to become narrower at the water line, for instance, have lower-magnitude and smoother variations in water level compared to cylindrical moulins, whereas moulins with small surface openings that widen toward the water line have larger and peakier water-level variations (~~Trunz et al., in review~~);(Trunz, 2021). Thus, when the shape of a moulin is explicitly resolved, any assumed linear relationship between melt

input rates and the range or pattern of oscillations in water level and subglacial channel size breaks down. The relationship also changes with the water level in the moulin; hence it varies in time.

MouSh demonstrates that moulin capacity can vary greatly both seasonally and during short periods of large variability in supraglacial input. Moulin growth rates are largest particularly when water levels are above flotation, maximizing turbulent melting and outward visco-elastic deformation. Our results show that moulin capacity changes by ~20% daily (Fig. 10) and ~50–100% over the melt season (Fig. 7e6c and Fig. 8b), with larger changes during periods of large supraglacial input variability and at locations with thicker ice. These variations in moulin shape and size may explain difficulties with modeling subglacial behavior during melt events (Cowton et al., 2016), which are sometimes addressed by temporarily increasing englacial storage (Hoffman et al., 2016). Our results with MouSh lead us to recommend that moulin shape and size be modeled alongside the evolution of the subglacial system, especially during periods of large meltwater variability, in order to more accurately predict subglacial water pressures and ice motion.

Practical limits on model complexity or computational costs may preclude fully time-evolving moulin geometries. While not ideal, a static shape is still preferable to a static cylinder (Trunz et al., in review). Therefore, we interpret our moulin shape results (Fig. 8) While not ideal, an arbitrary static shape is still preferable to a static cylinder (Trunz, 2021). Therefore, we interpret our moulin shape results (Fig. 7) to recommend a representative shape for a static moulin. Below the water line, a cylinder is a reasonable approximation, especially in thinner ice or for newly made moulins, for which full-column ice deformation is minimized. Above the water line, moulin shape is widely variable in time, by location, and across parameter combinations. It is especially sensitive to the friction parameter for open-channel flow (Fig. 43m and Fig. 4m), with low friction values making bottle-shaped moulins that have narrow necks above the water line and larger chambers below the water line, and high friction values making goblet-shaped moulins with open rooms and amphitheatres above the water line atop a narrower geometry below the water line. Exploration of Greenland moulins to date has uncovered multiple goblet-shaped moulins and a few instances of near-cylindrical moulins, but no bottle-shaped moulins (Covington et al., 2020; Moreau, 2009; Trunz et al., in review). Overall, our MouSh results support goblet-shaped moulins, although with great variation in the height and width of the upper chamber.

4.5 The role of Limitations of elastic deformation in moulin geometry

~~Our model results indicate that the equilibrium moulin geometry is dictated by a balance of visco-elastic deformation and turbulence driven melting (Fig. 6 and 9). In both the sensitivity study and realistic model runs, visco-elastic deformation generally closes the moulin, while melting of the surrounding ice consistently opens the moulin. The exception is when moulin water levels exceed flotation, in which case all three mechanisms open the moulin. In all model runs, we find that elastic and viscous deformation are of the same order of magnitude, and that the elastic mode can be between 40% and 80% of the viscous deformation (Fig. 6g and 9). The importance of elastic deformation holds even in the bottom few hundred meters of the ice~~

column, where stress conditions are similar to those in subglacial models (Fig. 6f). However, the relative importance of viscous and elastic deformation in closing the moulin is also dependent on the values of Young's modulus and viscous enhancement factor (Fig. 6 & 9). Despite extensive study of these parameters, their values are difficult to constrain. Currently, the space of viscous and elastic parameter values could conceivably allow either elastic or viscous deformation to dominate the closure of a moulin. This underscores the importance of including both modes in the MouSh model.

Current subglacial hydrology models represent subglacial channel development (opening) by turbulent energy dissipation and destruction (closing) by viscous deformation alone. Some more recent work involving elastically responding storage elements or elastic flexure of the ice sheet has occurred (Clarke, 1996; Dow et al., 2015), and there have been efforts to use elastic deformation or fluid compressibility to improve numeric stability of channel equations (Clarke, 2003; Spring and Hutter, 1981, 1982). Interestingly, Clarke (2003) chose to use fluid compressibility due to model integration times. Yet, elastic deformation has generally been omitted from current models of subglacial channelization, even when modeling rapid changes in meltwater inputs (≤ 1 day; e.g., Hewitt, 2013; Hoffman et al., 2016; Werder et al., 2013). This choice is likely because the role of elastic deformation was considered negligible over timescales of subglacial evolution (e.g., days to weeks). However, the importance of elastic deformation in diurnally closing moulins, particularly in thinner ice (Fig. 9b), suggests that its exclusion from subglacial channel models could result in the underestimation of channel closure rates when water levels are below flotation.

This leads us to ask why elastic deformation is absent from subglacial models, particularly because its importance relative to viscous deformation is difficult to constrain given the current range of observed Young's modulus (e.g., Vaughan, 1995). Hypothetical subglacial channel models that included elastic deformation alongside viscous deformation would show less temporal asymmetry, particularly in thinner ice, where channel closure may be strongly dictated by elastic deformation. Elastic incorporating models would also likely predict larger diurnal variations in channel size and moulin water level. This in turn would incite stronger local pressure gradients at the bed, increasing connectivity between the channel and the surrounding distributed system.

4.6 Potential coupled the current MouSh englacial -- subglacial hydrology models model

Moulins occupy a moderate fraction of the channelized englacial-subglacial system and moulins persistently form along subglacial flow pathways (Fig. 3). The subglacial component of MouSh includes an optional baseflow term, which is necessary to produce realistic equilibrium water levels with the realistic supraglacial inputs we prescribed (Fig. 7). The baseflow value we used does not accurately represent any process because our model runs resolve only a single moulin connected to a single channel, whereas in the real world, multiple moulins feed a network of channels. Though idealized, the baseflow term conceptually connects to a number of potential water sources. These include (1) basal melting from geothermal and frictional heating, (2) supraglacial water delivered via nearby moulins that are connected to the same subglacial channel, and (3) water

that moves from the channelized system to the surrounding inefficient system at high pressures and then flows back into the subglacial channel at lower water pressures (Hoffman et al., 2016; Mair et al., 2002, 2002; Tedstone et al., 2015).

Baseflow maintains a larger, less variable subglacial channel. This can alternately be achieved by lessening the local hydraulic gradient, thus increasing the mean water pressure along a given reach. This may locally occur where one subglacial channel enters another in an arborescent network (Fountain and Walder, 1998). MouSh currently does not have an interconnected network of channels; however, this is under development (Trunz et al., 2020).

Water inputs from surrounding moulins is likely the primary origin of base flow at low and moderate elevations. A substantial amount of baseflow, however, is needed to maintain high pressures necessary to keep a moulin open at high elevations (Fig. 7a): up to $30 \text{ m}^3 \text{ s}^{-1}$. There are two potential explanations for this. First, supraglacial drainage basins at higher elevations are generally larger due to muted transmission of basal topography through thick ice (Crozier et al., 2018; Gudmundsson, 2003; Karlstrom and Yang, 2016). Thus, neighboring moulins deliver high fluxes to the subglacial channel. However, the wider spacing of inland moulins reduces their density within the network. The more likely explanation is that a channelized system is a poor representation of subglacial drainage under thick ice, where lower surface slopes reduce the growth rate of subglacial channels, forcing most water flow through the inefficient system (Andrews et al., 2018; Chandler et al., 2013; Dow et al., 2014; Meierbaechtol et al., 2013). Coupling MouSh to a more sophisticated subglacial hydrologic model that includes both the efficient and inefficient systems may therefore be necessary to accurately model moulins in thick ice.

Moulins are a dynamic component of the channelized englacial–subglacial system, and explicitly modeling their evolution can therefore improve the accuracy of englacial–subglacial hydrology models (Sect. 3.4). MouSh currently uses a single subglacial channel to represent the entire subglacial system, limiting its accuracy. An optional baseflow term, which parametrizes subglacial water flow from surrounding regions, improves MouSh performance. This base flow, added directly to the subglacial channel, is necessary to produce realistic equilibrium water levels with the realistic supraglacial inputs we prescribed (Fig. 6a). The baseflow value we used does not explicitly represent any specific process because our model runs resolve only a single moulin connected to a single channel, whereas in the real world, multiple moulins feed a network of channels. The idealized baseflow term conceptually connects to multiple potential water sources, including (1) basal melting from geothermal and frictional heating, (2) supraglacial water delivered via nearby moulins that are connected to the same subglacial channel, and (3) water that moves from the channelized system to the surrounding inefficient system at high pressures and then flows back into the subglacial channel at lower water pressures (Hoffman et al., 2016; Mair et al., 2001, 2002; Tedstone et al., 2015).

The addition of baseflow maintains a larger, less variable subglacial channel. This can alternately be achieved by lessening the local hydraulic gradient, thus increasing the mean water pressure along a given reach. This may locally occur where one subglacial channel enters another in an arborescent network (Fountain and Walder, 1998). MouSh currently does not have an interconnected network of channels; however, this is under development (Trunz, 2021).

We use a highly simplified model of the subglacial hydrology system: a single channel that connects the moulin to the ice-sheet margin. Yet, MouSh results clearly indicate that including and evolving englacial component a moulin can reduce diurnal and long-term subglacial pressures by allowing moulins to behave as a via time-varying storage mechanism, which in

the moulin (Fig. 10a). This has implications for subglacial channel growth and size (Fig. 10c). Nevertheless, this model MouSh currently lacks a distributed system, which limits its fidelity for assimilating daily meltwater volumes into the subglacial system. Realistically, the channelized subglacial system cannot always accommodate the full volume of meltwater produced during summer days, and a portion of this water goes into the distributed system (e.g., Mair et al., 2001, 2002) (e.g., Mair et al., 2001, 2002). In our model, however, when the channelized system is overwhelmed, the water level in the moulin rises above what is typically observed, and sometimes even exceeds the height of the ice (Figs. 6b, S2b, S3b). The melt-driven opening and creep closure processes in the subglacial model explain this behavior: A lower water input to the moulin (Q_m) lowers the water flux into the subglacial system (Q_{out}), which lowers the melt rates that keep subglacial channels open, reducing the size of the subglacial channels and thus further reducing the subglacial water flux. This increases the water level in the moulin. Thus, a reduced rate of surface melt can counterintuitively raise the modeled water level (Fig. 6 around day 30), whereas in reality, much of that water would enter the inefficient subglacial hydrologic system when moulin water levels exceed flotation. If the moulin model were coupled to a two-component subglacial model that represents the inefficient system alongside the channelized system, we would anticipate a much-improved ability to assimilate a wide range of meltwater input rates.

5 Conclusions

First results Results from the MouSh model show that moulins are not static cylinders. Their shapes oscillate daily by some 30% around an equilibrium value reached within the first week of the melt season. These daily diurnally oscillating inputs. Daily fluctuations change the water volume held in the englacial hydrologic system, which in turn influences the evolution of the subglacial channels that moulins feed. When we represent a moulin as a static cylinder in our englacial–subglacial hydrology model, these daily fluctuations go absent, and can be substantially over estimated or underestimated, affecting the overall volume hydraulic gradient of water stored in the englacial subglacial system is underestimated by 50–100%. Modeled moulin size and shape may provide a more realistic representation of moulin water level and the englacial void ratio commonly used in subglacial hydrology models, particularly with future efforts to improve the parameterization of moulin development above the water line. This could be achieved by using an englacial hydrology – channelized subglacial system model, such as the MouSh model we present here, to characterize variability in moulin size and shape, or by coupling moulin models to more complete models of the subglacial system (channelized, distributed, and optionally weakly connected) to make a unified englacial–subglacial hydrology model system. Improving the representation of the englacial–subglacial system to explicitly include moulins would have greatest efficacy during periods of rapidly varying supraglacial input (e.g., during the beginning and end of the melt season and during melt events) and in inland areas, with thick ice and high overburden pressures. These are coincident with situations where subglacial models without moulins, or with implicitly static moulins, tend to perform poorly.

Code and Data availability. The Moulin Shape model is publicly available at <https://github.com/kpoinar/moulin-physical-model/releases/tag/v1.0-MouSh-beta>. <https://github.com/kpoinar/moulin-physical-model/tree/MouSh-beta-revisions> (we will make a release when revisions are complete). The model results used in the analysis presented here are ~~available within the above GitHub repository and are~~ archived at the University at Buffalo Libraries at <http://hdl.handle.net/10477/82587>. <http://hdl.handle.net/10477/82587>.

Author contributions. L.C.A. and K.P. jointly conceived of and developed the MouSh model. Both L.C.A. and K.P. designed the study, executed the model runs, analyzed the data, produced the figures, and wrote the manuscript. C.T. implemented the subglacial module, participated in discussions, and edited the manuscript.

Acknowledgements. This work was supported by NASA Cryosphere grant 80NSSC19K0054 (L.C.A. and K.P.), the Global Modeling and Assimilation Office at NASA Goddard Space Flight Center funded under the NASA Modeling, Analysis, and Prediction (MAP) program (L.C.A.), the Research and Education in eNergy, Environment and Water (RENEW) Institute at the University at Buffalo (K.P.), and the United States National Science Foundation award number NSF-ANS 1603835 (C.T.). We acknowledge DigitalGlobe, Inc. for providing WorldView images via the Enhanced View Web Hosting Services and the support therein provided by the Polar Geospatial Center under NSF-OPP awards 1043681 and 1559691. We thank two anonymous reviewers and editor Dr. Elizabeth Bagshaw for constructive feedback which substantially improved manuscript clarity and completeness.

Competing interests. An author is a member of the editorial board of The Cryosphere. The peer-review process was guided by an independent editor, and the authors declare that they have no conflicts of interest other competing interests to declare.

References

- ~~Alley, R. B.: Flow law hypotheses for ice sheet modeling, Journal of Glaciology, 38(129), 245–256, <https://doi.org/10.3189/S0022143000003658>, 1992.~~
- ~~Alley, R. B., Clark, P. U., Huybrechts, P. and Joughin, I.: Ice Sheet and Sea-Level Changes, Science, 310(5747), 456–460, <https://doi.org/10.1126/science.1114613>, 2005.~~
- ~~Amadei, B.: Rock Anisotropy and the Theory of Stress Measurements, Springer-Verlag, Berlin, New York., 1983.~~
- ~~Andrews, L. C.: Spatial and Temporal Evolution of the Glacial Hydrologic System of the Western Greenland Ice Sheet: Observational and Remote Sensing Results, Ph.D., The University of Texas at Austin, Austin, Texas, 2015.~~
- ~~Andrews, L. C., Catania, G. A., Hoffman, M. J., Gulley, J. D., Lüthi, M. P., Ryser, C., Hawley, R. L. and Neumann, T. A.: Direct observations of evolving subglacial drainage beneath the Greenland Ice Sheet, Nature, 514(7520), 80–83, <https://doi.org/10.1038/nature13796>, 2014.~~

Andrews, L. C., Hoffman, M. J., Neumann, T. A., Catania, G. A., Lüthi, M. P., Hawley, R. L., Schild, K. M., Ryser, C. and Morriss, B. F.: Seasonal Evolution of the Subglacial Hydrologic System Modified by Supraglacial Lake Drainage in Western Greenland, *Journal of Geophysical Research: Earth Surface*, 123(6), 1479–1496, <https://doi.org/10.1029/2017JF004585>, 2018.

Banwell, A. F., Willis, I. C. and Arnold, N. S.: Modeling subglacial water routing at Paakitsoq, W Greenland, *J. Geophys. Res. Earth Surf.*, 118(3), 1282–1295, <https://doi.org/10.1002/jgrf.20093>, 2013.

Banwell, A. F., Hewitt, I., Willis, I. and Arnold, N.: Moulin density controls drainage development beneath the Greenland ice sheet, *J. Geophys. Res. Earth Surf.*, 2015JF003801, <https://doi.org/10.1002/2015JF003801>, 2016.

Bartholomew, T. C., Anderson, R. S. and Anderson, S. P.: Growth and collapse of the distributed subglacial hydrologic system of Kennicott Glacier, Alaska, USA, and its effects on basal motion, *Journal of Glaciology*, 57(206), 985–1002, <https://doi.org/10.3189/002214311798843269>, 2011.

Bartholomew, I. D., Nienow, P., Sole, A., Mair, D., Cowton, T. and King, M. A.: Short-term variability in Greenland Ice Sheet motion forced by time varying meltwater drainage: Implications for the relationship between subglacial drainage system behavior and ice velocity, *J. Geophys. Res.*, 117(F3), F03002, <https://doi.org/10.1029/2011JF002220>, 2012.

Bathurst, J. C.: Flow Resistance Estimation in Mountain Rivers, *Journal of Hydraulic Engineering*, 111(4), 625–643, [https://doi.org/10.1061/\(ASCE\)0733-9429\(1985\)111:4\(625\)](https://doi.org/10.1061/(ASCE)0733-9429(1985)111:4(625)), 1985.

Bell, R. E.: The role of subglacial water in ice sheet mass balance, *Nature Geosci.*, 1(5), 297–304, <https://doi.org/10.1038/ngeo186>, 2008.

Benn, D. I., Thompson, S., Gulley, J., Mertes, J., Luckman, A. and Nicholson, L.: Structure and evolution of the drainage system of a Himalayan debris-covered glacier, and its relationship with patterns of mass loss, *The Cryosphere*, 11, 2247–2264, <https://doi.org/10.5194/te-11-2247-2017>, 2017.

Boulton, G. S., Lunn, R., Vidstrand, P. and Zatsepin, S.: Subglacial drainage by groundwater channel-coupling, and the origin of esker systems: part II—theory and simulation of a modern system, *Quaternary Science Reviews*, 26(7–8), 1091–1105, <https://doi.org/10.1016/j.quascirev.2007.01.006>, 2007.

Catania, G. A. and Neumann, T. A.: Persistent englacial drainage features in the Greenland Ice Sheet, *Geophys. Res. Lett.*, 37(2), L02501, <https://doi.org/10.1029/2009GL041108>, 2010.

Catania, G. A., Neumann, T. A. and Price, S. F.: Characterizing englacial drainage in the ablation zone of the Greenland Ice Sheet, *Journal of Glaciology*, 54(187), 567–578, <https://doi.org/10.3189/002214308786570854>, 2008.

Chandler, D. M., Wadham, J. L., Lis, G. P., Cowton, T., Sole, A., Bartholomew, I., Telling, J., Nienow, P., Bagshaw, E. B., Mair, D., Vinen, S. and Hubbard, A.: Evolution of the subglacial drainage system beneath the Greenland Ice Sheet revealed by tracers, *Nature Geosci.*, 6(3), 195–198, <https://doi.org/10.1038/ngeo1737>, 2013.

Christoffersen, P., Bougamont, M., Hubbard, A., Doyle, S. H., Grigsby, S. and Pettersson, R.: Cascading lake drainage on the Greenland Ice Sheet triggered by tensile shock and fracture, *Nature Communications*, 9(1), 1064, <https://doi.org/10.1038/s41467-018-03420-8>, 2018.

- Chu, V. W.: Greenland Ice Sheet hydrology: A review, *Progress in Physical Geography*, 38(1), 19–54, <https://doi.org/10.1177/0309133313507075>, 2014.
- Chu, W., Creyts, T. T. and Bell, R. E.: Rerouting of Subglacial Water Flow between Neighboring Glaciers in West Greenland, *J. Geophys. Res. Earth Surf.*, 2015JF003705, <https://doi.org/10.1002/2015JF003705>, 2016.
- Chudley, T. R., Christoffersen, P., Doyle, S. H., Bougamont, M., Schoonman, C. M., Hubbard, B. and James, M. R.: Supraglacial lake drainage at a fast flowing Greenlandic outlet glacier, *PNAS*, 116(51), 25468–25477, <https://doi.org/10.1073/pnas.1913685116>, 2019.
- Clarke, G. K. C.: Lumped-element analysis of subglacial hydraulic circuits, *J. Geophys. Res.*, 101(B8), 17547–17559, <https://doi.org/10.1029/96JB01508>, 1996.
- Clarke, G. K. C.: Hydraulics of subglacial outburst floods: new insights from the Spring Hutter formulation, *Journal of Glaciology*, 49(165), 299–313, <https://doi.org/10.3189/172756503781830728>, 2003.
- Colgan, W. and Steffen, K.: Modelling the spatial distribution of moulins near Jakobshavn, Greenland, *IOP Conf. Ser.: Earth Environ. Sci.*, 6(1), 012022, <https://doi.org/10.1088/1755-1307/6/1/012022>, 2009.
- Colgan, W., Steffen, K., McLamb, W. S., Abdalati, W., Rajaram, H., Motyka, R., Phillips, T. and Anderson, R.: An increase in crevasse extent, West Greenland: Hydrologic implications, *Geophys. Res. Lett.*, 38(18), L18502, <https://doi.org/10.1029/2011GL048491>, 2011a.
- Colgan, W., Rajaram, H., Anderson, R., Steffen, K., Phillips, T., Joughin, I., Zwally, H. J. and Abdalati, W.: The annual glaciology cycle in the ablation zone of the Greenland ice sheet: Part 1. Hydrology model, *Journal of Glaciology*, 57(204), 697–709, <https://doi.org/10.3189/002214311797409668>, 2011b.
- Covington, M. D., Banwell, A. F., Gulley, J., Saar, M. O., Willis, I. and Wicks, C. M.: Quantifying the effects of glacier conduit geometry and recharge on proglacial hydrograph form, *Journal of Hydrology*, 414–415, 59–71, <https://doi.org/10.1016/j.jhydrol.2011.10.027>, 2012.
- Covington, M. D., Gulley, J. D., Trunz, C., Mejia, J. and Gadd, W.: Moulin Volumes Regulate Subglacial Water Pressure on the Greenland Ice Sheet, *Geophysical Research Letters*, 47(20), e2020GL088901, <https://doi.org/10.1029/2020GL088901>, 2020.
- Cowton, T., Nienow, P., Sole, A., Wadham, J., Lis, G., Bartholomew, I., Mair, D. and Chandler, D.: Evolution of drainage system morphology at a land terminating Greenlandic outlet glacier, *J. Geophys. Res. Earth Surf.*, 118(1), 29–41, <https://doi.org/10.1029/2012JF002540>, 2013.
- Cowton, T., Nienow, P., Bartholomew, I. and Mair, D.: Variability in ice motion at a land-terminating Greenlandic outlet glacier: the role of channelized and distributed drainage systems, *Journal of Glaciology*, FirstView, 1–16, <https://doi.org/10.1017/jog.2016.36>, 2016.
- Crozier, J., Karlstrom, L. and Yang, K.: Basal control of supraglacial meltwater catchments on the Greenland Ice Sheet, *The Cryosphere*, 12(10), 3383–3407, <https://doi.org/10.5194/te-12-3383-2018>, 2018.
- Cuffey, K. M. and Paterson, W. S. B.: *References: The Physics of Glaciers*, Elsevier., 2010.

Das, S. B., Joughin, I., Behn, M. D., Howat, I. M., King, M. A., Lizarralde, D. and Bhatia, M. P.: Fracture Propagation to the base of the Greenland Ice Sheet during supraglacial lake drainage, *Science*, 320(5877), 778–781, <https://doi.org/10.1126/science.1153360>, 2008.

Dow, C. F., Kulessa, B., Rutt, I. C., Doyle, S. H. and Hubbard, A.: Upper bounds on subglacial channel development for interior regions of the Greenland ice sheet, *Journal of Glaciology*, 60(224), 1044–1052, <https://doi.org/10.3189/2014JG14J093>, 2014.

Dow, C. F., Kulessa, B., Rutt, I. C., Tsai, V. C., Pimentel, S., Doyle, S. H., van As, D., Lindbäck, K., Pettersson, R., Jones, G. A. and Hubbard, A.: Modeling of subglacial hydrological development following rapid supraglacial lake drainage, *J. Geophys. Res. Earth Surf.*, 120(6), 2014JF003333, <https://doi.org/10.1002/2014JF003333>, 2015.

Everett, A., Murray, T., Selmes, N., Rutt, I. C., Luckman, A., James, T. D., Clason, C., O’Leary, M., Karunarathna, H., Moloney, V. and Reeve, D. E.: Annual down-glacier drainage of lakes and water-filled crevasses at Helheim Glacier, southeast Greenland, *J. Geophys. Res. Earth Surf.*, 121(10), 2016JF003831, <https://doi.org/10.1002/2016JF003831>, 2016.

Flowers, G. E.: Subglacial modulation of the hydrograph from glacierized basins, *Hydrol. Process.*, 22(19), 3903–3918, <https://doi.org/10.1002/hyp.7095>, 2008.

Flowers, G. E.: Hydrology and the future of the Greenland Ice Sheet, *Nature Communications*, 9(1), 2729, <https://doi.org/10.1038/s41467-018-05002-0>, 2018.

Flowers, G. E. and Clarke, G. K. C.: A multicomponent coupled model of glacier hydrology 1. Theory and synthetic examples, *J. Geophys. Res.*, 107(B11), 2287, <https://doi.org/10.1029/2001JB001122>, 2002.

Forster, R. R., Box, J. E., van den Broeke, M. R., Miège, C., Burgess, E. W., van Angelen, J. H., Lenaerts, J. T. M., Koenig, L. S., Paden, J., Lewis, C., Gogineni, S. P., Leuschen, C. and McConnell, J. R.: Extensive liquid meltwater storage in firn within the Greenland ice sheet, *Nature Geosci.*, 7(2), 95–98, <https://doi.org/10.1038/ngeo2043>, 2014.

Fountain, A. G. and Walder, J. S.: Water flow through temperate glaciers, *Rev. Geophys.*, 36(3), 299–328, <https://doi.org/10.1029/97RG03579>, 1998.

Fried, M. J., Catania, G. A., Bartholomäus, T. C., Dunean, D., Davis, M., Stearns, L. A., Nash, J., Shroyer, E. and Sutherland, D.: Distributed subglacial discharge drives significant submarine melt at a Greenland tidewater glacier, *Geophys. Res. Lett.*, 42(21), 2015GL065806, <https://doi.org/10.1002/2015GL065806>, 2015.

Germain, S. L. S. and Moorman, B. J.: Long term observations of supraglacial streams on an Arctic glacier, *Journal of Glaciology*, 65(254), 900–911, <https://doi.org/10.1017/jog.2019.60>, 2019.

Goodman, R. E.: *Introduction to Rock Mechanics*, 2nd ed., Wiley, New York., 1989.

Gudmundsson, G. H.: Transmission of basal variability to a glacier surface, *Journal of Geophysical Research*, 108(B5), <https://doi.org/10.1029/2002JB002107>, 2003.

Gulley, J. D., Benn, D. I., Sreaton, E. and Martin, J.: Mechanisms of englacial conduit formation and their implications for subglacial recharge, *Quaternary Science Reviews*, 28(19–20), 1984–1999, <https://doi.org/10.1016/j.quascirev.2009.04.002>, 2009.

Gulley, J. D., Spellman, P. D., Covington, M. D., Martin, J. B., Benn, D. I. and Catania, G.: Large values of hydraulic roughness in subglacial conduits during conduit enlargement: implications for modeling conduit evolution, *Earth Surf. Process. Landforms*, 39(3), 296–310, <https://doi.org/10.1002/esp.3447>, 2014.

Hewitt, I. J.: Seasonal changes in ice sheet motion due to melt water lubrication, *Earth and Planetary Science Letters*, 371–372, 16–25, <https://doi.org/10.1016/j.epsl.2013.04.022>, 2013.

Hoffman, M. J. and Price, S.: Feedbacks between coupled subglacial hydrology and glacier dynamics, *J. Geophys. Res. Earth Surf.*, 119(3), 414–436, <https://doi.org/10.1002/2013JF002943>, 2014.

Hoffman, M. J., Catania, G. A., Neumann, T. A., Andrews, L. C. and Rumrill, J. A.: Links between acceleration, melting, and supraglacial lake drainage of the western Greenland Ice Sheet, *J. Geophys. Res.*, 116(F4), F04035, <https://doi.org/10.1029/2010JF001934>, 2011.

Hoffman, M. J., Andrews, L. C., Price, S. A., Catania, G. A., Neumann, T. A., Lüthi, M. P., Gulley, J., Ryser, C., Hawley, R. L. and Morriss, B.: Greenland subglacial drainage evolution regulated by weakly connected regions of the bed, *Nature Communications*, 7, 13903, <https://doi.org/10.1038/ncomms13903>, 2016.

Hoffman, M. J., Perego, M., Andrews, L. C., Price, S. F., Neumann, T. A., Johnson, J. V., Catania, G. and Lüthi, M. P.: Widespread Moulin Formation During Supraglacial Lake Drainages in Greenland, *Geophysical Research Letters*, <https://doi.org/10.1002/2017GL075659>, 2018.

Holmlund, P.: Internal geometry and evolution of moulins, *Journal of Glaciology*, 34(117), 242–248, 1988.

Howat, I. M., Negrete, A. and Smith, B. E.: The Greenland Ice Mapping Project (GIMP) land classification and surface elevation data sets, *The Cryosphere*, 8(4), 1509–1518, <https://doi.org/10.5194/te-8-1509-2014>, 2014.

Iken, A.: Measurements of water pressure in moulins as part of a movement study of the White Glacier, Axel Heiberg Island, Northwest Territories, Canada, *Journal of Glaciology*, 11(61), 53–58, 1972.

Iken, A. and Bindenschadler, R.: Combined measurements of subglacial water pressure and surface velocity of Findelengletscher, Switzerland: conclusions about drainage system and sliding mechanism, *J. Glac.*, 32(110), 101–119, <https://doi.org/10.3189/S0022143000006936>, 1986.

Iken, A., Echelmeyer, K., Harrison, W. and Funk, M.: Mechanisms of fast flow in Jakobshavns Isbræ, West Greenland: Part I. Measurements of temperature and water level in deep boreholes, *Journal of Glaciology*, 39(131), 15–25, <https://doi.org/10.1017/S0022143000015689>, 1993.

Jarosch, A. H. and Gudmundsson, M. T.: A numerical model for meltwater channel evolution in glaciers, *The Cryosphere*, 6(2), 493–503, <https://doi.org/10.5194/te-6-493-2012>, 2012.

Karlstrom, L. and Yang, K.: Fluvial supraglacial landscape evolution on the Greenland Ice Sheet, *Geophys. Res. Lett.*, 43(6), 2016GL067697, <https://doi.org/10.1002/2016GL067697>, 2016.

Kirsch, G.: *Die Theorie der Elastizität und die Bedürfnisse der Festigkeitslehre*, Springer, Berlin., 1898.

Lüthi, M. P., Ryser, C., Andrews, L. C., Catania, G. A., Funk, M., Hawley, R. L., Hoffman, M. J. and Neumann, T. A.: Heat sources within the Greenland Ice Sheet: dissipation, temperate paleo-firn and cryo-hydrologic warming, *The Cryosphere*, 9(1), 245–253, <https://doi.org/10.5194/te-9-245-2015>, 2015.

MacFerrin, M., Machguth, H., As, D. van, Charalampidis, C., Stevens, C. M., Heilig, A., Vandeerux, B., Langen, P. L., Mottram, R., Fettweis, X., Broeke, M. R. van den, Pfeffer, W. T., Moussavi, M. S. and Abdalati, W.: Rapid expansion of Greenland's low permeability ice slabs, *Nature*, 573(7774), 403–407, <https://doi.org/10.1038/s41586-019-1550-3>, 2019.

MacGregor, J. A., Fahnestock, M. A., Colgan, W. T., Larsen, N. K., Kjeldsen, K. K. and Welker, J. M.: The age of surface-exposed ice along the northern margin of the Greenland Ice Sheet, *Journal of Glaciology*, 66(258), 667–684, <https://doi.org/10.1017/jog.2020.62>, 2020.

Mair, D., Nienow, P., Willis, I. and Sharp, M.: Spatial patterns of glacier motion during a high-velocity event: Haut Glacier d'Arolla, Switzerland, *Journal of Glaciology*, 47(156), 9–20, <https://doi.org/10.3189/172756501781832412>, 2001.

Mair, D., Nienow, P. W., Sharp, M. J., Wohlleben, T. and Willis, I.: Influence of subglacial drainage system evolution on glacier surface motion: Haut Glacier d'Arolla, Switzerland, *Journal of Geophysical Research*, 107(B8), <https://doi.org/10.1029/2001JB000514>, 2002.

Mankoff, K. D., Gulley, J. D., Tulaczyk, S. M., Covington, M. D., Liu, X., Chen, Y., Benn, D. I. and Glowacki, P. S.: Roughness of a subglacial conduit under Hansbreen, Svalbard, *Journal of Glaciology*, 1–13, <https://doi.org/10.1017/jog.2016.134>, 2017.

McGrath, D., Colgan, W., Steffen, K., Lauffenburger, P. and Balog, J.: Assessing the summer water budget of a moulin basin in the Sermeq Avannarleq ablation region, Greenland ice sheet, *Journal of Glaciology*, 57(205), 954–964, <https://doi.org/10.3189/002214311798043735>, 2011.

Meierbachtol, T. W., Harper, J. and Humphrey, N.: Basal Drainage System Response to Increasing Surface Melt on the Greenland Ice Sheet, *Science*, 341(6147), 777–779, <https://doi.org/10.1126/science.1235905>, 2013.

Miège, C., Forster, R. R., Brucker, L., Koenig, L. S., Solomon, D. K., Paden, J. D., Box, J. E., Burgess, E. W., Miller, J. Z., McNerney, L., Brautigam, N., Fausto, R. S. and Gogineni, S.: Spatial extent and temporal variability of Greenland firn aquifers detected by ground and airborne radars, *J. Geophys. Res. Earth Surf.*, 121(12), 2016JF003869, <https://doi.org/10.1002/2016JF003869>, 2016.

Moon, T., Joughin, I., Smith, B., van den Broeke, M. R., van de Berg, W. J., Noël, B. and Usher, M.: Distinct patterns of seasonal Greenland glacier velocity, *Geophys. Res. Lett.*, 41(20), 2014GL061836, <https://doi.org/10.1002/2014GL061836>, 2014.

Moreau, L.: L'exploration du cryokarst glaciaire et son intérêt scientifique pour l'étude du drainage des eaux de fonte : porches, cavités, crevasses, bédrières et moulins., *Collection EDYTEM. Cahiers de géographie*, 8, 163–170, 2009.

Morlighem, M., Williams, C. N., Rignot, E., An, L., Arndt, J. E., Bamber, J. L., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L., Mouginot, J., Noël, B. P. Y., O'Cofaigh, C., Palmer, S., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F.,

van den Broeke, M. R., Weinrebe, W., Wood, M. and Zinglensen, K. B.: BedMachine v3: Complete Bed Topography and Ocean Bathymetry Mapping of Greenland From Multibeam Echo Sounding Combined With Mass Conservation, *Geophysical Research Letters*, 44(21), 11,051–11,061, <https://doi.org/10.1002/2017GL074954>, 2017.

Morlighem, M., Rignot, E., Mouginot, J., Seroussi, H. and Larour, E.: IceBridge BedMachine Greenland, Version 3, NASA National Snow and Ice Data Distributed Active Archive Center, Boulder, Colorado USA. doi.org/10.5067/2CIX82HUV88Y, last access: 16 November 2020, 2018.

Müller, F. and Iken, A.: Velocity fluctuations and water regime of Arctic valley glaciers, *International Association of Scientific Hydrology Publication*, 95, 165–182, 1973.

Naruse, R., Okuhira, F., Ohmae, H., Kawada, K. and Nakawo, M.: Closure Rate of a 700 m Deep Bore Hole at Mizuho Station, East Antarctica, *Annals of Glaciology*, 11, 100–103, <https://doi.org/10.3189/S0260305500006406>, 1988.

Nossokoff, A.: Using Small Scale Physical Experiments to Improve Enthalpy Based Models of Ice Sheets, M.S., University of Colorado, Boulder, 2013.

Palmer, S., Shepherd, A., Nienow, P. and Joughin, I.: Seasonal speedup of the Greenland Ice Sheet linked to routing of surface water, *Earth and Planetary Science Letters*, 302(3–4), 423–428, <https://doi.org/10.1016/j.epsl.2010.12.037>, 2011.

Paterson, W. S. B.: Secondary and tertiary creep of glacier ice as measured by borehole closure rates, *Reviews of Geophysics*, 15(1), 47–55, <https://doi.org/10.1029/RG015i001p00047>, 1977.

Phillips, T., Leyk, S., Rajaram, H., Colgan, W., Abdalati, W., McGrath, D. and Steffen, K.: Modeling moulin distribution on Sermeq Avannarleq glacier using ASTER and WorldView imagery and fuzzy set theory, *Remote Sensing of Environment*, 115(9), 2292–2301, <https://doi.org/10.1016/j.rse.2011.04.029>, 2011.

Pitcher, L. H. and Smith, L. C.: Supraglacial Streams and Rivers, *Annual Review of Earth and Planetary Sciences*, 47(1), 421–452, <https://doi.org/10.1146/annurev-earth-053018-060212>, 2019.

Poinar, K.: The Influence of Meltwater on the Thermal Structure and Flow of the Greenland Ice Sheet, Ph.D., University of Washington, Seattle <https://digital.lib.washington.edu/researchworks/handle/1773/35062>, 2015.

Poinar, K. and Andrews, L. C.: Challenges in predicting Greenland supraglacial lake drainages at the regional scale, *The Cryosphere Discussions*, 1–46, <https://doi.org/10.5194/te-2020-251>, 2020.

Poinar, K., Joughin, I., Das, S. B., Behn, M. D., Lenaerts, J. T. M. and van den Broeke, M. R.: Limits to future expansion of surface melt enhanced ice flow into the interior of western Greenland, *Geophys. Res. Lett.*, 42(6), 2015GL063192, <https://doi.org/10.1002/2015GL063192>, 2015.

Poinar, K., Joughin, I., Lenaerts, J. T. M. and Broeke, M. R. V. D.: Englacial latent heat transfer has limited influence on seaward ice flux in western Greenland, *Journal of Glaciology*, 1–16, <https://doi.org/10.1017/jog.2016.103>, 2016.

Poinar, K., Joughin, I., Lilien, D., Brueker, L., Kehrl, L. and Nowicki, S.: Drainage of Southeast Greenland Firn Aquifer Water through Crevasses to the Bed, *Front. Earth Sci.*, 5, <https://doi.org/10.3389/feart.2017.00005>, 2017.

Priest, S. D.: Discontinuity Analysis for Rock Engineering, 1st ed., Chapman & Hall, London; New York., 1993.

Rippin, D., Willis, I., Arnold, N., Hodson, A., Moore, J., Kohler, J. and Björnsson, H.: Changes in geometry and subglacial drainage of Midre Lovénbreen, Svalbard, determined from digital elevation models, *Earth Surf. Process. Landforms*, 28(3), 273–298, <https://doi.org/10.1002/esp.485>, 2003.

Ryser, C., Lüthi, M. P., Andrews, L. C., Hoffman, M. J., Catania, G. A., Hawley, R. L., Neumann, T. A. and Kristensen, S. S.: Sustained high basal motion of the Greenland ice sheet revealed by borehole deformation, *Biocontrol Science and Technology*, 60(222), 647–660, <https://doi.org/10.3189/2014JcG13J196>, 2014.

Scheingross, J. S. and Lamb, M. P.: A Mechanistic Model of Waterfall Plunge Pool Erosion into Bedrock, *Journal of Geophysical Research: Earth Surface*, 122(11), 2079–2104, <https://doi.org/10.1002/2017JF004195>, 2017.

Schoof, C.: Ice-sheet acceleration driven by melt supply variability, *Nature*, 468(7325), 803–806, <https://doi.org/10.1038/nature09618>, 2010.

Selmes, N., Murray, T. and James, T. D.: Fast draining lakes on the Greenland Ice Sheet, *Geophys. Res. Lett.*, 38(15), L15501, <https://doi.org/10.1029/2011GL047872>, 2011.

Shreve, R. L.: Movement of water in glaciers, *Journal of Glaciology*, 11(62), 205–214, <https://doi.org/doi:10.1017/S002214300002219X>, 1972.

Smith, L. C., Chu, V. W., Yang, K., Gleason, C. J., Pitcher, L. H., Rennermalm, A. K., Legleiter, C. J., Behar, A. E., Overstreet, B. T., Moustafa, S. E., Tedesco, M., Forster, R. R., LeWinter, A. L., Finnegan, D. C., Sheng, Y. and Balog, J.: Efficient meltwater drainage through supraglacial streams and rivers on the southwest Greenland ice sheet, *PNAS*, 112(4), 1001–1006, <https://doi.org/10.1073/pnas.1413024112>, 2015.

Smith, L. C., Yang, K., Pitcher, L. H., Overstreet, B. T., Chu, V. W., Rennermalm, A. K., Ryan, J. C., Cooper, M. G., Gleason, C. J., Tedesco, M., Jeyaratnam, J., As, D. van, Broeke, M. R. van den, Berg, W. J. van de, Noël, B., Langen, P. L., Cullather, R. I., Zhao, B., Willis, M. J., Hubbard, A., Box, J. E., Jenner, B. A. and Behar, A. E.: Direct measurements of meltwater runoff on the Greenland ice sheet surface, *PNAS*, 114(50), E10622–E10631, <https://doi.org/10.1073/pnas.1707743114>, 2017.

Sommers, A. N. and Rajaram, H.: Energy Transfer by Turbulent Dissipation in Glacial Conduits, *Journal of Geophysical Research: Earth Surface*, n/a(n/a), e2019JF005502, <https://doi.org/10.1029/2019JF005502>, 2020.


Spring, U. and Hutter, K.: Numerical studies of Jökulhlaups, *Cold Regions Science and Technology*, 4(3), 227–244, [https://doi.org/10.1016/0165-232X\(81\)90006-9](https://doi.org/10.1016/0165-232X(81)90006-9), 1981.

Spring, U. and Hutter, K.: Conduit flow of a fluid through its solid phase and its application to intraglacial channel flow, *International Journal of Engineering Science*, 20(2), 327–363, [https://doi.org/10.1016/0020-7225\(82\)90029-5](https://doi.org/10.1016/0020-7225(82)90029-5), 1982.

Stevens, L. A., Behn, M. D., McGuire, J. J., Das, S. B., Joughin, I., Herring, T., Shean, D. E. and King, M. A.: Greenland supraglacial lake drainages triggered by hydrologically induced basal slip, *Nature*, 522(7554), 73–76, <https://doi.org/10.1038/nature14480>, 2015.

Tedstone, A. J., Nienow, P. W., Gourmelen, N., Dehecq, A., Goldberg, D. and Hanna, E.: Decadal slowdown of a land-terminating sector of the Greenland Ice Sheet despite warming, *Nature*, 526(7575), 692–695, <https://doi.org/10.1038/nature15722>, 2015.

- Trunz, C., Covington, M. D., Poinar, K., Andrews, L. C., Mejia, J. Z. and Gulley, J. D.: Modeling the impact of moulin shape on englacial hydrology, *JGR: Earth Surface*, in review.
- Trunz, C., Poinar, K., Andrews, L. C., Covington, M. D., Mejia, J. Z., Gulley, J. D., Siegel, V. L. and Breithaupt, C.: Combined Modeled and Explored Moulin Shape Informs Subglacial Pressure Dynamics in Western Greenland, <https://agu.confex.com/agu/fm20/webprogram/Paper675174.html>, 2020.
- Tureotte, D. L. and Schubert, G.: *Geodynamics*, Cambridge University Press., 2002.
- Vaughan, D. G.: Tidal flexure at ice shelf margins, *Journal of Geophysical Research: Solid Earth*, 100(B4), 6213–6224, <https://doi.org/10.1029/94JB02467>, 1995.
- Werder, M. A., Schuler, T. V. and Funk, M.: Short term variations of tracer transit speed on alpine glaciers, *The Cryosphere*, 4(3), 381–396, 2010.
- Werder, M. A., Hewitt, I. J., Schoof, C. G. and Flowers, G. E.: Modeling channelized and distributed subglacial drainage in two dimensions, *Journal of Geophysical Research: Earth Surface*, 118(4), 2140–2158, <https://doi.org/10.1002/jgrf.20146>, 2013.
- Williams, J. J., Gourmelen, N. and Nienow, P.: Dynamic response of the Greenland ice sheet to recent cooling, *Scientific Reports*, 10(1), 1647, <https://doi.org/10.1038/s41598-020-58355-2>, 2020.
- Wright, P. J., Harper, J. T., Humphrey, N. F. and Meierbachtol, T. W.: Measured basal water pressure variability of the western Greenland Ice Sheet: Implications for hydraulic potential, *J. Geophys. Res. Earth Surf.*, 121(6), 2016JF003819, <https://doi.org/10.1002/2016JF003819>, 2016.
- Yang, K. and Smith, L. C.: Internally drained catchments dominate supraglacial hydrology of the southwest Greenland Ice Sheet, *J. Geophys. Res. Earth Surf.*, 2016JF003927, <https://doi.org/10.1002/2016JF003927>, 2016.
- Yang, K., Smith, L. C., Chu, V. W., Pitcher, L. H., Gleason, C. J., Rennermalm, A. K. and Li, M.: Fluvial morphometry of supraglacial river networks on the southwest Greenland Ice Sheet, *GIScience & Remote Sensing*, 53(4), <https://doi.org/10.1080/15481603.2016.1162345>, 2016.
- Zwally, H. J., Abdalati, W., Herring, T., Larson, K., Saba, J. and Steffen, K.: Surface Melt-Induced Acceleration of Greenland Ice-Sheet Flow, *Science*, 297(5579), 218–222, <https://doi.org/10.1126/science.1072708>, 2002.



[Aadnøy, B. S.: A complete elastic model for fluid-induced and in-situ generated stresses with the presence of a borehole, Energy Sources, 9, 239–259, 1987.](#)

[Alley, R. B.: Flow-law hypotheses for ice-sheet modeling, J. Glac., 38\(129\), 245–256, https://doi.org/10.3189/S0022143000003658, 1992.](#)

[Alley, R. B., Dupont, T. K., Parizek, B. R., and Anandakrishnan, S.: Access of surface meltwater to beds of sub-freezing glaciers: preliminary insights, Ann. Glac., 40, 8–14, https://doi.org/10.3189/172756405781813483, 2005.](#)

[Amadei, B.: Rock Anisotropy and the Theory of Stress Measurements, Springer-Verlag, Berlin, New York., 1983.](#)

[Andrews, L. C., Catania, G. A., Hoffman, M. J., Gulley, J. D., Lüthi, M. P., Ryser, C., Hawley, R. L. and Neumann, T. A.: Direct observations of evolving subglacial drainage beneath the Greenland Ice Sheet, Nature, 514\(7520\), 80–83, https://doi.org/10.1038/nature13796, 2014.](#)

[Andrews, L. C., Hoffman, M. J., Neumann, T. A., Catania, G. A., Lüthi, M. P., Hawley, R. L., Schild, K. M., Ryser, C. and Morriss, B. F.: Seasonal Evolution of the Subglacial Hydrologic System Modified by Supraglacial Lake Drainage in Western Greenland, J. Geophys. Res. Earth Surf., 123\(6\), 1479–1496, https://doi.org/10.1029/2017JF004585, 2018.](#)

[Banwell, A. F., Willis, I. C. and Arnold, N. S.: Modeling subglacial water routing at Paakitsoq, W Greenland, J. Geophys. Res. Earth Surf., 118\(3\), 1282–1295, https://doi.org/10.1002/jgrf.20093, 2013.](#)

[Banwell, A. F., Hewitt, I., Willis, I. and Arnold, N.: Moulin density controls drainage development beneath the Greenland ice sheet, J. Geophys. Res. Earth Surf., 2015JF003801, https://doi.org/10.1002/2015JF003801, 2016.](#)

- Bartholomaeus, T. C., Anderson, R. S. and Anderson, S. P.: Growth and collapse of the distributed subglacial hydrologic system of Kennicott Glacier, Alaska, USA, and its effects on basal motion, J. Glac., 57(206), 985–1002, <https://doi.org/10.3189/002214311798843269>, 2011.
- Bartholomew, I. D., Nienow, P., Sole, A., Mair, D., Cowton, T. and King, M. A.: Short-term variability in Greenland Ice Sheet motion forced by time-varying meltwater drainage: Implications for the relationship between subglacial drainage system behavior and ice velocity, J. Geophys. Res. Earth Surf., 117(F3), F03002, <https://doi.org/10.1029/2011JF002220>, 2012.
- Bell, R. E.: The role of subglacial water in ice-sheet mass balance, Nature Geosci, 1(5), 297–304, <https://doi.org/10.1038/ngeo186>, 2008.
- Benn, D. I., Thompson, S., Gulley, J., Mertes, J., Luckman, A. and Nicholson, L.: Structure and evolution of the drainage system of a Himalayan debris-covered glacier, and its relationship with patterns of mass loss, TC, 11, 2247–2264, <https://doi.org/10.5194/tc-11-2247-2017>, 2017.
- Boulton, G. S., Lunn, R., Vidstrand, P. and Zatzepin, S.: Subglacial drainage by groundwater–channel coupling, and the origin of esker systems: part II—theory and simulation of a modern system, Quat. Sci. Rev., 26(7–8), 1091–1105, <https://doi.org/10.1016/j.quascirev.2007.01.006>, 2007.
- Catania, G. A. and Neumann, T. A.: Persistent englacial drainage features in the Greenland Ice Sheet, Geophys. Res. Lett., 37(2), L02501, <https://doi.org/10.1029/2009GL041108>, 2010.
- Catania, G. A., Neumann, T. A. and Price, S. F.: Characterizing englacial drainage in the ablation zone of the Greenland Ice Sheet, J. Glac., 54(187), 567–578, <https://doi.org/10.3189/002214308786570854>, 2008.
- Chandler, D. M., Wadham, J. L., Lis, G. P., Cowton, T., Sole, A., Bartholomew, I., Telling, J., Nienow, P., Bagshaw, E. B., Mair, D., Vinen, S. and Hubbard, A.: Evolution of the subglacial drainage system beneath the Greenland Ice Sheet revealed by tracers, Nature Geosci, 6(3), 195–198, <https://doi.org/10.1038/ngeo1737>, 2013.
- Chu, V. W.: Greenland Ice Sheet hydrology: A review, Prog. Phys. Geogr., 38(1), 19–54, <https://doi.org/10.1177/0309133313507075>, 2014.
- Chudley, T. R., Christoffersen, P., Doyle, S. H., Bougamont, M., Schoonman, C. M., Hubbard, B. and James, M. R.: Supraglacial lake drainage at a fast-flowing Greenlandic outlet glacier, PNAS, 116(51), 25468–25477, <https://doi.org/10.1073/pnas.1913685116>, 2019.
- Church, G., Grab, M., Schmelzbach, C., Bauder, A., and Maurer, H.: Monitoring the seasonal changes of an englacial conduit network using repeated ground-penetrating radar measurements, TC, 14, 3269–3286, <https://doi.org/10.5194/tc-14-3269-2020>, 2020.
- Clarke, G. K. C.: Lumped-element analysis of subglacial hydraulic circuits, J. Geophys. Res., 101(B8), 17547–17559, <https://doi.org/10.1029/96JB01508>, 1996.
- Clarke, G. K. C.: Hydraulics of subglacial outburst floods: new insights from the Spring–Hutter formulation, J. Glac., 49(165), 299–313, <https://doi.org/10.3189/172756503781830728>, 2003.
- Colgan, W. and Steffen, K.: Modelling the spatial distribution of moulins near Jakobshavn, Greenland, IOP Conf. Ser.: Earth Environ. Sci., 6(1), 012022, <https://doi.org/10.1088/1755-1307/6/1/012022>, 2009.
- Colgan, W., Rajaram, H., Anderson, R., Steffen, K., Phillips, T., Joughin, I., Zwally, H. J. and Abdalati, W.: The annual glaciohydrology cycle in the ablation zone of the Greenland ice sheet: Part 1. Hydrology model, J. Glac., 57(204), 697–709, <https://doi.org/10.3189/002214311797409668>, 2011.

- Covington, M. D., Banwell, A. F., Gulley, J., Saar, M. O., Willis, I. and Wicks, C. M.: Quantifying the effects of glacier conduit geometry and recharge on proglacial hydrograph form, *J. Hydro.*, 414–415, 59–71, <https://doi.org/10.1016/j.jhydrol.2011.10.027>, 2012.
- Covington, M. D., Gulley, J. D., Trunz, C., Mejia, J. and Gadd, W.: Moulin Volumes Regulate Subglacial Water Pressure on the Greenland Ice Sheet, *Geophys. Res. Lett.*, 47(20), e2020GL088901, <https://doi.org/10.1029/2020GL088901>, 2020.
- Cowton, T., Nienow, P., Sole, A., Wadham, J., Lis, G., Bartholomew, I., Mair, D. and Chandler, D.: Evolution of drainage system morphology at a land-terminating Greenlandic outlet glacier, *J. Geophys. Res. Earth Surf.*, 118(1), 29–41, <https://doi.org/10.1029/2012JF002540>, 2013.
- Cowton, T., Nienow, P., Bartholomew, I. and Mair, D.: Variability in ice motion at a land-terminating Greenlandic outlet glacier: the role of channelized and distributed drainage systems, *J. Glac.*, 62(233), 451–466, <https://doi.org/10.1017/jog.2016.36>, 2016.
- Cuffey, K. M. and Paterson, W. S. B.: *The Physics of Glaciers*, Elsevier., 2010.
- Das, S. B., Joughin, I., Behn, M. D., Howat, I. M., King, M. A., Lizarralde, D. and Bhatia, M. P.: Fracture Propagation to the base of the Greenland Ice Sheet during supraglacial lake drainage, *Science*, 320(5877), 778–781, <https://doi.org/10.1126/science.1153360>, 2008.
- Dow, C. F., Kulesa, B., Rutt, I. C., Tsai, V. C., Pimentel, S., Doyle, S. H., van As, D., Lindbäck, K., Pettersson, R., Jones, G. A. and Hubbard, A.: Modeling of subglacial hydrological development following rapid supraglacial lake drainage, *J. Geophys. Res. Earth Surf.*, 120(6), 2014JF003333, <https://doi.org/10.1002/2014JF003333>, 2015.
- Downs, J. Z., Johnson, J. V., Harper, J. T., Meierbachtol, T., and Werder, M. A.: Dynamic hydraulic conductivity reconciles mismatch between modeled and observed winter subglacial water pressure, *J. Geophys. Res. Earth Surf.*, 123(4), 818–836, <https://doi.org/10.1002/2017JF004522>, 2018.
- Flowers, G. E.: Subglacial modulation of the hydrograph from glacierized basins, *Hydrol. Process.*, 22(19), 3903–3918, <https://doi.org/10.1002/hyp.7095>, 2008.
- Flowers, G. E.: Modelling water flow under glaciers and ice sheets, *Proc. Math. Phys.* 471, 20140907–20140907, <https://doi.org/10.1098/rspa.2014.0907>, 2015.
- Flowers, G. E.: Hydrology and the future of the Greenland Ice Sheet, *Nature Comm.*, 9(1), 2729, <https://doi.org/10.1038/s41467-018-05002-0>, 2018.
- Flowers, G. E. and Clarke, G. K. C.: A multicomponent coupled model of glacier hydrology 1. Theory and synthetic examples, *J. Geophys. Res.*, 107(B11), 2287, <https://doi.org/10.1029/2001JB001122>, 2002.
- Forster, R. R., Box, J. E., van den Broeke, M. R., Miège, C., Burgess, E. W., van Angelen, J. H., Lenaerts, J. T. M., Koenig, L. S., Paden, J., Lewis, C., Gogineni, S. P., Leuschen, C. and McConnell, J. R.: Extensive liquid meltwater storage in firn within the Greenland ice sheet, *Nature Geosci.*, 7(2), 95–98, <https://doi.org/10.1038/ngeo2043>, 2014.
- Fountain, A. G. and Walder, J. S.: Water flow through temperate glaciers, *Rev. Geophys.*, 36(3), 299–328, <https://doi.org/10.1029/97RG03579>, 1998.

- Gajek, W., Gräff, D., Hellmann, S., Rempel, A. W., and Walter, F.: Diurnal expansion and contraction of englacial fracture networks revealed by seismic shear wave splitting, *Comm. Earth. Environ.*, 2, 1–8, <https://doi.org/10.1038/s43247-021-00279-4>, 2021.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *J. Climate*, 30, 5419–5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- Germain, S. L. S. and Moorman, B. J.: Long-term observations of supraglacial streams on an Arctic glacier, *J. Glac.*, 65(254), 900–911, <https://doi.org/10.1017/jog.2019.60>, 2019.
- Goodman, R. E.: Introduction to Rock Mechanics, 2nd ed., Wiley, New York., 1989.
- Gulley, J. D., Benn, D. I., Screatton, E. and Martin, J.: Mechanisms of englacial conduit formation and their implications for subglacial recharge, *Quat. Sci. Rev.*, 28(19–20), 1984–1999, <https://doi.org/10.1016/j.quascirev.2009.04.002>, 2009.
- Gulley, J. D., Spellman, P. D., Covington, M. D., Martin, J. B., Benn, D. I. and Catania, G.: Large values of hydraulic roughness in subglacial conduits during conduit enlargement: implications for modeling conduit evolution, *Earth Surf. Process. Landforms*, 39(3), 296–310, <https://doi.org/10.1002/esp.3447>, 2014.
- Hewitt, I. J.: Seasonal changes in ice sheet motion due to melt water lubrication, *EPSL*, 371–372, 16–25, <https://doi.org/10.1016/j.epsl.2013.04.022>, 2013.
- Hoffman, M. J. and Price, S.: Feedbacks between coupled subglacial hydrology and glacier dynamics, *J. Geophys. Res. Earth Surf.*, 119(3), 414–436, <https://doi.org/10.1002/2013JF002943>, 2014.
- Hoffman, M. J., Catania, G. A., Neumann, T. A., Andrews, L. C. and Rumrill, J. A.: Links between acceleration, melting, and supraglacial lake drainage of the western Greenland Ice Sheet, *J. Geophys. Res. Earth Surf.*, 116(F4), F04035, <https://doi.org/10.1029/2010JF001934>, 2011.
- Hoffman, M. J., Andrews, L. C., Price, S. A., Catania, G. A., Neumann, T. A., Lüthi, M. P., Gulley, J., Ryser, C., Hawley, R. L. and Morriss, B.: Greenland subglacial drainage evolution regulated by weakly connected regions of the bed, *Nature Comm.*, 7, 13903, <https://doi.org/10.1038/ncomms13903>, 2016.
- Holmlund, P.: Internal geometry and evolution of moulins, *J. Glac.*, 34(117), 242–248, 1988.
- Iken, A.: Measurements of water pressure in moulins as part of a movement study of the White Glacier, Axel Heiberg Island, Northwest Territories, Canada, *J. Glac.*, 11(61), 53–58, 1972.
- Iken, A. and Bindshadler, R.: Combined measurements of subglacial water pressure and surface velocity of Findelengletscher, Switzerland: conclusions about drainage system and sliding mechanism, *J. Glac.*, 32(110), 101–119, <https://doi.org/10.3189/S0022143000006936>, 1986.
- Iken, A., Echelmeyer, K., Harrison, W. and Funk, M.: Mechanisms of fast flow in Jakobshavns Isbræ, West Greenland: Part I. Measurements of temperature and water level in deep boreholes, *J. Glac.*, 39(131), 15–25, <https://doi.org/10.1017/S0022143000015689>, 1993.
- Jarosch, A. H. and Gudmundsson, M. T.: A numerical model for meltwater channel evolution in glaciers, *TC*, 6(2), 493–503, <https://doi.org/10.5194/tc-6-493-2012>, 2012.

- Kirsch, G.: Die Theorie der Elastizität und die Bedürfnisse der Festigkeitslehre, Springer, Berlin., 1898.
- Krawczynski, M. J., Behn, M. D., Das, S. B., and Joughin, I.: Constraints on the lake volume required for hydro-fracture through ice sheets, Geophys. Res. Lett., 36, L10501, <https://doi.org/10.1029/2008GL036765>, 2009.
- Lüthi, M. P., Ryser, C., Andrews, L. C., Catania, G. A., Funk, M., Hawley, R. L., Hoffman, M. J. and Neumann, T. A.: Heat sources within the Greenland Ice Sheet: dissipation, temperate paleo-firn and cryo-hydrologic warming, TC, 9(1), 245–253, <https://doi.org/10.5194/tc-9-245-2015>, 2015.
- MacFerrin, M., Machguth, H., As, D. van, Charalampidis, C., Stevens, C. M., Heilig, A., Vandecrux, B., Langen, P. L., Mottram, R., Fettweis, X., Broeke, M. R. van den, Pfeffer, W. T., Moussavi, M. S. and Abdalati, W.: Rapid expansion of Greenland’s low-permeability ice slabs, Nature, 573(7774), 403–407, <https://doi.org/10.1038/s41586-019-1550-3>, 2019.
- MacGregor, J. A., Fahnestock, M. A., Colgan, W. T., Larsen, N. K., Kjeldsen, K. K. and Welker, J. M.: The age of surface-exposed ice along the northern margin of the Greenland Ice Sheet, J. Glac., 66(258), 667–684, <https://doi.org/10.1017/jog.2020.62>, 2020.
- Mair, D., Nienow, P., Willis, I. and Sharp, M.: Spatial patterns of glacier motion during a high-velocity event: Haut Glacier d’Arolla, Switzerland, J. Glac., 47(156), 9–20, <https://doi.org/10.3189/172756501781832412>, 2001.
- Mair, D., Nienow, P. W., Sharp, M. J., Wohlleben, T. and Willis, I.: Influence of subglacial drainage system evolution on glacier surface motion: Haut Glacier d’Arolla, Switzerland, J. of Geophys Res. Solid Earth, 107(B8), <https://doi.org/10.1029/2001JB000514>, 2002.
- Mankoff, K. D., Gulley, J. D., Tulaczyk, S. M., Covington, M. D., Liu, X., Chen, Y., Benn, D. I. and Głowacki, P. S.: Roughness of a subglacial conduit under Hansbreen, Svalbard, J. Glac., 1–13, <https://doi.org/10.1017/jog.2016.134>, 2017.
- McGrath, D., Colgan, W., Steffen, K., Lauffenburger, P. and Balog, J.: Assessing the summer water budget of a moulin basin in the Sermeq Avannarleq ablation region, Greenland ice sheet, J. Glac., 57(205), 954–964, <https://doi.org/10.3189/002214311798043735>, 2011.
- McQuillan, M. and Karlstrom, L.: Fluid resonance in elastic-walled englacial transport networks, J. Glac., 1–14, <https://doi.org/10.1017/jog.2021.48>, 2021.
- Meierbachtol, T. W., Harper, J. and Humphrey, N.: Basal Drainage System Response to Increasing Surface Melt on the Greenland Ice Sheet, Science, 341(6147), 777–779, <https://doi.org/10.1126/science.1235905>, 2013.
- Mejia, J. Z., Gulley, J. D., Trunz, C., Covington, M. D., Bartholomaeus, T. C., Xie, S., and Dixon, T. H.: Isolated Cavities Dominate Greenland Ice Sheet Dynamic Response to Lake Drainage, Geophys. Res. Lett., 48, e2021GL094762, <https://doi.org/10.1029/2021GL094762>, 2021.
- Miège, C., Forster, R. R., Brucker, L., Koenig, L. S., Solomon, D. K., Paden, J. D., Box, J. E., Burgess, E. W., Miller, J. Z., McNerney, L., Brautigam, N., Fausto, R. S. and Gogineni, S.: Spatial extent and temporal variability of Greenland firn aquifers detected by ground and airborne radars, J. Geophys. Res. Earth Surf., 121(12), 2016JF003869, <https://doi.org/10.1002/2016JF003869>, 2016.
- Moon, T., Joughin, I., Smith, B., van den Broeke, M. R., van de Berg, W. J., Noël, B. and Usher, M.: Distinct patterns of seasonal Greenland glacier velocity, Geophys. Res. Lett., 41(20), 2014GL061836, <https://doi.org/10.1002/2014GL061836>, 2014.

- Moreau, L.: L'exploration du cryokarst glaciaire et son intérêt scientifique pour l'étude du drainage des eaux de fonte : porches, cavités, crevasses, bédrières et moulins., Collection EDYTEM. Cahiers de géographie, 8, 163–170, 2009.
- Müller, F. and Iken, A.: Velocity fluctuations and water regime of Arctic valley glaciers, IAHS, 95, 165–182, 1973.
- Naruse, R., Okuhira, F., Ohmae, H., Kawada, K. and Nakawo, M.: Closure Rate of a 700 m Deep Bore Hole at Mizuho Station, East Antarctica, Ann. Glac., 11, 100–103, <https://doi.org/10.3189/S0260305500006406>, 1988.
- Nossokoff, A.: Using Small Scale Physical Experiments to Improve Enthalpy Based Models of Ice Sheets, M.S., University of Colorado, Boulder, 2013.
- Paterson, W. S. B.: Secondary and tertiary creep of glacier ice as measured by borehole closure rates, Rev. Geophys., 15(1), 47–55, <https://doi.org/10.1029/RG015i001p00047>, 1977.
- Pitcher, L. H. and Smith, L. C.: Supraglacial Streams and Rivers, Ann. Rev. Earth Planetary Sci., 47(1), 421–452, <https://doi.org/10.1146/annurev-earth-053018-060212>, 2019.
- Poinar, K. and Andrews, L. C.: Challenges in predicting Greenland supraglacial lake drainages at the regional scale, TC, 15, 1455–1483, <https://doi.org/10.5194/tc-15-1455-2021>, 2021.
- Poinar, K., Joughin, I., Das, S. B., Behn, M. D., Lenaerts, J. T. M. and van den Broeke, M. R.: Limits to future expansion of surface-melt-enhanced ice flow into the interior of western Greenland, Geophys. Res. Lett., 42(6), 2015GL063192, <https://doi.org/10.1002/2015GL063192>, 2015.
- Poinar, K., Joughin, I., Lenaerts, J. T. M. and Broeke, M. R. V. D.: Englacial latent-heat transfer has limited influence on seaward ice flux in western Greenland, J. Glac., 1–16, <https://doi.org/10.1017/jog.2016.103>, 2016.
- Poinar, K., Joughin, I., Lilien, D., Brucker, L., Kehrl, L. and Nowicki, S.: Drainage of Southeast Greenland Firn Aquifer Water through Crevasses to the Bed, Front. Earth Sci., 5, <https://doi.org/10.3389/feart.2017.00005>, 2017.
- Priest, S. D.: Discontinuity Analysis for Rock Engineering, 1st ed., Chapman & Hall, London; New York., 1993.
- Röösli, C., Walter, F., Ampuero, J.-P., and Kissling, E.: Seismic moulin tremor, J. Geophys. Res. Solid Earth 121, 5838–5858, <https://doi.org/10.1002/2015JB012786>, 2016.
- Ryser, C., Lüthi, M. P., Andrews, L. C., Hoffman, M. J., Catania, G. A., Hawley, R. L., Neumann, T. A. and Kristensen, S. S.: Sustained high basal motion of the Greenland ice sheet revealed by borehole deformation, J. Glac., 60(222), 647–660, <https://doi.org/10.3189/2014JG13J196>, 2014.
- Scheingross, J. S. and Lamb, M. P.: A Mechanistic Model of Waterfall Plunge Pool Erosion into Bedrock, J. Geophys. Res. Earth Surf., 122(11), 2079–2104, <https://doi.org/10.1002/2017JF004195>, 2017.
- Schoof, C.: Ice-sheet acceleration driven by melt supply variability, Nature, 468(7325), 803–806, <https://doi.org/10.1038/nature09618>, 2010.
- Selmes, N., Murray, T. and James, T. D.: Fast draining lakes on the Greenland Ice Sheet, Geophys. Res. Lett., 38(15), L15501, <https://doi.org/10.1029/2011GL047872>, 2011.
- Shreve, R. L.: Movement of water in glaciers, J. Glac., 11(62), 205–214, <https://doi.org/10.1017/S002214300002219X>, 1972.
- Smith, L. C., Chu, V. W., Yang, K., Gleason, C. J., Pitcher, L. H., Rennermalm, A. K., Legleiter, C. J., Behar, A. E., Overstreet, B. T., Moustafa, S. E., Tedesco, M., Forster, R. R., LeWinter, A. L., Finnegan, D. C., Sheng, Y. and Balog, J.: Efficient

[meltwater drainage through supraglacial streams and rivers on the southwest Greenland ice sheet, PNAS, 112\(4\), 1001–1006, https://doi.org/10.1073/pnas.1413024112, 2015.](https://doi.org/10.1073/pnas.1413024112)

[Smith, L. C., Yang, K., Pitcher, L. H., Overstreet, B. T., Chu, V. W., Rennermalm, Å. K., Ryan, J. C., Cooper, M. G., Gleason, C. J., Tedesco, M., Jeyaratnam, J., As, D. van, Broeke, M. R. van den, Berg, W. J. van de, Noël, B., Langen, P. L., Cullather, R. I., Zhao, B., Willis, M. J., Hubbard, A., Box, J. E., Jenner, B. A. and Behar, A. E.: Direct measurements of meltwater runoff on the Greenland ice sheet surface, PNAS, 114\(50\), E10622–E10631, https://doi.org/10.1073/pnas.1707743114, 2017.](https://doi.org/10.1073/pnas.1707743114)

[Spring, U. and Hutter, K.: Numerical studies of Jökulhlaups, Cold Reg Sci Technol., 4\(3\), 227–244, https://doi.org/10.1016/0165-232X\(81\)90006-9, 1981.](https://doi.org/10.1016/0165-232X(81)90006-9)

[Spring, U. and Hutter, K.: Conduit flow of a fluid through its solid phase and its application to intraglacial channel flow, Int. J. Eng. Sci., 20\(2\), 327–363, https://doi.org/10.1016/0020-7225\(82\)90029-5, 1982.](https://doi.org/10.1016/0020-7225(82)90029-5)

[Tedstone, A. J., Nienow, P. W., Gourmelen, N., Dehecq, A., Goldberg, D. and Hanna, E.: Decadal slowdown of a land-terminating sector of the Greenland Ice Sheet despite warming, Nature, 526\(7575\), 692–695, https://doi.org/10.1038/nature15722, 2015.](https://doi.org/10.1038/nature15722)

[Trunz, C.: Modeling and Measuring Water Level Fluctuations in the Greenland Ice Sheet: How Moulin Life Cycle and Shape can Inform us on the Subglacial Drainage System, Ph.D., University of Arkansas, Fayetteville, Arkansas, 156 pp., 2021.](#)

[Turcotte, D. L. and Schubert, G.: Geodynamics, Cambridge University Press., 2002.](#)

[Vaňková, I., Voytenko, D., Nicholls, K. W., Xie, S., Parizek, B. R., and Holland, D. M.: Vertical Structure of Diurnal Englacial Hydrology Cycle at Helheim Glacier, East Greenland, Geophys. Res. Lett., 45, 8352–8362, https://doi.org/10.1029/2018GL077869, 2018.](https://doi.org/10.1029/2018GL077869)

[Vaughan, D. G.: Tidal flexure at ice shelf margins, J. Geophys. Res. Solid Earth, 100\(B4\), 6213–6224, https://doi.org/10.1029/94JB02467, 1995.](https://doi.org/10.1029/94JB02467)

[Weertman, J.: Theory of water-filled crevasses in glaciers applied to vertical magma transport beneath oceanic ridges, J. Geophys. Res., 76, 1171–1183, https://doi.org/10.1029/JB076i005p01171, 1971.](https://doi.org/10.1029/JB076i005p01171)

[Weertman, J.: Can a water-filled crevasse reach the bottom surface of a glacier?, IASH, 95, 139–145, 1973.](#)

[Weertman, J.: Dislocation Based Fracture Mechanics, World Scientific Publishing Company, Singapore, 1996.](#)

[Werder, M. A.: The hydrology of subglacial overdeepenings: A new supercooling threshold formula, Geophys. Res. Lett., 43, 2015GL067542, https://doi.org/10.1002/2015GL067542, 2016.](https://doi.org/10.1002/2015GL067542)

[Werder, M. A., Schuler, T. V. and Funk, M.: Short term variations of tracer transit speed on alpine glaciers, TC, 4\(3\), 381–396, 2010.](#)

[Werder, M. A., Hewitt, I. J., Schoof, C. G. and Flowers, G. E.: Modeling channelized and distributed subglacial drainage in two dimensions, J. Geophys. Res. Earth Surf., 118\(4\), 2140–2158, https://doi.org/10.1002/jgrf.20146, 2013.](https://doi.org/10.1002/jgrf.20146)

[Williams, J. J., Gourmelen, N. and Nienow, P.: Dynamic response of the Greenland ice sheet to recent cooling, Sci. Rep., 10\(1\), 1647, https://doi.org/10.1038/s41598-020-58355-2, 2020.](https://doi.org/10.1038/s41598-020-58355-2)

[Wright, P. J., Harper, J. T., Humphrey, N. F. and Meierbachtol, T. W.: Measured basal water pressure variability of the western Greenland Ice Sheet: Implications for hydraulic potential, J. Geophys. Res. Earth Surf., 121\(6\), 2016JF003819, https://doi.org/10.1002/2016JF003819, 2016.](https://doi.org/10.1002/2016JF003819)

Yang, K. and Smith, L. C.: Internally drained catchments dominate supraglacial hydrology of the southwest Greenland Ice Sheet, J. Geophys. Res. Earth Surf., 2016JF003927, <https://doi.org/10.1002/2016JF003927>, 2016.

Yang, K., Smith, L. C., Chu, V. W., Pitcher, L. H., Gleason, C. J., Rennermalm, A. K. and Li, M.: Fluvial morphometry of supraglacial river networks on the southwest Greenland Ice Sheet, GISci. Remote Sens., 53(4), <https://doi.org/10.1080/15481603.2016.1162345>, 2016.

Figure 1




Figure 1. Processes included in the MouSh model. Black lines show a base moulin geometry that each process acts on, and colored lines show the change in moulin geometry (not to scale) due to that process alone. From left to right: changes in moulin geometry due to viscous deformation; elastic deformation; melting by turbulent energy dissipation of flowing water inside the moulin; melting by open-channel water flow along bare ice; refreezing over winter inside the moulin; and deformation due to ice motion prescribed by Glen's flow law. Unlike the other components, elastic deformation is instantaneous, but applied over the model timestep (Sect. 2.2.1; Supplement S2). The right-most moulin shows the moulin geometry before (dashed black lines) and after (solid black lines and blue water) a hypothetical model timestep, i.e., the sum of all processes shown in the preceding panels. Changes are not to scale.

Figure 2. **Surface**




Figure 2. MouSh geometry and surface expression of a moulin and its reflection in the MouSh model. (a) **Mosaic** Schematic of Sentinel 2 scenes from August 2019 (MacGregor et al., 2020) across the western Greenland Ice Sheet. Pâkitsoq region (green), Russell Glacier area (purple), and location of example MouSh geometry and inputs. Inflow and outflow of the system are indicated by Q_{in} , Q_{out} , and Q_{base} . Time evolving moulin (yellow) are shown, and subglacial parameters include moulin radii (r_1 , r_2), moulin water level (h_w), and subglacial cross-sectional area (S). r_1 and r_2 are evolved by dr_{oc} , dr_v , dr_e , dr_f , and dr_r (open channel melting, viscous deformation, elastic deformation, refreezing, and turbulent melting, respectively; colored as in Fig. 1). u_d shears the moulin as prescribed by Glen's Flow Law. Ice thickness and subglacial path length are indicated by H_i and L , respectively. Ice flow is from left to right. Further details are in Sect. 2. Modified from Trunz (2021). (b) WorldView-2 scene from July 2010 of an approximately 1.2×0.8 km region surrounding the example moulin (yellow) formed by a drained supraglacial lake. (c) Detail of panel b, with the inflow stream and moulin indicated. (d) Detail of panel c, showing the moulin minor radius r_1 , major radius r_2 , and water input Q_{in} from the inflow stream, as represented in the MouSh model. WorldView image © 2010 DigitalGlobe, Inc.




Figure 3. Remotely sensed moulin locations and modeled subglacial flow pathways. Subglacial flow paths predicted by Eq. (1) for (a) Russell glacier region (Smith et al., 2015) and (b) Pâkitsoq (Andrews, 2015; Hoffman et al., 2018) regions and (c) BedMachine (Morlighem et al., 2017, 2018) ice thickness at moulin locations. Background for a–b is the same as in Fig. 2a. Yellow outline in (a) is not included in subglacial pathway length calculations for Russell due to lack of surface imagery and moulin identification. Moulins occupy between 10% (Russell) and 14% (Pâkitsoq) of the englacial-subglacial efficient pathways below 1300 m elevation, suggesting that moulin geometry and evolution may be important to subglacial processes.

Figure 3




Figure 43. Results of parameter sensitivity studies for 10-day MouSh model runs. Shown are the sensitivity of moulin size to initial condition for moulin radius (a–b), enhancement factor for englacial ice (c–d), ice temperature scenario (e–f), Young’s modulus (g–h), softness of basal ice (i–j), friction factor for water flow beneath the water line (k–l), friction factor for water flow above the water line (m–n), and ice thickness (o–p). The left column shows the moulin radii (black and grey) at the equilibriummean water level and the equilibriummean subglacial channel radius (purple) averaged over the final 24-hour period of the ten-day model run. The right column shows the equilibrium water level (blue), moulin volume (red), and volume of water in the moulin (gold) averaged over the same 24-hour period. Overall, moulin radius is most sensitive to the friction factors, while moulin water level and volume are most sensitive to ice thickness H_i and basal ice softness A .

|

|

Figure 54




Figure 4. Diurnal variations in moulin sizes in 10-day parameter sensitivity runs. Shown are the sensitivity of diurnal variation in moulin size and water storage metrics to initial condition for moulin radius (a–b), enhancement factor for englacial ice (c–d), ice temperature scenario from coldest to warmest ice (e–f), Young’s modulus (g–h), softness of basal ice (i–j), friction factor for water flow beneath the water line (k–l), friction factor for water flow above the water line (m–n), and ice thickness (o–p). The left column shows diurnal variations in moulin radii (black and grey) at the equilibrium water level and the subglacial channel radius (purple) in the final 24-hour period of the ten-day model run. The right column shows the diurnal variation in water level (blue), moulin volume (red), and volume of water in the moulin (gold) within the same 24-hour period.

Figure 65




Figure 5. Contributions of viscous deformation, elastic deformation, and phase changes to moulin geometry. (a–e) Equilibrium geometries of five moulins in ice of different ice thicknesses H (same as Fig. 6o–p) averaged over the final 24-hour period of a 10-day model run. (f) Vertical variation of viscous deformation (blue), elastic deformation (red), and phase change (green) contributions to moulin geometry averaged over the same 24-hour period. Negative values indicate contributions to moulin closure; positive values open the moulin. Darkening shades of each color map to moulins of increasing ice thickness. Closure and opening rates are greatest at the minimum daily water level (which is inferable by the lower notch in the moulin wall). (g) Time series of the components shown in panel f (colors the same) at the mean water level over the entire ten-day model run. The greater diurnal range in water level in moulins in thick ice drives the observed larger diurnal variations in viscous and elastic deformation.

Figure 76




Figure 6. MouSh model runs with realistic supraglacial and ice conditions. The model runs are for a low-elevation basin (553 m ice thickness; black lines), mid-elevation basin (741 m ice thickness; purple lines), and high-elevation basin (1315 m ice thickness; grey lines). (a) Supraglacial discharge into the moulin Q_{in} and prescribed base flow Q_{base} . (b) Moulin water level as a fraction of overburden. Note that the highest elevation moulin exceeds the ice surface most days. (c) Moulin capacity, or the total moulin volume. (d) Subglacial channel cross-sectional area. Colored vertical lines indicate times in Fig. 87.

Figure 87




Figure 7. Evolution of moulin geometry over the melt season. Colored boxes correspond to the times indicated in Fig. 76. (a–f) Basin 1 with ice thickness of 553 m. (g–l) Basin 2 with ice thickness of 741 m. (m–r) Basin 3 with ice thickness of 1315 m. Axes are not to scale.

Figure 98




Figure 8. Time series of viscous, elastic and phase change components of moulin evolution and their relative importance in determining moulin geometry. (a) Time varying viscous (blues), elastic (reds), and phase change (melting, greens) components of moulin geometry. (b) The ratio of elastic to viscous deformation (greys) indicates the relative importance of the two deformational processes in moulin evolution. All values are lower than 1, indicating that viscous deformation is always greater. The ratio of the total amount of phase change (melting above and below the water line) to total deformation (elastic plus viscous) (purple). Values above 1 indicate that melting dominates; values below 1 indicate that deformation dominates. Data is smoothed over 24h. For both panels, light colors are for Basin 1 (H_{1c} =553 m), medium colors for Basin 2 (H_{2c} =741 m), and dark colors for Basin 3 (H_{3c} =1315 m).

Figure 109




Figure 9. Daily percentage change in moulin variables relative to the daily mean value.- (a) Daily percentage change in moulin water level relative to the daily mean water level for Basins 1, 2, and 3 (black, purple, and grey lines, respectively).- (b) Daily percentage change in moulin capacity relative to the daily mean moulin capacity. (c) Daily percentage change in the subglacial channel cross-sectional area relative to the daily mean value. For (b-c), colors are as in (a).

Figure 10




Figure 10. Difference between variable and fixed moulin geometries for Basin 1 and 2 (ice thicknesses of 553 m and 741 m, respectively). The fixed moulins are cylinders with a fixed radius of 1.4 m (Basin 1) and 2 m (Basin 2), which are the time-mean radius at the equilibrium water level for the variable moulins. In all instances, the difference is calculated as $(variable - cylindrical - variable)$ with instances of percentage difference calculated as $2(variable - (cylindrical - variable)) / (variable + cylindrical)$. (a, b) Difference in moulin water level for Basin 1 (black) and Basin 2 (purple). Negative values indicate periods where the variable moulin water levels are lower/higher than those of the fixed cylindrical moulin. (c, d) Percentage difference in moulin capacity. When values are negative, the variable moulin is smaller/larger than the fixed cylindrical moulin. (e, f) Percentage difference in subglacial channel cross-sectional area. These values are persistently positive/negative, indicating that the subglacial channel is larger with a variable moulin. (d) Percentage difference in meltwater input. This value is always positive because the addition of water from turbulent melting increases the total amount of water in the system.

Table 1. MouSh model constants and parameter ranges. During realistic runs (Section 2.4) Median values were generally used. In instances where values used differ from the median value, the values used is indicated in parentheses.

Constant		Value	Units	
ρ_i	Ice density	910	kg m ⁻³	
ρ_w	Water density	1000	kg m ⁻³	
g	Gravitational acceleration	9.81	m s ⁻²	
L_f	Latent heat of fusion	335000	J kg ⁻¹	
M_u	Dynamic viscosity (liquid water)	0.0017916	Pa s	
K_w	Thermal conductivity (liquid water)	0.555	J (m K s) ⁻¹	
C_w	Heat capacity (liquid water)	4210	J (K kg) ⁻¹	
C_p	Heat capacity (ice)	2115	J (K kg) ⁻¹	
Parameter		Median value	Range	Units
R_0	Initial moulin radius	2.4 (3)	0.5 to 5	m
E	Ice deformation enhancement factor	5	1 to 9	-
T(z)	Ice temperature	-6 (FOXX profile)	-23 to 0	°C
Y	Young's modulus	5 (9)	1 to 9	GPa
A	Basal ice softness	6×10^{-24}	5×10^{-25} to 5×10^{-23}	Pa ⁻³ s ⁻¹
f_M	Friction factor (under water)	0.1	0.01 to 1	-

foc	Friction factor (subaerial / open channel)	1 (0.8)	0.01 to 1	-
H	Ice thickness	1058 (553, 741, 1315)	669 to 1569	m

Table 2. General ice and moulin input parameters for realistic runs

Parameter	Basin 1	Basin 2	Basin 3
Ice thickness (m)	553	741	1315
Distance from terminus (km)	13.6	24.5	77.1
Catchment size (km ²)	19.8	18.4	55.5
Moulin input, mean diurnal range (m·s ⁻³)	11.5	6.7	2.5
Moulin input, maximum value (m·s ⁻³)	22.1	12.8	6.3
Baseflow, mean value (m·s ⁻³)	20.2	21.2	6.2