

# 1 Controls on Greenland moulin geometry and evolution from the

## 2 Moulin Shape model

3 Lauren C. Andrews<sup>1</sup>, Kristin Poinar<sup>2,3</sup>, Celia Trunz<sup>4,5</sup>

4 <sup>1</sup>Global Modeling and Assimilation Office, NASA Goddard Space Flight Center, Greenbelt, MD, 20771, USA

5 <sup>2</sup>Department of Geology, University at Buffalo, Buffalo, NY, 14260, USA

6 <sup>3</sup>Research and Education in eNergy, Environment and Water (RENEW) Program, University at Buffalo, Buffalo, NY, 14260,  
7 USA

8 <sup>4</sup>Geosciences Department, University of Arkansas, Fayetteville, AR, 72701, USA

9 <sup>5</sup>Department of Applied Geomatics, Université de Sherbrooke, Quebec, J1K 2R1, Canada

10 **Correspondence:** Lauren C. Andrews (lauren.c.andrews@nasa.gov)

11 **Abstract.** Nearly all meltwater from glaciers and ice sheets is routed englacially through moulins. Therefore, the geometry  
12 and evolution of moulins has the potential to influence subglacial water pressure variations, ice motion, and the runoff  
13 hydrograph delivered to the ocean. We develop the *Moulin Shape* (MouSh) model, a time-evolving model of moulin geometry.  
14 MouSh models ice deformation around a moulin using both viscous and elastic rheologies and melting within the moulin  
15 through heat dissipation from turbulent water flow, both above and below the water line. We force MouSh with idealized and  
16 realistic surface melt inputs. Our results show that, under realistic surface melt inputs, variations in surface melt change the  
17 geometry of a moulin by approximately 10% daily and over 100% seasonally. These size variations cause observable  
18 differences in moulin water storage capacity and moulin water levels compared to a static, cylindrical moulin. Our results  
19 suggest that moulins are important storage reservoirs for meltwater, with storage capacity and water levels varying over  
20 multiple timescales. Implementing realistic moulin geometry within subglacial hydrologic models may therefore improve the  
21 representation of subglacial pressures, especially over seasonal periods or in regions where overburden pressures are high.

## 22 1 Introduction

23 Surface-sourced meltwater delivered to the glacier bed drives the evolution of the subglacial hydrologic system and associated  
24 subglacial pressures (e.g., Iken and Bindshadler, 1986; Müller and Iken, 1973). The efficiency of the subglacial system, in  
25 turn, changes the flow patterns of the overlying ice on daily, seasonal, and multi-annual timescales (e.g., Hoffman et al., 2011;  
26 Iken and Bindshadler, 1986; Moon et al., 2014; Tedstone et al., 2015; Williams et al., 2020). Thus, glacial hydrology is a  
27 crucial factor in short-term changes to glacier and ice sheet dynamics (Bell, 2008; Flowers, 2018).

28 On the Greenland Ice Sheet (GIS), surface meltwater can take multiple paths, depending on its origin. In the  
29 accumulation zone, meltwater may percolate through snow and firn, remaining liquid (Forster et al., 2014) or refreezing

30 (MacFerrin et al., 2019). In the ablation zone, meltwater runs over bare ice, coalesces into supraglacial streams, and pools into  
31 supraglacial lakes (e.g., Smith et al., 2015). These surficial water features – rivers, streams, lakes, aquifers, etc. – direct  
32 meltwater into englacial features that can deliver the water to the bed of the ice sheet (Andrews et al., 2014; Das et al., 2008;  
33 Miège et al., 2016; Poinar et al., 2017; Smith et al., 2015). Englacial features include moulins, which are near-vertical shafts  
34 with large surface catchments ( $\sim 1\text{--}5\text{ km}^2$  per moulin, Banwell et al., 2016; Colgan and Steffen, 2009; Yang and Smith, 2016),  
35 and crevasses, which are linear features with limited local catchments ( $\sim 0.05\text{ km}^2$  per crevasse, Poinar et al., 2017). Together,  
36 moulins and crevasses constitute a substantial fraction of the englacial hydrologic system in the ablation zone of the GIS.

37 Water fluxes through the englacial system, and therefore to the subglacial system, are non-uniform in space and time.  
38 Quantifying these temporal variations in water fluxes to the glacier bed requires understanding the time evolution of the  
39 supraglacial and englacial water systems that deliver it. Ongoing research is making great strides in characterizing the  
40 supraglacial water network (Germain and Moorman, 2019; Smith et al., 2017; Yang et al., 2016). For instance, field  
41 observations from Greenland indicate that much of the supraglacial water network terminates into crevasses and moulins  
42 (McGrath et al., 2011; Smith et al., 2015) and that moulins are important modulators of surface melt inputs to the ice sheet bed  
43 (Andrews et al., 2014; Cowton et al., 2013; Mejia et al., 2021; Smith et al., 2021).

44 Our knowledge of moulin sizes, scales, and time evolution has largely been informed by exploration and mapping of  
45 the top tens to hundred meters of a few moulins (Benn et al., 2017; Covington et al., 2020; Gulley et al., 2009; Holmlund,  
46 1988; Moreau, 2009). These sparse field data indicate that moulin shapes deviate greatly from simple cylinders. Furthermore,  
47 deployments of tethered sensors into Greenland moulins have encountered irregularities including apparent ledges and plunge  
48 pools (Andrews et al., 2014; Covington et al., 2020; Cowton et al., 2013), and seismic (Röösli et al., 2016) and radar (Catania  
49 et al., 2008) studies suggest constrictions below the depths of human exploration. These direct near-surface and indirect deep  
50 observations suggest that moulin geometry evolves a high degree of complexity at all depths.

51 State-of-the-art subglacial hydrology models are forced by meltwater inputs that enter the system through crevasses  
52 or moulins. These models generally represent the geometry of moulins in a simplified and time-independent manner, for  
53 instance as a static vertical cylinder (e.g., Hewitt, 2013; Hoffman et al., 2016; Werder et al., 2013) or cone (Clarke, 1996;  
54 Flowers and Clarke, 2002; Werder et al., 2010). The basis for the cylindrical simplification arises from the assumption that  
55 depth-dependent variations in moulin size are small relative to the vertical scale of the moulin. The basis for time independence  
56 is the assumption that the moulin capacity is, again, small relative to that of the subglacial system. However, neither of these  
57 assumptions have been tested. Here, we explore the extent to which an evolving moulin geometry can impact moulin water  
58 level, capacity, and water volume, each of which can impact the evolution of the subglacial system.

59 We present the Moulin Shape (MouSh) model, a new, physically based numeric model that evolves moulin geometry  
60 over diurnal and seasonal periods. The MouSh model can be coupled to subglacial hydrology models to more completely  
61 characterize the time evolution of the englacial and subglacial hydrologic systems, which are intimately linked.

## 62 2 Moulin physical model

63 We develop the Moulin Shape (MouSh) model, a numeric model of moulin evolution that considers ice deformation and ice  
64 melt associated with the dissipation of energy from turbulently flowing meltwater (Fig. 1). We include here a detailed  
65 description of the model framework and each module that influences the time-evolving geometry of the modeled moulin (Fig.  
66 2a).

### 67 2.1 Moulin geometry coordinate system

68 We discretize our model in the vertical ( $z$ ) and radial ( $r_1$  and  $r_2$ , or generally  $r_m$ ) directions, treating the moulin as a stack of  
69 egg-shaped (semi-circular, semi-elliptical) holes in the ice that both change in size and move laterally relative to each other.  
70 We calculate moulin geometry (elliptical radii  $r_1$  and  $r_2$ ) and water level  $h_w$  with a 5-minute timestep  $dt$ . Model calculations  
71 are performed in cylindrical coordinates, where  $\Pi(z)$  is the perimeter of the semi-circular, semi-elliptical moulin, using  
72 Ramanujan's approximation:

$$73 \quad \Pi \approx \pi r_1 + \frac{1}{2}\pi[3(r_1 + r_2) - \sqrt{(3r_1 + r_2)(r_1 + 3r_2)}] \quad (1)$$

74 Here,  $r_1$  and  $r_2$  are the minor and major radii, respectively, for each node in the vertical direction. The minor radius  $r_1$  is also  
75 the radius of the half-circle.

76 We calculate the cross-sectional area  $A_m$  of the semi-circular, semi-elliptical moulin as follows:

$$77 \quad A_m = \frac{\pi r_1}{2}(r_2 + r_1) \quad (2)$$

78 The plan-view orientation of the radii and the coordinate system, as detailed on a remotely sensed moulin, are indicated in  
79 Fig. 2b-d. The elliptical shape was chosen to reflect the observation that supraglacial meltwater flows into a moulin along a  
80 single side above the water line. This asymmetry leads to a nonuniform, noncircular geometry above the water level. This  
81 choice is in line with observations of a GIS moulin becoming more elliptical over time (Röösli et al., 2016). For simplicity,  
82 MouSh also contains an option to set the moulin cross-sectional geometry to a circle, rather than an egg (see Supplement  
83 S2.2.2).

84 Each module is also dependent on the depth varying hydrostatic and cryostatic pressures. We subtract the cryostatic  
85 pressure  $P_i$  from the hydrostatic pressure  $P_w$  to calculate the total depth-dependent pressure  $P$  at all vertical levels  $z$  within  
86 the moulin:

$$87 \quad P_i = \rho_i g(H_i - z) \quad (3a)$$

$$88 \quad P_w = \rho_w g(h_w - z) \quad (3b)$$

$$89 \quad P = P_w - P_i \quad (3c)$$

90 where  $H_i$  is the ice thickness;  $h_w$  is the height of the water above the bed (moulin water level);  $z$  is the vertical  
91 coordinate;  $\rho_i$  and  $\rho_w$  are ice and water density, respectively; and  $g$  is gravitational acceleration (Table 1). Note that  $P$  is not  
92 effective pressure, which is defined as  $P = P_i - P_w$  (Cuffey and Paterson, 2010). In our formulation, positive pressures cause

93 outward expansion of the moulin walls (radial growth), and negative pressures reduces the size of the moulin (radial closure).  
94 We use a flat bed at sea level for all model runs presented here; bed elevation is  $z = b = 0$ .

## 95 **2.2 Ice deformation modules**

96 We represent the deformation of the ice with the simplest possible combination of elastic and viscous components: a Maxwell  
97 rheology, where elastic and viscous deformation occur independently, without interaction (Turcotte and Schubert, 2002). The  
98 Maxwell model comprises an elastic element (a spring) and a viscous element (a dashpot) in series and is standard in  
99 geophysical modeling. The response timescale in our Maxwell model is equal to  $(E \times A \times \tau^2)^{-1}$  where  $E$  is Young's modulus,  
100  $A$  is the viscous flow law parameter, and  $\tau$  is stress (Table 1; Turcotte and Schubert, 2002). The Maxwell timescale is thus  
101 roughly 10–100 hours for typical Greenland ice.

102 Elastic deformation is described in Sect. 2.2.1. We represent total viscous deformation in two modes: (1) radial  
103 opening and closure of the moulin, which changes the size of the moulin (Sect. 2.2.2), and (2) vertical shear of the moulin,  
104 which changes the shape but not the size of the moulin (Sect. 2.2.3).

### 105 **2.2.1 Elastic deformation**

106 Field measurements indicate that, nearly universally during the melt season, the water level in a moulin varies at a sub-hourly  
107 timescale (Andrews et al., 2014; Covington et al., 2020; Cowton et al., 2013; Iken, 1972). This variability is shorter than, but  
108 comparable to, the Maxwell timescale for ice (10–100 hours; see Sect. 2.2); therefore, we must assume that elastic deformation  
109 plays a role in the response of the ice to variations in moulin water level.

110 Weertman (1971, 1973, 1996) applied dislocation fracture mechanics principles to vertical glaciological features:  
111 water-filled crevasses. These equations have applied to supraglacial lake drainages (Krawczynski et al., 2009) and slow ice  
112 hydrofracture (Poinar et al., 2017). However, these problems are Cartesian (linear), not cylindrical, so their solutions are not  
113 readily adaptable to a moulin. The stress and deformational patterns around cylindrical boreholes have been well studied in  
114 the rock mechanics literature (Amadei, 1983; Goodman, 1989; Priest, 1993). We therefore base our description of the stress  
115 field surrounding the moulin on that of a fluid-filled borehole in a porous rock medium, described by Aadnøy (1987) and based  
116 on the Kirsch equations, which describe stresses surrounding a circular hole in a rigid plate (Kirsch, 1898). We assume plane  
117 strain and approximate our moulin as a stack of such plates with analogous holes (Goodman, 1989). A subtle difference is that  
118 our moulin shape is not circular, but egg-shaped: half circular, half elliptical.

119 At each vertical level  $z$  in the moulin, we apply Hooke's Law to the stress field to calculate the strain, in horizontal  
120 cross-section, at all points on the moulin wall and in the surrounding ice for both radii  $r_1$  and  $r_2$ . We then integrate these strains  
121 from an infinite distance (cylindrical coordinate  $r = \infty$ ) to the moulin wall ( $r_1, r_2 = r_m$ ). A full derivation, based on the stress  
122 states in a borehole described by Aadnøy (1987), is in Supplement S1. We express the total radial elastic deformation  $r_e$  of a  
123 moulin segment as:

$$r_e = \frac{r_m}{E} \left[ (1 + \nu) (\Delta P - \frac{1}{2} (\Delta \sigma_x + \Delta \sigma_y)) + \frac{1}{4} (\Delta \sigma_x - \Delta \sigma_y) (1 - 3\nu - 4\nu^2) + \frac{1}{4} \Delta \tau_{xy} (2 - 3\nu - 8\nu^2) \right] \quad (4)$$

Here,  $\Delta P$  is the change in cryo-hydrostatic pressure (Eq. 3c) over a time interval,  $\nu$  is Poisson's ratio;  $r_m$  is used to refer to  $r_1$  or  $r_2$ ; and  $\Delta \sigma_x$ ,  $\Delta \sigma_y$ , and  $\Delta \tau_{xy}$  are changes in background deviatoric and shear stresses that describe the regional setting of the moulin. The model is designed to accept user-defined deviatoric and shear stresses; however, we choose a neutral surface stress state ( $\Delta \sigma_x = \Delta \sigma_y = \Delta \tau_{xy} = 0 \text{ kPa}$ ) for experimental simplicity and because these stresses and their changes over time are poorly constrained. This simplification reduces elastic deformation  $r_e$ :

$$r_e = \frac{r_m}{E} (1 + \nu) \Delta P \quad (5)$$

Unlike viscous deformation and melting, elastic deformation is instantaneous. However, we take advantage of the observation that elastic deformation is driven by changes in the cryostatic and hydrostatic pressures (Supplement S1.5). Therefore, we express Eq. 4 and Eq 5 as an elastic 'deformation rate' for varying (Eq. 6) and constant (Eq. 7) surface stresses:

$$\frac{dr_e}{dt} = \frac{1}{E} \left( r_m (1 + \nu) \frac{dP}{dt} + [(1 + \nu) \left( -\frac{1}{2} \left( \frac{d\sigma_x}{dt} + \frac{d\sigma_y}{dt} \right) + \frac{1}{4} \left( \frac{d\sigma_x}{dt} - \frac{d\sigma_y}{dt} \right) (1 - 3\nu - 4\nu^2) + \frac{1}{4} \frac{d\tau_{xy}}{dt} (2 - 3\nu - 8\nu^2) \right] \right) \quad (6)$$

$$dr_e = \frac{r_m}{E} (1 + \nu) \left( \frac{dP}{dt} \right) dt \quad (7)$$

Equations 6 and 7 assume that both effective pressure and moulin radius vary smoothly over the time interval in question, which is generally true for small timesteps (5-minutes in our model). The dominant term in Eq. 6 is the first term, since  $\frac{dP}{dt}$  (~1 kPa over a typical hour during the melt season) greatly exceeds the rate of change of the surface stresses (~1 kPa over a year), as explained in the Supplement S1. Equation 7 is commonly used for dilatometer testing in rock mechanics (Goodman, 1989).

### 2.2.2 Viscous radial opening and closure

Moulins close when they lose their water source at the end of a melt season (Catania and Neumann, 2010). Similarly, boreholes close if they are not filled with drilling fluid with a density like ice (Alley, 1992). Our modeled moulin is intermediate to these edge cases because it typically contains water. When the moulin is filled with water to the flotation level, it will stay open at its base and viscously close at and below the water level. The moulin will viscously open in regions where hydrostatic pressure exceeds the cryostatic pressure. When the water level is below flotation, which is the typical case, viscous deformation shrinks the moulin at all depths.

We calculate strain rate  $\dot{\epsilon}$  from the total depth-dependent pressure  $P$  (Eq. 3c) using Glen's Flow Law:

$$\dot{\epsilon} = F^* A(T_i, P_i) \cdot \left( \frac{1}{3} P \right)^n \quad (8)$$

152 where  $F^*$  is the flow law enhancement factor;  $A(T_i, P_i)$  is the flow law parameter; and  $n$  is Glen's Flow Law exponent. For the  
 153 flow law parameter, we use the standard relationship from Cuffey and Paterson (2010, Eq. 3.35), which is a function of ice  
 154 temperature  $T_i$  and ice pressure  $P_i$ .

155 We follow borehole studies by Naruse et al (1988) and Paterson (1977) to write strain,  $\varepsilon$ , in the radial direction as

$$156 \quad \varepsilon = \ln\left(\frac{r_f}{r_0}\right) \quad (9)$$

157 where a moulin with initial radius  $r_0$  and final radius  $r_f$  underwent radial strain of  $\varepsilon$ .

158 We use the time derivative of Eq. (9) to calculate the change in moulin radius due to viscous deformation:

$$159 \quad dr_v = r_m \exp(\dot{\varepsilon} dt) - r_m \quad (10)$$

160 with strain rate given by Eq. (8). This is the same relationship used by Catania and Neumann (2010).

### 161 2.2.3 Shear deformation

162 We use Glen's Flow Law to calculate the change in shape of the moulin due to regional-scale ice flow. This deforms the entire  
 163 moulin in bulk, shearing it in the vertical and shifting it laterally downstream, without changing its radii. Basal sliding is not  
 164 currently included in the model. To represent deformation, we discretize the moulin as a stack of plates with elliptical (or  
 165 circular) holes with a thickness  $dz$  and represent deformational ice flow as displacement between these plates.

166 We calculate the rate of deformational ice flow  $u_d$  in the downstream direction from ice temperature  $T_i$  and pressure  
 167  $P_i$ , surface slope  $\alpha$ , a constant  $F^*$ , and  $H_i$ , using Glen's Flow Law (Cuffey and Paterson, 2010):

$$168 \quad u_d = 2F^* (\rho_i g \alpha)^n \cdot \int_b^{H_i} A(T_i, P_i) (H_i - z)^n dz \quad (11)$$

169 where  $b$  is the ice sheet bed. We obtain ice deformation rates of  $\sim 20$  m yr<sup>-1</sup>, which is typical of the ablation zone in western  
 170 Greenland (Ryser et al., 2014).

## 171 2.3 Phase change modules

172 The second mode that changes the geometry of the moulin is ice ablation from or accretion to the moulin walls. During the  
 173 melt season, the flow of water into and through the moulin generates turbulence, which as it dissipates acts to melt back the  
 174 moulin walls, expanding the size of the moulin. There is also a small component of melting due to temperature differences  
 175 between the water and surrounding ice. Outside the melt season, conduction of latent heat into the surrounding ice causes  
 176 stagnant water to freeze back onto the moulin walls, contracting the size of the moulin.

### 177 2.3.1 Refreezing

178 Refreezing occurs in cold ice when water flow is absent or slow enough that the rate of heat conduction into the  
 179 surrounding ice drops the water temperature to the freezing point. These conditions occur primarily outside the melt season.  
 180 When these conditions are met, we apply a radial freezing  $r_f$ , which is parameterized economically, following Alley (2005):

$$dr_f = 2 \frac{T_i - T_{pmp}}{L_f} \sqrt{\frac{K_i C_p}{\pi \rho_i}} (\sqrt{t_t} - \sqrt{t_t - dt}) \quad (12)$$

Here,  $T_i - T_{pmp}$  is the depth-varying difference between the far-field temperature (prescribed as from borehole temperature observations) and the moulin water temperature, which is taken as the pressure melting temperature  $T_{pmp}$ .  $L_f$  is the latent heat of fusion;  $K_i$  is water's thermal conductivity;  $C_p$  is the specific heat capacity of ice. The refreezing rates evolve exclusively based on the elapsed time since the cessation of turbulent flow  $t_t$ .

We calculate the change in moulin water volume from freezing,  $V_{frz}$ , by summing the refrozen ice thickness in a timestep,  $dr_f$ , around the perimeter of the moulin at all depths  $z$ , and converting ice volume to water volume:

$$V_{frz} = \frac{\rho_i}{\rho_w} \int_b^{h_w} \Pi(z) r_f(z) dz \quad (13).$$

### 2.3.2 Moulin wall melting

During the melt season, turbulent energy dissipation from water flowing through the moulin melts back the moulin walls. The dissipation of turbulent energy and the associated melting of the surrounding ice will increase the local moulin radius. We parameterize turbulence in two separate spatial domains: (1) within the water column of the moulin, where  $r_1$  and  $r_2$  are evolved uniformly, and (2) above the water level along the side of the moulin, as supraglacial input falls to the water level, where only  $r_2$  is evolved.

The parameterizations of turbulently driven melting we use in both regimes rely on three simplifications. First, the volume of water moving through each vertical model node is constant within each time step. This ensures that water mass is conserved and that all model elements below the water line are water filled; however, this eliminates the potential long-term storage of meltwater within plunge pools caused by non-uniform incision into the ice. Second, all energy generated from turbulent dissipation is instantaneously applied to melting the surrounding ice. This neglects any heat transport within the water, which is a common approximation in subglacial models (e.g., Hewitt, 2013; Schoof, 2010; Werder et al., 2013). Third, we also make the simplifying assumption that meltwater entering the moulin is at  $0^\circ\text{C}$  and at the pressure melting temperature  $T_{pmp}$  at all points below the water line, although we do not model the impact of this temperature change on melting because moulin water temperatures are unknown.

*Submerged zone:* Below the water line, the vertical velocity of the water is dictated by the hydraulic gradient within the system and the local cross-sectional area of the moulin. Under such conditions, head loss – the departure of the hydraulic head from that calculated by Bernoulli's equation – reflects the energy dissipated as heat. We parameterize head loss using the Darcy–Weisbach equation, which relates water velocity  $u_w$  to changes in the hydraulic gradient  $dh_w/dl$  (head loss per unit length along flow), via the hydraulic radius  $R_h$  and a dimensionless friction factor  $f$ . Because water velocity is constrained by mass balance within the system, we calculate the head loss  $dh_w/dl$  as follows:

$$\frac{dh_w}{dl} = \frac{u_w^2 f}{8R_h g} \quad (14)$$

212 The differential element  $dl$  represents the path length over which the water experiences head loss:  $dl = \sqrt{dx^2 + dz^2}$  for  
 213 horizontal distance  $dx$  and vertical drop  $dz$ . The friction factor  $f$  is a unitless model parameter that controls the rate of head loss  
 214 within the system. Its value thus directly affects the amount of melting. Most subglacial models fix the Darcy–Weisbach  
 215 friction factor, with values ranging from 0.01 to 0.5 (e.g., Colgan et al., 2011b; Schoof, 2010; Spring and Hutter, 1981) or use  
 216 equivalent values of Manning’s  $n$  (e.g., Hewitt, 2013; Hoffman and Price, 2014). Alternatively, other models parameterize  
 217 channel roughness using a geometry-dependent friction factor (e.g., Boulton et al., 2007; Clarke, 2003; Flowers, 2008). Thus,  
 218 MouSh has options for fixed or variable  $f$ .

219 The friction factor within the submerged zone is indicated by  $f_m$  and in the open channel zone by  $f_{oc}$ . To explore the  
 220 impact of the chosen friction factor, we complete a sensitivity study (Sect. 2.5.2 and 3.2). We use a constant  $f_m = 0.1$  for all  
 221 other model runs presented.

222 Because we approximate the moulin as a half-circular, half-elliptical cylinder with perimeter  $\Pi$ , the hydraulic radius  
 223  $R_h$  of a water filled node is:

$$224 \quad R_h = \frac{A_m}{\Pi} \quad (15).$$

225 To calculate moulin wall melting, we use a simple energy balance equation, following previous work (e.g., Gulley et al., 2014;  
 226 Jarosch and Gudmundsson, 2012; Nossokoff, 2013):

$$227 \quad \rho_i C_w (T_{pmp} - T_i) \frac{dA_m}{dt} + \rho_i L_f \frac{dA_m}{dt} = Q_{out} \left( \rho_w g \frac{dh_w}{dt} \right) \quad (16)$$

228 where  $C_w$  is the heat capacity of water. The first term represents the energy needed to warm the surrounding ice to the pressure  
 229 melting temperature of water  $T_{pmp}$ . Equation (16) can be rearranged and combined with equation (14) to provide the area of  
 230 ice melted:

$$231 \quad dA_t = Q_{out} \left( \rho_w g \frac{u_w^2 f}{4R_h g} \right) \left( \rho_i C_w (T_{pmp} - T_i) + \rho_i L_f \right)^{-1} dt \quad (17)$$

232 where  $Q_{out}$  is the discharge from the moulin-subglacial system as dictated by the subglacial model component (Sect. 2.4.2);  
 233 and  $T_{pmp} - T_i$  is the temperature difference between the water (prescribed to be at the pressure melting point) and the  
 234 surrounding ice. We vary  $T_i$  based on observations as described in Table 1 and Sect. 2.5.2. Note that Eq. 17 determines the  
 235 area of ice that is removed through melting. For each time step, we reframe Eq. 17 into radial melting within an egg-shaped  
 236 moulin using information about the previous geometry and the assumption that melting occurs uniformly around the perimeter:

$$237 \quad dr_t = 2dA_t / \left[ \pi(5r_1 + 3r_2 - \sqrt{(3r_1 + r_2)(r_1 + 3r_2)}) \right] \quad (18).$$

238 Equation 18 is simplified when considering a circular geometry ( $r_1 = r_2$ ).

239  
 240 *Unsubmerged zone:* Above the water line, a variety of complex processes drive melting. A first-principles approach would be  
 241 to quantify melting due to the potential energy loss of falling water, following the work on terrestrial waterfalls (e.g.,  
 242 Scheingross and Lamb, 2017). However, nearly all waterfall-based parameterizations rely on abrasion between waterborne



243 sediment and the substrate as the primary mechanism of erosion. Instead, we implement a simple parameterization for open-  
 244 channel flow with the understanding that the complexities of thermal erosion are not completely captured. In our model, open-  
 245 channel melting occurs only on the up-glacier wall of the moulin and follows two ad-hoc rules based on the slope between the  
 246 vertical nodes: (1) open-channel turbulent melting is applied if the slope of the upstream moulin wall allows water to flow  
 247 over it; and (2) a small, prescribed amount of melting is applied when the upstream wall slope is vertical or overhung, because  
 248 while water cannot flow directly along the ice, spray and other processes likely drive some amount of melting. These cases are  
 249 respectively (1) the open-channel zone and (2) the falling water zone.

250 In the open-channel zone, we use a similar approach as for melting below the water line. However, the hydraulic  
 251 radius  $R_h$  is adjusted to reflect the observation that water runs down only one wall of the moulin, and a higher friction factor is  
 252 used to parameterize complex geometries. Due to the presence of a discontinuity between open-channel and water-filled  
 253 regions (at the water line), we parameterize the hydraulic radius of open channel flow as  $R_{h_{open}} = 0.5r_2$ . We also use a higher  
 254 open channel friction factor  $f_{oc}$  of 0.8 to parameterize observed extensive scalloping (e.g., Gully et al., 2014; Covington et al.,  
 255 2020). We apply melting to only the elliptical side of the moulin, defined by  $r_2$  using Eq. 18. Note that the hydraulic radius  
 256 prescribed for open-channel flow is likely larger than the small region over which water is flowing in the natural system (Fig.  
 257 2). Further, the resulting open channel melt  $dA_{oc}$  is applied only to the major radius to calculate the change in open channel  
 258 radius  $dr_{oc}$ .

259 In the falling water zone, there is very limited interaction between the moulin walls and the water. For simplicity, we  
 260 assume that a small fraction,  $f_p$ , of the potential energy lost as water falls is deposited into the moulin walls, perhaps as the  
 261 kinetic energy of spray. The change in radius due to this process is as follows:

$$262 \quad dr_{mf} = f_p \frac{(\rho_w/\rho_i)gQ_{out}}{L_f\Pi} dt \quad (19)$$

263 We set  $f_p$  to 0.1 for all model runs presented here.

264 We add the volume of ice melted to the water already in the moulin, similarly to Eq. 12 for  $V_{jz}$ . We calculate the  
 265 change in moulin water volume from melting by summing the melted ice thickness,  $r_{mf}$ , around the perimeter of the moulin at  
 266 all depths  $z$ , and converting ice volume to water volume:

$$267 \quad V_{wallmelt} = \frac{\rho_i}{\rho_w} dt \int_b^{H_i} (\Pi(z) dr_{mf}(z) + A_{oc}(z) + A_t(z)) dz \quad (20).$$

## 268 2.4 Water flux into and out of the moulin (Mass conservation)

269 Water balance within the moulin and the subglacial channel is dictated by recharge from a supraglacial stream ( $Q_{in}$ , Sect.  
 270 2.4.1), discharge through a subglacial channel ( $Q_{out}$ ,  $Q_{base}$ ; Sect. 2.4.2) and any change in volume due to melting or refreezing,  
 271 such that the volume of water in the system ( $V$ ) is:

$$272 \quad \frac{dV}{dt} = Q_{in} - Q_{out} + Q_{base} + \frac{(dV_{wallmelt} - dV_{frz})}{dt} \quad (21).$$

273 The final term varies in space and time, with high rates of volume lost to melt above the water line during the melt  
 274 season (when  $Q_{in} > 0$ ), and moderate rates of volume lost to melt at and below the water line during and after the melt season,  
 275 when there is water flow through the moulin ( $Q_{out} > 0$ ) and refreezing below the water line throughout the winter (when  $Q_{in}$   
 276  $= Q_{out} = 0$ ). The MouSh model can also accept an additional prescribed baseflow  $Q_{base}$  directly to the subglacial module. We  
 277 design baseflow as a loose approximation of additional subglacial water inputs from varied upstream sources, including other  
 278 moulins on the same subglacial channel, regional basal melt, and the addition and removal of meltwater from subglacial  
 279 storage. Baseflow is generally required to maintain realistic moulin water levels. In the moulin runs forced by realistic  $Q_{in}$ , we  
 280 represent subglacial flow from  $\sim 5$  surrounding moulins by prescribing baseflow as five times the running 5-day mean of  $Q_{in}$ .  
 281 In other model runs, we do not include baseflow. The addition of baseflow is designed to represent the widespread seasonal  
 282 evolution of surface melt; its inclusion maintains a slightly larger subglacial channel than would otherwise occur, which  
 283 reduces otherwise unrealistically large daily swings in modeled moulin water level (Supplement 2.2.5).

#### 284 **2.4.1 Meltwater runoff from the ice-sheet surface**

285 We force the MouSh model with time-varying water inputs from the supraglacial environment,  $Q_{in}$ . We use two different  $Q_{in}$   
 286 scenarios: a simple diurnal cosine with a mean discharge of  $5 \text{ m}^3 \text{ s}^{-1}$ , in rough agreement with observations near the margins of  
 287 the GIS (Eq. 22, Chandler et al., 2013; McGrath et al., 2011; Smith et al., 2017); and realistic supraglacial discharge over a  
 288 melt season, determined by using in-situ surface melting data and internally drained catchment size and geometry (Yang and  
 289 Smith, 2016).

290 We use the following cosine curve to represent our simplest form of supraglacial discharge into the moulin during  
 291 sensitivity studies:

$$292 \quad Q_{in} = \cos(\pi(t - 19.5)/12) + 5 \quad (22)$$

293 Here,  $t$  is time in hours and  $Q_{in}$  is in  $\text{m}^3 \text{ s}^{-1}$ . This function has its daily peak at 19:30 hours and a daily minimum at 07:30.

294 To examine a set of realistic moulins, we select three supraglacial basins from Yang and Smith (2016) and extract  
 295 their size, distance from terminus from information provided therein (Basin 1-3; Table 2). We derive surface runoff from  
 296 MERRA-2 reanalysis (Gelaro et al., 2017; Smith et al., 2017). Further details on supraglacial characteristics are included in  
 297 Sect. 2.5.3.

#### 298 **2.4.2 Water flow from the subglacial system**

299 We couple the moulin model and a single evolving subglacial channel controlled by melt opening and creep closure (Covington  
 300 et al., 2020; Schoof, 2010) using a reservoir-constriction model (Covington et al., 2012) that simulate flows between the moulin  
 301 and subglacial channel. Following Covington et al. (2020), the rate of change of moulin water level  $h_w$  is

$$302 \quad dh_w = \frac{1}{A_m(h_w)} dV \quad (23)$$

303 With the change in water volume within the system being  $dV$  and the volume of the moulin-subglacial system is related to the  
 304 channel  $S$  and the moulin cross-sectional area  $A_m$ . The water volume is related to  $Q_{in}$ ,  $Q_{base}$  and  $Q_{out}$ , where  $Q_{out}$  is the  
 305 meltwater output from the subglacial channel, defined as follows:

$$306 \quad Q_{out} = c_3 S^{5/4} \Psi / \sqrt{|\Psi|} \quad (24)$$

307 The hydraulic gradient  $\Psi = -\rho_i g \frac{d(h_w - b)}{dL}$  is a linear gradient in  $h_w$  to the outlet at a horizontal distance  $L$ , where the pressure  
 308 head is zero. In our calculations, the bed elevation  $b$  is zero. Finally,  $c_3$  is a flux parameter:

$$309 \quad c_3 = \frac{2^{5/4}}{\pi^{1/4}} \sqrt{\frac{\pi}{(\pi+2)\rho_w f_{sub}}}. \quad (25)$$

310 Equation (25) for  $c_3$  follows Covington et al. (2020), who corrected a small error from the original Schoof (2010) formulation.

311 We use an equation from Schoof (2010) for the time rate of change in subglacial channel cross-section area  $S$ , with  
 312 the first part describing the turbulent melting of the subglacial channel walls, and the second term describing closure due to  
 313 the pressure of the overlying ice, which is dependent on effective pressure  $N = P_i - P_w$ :

$$314 \quad dS = (c_1 Q_{out} \Psi - c_2 N^n S) dt \quad (26)$$

315 Here, the constant  $c_1 = \frac{1}{\rho_i L_f}$  and the constant  $c_2 = 2A_{sub} n^{-n}$  with the Glen's flow law parameters for the subglacial component  
 316 defined as  $A_{sub} = 6 \cdot 10^{-24} \text{Pa}^{-3} \text{s}^{-1}$ .

317 Replacing  $Q_{out}$ ,  $\Psi$ , and  $N$  in Eq. (26) yields

$$318 \quad dS = c_1 c_3 S^{5/4} \left(\frac{\rho_w g h_w}{L}\right)^{3/2} - c_2 (P_i - \rho_w g h_w)^n S) dt \quad (27)$$

319 Equations (23) and (27) are numerically solved simultaneously, as in Schoof (2010) and Covington et al. (2021). The  
 320 parameters used in this module are included in Table 1 and are the same as those used in the englacial component of MouSh,  
 321 apart from the flow law parameter  $A_{sub}$ . In the englacial system,  $A$  is calculated from local temperature within the ice column,  
 322 which can be as cold as  $-23^\circ\text{C}$  in western Greenland (Iken et al., 1993). This contrasts with the temperature at the ice-bed  
 323 interface, which must be at the melting point; the subglacial component of MouSh uses a fixed  $A_{sub}$  value.

324 In its current configuration, the subglacial module provides a single set of outputs representative of conditions at the  
 325 moulin. This is primarily because this study focuses on the evolution of a moulin and is not representative of a channel running  
 326 from a moulin to the terminus in a natural system. A more complex subglacial model would more accurately resolve the spatial  
 327 changes in subglacial channel geometry and flow.

## 328 2.5 Suites of model experiments

329 To examine the sensitivity of the MouSh model to uncertain parameters, ice and meltwater characteristics, and model choices,  
 330 and difference from previous moulin parameterizations, we run four suites of experiments. While these experiments do not  
 331 cover the complete range of possibilities, they were designed to address primary uncertainties in the MouSh model and examine  
 332 how moulin geometry might vary spatially and temporally.

### 333 2.5.1 Quasi-equilibrium and the impact of diurnal supraglacial variability

334 Under steadily varying conditions such as a repeating diurnal variation, the modeled moulin reaches a quasi-equilibrium state  
335 independent of initial conditions with melting opposing viscous and elastic deformation and the only change being driven by  
336 shear deformation. Moulin water level and shape respond to these patterns of variability. To examine the impact of  $Q_{in}$   
337 magnitude (mean) and  $Q_{in}$  amplitude (variability), we perform a series of model runs that vary the magnitude of a cosine curve  
338 between 1 and 20  $\text{m}^3 \text{s}^{-1}$  with a fixed amplitude of 0.5  $\text{m}^3 \text{s}^{-1}$  and a series of runs that vary the amplitude of a cosine curve  
339 between 0 and 2  $\text{m}^3 \text{s}^{-1}$  with a fixed magnitude of 5.0  $\text{m}^3 \text{s}^{-1}$ . The amplitude is one half the diurnal range. These runs use Basin  
340 1 ice conditions (Table 2; Sect. 2.5.3). Further details can be found in Supplement S2.1 and Figures S2-4.

### 341 2.5.2 Sensitivity to uncertain parameters

342 We explored the sensitivity of our results to the values of seven parameters, shown in Figs. 3-5, with the prescribed ranges  
343 shown in Table 1. We examined the effect on the water level, the moulin radius at the equilibrium water level, the volume and  
344 water storage of the moulin, and the cross-sectional area of the subglacial channel at the end of a 40-day model run. These  
345 values reach equilibrium, with daily oscillations superimposed, after  $\sim 15$  days. We also tested the dependence of our results  
346 on the initial moulin radius,  $r_0$ , which we varied across an order of magnitude from 0.65 to 5.0 meters.

347 We varied the value of a uniform deformation enhancement factor  $F^*$  over an order of magnitude ( $F^* = 1$  to 9), which  
348 affects viscous flow of the ice surrounding the moulin. While the range of enhancement factors tested here cover a variety of  
349 ice conditions, including ice shelves and temperate glaciers, the GIS likely has values between 4 and 6 (e.g., Cuffey and  
350 Paterson, 2010). Outside of testing the model sensitivity to the enhancement factor, we assigned  $F^* = 5$ . We also tested the  
351 effect of ice temperature, independent of the enhancement factor. We used five different temperature profiles: cold ice  
352 temperatures (mean  $\sim -15^\circ\text{C}$ , range  $-23.1^\circ\text{C}$  to the pressure melting point) measured in the center of Jakobshavn Isbræ (Iken  
353 et al., 1991); moderate ice temperatures (mean  $\sim -7^\circ\text{C}$ , range  $-13.5^\circ\text{C}$  to the pressure melting point) measured at the GULL  
354 site in Pâkitsoq (Lüthi et al., 2015; Ryser et al., 2014); warmer ice temperatures (mean  $\sim -5^\circ\text{C}$ , range  $-9.3^\circ\text{C}$  to the pressure  
355 melting point) measured at the FOXX site in Pâkitsoq (Lüthi et al., 2015; Ryser et al., 2014); a hypothetical linear profile from  
356  $-5^\circ\text{C}$  at the surface to  $0^\circ\text{C}$  at the bed; and, finally, a fully temperate ice column. These different ice temperature scenarios  
357 affected the creep closure rates of ice through the temperature-dependent softness parameter  $A$  by approximately a factor of 6  
358 from the coldest profile (Iken et al., 1993) compared to the fully temperate column.

359 We also examined moulin sensitivity to elastic deformation by varying Young's modulus ( $E$ ) of the ice column  
360 between 1–9 GPa (Vaughan, 1995) and the sensitivity to the values of friction factors for the moulin walls. MouSh has two  
361 friction factors:  $f_m$  (below the water line) and  $f_{oc}$  (above the water line). We varied these friction factors across two orders of  
362 magnitude (0.01 to 1). We did not vary the subglacial channel friction factor. Finally, we varied values for basal ice softness  
363  $A_{sub}$  over two orders of magnitude ( $5\text{e-}25$  to  $5\text{e-}23$ ) and independently examined moulins over a range of ice thicknesses (670–

364 1570 m) and corresponding distance from the terminus (~20–110 km), which in combination results in variations in hydraulic  
365 gradient.

### 366 **2.5.3 Sensitivity to local conditions**

367 We examined moulins over a range of ice thicknesses and corresponding distances from the terminus (Table 2). Each moulin  
368 is associated with a supraglacial basin derived by Yang and Smith (2016). The moulins were selected based on ice thicknesses  
369 that broadly represent the range of ice thicknesses within the ablation zone of the western GIS and supraglacial drainage basin  
370 sizes and geometries that were visually similar to nearby drainage basins and approximately representative of the mean  
371 supraglacial drainage basin area for the given ice thicknesses (553m, 741m, and 1315m). To derive broadly representative  $Q_{in}$   
372 values for each basin, we integrate 3-hourly modeled surface melting from a downscaled version of MERRA-2 (Gelaro et al.,  
373 2017) over the surface area of each moulin surface drainage basin. We then use synthetic unit hydrograph parameters derived  
374 for a supraglacial basin from western Greenland during the middle of the 2019 melt season (Smith et al., 2017) to estimate  
375 supraglacial discharge into each moulin. Surface runoff values for the 2019 melt season were modified using a synthetic unit  
376 hydrograph derived for the ablation zone and parameters appropriate for western GIS (Smith et al., 2017) with manual  
377 dampening of diurnal variability to minimize long periods of no surface melt during the beginning and end of the season. We  
378 apply this dampening because the parameters for the unit hydrograph were determined during the middle of the melt season  
379 and therefore may inaccurately represent routing delays at the beginning and end of the melt season.

380 The supraglacial discharge curves for each moulin are only meant to capture the seasonal change in discharge rates  
381 and diurnal variability and occasional increases in runoff due to surface melt events during the 2019 melt season. The primary  
382 goal of this exercise is to examine season-long and daily differences in model outputs, the variation in each model component  
383 (viscous, elastic and phase change), and the relative importance of each component in driving moulin geometry and water level  
384 change at different representative locations of the western GIS (Figs. 6-9).

### 385 **2.5.4 Comparison to a cylindrical moulin**

386 Subglacial models generally use a time-invariant vertical cylinder to represent moulins. To investigate and quantify the efficacy  
387 of our time-evolving moulin shape model, we drove MouSh and a static cylinder with the same meltwater inputs. We use the  
388 time-mean radius at the water level as the radius of the static cylinder; this is 1.6 m and 1.4 m for Basin 1 and Basin 2,  
389 respectively. We compared the resulting moulin water level, moulin capacity, subglacial cross-sectional area and meltwater  
390 input difference (due to melt generated within the model itself) across these runs. We compared the moulin water level values  
391 directly (*cylindrical water level – variable water level*) and moulin capacity by percentage difference (*cylindrical – variable*)  
392 / (*variable*); differences are presented in Figure 10.

### 393 2.5.5 Sensitivity to model choices

394 As part of MouSh development, we made several decisions about how to represent moulin geometry, water inputs, and the  
395 associated subglacial system that can directly impact the shape and water level of a modeled moulin. We test the impact of  
396 these decisions in a series of experiments, including (1) representing moulin cross-sectional area as a semi-elliptical, semi-  
397 circular “egg” instead of as a circle (Sect. 2.1); (2) the inclusion of elastic deformation (Sect. 2.2.1); (3) the use of a parabolic  
398 ice sheet profile to determine the surface slope and distance to terminus for a given ice thickness (Cuffey and Paterson, 2010);  
399 (4) the use of prescribed baseflow into the subglacial component of the model (Sect. 2.4); and (5) the use of a time-evolving  
400 subglacial channel (Sect. 2.4.2).

401 To explore the impact of our model choices for experiments 1-4, we perform a series of comparisons against a slightly  
402 modified seasonal run for Basin 1. This allows us to capture the effect of our choices during periods of increasing and  
403 decreasing  $Q_{in}$ . We change only the parameter of interest to isolate the effect on moulin water level and moulin capacity, the  
404 two variables that most directly affect water flow within the subglacial system. Further description of these runs is included in  
405 Supplement S2.2 and resulting differences are highlighted in Figures S5.

406 The first two choices pertain to the complexity of the model, with our choices being more complex; simplification  
407 may be beneficial in some circumstances. In experiment 1, the model is initialized with the same circular geometry as the  
408 control run (Supplement S2.2.1) but melting above the water line is uniformly distributed around the moulin perimeter, thus  
409 there is only one radius to evolve (Supplement S2.2.2). In experiment 2, we test model sensitivity to the inclusion of elastic  
410 deformation (Supplement S2.2.3).

411 Experiments 3 - 5 reflect the simplicity of the current subglacial hydrologic model and would be eliminated if MouSh  
412 was configured to function with either specific observational data or with a more comprehensive subglacial model. In  
413 experiment 4, we test using a subglacial channel length of one half, and one and one half the length defined in the control run  
414 (Supplement S2.2.4). In experiment 4, we prescribe a lower baseflow (Supplement S2.2.5).

415 In experiment 5, we examine the effect of an evolving versus a fixed-geometry subglacial channel (Supplement  
416 S2.2.6). The fixed subglacial channel cross-sectional area is set to 1.95 m<sup>2</sup>. For these runs we use a simpler  $Q_{in}$ , the co-  
417 sinusoidally varying function described in Sect. 2.4.1. Details about this simplification are described in Supplement S2.2.6  
418 with results in Figure S6.

## 419 3 Results

### 420 3.1 Quasi-equilibrium and dependence on $Q_{in}$

421 Under a constant supraglacial input, the moulin water level, radius, and water capacity reach equilibrium within 15  
422 days (red line, Fig. S2c). However, supraglacial inputs are rarely, if ever, uniform, so under constantly varying conditions, the  
423 moulin will reach a ‘quasi-equilibrium’ state. This is a mean state (geometry, water level, deformation rates) with superimposed

424 variability on the timescale of variations in  $Q_{in}$  alone. Therefore, if the forcing is diurnal, the moulin will exhibit diurnal  
425 variability from a mean state. The quasi-equilibrium state is dependent on model characteristics and parameters (Sect. 3.2;  
426 Supplement S2.2).

427 The magnitude and amplitude of  $Q_{in}$  alter the moulin water level and major radius at the mean water level (a proxy  
428 for moulin geometry) in predictable ways (Fig. S2 and Fig. S3). Increasing the diurnal amplitude of  $Q_{in}$  increases the diurnal  
429 variability and mean moulin water level (Fig. S2b, Fig. S4). This occurs due to the disparate timescales of ice deformation  
430 versus melting. The daily increase in  $Q_{in}$  raises the water level quickly because the moulin and subglacial channel is slow to  
431 expand by melting. Conversely, the nightly fall in  $Q_{in}$  is muted by a fast viscous contraction of the moulin and subglacial  
432 channel. This behavior drives the daily peak in moulin water level higher above the mean water level than daily minimum  
433 water level falls below it (Fig. S2b). The “extra” time spent with higher water levels reduces the visco-elastic closure of the  
434 moulin while also increasing turbulent melting, resulting in a larger moulin, as indicated by the moulin radius at the mean  
435 water level (Fig. S2c). Higher diurnal amplitudes in  $Q_{in}$  magnify this effect.

436 As the  $Q_{in}$  magnitude increases, both the mean water level and its diurnal variability decrease (Fig. S3a-b). This occurs  
437 because the moulin becomes larger in response to increasing  $Q_{in}$  and subsequent increases in subglacial discharge. As the  
438 moulin and subglacial channel widen, they can readily accommodate the fluctuations in  $Q_{in}$  with lower variations in moulin  
439 water level. This accommodation is evident in the moulin radius at the mean water level (Fig. S3c). Higher  $Q_{in}$  magnitude  
440 drives a linear increase in melt rates within the moulin alongside nonlinear increases in visco-elastic deformation, causing an  
441 overall nonlinear increase in mean moulin water level (Fig. S4). However, when moulin water levels exceed flotation, the  
442 moulin grows due to both visco-elastic deformation and melting, resulting in a moulin larger than would be expected based on  
443 the equilibrium water level (blue line, Fig. S3c).

### 444 3.2 Sensitivity of MouSh to parameter values and deformational processes

445 A range of ice characteristics affect the time evolution of moulin geometry. These include the initial moulin size, temperature  
446 and viscosity of the ice column, viscosity of basal ice, friction factors, and ice thickness. Some of these factors are highly  
447 spatially variable (e.g., ice thickness) and others are poorly known (e.g., basal ice viscosity). We quantify the effect of these  
448 factors on moulin water level and moulin volume, moulin geometry, and subglacial channel cross-sectional area over both  
449 multi-day and diurnal timescales by performing multiple independent sensitivity studies (Sect. 2.3).

450 We find that moulins reach a quasi-equilibrium, where the mean moulin water level and the moulin radius at this  
451 location oscillate consistently around a daily mean value, within 15-20 days of model initialization. The quasi-equilibrium  
452 value is independent of the initial moulin radius (Fig. 3a–b, Fig 4a–b).

453 Two primary parameters affect the degree of viscous deformation in the moulin: the ice flow enhancement factor  $F^*$   
454 and the ice temperature profile  $T_i(z)$ . We tested a span of reasonable values representative of glacier and ice sheet ice (Table  
455 1) and found a limited effect on moulin geometry. Equilibrium moulin water level, subglacial channel area, and their diurnal  
456 variabilities remain constant (<0.1% change) over the tested range of these parameters (Fig. 3d,f,h and 4d,f,h). Moulin capacity

457 and water storage show high sensitivity ( $\sim 100\text{--}150\%$  in equilibrium value and  $\sim 100\text{--}200\%$  in diurnal range) across the range  
458 of  $F^*$  and  $T_i$  scenarios tested; a decrease in moulin capacity and water storage pair with an increase in the diurnal variability  
459 for these variables. For instance, varying  $F^*$  across an order of magnitude grew the equilibrium major radius by 26% and  
460 shrank the equilibrium minor radius by 72%, with a net effect that moulins had 65% less volume and 58% less water storage  
461 capacity in softer ice ( $F^* = 9$ ) compared to harder ice ( $F^* = 1$ ) (Fig. 3c–d). Similarly, the different ice temperature profiles we  
462 tested caused variations of 29% in moulin major radius, 65% in moulin minor radius, 63% in moulin capacity and 73% in  
463 moulin water storage, with warmer ice hosting smaller moulins (Fig. 3e–f). We also varied Young’s modulus  $E$  across one  
464 order of magnitude, but this affected moulin radius, water volume, and moulin capacity by  $\sim 0.01\%$ . This is due to the low  
465 magnitude of elastic deformation overall compared to viscous deformation (Fig. 5g).

466 We find that moulin geometry is strongly sensitive to the choice of basal ice softness and the friction factors used  
467 within the moulin ( $f_m$  and  $f_{oc}$ ). Melting due to the dissipation of turbulent energy is partially controlled by the friction factors  
468 chosen for the moulin walls. The friction factor above the water line ( $f_{oc}$ , “open channel”) does not significantly affect moulin  
469 water level ( $<0.1\%$  change for  $f_{oc}$  variations over two orders of magnitude), moulin volume (6%), moulin water storage (0.1%),  
470 or subglacial channel area ( $<0.1\%$ ) over either long or diurnal timescales (Fig. 3m–n and 4m–n). However, like the  
471 deformational parameters, the open channel friction factor does affect moulin radii, with the major radius growing by 50% as  
472 the open channel friction factor increases over two orders of magnitude, and the minor radius decreasing by 24%. This dampens  
473 the diurnal variability in the major and minor radii by 70% and 24%, respectively (Fig. 4m).

474 Increasing the friction factor below the water line ( $f_m$ ) had similar effects to changing  $f_{oc}$ . Increasing  $f_m$  by two orders  
475 of magnitude increased the cross-sectional area of the moulin by 106%, via a 10% increase in the major radius and a 93%  
476 increase in the minor radius. The water volume increased by 127% and the storage capacity increased by 74% (Fig. 3k–l) while  
477 the equilibrium water level and the subglacial channel area changed by  $<0.1\%$ . Increasing  $f_m$  also increased the diurnal  
478 variability of the moulin capacity and water storage by 130% and 126%, respectively, by increasing the diurnal differential  
479 melt rate (Fig. 4k–l).

480 The two parameters which have the largest impact on moulin water level are the basal ice softness  $A_{sub}$  and the moulin  
481 location on the ice sheet, described jointly by the ice thickness ( $H_i$ ) and distance from the terminus ( $L$ ). This sensitivity indicates  
482 an interplay among these parameters, the subglacial hydraulic gradient, and moulin water level.

483 We varied basal ice softness  $A_{sub}$  by two orders of magnitude. Softer basal ice increased the size and storage capacity  
484 of the moulin: the major radius by 23%, the minor radius by 23%, the total capacity by 41%, and the stored water volume by  
485 88% (Fig. 3i–j). These changes also increased the equilibrium water level by 34% and the subglacial channel area by 14%,  
486 unlike tests on englacial parameters ( $F^*$ ,  $T_i$ , and  $E$ ), which did not affect the water level or subglacial channel area. These  
487 changes occur because softer basal ice increases the rate of subglacial creep closure, which reduces subglacial channel cross-  
488 sectional area, which reduces water throughflow in the moulin and increases water level, which in turn reduces the amount of  
489 viscous and elastic radial closure in the moulin. Increasing the basal ice softness to approximately  $10^{-23} \text{ Pa}^{-3}\text{s}^{-1}$  increases the  
490 diurnal variability in the sizes of the subglacial channel and moulin (Fig. 4i–j); however, increasing  $A_{sub}$  above this value



491 causes moulin water levels to rise high enough that diurnal fluctuations are truncated by the ice thickness resulting in an  
492 observed decrease in diurnal range that would not be present in thicker ice (Fig. 4j).

493 We co-varied ice thickness and distance from terminus using a parabolic approximation for a perfectly plastic ice  
494 surface profile (Cuffey and Paterson, 2010); this covariance alters the hydraulic gradient of the system. Changes in ice  
495 thickness from 670 to 1570 m (80%) increase the equilibrium subglacial channel area by 24% and increase equilibrium water  
496 levels by 203% (Fig. 3o–p). Increasing ice thickness and distance from the terminus increases the moulin major and minor  
497 radii by 7%, increases moulin volume by 93%, and increases moulin water storage by 235% (Fig. 4p). We also find significant  
498 increases in diurnal variability in subglacial channel size (29%), water level (178%), moulin radii (major radius 84% and minor  
499 radius 24%), moulin volume (130%), and moulin water storage (750%) in thicker ice farther from the terminus (Fig. 4o–p).

500 Overall, we find that MouSh-modeled moulins are primarily sensitive to the friction factors for water flow through  
501 the moulin, basal ice softness, and location on the ice sheet (ice thickness and distance from the terminus). The results are less  
502 sensitive to englacial material factors that govern elastic and viscous deformation. The observed sensitivity to the ice thickness  
503 and distance from terminus signals that moulin geometry can vary spatially. The sensitivity to friction factors and basal ice  
504 softness indicates that the values of these poorly constrained parameters should be carefully chosen and kept in mind when  
505 interpreting model output.

### 506 3.2.1 Contributions to moulin shape

507 Figure 5 illustrates the role of each process, phase change, viscous deformation, and elastic deformation, in determining moulin  
508 radius under different hydraulic potential gradients with median model values (Table 1). Elastic deformation has little impact  
509 on moulin shape or variability (Fig. 5f,g) and is persistently an order of magnitude smaller than either viscous deformation or  
510 radius evolution due to phase change. Viscous deformation and phase change due to melting peak near the daily maximum  
511 water line, with the daily mean of each increasing with increasing ice thickness (Figure 5f); however, the opposite effect is  
512 observed near the bed, where lower mean water levels in moulins in thinner ice increase viscous deformation at the bed;  
513 melting also increases in response to the higher hydraulic potential gradient.

514 At any given depth, viscous deformation and phase change due to melting are similar below the waterline; however,  
515 the diurnal variation in these parameters is quite different (Fig. 5g). At the mean water level, moulin growth due to melting  
516 varies less than  $0.04 \text{ m day}^{-1}$ , with the shape of the diurnal variability dependent on the parameterization of melting both above  
517 and below the water line. In contrast, viscous deformation displays diurnal variations between  $0.08 \text{ m day}^{-1}$  in the thinnest ice  
518 and more than  $0.21 \text{ m day}^{-1}$  in the thickest ice.

### 519 3.3 Moulin shape in different environments

520 We modeled the seasonal growth and collapse of moulins in a range of environments across the GIS using realistic melt  
521 forcings derived for the 2019 melt season (Sect. 2.4.1 and Sect. 2.5.3). These model runs varied with respect to ice thickness,  
522 moulin distance from the terminus, baseflow, and the magnitude, diurnal range, and seasonal evolution of supraglacial inputs

523 (Table 2; Fig. 6a). Overall, we find that moulin setting affects the scale of diurnal and seasonal variability in the size and water  
524 capacity of moulins as well as the evolution of subglacial channels (Fig. 6 and 7).

525 The sizes of all three modeled moulins reach equilibrium with the melt forcing within ~15 days of the onset of the  
526 melt seasons (Fig. 6b–c). As the water flux increases over the next few weeks, each moulin grows in response to increasing  
527 supraglacial inputs, both diurnally and with a long-term trend, although this growth is more significant in thicker ice (Fig. 6c  
528 and Fig. 7). The subglacial channel grows with a similar pattern, but interestingly, the setting and fluxes of Basin 1 and Basin  
529 2 result in very similar subglacial channel cross-sectional areas despite different moulin water levels and capacities (Fig. 6d).

530 Although the three moulins all evolve in a similar fashion, there are differences in moulin capacity, water level (Fig.  
531 6), overall moulin geometry (Fig. 7), and the magnitude of englacial deformation (Fig. 8). Basin 3 exhibits the largest seasonal  
532 change in moulin capacity in part because a lower supraglacial input and subglacial hydraulic gradient results in a smaller  
533 subglacial channel and periods where moulin water level is above flotation (Fig. 6). This causes substantial variability of  
534 viscous deformation while limiting variations in melt due to changing moulin water level (Fig. 8a). One of the largest periods  
535 of Basin 3 moulin growth occurs starting at day 30. During this period, supraglacial inputs experience a step change (Fig. 7a);  
536 moulin water levels stayed near flotation and were less variable for several days (Fig. 7b), keeping effective pressure near zero  
537 and retarding deformation (Fig. 8a). In this case, viscous deformation hovers around zero and causes moulin opening, resulting  
538 in a high ratio of elastic to viscous deformation and a high ratio of phase change to viscous deformation (purple line in Fig.  
539 8b). Similar behavior also occurs around day 110. Basins 1 and 2 exhibit smaller seasonal variations in moulin capacity because  
540 the ratio of melting to deformation stays near one until near the end of the season (Fig. 8b). This occurs because viscous  
541 deformation in Basins 1 and 2 is only slightly lower than in Basin 3 and melt rates tend to be higher (Fig. 8a) due to increased  
542 subglacial discharge associated with a higher hydraulic gradient. Further, there are fewer periods where water levels above  
543 flotation drive viscous opening.

544 Each moulin has a different daily mean capacity (Fig. 7c). This, in addition to differences in supraglacial inputs,  
545 ensures that daily moulin water level variations are substantially different across moulins. Basin 1 exhibits the largest variation  
546 in daily moulin water level, followed by Basin 2 (Fig. 9a). Basin 3 shows the lowest daily change; however, this is due at least  
547 in part to the fact that water overtops the moulin nearly daily (Fig. 6b and 7m–n). Changing water levels drive changes in  
548 moulin and subglacial capacity. Over the melt season, daily change in moulin capacity can be as low as 2% during lulls in  
549 diurnal melt variability (Basin 3) or as high as 12% following a recovery from a low melt day (Basin 1; Fig. 9b). However, in  
550 general all moulins display a similar daily change in capacity of ~5-10%, with peak values of 12 to 13%.

551 The subglacial system undergoes diurnal variations in channel size between 1 and 20% (Fig. 9c). These changes are  
552 similar in magnitude to daily capacity changes within the moulin but exhibit more variability across ice thicknesses. Like  
553 changes in moulin capacity, these variations are related to the daily changes in moulin water level (Fig. 9a). This suggests that  
554 the time evolution of moulin geometry dampen the diurnal pressure fluctuations that drive subglacial channel growth and  
555 collapse. Evidence for this can be seen in the temporal pattern of moulin water level and subglacial channel cross-sectional  
556 area (Fig. 9a,c).

### 557 3.4 Comparison to cylindrical moulins

558 To examine the role moulin evolution plays in modifying the subglacial hydrologic system, we compared moulin water levels,  
559 moulin capacity, and subglacial channel size between model runs with a fully evolving moulin and runs with a static cylindrical  
560 moulin. We performed these tests with realistic melt inputs based on the 2019 melt season (Sect. 2.4.1, 2.5.3), at moulins with  
561 low and moderate ice thicknesses (553 m – Basin 1 and 741 m – Basin 2). We defined the radius of the static cylinder as the  
562 mean radius at the mean water level: 1.6 m and 1.4 m for Basin 1 and 2, respectively. This results in fixed moulin cross-  
563 sectional areas ( $\sim 6$  to  $8 \text{ m}^2$ ) within the range of spatially invariant moulin cross-sectional areas  $\sim 2\text{--}10 \text{ m}^2$  often prescribed in  
564 subglacial models (e.g., Andrews et al., 2014; Banwell et al., 2013; Bartholomew et al., 2012; Cowton et al., 2016;  
565 Meierbachtol et al., 2013; Werder et al., 2013).

566 Comparison of moulin water level and capacity between static cylindrical and evolving moulins show differences on  
567 both the diurnal and seasonal times scales (Fig. 10). The differences in moulin water level (both positive and negative) are  
568 generally great during lower supraglacial inputs at the beginning and end of the melt season, with the relatively limited  
569 differences occurring during the highest discharges (Fig. 10a-b). These values are both positive, indicating that the static radius  
570 moulin has higher water levels, and negative, indicating that the evolving moulin has higher water levels. Differences in moulin  
571 water level can reach nearly 20 m, but are most commonly below 10 m. The seasonal mean water level difference between the  
572 static cylindrical and evolving moulin in both basins is less than 1 m.

573 Moulin capacity also displays a clear seasonal pattern; in both basins, the static cylindrical moulin larger than the  
574 evolving moulin at the beginning of the melt season with the evolving moulin gradually growing larger as the melt season  
575 progresses (Fig. 10c-d). After peak melt (day  $\sim 60$ ), the evolving moulin begins to viscously close and gradually becomes  
576 smaller than the static cylindrical moulin. The static cylindrical moulin can be more than 100% larger than the variable moulin  
577 during the tails of the melt season with the evolving moulin becoming 36% and 42% larger than the static cylindrical moulin  
578 during mid-melt season. Overall, the mean capacity difference between the static cylindrical and evolving moulin is less than  
579 5%, with the static cylindrical moulin being slightly larger.

580 The radius of the cylindrical moulin was chosen to minimize differences with the evolving moulin. This is evident by  
581 the limited long-term differences between the two moulins in both Basin 1 and 2. As such, there is limited differences ( $<1\%$ )  
582 between the modeled subglacial channels. We expect difference in moulin water level, moulin capacity, and subglacial  
583 geometry to change if the static cylindrical moulin geometry is poorly chosen; if the different or different experimental  
584 parameters are used; or the setting changes (e.g., different hydraulic gradients). For example, we use commonly used values  
585 of ice softness  $A$  for both the moulin and subglacial channel; however, these values are poorly known, and their choice can  
586 directly impact the relative importance of moulin shape in dictating moulin water levels and subglacial channel size (Fig. 4).

### 587 3.5 Impact of model choices on moulin geometry

588 Chosen parameterizations have the potential to impact the representation of moulin water level and capacity (Supplement S2).  
589 Overall, we find that a circular geometry has limited impact on moulin water level with the circular geometry having water  
590 levels that are less than 3 m higher than the egg-shaped geometry, although in nearly all instances the difference is less than  
591 0.5 m (Fig. S5a); however, the impact on capacity is slightly larger (the circular moulin is up to 31% smaller) and displays a  
592 seasonal trend as the egg-shaped moulin elongates along its elliptical axis (Fig. S5b).

593 Elastic deformation within the moulin is small (Supplement S1 and S2.2.3; Figure 8a). Excluding elastic deformation  
594 has a negligible impact on moulin water levels and moulin capacity ( $< 1\%$ ; Figure S5c-d).

595 In contrast to the previous choices, the distance from the terminus  $L$  and the prescribed baseflow  $Q_{base}$  can have a  
596 substantial impact on moulin water level and capacity (Fig. S5e-h). Distance from the terminus is defined by the position of a  
597 given moulin on the ice sheet, and as such is not a choice or parameter per se; however, does directly influence the hydraulic  
598 gradient. A shorter  $L$  increases the hydraulic gradient and reduces both moulin water levels and capacities (Fig. S5e-f).  
599 Baseflow is used here to mitigate the use of a simplistic subglacial hydrology model. Reducing the baseflow within the  
600 subglacial system increases moulin water levels and reduces moulin capacity (Fig. S5g-h).

601 Finally, we examine the impact of fixing the subglacial channel cross-sectional area  $S$ . Experimental results using a  
602 fixed  $S$  and a seasonally evolving melt curve resulted in unrealistically low or zero water levels during low, early season  $Q_{in}$   
603 and complete viscous collapse of the moulin if the subglacial channel size was prescribed to be too large, or persistently high  
604 (always above the ice thickness) water levels and runaway moulin growth if the subglacial channel was prescribed to be too  
605 small. Therefore, we explore the impact of fixing  $S$  using a constant mean  $Q_{in}$  with an overlaid diurnal variability (Supplement  
606 S2.2.6). With constant variability, we can easily prescribe the fixed  $S$  to be the mean value of the time-varying subglacial  
607 channel  $S$  (1.95 m). In this instance, the fixed  $S$  experiment displays a similar mean moulin water level, but lower diurnal  
608 variability than the experiment with a time-varying  $S$  (Fig. S6). Further details are included in the Supplement S2.

## 609 4 Discussion

### 610 4.1 Timescales of moulin formation and evolution

611 We consider the formation timescales of moulins in the context of the shape evolution of a mature moulin. Using MouSh, we  
612 find that in the absence of external forcing, such as time-variable  $Q_{ins}$ , the size of a moulin reaches its equilibrium value in  $\sim 15$   
613 days depending on ice and supraglacial input conditions and initial moulin geometry (Fig. 5g, Fig. S2 and Fig. S3). This  
614 relaxation time is comparable to the Maxwell time for ice (10–100 hours), as expected for a linear visco-elastic system. Our  
615 relaxation time also compares well to the equilibration timescale defined by Covington et al. (2020) for their modeled moulin  
616 – subglacial conduit system, which Trunz (2021) found to be 1–20 days. The most realistically sized moulins in Trunz (2021)

617 had relaxation times closer to 1 day. Their modeled system was governed solely by melt and viscous deformation and lacked  
618 elastic deformation; however, elastic deformation in MouSh is small explaining why our relaxation times are comparable.

619 If the process of moulin formation occurs on a timescale shorter than the 15-day relaxation time, the formation process  
620 likely will not influence the overall form of the englacial system at equilibrium. This time range includes hydrofracture during  
621 rapid lake drainage (~2 hours) and slow lake drainage (<~6 days, e.g., Selmes et al., 2011), and likely also the reactivation of  
622 existing moulins in ensuing melt seasons, which, based on the timing difference between surface melt onset and ice  
623 acceleration, occurs over multiple days (Andrews et al., 2018; Hoffman et al., 2011). On the other hand, moulin formation by  
624 cut-and-closure occurs over years to decades (Gulley et al., 2009), well above the MouSh relaxation time and the Maxwell  
625 time for ice, are more likely to create subvertical englacial channels. The interdependence of formation and evolution of these  
626 moulins gives us less confidence in applying our model to moulins with cut-and-closure origins. Those moulins primarily  
627 occur in temperate near-surface ice within polythermal glaciers (Gulley et al., 2009) and have not been reported on the GIS.

#### 628 **4.2 Comparison of modeled and observed moulin geometries**

629 Field observations suggest that moulin geometry evolves a high degree of complexity. Observations include anecdotes of  
630 difficulty deploying sensors to the bottom of a moulin, which suggests the presence of kinks, ledges, knickpoints, and other  
631 twists (Andrews et al., 2014; Covington et al., 2020; Cowton et al., 2013). Complex geometry revealed during mapping moulins  
632 above the water line further suggests that moulins are not simply vertical cylindrical shafts (Covington et al., 2020; Moreau,  
633 2009).

634 The MouSh model suggests that the energy transfer from turbulent meltwater entering the moulin to the surrounding  
635 ice drives highly spatially variable melt rates above the water line. We incorporated the open-channel melt module to allow a  
636 large opening to emerge above the water line (Fig. 5a–e and 7). When we run MouSh without the open-channel module, the  
637 surface expression of the moulin becomes dependent on surface stresses and can in some instances pinch closed. The open  
638 channel module also permits the development of an egg-shaped geometry, which is supported by seismic observations and a  
639 resonance model of a moulin which suggests that moulins may increase in ellipticity over time (Rössli et al., 2016).

640 The value of the open-channel friction factor and the size of the spatial footprint over which melting occurs directly  
641 affects the size of the upper, air-filled chamber of the moulin, which differs from when the moulin is modeled as circular and  
642 open-channel melting is applied uniformly around the perimeter (Fig. S5b). MouSh can predict ledges at the top and bottom  
643 of a consistent diurnal range in water level. Thus, we infer that energetic subaerial water flow drives formation of moulin  
644 complexity above the water line, and diurnal fluctuations around a steady multi-day water level drive ledge formation through  
645 a differential in melting and viscous deformation above and below the water line. Energetic water flow is commonly observed  
646 at stream-fed moulins near the peak of the melt season (Pitcher and Smith, 2019) or during and immediately following rapid  
647 lake drainage (Chudley et al., 2019). This suggests that complex moulin geometries form during periods of relatively consistent  
648 water supply. Conversely, multi-day rises in water level, driven by either the surface water supply or the basal water supply  
649 (baseflow), can erase geometric complexities such as ledges, as seen in MouSh results during a melt event (Fig. 7).

650 Above the water line, explored moulins in Greenland show highly variable shapes from moulin to moulin (e.g.,  
651 Covington et al., 2020). Some moulins, for example the FOXX moulin, are nearly cylindrical within the explored depth (~100  
652 m), with radii comparable to what we model (~1.5 m). Others open some tens of meters below the surface to large caverns  
653 with radii approaching 10 meters, a similar morphology to karst caves with narrow entrance shafts (Covington et al., 2020).  
654 MouSh can produce large openings above the water line if we use a suitably large open channel friction parameter, although  
655 we lack a narrow entrance shaft and substantial vertical variability. These differences are due to the inability of model  
656 parameterizations to represent complex geometries such as scalloping, plunge pools, and knickpoint migration (Gulley et al.,  
657 2014; Mankoff et al., 2017). Indeed, instead of modeling processes above the water line as turbulent open flow, they could, in  
658 the future, be modeled using geomorphic parameterizations to model waterfall migration, perhaps resulting in the clearer  
659 development of steps and plunge pools. This would require development and inclusion of a supraglacial channel model as  
660 well.

661 Below the water line, MouSh results indicate that a cylinder is a reasonable representation for newly formed moulins  
662 in Greenland. However, there are two caveats. First, moulin cross-sectional area, and thus water storage capacity, can vary  
663 substantially over the course of a day or season (Fig. 6c. and 9b) and features such as englacial crevasses and reservoirs may  
664 be present (e.g., McQuillan and Karlstrom, 2021). Second, in instances where moulins are reactivated over multiple melt  
665 seasons (Chu, 2014; Smith et al., 2017), there may be substantial deformation, as suggested by cable breakage in boreholes  
666 (Ryser et al., 2014; Wright et al., 2016).

667 Observations show a wide range of moulin volumes above the water line, and moulin volumes predicted by MouSh  
668 are sensitive to the consideration of turbulent melting and associated parameter choices. Given the flexibility of model results,  
669 we should continue to rely on field exploration to measure moulin size and geometry above the water line and make efforts to  
670 constrain the parameters that affect sub-seasonal growth and collapse. MouSh results below the water line are less sensitive to  
671 uncertain parameter values, so direct observations of underwater geometry would be less relevant for model validation than  
672 subaerial observations. Overall, results from the MouSh model demonstrate that moulin geometry evolves substantially over  
673 diurnal to seasonal timescales and varies with ice conditions.

#### 674 **4.3 Diurnal water level oscillations and moulin size**

675 Moulin geometry can directly alter the relationship between meltwater inputs and moulin water level changes – the primary  
676 driver of subglacial channel evolution (Andrews et al., 2014; Cowton et al., 2013). Field measurements of moulin water levels  
677 indicate diurnal oscillations of 3–12% (Covington et al., 2020), ~25% (Andrews et al., 2014), and >20% (Cowton et al., 2013)  
678 of overburden pressure with mean water levels of ~70% of overburden. These diurnal fluctuations are larger than those  
679 observed in boreholes, which are generally, though not always, thought to sample inefficient components of the subglacial  
680 hydrologic system (Andrews et al., 2014; Meierbachtol et al., 2013; Wright et al., 2016).

681 Our model results agree well with observations of moulin water level: diurnal fluctuations of approximately 25 to  
682 50% of overburden pressure, with larger absolute oscillations occurring in thicker ice. To explain larger-than-expected daily

683 oscillations (~10%) in thinner ice, Covington et al. (2020) incorporated moulin cross-sectional area as a free parameter into  
684 their model. Matching field measurements of water level required a modeled moulin radius of ~5 m (~75 m<sup>2</sup> cross-sectional  
685 area) at ice thickness 500 m and a much larger moulin (radius ~20 m and cross-sectional area ~1500 m<sup>2</sup>) at ice thickness 700  
686 m (Covington et al., 2020). For comparison, MouSh predicts average radii ~1.4 m (~6 m<sup>2</sup> cross-sectional area) in similar ice  
687 thicknesses using parameters described in Table 2, including substantially larger meltwater inputs compared to Covington et  
688 al. (2020). The drastic differences in moulin size despite similar variations in diurnal water level between our study and  
689 Covington et al. (2020) cannot easily be attributed to a single factor but may be explained by our limited ability to model  
690 processes above the water line, our inclusion of baseflow (Fig. S5g–f), substantial differences in meltwater input (e.g., Fig. S2  
691 and S3), fluctuations in moulin capacity (Covington et al. (2020) use a fixed moulin geometry), or that their measured water  
692 levels were not from the same moulin they mapped englacially. However, our results suggest that an evolving moulin capacity  
693 may be important to represent realistic moulin water levels and capacity (Fig. 10). Thus, to match observed moulin water level  
694 fluctuations without evolving the moulin geometry, a fixed cross-sectional area substantially larger than the associated  
695 subglacial channel may be necessary, as reported in Covington et al. (2020).

#### 696 **4.4 Magnitude of viscous moulin deformation**

697 Viscous and elastic deformation drive moulin closure. The role of elastic deformation in the glacial hydrologic system is  
698 discussed below (Sect. 4.4); viscous deformation is the primary closure mechanism of moulins, boreholes and subglacial  
699 channels (e.g., Catania and Neuman, 2010; Paterson, 1977, Shreve, 1972), with viscous deformation dependent on local  
700 effective pressure, ice characteristics, and the geometry of the feature of interest (Flowers, 2015). Viscous deformation within  
701 our moulin varies in response to meltwater inputs (Fig. 5g and Fig. 8a) with the highest deformation rates occurring at the  
702 water line (Fig. 5f) because at the water line, inward cryostatic pressure is least offset by outward hydrostatic pressure (see Eq.  
703 3).

704 During our realistic runs, viscous deformation can exceed 0.25 m day<sup>-1</sup> for short periods of the day at all three moulin  
705 locations (Fig. 8a). These deformation rates are substantially larger than measured borehole deformation rates for the primary  
706 reasons that boreholes are often at or above flotation due to high subglacial water pressures (e.g., Ryser et al., 2014) or because  
707 creep measurements are recorded in much smaller boreholes in colder ice (e.g., Paterson, 1997).

708 A previous moulin modeling effort focused on understanding moulin closure rates (Catania and Neumann, 2010).  
709 Their results indicate that an air-filled moulin will close within a single day at the bed. However, in this instance there is no  
710 opposing hydrostatic pressure. While our modeled closure rates are similar to those calculated by Catania and Neuman (2010)  
711 near the surface, the moulins modeled here always contain water even at the end of the melt season (Fig. 6b). This continued  
712 retention of meltwater is in line with borehole observations that subglacial pressures tend to be highest outside the melt season  
713 (Downs et al., 2018) and preclude the presence of completely air-filled moulins in areas where viscous deformation rapidly  
714 shuts down the hydrologic system as supraglacial inputs fall.

#### 715 **4.5 The role of elastic deformation in ice sheet hydrology**

716 Our model results indicate that the equilibrium moulin geometry is dictated by a balance of visco-elastic deformation and  
717 turbulence-driven melting (Fig. 5 and Fig. 8). In both the sensitivity study and realistic model experiments, visco-elastic  
718 deformation generally closes the moulin, while melting of the surrounding ice consistently opens the moulin. The exception is  
719 when moulin water levels exceed flotation, in which case all three mechanisms open the moulin. In all model runs, we find  
720 that the rates of viscous deformation exceed elastic deformation by three to four orders of magnitude (Fig. 5g and Fig. 8).  
721 Elastic deformation rates are greatest near the water line and at the bottom of the ice column, where stress conditions are  
722 similar to those in subglacial models, at a few centimeters per year of closure within a moulin of radius  $\sim 1$  meter. This moulin  
723 size is comparable to that of a typical subglacial channel in our model ( $A \sim 2 \text{ m}^2$ , or radius 1.1 m), implying that elastic closure  
724 of a subglacial channel would also amount to a few centimeters per year. Elastic closure rates scale linearly with moulin  
725 radius; thus, larger moulins or channels would undergo commensurately faster elastic closure. The contribution of elastic  
726 deformation relative to viscous deformation increases with increasing ice thickness (Fig. 5f); at  $H = 670$  meters, viscous  
727 deformation is 4000 times larger than elastic deformation at the water line, while at  $H = 1570$  meters, it is 2000 times larger.

728 Current subglacial hydrology models represent subglacial channel development (opening) by turbulent energy  
729 dissipation and destruction (closing) by viscous deformation alone. However, work involving elastically responding storage  
730 elements or elastic flexure of the ice sheet has occurred (Clarke, 1996; Dow et al., 2015), and there have been efforts to use  
731 elastic deformation or fluid compressibility to improve numeric stability of channel equations (Clarke, 2003; Spring and Hutter,  
732 1981, 1982). Interestingly, Clarke (2003) chose to use fluid compressibility due to model integration times. Yet, elastic  
733 deformation has generally been omitted from current models of glacial hydrology, even when modeling rapid changes in  
734 meltwater inputs ( $< 1$  day; e.g., Hewitt, 2013; Hoffman et al., 2016; Werder et al., 2013). Our investigation of the role of elastic  
735 deformation in diurnally closing moulins, particularly in thicker ice (Fig. 8b and S5c,d), suggests that its exclusion from  
736 subglacial channel models should cause errors of  $< 0.1\%$  and is warranted. On length scales considerably larger than  $\sim 1$  meter  
737 moulins, as well as in problems where elastic flexure is more central to the geometry, elastic deformation remains important.  
738 These applications include ice shelves (e.g., Reeh et al, 2003; Walker et al., 2016), large marine-terminating glaciers  
739 (Christmann et al., 2021), crevasse opening (Poinar et al., 2017), and rapid supra- and subglacial lake drainage (Dow et al.,  
740 2016; Dow et al., 2015; Lai et al., 2021).

#### 741 **4.6 Moulin geometry and the englacial void ratio**

742 Subglacial hydrology models use an englacial void ratio parameter to represent bulk storage and release of meltwater in the  
743 englacial system (see Flowers and Clarke (2002) for the best description). Because the englacial void ratio acts as short term,  
744 pressure dependent, storage for subglacial models, it can improve the representation of diurnal water pressure fluctuations in  
745 subglacial models (Flowers and Clarke, 2002). This parameter represents bulk behavior and is usually tuned by comparing to  
746 local observations (e.g., Bartholomaus et al., 2011; Hoffman et al., 2016; Werder et al., 2013). The inclusion of time-varying



747 moulin geometry, potentially in addition to time varying representation of englacial fractures (Gajek et al., 2021), that evolve  
748 in response to meltwater inputs and subglacial pressures could reduce subglacial model dependence on this parameterized  
749 englacial storage, particularly in light of observations of time varying englacial features (Church et al., 2020) and meltwater  
750 content (e.g., Vankova et al, 2018).

751 MouSh demonstrates that moulin capacity can vary both seasonally and during large variations in supraglacial input.  
752 Moulin growth rates are largest particularly when water levels are above flotation, when both melting and viscous deformation  
753 work to increase moulin capacity. Our results show that moulin capacity changes by up to 13% daily under realistic conditions  
754 (Fig. 9b) and 87 to 138% over the melt season (Fig. 6c). These variations in moulin shape and size may explain difficulties  
755 with modeling subglacial behavior during melt events (Cowton et al., 2016), which are sometimes addressed by temporarily  
756 increasing englacial storage (Hoffman et al., 2016). Our results with MouSh suggest that modeling moulin shape alongside the  
757 evolution of the subglacial system could potentially improve the representation of subglacial pressures, especially during  
758 periods of large meltwater variability; however, additional development is necessary to explore the impact of multiple moulins  
759 evolving along with the subglacial system.

760 Practical limits on model complexity or computational costs may preclude fully time-evolving moulin geometries.  
761 While not ideal, an arbitrary static shape may preferable to a static cylinder (Trunz, 2021). Therefore, we interpret our moulin  
762 shape results (Fig. 7) to recommend a representative shape for a static moulin. Below the water line, a cylinder is a reasonable  
763 approximation, especially in thinner ice or for newly made moulins, for which full-column ice deformation is minimized.  
764 Above the water line, moulin shape is widely variable in time, by location, and across parameter combinations. It is especially  
765 sensitive to the friction parameter for open-channel flow (Fig. 3m and Fig. 4m), with low friction values making bottle-shaped  
766 moulins that have narrow necks above the water line and larger chambers below the water line, and high friction values making  
767 goblet-shaped moulins with open rooms and amphitheaters above the water line atop a narrower geometry below the water  
768 line. Exploration of Greenland moulins to date has uncovered multiple goblet-shaped moulins and a few instances of near-  
769 cylindrical moulins, but no bottle-shaped moulins (Covington et al., 2020; Moreau, 2009; Trunz, 2021). Overall, our MouSh  
770 results support goblet-shaped moulins, although with great variation in the height and width of the upper chamber.

#### 771 **4.7 Limitations of the current MouSh englacial – subglacial model**

772 Moulins are a dynamic component of the channelized englacial–subglacial system; therefore, explicitly modeling their  
773 evolution can improve the accuracy of glacial hydrology models. MouSh currently uses a single subglacial channel to represent  
774 the entire subglacial system, limiting its accuracy. An optional baseflow term, which parametrizes subglacial water flow from  
775 surrounding regions, improves MouSh performance. This baseflow, added directly to the subglacial channel, is necessary to  
776 produce realistic equilibrium water levels with the realistic supraglacial inputs we prescribed (Fig. 6a). The baseflow value we  
777 used does not explicitly represent any specific process because our model runs resolve only a single moulin connected to a  
778 single channel, whereas in the real world, multiple moulins feed a network of channels. The idealized baseflow term  
779 conceptually connects to multiple potential water sources, including (1) basal melting from geothermal and frictional heating,

780 (2) supraglacial water delivered via nearby moulins that are connected to the same subglacial channel, and (3) water that moves  
781 from the channelized system to the surrounding inefficient system at high pressures and then flows back into the subglacial  
782 channel at lower water pressures (Hoffman et al., 2016; Mair et al., 2001, 2002; Tedstone et al., 2015).

783 The addition of baseflow maintains a larger, less variable subglacial channel. This can alternately be achieved by  
784 lessening the local hydraulic gradient, thus increasing the mean water pressure along a given reach. This may locally occur  
785 where one subglacial channel enters another in an arborescent network (Fountain and Walder, 1998). MouSh currently does  
786 not have an interconnected network of channels; however, this is under development (Trunz, 2021).

787 We use a highly simplified model of the subglacial hydrology system: a single channel that connects the moulin to  
788 the ice-sheet margin. Yet, MouSh results clearly indicate that including and evolving a moulin can alter the hydraulic gradient  
789 of the subglacial system via time-varying storage in the moulin (Fig. 10), though in our current single moulin configuration,  
790 there is limited impact on subglacial channel geometry. Further, MouSh currently lacks a distributed system, which limits its  
791 fidelity for assimilating daily meltwater volumes into the subglacial system; realistically, the channelized subglacial system  
792 cannot always accommodate the full volume of meltwater produced during summer days, and a portion of this water goes into  
793 the distributed system (e.g., Mair et al., 2001, 2002). In our model, however, when the system is overwhelmed, the water level  
794 in the moulin rises above what is typically observed, and sometimes even exceeds the height of the ice (Figs. 6b, S4b). The  
795 melt-driven opening and creep closure processes in the subglacial model explain this behavior: A lower water input to the  
796 moulin ( $Q_{in}$ ) lowers the water flux into the subglacial system ( $Q_{out}$ ), which lowers the melt rates that keep subglacial channels  
797 open, reducing the size of the subglacial channels and thus further reducing the subglacial water flux. This increases the water  
798 level in the moulin. Thus, a reduced rate of surface melt can counterintuitively raise the modeled water level (Fig. 6 around  
799 day 30), whereas in reality, much of that water would enter the inefficient subglacial hydrologic system when moulin water  
800 levels exceed flotation. If the moulin model were coupled to a two-component subglacial model that represents the inefficient  
801 system alongside the channelized system, we would anticipate a much-improved ability to assimilate a wide range of meltwater  
802 input rates.

## 803 **5 Conclusions**

804 Results from the MouSh model show that moulins are not static cylinders. Daily fluctuations in moulin capacity change the  
805 water volume held in the englacial hydrologic system, which in turn influences the evolution of the subglacial channels that  
806 moulins feed. When we represent a moulin as a static cylinder in our englacial–subglacial hydrology model, these daily  
807 fluctuations can be overestimated or underestimated, affecting the volume of water stored englacially and the hydraulic  
808 gradient of the subglacial system. Modeled moulin size and shape may provide a more realistic representation of moulin water  
809 level and the englacial void ratio commonly used in subglacial hydrology models, particularly with future efforts to improve  
810 the parameterization of moulin development above the water line. This could be achieved by using an englacial hydrology –  
811 channelized subglacial system model, such as the MouSh model we present here, to characterize variability in moulin size and

812 shape, or by coupling moulin models to more complete models of the subglacial system (channelized, distributed, and  
813 optionally weakly connected) to make a unified englacial–subglacial hydrology model system. Improving the representation  
814 of the englacial–subglacial system to explicitly include moulins would have greatest efficacy during periods of rapidly varying  
815 supraglacial input (e.g., during the beginning and end of the melt season and during melt events) and in inland areas with thick  
816 ice and high overburden pressures. These are coincident with situations where subglacial models without moulins, or with  
817 implicitly static moulins, tend to perform poorly.

818

819 *Code and Data availability.* The Moulin Shape model is publicly available at [https://github.com/kpoinar/moulin-physical-](https://github.com/kpoinar/moulin-physical-model)  
820 [model](https://github.com/kpoinar/moulin-physical-model) (we will make a release when revisions are complete). The model results used in the analysis presented here are archived  
821 at the University at Buffalo Libraries at <http://hdl.handle.net/10477/82587>.

822

823 *Author contributions.* L.C.A. and K.P. jointly conceived of and developed the MouSh model. Both L.C.A. and K.P. designed  
824 the study, executed the model runs, analyzed the data, produced the figures, and wrote the manuscript. C.T. implemented the  
825 subglacial module, participated in discussions, and edited the manuscript.

826

827 *Acknowledgements.* This work was supported by NASA Cryosphere grant 80NSSC19K0054 (L.C.A. and K.P.), the Global  
828 Modeling and Assimilation Office at NASA Goddard Space Flight Center funded under the NASA Modeling, Analysis, and  
829 Prediction (MAP) program (L.C.A.), the Research and Education in eNergy, Environment and Water (RENEW) Institute at  
830 the University at Buffalo (K.P.), and the United States National Science Foundation award number NSF-ANS 1603835 (C.T.).  
831 We acknowledge DigitalGlobe, Inc. for providing WorldView images via the Enhanced View Web Hosting Services and the  
832 support therein provided by the Polar Geospatial Center under NSF-OPP awards 1043681 and 1559691. We thank two  
833 anonymous reviewers and editor Dr. Elizabeth Bagshaw for constructive feedback which substantially improved manuscript  
834 clarity and completeness.

835

836 *Competing interests.* An author is a member of the editorial board of The Cryosphere. The peer-review process was guided by  
837 an independent editor, and the authors have no other competing interests to declare.

## 838 **References**

839 Aadnøy, B. S.: A complete elastic model for fluid-induced and in-situ generated stresses with the presence of a borehole,  
840 *Energy Sources*, 9, 239–259, 1987.

841 Alley, R. B.: Flow-law hypotheses for ice-sheet modeling, *J. Glac.*, 38(129), 245–256,  
842 <https://doi.org/10.3189/S0022143000003658>, 1992.

843 Alley, R. B., Dupont, T. K., Parizek, B. R., and Anandakrishnan, S.: Access of surface meltwater to beds of sub-freezing  
844 glaciers: preliminary insights, *Ann. Glac.*, 40, 8–14, <https://doi.org/10.3189/172756405781813483>, 2005.

- 845 Amadei, B.: *Rock Anisotropy and the Theory of Stress Measurements*, Springer-Verlag, Berlin, New York., 1983.
- 846 Andrews, L. C., Catania, G. A., Hoffman, M. J., Gulley, J. D., Lüthi, M. P., Ryser, C., Hawley, R. L. and Neumann, T. A.:  
847 Direct observations of evolving subglacial drainage beneath the Greenland Ice Sheet, *Nature*, 514(7520), 80–83,  
848 <https://doi.org/10.1038/nature13796>, 2014.
- 849 Andrews, L. C., Hoffman, M. J., Neumann, T. A., Catania, G. A., Lüthi, M. P., Hawley, R. L., Schild, K. M., Ryser, C. and  
850 Morriss, B. F.: Seasonal Evolution of the Subglacial Hydrologic System Modified by Supraglacial Lake Drainage in Western  
851 Greenland, *J. Geophys. Res. Earth Surf.*, 123(6), 1479–1496, <https://doi.org/10.1029/2017JF004585>, 2018.
- 852 Banwell, A. F., Willis, I. C. and Arnold, N. S.: Modeling subglacial water routing at Paakitsoq, W Greenland, *J. Geophys. Res.*  
853 *Earth Surf.*, 118(3), 1282–1295, <https://doi.org/10.1002/jgrf.20093>, 2013.
- 854 Banwell, A. F., Hewitt, I., Willis, I. and Arnold, N.: Moulin density controls drainage development beneath the Greenland ice  
855 sheet, *J. Geophys. Res. Earth Surf.*, 2015JF003801, <https://doi.org/10.1002/2015JF003801>, 2016.
- 856 Bartholomaeus, T. C., Anderson, R. S. and Anderson, S. P.: Growth and collapse of the distributed subglacial hydrologic system  
857 of Kennicott Glacier, Alaska, USA, and its effects on basal motion, *J. Glac.*, 57(206), 985–1002,  
858 <https://doi.org/10.3189/002214311798843269>, 2011.
- 859 Bartholomew, I. D., Nienow, P., Sole, A., Mair, D., Cowton, T. and King, M. A.: Short-term variability in Greenland Ice Sheet  
860 motion forced by time-varying meltwater drainage: Implications for the relationship between subglacial drainage system  
861 behavior and ice velocity, *J. Geophys. Res. Earth Surf.*, 117(F3), F03002, <https://doi.org/10.1029/2011JF002220>, 2012.
- 862 Bell, R. E.: The role of subglacial water in ice-sheet mass balance, *Nature Geosci.*, 1(5), 297–304,  
863 <https://doi.org/10.1038/ngeo186>, 2008.
- 864 Benn, D. I., Thompson, S., Gulley, J., Mertes, J., Luckman, A. and Nicholson, L.: Structure and evolution of the drainage  
865 system of a Himalayan debris-covered glacier, and its relationship with patterns of mass loss, *TC*, 11, 2247–2264,  
866 <https://doi.org/10.5194/tc-11-2247-2017>, 2017.
- 867 Boulton, G. S., Lunn, R., Vidstrand, P. and Zatzepin, S.: Subglacial drainage by groundwater–channel coupling, and the origin  
868 of esker systems: part II—theory and simulation of a modern system, *Quat. Sci. Rev.*, 26(7–8), 1091–1105,  
869 <https://doi.org/10.1016/j.quascirev.2007.01.006>, 2007.
- 870 Catania, G. A. and Neumann, T. A.: Persistent englacial drainage features in the Greenland Ice Sheet, *Geophys. Res. Lett.*,  
871 37(2), L02501, <https://doi.org/10.1029/2009GL041108>, 2010.
- 872 Catania, G. A., Neumann, T. A. and Price, S. F.: Characterizing englacial drainage in the ablation zone of the Greenland Ice  
873 Sheet, *J. Glac.*, 54(187), 567–578, <https://doi.org/10.3189/002214308786570854>, 2008.
- 874 Chandler, D. M., Wadham, J. L., Lis, G. P., Cowton, T., Sole, A., Bartholomew, I., Telling, J., Nienow, P., Bagshaw, E. B.,  
875 Mair, D., Vinen, S. and Hubbard, A.: Evolution of the subglacial drainage system beneath the Greenland Ice Sheet revealed  
876 by tracers, *Nature Geosci.*, 6(3), 195–198, <https://doi.org/10.1038/ngeo1737>, 2013.
- 877 Christmann, J., Helm, V., Khan, S. A., Kleiner, T., Müller, R., Morlighem, M., Neckel, N., Rückamp, M., Steinhage, D.,  
878 Zeising, O., and Humbert, A.: Elastic deformation plays a non-negligible role in Greenland’s outlet glacier flow, *Commun*  
879 *Earth Environ.*, 2, 1–12, <https://doi.org/10.1038/s43247-021-00296-3>, 2021.  
880
- 881 Chu, V. W.: Greenland Ice Sheet hydrology: A review, *Prog. Phys. Geogr.*, 38(1), 19–54,  
882 <https://doi.org/10.1177/0309133313507075>, 2014.

- 883 Chudley, T. R., Christoffersen, P., Doyle, S. H., Bougamont, M., Schoonman, C. M., Hubbard, B. and James, M. R.:  
884 Supraglacial lake drainage at a fast-flowing Greenlandic outlet glacier, *PNAS*, 116(51), 25468–25477,  
885 <https://doi.org/10.1073/pnas.1913685116>, 2019.
- 886 Church, G., Grab, M., Schmelzbach, C., Bauder, A., and Maurer, H.: Monitoring the seasonal changes of an englacial conduit  
887 network using repeated ground-penetrating radar measurements, *TC*, 14, 3269–3286, <https://doi.org/10.5194/tc-14-3269-2020>,  
888 2020.
- 889 Clarke, G. K. C.: Lumped-element analysis of subglacial hydraulic circuits, *J. Geophys. Res.*, 101(B8), 17547–17559,  
890 <https://doi.org/10.1029/96JB01508>, 1996.
- 891 Clarke, G. K. C.: Hydraulics of subglacial outburst floods: new insights from the Spring–Hutter formulation, *J. Glac.*, 49(165),  
892 299–313, <https://doi.org/10.3189/172756503781830728>, 2003.
- 893 Colgan, W. and Steffen, K.: Modelling the spatial distribution of moulins near Jakobshavn, Greenland, *IOP Conf. Ser.: Earth  
894 Environ. Sci.*, 6(1), 012022, <https://doi.org/10.1088/1755-1307/6/1/012022>, 2009.
- 895 Colgan, W., Rajaram, H., Anderson, R., Steffen, K., Phillips, T., Joughin, I., Zwally, H. J. and Abdalati, W.: The annual  
896 glaciohydrology cycle in the ablation zone of the Greenland ice sheet: Part 1. Hydrology model, *J. Glac.*, 57(204), 697–709,  
897 <https://doi.org/10.3189/002214311797409668>, 2011.
- 898 Covington, M. D., Banwell, A. F., Gulley, J., Saar, M. O., Willis, I. and Wicks, C. M.: Quantifying the effects of glacier  
899 conduit geometry and recharge on proglacial hydrograph form, *J. Hydro.*, 414–415, 59–71,  
900 <https://doi.org/10.1016/j.jhydrol.2011.10.027>, 2012.
- 901 Covington, M. D., Gulley, J. D., Trunz, C., Mejia, J. and Gadd, W.: Moulin Volumes Regulate Subglacial Water Pressure on  
902 the Greenland Ice Sheet, *Geophys. Res. Lett.*, 47(20), e2020GL088901, <https://doi.org/10.1029/2020GL088901>, 2020.
- 903 Cowton, T., Nienow, P., Sole, A., Wadham, J., Lis, G., Bartholomew, I., Mair, D. and Chandler, D.: Evolution of drainage  
904 system morphology at a land-terminating Greenlandic outlet glacier, *J. Geophys. Res. Earth Surf.*, 118(1), 29–41,  
905 <https://doi.org/10.1029/2012JF002540>, 2013.
- 906 Cowton, T., Nienow, P., Bartholomew, I. and Mair, D.: Variability in ice motion at a land-terminating Greenlandic outlet  
907 glacier: the role of channelized and distributed drainage systems, *J. Glac.*, 62(233), 451–466,  
908 <https://doi.org/10.1017/jog.2016.36>, 2016.
- 909 Cuffey, K. M. and Paterson, W. S. B.: *The Physics of Glaciers*, Elsevier., 2010.
- 910 Das, S. B., Joughin, I., Behn, M. D., Howat, I. M., King, M. A., Lizarralde, D. and Bhatia, M. P.: Fracture Propagation to the  
911 base of the Greenland Ice Sheet during supraglacial lake drainage, *Science*, 320(5877), 778–781,  
912 <https://doi.org/10.1126/science.1153360>, 2008.
- 913 Dow, C. F., Kulessa, B., Rutt, I. C., Tsai, V. C., Pimentel, S., Doyle, S. H., van As, D., Lindbäck, K., Pettersson, R., Jones, G.  
914 A. and Hubbard, A.: Modeling of subglacial hydrological development following rapid supraglacial lake drainage, *J. Geophys.  
915 Res. Earth Surf.*, 120(6), 2014JF003333, <https://doi.org/10.1002/2014JF003333>, 2015.
- 916 Dow, C. F., Werder, M. A., and Walker, R. T.: Modeling Antarctic subglacial lake filling and drainage cycles, 10, 1381–1393,  
917 <http://dx.doi.org/10.5194/tc-10-1381-2016>, 2016.  
918
- 919 Downs, J. Z., Johnson, J. V., Harper, J. T., Meierbachtol, T., and Werder, M. A.: Dynamic hydraulic conductivity reconciles  
920 mismatch between modeled and observed winter subglacial water pressure, *J. Geophys. Res. Earth Surf.*, 123(4), 818–836,  
921 <https://doi.org/10.1002/2017JF004522>, 2018.

- 922 Flowers, G. E.: Subglacial modulation of the hydrograph from glacierized basins, *Hydrol. Process.*, 22(19), 3903–3918,  
923 <https://doi.org/10.1002/hyp.7095>, 2008.
- 924 Flowers, G. E.: Modelling water flow under glaciers and ice sheets, *Proc. Math. Phys.* 471, 20140907–20140907,  
925 <https://doi.org/10.1098/rspa.2014.0907>, 2015.
- 926 Flowers, G. E.: Hydrology and the future of the Greenland Ice Sheet, *Nature Comm.*, 9(1), 2729,  
927 <https://doi.org/10.1038/s41467-018-05002-0>, 2018.
- 928 Flowers, G. E. and Clarke, G. K. C.: A multicomponent coupled model of glacier hydrology 1. Theory and synthetic examples,  
929 *J. Geophys. Res.*, 107(B11), 2287, <https://doi.org/10.1029/2001JB001122>, 2002.
- 930 Forster, R. R., Box, J. E., van den Broeke, M. R., Miège, C., Burgess, E. W., van Angelen, J. H., Lenaerts, J. T. M., Koenig,  
931 L. S., Paden, J., Lewis, C., Gogineni, S. P., Leuschen, C. and McConnell, J. R.: Extensive liquid meltwater storage in firn  
932 within the Greenland ice sheet, *Nature Geosci.*, 7(2), 95–98, <https://doi.org/10.1038/ngeo2043>, 2014.
- 933 Fountain, A. G. and Walder, J. S.: Water flow through temperate glaciers, *Rev. Geophys.*, 36(3), 299–328,  
934 <https://doi.org/10.1029/97RG03579>, 1998.
- 935 Gajek, W., Gräff, D., Hellmann, S., Rempel, A. W., and Walter, F.: Diurnal expansion and contraction of englacial fracture  
936 networks revealed by seismic shear wave splitting, *Comm. Earth. Environ.*, 2, 1–8, [https://doi.org/10.1038/s43247-021-00279-](https://doi.org/10.1038/s43247-021-00279-4)  
937 4, 2021.
- 938 Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G.,  
939 Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim,  
940 G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S.  
941 D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2  
942 (MERRA-2), *J. Climate*, 30, 5419–5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- 943 Germain, S. L. S. and Moorman, B. J.: Long-term observations of supraglacial streams on an Arctic glacier, *J. Glac.*, 65(254),  
944 900–911, <https://doi.org/10.1017/jog.2019.60>, 2019.
- 945 Goodman, R. E.: *Introduction to Rock Mechanics*, 2nd ed., Wiley, New York., 1989.
- 946 Gulley, J. D., Benn, D. I., Screaton, E. and Martin, J.: Mechanisms of englacial conduit formation and their implications for  
947 subglacial recharge, *Quat. Sci. Rev.*, 28(19–20), 1984–1999, <https://doi.org/10.1016/j.quascirev.2009.04.002>, 2009.
- 948 Gulley, J. D., Spellman, P. D., Covington, M. D., Martin, J. B., Benn, D. I. and Catania, G.: Large values of hydraulic roughness  
949 in subglacial conduits during conduit enlargement: implications for modeling conduit evolution, *Earth Surf. Process.*  
950 *Landforms*, 39(3), 296–310, <https://doi.org/10.1002/esp.3447>, 2014.
- 951 Hewitt, I. J.: Seasonal changes in ice sheet motion due to melt water lubrication, *EPSL*, 371–372, 16–25,  
952 <https://doi.org/10.1016/j.epsl.2013.04.022>, 2013.
- 953 Hoffman, M. J. and Price, S.: Feedbacks between coupled subglacial hydrology and glacier dynamics, *J. Geophys. Res. Earth*  
954 *Surf.*, 119(3), 414–436, <https://doi.org/10.1002/2013JF002943>, 2014.
- 955 Hoffman, M. J., Catania, G. A., Neumann, T. A., Andrews, L. C. and Rumrill, J. A.: Links between acceleration, melting, and  
956 supraglacial lake drainage of the western Greenland Ice Sheet, *J. Geophys. Res. Earth Surf.*, 116(F4), F04035,  
957 <https://doi.org/10.1029/2010JF001934>, 2011.

- 958 Hoffman, M. J., Andrews, L. C., Price, S. A., Catania, G. A., Neumann, T. A., Lüthi, M. P., Gulley, J., Ryser, C., Hawley, R.  
959 L. and Morriss, B.: Greenland subglacial drainage evolution regulated by weakly connected regions of the bed, *Nature Comm.*,  
960 7, 13903, <https://doi.org/10.1038/ncomms13903>, 2016.
- 961 Holmlund, P.: Internal geometry and evolution of moulins, *J. Glac.*, 34(117), 242–248, 1988.
- 962 Iken, A.: Measurements of water pressure in moulins as part of a movement study of the White Glacier, Axel Heiberg Island,  
963 Northwest Territories, Canada, *J. Glac.*, 11(61), 53–58, 1972.
- 964 Iken, A. and Bindschadler, R.: Combined measurements of subglacial water pressure and surface velocity of Findelengletscher,  
965 Switzerland: conclusions about drainage system and sliding mechanism, *J. Glac.*, 32(110), 101–119,  
966 <https://doi.org/10.3189/S0022143000006936>, 1986.
- 967 Iken, A., Echelmeyer, K., Harrison, W. and Funk, M.: Mechanisms of fast flow in Jakobshavns Isbræ, West Greenland: Part  
968 I. Measurements of temperature and water level in deep boreholes, *J. Glac.*, 39(131), 15–25,  
969 <https://doi.org/10.1017/S0022143000015689>, 1993.
- 970 Jarosch, A. H. and Gudmundsson, M. T.: A numerical model for meltwater channel evolution in glaciers, *TC*, 6(2), 493–503,  
971 <https://doi.org/10.5194/tc-6-493-2012>, 2012.
- 972 Kirsch, G.: *Die Theorie der Elastizität und die Bedürfnisse der Festigkeitslehre*, Springer, Berlin., 1898.
- 973 Krawczynski, M. J., Behn, M. D., Das, S. B., and Joughin, I.: Constraints on the lake volume required for hydro-fracture  
974 through ice sheets, *Geophys. Res. Lett.*, 36, L10501, <https://doi.org/10.1029/2008GL036765>, 2009.
- 975 Lai, C.-Y., Stevens, L. A., Chase, D. L., Creyts, T. T., Behn, M. D., Das, S. B., and Stone, H. A.: Hydraulic transmissivity  
976 inferred from ice-sheet relaxation following Greenland supraglacial lake drainages, *Nat Commun*, 12, 3955,  
977 <https://doi.org/10.1038/s41467-021-24186-6>, 2021.
- 978
- 979 Lüthi, M. P., Ryser, C., Andrews, L. C., Catania, G. A., Funk, M., Hawley, R. L., Hoffman, M. J. and Neumann, T. A.: Heat  
980 sources within the Greenland Ice Sheet: dissipation, temperate paleo-firn and cryo-hydrologic warming, *TC*, 9(1), 245–253,  
981 <https://doi.org/10.5194/tc-9-245-2015>, 2015.
- 982 MacFerrin, M., Machguth, H., As, D. van, Charalampidis, C., Stevens, C. M., Heilig, A., Vandecrux, B., Langen, P. L.,  
983 Mottram, R., Fettweis, X., Broeke, M. R. van den, Pfeffer, W. T., Moussavi, M. S. and Abdalati, W.: Rapid expansion of  
984 Greenland's low-permeability ice slabs, *Nature*, 573(7774), 403–407, <https://doi.org/10.1038/s41586-019-1550-3>, 2019.
- 985 MacGregor, J. A., Fahnestock, M. A., Colgan, W. T., Larsen, N. K., Kjeldsen, K. K. and Welker, J. M.: The age of surface-  
986 exposed ice along the northern margin of the Greenland Ice Sheet, *J. Glac.*, 66(258), 667–684,  
987 <https://doi.org/10.1017/jog.2020.62>, 2020.
- 988 Mair, D., Nienow, P., Willis, I. and Sharp, M.: Spatial patterns of glacier motion during a high-velocity event: Haut Glacier  
989 d'Arolla, Switzerland, *J. Glac.*, 47(156), 9–20, <https://doi.org/10.3189/172756501781832412>, 2001.
- 990 Mair, D., Nienow, P. W., Sharp, M. J., Wohlleben, T. and Willis, I.: Influence of subglacial drainage system evolution on  
991 glacier surface motion: Haut Glacier d'Arolla, Switzerland, *J. of Geophys Res. Solid Earth*, 107(B8),  
992 <https://doi.org/10.1029/2001JB000514>, 2002.
- 993 Mankoff, K. D., Gulley, J. D., Tulaczyk, S. M., Covington, M. D., Liu, X., Chen, Y., Benn, D. I. and Głowacki, P. S.:  
994 Roughness of a subglacial conduit under Hansbreen, Svalbard, *J. Glac.*, 1–13, <https://doi.org/10.1017/jog.2016.134>, 2017.

- 995 McGrath, D., Colgan, W., Steffen, K., Lauffenburger, P. and Balog, J.: Assessing the summer water budget of a moulin basin  
996 in the Sermeq Avannarleq ablation region, Greenland ice sheet, *J. Glac.*, 57(205), 954–964,  
997 <https://doi.org/10.3189/002214311798043735>, 2011.
- 998 McQuillan, M. and Karlstrom, L.: Fluid resonance in elastic-walled englacial transport networks, *J. Glac.*, 1–14,  
999 <https://doi.org/10.1017/jog.2021.48>, 2021.
- 1000 Meierbachtol, T. W., Harper, J. and Humphrey, N.: Basal Drainage System Response to Increasing Surface Melt on the  
1001 Greenland Ice Sheet, *Science*, 341(6147), 777–779, <https://doi.org/10.1126/science.1235905>, 2013.
- 1002 Mejia, J. Z., Gulley, J. D., Trunz, C., Covington, M. D., Bartholomaeus, T. C., Xie, S., and Dixon, T. H.: Isolated Cavities  
1003 Dominate Greenland Ice Sheet Dynamic Response to Lake Drainage, *Geophys. Res. Lett.*, 48, e2021GL094762,  
1004 <https://doi.org/10.1029/2021GL094762>, 2021.
- 1005 Miège, C., Forster, R. R., Brucker, L., Koenig, L. S., Solomon, D. K., Paden, J. D., Box, J. E., Burgess, E. W., Miller, J. Z.,  
1006 McNerney, L., Brautigam, N., Fausto, R. S. and Gogineni, S.: Spatial extent and temporal variability of Greenland firn aquifers  
1007 detected by ground and airborne radars, *J. Geophys. Res. Earth Surf.*, 121(12), 2016JF003869,  
1008 <https://doi.org/10.1002/2016JF003869>, 2016.
- 1009 Moon, T., Joughin, I., Smith, B., van den Broeke, M. R., van de Berg, W. J., Noël, B. and Usher, M.: Distinct patterns of  
1010 seasonal Greenland glacier velocity, *Geophys. Res. Lett.*, 41(20), 2014GL061836, <https://doi.org/10.1002/2014GL061836>,  
1011 2014.
- 1012 Moreau, L.: L'exploration du cryokarst glaciaire et son intérêt scientifique pour l'étude du drainage des eaux de fonte : porches,  
1013 cavités, crevasses, bédrières et moulins., *Collection EDYTEM. Cahiers de géographie*, 8, 163–170, 2009.
- 1014 Müller, F. and Iken, A.: Velocity fluctuations and water regime of Arctic valley glaciers, *IAHS*, 95, 165–182, 1973.
- 1015 Naruse, R., Okuhira, F., Ohmae, H., Kawada, K. and Nakawo, M.: Closure Rate of a 700 m Deep Bore Hole at Mizuho Station,  
1016 East Antarctica, *Ann. Glac.*, 11, 100–103, <https://doi.org/10.3189/S0260305500006406>, 1988.
- 1017 Nossokoff, A.: Using Small Scale Physical Experiments to Improve Enthalpy Based Models of Ice Sheets, M.S., University  
1018 of Colorado, Boulder, 2013.
- 1019 Paterson, W. S. B.: Secondary and tertiary creep of glacier ice as measured by borehole closure rates, *Rev. Geophys.*, 15(1),  
1020 47–55, <https://doi.org/10.1029/RG015i001p00047>, 1977.
- 1021 Pitcher, L. H. and Smith, L. C.: Supraglacial Streams and Rivers, *Ann. Rev. Earth Planetary Sci.*, 47(1), 421–452,  
1022 <https://doi.org/10.1146/annurev-earth-053018-060212>, 2019.
- 1023 Poinar, K. and Andrews, L. C.: Challenges in predicting Greenland supraglacial lake drainages at the regional scale, *TC*, 15,  
1024 1455–1483, <https://doi.org/10.5194/tc-15-1455-2021>, 2021.
- 1025 Poinar, K., Joughin, I., Das, S. B., Behn, M. D., Lenaerts, J. T. M. and van den Broeke, M. R.: Limits to future expansion of  
1026 surface-melt-enhanced ice flow into the interior of western Greenland, *Geophys. Res. Lett.*, 42(6), 2015GL063192,  
1027 <https://doi.org/10.1002/2015GL063192>, 2015.
- 1028 Poinar, K., Joughin, I., Lenaerts, J. T. M. and Broeke, M. R. V. D.: Englacial latent-heat transfer has limited influence on  
1029 seaward ice flux in western Greenland, *J. Glac.*, 1–16, <https://doi.org/10.1017/jog.2016.103>, 2016.
- 1030 Poinar, K., Joughin, I., Lilien, D., Brucker, L., Kehrl, L. and Nowicki, S.: Drainage of Southeast Greenland Firn Aquifer Water  
1031 through Crevasses to the Bed, *Front. Earth Sci.*, 5, <https://doi.org/10.3389/feart.2017.00005>, 2017.




- 1032 Priest, S. D.: *Discontinuity Analysis for Rock Engineering*, 1st ed., Chapman & Hall, London; New York., 1993.
- 1033 Reeh, N., Christensen, E. L., Mayer, C., and Olesen, O. B.: Tidal bending of glaciers: a linear viscoelastic approach, 37, 83–  
1034 89, <https://doi.org/10.3189/172756403781815663>, 2003.
- 1035
- 1036 Rösli, C., Walter, F., Ampuero, J.-P., and Kissling, E.: Seismic moulin tremor, *J. Geophys. Res. Solid Earth* 121, 5838–5858,  
1037 <https://doi.org/10.1002/2015JB012786>, 2016.
- 1038 Ryser, C., Lüthi, M. P., Andrews, L. C., Hoffman, M. J., Catania, G. A., Hawley, R. L., Neumann, T. A. and Kristensen, S. S.:  
1039 Sustained high basal motion of the Greenland ice sheet revealed by borehole deformation, *J. Glac.*, 60(222), 647–660,  
1040 <https://doi.org/10.3189/2014JoG13J196>, 2014.
- 1041 Scheingross, J. S. and Lamb, M. P.: A Mechanistic Model of Waterfall Plunge Pool Erosion into Bedrock, *J. Geophys. Res.*  
1042 *Earth Surf.*, 122(11), 2079–2104, <https://doi.org/10.1002/2017JF004195>, 2017.
- 1043 Schoof, C.: Ice-sheet acceleration driven by melt supply variability, *Nature*, 468(7325), 803–806,  
1044 <https://doi.org/10.1038/nature09618>, 2010.
- 1045 Selmes, N., Murray, T. and James, T. D.: Fast draining lakes on the Greenland Ice Sheet, *Geophys. Res. Lett.*, 38(15), L15501,  
1046 <https://doi.org/10.1029/2011GL047872>, 2011.
- 1047 Shreve, R. L.: Movement of water in glaciers, *J. Glac.*, 11(62), 205–214, <https://doi.org/10.1017/S002214300002219X>, 1972.
- 1048 Smith, L. C., Chu, V. W., Yang, K., Gleason, C. J., Pitcher, L. H., Rennermalm, A. K., Legleiter, C. J., Behar, A. E., Overstreet,  
1049 B. T., Moustafa, S. E., Tedesco, M., Forster, R. R., LeWinter, A. L., Finnegan, D. C., Sheng, Y. and Balog, J.: Efficient  
1050 meltwater drainage through supraglacial streams and rivers on the southwest Greenland ice sheet, *PNAS*, 112(4), 1001–1006,  
1051 <https://doi.org/10.1073/pnas.1413024112>, 2015.
- 1052 Smith, L. C., Yang, K., Pitcher, L. H., Overstreet, B. T., Chu, V. W., Rennermalm, Å. K., Ryan, J. C., Cooper, M. G., Gleason,  
1053 C. J., Tedesco, M., Jeyaratnam, J., As, D. van, Broeke, M. R. van den, Berg, W. J. van de, Noël, B., Langen, P. L., Cullather,  
1054 R. I., Zhao, B., Willis, M. J., Hubbard, A., Box, J. E., Jenner, B. A. and Behar, A. E.: Direct measurements of meltwater runoff  
1055 on the Greenland ice sheet surface, *PNAS*, 114(50), E10622–E10631, <https://doi.org/10.1073/pnas.1707743114>, 2017.
- 1056 Smith, L. C., Andrews, L. C., Pitcher, L. H., Overstreet, B. T., Rennermalm, Å. K., Cooper, M. G., Cooley, S. W., Ryan, J. C.,  
1057 Miège, C., Kershner, C., and Simpson, C. E.: Supraglacial River Forcing of Subglacial Water Storage and Diurnal Ice Sheet  
1058 Motion, 48, e2020GL091418, <https://doi.org/10.1029/2020GL091418>, 2021.
- 1059
- 1060 Spring, U. and Hutter, K.: Numerical studies of Jökulhlaups, *Cold Reg Sci Technol.*, 4(3), 227–244,  
1061 [https://doi.org/10.1016/0165-232X\(81\)90006-9](https://doi.org/10.1016/0165-232X(81)90006-9), 1981.
- 1062 Spring, U. and Hutter, K.: Conduit flow of a fluid through its solid phase and its application to intraglacial channel flow, *Int.*  
1063 *J. Eng. Sci.*, 20(2), 327–363, [https://doi.org/10.1016/0020-7225\(82\)90029-5](https://doi.org/10.1016/0020-7225(82)90029-5), 1982.
- 1064 Tedstone, A. J., Nienow, P. W., Gourmelen, N., Dehecq, A., Goldberg, D. and Hanna, E.: Decadal slowdown of a land-  
1065 terminating sector of the Greenland Ice Sheet despite warming, *Nature*, 526(7575), 692–695,  
1066 <https://doi.org/10.1038/nature15722>, 2015.
- 1067 Trunz, C.: *Modeling and Measuring Water Level Fluctuations in the Greenland Ice Sheet: How Moulin Life Cycle and Shape*  
1068 *can Inform us on the Subglacial Drainage System*, Ph.D., University of Arkansas, Fayetteville, Arkansas, 156 pp., 2021.
- 1069 Turcotte, D. L. and Schubert, G.: *Geodynamics*, Cambridge University Press., 2002.

- 1070 Vaňková, I., Voytenko, D., Nicholls, K. W., Xie, S., Parizek, B. R., and Holland, D. M.: Vertical Structure of Diurnal Englacial  
1071 Hydrology Cycle at Helheim Glacier, East Greenland, *Geophys. Res. Lett.*, 45, 8352–8362,  
1072 <https://doi.org/10.1029/2018GL077869>, 2018.
- 1073 Vaughan, D. G.: Tidal flexure at ice shelf margins, *J. Geophys. Res. Solid Earth*, 100(B4), 6213–6224,  
1074 <https://doi.org/10.1029/94JB02467>, 1995.
- 1075 Weertman, J.: Theory of water-filled crevasses in glaciers applied to vertical magma transport beneath oceanic ridges, *J.*  
1076 *Geophys. Res.*, 76, 1171–1183, <https://doi.org/10.1029/JB076i005p01171>, 1971.
- 1077 Weertman, J.: Can a water-filled crevasse reach the bottom surface of a glacier?, *IASH*, 95, 139–145, 1973.
- 1078 Weertman, J.: *Dislocation Based Fracture Mechanics*, World Scientific Publishing Company, Singapore, 1996.
- 1079 Werder, M. A.: The hydrology of subglacial overdeepenings: A new supercooling threshold formula, *Geophys. Res. Lett.*, 43,  
1080 2015GL067542, <https://doi.org/10.1002/2015GL067542>, 2016.
- 1081 Werder, M. A., Schuler, T. V. and Funk, M.: Short term variations of tracer transit speed on alpine glaciers, *TC*, 4(3), 381–  
1082 396, 2010.
- 1083 Werder, M. A., Hewitt, I. J., Schoof, C. G. and Flowers, G. E.: Modeling channelized and distributed subglacial drainage in  
1084 two dimensions, *J. Geophys. Res. Earth Surf.*, 118(4), 2140–2158, <https://doi.org/10.1002/jgrf.20146>, 2013.
- 1085 Williams, J. J., Gourmelen, N. and Nienow, P.: Dynamic response of the Greenland ice sheet to recent cooling, *Sci. Rep.*,  
1086 10(1), 1647, <https://doi.org/10.1038/s41598-020-58355-2>, 2020.
- 1087 Wright, P. J., Harper, J. T., Humphrey, N. F. and Meierbachtol, T. W.: Measured basal water pressure variability of the western  
1088 Greenland Ice Sheet: Implications for hydraulic potential, *J. Geophys. Res. Earth Surf.*, 121(6), 2016JF003819,  
1089 <https://doi.org/10.1002/2016JF003819>, 2016.
- 1090 Yang, K. and Smith, L. C.: Internally drained catchments dominate supraglacial hydrology of the southwest Greenland Ice  
1091 Sheet, *J. Geophys. Res. Earth Surf.*, 2016JF003927, <https://doi.org/10.1002/2016JF003927>, 2016.
- 1092 Yang, K., Smith, L. C., Chu, V. W., Pitcher, L. H., Gleason, C. J., Rennermalm, A. K. and Li, M.: Fluvial morphometry of  
1093 supraglacial river networks on the southwest Greenland Ice Sheet, *GISci. Remote Sens.*, 53(4),  
1094 <https://doi.org/10.1080/15481603.2016.1162345>, 2016.

1095

1096

1097



1099

1100 **Figure 1. Processes included in the MouSh model.** Black lines show a base moulin geometry that each process acts on, and colored lines  
 1101 show the change in moulin geometry (not to scale) due to that process alone. From left to right: changes in moulin geometry due to viscous  
 1102 deformation; elastic deformation; melting by turbulent energy dissipation of flowing water inside the moulin; melting by open-channel water  
 1103 flow along bare ice; refreezing over winter inside the moulin; and deformation due to ice motion prescribed by Glen’s flow law. Unlike the  
 1104 other components, elastic deformation is instantaneous, but applied over the model timestep (Sect. 2.2.1; Supplement S2). The right-most  
 1105 moulin shows the moulin geometry before (dashed black lines) and after (solid black lines and blue water) several hypothetical model  
 1106 timesteps, i.e., the sum of all processes shown in the preceding panels. Changes are not to scale.

1107

1108

1109

1110

1111

1112


1113

1114

1115

1116

1117



1119

1120 **Figure 2. MouSh geometry and surface expression of a moulin and its reflection in the MouSh model.** (a) Schematic of MouSh  
 1121 geometry and inputs. Inflow and outflow of the system are indicated by  $Q_{in}$ ,  $Q_{out}$ , and  $Q_{base}$ . Time evolving moulin and subglacial parameters  
 1122 include moulin radii ( $r_1$ ,  $r_2$ ), moulin water level ( $h_w$ ), and subglacial cross-sectional area ( $S$ ).  $r_1$  and  $r_2$  are evolved by  $dr_{oc}$ ,  $dr_v$ ,  $dr_e$ ,  $dr_f$ , and  
 1123  $dr_t$  (open channel melting, viscous deformation, elastic deformation, refreezing, and turbulent melting, respectively; colored as in Fig. 1).  $u_d$   
 1124 shears the moulin as prescribed by Glen's Flow Law. Ice thickness and subglacial path length are indicated by  $H_i$  and  $L$ , respectively. Ice  
 1125 flow is from left to right. Further details are in Sect. 2. Modified from Trunz (2021). (b) WorldView-2 scene from July 2010 of an  
 1126 approximately  $1.2 \times 0.8$  km region surrounding the example moulin (yellow) formed by a drained supraglacial lake. (c) Detail of panel b,  
 1127 with the inflow stream and moulin indicated. (d) Detail of panel c, showing the moulin minor radius  $r_1$ , major radius  $r_2$ , and water input  $Q_{in}$   
 1128 from the inflow stream, as represented in the MouSh model. Maps generated by authors. WorldView image © 2010 DigitalGlobe, Inc.

1129

1130

1131


1132

1133

1134


1135

1136



1138

1139 **Figure 3. Results of parameter sensitivity studies for 40-day MouSh model runs.** Shown are the sensitivity of moulin size to initial  
 1140 condition for moulin radius (a–b), enhancement factor for englacial ice (c–d), ice temperature scenario (e–f), Young's modulus (g–h), softness  
 1141 of basal ice (i–j), friction factor for water flow beneath the water line (k–l), friction factor for water flow above the water line (m–n), and ice  
 1142 thickness (o–p). The left column shows the moulin radii (black and grey) at the mean water level and the mean subglacial channel radius  
 1143 (purple) averaged over the final 24-hour period of the 40-day model run. The right column shows the equilibrium water level (blue), moulin  
 1144 capacity (red), and volume of water in the moulin (green) averaged over the same 24-hour period. Overall, moulin radius is most sensitive  
 1145 to the friction factors, while moulin water level and volume are most sensitive to ice thickness  $H_i$  (also an indicator of the hydraulic potential  
 1146 gradient) and basal ice softness  $A$ . Note the different y-axis range in panels (c) and (e).




1148

1149 **Figure 4. Diurnal variations in moulin sizes in 40-day parameter sensitivity runs.** Shown are the sensitivity of diurnal variation in  
 1150 moulin size and water storage metrics to initial condition for moulin radius (a–b), enhancement factor for englacial ice (c–d), ice temperature  
 1151 scenario from coldest to warmest ice (e–f), Young's modulus (g–h), softness of basal ice (i–j), friction factor for water flow beneath the water  
 1152 line (k–l), friction factor for water flow above the water line (m–n), and ice thickness (o–p). The left column shows diurnal variations in  
 1153 moulin radii (black and grey) at the equilibrium water level and the subglacial channel radius (purple) in the final 24-hour period of the 40-  
 1154 day model run. The right column shows the diurnal variation in water level (blue), moulin volume (red), and volume of water in the moulin  
 1155 (green) within the same 24-hour period. Note the right y-axis difference in panel (p).

1156


1157



1159

1160 **Figure 5. Contributions of viscous deformation, elastic deformation, and phase changes to moulin geometry.** (a–e) Equilibrium  
 1161 geometries of five moulins in ice of different ice thicknesses  $H$  and different distances from the terminus (same as Fig. 6o–p) averaged over  
 1162 the final 24-hour period of a 40-day model run. (f) Vertical variation of viscous deformation (blue), elastic deformation (red), and phase  
 1163 change (green) contributions to moulin geometry averaged over the same 24-hour period. Negative values indicate contributions to moulin  
 1164 closure; positive values open the moulin. Darkening shades of each color map to moulins of increasing ice thickness. Closure and opening  
 1165 rates are greatest at the minimum daily water level (which is inferable by the lower notch in the moulin wall). (g) Time series of the  
 1166 components shown in panel f (colors the same) at the mean water level over the entire 40-day model run. The greater diurnal range in water  
 1167 level in moulins in thick ice drives the observed larger diurnal variations in viscous deformation. (h) For reference, moulin water level as  
 1168 fraction of overburden for different ice thicknesses  $H_i$ . Lighter greys indicate thinner ice; blue dashed line indicates where fraction of  
 1169 overburden = 1.

1170



1172

1173 **Figure 6. MouSh model runs with realistic supraglacial and ice conditions.** The model runs are for a low-elevation Basin 1 (553 m ice  
 1174 thickness; black lines), mid-elevation Basin 2 (741 m ice thickness; purple lines), and high-elevation Basin 3 (1315 m ice thickness; grey  
 1175 lines). (a) Supraglacial discharge into the moulin  $Q_{in}$  and prescribed base flow  $Q_{base}$ . (b) Moulin water level as a fraction of overburden. Note  
 1176 that the highest elevation moulin exceeds the ice surface most days. (c) Moulin capacity, or the total moulin volume. (d) Subglacial channel  
 1177 cross-sectional area. Colored vertical lines indicate times in Fig. 7. Note x-axes start on day 5.

1178


1179

1180

1181

1182





1184

1185 **Figure 7. Evolution of moulin geometry over the melt season.** Colored boxes correspond to the times indicated with colored vertical lines

1186 in Fig. 6. (a–f) Basin 1 with ice thickness of 553 m. (g–l) Basin 2 with ice thickness of 741 m. (m–r) Basin 3 with ice thickness of 1315 m.

1187 Axes are not to scale.

1188

1189

1190


1191

1192

1193

1194

1195



1197

1198 **Figure 8. Time series of viscous, elastic and phase change components of moulin evolution and their relative importance in**  
 1199 **determining moulin geometry.** (a) Time varying viscous (blues), elastic (reds), and phase change (melting, greens) components of moulin  
 1200 geometry. (b) The daily ratio of the total amount of phase change (melting above and below the water line) to total deformation (elastic plus  
 1201 viscous; purples). Values above 1 indicate that melting dominates; values below 1 indicate that deformation dominates. Data is smoothed  
 1202 over 24 h. For both panels, light colors are for Basin 1 ( $H_i=553$  m), medium colors for Basin 2 ( $H_i=741$  m), and dark colors for Basin 3  
 1203 ( $H_i=1315$  m). Note x-axes start on day 5.

1204

1205

1206

1207

1208

1209

1210

1211


1212

1213

1214

1215

1216 FIGURE 9



1217

1218 **Figure 9. Daily percentage change in moulin variables relative to the daily mean value.** (a) Daily percentage change in moulin water  
1219 level relative to the daily mean water level for Basins 1, 2, and 3 (black, purple, and grey lines, respectively). (b) Daily percentage change  
1220 in moulin capacity relative to the daily mean moulin capacity. (c) Daily percentage change in the subglacial channel cross-sectional area  
1221 relative to the daily mean value. For (b–c), colors are as in (a).

1222

1223


1224

1225

1226

1227

1228



1231

1232 **Figure 10. Difference between variable and fixed moulin geometries for Basin 1 and 2** (ice thicknesses of 553 m and 741 m,  
 1233 respectively). The fixed moulins are cylinders with a fixed radius of 1.6 and 1.4 m for Basin 1 and 2 respectively, which are the time-mean  
 1234 radii at the mean water level for the variable moulins. In all instances, the difference is calculated as  $(cylindrical - variable)$  with instances  
 1235 of percentage difference calculated as  $(cylindrical - variable) / (variable)$ . (a, b) Difference in moulin water level for Basin 1 and Basin 2  
 1236 respectively, plotted every hour. Negative values indicate periods where the variable moulin water levels are higher than those of the fixed  
 1237 cylindrical moulin. (c, d) Percentage difference in moulin capacity plotted every 2 hours for clarity. When values are negative, the variable  
 1238 moulin is larger than the fixed cylindrical moulin.

1239

1240

1241

1242

1243 **Table 1. MouSh model constants and parameter ranges.** During realistic runs (Sect. 2.4) Values used during realistic model  
 1244 experiments are generally the median value of the sensitivity experiment.

Constant	Description	Value	Units	
$\rho_i$	Ice density	910	kg m <sup>-3</sup>	
$\rho_w$	Water density	1000	kg m <sup>-3</sup>	
$\nu$	Poisson's ratio	0.3	-	
$C_p$	Heat capacity (ice)	2115	J (K kg) <sup>-1</sup>	
$C_w$	Heat capacity (liquid water)	4210	J (K kg) <sup>-1</sup>	
$g$	Gravitational acceleration	9.81	m s <sup>-2</sup>	
$K_i$	Thermal conductivity (ice)	2.1	J (m K s) <sup>-1</sup>	
$K_w$	Thermal conductivity (liquid water)	0.555	J (m K s) <sup>-1</sup>	
$L_f$	Latent heat of fusion	335000	J kg <sup>-1</sup>	
Parameter	Description	Realistic run value	Range	Units
A	A	<i>T<sub>i</sub> &amp; F* dependent</i>		Pa <sup>-3</sup> s <sup>-1</sup>
	$A_{sub}$	6 x 10 <sup>-24</sup>	5 x 10 <sup>-25</sup> to 5 x 10 <sup>-23</sup>	
E	Young's modulus	5	1 to 9	GPa
F *	Ice deformation enhancement factor	5	1 to 9	-
f	$f_{oc}$	0.1	0.01 to 1	-
	$f_m$	0.8	0.01 to 1	
$H_i$	Ice thickness*	553, 741, 1315	669 to 1569	m
n	Glen's Flow Law exponent	3	-	-
$R_0$	Initial moulin radius	2	0.5 to 5	m
$T_i(z)$	Ice temperature	-6 (FOXX profile)	-23 to 0 **	°C

\* $H_i$  defines distance from terminus  $L$  and surface slope  $\alpha$  based on a perfectly plastic ice surface profile

\*\*including Iken et al. (1993), Lüthi et al. (2015), and Ryser et al. (2014)

1246

1247

1248 **Table 2. General ice and moulin input parameters for realistic runs**

<b>Parameter</b>	<b>Basin 1</b>	<b>Basin 2</b>	<b>Basin 3</b>
Ice thickness (m)	553	741	1315
Distance from terminus (km)	13.6	24.5	77.1
Catchment size (km <sup>2</sup> )	19.8	18.4	55.5
Moulin input, mean diurnal range (m·s <sup>-3</sup> )	11.5	6.7	2.5
Moulin input, maximum value (m·s <sup>-3</sup> )	19.3	12.8	3.8
Baseflow, mean value (m·s <sup>-3</sup> )	20.2	17.7	6.2

1249