

Multi-decadal (1953 – 2017) rock glacier kinematics analysed by high-resolution topographic data in the Upper Kauner Valley, Austria

Fabian Fleischer¹, Florian Haas¹, Livia Piermattei², Madlene Pfeiffer³, Tobias Heckmann¹, Moritz Altmann¹, Jakob Rom¹, Manuel Stark¹, Michael H. Wimmer⁴, Norbert Pfeifer⁴ and Michael Becht¹

- 5
¹ Chair of Physical Geography, Catholic University of Eichstätt-Ingolstadt, 95072 Eichstätt, Germany
² Department of Geosciences, University of Oslo, 0316 Oslo, Norway
³ Institute of Geography, University of Bremen, 28359 Bremen, Germany
⁴ Department of Geodesy and Geoinformation, TU Wien, 1040 Vienna, Austria

10 *Correspondence to:* Fabian Fleischer (Fabian.fleischer@ku.de)

Abstract. Permafrost is being degraded worldwide due to the change in external forcing caused by climate change. This has also been shown to affect the morphodynamics of active rock glaciers. We studied these changes, depending on the analysis, on nine or eight active rock glaciers respectively with different characteristics in multiple epochs between 1953 and 2017 in Kauner Valley, Austria. A combination of historical aerial photographs and airborne laser scanning data and their derivatives were used to analyse surface movement and surface elevation change. In general, the studied landforms showed a significant acceleration of varying magnitude in the epoch 1997–2006 and a volume loss to variable degrees throughout the investigation period. Rock glaciers related to glacier forefields showed significantly higher rates of subsidence than talus-connected ones. Besides, we detected two rock glaciers with deviating behaviour and one that showed an inactivation of its terminal part. By analysing meteorological data (temperature, precipitation and snow cover onset and duration), we were able to identify possible links to these external forcing parameters. The catchment-wide survey further revealed that, despite the general trend, timing, magnitude and temporal peaks of morphodynamic changes indicate a slightly different sensitivity, response or response time of individual rock glaciers to fluctuations and changes in external forcing parameters.

1 Introduction

Rock glaciers are a downslope creep phenomenon of mountain permafrost and are responsible for extensive mass transport in alpine environments (Barsch, 1996). As it has been shown that rock glacier kinematics are affected by changing external forcing factors, the aim of this study was to investigate how rock glaciers in the same catchment react to these changes on a multi-decadal time scale. Active rock glaciers consist of a generally coarse debris layer (active layer), covering ice supersaturated debris and form lobate or tongue shaped landforms (Haeberli et al., 2006). They can be found in most cold mountain regions of the earth (Jones et al., 2019). Although permafrost is the primary condition for rock glaciers to form, the deforming ice and debris may be of both glacial and periglacial origin (Berthling, 2011). Borehole cores and inclinometer measurements from different rock glaciers reveal the internal structure and deformation of these landforms (Arenson et al.,

2002; Krainer et al., 2015; Buchli et al., 2018). They suggest rock glaciers to be similarly composed by such layers, although structure, composition and thickness of these layers differ to a certain extent also within individual rock glaciers. The few meters thick topmost active layer consists of unconsolidated debris and isolates the underlying ice-rich frozen body. The ice-rich permafrost body constitutes the main layer of the rock glacier, with a smaller proportion of coarser debris per volume than the active layer and volumetric ice contents between 40% - 90% (Arenson et al., 2002; Haeberli et al., 2006; Hausmann et al., 2012). The thermal regime of the permafrost layer is mainly controlled by heat conduction, therefore the temperature signal from the surface is linearly delayed and its amplitude exponentially decreases with depth (Haeberli et al., 2006). As a result, changes in temperature of the permafrost body below the depth of zero amplitude require long-term changes of the thermal forcing. In the permafrost layer, internal plastic deformation is the main component of deformation, which is governed by temperature and the structure of the debris-ice mixture. A large part of the horizontal deformation (50% -97 %) takes place in one or more shear zones at the base of the ice-rich permafrost body, which are maximally a few meters thick (Arenson et al., 2002). The dynamics of rock glaciers vary at different temporal scales: decadal, inter-annual and intra-annual (Delaloye et al., 2008; Delaloye et al., 2010; Wirz et al., 2016; Kenner et al., 2017). Studies investigating the decadal-scale variability of rock glacier morphodynamics in the European Alps reveal a significant increase in flow velocities starting in the 1990s, while some studies also report phases of stable flow velocities and velocity decrease since then (Roer, 2005; Kellerer-Pirklbauer and Kaufmann, 2012; Scapozza et al., 2014; Hartl et al., 2016; Kellerer-Pirklbauer and Kaufmann, 2018; Kenner et al., 2020; Marcer et al., 2021). The increase in flow velocities has been explained by rising mean annual air and ground temperatures and mechanisms of heat conduction and melt water advection (Roer, 2005; Käab et al., 2007; Ikeda et al., 2008; Delaloye et al., 2010). More recent studies highlighted the role of liquid water, especially in the shear horizon, and attribute little or no significance to the change in permafrost temperature to explain the deformation variations on a multi-annual, inter-annual, seasonal and short-term scale (Wirz et al., 2016; Kenner et al., 2017; Buchli et al., 2018; Cicoira et al., 2019). Kenner et al. (2020) synthesise these findings by showing that water availability in the rock glacier is governed by ground temperature which is a function of mean annual air temperature and onset as well as duration of snow cover and thus correlate with rock glacier deformation as well. Besides, rock glacier dynamics can also be influenced by other factors like topography, temporal and vertical variations in ice content, rheology of the ice-debris mixture, thickness, and input of ice and debris to the system. Although rock glaciers normally move at rates ranging from a few centimetres per year to a few meters per year, some studies show a destabilization of rock glaciers, which results in a landslide like movement with displacement rates of up to several tens of meters per year (Roer et al., 2008; Scotti et al., 2017; Vivero and Lambiel, 2019; Marcer et al., 2021).

The present and former response of rock glacier morphodynamics to atmospheric warming and climate change observed in many high mountain regions (Hock et al., 2019) is of large scientific interest for climate change projections and landscape evolution models. But an understanding of these landforms has also implications for natural hazard protection (Schoeneich et al., 2015), future water availability (Jones et al., 2019) or sediment connectivity (Kummert and Delaloye, 2018). Although there are several studies on rock glacier kinematics covering different time scales, the number of studies is low compared to ice glaciers. Apart from Roer et al. (2005), who investigated multi-decadal catchment wide rock glacier kinematics in Turtman

Valley, Swiss Alps, most studies on a decadal time scale investigate just one or two large and prominent rock glaciers (e.g. Scapozza et al., 2014; Scotti et al., 2017; Kellerer-Pirklbauer et al., 2018; Kellerer-Pirklbauer and Kaufmann, 2018; Kaufmann et al., 2019; Kenner et al., 2020).

Therefore, we focus on the long-term (1953-2017) kinematic investigations of rock glaciers, located in the Upper Kauner Valley, Ötztal Alps, Austria, displaying different characteristics. By doing so we want to understand their reaction to climate change under similar climatic forcing. We do this by analysing surface movement (flow velocity) of eight rock glaciers by means of image-correlation techniques on the base of orthoimages and hillshades. In addition, multitemporal surface elevation changes are derived for one additional rock glacier (nine) by a DEM of Difference (DoD) analysis using photogrammetric as well as Airborne Laser Scanning (ALS) data. The identified changes in rock glacier kinematics will be discussed with regard to rock glacier characteristics and changes in the meteorological forcing, by investigating different climate parameters (temperature, precipitation and snow cover) recorded directly in the catchment and nearby meteorological stations.

2 Study Area

The investigated rock glaciers are located in the catchment area of the Upper Kauner Valley within the Ötztal Alps, Austria (Fig. 1). The study area is more precisely defined as the hydrological contributing area at the inlet of the Fagge river into the Gepatsch reservoir. It has an area of ~62 km² and a relief of 1773 m ranging from 1810 m at the inlet to 3583 m at the summit of Hochvernagtspitze.




Figure 1. Location and relative size of the investigated rock glaciers and location of the analysed meteorological stations. The stations Obergurgl and Obergurgl-Vent are marked as one station. The coordinate system used is ETRS89 / UTM zone 32N EPSG:25832 with the unit meter. Background was created using Copernicus data and information funded by the European Union - EU-DEM layers (<https://land.copernicus.eu/imagery-in-situ/eu-dem>).

Geologically, the study area belongs to the Eastern Alps crystalline zone and the polymetamorphic Stubai complex. Paragneiss and Orthogneiss are dominant, but Amphibolit and Mica-schist occur subordinately (Hoinkes and Thöni, 1993). The study area is climatically characterized by the central-alpine dry region (Fliri, 1975). At Weißsee meteorological station (2470 m), a mean annual air temperature (MAAT) of -0.11 °C (2007-2019) and annual precipitation ranging from 731 mm to 1118 mm were recorded (data source: Tiroler Wasserkraft AG (TIWAG)). Detailed analysis of climate parameters of the study area is provided in Sect. 5.3 of this paper. According to a permafrost probability map (Otto et al., 2020), 38% of the study area could be underlain by permafrost and 30% was covered by glaciers in 2015 (Buckel and Otto, 2018). The most prominent, Gepatschferner and Weißseeferner are located in the southern part of the catchment. Due to the glacier road, which makes the valley accessible by car, many studies by geo- and bioscience have been carried out in this area (Dusik et al., 2015; Groh and Blöthe, 2019; Altmann et al., 2020). But the road, which intersects RG03 and the associated ski area also cause anthropogenic influences on natural systems, which have to be considered.

The study of rock glaciers has a long tradition in the Kauner Valley, especially at the rock glacier Innere Ölgrube, which is named RG01 in this study (Fig.2). The first velocity studies were already carried out by Finsterwalder (1928) and Pillewizer (1957) and more recent studies continue their research and contribute additional information about the hydrology, internal and external structure and morphodynamics of this rock glacier (Berger et al., 2004; Krainer and Mostler, 2006; Hausmann et al., 2012; Groh and Blöthe, 2019). Groh and Blöthe (2019), who investigated the recent development of flow velocities of rock glaciers in Kauner Valley, ascertained a slightly increase and noted that the velocities of rock glaciers in the study area mainly depend on parameters describing the general inclination and that their activity status is controlled by their size and the topoclimate.




Figure 2. Images of two of the studied rock glaciers. (a) The north-exposed rock glacier RG08, located at the entrance to the Krummgampen Valley and covering an area of 0.088 km². (b) The well-studied, west-exposed Ölgruben rock glacier RG01 covering an area of 0.237 km².

We investigated eight active rock glaciers representing different characteristics and conditions in detail regarding flow velocities and one more rock glacier regarding the surface elevation change analysis (Fig. 1). These are described in more detail in Sect. 5.1.

3 Materials

3.1 Aerial photographs

In order to quantify the kinematic changes of the rock glaciers that occurred in the previous century, we used historical aerial photos to derive orthoimages and point clouds. The historical aerial photos used in this study were scanned and provided in tiff-format by "Office of the Tyrolean Government-Department of Geoinformation" (<https://www.tirol.gv.at/en/>) and "Austrian Federal Office of Surveying and Metrology" (BEV) (<https://www.bev.gv.at>) along with the camera calibration protocols if available (Table 1). The aerial photos for the Kauner Valley catchment were collected at six separate epochs between 1953 and 1997. The epochs were chosen based on data availability, similarity in acquisition date (i.e. late summer), image quality, and sufficient image overlap.

Table 1. Overview of the acquired historical aerial image flights used to generate point clouds and orthoimages in Agisoft Metashape Professional (v.1.6.1).

Flight date	Number of images*	Purpose	Source	Camera	Focal length [mm]	Scanning resolution [µm]	Flight altitude [m]	Resolution orthoimage [m]	Point density*** [pts/m ²]
Jun. 5 th Aug. 31 st Sep. 8 th 1953**	124	Forest condition estimation; Flight C	BEV	unknown	210.11	15	3210	0.225	6.6
Sep. 1 st 1954	36	Forest condition estimation; Flight D	BEV	unknown	210.11	15	3710	0.225	7.0
Sep. 29 th 1970	32	Tyrolean state surveying flight	Land Tirol	Wild RC5/RC8	210.43	12	3590	0.2	10.1

Aug. 18th 1971	91	Tyrolean state surveying flight	Land Tirol	Wild RC5/RC8	209.48	12	3120	0.2	12.0
Sep. 13th 1982	34	Tyrolean state high altitude surveying flight	BEV	Wild RC10	152.58	15	6030	0.5	1.2
Sep. 11th 1997	25	KF 173	BEV	Wild RC10	152.70	15	5860	0.5	1.2

* Number of images used to reconstruct the whole catchment

** Three Dates were processed as one dataset

125 *** Mean point density on studied rock glaciers with snow patches excluded

3.2 Airborne Laser Scanning (ALS) data

To analyse rock glacier flow velocities on hillshades (see Sect 4.3) and surface elevation change using DoDs (see Sect. 4.4) in the two most recent epochs 2006-2012 and 2012-2017, we used data from different ALS campaigns (Table 2). The most recent one was acquired on the 5th of June 2017 by a helicopter and a mounted mobile laser scanning system VuxSys-LR from Riegl
130 (www.riegl.com). This ALS flight mission was carried out by the Chair of Physical Geography at the Catholic University of Eichstätt-Ingolstadt during the DFG-funded project (PROSA), achieving a mean point density of 20.0 pts/m² on the studied rock glaciers. Due to weather conditions and time constrains of the contracted company, which made an area-wide data acquisition on one day impossible, the 2012 ALS data were recorded (also during the PROSA project) on the 4th and 18th of July.

135 **Table 2. Overview of ALS data used for flow velocity analysis and surface elevation change analysis. Average point density is given for the areas of the investigated rock glaciers without snow patches.**

Flight date	Scanner model	Wave length [nm]	Average point density* [pts/m ²]	Operator
Sep. 5 th 2006	unknown	999	5.00	TIWAG 140
Jul. 4 th 2012	Riegl LMS Q680i-S	1064	12.7	Milan Geoservice GmbH (commissioned)
Jul. 18 th 2012	Riegl LMS Q680i-S	1064	12.3	Milan Geoservice GmbH (commissioned)
Jun. 5 th 2017	Riegl VuxSys-LR	1550	20.0	Chair of Physical Geography University of Eichstätt- Ingolstadt ¹⁴⁵

An LMS Q680i-S laser scanner from Riegl mounted on a helicopter was used for data recording. Depending on the date of recording, the average point density ranges between 12.3 pts/m² and 12.7 pts/m². Furthermore, an additional ALS data set from the 5th of September 2006 with an average point density of 5.0 pts/m² was provided by the TIWAG. All datasets were georeferenced with parameters optimised by an automatic strip adjustment (Glira et al., 2015).

3.3 Meteorological data

For the analysis of rock glacier morphodynamics over a decadal time period, a reference to climatic conditions that influence such systems in various ways is indispensable. However, long time series data in the high alpine areas are only very sporadically available, as early climate monitoring stations tended to be located in population centres. This also applies to our catchment area, where the meteorological station Weißsee (2540 m) (data source: TIWAG) is recording data since 2006. For this reason, we used additional data from nearby meteorological stations, which have longer time series available, to provide information on the approximate climatic development in the catchment area. The locations of these stations are shown in Fig. 1, while an overview of the stations and the used data is given in Table 3.

Table 3. Overview of the meteorological stations used. The distance is given to the center of the study area in km. T - Temperature; Pr - Precipitation, SC - Snow cover. The data were provided by the “Federal Ministry of Agriculture, Regions and Tourism“ (BMLRT), the “Central Institute for Meteorology and Geodynamics“ (ZAMG), “Historical Instrumental Climatological Surface Time Series of the Greater Alpine Region“ (HISTALP), “Autonomous Province of Bozen/Bolzano “ (Province BZ) and TIWAG.

Station Name	Recording started	Distance [km]	Temporal resolution	Elevation [m]	Parameters analysed	Provider
Kaunertal-Vergötschen	1895	18.1	monthly, daily	1269	Pr, SC	BMLRT
Nauders	1895	19	daily	1330	T, Pr, SC	ZAMG
Obergurgl	1953	21	daily	1942	T, Pr, SC	ZAMG
Obergurgl-Vent	1851	21.4	monthly	1938	T	HISTALP
Plangeroß	1895	16	daily	1605	Pr, SC	BMLRT
Vernago	1953	17	daily	1700	Pr	Province BZ
Weißsee	2006	2.4	15-min	2540	T, Pr, SC	TIWAG

4. Methods

4.1 Rock glacier inventory

Although manual mapping of rock glacier landforms is shown to be highly subjective (Brardinoni et al., 2019), we tried to minimize the heterogeneity in the inventory by incorporating the guide lines for inventorying rock glaciers ([IPA Action Group Rock glacier inventories and kinematics 2020](#)) and only mandate one operator to compile the inventory on the basis of the inventory by Krainer & Ribis (2012). Rock glacier outlines were corrected and additional landforms were mapped on the basis of the most recent hillshade derived from the 2017 ALS campaign of the PROSA project and an orthoimage of 2015 (data

170 source: Land Tirol - data.tirol.gv.at). Activity status was assigned according to morphological characteristics in combination
with a DoD of the 2012 and 2017 ALS campaigns to detect surface elevation change and therefore ice melting. Image
correlation analysis on the derived hillshades was used to derive information on flow velocity. For the inventory of the entire
study area, both analyses were carried out without local alignment of the data. The classification is based on the common
division into active - contains ice and is in motion, inactive – (almost) no motion but still contains ice, and fossil - contains
175 (almost) no more ice and is not in motion (e.g. Krainer and Ribis, 2012). We then derived rock glacier characteristics such as
aspect, slope, and elevation from the elevation model of the 2017 ALS campagne and classified them according to their spatial
connection to the upslope unit. We further determined whether the area of the investigated rock glaciers were covered by a
glacier during the Little Ice Age (LIA) with the help of a glacier inventory (Fischer et al., 2015).

4.2 Processing of aerial images

180 To derive point clouds and orthoimages from the aerial images, these were processed using advances in digital
photogrammetry, particularly Structure from Motion (SfM) with Multi-View Stereo (MVS). The reconstruction of 3D
information in form of point cloud from scanned historical photos does not require specialized knowledge when important
basic principles are followed (Bakker and Lane, 2017; Fawcett et al., 2019). The aerial images were processed in Agisoft
Metashape (v.1.6.1) using the film camera tool, which estimates the camera calibration parameters based on the fiducial marks.
185 The software automatically derived the locations of the fiducial mark in the images. Their distance in mm and the focal length
were available from the calibration protocols. Having defined the camera interior orientation, the camera exterior orientation,
the 3D point cloud reconstruction and the orthoimage generation followed the standard SfM-MVS workflow. This includes
ground control points (GCPs) measurement for georeferencing and dense image matching. The 3D coordinates of 101 GCPs
were chosen from the ALS 2017 point cloud on stable terrain and were evenly distributed throughout the catchment. In the
190 case of the 1953 data set, three different aerial image flights from this year had to be processed together, as there was
insufficient image overlap in the individual flights.

The resulting average point density on the studied rock glaciers varied from 1.2 pts/m² to 11.9 pts/m² (Table 1). Therefore,
only the point clouds of the years 1953/54 and 1970/71 with a point density between 6.6 pt/m² and 12 pt/m² were used in the
DoD analysis. The ground resolution of the orthoimages varied between 0.2 m and 0.5 m.

195 4.3 Horizontal flow velocities

Horizontal flow velocities of the rock glaciers were calculated for the six processed time steps between 1953 and 2017. For
this purpose, an image correlation approach was chosen, which is a common method to derive glacier and rock glacier velocity
from orthoimage, hillshades and satellite images (Scambos et al., 1992; Kääb and Vollmer, 2000; Heid and Kääb, 2012;
Monnier and Kinnard, 2017; Kellerer-Pirklbauer and Kaufmann, 2018). In this study, orthoimage and hillshade image pairs
200 were utilized. In order to ensure their comparability and to account for possible shifts in the orthoimages, we resampled them
at a resolution of 0.5 m and locally co-registered all individual rock glaciers for each epoch to the 1953/54 orthoimage. We

used 9 to 29 co-registration points equally distributed around the rock glaciers, obtaining co-registration root mean squared errors (RSMEs) between 0.23 m and 0.55 m with an average of 0.32 m.

For the time step 1997 – 2006, where orthoimages and hillshades were used in combination, illumination was chosen for the 2006 DEM according to the exact position of the sun in the 1997 orthoimage. To calculate flow velocity vectors, the image correlation algorithm IMCORR (Scambos et al., 1992) within SAGA-GIS software was applied. The algorithm attempts to match small sub-scenes from two images by applying a fast Fourier transform-based version of a cross-correlation. It can locally adjust the intensity values between two image pairs and therefore compensate for differences in illumination. Using this algorithm, sub-pixel precision of displacement vectors can be achieved. We used search and reference chip size combinations of 64/32, 128/64 and 256/128 with a fixed spacing of 5 m. The combinations were calculated for all image pairs and the most reasonable was chosen for further analysis. This was done by visually analysing the resulting displacement vectors in combination with the input data. In general, larger chip sizes were chosen for faster moving rock glaciers and/or long time spans between the image pairs. The resulting raw vector maps can contain erroneous displacement measurements or decorrelation, where no measurement is possible. Image decorrelation might be caused by snow, strong shading effects, areas where displacements are dominated by rock fall and large displacements, which cause a change of texture. These vectors were excluded manually for all time steps with the help of the matching orthoimages or hillshades. Subsequently, a mask was created for the areas where measurements were possible in all time steps and just measurements in these areas were used for further analysis to make the individual time steps comparable. Since it is therefore not possible to measure the flow velocity over the entire rock glacier, the calculated statistics are only valid for the area in which measurements were possible in all epochs.

The combination of orthoimages and hillshades had to be chosen because low point densities in some of the aerial images derived point clouds resulted in low details in the resulting DEMs. Tests regarding image correlation on these DEMs showed very poor results. We are aware that the low point densities also affected the accuracy of the resulting orthoimages and outline the variable errors in Sect. 5.2. On the other hand, we decided not to use orthoimages for the more recent epochs from 2006 to 2017, available from “Office of the Tyrolean Government-Department of Geoinformation” (<https://www.tirol.gv.at/en/>) for the reason that they were orthorectified utilizing the most up to date DEM with a resolution of 5 m, which could result in erroneous displacement measurements. If a non-contemporary DEM is used, it would lead to orthorectification errors particularly on moving landforms, like rock glaciers (Kaufmann and Kellerer-Pirklbauer, 2015).

The measurement of horizontal flow velocities of rock glaciers on remote sensing data, especially when using historical aerial images and their derivatives, is prone to errors. As described by Käab et al. (2020), the error budget is composed of the following components: 1) overall shifts between the orthorectified data 2) lateral shifts in the orthoimages due to errors in the DEM used for orthorectification 3) distortions in the aerial images or in the sensor model that propagate into the orthoimages 4) image matching uncertainties and errors. We minimized the shifts between the orthoimages by a local co-registration of the orthoimages. By using the matching DEMs of the individual years for orthorectification, we addressed error type 2). However, quality of the DEMs varied locally in a single epoch and more crucially between the epochs and therefore might influence the accuracy of the orthorectification. The DEMs with the lowest quality were the epochs 1982 and 1997. These were also the years

with the worst quality of the raw aerial images (error type 3). Another source of error when working with historical aerial images are scratches and alterations on the original image film caused by storage and age. These can lead to problems in the processing and thus were masked out before processing. Errors of type 4) contain errors caused by the image correlation method itself. The measurement errors as consequence of image correlation vary with the image quality like resolution, shadow, contrast and noise of the image pairs (Kääb et al., 2020). We removed both directional and magnitudinal gross outliers manually by counterchecking the resulting displacement vectors with the corresponding orthoimage and hillshade pairs. To quantify the overall error budget for horizontal flow velocities, we mapped close stable areas of similar texture/roughness and exposition on the single rock glaciers for all time steps. Due to snow and shading effects, these stable areas had to be adjusted slightly for some epochs. Subsequently, displacement vectors in these areas were analysed for all individual epochs and rock glaciers. As no gross outliers were found in these areas, we used the mean value ($disp\bar{x}$) added by two times the standard deviation ($disp\sigma$) as measure for error budget ($error_{disp}$) of flow velocity measurements.

$$error_{disp} = disp\bar{x}_{\Delta epoch1-epoch2} + 2disp\sigma_{\Delta epoch1-epoch2} \quad (1)$$

This measure was also applied by Fey and Krainer (2020) to determine a level of detection (LoD) for rock glacier flow velocity and mean and standard deviation in stable areas are recommended as a statistical measure of flow velocity error by Paul et al. (2017).

4.4 Surface elevation and volume change

For the analysis of surface elevation and volume change on the rock glaciers we used both photogrammetric and ALS point clouds in a DoD analysis. All processing steps were performed in SAGA LIS (Laser Information System; Laserdata.at). The epochs for this analysis differ slightly from the datasets used for the flow velocity analysis as the processing of the aerial images did not lead to sufficient point cloud resolutions for the datasets 1982 and 1997. Therefore, four epochs between 1953 - 2017 were analysed. All point clouds (photogrammetrically derived and ALS) were thinned (0.5 m) to produce homogeneous point densities for all epochs and outliers were removed. In order to minimize inherent systematic errors (Bakker and Lane, 2017) all point clouds were co-registered with the reference 2017 ALS data by using an iterative closest point (ICP) algorithm (Besl and McKay, 1992) on mapped stable areas around the rock glaciers. Fine registration was performed for all individual rock glaciers and epochs separately to account for any variable errors throughout the catchment. The point clouds were then converted into a DEM with a cell size of 1 m. The elevation change for each epoch and rock glacier was derived by subtracting the earlier DEM from the newer DEM (DoD). These values were summed and multiplied by the squared cell size to obtain the volume change. In order to be able to compare epochs, annual values were calculated. To avoid surface elevation changes and volume changes being influenced by snow, snow masks were created and these areas were excluded from the analysis (Fig.10). This implies that in some cases the entire landform cannot be considered for the mass balances, so the results of these must be interpreted with caution.

For the uncertainty analysis we followed the approach proposed by Anderson (2019), who presented an error propagation
270 methods to derive uncorrelated, correlated and systematic errors and combine them to a total uncertainty without applying a
minimum LoD. We carried out this procedure on stable areas that we mapped in proximity to the rock glaciers. For
visualisation, however, we applied a LoD of 1.96 times the standard deviation in stable areas.

4.5 Meteorological data

In order to identify possible links between rock glacier kinematics and external forcing factors, we analysed the changes in
275 temperature, precipitation and snow cover during the study period from 1953 to 2017. In addition to a trend analysis of the
data, we also calculated the annual anomalies based on the reference period 1961 to 1990 and the 5-year running mean to
smooth out short-term fluctuations. In addition, we disaggregated these analyses by season, here, spring was defined as March
– May, summer as Juli – August, autumn as September – November and winter as December – February. The variables snow
cover onset and duration were determined according to Peng et al. (2013) From the depth of snow data of the individual station
280 we also calculated the dates when a significant snow cover (> 50 cm) is established, allowing for ground isolation. Besides
this we derived the date of complete snow melt in spring.

5 Results

5.1 Rock glacier inventory and investigated rock glaciers

A rock glacier inventory was compiled for the Upper Kauner Valley. Within the catchment, 40 rock glaciers were found, which
285 were classified as active (15), inactive (11) and fossil (14) (Fig. 3). Due to poor image quality or snow cover and the activity
status, the vast majority of these rock glaciers had to be excluded from the following analyses.




Figure 3. Roseplot of the rock glacier inventory of the Upper Kauner Valley. The radial axis gives the mean elevation of the rock glaciers [m], the angular axis gives the exposition of the rock glaciers [°]

290 Nevertheless, eight active rock glaciers representing different characteristics and conditions were investigated in detail regarding flow velocities and one more regarding surface elevation change. The most prominent of those is the well-studied and largest (0.237 km²) rock glacier Innere Ölgrube (RG01) (Fig. 2). The area of these rock glaciers ranges from 0.02 km² (RG07) to 0.237 km² (RG01). They show expositions of N, NE, E and W, with minimum elevation ranging from 2446 m to 2727 m. Without geophysical, geochemical or petrographic information, interpretations about the genesis and internal structure are difficult (Berthling, 2011; Clark et al., 1998). In the case of the rock glaciers RG03, RG04 and RG09 a complete or partial covering of the rock glaciers by the LIA glacial extent (Fischer et al., 2015) suggested a glacial genesis after 1850 or at least a glacial influence during and after this time as described by Dusik et al. (2015) for RG09. In the case of RG06, a thermokarst lake suggested a glacier coverage of the root zone during LIA and the presence of massive sedimentary ice.

300 **Table 4. Characteristics of the rock glaciers studied. Permafrost occurrence gives the pseudo-probability of permafrost (Otto et al. 2020). Area covered by 1850 glacier extent is ascertained according to LIA glacier extends (Fischer et al. 2015) if not specified otherwise.**

Rock glacier	RG 01	RG 02	RG 03	RG 04	RG 05	RG 06	RG 07	RG 08	RG 09
Exposition	W	NE	N	NE	W	E	E	N	NE
Area [km²]	0.237	0.058	0.036	0.036	0.059	0.182	0.02	0.088	0.171
Slope [°]	25	26	26	20	31	24	30	30	22
Elevation [m]									
min	2446	2 615	2596	2 727	2702	2 695	2709	2 510	2627
max	2780	2 755	2787	2 812	3093	2 948	2861	2 761	2925
Permafrost occurrence [%]									
min	0	0	16.97	0	0	0	0	0	0
max	65.41	57.95	45.68	70.53	76.04	81.42	50.52	60.52	78.57
mean	14.49	33.22	48.99	47.08	48.24	44.04	20.54	38.33	37.57
Connection to the upslope unit	GFC; TC	TC	GFC	GFC	TC	TC	TC	TC	GFC; TC
Area covered by 1850 glacier extent	Yes*	No	Yes	Yes	No	No***	No	No	Yes**

*As described in (Berger et al. 2004);

** As described in (Dusik et al. 2015);

*** A thermokarst lake suggests massive ice and a covering of the root zone by LIA glacier.


305 Concerning the connection to the upslope unit both glacier forefield connected (GFC) and talus connected (TC) rock glaciers were included in the further analysis. Detailed characteristics of the studied rock glaciers can be found in Table 4 and a KMZ-file containing the locations of the investigated rock glaciers is included in the supporting materials.

5.2 Errors and uncertainties

310 Although the workflow was chosen to minimise errors during data processing, they are still present to some extent in the data. Wherever necessary and possible, we indicated the uncertainties of the values in the text and in the figures.

For flow velocity measurements, error values ranged between 0.01 m/yr and 0.09 m/yr. The distribution of these values from individual rock glaciers per image pair combination are displayed in Fig. 4. In general, best results were obtained for epochs where hillshade image pairs were utilized. The largest errors on average were determined for the epoch 1997-2006, in which
315 a combination of orthoimages and hillshade was used. We attribute this to the poor image quality of the 1997 dataset and the use of orthoimages in combination with hillshades. Although the illumination of the hillshades was adapted to the orthoimages, only contrasts and patterns, which are caused by the illumination, can be used to determine the flow velocity by image correlation. On the contrary, contrasts and patterns that are caused by different colours in the orthoimage are not taken into account or can even lead to incorrect measurements. In other cases, the variability of the errors may be related to the accuracy
320 of the co-registration, but also to differences in the quality of the image in terms of contrast, illumination, and resolution.

To assert the validity of our results we performed a qualitative comparison with dGPS measurements, which were taken by Krainer and Mostler (2006) between 2002 and 2004 for Ölgruben rock glacier (RG01). This confirmed the magnitude, flow direction and pattern that were determined by our method. In addition, a comparison of flow velocities derived by Groh and Blöthe (2019) by feature tracking for Kauner Valley rock glaciers between 2001 and 2015, showed good agreement. Although
325 other time-steps were used, span, mean and median values showed similar and plausible results when compared to the values of the two most recent time steps of our study.



330 **Figure 4. (a) Flow velocity errors derived from stable areas in proximity to the individual rock glaciers for all image pair combinations. Like all subsequent boxplots, these show summary statistics of the data, with the bottom of the box representing the 25th percentile, the top of the box the 75th percentile and the black bar the median. The lines above and below the boxes indicate the minimum and maximum values. (b) Example of a stable area displaying similar exposition, texture/roughness used to analyse error for RG08.**

For the surface elevation change and volume change analysis, the uncertainties were largely determined by the magnitude of the systematic error of the DoD. The analysis of the errors revealed that a systematic error mostly occurred if one or both DEMs

335 used to calculate the DoD were photogrammetrically created from historical imagery. This is reasoned by the fact that, due to
the quality of the data, these point clouds could not be matched to the reference dataset by the ICP algorithm with the same
accuracy as, for example, high-resolution and high-precision modern ALS datasets. Since the data quality of the point clouds
generated from historical images varies spatially, a systematic error did not occur to the same severity in all investigated rock
glaciers of an epoch. The spatially correlated random error, on the other hand, always occurred in DoDs with at least one DEM
340 generated from historical images, whereas it was not present in DoDs created from ALS data. However, similar to the
uncorrelated random error, this contributed only little to the total uncertainty in case a systematic error was present.
Since areas of snow cover, strong shadows or decorrelation in the image correlation would bias the results, masks were created
comprising areas in which an unbiased analysis was possible in all epochs in order to ensure comparability. Therefore, the
analysis of the flow velocity could be conducted on 27.39% to 80.00% of the rock glacier area, while, regarding the surface
345 elevation change analysis, a coverage of 50.50% to 95.67% could be achieved (Table 5).

Table 5. Percentage of the rock glacier area in which measurements were possible in all time steps and which were therefore considered for the analysis of flow velocity and surface elevation change. Reasons for exclusion were snow cover, shading effects or decorrelation in the image correlation.


Rock glacier		RG01	RG02	RG03	RG04	RG05	RG06	RG07	RG08	RG09
Area [%]	Flow velocity	41.97	75.94	45.42	65.72	27.39	39.12	70.62	80	-
	Surface elevation change	92.09	87.72	70.93	90.71	81.1	64.4	92.9	95.67	50.5

5.3 Changes in meteorological forcing

350 All stations showed similar patterns of change, even if the manifestation of the anomalies was slightly different in some cases.
We note that the positive trend of temperature increase was slightly higher for stations of higher elevation in the study period.
In the case of temperature and snow cover, we mainly present data from the stations Obergurgl-Vent (1938 m a.s.l.) and
Obergurgl (1942 m a.s.l.), as these are located at the highest elevation and only about 21 km away from the centre of our study
area. In the case of precipitation, we mainly present data from the Plangeroß station (1605 m a.s.l.), because although it is
355 located at a significantly lower elevation than the Weißsee station (2540 m a.s.l.) and the studied rock glaciers, it offered the
best agreement with the Weißsee station data in terms of monthly precipitation ($r=0.892$, $p<0.001$).

During the period of investigation (1953-2017), the temperature trend showed an increase of 1.92 °C in 65 years at the
Obergurgl-Vent (1938 m a.s.l.) station (Fig. 5). This was a stronger increase than at the lower elevated station Nauders (1330
m a.s.l.), where the increase was 1.29 °C. Possible reasons and explanations for the elevation-dependent warming are given
360 in Rangwalla and Miller (2012). In general, the temperature increase goes well in line with the alpine wide temperature
increase, which has a significantly higher amplitude than the global average (Beniston, 2006). The seasonal development of
the temperature trends showed a strong increase in temperatures in spring and summer of 2.73 °C and 2.64 °C, respectively.
This trend was clearly weakened for winter and autumn temperatures and amounted to 1.6 °C for winter and 0.69 °C for

autumn. The derived temperature anomalies showed exclusively positive anomalies from the beginning of the 1990s onwards
 365 for spring and summer temperatures (Fig. 12). The summers of 2015, 2017 and in particular the summer of 2003 were
 conspicuous for their high temperatures and are known to have brought profound change to permafrost-affected systems
 (Ravel et al., 2017). For the period before the beginning of the 1990s, periods/years with both positive and negative
 anomalies occurred, whereby the strong anomalies were mostly in the negative range.



370 **Figure 5. Changes in meteorological forcing during the study period between 1953 and 2017. (a) Mean annual precipitation (MAP)**
for the station Plangeroß (1605 m a.s.l.) and mean annual air temperature (MAAT) for the station Obergurgl-Vent (1938 m a.s.l.);
the dashed lines indicate the values for the Weißsee station (2540 m a.s.l.). (b) Snow cover onset and end for the station Obergurgl
(1942 m a.s.l.). The black lines show the trend of the respective parameters.

All considered meteorological stations, except for the station Nauders (1330 m a.s.l.), recorded a slightly positive trend in
 375 terms of precipitation, translating to an increase between 53 mm and 241 mm during the investigation period. The positive
 trend for the station Plangeroß (1605 m a.s.l.) was expressed in an increase in precipitation of 152 mm per 65 years (Fig.5).
 The mean annual precipitation accounted to 931 mm/yr during the period of investigation at the station Plangeroß (1605 m
 a.s.l.) and to 957 mm/yr at the Weißsee station (2540 m a.s.l.) in the period from 2007 to 2017. The greatest positive trend was
 recorded in autumn (52 mm per 65 years), the least in winter (18 mm per 65 years). On average, most precipitation fell in the
 380 summer months, with an average of 362 mm/yr, and the least in winter, with an average of 171 mm/yr.

Precipitation anomalies showed a clear increase from the mid-1990s for positive summer and autumn anomalies, which is
 particularly pronounced in the period 1995-2002 (Fig. 12).

The derived snow cover parameters are difficult to transfer to the study area, as the station Obergurgl (1942 m a.s.l.) is located
 at a lower elevation than the rock glaciers studied but could provide an indication of general trends and anomalies in snow
 385 cover onset and duration (Fig. 5). For the entire study period from 1953 to 2017, no significant trend could be observed for the
 onset of the snow cover, significant snow cover and snow cover duration. This fits in with the results of Olefs et al. (2020),
 who found an elevation-dependent reduction in snow depth and duration in Austria between 1961 and 2020, but this only
 applies to elevations below 2000 m. However, there were some periods and years showing a particularly early or late onset of
 snow cover and significant snow cover respectively (Fig. 12). On the other hand, a significant negative trend of -17 days was
 390 observed for the end of snow melt over the study period, meaning the snow melt takes place earlier with time (Fig.5).

Although this was not always the case, in general, the data showed that when the onset was earlier, the duration of the snow cover was also longer and vice versa.

5.4 Flow velocities

For the whole period of investigation, we derived maximum flow velocities ranging between $0.09 \text{ m/yr} \pm 0.03 \text{ m/yr}$ and $1.72 \text{ m/yr} \pm 0.01 \text{ m/yr}$ and mean flow velocities ranging between $0.08 \text{ m/yr} \pm 0.03 \text{ m/yr}$ and $0.60 \text{ m/yr} \pm 0.01 \text{ m/yr}$ for the eight individual rock glaciers (Fig. 6). The mean values were calculated for values above the maximum LoD of the respective rock glacier in order to exclude marginal areas, showing values close to zero, that do not move homogeneously with the rock glacier and to ensure comparability between the epochs. All of the investigated rock glaciers showed the highest mean and/or maximum values either in the epoch 1997-2006 or 2012-2017.





Figure 6. Flow velocities for individual rock glaciers for six distinct epochs from 1953 to 2017. Rock glacier RG09 was not included in the flow velocity analysis and rock glacier RG 08 had to be excluded from epoch 1 due to extensive decorrelations in the frontal part because of large displacements. The mean value (red dot) was calculated for values above the maximum LoD of the respective rock glaciers. The respective LoDs of the epochs are shown as red boxes. The black dots represent outliers, which by definition are located at least 1.5 times the box length (interquartile range) above the box.

An exception to this was RG08, which had the highest maximum flow velocity of 1.66 m/yr in the epoch 1953 to 1971. In this epoch, flow velocities for RG08 could only be determined by manual mapping of a few blocks, as image correlation resulted in decorrelation due to excessive changes in surface structure. RG01, RG02, RG03, RG06, RG07 and RG08 showed a distinct acceleration of flow velocities beginning in the epoch 1997-2006. All of the investigated rock glaciers of the Kauner Valley showed an increase in flow velocities in the most recent epoch, 2012-2017 compared to the previous 2006-2012 epoch. Exceptions to the general trend could particularly be seen in the case of RG04, as it was characterised by relatively constant mean flow velocities well below 0.15 m/yr during the whole investigation period.






Figure 7. Maps and violin plots showing the flow velocities and its distribution for the individual rock glaciers for six epochs between 1953 and 2017. The units of the colour scales are m/yr. Areas with flow velocities below LoD are marked with a black grid. The background image is always the older date of the timespan. The colour of the violin plots corresponds to the colour of the respective time period in the maps. Violin plots show the kernel probability density of the data at different values.

420 In the first epoch, all rock glaciers, except for RG01 ($0.49 \text{ m/yr} \pm 0.03 \text{ m/yr}$) and RG08 (max. 1.66 m/yr by manual mapping), displayed similar, low mean values ranging from $0.08 \text{ m/yr} \pm 0.03 \text{ m/yr}$ to $0.16 \text{ m/yr} \pm 0.02 \text{ m/yr}$ and value ranges (Fig. 6). If the maximum flow velocities are taken into account, this similarity has to be put into perspective, as the range here was significantly greater and lies between $0.09 \text{ m/yr} \pm 0.03 \text{ m/yr}$ and $0.41 \text{ m/yr} \pm 0.03 \text{ m/yr}$, due to the higher flow rates of RG 03-05 at their terminal part (Fig. 7).

425 Compared to the first epoch, the range of the mean values increased significantly in the epoch 2012 - 2017 and was now between $0.12 \text{ m/yr} \pm 0.03 \text{ m/yr}$ and $0.49 \text{ m/yr} \pm 0.03 \text{ m/yr}$ and $0.60 \text{ m/yr} \pm 0.01 \text{ m/yr}$ for RG01. The strongest proportional increase could be seen in rock glaciers RG07 and RG02, where mean flow velocities increased by 485% and 251% and maximum flow velocities increased by 799% and 383%, respectively. In contrast, in the case of RG04, the average and maximum flow velocity was reduced by -11% and -47%, respectively.

430 The relative changes between the first and the last epoch regarding the remaining rock glaciers differed clearly and ranged between 23% and 141% for mean flow velocity and 22% and 348% for maximum flow velocity (Fig.6).

Some rock glaciers also showed a local temporal peak in one or more epochs, which was characterised by higher flow rates. On rock glaciers RG01 and RG08, higher mean flow velocities were measured between 1953/54 and 1970/71 compared to the subsequent two epochs. Higher mean flow velocities were also observed on rock glaciers RG01, RG03, RG06 and RG08 in
435 the epoch 1997 to 2006, compared to the epochs before and after, although caution is required in the interpretation due to the higher error values (mean error 0.08 m/yr) in this epoch.

Furthermore, rock glaciers did not move uniformly, but flow velocities varied spatially (Fig. 7). The zones of higher flow velocity were usually, but not always, located in the terminal section of the rock glacier. Exceptions were mainly RG03, which showed a gradual change in its flow pattern over time and RG06 which showed its maximum flow velocities in an area above
440 a thermokarst depression in particular in the latest epoch. On RG08 maximum flow velocities were measured in the terminal part, which gradually decreased and were close to zero in the most recent epoch, whereas the rest of the rock glacier accelerated from the epoch 1997–2006 onwards.

For all landforms, there were areas of varying extent that showed very little to no measurable movement over the entire study period. Apart from these areas, most of the remaining rock glacier areas accelerated especially from the epoch 1997 - 2006,
445 though to varying degrees. The taper (entire area accelerated) and thickening (individual areas of different velocities) of the violin plots as well as the observation of the flow velocity maps give an indication for the acceleration of different zones of the individual rock glaciers studied (Fig. 7).

5.5 Surface elevation and volume changes

The calculation of the surface elevation and volume change could be carried out on a total of nine rock glaciers in four epochs.
450 The volume changes of the investigated rock glaciers ranged from $-9931 \text{ m}^3/\text{yr} \pm 57\text{m}^3/\text{yr}$ (RG1, epoch 2012 – 2017) to $235 \text{ m}^3/\text{yr} \pm 121 \text{ m}^3/\text{yr}$ (RG05, epoch 1953 – 1970) (Fig. 9). Since the mean annual surface elevation changes allow for a better comparison of rock glaciers of different sizes, we will describe them in more detail in the text. However, the volumetric

changes of the individual rock glaciers are also shown in Fig. 9. Since snow-covered areas had to be excluded from the analysis and only the rock glacier body was considered, it is not possible to speak of a true mass balance. Possible drawbacks and consequences are discussed in chapter 6.5.




Figure 8. Boxplots showing annual surface elevation changes on the single rock glaciers and corresponding stable areas (greyed out) for the four epochs between 1953 and 2017.

The mean values ranged from $0.005 \text{ m/yr} \pm 0.003 \text{ m/yr}$ (RG05, epoch 1953 - 1970) to $-0.047 \text{ m/yr} \pm 0.001 \text{ m/yr}$ (RG04, epoch 2006-2012) (Fig. 9). In the entire study period, the values were predominantly in the negative range, with only 3 measurements showing positive values very close to zero. All rock glaciers showed values of surface elevation change in the negative and the positive range (Fig. 8). A clear difference could be seen in the temporal development of the surface and volume changes of different the rock glaciers. On RG01, RG03, RG04 and RG09, negative values ranging between $-0.022 \text{ m/yr} \pm 0.005 \text{ m/yr}$ and $-0.008 \text{ m/yr} \pm 0.002 \text{ m/yr}$ already occurred in the first epoch and became increasingly negative in the following epochs, reaching values ranging between $-0.046 \text{ m/yr} \pm 0.001 \text{ m/yr}$ (RG04) and $-0.021 \text{ m/yr} \pm 0.001 \text{ m/yr}$ (RG03) in the last epoch. All of these rock glaciers were classified as GFC and were covered or partially covered by glaciers during the LIA (Table 4). On the other rock glaciers, which are of TC-type, considering the uncertainties, there were no negative mean surface elevation changes or volume changes measurable in the first epoch. For RG02, RG05 and RG07 they only occurred in the last epoch between 2012 and 2017 and ranged from $-0.015 \text{ m/yr} \pm 0.001 \text{ m/yr}$ (RG05) to $-0.003 \text{ m/yr} \pm 0.001 \text{ m/yr}$ (RG07). In contrast for RG06 and RG08, these already became increasingly negative from the epoch 1970 - 2006 onwards and in the last epoch showed values of $-0.032 \text{ m/yr} \pm 0.001 \text{ m/yr}$ (RG06) and $-0.017 \text{ m/yr} \pm 0.001 \text{ m/yr}$ (RG08).

The investigated rock glaciers differed not only in their temporal development but also in the magnitude of the change in surface elevation. RG01 and RG04 showed similar and very high values, and RG06, RG08 and RG 09 were in a similar, middle of the range. RG02, RG05 and RG07 showed values close to zero, with the exception of RG05 which had clearly negative values in the most recent epoch (Fig. 9). Figure 10 shows the spatial patterns of the surface changes of the rock glaciers and their temporal development. It can be seen that for a large part of the rock glaciers the spatial patterns of positive and negative

surface change remained relatively consistent and the magnitude, mostly of the negative values, increased as time progresses. Exceptions to this spatial development are RG02, RG05 and RG07, where hardly any significant surface change was measured in the first epoch and therefore a spatial pattern only emerged from 1970/71 onwards.






Figure 9. Mean annual volume change and mean annual surface elevation change of the individual rock glaciers for four epochs between 1953 and 2017. Black error bars give the uncertainty of measurements.

RG04 only showed negative values of surface elevation change in the upper part in the first epoch, which were measured on the entire rock glacier from the second epoch onwards. Probably the greatest exception in the case of spatial patterns was 485 RG08. After high values of both positive and negative surface elevation change were measured in the terminal area between 1953 and 2006, only small negative changes could be observed in this area thereafter. From 1971 onwards, however, the upper part of the rock glacier showed increasingly negative values.





500 **Figure 10. Mean annual surface elevation change maps for all individual rock glaciers and four epochs between 1953 and 2017. A LoD of twice the standard deviation of measurements in stable areas was applied for reasons of visualisation. The background image is a hillshade of the more recent DEM of the epoch. The white area with blue dots represents the snow mask. The black boxes in the maps of epoch 2012 – 2017 represent the respective subareas used for Fig. 11.**

5.6 Interaction of surface elevation change and flow velocity

To be able to relate and compare both the changes in flow velocity and the surface elevation change, we have plotted their mean values for subareas of the individual rock glaciers in Fig. 11, with the subareas shown in Fig. 10. The subareas were selected according to the following criteria: (1) representative of the rock glacier, (2) both surface elevation change and flow velocity measurements available, (3) areas above the LoD of flow velocity, (4) homogeneous flow velocities, (5) no influence of obvious thermokarst deepening. To make the two variables comparable for the entire study period, the epoch division of the surface elevation change analysis, which is not as highly resolved in terms of temporal resolution, was used. Although the general pattern of the previous analysis of the entire rock glaciers of increasing velocities and increasingly negative mean surface elevation change values was confirmed, the subarea analysis showed a slightly different pattern for some rock glaciers in terms of values, magnitude of change and timing of change particularly for the surface elevation changes. This is to be expected due to the spatial heterogeneity and the different temporal development of both the flow velocities and the surface

510

elevation changes on some of the investigated rock glaciers as described in the previous chapters and illustrated in Fig. 7 and Fig. 10.




Figure 11. Mean flow velocity plotted against mean surface elevation change for subareas of the investigated rock glaciers (indicated in Fig. 10) and the four epochs between 1953 and 2017. The black crosses represent the uncertainty of the mean flow velocity and the mean surface elevation change, respectively.

In the case of the subareas of the rock glaciers RG02, RG03, RG06, RG07 and RG08, the changes tended to be proportional to one another to a certain degree. Although the values of the subareas were obviously different from the mean values of the entire rock glaciers, both showed a similar temporal development. Whereas the values for RG02, RG03, RG06 and RG07 increased significantly in both epochs after 2006, RG08 showed a significant increase in both variables only between the epochs 1970 - 2006 and 2006 - 2012. Rock glaciers RG01, RG04 and RG05 showed a clear deviation from the behaviour described above, since mean surface elevation change and flow velocity did not behave similarly. In addition, the observations in the subareas for RG01 and RG05 also differed from the observations made when considering the entire rock glaciers. RG04 showed no significant change in flow velocity, whereas the surface elevation changes between the 3 epochs between 1953 and 2012 increased significantly. In the case of RG01, the mean surface elevation change showed the most negative value in the epoch 1953 - 1970, to become smaller again after a significant increase between 1970 - 2006. The flow velocity followed this trend, but showed a higher value in the epoch 2012 - 2017 than in the first epoch. RG05, together with RG04, was the only one that showed significantly more negative surface elevation change values from epochs 1 to 2, whereas the flow velocity hardly increased. Between the epochs 2006 - 2012 and 2012 - 2017, the surface elevation changes became significantly more

negative and the flow velocity also increased sharply. In summary, in the case of the rock glacier subareas, the change in flow velocity and surface elevation change followed a similar trend. Exceptions to this could be observed in particular on rock glaciers RG04 and RG05.

535 6. Discussion

6.1 Development of flow velocities in a regional context

The magnitude of the flow velocity of the studied rock glaciers was well within the range of the common movement of a few centimetres per year to a few meters per year over the entire study period (Delaloye et al., 2010). The timing of the acceleration of most rock glaciers in the 1997-2006 epoch also agreed well with other studies that have investigated the kinematics of rock glaciers in the European Alps over a multi-decadal period. All of these studies observed a significant acceleration of flow velocities in the early to late 1990s (Roer et al., 2005; Kellerer-Pirklbauer and Kaufmann, 2012; Micheletti et al., 2015; Scapozza et al., 2014; Hartl et al., 2016; Kaufmann et al., 2018; Kellerer-Pirklbauer and Kaufmann, 2018; Kellerer-Pirklbauer et al., 2018; Kenner et al., 2020). This is not surprising as an almost synchronicity of rock glacier flow velocity on an annual and multi-decadal scale has been ascertained for the European Alps (Delaloye et al., 2008; Delaloye et al., 2010).

540 Kellerer-Pirklbauer and Kaufmann (2012) found values of 150% to 320% for relative changes in mean flow velocities for a similar period of study for three rock glaciers located in the Hohe Tauern Range, Austria. Furthermore, Roer (2005) determined values ranging from 16% to 350% investigating a multitude of rock glaciers located in the Turtmann Valley, Switzerland. Therefore, our value range of 23% to 251%, except for RG04 (-11%) and RG07 (485%), agree well with the previously determined multi-decadal relative velocity increases of rock glaciers. Conversely, rock glacier destabilisation, which is

545 increasingly observed on rock glaciers in recent years (Roer et al., 2008; Scotti et al., 2017; Vivero and Lambiel, 2019; Marcer et al., 2021), could not be identified on any of the studied rock glaciers.

6.2 Atypical development of flow velocities

Although the magnitude of the change spatiotemporally varied both between individual rock glaciers and on different areas of the same rock glacier, the general trends, observed throughout the Alps could be confirmed for most of the studied landforms.

555 On the other hand, a few exhibited special characteristics or an atypical behaviour could be detected. RG 08 is an example of how the lower part of a rock glacier can become inactive. As flow velocities exceeded 1.5 m/yr in 1953-1971 in this area, progressively decreased until 2006 and were close to zero in the two most recent epochs. In the 2006-2017 epoch, a slight subsidence was still visible in this area, with a decreasing trend in the latest epoch. In contrast, as for most of the rock glaciers in our catchment, an acceleration of flow velocities was observed in the upper part of this rock glacier since the epoch 1997-

560 2006. Here, we assume that the topography (terminal part overflowed escarpment and reached low slope valley floor) and altitude of the front of just 2510 m in combination with the separation of the upper part favoured the inactivation of the terminal part.

RG04 showed a clearly atypical temporal course of the flow velocities. These showed low mean values of well below 0.15 m/yr, no acceleration and back creeping-movement in the upper part, which indicates that RG04 is a push-moraine. Constant or even decreasing flow velocities on rock glaciers have also been observed by Roer (2005) on a multi-decadal scale. In recent years, for example, the Aget rock glacier, which is also a push-moraine, has followed such a course (PERMOS, 2019). In contrast, RG07 and RG02 showed flow velocities barely above LoD (mean below 0.1 m/yr) in the epochs between 1953 and 1997, followed by the strongest proportional acceleration observed in the catchment. Such a behaviour of reactivation has already been described by Michelletti et al. (2015) for a small rock glacier in the Hérens Valley (Switzerland). Beside rock glacier destabilisation, the proportional velocity increase of RG07 of 485% was one of the highest compared to other studies, although its absolute speeds were not particularly high (c.f. Roer, 2005; Kellerer-Pirklbauer and Kaufmann, 2012, 2018; PERMOS, 2019). Since rock glaciers RG 02, RG 07 and RG 04 show similar exposition, size, and elevation ranges, we assume that the different behaviour might be explained by a topographic (RG04 has a significantly lower mean slope), structural control (significant volume loss and LIA glacier cover of RG04 indicates high ice content, whereas RG02 and RG07 are of TC-type and with the exception of the most recent epoch, showed no or hardly any volume loss) or change in the mechanism of flow (creation or reactivation of a sheare zone).

Another atypical development of flow velocities was observed in the case of RG05. Although a slight increase in flow velocity could already be ascertained in the terminal part during the epoch 1997 - 2006, a significant acceleration of whole rock glacier was only observed in the epoch 2012 - 2017. This delayed reaction could be related to the relatively high altitude of the rock glacier (2702 m - 3093 m) in combination with its location between two steep slopes. Although we have no annual data on this, all historical images show avalanche deposition, which have their origin in the steep adjacent slopes, over a large part of the rock glacier area (Fig. 10). This may have prevented effective heat transfer to the permafrost body of the rock glacier during the otherwise snow-free period until the last epoch, and therefore acceleration took place significantly later.

6.3 Possible implications of changes in external forcing for rock glacier flow velocities

Kenner et al. (2017) synthesise findings for external factors controlling rock glacier flow velocity. According to them, an increase in the permafrost temperature, which changes the viscosity, hardness, and shear- and crushing strength of the permafrost ice, can thus increase its internal plastic deformation. Another factor would be the increase in water availability and water pressure, which reduces the friction resistance in the shear zone. The former is primarily determined by changes in air temperature leading to changes in ground temperature and the timing and duration of snow cover. The latter can be controlled by precipitation, snowmelt, the formation of new drainage systems and melting permafrost ice. The following sections describe possible implications of changes in the meteorological forcing (Sect. 5.3) based on the development of the flow velocity for the six epochs between 1953 and 2017 (Sect. 5.4).


6.3.1 Temperature

As described by numerous studies, the development of temperatures, in particular ground surface temperatures, fits well with the development of flow velocities (e.g. Roer, 2005; Kääh et al., 2007; Delaloye et al., 2010; Scapozza et al., 2014; Hartl et al., 2016; Kenner et al., 2017; Kenner et al., 2020). Even though not covered in Fig. 12, we observed exclusively positive temperature anomalies ranging between 0.5 °C and 1°C between 1946-1951. Relatively warm temperatures were measured throughout the Alps during this period (Beniston, 2006). This could be a possible explanation for the local peak in flow velocities of RG01 and RG08 between 1953 and 1971. This phenomenon has also been observed in the Austrian Alps by Kellerer-Pirklbauer and Kaufmann (2012) and Kellerer-Pirklbauer et al. (2018) and for the Swiss Alps by Kenner et al. (2020) and is explained by decennial variations in mean annual air temperature (Delaloye et al., 2010). Since only the two rock glaciers that are located at the lowest elevation showed this reaction, this might indicate that the remaining rock glaciers have not yet reached a certain system state and have therefore hardly or not at all reacted to the increased temperatures of this period.


Looking at the strong increase in flow velocities from 1997 onwards, it appears that mainly spring and summer temperatures were likely to be responsible for the increase, as the average winter temperature actually decreased and the mean autumn temperature remained constant in the case of the sharp increase in the epoch 1997-2006. This is also supported by the fact that years with exceptional high positive winter anomalies between 1982 and 1997 did not lead to an increase in flow velocities on the studied rock glaciers. As the positive spring and summer anomalies already began in the first half of the epoch 1982-1997, while a sharp increase in flow velocities was only evident from 1997 onwards, it shows that the increase in flow velocity, if simply controlled by temperature, is slightly delayed. This might be due to a delayed warming of the permafrost ice or to the duration of the formation of new drainage systems (Kenner et al., 2017; Kenner et al., 2020), which also might explain the varying magnitude of the increases. The local peak of some rock glaciers between 1997 and 2006 could be explained by the particularly strong increase in spring temperatures or by the heatwave in the summer of 2003, which has also led to flow velocity peaks in annual studies (e.g. Kellerer-Pirklbauer and Kaufmann, 2012, 2018; PERMOS, 2019). The further increase in flow velocities in the epoch 2012-2017 could be due to the fact that, in addition to spring and summer temperatures, winter and autumn temperatures also showed exclusively positive anomalies in this time step. The low, otherwise constant or falling flow velocities in the period 1970/71 to 1982, on the other hand, fit in well with the relatively low summer and autumn temperatures.

6.3.2 Precipitation

While many studies, especially recent ones, emphasise the role of liquid water in rock glacier movement (Ikeda et al., 2008), especially in the shear horizon (Kenner et al., 2017; Cicoira et al., 2019; Kenner et al., 2020), only some show a correlation between precipitation and movement (Micheletti et al., 2015; Hartl et al., 2016; Eriksen et al., 2018), while others find no or only a weak connection (Kenner et al., 2017; Kenner et al., 2020). Kenner et al. (2020) give an increase in runoff efficiency, due to the formation of new drainage pathways in the permafrost body, as a main reason for the observed acceleration.



Rock glacier



In our study, the development of flow velocities also corresponded well with the development of precipitation. Although a direct connection cannot be proven by the qualitative analysis. In the epoch from 1953 to 1997, no clear accumulation of positive or negative anomalies can be observed in the individual time steps. In case of the epoch 1971 to 1997, the negative summer anomalies were balanced out by positive autumn anomalies and vice versa. In the epoch from 1997 to 2006 and between 2012 and 2017, positive precipitation anomalies occurred for spring as well as for summer and autumn. This did not apply to the epoch from 2006 to 2012, where only positive summer anomalies were increasingly observed. As this development fits in well with the determined flow velocities the availability of water by liquid precipitation might influence acceleration in our catchment, as suggested by Micheletti et al. (2015) for rock glaciers in the Hérens Valley (Switzerland).

640 **6.3.4 Snow Cover**

The onset and duration of snow cover have been shown to be important factors for the development of rock glacier flow velocities, as they control the time span of liquid water availability as well as the temperature in the subsurface due to the winter cooling intensity (Kenner et al., 2017; Kenner et al., 2020). As for the other two parameters, temperature and precipitation, links could be found between the temporal development of snow cover and the evolution of flow velocities in the rock glaciers studied. In the last three epochs from 1997 to 2017 and in the case of significant snow cover (>50 cm) in the last two epochs between 2006 - 2017, the snow cover set in relatively early. In combination with the amount of snow and the temperature, this could decrease the rock glacier deceleration in winter, by isolating the rock glacier from cold winter temperatures, which in turn favours warming in spring and summer (Wirz et al., 2016). This could have led to an increase in the flow velocity, especially in the most recent time period. This is also suggested as an explanation for the velocity peak in 2015, which was recorded on many rock glaciers in the Alps in annual measurements (PERMOS, 2019). Although we cannot resolve this in terms of temporal resolution, the epoch 2012 - 2017 showed higher values on all rock glaciers than 2006 - 2012 and both winters of 2013/14 and 2014/15 showed a very early development of significant snow cover, although the following winters showed a contrasting development (Fig.12).

Another factor that could be identified from the analysis of snow data is the end of snow melt, which showed a negative trend over the study period and, especially from the beginning of the 1990s, almost exclusively negative anomalies. This is of great importance for the development of the velocity of rock glaciers, as Wirz et al. (2016) show that seasonal rock glacier acceleration coincides with zero curtain during snow melt. Kenner et al. (2020) give the end of active layer freezing and start of snow melt as one of the most important factors controlling rock glacier flow velocities. Therefore, the timing of the end of snow melt in the catchment might be seen as another factor for the acceleration observed.

In the period before 1997, it was more difficult to establish a connection. This may be due to the fact that the time periods were longer and thus positive and negative anomalies balance each other out, but possibly also to the fact that the factor snow cover must always be seen in connection with the temperature, which only changed drastically from the beginning of the 1990s. This may have potentially led to the formation of new drainage systems, causing a tipping point of flow velocities to a

higher level, which in turn might have changed the value on the influence of the snow cover and precipitation on the flow
665 velocities.

6.4 Surface elevation changes in a regional context

The mean surface elevation changes ranged from $-0.047 \text{ m/yr} \pm 0.001 \text{ m/yr}$ to $0.005 \text{ m/yr} \pm 0.003 \text{ m/yr}$ for all rock glaciers during the whole period of investigation. Although, to the best of our knowledge, there are only a few studies that examine surface elevation change and volume change of rock glaciers on a comparable temporal scale, their results are in good
670 agreement with this study in terms of the magnitude of the values. For example, the Hintere Langtalkar rock glacier, located in the Hohe Tauern Range, Austria, which was partially covered by LIA glaciers, showed increasingly negative values between 0.016 m/yr and 0.058 m/yr and a sharp increase in the period 2002 - 2006 (Kellerer-Pirklbauer and Kaufmann, 2018). In terms of temporal development and magnitude, this compares well with RG01 and RG04 of this study.

Kaufmann et al. (2018) derived values of -0.0008 m/yr to -0.013 m/yr for Tschadinhorn rock glacier (Hohe Tauern Range,
675 Austria). Although we found similar values ranges in our study, the temporal development differed, since after an increase in negative values from 1953 - 2009, values very close to zero were recorded in the most recent epoch 2009 - 2015. Whereas for the rock glaciers with comparable value ranges in our study, there was either a steady increase in negative values (RG03, RG06, RG08, RG 09) or values very close to zero followed by a negative surface elevation change in the last epoch between 2012 – 2017 (RG05).

680 6.5 Interpretation and implications of surface elevation changes

Although an attempt was made to cover the entire area of the rock glaciers by analysing the surface elevation change, snow-covered areas had to be masked out for all epochs in order to increase comparability. Therefore, one cannot speak of a mass balance in the strictest sense, nevertheless, the temporal comparison of the rock glaciers allowed for interpretations to be made. The surface elevation change of rock glaciers reflects the sum of the downslope movement, the strain pattern
685 (compression/extension), compaction, debris input and aggradation/melting of ice. Therefore, surface elevation changes close to zero result either because no significant surface elevation change has occurred (e.g. RG02 1953 - 1971) or because the aforementioned processes that lead to positive and negative changes compensate each other (e.g. RG08 1953 - 1971). The only area of substantial surface elevation gains on most rock glaciers was the front indicating a rock glacier advance or the front of flow bulges, particularly pronounced on RG01 and RG06, which was also largely due to the movement of the rock glacier
690 (Frehner et al., 2015). Major debris mass input onto the rock glacier transport system in the rooting zone of the rock glacier due to larger rockfalls could not be observed. Masking out the patches of snow that often cover the root zones could however prevent such detection on some rock glaciers. These are particularly pronounced on TC-type rock glaciers RG05, RG06, RG09 and might be an indication of both snow and debris input from avalanches in the rooting zone of these rock glaciers.

Over time, there was a clear tendency towards negative values of varying degrees on the individual rock glaciers, indicating a
695 successive subsidence of the landforms at a rate of only a few cm/yr. Although we could not clearly separate the individual

processes that led to this in this study, we argue that this was mainly due to the melting of ice or loss of pore space/compaction of debris, which clearly outweighed the processes that led to positive changes.

If the negative mean surface elevation changes were mainly due to ice melt, the melt rate increased more and more on a majority of the rock glaciers during the investigated period. It is striking that the TC-type rock glaciers showed lower melt rates and that the mean surface elevation changes in three cases (RG02, RG05 and RG07) was only significantly negative in the most recent epoch. In contrast, GFC-type rock glaciers that were at least partially covered by LIA glaciers showed significantly higher values. As these rock glaciers vary in size, exposure, elevation and slope, we explain this on the one hand by the fact that GFC-type rock glaciers have no connection with an active feeding mechanism, therefore mass input can just occur by congelation of water. On the other hand, they are characterised by a high ice content and contain large amounts of massive sedimentary ice due to the LIA glacier cover, respectively. This is proven for RG01 (Hausmann et al., 2012) and RG09 (Dusik et al., 2015) by geophysical investigations and very likely due to the occurrence of thermokarst depressions on RG04 and a thermokarst lake on RG06. On the other hand, the fact that the mean surface elevation change for rock glaciers RG02, RG05 and RG07 was significantly negative only in the last epoch indicated an active feeding mechanism or ice aggregation, a low ice content or efficient isolation of the ice contained. Although the exclusion of the snow-covered areas weakens the validity of the statement to some extent, especially for RG05, RG06 and RG09, many observations indicate that the melt of ice was exceeding the feeding of the rock glaciers or there is no connection with any active feeding mechanism.

7 Conclusions

We conclude that the combination of different remote sensing techniques used in this study is well suited for the detection of flow velocity and surface elevation change of rock glaciers back to the mid of the 20th century. Thus, long-term trends in rock glacier kinematics could be identified and linked to climate changes over such a long period.

As a general result, we were able to demonstrate a significant increase in flow velocities in the epoch 1997 to 2006 and an increase in subsidence to varying degrees over the entire study period. Both observations can be explained by changes in external forcing. The sharp increase especially in spring and summer temperatures since the 1990s led to a change in the flow properties of the permafrost body due to a warming of the permafrost ice. Although thawing of the permafrost ice could not be distinguished from compaction due to a loss of pore space, trends to negative mass balances suggest a progressive thaw of the permafrost body throughout the study period, which is more pronounced on GFC rock glaciers. Furthermore, the melting of the ice body might create new drainage systems. This results in more water being available to the system, which is crucial for horizontal movement in shear zones. Flow velocity in this catchment area could also be linked to changes in precipitation pattern, which again governs water availability and the onset and duration of snow cover, which controls the time span of liquid water availability as well as the temperature in the subsurface due to the winter cooling intensity.

Although we were able to identify a general trend in rock glacier kinematics, the catchment wide view also showed a slightly different response of individual rock glaciers to similar external forcing regarding timing, magnitude and local temporal peaks.

No characteristic could be identified that explains the different responses to external forcing over the entire study period. The different behaviour could be explained by different sensitivity, response or response time of individual rock glaciers to intra-
730 annual, inter-annual or multi-annual fluctuations and changes in external forcing parameters. For some rock glaciers internal structure and topography might explain different reactions, as two rock glaciers of similar size, exposition, elevation and elevation range showed contrasting reactions of reactivation and constant flow velocities.

In addition, we were able to detect a rock glacier which shows an inactivation of its terminal part. By observing the surface elevation changes of all rock glaciers we conclude that the rates of rock glacier mass transport and volumetric losses of the
735 rock glacier due to ice melt are much higher than debris and ice input, at least in those cases where an almost complete analysis of the entire rock glacier surface was possible. Therefore, the investigated rock glaciers are prone to starvation which will eventually lead to rock glacier inactivation. However, since the current ice content of the rock glaciers is not known and the analysis of future flow rates, melt rates and ice and debris input was not the aim of this study, it is not possible to predict the timing of this inactivation.

We can also conclude that future studies are necessary to better understand the climate forcing on rock glacier
740 morphodynamics. Therefore, the analysis should be transferred to other catchments in order to identify differences and similarities within the Alps. Surface elevation change and flow velocity analysis should be combined with downscaled reanalysis data to better understand catchment wide differences in external forcing on a longer timescale. If possible, future studies should combine borehole measurements or geophysical investigation to shed light on the internal structure of rock
745 glaciers and clarify some of the assumptions made in this study and possible explanations of their behavior.

As a final important consideration, historical terrestrial images (if available) should be used with monoplottting tools. Mapping on such images would help to shorten the time span of the individual epochs which is crucial to better differentiate the influence of individual forcing parameters, as it is very likely that there are changes within our analysed epochs. Beside this, historical
750 terrestrial images would offer the opportunity to expand the analysis back to the 19th century and thus closer to the LIA in order to study an important period in terms of massive system changes in the glacial and periglacial regions of the Alps.

Code availability

The image correlation algorithm (IMCORR) is used for the calculation of the rock glacier flow velocities is implemented in the open source geoinformation system SAGA GIS. Furthermore, some modules of the commercial SAGA GIS extension
755 SAGA LIS PRO 3D were used to calculate the surface elevation change of the rock glaciers. The software which was used to create digital elevation models and orthoimagess from historical aerial images was the commercial software Agisoft Metashape.

Data availability

760 The analysed metrological data is available from the “Federal Ministry of Agriculture, Regions and Tourism“ (BMLRT), the “Central Institute for Meteorology and Geodynamics“ (ZAMG), the “Historical Instrumental Climatological Surface Time Series of the Greater Alpine Region“ (HISTALP), the “Autonomous Province of Bozen/Bolzano“ and “Tyrolean Hydropower AG “ (TIWAG). The aerial images used to create digital elevation models and orthoimages are available from the "Office of the Tyrolean Government-Department of Geoinformation" (<https://www.tirol.gv.at/en/>) and the "Austrian Federal Office of Surveying and Metrology" (BEV) (<https://www.bev.gv.at>). The self-collected ALS data will presumably be made available after completion of the SEHAG (“Sensitivity of High Alpine Geosystems to Climate Change Since 1850”) research project.

765 Author Contributions

Conceptualization, FF, FH and MB; data curation, FF, LP, MA, JR, MP, MS, FH and MW; formal analysis, FF, FH, MP; funding acquisition, MB, FH, TH and NP; investigation, FF, FH; methodology, FF, FH, LP, MP, TH, MA, JR, MS, MW, NP and MB; project administration, MB, FH, TB and NP; supervision, FH, TH and MB; writing—original draft, FF, FH; writing—review and editing, FF, FH, LP, MP, MA, JR, MW, NP. All authors have read and agreed to the published version of the
770 manuscript.

Conflicts of Interest

The authors declare that they have no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Acknowledgments

775 The study was part of the SEHAG project (“Sensitivity of High Alpine Geosystems to Climate Change Since 1850”) and financially supported by the German Research Foundation (DFG) and the Austrian ScienceFund (FWF). We gratefully acknowledge the DFG and the FWF for the financial support. We would also like to acknowledge the Tyrolean Hydropower AG (TIWAG, Austria, Innsbruck), the Federal Ministry of Agriculture, Regions and Tourism (BMLRT, Austria Vienna), Central Institute for Meteorology and Geodynamics (ZAMG, Austria, Vienna) and the Autonomous Province of
780 Bozen/Bolzano for providing the meteorological data. We would like to acknowledge the Office of the Tyrolean Government (Department of Geoinformation, Austria, Innsbruck) for providing the historical images 1970 and 1971 as well as the corresponding camera-calibration certificates. We would also like to thank the Austrian Federal Office of Surveying and Metrology (BEV, Austria, Vienna) for providing the historical aerial images of 1953/54, 1982 and 1997. We acknowledge the

Kaunertaler Gletscherbahnen GmbH for the free use of the toll road in the UpperKauner Valley. The open access publication
785 of this article was supported by the Open Access Fund of the Catholic University of Eichstätt-Ingolstadt.

References

- Altmann, M., Piermattei, L., Haas, F., Heckmann, T., Fleischer, F., Rom, J., Betz-Nutz, S., Knoflach, B., Müller, S.,
Ramskogler, K., Pfeiffer, M., Hofmeister, F., Ressler, C., and Becht, M.: Long-Term Changes of Morphodynamics on
Little Ice Age Lateral Moraines and the Resulting Sediment Transfer into Mountain Streams in the Upper Kauner
790 Valley, Austria, *Water*, 12, 3375, <https://doi.org/10.3390/w12123375>, 2020.
- Anderson, S. W.: Uncertainty in quantitative analyses of topographic change: error propagation and the role of thresholding,
Earth Surf. Process. Landforms, 44, 1015–1033, <https://doi.org/10.1002/esp.4551>, 2019.
- Arenson, L., Hoelzle, M., and Springman, S.: Borehole deformation measurements and internal structure of some rock
glaciers in Switzerland, *Permafrost Periglac. Process.*, 13, 117–135, <https://doi.org/10.1002/ppp.414>, 2002.
- 795 Bakker, M. and Lane, S. N.: Archival photogrammetric analysis of river-floodplain systems using Structure from Motion
(SfM) methods, *Earth Surf. Process. Landforms*, 42, 1274–1286, <https://doi.org/10.1002/esp.4085>, 2017.
- Barsch, D.: *Rockglaciers: Indicators for the Present and Former Geoecology in High Mountain Environments*, Springer
Series in Physical Environment, 16, Springer Berlin Heidelberg, Berlin, Heidelberg, 331 pp., 1996.
- Beniston, M.: Mountain Weather and Climate: A General Overview and a Focus on Climatic Change in the Alps,
800 *Hydrobiologia*, 562, 3–16, <https://doi.org/10.1007/s10750-005-1802-0>, 2006.
- Berger, J., Krainer, K., and Mostler, W.: Dynamics of an active rock glacier (Ötztal Alps, Austria), *Quat. res.*, 62, 233–242,
<https://doi.org/10.1016/j.yqres.2004.07.002>, 2004.
- Berthling, I.: Beyond confusion: Rock glaciers as cryo-conditioned landforms, *Geomorphology*, 131, 98–106,
<https://doi.org/10.1016/j.geomorph.2011.05.002>, 2011.
- 805 Besl, P. J. and McKay, N. D.: A method for registration of 3-D shapes, *IEEE Trans. Pattern Anal. Mach. Intell.*, 14, 239–
256, <https://doi.org/10.1109/34.121791>, 1992.
- Brardinoni, F., Scotti, R., Sailer, R., and Mair, V.: Evaluating sources of uncertainty and variability in rock glacier
inventories, *Earth Surf. Process. Landforms*, <https://doi.org/10.1002/esp.4674>, 2019.
- Buchli, T., Kos, A., Limpach, P., Merz, K., Zhou, X., and Springman, S. M.: Kinematic investigations on the
810 Furggwanhorn Rock Glacier, Switzerland, *Permafrost and Periglac Process*, 29, 3–20,
<https://doi.org/10.1002/ppp.1968>, 2018.
- Buckel, J. and Otto, J.-C.: The Austrian Glacier Inventory GI 4 (2015) in ArcGis (shapefile) format, supplement to: Buckel,
Johannes; Otto, Jan-Christoph; Prasicsek, Günther; Keuschnig, Markus (2018): Glacial lakes in Austria - Distribution and
formation since the Little Ice Age. *Global and Planetary Change*, 164, 39-51, 2018.

- 815 Cicoira, A., Beutel, J., Faillettaz, J., and Vieli, A.: Water controls the seasonal rhythm of rock glacier flow, *Earth and Planetary Science Letters*, 528, 115844, <https://doi.org/10.1016/j.epsl.2019.115844>, 2019.
- Clark, D. H., Steig, E. J., Potter, J. N., and Gillespie, A. R.: Genetic variability of rock glaciers, *Geografiska Annaler: Series A, Physical Geography*, 80, 175–182, <https://doi.org/10.1111/j.0435-3676.1998.00035.x>, 1998.
- Delaloye, R., Lambiel, C., and Gärtner-Roer, I.: Overview of rock glacier kinematics research in the Swiss Alps, *Geogr. Helv.*, 65, 135–145, <https://doi.org/10.5194/gh-65-135-2010>, 2010.
- 820 Delaloye, R., Perruchoud, E., Bodin, X., Avian, M., Hausmann, H., Kellerer-Pirklbauer, A., Kaufmann, V., Mihajlovic, D., Lambiel, C., Ikeda, A., Roer, I., Krainer, K., Käab, A., Staub, B., and Thibert, E.: Recent interannual variations of rock glacier creep in the European Alps, s.n., 7 pp., <https://doi.org/10.5167/UZH-7031>, 2008.
- Dusik, J.-M., Leopold, M., Heckmann, T., Haas, F., Hilger, L., Morche, D., Neugirg, F., and Becht, M.: Influence of glacier advance on the development of the multipart Riffeltal rock glacier, Central Austrian Alps, *Earth Surf. Process. Landforms*, 40, 965–980, <https://doi.org/10.1002/esp.3695>, 2015.
- 825 Eriksen, H. Ø., Rouyet, L., Lauknes, T. R., Berthling, I., Isaksen, K., Hindberg, H., Larsen, Y., and Corner, G. D.: Recent Acceleration of a Rock Glacier Complex, Ádjet, Norway, Documented by 62 Years of Remote Sensing Observations, *Geophys. Res. Lett.*, 45, 8314–8323, <https://doi.org/10.1029/2018GL077605>, 2018.
- 830 Fawcett, D., Blanco-Sacristán, J., and Benaud, P.: Two decades of digital photogrammetry: Revisiting Chandler’s 1999 paper on “Effective application of automated digital photogrammetry for geomorphological research” – a synthesis, *Progress in Physical Geography: Earth and Environment*, 43, 299–312, <https://doi.org/10.1177/0309133319832863>, 2019.
- Fey, C. and Krainer, K.: Analyses of UAV and GNSS based flow velocity variations of the rock glacier Lazaun (Ötztal Alps, South Tyrol, Italy), *Geomorphology*, 365, 107261, <https://doi.org/10.1016/j.geomorph.2020.107261>, 2020.
- 835 Finsterwalder, S.: Begleitworte zur Karte des Gepatschferners, *Zeitschrift für Gletscherkunde*, XVI, 20–41, 1928.
- Fischer, A., Seiser, B., Stocker-Waldhuber, M., Mitterer, C., and Abermann, J.: The Austrian Glacier Inventories GI 1 (1969), GI 2 (1998), GI 3 (2006), and GI LIA in ArcGIS (shapefile) format, supplement to: Fischer, Andrea; Seiser, Bernd; Stocker-Waldhuber, Martin; Mitterer, Christian; Abermann, Jakob (2015): Tracing glacier changes in Austria from the Little Ice Age to the present using a lidar-based high-resolution glacier inventory in Austria. *The Cryosphere*, 9(2), 753-766, 2015.
- 840 Fliri, F.: *Das Klima der Alpen im Raume von Tirol*, Monographien zur Landeskunde Tirol, Folge 1, Universitätsverlag, Innsbruck, 454 pp., 1975.
- Frehner, M., Ling, A. H. M., and Gärtner-Roer, I.: Furrow-and-Ridge Morphology on Rockglaciers Explained by Gravity-Driven Buckle Folding: A Case Study From the Murtèl Rockglacier (Switzerland), *Permafrost and Periglac Process*, 26, 57–66, <https://doi.org/10.1002/ppp.1831>, 2015.
- 845

- Glira, P., Pfeifer, N., Briese, C., and Ressler, C.: A Correspondence Framework for ALS Strip Adjustments based on Variants of the ICP Algorithm
Korrespondenzen für die ALS-Streifenausgleichung auf Basis von ICP, Photogrammetrie - Fernerkundung - Geoinformation, 2015, 275–289, <https://doi.org/10.1127/pfg/2015/0270>, 2015.
- 850 Groh, T. and Blöthe, J. H.: Rock Glacier Kinematics in the Kaunertal, Ötztal Alps, Austria, *Geosciences*, 9, 373, <https://doi.org/10.3390/geosciences9090373>, 2019.
- Haeberli, W., Hallet, B., Arenson, L., Elconin, R., Humlum, O., Kääb, A., Kaufmann, V., Ladanyi, B., Matsuoka, N., Springman, S., and Mühll, D. V.: Permafrost creep and rock glacier dynamics, *Permafrost Periglac. Process.*, 17, 189–214, <https://doi.org/10.1002/ppp.561>, 2006.
- 855 Hartl, L., Fischer, A., Stocker-waldhuber, M., and Abermann, J.: Recent speed-up of an alpine rock glacier: an updated chronology of the kinematics of outer hochebenkar rock glacier based on geodetic measurements, *Geografiska Annaler: Series A, Physical Geography*, 98, 129–141, <https://doi.org/10.1111/geoa.12127>, 2016.
- Hausmann, H., Krainer, K., Brückl, E., and Ullrich, C.: Internal structure, ice content and dynamics of Ölgrube and Kaiserberg rock glaciers (Ötztal Alps, Austria) determined from geophysical surveys., *AJES*, 105, 12–31, 2012.
- 860 Heid, T. and Kääb, A.: Evaluation of existing image matching methods for deriving glacier surface displacements globally from optical satellite imagery, *Remote Sensing of Environment*, 118, 339–355, <https://doi.org/10.1016/j.rse.2011.11.024>, 2012.
- Hock, R., Rasul, G., Adler, C., Caceres, S., Gruber, S., Hirabayashi, Y., Jackson, M., Kääb, A., Kang, S., Kutuzov, S., Milner, A., Molau, U., Morin, S., Orlove, B., and Steltzer, H.: High Mountain Areas, in: *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*, edited by: Pörtner, H.-O., Roberts, D.C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., Mintenbeck, K., Alegria, A., Nicolai, M., Okem, A., Petzold, J., Rama, B., and Weyer, N.M., 2019.
- 865 Hoinkes, G. and Thöni, M.: Evolution of the Ötztal-Stubai, Scarl-Campo and Ulten Basement Units, in: *Pre-Mesozoic Geology in the Alps*, edited by: Raumer, J. F. von and Neubauer, F., Springer Berlin Heidelberg, Berlin, Heidelberg, 485–494, https://doi.org/10.1007/978-3-642-84640-3_29, 1993.
- 870 Ikeda, A., Matsuoka, N., and Kääb, A.: Fast deformation of perennially frozen debris in a warm rock glacier in the Swiss Alps: An effect of liquid water, *J. Geophys. Res.*, 113, 212, <https://doi.org/10.1029/2007JF000859>, 2008.
- Jones, D. B., Harrison, S., Anderson, K., and Whalley, W. B.: Rock glaciers and mountain hydrology: A review, *Earth-Science Reviews*, 193, 66–90, <https://doi.org/10.1016/j.earscirev.2019.04.001>, 2019.
- 875 Kääb, A. and Vollmer, M.: Surface Geometry, Thickness Changes and Flow Fields on Creeping Mountain Permafrost: Automatic Extraction by Digital Image Analysis, *Permafrost Periglac. Process.*, 11, 315–326, [https://doi.org/10.1002/1099-1530\(200012\)11:4<315:AID-PPP365>3.0.CO;2-J](https://doi.org/10.1002/1099-1530(200012)11:4<315:AID-PPP365>3.0.CO;2-J), 2000.
- Kääb, A., Chiarle, M., Raup, B., and Schneider, C.: Climate change impacts on mountain glaciers and permafrost, *Global and Planetary Change*, 56, vii–ix, <https://doi.org/10.1016/j.gloplacha.2006.07.008>, 2007.

- 880 Kääh, A., Strozzi, T., Bolch, T., Caduff, R., Trefall, H., Stoffel, M., and Kokarev, A.: Inventory, motion and acceleration of rock glaciers in Ile Alatau and Kungöy Ala-Too, northern Tien Shan, since the 1950s, 37 pp., 2020.
- Kaufmann, V., Seier, G., Sulzer, W., Wecht, M., Liu, Q., Lauk, G., and Maurer, M.: Rock glacier monitoring using aerial photographs: conventional vs. UAV-based Mapping - a comparative study, *Int. Arch. Photogramm. Remote Sens. Spatial Inf. Sci.*, XLII-1, 239–246, <https://doi.org/10.5194/isprs-archives-XLII-1-239-2018>, 2018.
- 885 Kaufmann, V. and Kellerer-Pirklbauer, A.: Regional quantification of rock glacier movement in Austria using governmental GIS data, in: *Geomorphometry for geosciences: [this vol. is a contribution to the 4th International Conference on Geomorphometry ; Geomorphometry 2015: Conference and Workshops ; Geomorphometry for natural hazards geomodelling, Poznan (Poland), June 22 - 26, 2015]*, edited by: Jasiewicz, J., 165–168, 2015.
- Kaufmann, V., Sulzer, W., Seier, G., and Wecht, M.: Panta Rhei: Movement Change of Tschadinhorn Rock Glacier (Hohe
890 Tauern Range, Austria), 1 954–201 7, *Kartogr. geoinf. (Online)*, 18, 4–24, <https://doi.org/10.32909/kg.18.31.1>, 2019.
- Kellerer-Pirklbauer, A. and Kaufmann, V.: Deglaciation and its impact on permafrost and rock glacier evolution: New insight from two adjacent cirques in Austria, *The Science of the total environment*, 621, 1397–1414, <https://doi.org/10.1016/j.scitotenv.2017.10.087>, 2018.
- Kellerer-Pirklbauer, A. and Kaufmann, V.: About the relationship between rock glacier velocity and climate parameters in
895 central Austria, *AJES*, 105, 94–112, 2012.
- Kellerer-Pirklbauer, A., Lieb, G. K., and Kaufmann, V.: The Dösen Rock Glacier in Central Austria: A key site for multidisciplinary long-term rock glacier monitoring in the Eastern Alps, *AJES*, 110, <https://doi.org/10.17738/ajes.2017.0013>, 2018.
- Kenner, R., Pruessner, L., Beutel, J., Limpach, P., and Phillips, M.: How rock glacier hydrology, deformation velocities and
900 ground temperatures interact: Examples from the Swiss Alps, *Permafrost and Periglac Process*, 31, 3–14, <https://doi.org/10.1002/ppp.2023>, 2020.
- Kenner, R., Phillips, M., Beutel, J., Hiller, M., Limpach, P., Pointner, E., and Volken, M.: Factors Controlling Velocity Variations at Short-Term, Seasonal and Multiyear Time Scales, Ritigraben Rock Glacier, Western Swiss Alps, *Permafrost and Periglac. Process.*, 28, 675–684, <https://doi.org/10.1002/ppp.1953>, 2017.
- 905 Krainer, K. and Ribis, M.: A rock glacier inventory of the Tyrolean Alps (Austria), *Austrian Journal of Earth Sciences*, 105, 32–57, 2012.
- Krainer, K. and Mostler, W.: Flow velocities of active rock glaciers in the Austrian Alps, *Geografiska Annaler: Series A, Physical Geography*, 88 A, 267–280, 2006.
- Krainer, K., Bressan, D., Dietre, B., Haas, J. N., Hajdas, I., Lang, K., Mair, V., Nickus, U., Reidl, D., Thies, H., and
910 Tonidandel, D.: A 10,300-year-old permafrost core from the active rock glacier Lazaun, southern Ötztal Alps (South Tyrol, northern Italy), *Quat. res.*, 83, 324–335, <https://doi.org/10.1016/j.yqres.2014.12.005>, 2015.
- Kummert, M. and Delaloye, R.: Mapping and quantifying sediment transfer between the front of rapidly moving rock glaciers and torrential gullies, *Geomorphology*, 309, 60–76, <https://doi.org/10.1016/j.geomorph.2018.02.021>, 2018.

- 915 Marcer, M., Cicoira, A., Cusicanqui, D., Bodin, X., Echelard, T., Obregon, R., and Schoeneich, P.: Rock glaciers throughout the French Alps accelerated and destabilised since 1990 as air temperatures increased, *Commun Earth Environ*, 2, 383, <https://doi.org/10.1038/s43247-021-00150-6>, 2021.
- Micheletti, N., Lambiel, C., and Lane, S. N.: Investigating decadal-scale geomorphic dynamics in an alpine mountain setting, *J. Geophys. Res. Earth Surf.*, 120, 2155–2175, <https://doi.org/10.1002/2015JF003656>, 2015.
- 920 Monnier, S. and Kinnard, C.: Pluri-decadal (1955–2014) evolution of glacier–rock glacier transitional landforms in the central Andes of Chile (30–33° S), *Earth Surf. Dynam.*, 5, 493–509, <https://doi.org/10.5194/esurf-5-493-2017>, 2017.
- Olefs, M., Koch, R., Schöner, W., and Marke, T.: Changes in Snow Depth, Snow Cover Duration, and Potential Snowmaking Conditions in Austria, 1961–2020—A Model Based Approach, *Atmosphere*, 11, 1330, <https://doi.org/10.3390/atmos11121330>, 2020.
- 925 Otto, J.-C., Schrott, L., and Keller, F.: Map of permafrost distribution for Austria, Europe, 2020.
- Paul, F., Bolch, T., Briggs, K., Kääb, A., McMillan, M., McNabb, R., Nagler, T., Nuth, C., Rastner, P., Strozzi, T., and Wuite, J.: Error sources and guidelines for quality assessment of glacier area, elevation change, and velocity products derived from satellite data in the Glaciers_cci project, *Remote Sensing of Environment*, 203, 256–275, <https://doi.org/10.1016/j.rse.2017.08.038>, 2017.
- 930 Peng, S., Piao, S., Ciais, P., Friedlingstein, P., Zhou, L., and Wang, T.: Change in snow phenology and its potential feedback to temperature in the Northern Hemisphere over the last three decades, *Environ. Res. Lett.*, 8, 14008, <https://doi.org/10.1088/1748-9326/8/1/014008>, 2013.
- PERMOS: Permafrost in Switzerland 2014/2015 to 2017/2018, Swiss Permafrost Monitoring Network, Glaciological Report (Permafrost), No. 16–19, 2019.
- 935 Pillewizer, W.: Untersuchungen an Blockstrfmen der Ötztaler Alpen, *Geomorphologische Abhandlungen des Geographischen Institutes derFU Berlin (Otto-Maull-Festschrift)*, 5, 37–50, 1957.
- Rangwala, I. and Miller, J. R.: Climate change in mountains: a review of elevation-dependent warming and its possible causes, *Climatic Change*, 114, 527–547, <https://doi.org/10.1007/s10584-012-0419-3>, 2012.
- Ravanel, L., Magnin, F., and Deline, P.: Impacts of the 2003 and 2015 summer heatwaves on permafrost-affected rock-walls 940 in the Mont Blanc massif, *The Science of the total environment*, 609, 132–143, <https://doi.org/10.1016/j.scitotenv.2017.07.055>, 2017.
- Roer, I., Haeberli, W., Avian, M., Kaufmann, V., Delaloye, R., Lambiel, C., and Kääb, A.: Observations and considerations on destabilizing active rock glaciers in the European Alps, 9th International Conference on Permafrost, Fairbanks, Alaska, 29 June 2008 - 03 July 2008, 1505–1510, 2008.
- 945 Roer, I.: Rockglacier kinematics in a high mountain geosystem, Dissertation, Mathematisch-Naturwissenschaftliche Fakultät, Rheinische Friedrich-Wilhelms-Untiversität Bonn, Bonn, 263 pp., 2005.

- Roer, I., Kääh, A., and Dikau, R.: Rockglacier acceleration in the Turtmann valley (Swiss Alps): Probable controls, *Norsk Geografisk Tidsskrift - Norwegian Journal of Geography*, 59, 157–163, <https://doi.org/10.1080/00291950510020655>, 2005.
- 950 Scambos, T.A., Dutkiewicz, M.J., Wilson, J.C., and Bindschadler, R.A.: Application of image cross-correlation to the measurement of glacier velocity using satellite image data, *Remote Sensing of Environment*, 42, 177–186, 1992.
- Scapozza, C., Lambiel, C., Bozzini, C., Mari, S., and Conedera, M.: Assessing the rock glacier kinematics on three different timescales: a case study from the southern Swiss Alps, *Earth Surf. Process. Landforms*, 39, 2056–2069, <https://doi.org/10.1002/esp.3599>, 2014.
- 955 Schoeneich, P., Bodin, X., Echelard, T., Kaufmann, V., Kellerer-Pirklbauer, A., Krysiecki, J.-M., and Lieb, G. K.: Velocity Changes of Rock Glaciers and Induced Hazards, in: *Engineering Geology for Society and Territory - Volume 1*, edited by: Lollino, G., Manconi, A., Clague, J., Shan, W., and Chiarle, M., Springer International Publishing, Cham, 223–227, https://doi.org/10.1007/978-3-319-09300-0_42, 2015.
- Scotti, R., Crosta, G. B., and Villa, A.: Destabilisation of Creeping Permafrost: The Plator Rock Glacier Case Study (Central
960 Italian Alps), *Permafrost and Periglac. Process.*, 28, 224–236, <https://doi.org/10.1002/ppp.1917>, 2017.
- Vivero, S. and Lambiel, C.: Monitoring the crisis of a rock glacier with repeated UAV surveys, *Geogr. Helv.*, 74, 59–69, <https://doi.org/10.5194/gh-74-59-2019>, 2019.
- Wirz, V., Gruber, S., Purves, R. S., Beutel, J., Gärtner-Roer, I., Gubler, S., and Vieli, A.: Short-term velocity variations at
965 three rock glaciers and their relationship with meteorological conditions, *Earth Surf. Dynam.*, 4, 103–123, <https://doi.org/10.5194/esurf-4-103-2016>, 2016.