

Geometric Controls of Tidewater Glacier Dynamics

Thomas Frank^{1,2}, Henning Åkesson^{3,4}, Basile de Fleurian¹, Mathieu Morlighem^{5,6}, and Kerim H. Nisancioglu^{1,7}

¹Department of Earth Science, University of Bergen and Bjercknes Centre for Climate Research, Bergen, Norway

²Department of Earth Sciences, Uppsala University, Sweden

³Department of Geological Sciences, Stockholm University, Sweden

⁴Bolin Centre for Climate Research, Stockholm, Sweden

⁵Department of Earth Sciences, Dartmouth College, Hanover NH, USA

⁶Department of Earth System Science, University of California, Irvine CA, USA

⁷Centre for Earth Evolution and Dynamics, University of Oslo, Oslo, Norway

Correspondence: Thomas Frank (thomas.frank@geo.uu.se)

Abstract. Retreat of marine outlet glaciers often initiates depletion of inland ice through dynamic adjustments of the upstream glacier. The local topography of a fjord may promote or inhibit such retreat, and therefore fjord geometry constitutes a critical control on ice sheet mass balance. To quantify the processes of ice-topography interactions and enhance the understanding of the dynamics involved, we analyze a multitude of topographic fjord settings and scenarios using the Ice-sheet and Sea-level System Model (ISSM). We systematically study glacier retreat through a variety of artificial fjord geometries and quantify the modeled dynamics directly in relation to topographic features. We find that retreat in an upstream widening or deepening fjord does not necessarily promote retreat, as suggested by previous studies. Conversely, it may stabilize a glacier because converging ice flow towards a constriction enhances lateral shear stress gradients. An upstream narrowing or shoaling fjord, in turn, may promote retreat since fjord walls or bed provide little stability to the glacier where ice flow diverges. Furthermore, we identify distinct quantitative relationships directly linking grounding line discharge and retreat rate to fjord topography, and transfer these results to a long-term study of the retreat of Jakobshavn Isbræ. These findings offer new perspectives on ice-topography interactions, and give guidance to an ad-hoc assessment of future topographically induced ice loss based on knowledge of the upstream fjord geometry.

Copyright statement. ©2021 all rights reserved

15 1 Introduction

Rates of ice discharge from the Greenland Ice Sheet are likely to exceed their Holocene (last 12,000 yrs) maxima this century (Briner et al., 2020; Kajanto et al., 2020), and parts of Antarctica are on the brink of irreversible mass loss (Garbe et al., 2020). Consequently, major natural and societal challenges related to changes in the terrestrial cryosphere of the high latitudes

lay ahead. An advanced understanding of the underlying processes of ice loss is paramount for fact-based decision making
20 (Oppenheimer et al., 2019).

About half of the current mass loss over Greenland (30 to 70%) is due to dynamic ice discharge related to thinning, speed-up and increased calving of outlet glaciers (Nick et al., 2009; Felikson et al., 2017; Haubner et al., 2018; Mouginitot et al., 2019; King et al., 2020). In Antarctica, dynamic instability of the West Antarctic Ice Sheet is considered a major driver of future sea-level rise (Pattyn and Morlighem, 2020), but there is also emerging evidence of changes in ice dynamics at some glaciers in
25 East Antarctica (Brancato et al., 2020; Miles et al., 2020). While outlet glaciers therefore are critical to ice-sheet mass balance and associated sea-level rise, considerable knowledge gaps on the processes governing their dynamics still exist.

Despite the general warming trend observed over the recent decades, we do not observe an overall synchronous pattern in outlet glacier evolution. This is clear for various settings in the Arctic, such as in Greenland (Warren and Glasser, 1992; Carr et al., 2013; Bunce et al., 2018; Catania et al., 2018), Svalbard (Schuler et al., 2020), Novaya Zemlya (Carr et al., 2014) and
30 North America (McNabb and Hock, 2014), as well as in Antarctica (Pattyn and Morlighem, 2020). Even adjacent glaciers with similar climatic and oceanic conditions can show strongly different behaviour (Carr et al., 2013; Catania et al., 2018; Bunce et al., 2018). The main proposed explanation is that differing bathymetry and glacier geometry significantly modulate glacier response to climate over a range of time scales (Warren and Glasser, 1992; Briner et al., 2009; Jamieson et al., 2012; Åkesson et al., 2018a; Catania et al., 2018). Broadly, there is a consensus that wide and deep parts of a fjord promote retreat,
35 while narrow and shallow areas tend to stabilize glacier termini. Moreover, kinematic wave theory indicates that the upstream propagation of a thinning signal is heavily influenced by bed topography (Felikson et al., 2017, 2021). Modelling of idealized settings (Enderlin et al., 2013; Åkesson et al., 2018b) and theoretical studies based on analytical calculations and numerical experiments further emphasize the potential of fjord geometry to modulate glacier retreat (Weertman, 1974; Raymond, 1996; Vieli et al., 2001; Schoof, 2007; Pfeffer, 2007; Gudmundsson et al., 2012; Gudmundsson, 2013).

40 It is therefore critical to advance our understanding of the influence of fjord geometry on glacier retreat to accurately predict sea-level rise, especially when extrapolating observations of a few well-monitored glaciers to those less studied (Nick et al., 2009). This knowledge is also pivotal to correctly infer past climate signals from glacier proximal landforms, because their formation may have been influenced by fjord topography and may not necessarily have been in equilibrium with the prevailing climate (Åkesson et al., 2018b; Steiger et al., 2018).

45 The most important suggested mechanisms behind geometric control of glacier retreat are: 1) friction, with glaciers in narrow fjords and grounded well above flotation being stabilized by fjord geometry, while the opposite is the case for glaciers in wide fjords and close to flotation (Raymond, 1996; Pfeffer, 2007; Enderlin et al., 2013; Åkesson et al., 2018b). Buttressing and lateral shear between an ice shelf and nearby islands and/or fjord walls can be an important factor as well (Gudmundsson et al., 2012; Gudmundsson, 2013; Jamieson et al., 2012, 2014); 2) area exposed to ocean melt, where a wider/deeper fjord induces
50 a larger cumulative melt flux (Straneo et al., 2013; Åkesson et al., 2018b); 3) the marine ice sheet instability (MISI), which is a feedback mechanism between increasing driving stress with increasing ice thickness at the grounding line (GL), where inland-sloping (retrograde) beds lead to self-accelerating ice loss (Weertman, 1974; Schoof, 2007).

While the main controls of ice-topography interactions are known, a quantitative understanding is still largely missing, especially on timescales beyond a few decades. In this context, in-situ observations of ice dynamics do not cover the full spectrum of ice-topography interactions, because they are limited in space and time. While remotely-sensed observations of ice dynamics over the past decades exist, the recent retreat in Greenland and elsewhere over this period is too short to allow for a complete assessment of geometry-glacier interactions (Carr et al., 2013; Catania et al., 2018; Bunce et al., 2018). In contrast, on paleo-time scales, retreat has occurred over large distances, but the temporal resolution of geomorphological studies is limited by the available geological data and key information is missing to discern different drivers of glacier retreat (Briner et al., 2009; Åkesson et al., 2020). Meanwhile, numerical studies that can address these issues have so far mostly used width- and depth-integrated flowline models, which carry many assumptions that do not hold in some settings (Nick et al., 2009; Enderlin et al., 2013; Åkesson et al., 2018b; Steiger et al., 2018). In particular, they parameterize or do not account for factors that are thought to be instrumental to explain ice-topography interactions, such as lateral drag, across-flow variations in glacier characteristics and viscosity changes due to variations in ice temperature. The latter, for instance, was found to be key to explain Jakobshavn Isbræ's recent retreat (Bondzio et al., 2017).

Here, we use a numerical ice-flow model resolving two horizontal dimensions, we include a suite of 21 experiments and present a systematic approach to compare the relative importance of basal and lateral fjord topography. This setup allows to assess how fjord topography controls glacier retreat on interannual to centennial time scales. We hypothesize that there are quantifiable relationships between glacier retreat and topography that apply to a wide range of glaciological settings. Such general relationships would yield substantial predictive power for a broad assessment of expected future outlet glacier retreat.

We create a large ensemble of artificial fjords that include geometric features (referred to as 'perturbations' in the following) typically found in glacial fjords, such as sills and overdeepenings (referred to as 'bumps' and 'depressions', respectively; together 'basal perturbations') as well as narrow and wide fjord sections (referred to as 'bottlenecks' and 'embayments', respectively; together 'lateral perturbations'). We then force synthetic glaciers to retreat through this variety of fjords by increasing ocean-induced melt rates, and assess key retreat metrics such as the grounding line retreat rate. The ice dynamics of each simulation are compared and quantitatively linked to the characteristics of the respective fjord geometry. Finally, we investigate whether the retreat dynamics and ice-topography interactions are transferable from the idealized setup to a long-term study on Jakobshavn Isbræ in western Greenland.

2 Methods

2.1 Ice sheet model

We use the Ice-sheet and Sea-level System Model (ISSM; Larour et al., 2012) with the shallow-shelf approximation (SSA; Morland, 1987; MacAyeal, 1989). A discussion on the appropriateness of this approximation compared to a Full-Stokes model is provided in sect. 4.2. Our domain is rectangular (80 km×10 km) with x and y representing the along-flow and across-flow coordinates, respectively (Fig. 1a). We create an unstructured mesh with a fixed resolution of 100 m close to the GL, comparable to other high-resolution studies of Greenlandic fjords (e.g. Morlighem et al., 2016, 2019). The temporal resolution

is set to $\Delta t = 0.01$ yrs (3.65 days) to satisfy the Courant-Friedrichs-Levy condition (Courant et al., 1928). We apply a sub-element GL migration scheme (Seroussi et al., 2014) and enable a moving calving front with the level-set method (Bondzio et al., 2016). We use a thermal model to account for the feedback between frictional heating and ice viscosity. The spin-up ice temperature is -5°C , representative of conditions in South/Central Greenland and the Southern Fennoscandian ice sheet (Nick et al., 2013; Åkesson et al., 2018a). To simulate calving, a von-Mises law is used (Morlighem et al., 2016), where the calving rate c is given as:

$$c = \|\mathbf{v}\| \frac{\tilde{\sigma}}{\sigma_{max}}, \quad (1)$$

where \mathbf{v} is the velocity vector, $\tilde{\sigma}$ a scalar quantity representing the tensile stress at the ice front, and σ_{max} is a stress threshold. This formulation, demonstrated to perform well in Greenlandic fjords (Choi et al., 2018), means that calving occurs when the tensile stress at the glacier front exceeds a fixed threshold. σ_{max} generally needs to be determined experimentally for studies on real-world glaciers. Here we fix σ_{max} to 1 MPa for grounded ice and 200 kPa for floating ice. These values yield a representative setup and are within the range suggested for Greenland outlet glaciers (Morlighem et al., 2016; Choi et al., 2018, 2021). Note that the choice of the calving law is an important, yet poorly constrained, control on the grounding line dynamics and ice front behaviour (Schoof et al., 2017; Haseloff and Sergienko, 2018). In the absence of a universal calving law, a reasonable assumption has to be made, and we justify the choice of the von-Mises law through its relatively good performance in real-world applications (Choi et al., 2018).

Basal sliding is parameterized with a Budd type friction law (Budd et al., 1984) of the form

$$\boldsymbol{\tau}_b = -k^2 N \mathbf{v}_b, \quad (2)$$

where $\boldsymbol{\tau}_b$ is basal drag, k is a friction parameter, \mathbf{v}_b is the basal velocity, and N is the effective pressure. N is given as $N = \rho_i g H - \rho_w g \max(0, -z_B)$ where ρ_i and ρ_w are the density of ice and salt water, respectively, g is gravitational acceleration, H is ice thickness and z_B is bed elevation with respect to sea level. N is thus the difference between the ice overburden and water pressure assuming perfect connectivity between the subglacial water layer and the ocean. The effective pressure in the friction law induces an elevation-dependence of the basal resistance to flow. This dependence is motivated through the assumption that weak sediments are more likely to be present in low-lying areas. Implications for our results are discussed in sect. 4.2. We set k spatially uniform to isolate the topographic signal of retreat in our results and thus to reduce the number of degrees of freedom for the interpretation. We fix $k = 40$ (Pa yr m^{-1})^{1/2}, which is mid-range among values typically found in glaciological settings resembling ours (Bondzio et al., 2017; Haubner et al., 2018; Åkesson et al., 2018a).

Melting under floating ice is parameterized through prescribed melt rates that are invariant of depth. The reference forcing used for model spin-up is a subshelf melt rate of 30 m yr^{-1} and a frontal rate of undercutting of 200 m yr^{-1} , where the latter acts on the vertical ice front if the terminus is grounded. Both values are on the lower end of observed melt rates (Motyka et al., 2003; Enderlin and Howat, 2013; Xu et al., 2013), thus reflecting a configuration prior to recent warming when glaciers were more in balance with the ambient climate than today (King et al., 2020).

As part of the idealized setup, the surface mass balance (SMB) is fixed to zero, except in the uppermost 10 km of the domain (Fig. 1a), where we set an accumulation rate of 55 m yr^{-1} . This creates a realistic fixed upstream ice flux and is not meant

120 to represent local SMB found on real glaciers. Additionally, mass is added to the model domain by fixing an ice velocity of $v_x = 50 \text{ m yr}^{-1}$ at the upper domain boundary, which creates an influx as a function of the glacier thickness. These two approaches to adding mass represent a total accumulation (A) which can be expressed as

$$A = C + \int_0^{y_{max}} v_x(0, y) H(0, y) dy, \quad (3)$$

125 where $C = (10 \times 10^3)^2 \times 55 = 5.5 \times 10^9 \text{ m}^3 \text{ yr}^{-1}$ is a constant accumulation. Through the thickness-dependent influx represented by the second term on the right-hand side, we account for a reduction in accumulation for a shrinking glacier, thus parameterizing the SMB - altitude feedback (Harrison et al., 2001; Oerlemans and Nick, 2005).

We impose free-slip boundaries at the lateral margins of the domain (where $y = 0$ and $y = y_{max}$), meaning that no mass can leave the system laterally. In summary, the only mass source is at the upstream end of the domain, while the only mass wasting occurs where the glacier is in contact with the ocean (either through calving or melting).

130 2.2 Fjord Geometries

Our reference geometry is a fjord sloping linearly towards the ocean with a wide section in the upstream area from which ice is funneled towards a 5 km wide outlet channel with parallel walls (Fig. 1a). The fjord topography is given by $B(x, y) = B_x(x) + B_y(x, y)$ with

$$B_x(x) = B_0 + x \times \alpha + \Theta(x) \quad (4)$$

135 and

$$B_y(x, y) = \frac{d_f - B_x(x)}{1 + e^{(m_f(y - \frac{1}{2}L_y + w_f + \Omega(x) + \Theta(x)))}} + \frac{d_f - B_x(x)}{1 + e^{(-m_f(y - \frac{1}{2}L_y - w_f - \Omega(x) - \Theta(x)))}}, \quad (5)$$

where $\Omega(0 < x < x_U) = \frac{x - x_U}{F}^2$. The parameter values and descriptions are found in Table 1. This formulation is inspired by the MISMIP setup (Gudmundsson et al., 2012; Asay-Davis et al., 2016), but adapted to our purpose.

140 To insert basal or lateral perturbations in the outlet channel and thus alter the fjords' depth or width in specific areas, we modify the parameter $\Theta(x)$ in either eq. 4 or eq. 5 such that

$$\Theta(x_B < x < x_E) = -\sin\left(\frac{2\pi}{\Lambda}\left(x - \frac{\Lambda}{4} - x_C\right)\right)\frac{\Gamma}{2} + \frac{\Gamma}{2}. \quad (6)$$

Altering $\Theta(x)$ only in one of the terms on the right-hand side of eq. 5 allows to produce fjords with one-sided lateral perturbations, thus making them asymmetric. Combined, these equations reproduce the typical U-shape of fjords (Fig. 1b) and yield a setting that is representative of a wide range of outlet glaciers.

145 The metric used to quantitatively link fjord shape with glacier retreat is the wetted area S : the submerged cross-sectional area of the fjord (Fig. 1b), which can be calculated at every point along an outlet channel. This metric combines information




Figure 1. Schematic of the experimental setup. a) Sketch of the domain (not to scale) with annotated dimensions and mass balance processes (gains: thickness-dependent influx and surface accumulation; losses: melt at the ice-ocean interface and calving). Red box symbolizes how the fjord geometry is changed in different experiments to include geometric perturbations (their center being referred to as x_C). b) Cross-sections through the linear fjord (black line) and geometric perturbations. Upper panel: basal perturbations (green: depression, grey: bump); Lower panel: lateral perturbations (blue: embayment, yellow: bottleneck). The wetted area, i.e., the cross-sectional area of the fjord below sea level, is shaded in blue for each geometry.

about the width and depth of a fjord and is thus a comprehensible parameter for comparing basal and lateral perturbations. Furthermore, it is straightforward to calculate its first derivative, dS , which yields information on the along-fjord change in width or depth.

150 Besides our reference setup, we test 20 fjord geometries (Table 2), each of which contains either a small, medium or large geometric perturbation. The magnitude of the perturbation is defined by how much the width or depth of the fjord deviates from the reference fjord. Our 'core experiment', which the results will focus on, comprises 12 fjords, each of which features one of the four perturbation types (depression, bump, bottleneck, embayment) of one of the three magnitude classes (small, medium, large). The depressions and embayments in each magnitude class increase the wetted area S at the center of the perturbation
 155 x_C by the same amount, while the bottlenecks and bumps in each magnitude class reduce S at x_C by the same amount. The along-flow horizontal extent of all perturbations in the core experiment is 20 km (Fig. 2).

In the eight simulations outside our core experiments, we test *asymmetric* and *longer* perturbations to verify if the results from our core experiment can be transferred to a wider range of settings. We test two asymmetric embayments, which have the

Table 1. Parameters for generating fjord geometries (in parentheses for *longer* geometric perturbations).

parameter	value	unit	description
B		m	bed elevation
B_0	-450	m	bed elevation at $x=0$
α	-0.002		slope of bed in x-direction
d_f	2000	m	depth of fjord relative to upland areas on the sides
L_y	10	km	width of domain in y-direction
L_x	85	km	length of domain in x-direction
w_f	2.5	km	half-width of fjord
m_f	$\frac{1}{300}$		factor for steepness of fjord walls
x_U	30	km	extent in x-direction of wide upstream area
F	300		factor for smooth transition between wide upstream area and parallel fjord
x_B	45 (35)	km	x-coordinate of upstream end of perturbation
x_E	65 (65)	km	x-coordinate of downstream end of perturbation
x_C	55 (50)	km	x-coordinate of center of perturbation
Λ	20 (30)	km	horizontal extent of perturbation in x-direction
Γ	<i>variable</i>	m	deviation of fjord half-width or depth relative to parallel fjord at x_C

same S at x_C as the small and medium embayments and depressions, as well as two asymmetric bottlenecks which have the
160 same S at x_C as the small and large bottlenecks and bumps. The longer perturbations have an along-flow horizontal extent of
30 km. We test one longer perturbation per perturbation type with S at x_C corresponding to the medium magnitude class.

2.3 Reference Glacier

All experiments start from an artificial reference glacier, which is produced by relaxing a rectangular block of ice in the
reference fjord. The spin-up is run until the relative ice volume ($(dV/yr)/V \ll 0.05\%$) and GL position are stable. The
165 length of the spun-up reference glacier is 82 km, its GL is at $x = 73$ km and the velocity at the GL along the central flow line
of the glacier $v_{GL} = 2.5$ km yr⁻¹. At steady-state, the total mass gain is ~ 6.1 km³ yr⁻¹ (≈ 5.6 Gt yr⁻¹), which is balanced
by mass losses through melting at the ice - ocean interface (~ 0.9 km³ yr⁻¹) and calving (~ 5.2 km³ yr⁻¹). In a sensitivity
experiment with doubled ocean melt rates (400 m yr⁻¹ undercutting and 60 m yr⁻¹ subshelf melting), mass loss through ocean
melt increases to ~ 2.1 km³ yr⁻¹, while calving reduces to ~ 4 km³ yr⁻¹. The GL remains largely stable, indicating that the
170 reference glacier is not very sensitive to ocean forcing due to compensating effects in the mass wasting processes.

The setup represents a medium sized fjord-glacier system, which has similar dimensions and dynamics as, for example, the
present-day Alison glacier in NW Greenland, where the fjord width is about 5 km, water depth is around 500 m and observed
ice discharge has increased from ~ 4 to ~ 8 Gt yr⁻¹ in the past 20 yrs (Mouginot et al., 2019). It is furthermore broadly

Table 2. Suite of experiments with name (extensions *_lon* and *_asy* refer to *longer* and *asymmetric* geometries), type of geometric perturbation, perturbation magnitude, the deviation of fjord width (two times Γ for symmetric lateral perturbations, one time Γ for asymmetric ones) or depth (one time Γ for basal perturbations) at the center of the perturbation relative to the linear reference fjord, S at x_C (i.e., the wetted area at the center of the perturbation) and forcings required to induce complete retreat through the entire geometric perturbation (/ if no complete retreat could be enforced).

Experiment	Perturbation type	Perturbation magnitude	Fjord width/depth deviation [m]	S at x_C [km ²]	Forcing for complete retreat (undercutting/subshelf melt rate) [m yr ⁻¹]
Ref	-	-	0	2.1	800/120
ByH450	embayment	small	900	2.6	1200/180
ByH900	embayment	medium	1800	3.1	1200/180
ByH1350	embayment	large	2700	3.6	1200/180
ByH-450	bottleneck	small	-900	1.6	800/120
ByH-675	bottleneck	medium	-1350	1.3	/
ByH-900	bottleneck	large	-1800	1.1	/
BuH-120	depression	small	-120	2.6	1200/180
BuH-240	depression	medium	-240	3.1	1000/150
BuH-360	depression	large	-360	3.6	800/120
BuH120	bump	small	120	1.6	/
BuH180	bump	medium	180	1.3	/
BuH240	bump	large	240	1.1	/
ByH900_lon	embayment	medium	1800	2.8	1200/180
ByH-675_lon	bottleneck	medium	-1350	1.5	1200/180
BuH-240_lon	depression	medium	-240	2.8	1200/180
BuH180_lon	bump	medium	180	1.5	/
ByH900_asy	embayment	small	900	2.6	1200/180
ByH1800_asy	embayment	medium	1800	3.1	1200/180
ByH-900_asy	bottleneck	small	-900	1.6	1200/180
ByH-1800_asy	bottleneck	large	-1800	1.1	/




Figure 2. Along-fjord profiles of the wetted area S (a: lateral; b: basal perturbations) and its derivative dS (c: lateral; d: basal) for fjords featuring different geometric perturbations of different magnitude classes. Note that the profiles of embayments and depressions, and likewise, bottlenecks and bumps, of the same magnitude class are largely congruent, thus allowing a straightforward comparison between basal and lateral perturbations.

representative of outlet glaciers from the Fennoscandian ice sheet during the last glacial, such as the Hardangerfjorden glacier
175 (Mangerud et al., 2013; Åkesson et al., 2020).

2.4 Retreat Experiments in variable fjords

We slightly modify the reference glacier to match the new fjord geometry when introducing geometric perturbations. For
embayments, we extrapolate the glacier surface laterally to fill the newly introduced lateral cavities. For depressions, we fill
the new basal cavity with ice, but keep the glacier surface the same. For bumps or bottlenecks, we remove ice while keeping
180 the glacier surface unaltered. Subsequently, we relax the glacier in each geometry for 50 yrs, resulting in an ice volume change
(dV/yr)/ $V < 0.5\%$ at the end of relaxation for every setup tested and a stable GL.

After relaxation, we increase the ocean forcing to trigger a retreat. We aim to force the GL to retreat through the entire
geometric perturbations. The ocean melt rates required to induce such a retreat depend on the fjord geometry, which we
elaborate on in the results section. To determine what melt rates are needed to force this complete retreat in a particular fjord,
185 we strengthen the ocean forcing using multiples of the reference forcing (200 m yr^{-1} frontal rate of undercutting, 30 m yr^{-1}
subshelf melt) until complete retreat takes place. In some cases (c.f. sect. 3.2), even unrealistically high values for the ocean
forcing (e.g. 20 times the reference forcing) did not trigger complete retreat, suggesting that glaciers in these geometries are
not sensitive to ocean melt.

Since we want to explore the response of outlet glaciers to melting at the ice-ocean interface, we keep the SMB constant
190 with time, and let the upstream ice flux vary only through our parameterized SMB - altitude feedback (Eq. 3).

We assess 16 glacier metrics during the retreat, which we expect to show a response to local topography (Table 3). All of
these can be observed in-situ or via remote sensing techniques (e.g. Mouginot et al., 2019; King et al., 2020), which means that
our results are readily transferable to real-world settings. The GL position (x_{GL}) and its derivative, the GL retreat rate (dGL),
the front position (x_{Fr}) and its derivative, the frontal retreat rate (dFr), but also the velocity at the GL (v_{GL}) and the shelf
195 length (L_S) are measured along the central flow-line of the glacier.

2.5 A real-world case study: Jakobshavn Isbræ

We want to verify to what degree the dynamics seen in our experiments are also prevalent in real-world settings. This is
challenging since we investigate decadal to centennial time scales. Specifically, we would need observations with high temporal
resolution on glacier metrics (Table 3) for a glacier that has retreated over tens of kilometers, through a fjord with variable and
200 known topography. There are perhaps only a handful of glaciers worldwide that may fulfil these requirements, and even so,
acquiring the necessary data is difficult and outside the scope of the present study.

To test our idealized results in a real-world setting, we instead turn to model simulations from Jakobshavn Isbræ's (JI)
evolution in the Holocene (Kajanto et al., 2020). We focus on a simulation of the retreat of JI from a sill at the fjord mouth
of Jakobshavn Isfjord, to a point inland of today's GL position. This model retreat is forced using a step reduction in the
205 equilibrium line altitude early in the Holocene (experiment SE_CM in Kajanto et al. (2020)). While this is a sensitivity exper-

Table 3. Glacier characteristics assessed during retreat for later correlation with fjord geometry. Parameters marked with * are assessed along the central flow-line of the glacier.

Glacier metric	Variable	Unit
Grounding line position*	x_{GL}	km
Grounding line retreat rate*	dGL	m yr^{-1}
Front position*	x_{Fr}	km
Front retreat rate*	dFr	m yr^{-1}
Grounding line mass flux	Q_{GL}	$\text{km}^3 \text{ yr}^{-1}$
Ice front mass flux	Q_{Fr}	$\text{km}^3 \text{ yr}^{-1}$
Flux through an upstream gate	Q_U	$\text{km}^3 \text{ yr}^{-1}$
Calving Flux	Q_C	$\text{km}^3 \text{ yr}^{-1}$
Velocity at the grounding line*	v_{GL}	m yr^{-1}
Maximum velocity	v_{max}	m yr^{-1}
Shelf length*	L_S	m
Floating Area	A_F	m^2
Grounded Area	A_G	m^2
Ice Volume	V	km^3
Ice Volume Above Flotation	V_{AF}	km^3
Maximum Ice Thickness	H_{max}	m

iment not meant to reflect the actual evolution of JI (Kajanto et al., 2020), it is convenient for our purposes since it produces a long-lasting, dynamic retreat.

Just like in our idealized experiments, we calculate S and dS for Jakobshavn Isfjord. We then assess whether the relationships found in our idealized settings are also prevalent in JI’s dynamics (sect. 3.5).

210 3 Results

3.1 Stagnant and ephemeral grounding line positions

We identify positions of GL stagnation (‘stagnant’ GL positions), i.e., where the GL rests for a sustained time (typically 50 to 200 yrs), or retreats slowly ($dGL > -100 \text{ m yr}^{-1}$), and areas where the GL retreats quickly (‘ephemeral’ GL positions; $dGL < -500 \text{ m yr}^{-1}$). Fig. 3 shows both stagnant and ephemeral positions for one representative run per perturbation type. For
215 a comparison of the GL retreat dynamics of all simulations within our core experiment, the reader is referred to Fig. SA1. Note that in the following, all terminology related to along-fjord changes in width or depth of the fjord (e.g. narrowing, deepening) will refer to the direction of glacier retreat.

Stagnant positions exist at the downstream end of embayments and depressions where the fjord becomes wider/deeper ($x \approx 62$ to 65 km, Fig. 3a,c). Ephemeral positions, associated with rapid retreat, are found in the remainder of both perturbations (220 $x \approx 45$ to 62 km). Retreat from the stagnant position at the downstream end of the embayment occurs gradually ($dGL \approx -22$ m yr⁻¹ while $x_{GL} \approx 62$ to 64 km), but is rapidly accelerating as the GL retreats further into the perturbation, accompanied by some lateral ungrounding. Retreat from the stagnant position in the depression occurs suddenly after a phase of near-stability ($dGL \approx -6$ m yr⁻¹, while $x_{GL} \approx 64.5$ to 65.5 km) as the glacier ungrounds where the fjord is deepest in the center of the perturbation ($x \approx 55$ km, Fig. 3c inset plot). The cavity formed under the glacier rapidly grows in size and expands (225 downstream until it eventually detaches the glacier from the bed also at the downstream end of the depression ($x \approx 65$ km). In bottlenecks and on bumps, stagnant positions are found where the fjord is narrow/shallow ($x \approx 55$ to 58 km, Fig. 3b,d). The stabilizing effect of bumps is, in fact, so large that no glacier could be forced to retreat over them within reasonable limits for the ocean forcing. However, we observe that retreat onto bumps occurs fast ($dGL \approx -500$ m yr⁻¹ for $x_{GL} \approx 58$ to 65 km). For bottlenecks, only the glacier situated in the fjord with a 'small' bottleneck (i.e., the bottleneck with the largest S among the ones (230 tested) could be forced to retreat completely. Noticeably, retreat at the downstream end of the bottleneck ($x \approx 56$ to 65 km), where the fjord narrows in, is fast (ephemeral) with $dGL \approx -900$ m yr⁻¹, whereas it is very slow (relatively stagnant) with $dGL \approx -25$ m yr⁻¹ upstream of the narrowest point, where the fjord is widening ($x \approx 45$ to 55 km; Fig. 3d).

In summary, stagnant GL positions are found where the fjord widens/deepens in the direction of glacier retreat (positive dS , Fig. 2) and rapid retreat occurs through areas where the fjord becomes narrower/shallower (negative dS). Thus, the along-fjord change in width or depth (dS) is a key control on GL retreat. However, glaciers in narrower or shallower fjords than (235 the reference fjord (bottlenecks and bumps) can also temporarily stagnate where S is small. This shows that the wetted area constitutes an additional important control on GL retreat. The experiments with asymmetric and longer perturbations confirm these findings (see Fig. SA2).

3.2 Forcings and timings of retreat

Now, we investigate how retreat from stagnant and ephemeral GL positions is correlated with fjord topography (i.e., S and dS). Two parameters are important in this context (Fig. 4): First, the amplitude of the forcings needed to induce complete retreat through the different geometric perturbations. As mentioned previously, distinctions exist both between the different perturbation types (bumps, depressions, bottlenecks, embayments) as well as the different magnitude classes (small, medium, large) for a given geometry type. Second, the approximate residence time of the GL in a stagnant position. The stronger the (245 GL is stabilized by a particular geometric perturbation, the longer it will be stagnating.

All glaciers in embayments require the same increase in forcing to retreat completely (6 times the spin-up forcing). This increase is larger than what is needed to induce retreat through the linear reference fjord (4 times the spin-up forcing). The residence time of the GL in the stagnant positions at the downstream end of the embayments are such that the glacier in the smallest embayment is the earliest to retreat (after 61 yrs of GL stagnation), and the one in the largest the latest (after 173 yrs) (250 Fig. 4). This implies that the larger the embayment, the more stability it provides to the glacier at its downstream end, before retreat through the entire perturbation is possible. A larger embayment means a locally larger along-flow change in wetted area




Figure 3. Annual glacier evolution (a, b: top down view on domain showing yearly grounding lines; c, d: yearly glacier profiles along central flow line) in fjords featuring different geometric perturbations (location of perturbations marked in grey) with the spacing between the lines indicating retreat velocity: a) medium embayment; b) small bottleneck; c) medium depression; d) small bump. Inset plot in c) shows profile (blue) in year 217 when glacier ungrounds in the central part of the depression which triggers further retreat. Red line is the level to which glacier needs to thin to reach flotation.




Figure 4. Forcing required to induce complete retreat in multiples of the reference forcing (200 m yr^{-1} undercutting rate, 30 m yr^{-1} subshelf melt) and approximate residence time of the GL in a stagnant position for different fjord geometries. A longer residence time and a larger forcing required indicate that fjord geometry provides larger stability. More stability is correlated with decreasing fjord depth for depressions (shades of green), and with increasing along-fjord change in wetted area dS for embayments (shades of blue). The simulations for which no retreat through the perturbation was observed have been omitted from the figure.

dS at its downstream end (Fig. 2). Thus, there is a positive correlation between GL stability and dS . This indicates that dS not only determines the location of stagnant GL positions in embayments, as shown before, but that it also quantitatively impacts how stagnant the GL is.

255 The glaciers in fjords with depressions require different forcings to retreat completely (small: $6\times$ the reference forcing, medium: $5\times$, large: $4\times$). The residence time also varies; retreat over small depressions occurs ~ 65 yrs later than over medium and large depressions, which retreat after about the same time (after 169 and 170 yrs, respectively). These findings indicate that the stabilizing effect of a depression declines the deeper it is (Fig. 4). Thus, there is a negative correlation between GL stagnation and S . In sect. 3.1, ungrounding in the central part of depressions was identified as the trigger of rapid retreat from

260 the temporary stillstands. Such GL retreat occurs more easily in a deeper fjord (larger S). Therefore, it is consequential that a deeper depression is less stabilizing. Note, however, that the fjord depth several kilometers upstream of the GL determines for how long the GL stagnates. There is no direct correlation between S or dS at the GL and the stability provided to the glacier by the fjord in our settings with depressions.

The glacier in a fjord with a 'small' bottleneck required a four-fold increase in oceanic melt rates, and retreated from its stagnant position after 126 yrs of stagnation. This is a weaker forcing than for the glaciers in the embayments, as well as for the medium and small depressions, and thus suggests that this bottleneck provides less stability than these geometries. This contrasts with the common pattern, where a small S and a positive dS should stabilize the glacier strongly. It is unclear why this is the case here. We hypothesize that it might be related to a combination of high driving stresses due to a steepened surface inside the bottleneck in conjunction with high modeled calving rates (not shown). The two experiments with glaciers in geometries with narrower bottlenecks ('medium' and 'large' bottleneck) did not retreat through the entire perturbation. This, in turn, aligns well with the general notion of a confined (low S) and downstream narrowing (positive dS) fjord yielding strong stability to the glacier. Likewise, none of the glaciers in fjords with bumps retreated completely, which follows the same concept. However, the strong stability that bumps provide to the glacier may also be related to the choice of model parameters (sect. 4.2).

3.3 Stress balance response to fjord geometry

To assess the underlying mechanisms behind the geometric controls described before, we now analyze the stress regimes across the studied geometries. We focus on lateral shear stress gradients for lateral perturbations, and longitudinal stress gradients for basal perturbations as given by the SSA in x-direction by

$$\rho g H \alpha_x = \tau_{bx} + \frac{\partial}{\partial x} (2H \sigma'_{xx} + H \sigma'_{yy}) + \frac{\partial}{\partial y} (H \sigma'_{xy}) \quad (7)$$

where σ' is the deviatoric stress and τ_b is basal drag. We interpret the second and third term on the right-hand side as longitudinal stress gradient and lateral shear stress gradient, respectively. With the imposed spatially uniform friction coefficient, variations in the investigated stress fields are largely caused by variable fjord topography, and are hence convenient to investigate for our purpose.

For embayments and bottlenecks, variations in lateral shear stress gradients can be seen along the fjord walls (Fig. 5a,b). Specifically, strongly negative shear stresses are found where ice is funneled in a downstream narrowing fjord. This occurs, for example, where $55 \text{ km} < x < 65 \text{ km}$ in embayments (Fig. 5a) and at $45 \text{ km} < x < 55 \text{ km}$ in bottlenecks (Fig. 5b). This indicates enhanced resistance to flow for the glacier originating from the fjord walls, which provides stability to the glacier. Where ice flow diverges in a widening section of the fjord (in the upstream half of embayments and the downstream half of bottlenecks, Fig 5a,b), lateral shear stress gradients are comparatively weak. This indicates that the glacier - fjord wall contact is reduced here, and that the fjord walls provide little support to the glacier in these areas.

For depressions and bumps, we see clear variations in longitudinal stress gradient along the glacier bed (Fig. 5c,d). In depressions, a band of negative values stretches across the full width of the outlet channel where the bed turns from being prograde to retrograde (at $x = 55 \text{ km}$, Fig. 5c), indicating that ice flow is being blocked here. Likewise, a marked reduction in positive longitudinal stresses is seen at the onset of the bump where the bed slope switches sign (at $x = 45 \text{ km}$, Fig. 5d). Together, the stress regimes in basal perturbations demonstrate that a retrograde glacier bed, tilted against the direction of flow,




Figure 5. Stress states in perturbations. lateral shear stress gradients for a, b with flow lines in grey, and longitudinal stress gradients for basal perturbations. Green line is grounding line, black line is glacier outline.

reduces longitudinal stress gradients considerably as it increases the basal resistance to flow, which ultimately stabilizes the glacier.

In summary, the stress analysis above suggests that increased lateral shear stress gradients or negative longitudinal stress gradients are found wherever ice flow is forced to converge, either horizontally or vertically, towards a narrowing or shoaling area downstream. Simulations using asymmetric as well as longer perturbations confirm that these findings are robust (see Fig. SA3). Through the convergent flow, the contact between the glacier and the fjord is enhanced, leading to increased resistance to flow. Overall, along-flow change in fjord width or depth (i.e., dS) is found to define areas of increased lateral shear gradients or negative longitudinal stress gradients, and thus GL stagnation.

3.4 A universal quantitative relationship for ice-topography interaction

We hypothesize that there is a quantitative relationship between fjord geometry and glacier retreat, valid across a range of different geometries. To test this, we correlate a variety of metrics indicative of glacier retreat (Table 3) against relevant metrics of fjord geometry, that is, the submerged cross-sectional area (S) and its derivative (dS). We restrict the data to those instances when the GL is located within a geometric perturbation (gray-shaded areas in Fig. 3). Among all combinations of

retreat and geometry metrics tested, including those of asymmetric and 'longer' perturbations (Table 2), the clearest and most
310 universal relationship found is a negative, close to linear correlation between the ratio of the GL flux and the submerged cross-
sectional area Q_{GL}/S over the change in submerged cross-sectional area dS (Fig. 6). This relation expresses that a widening
or deepening fjord in the downstream direction (negative dS) promotes a high GL flux per wetted area (Q_{GL}/S). Conversely,
a glacier retreating in a fjord that becomes narrower or shallower downstream (positive dS), will have a reduced Q_{GL}/S . Note
that the ratio Q_{GL}/S for basal geometry perturbations (grey to black and green colors in Fig. 6) is on average lower than for
315 lateral geometry perturbations. This means that basal perturbations generally inhibit ice flux across the GL more efficiently
than lateral perturbations (note that this may be influenced by our modelling choices (sect. 4.2)). Also, note that the GL flux
is the product of the velocity v_{GL} and the flux gate area at the GL A_{GL} , that is $Q_{GL} = v_{GL} \times A_{GL}$. The ratio Q_{GL}/S is thus
proportional to v_{GL} when there is hydrostatic equilibrium at the GL (because in that case, $S = 0.9 \times A_{GL}$), and so we find a
comparable, negative linear relationship between v_{GL} and dS (Fig. SA4).

320 We find an additional, yet less distinct, negative relationship between the wetted area S and the GL retreat rate dGL (Fig. 7a).
This shows that a wider or deeper fjord promotes faster GL retreat, while a narrower or shallower geometry stabilizes the
glacier. This relation is not as universal as the previous one since one value for dGL is not uniquely linked to one value for
 S across different geometries. Furthermore, it is not linear, but rather such that for a range of low S values, dGL does not
vary noticeably. Only above a certain threshold in S , the GL retreats markedly faster (Fig. 7b). This threshold varies between
325 different fjord geometries. However, we find that it is always associated with the location of GL stagnation (Fig 3.1). This
means that a relationship between GL retreat rate dGL and S only unfolds if a local stability position is passed. These stagnant
positions can be either where S is low, or where dS is high, as shown previously. For instance, dGL does not increase as the
GL retreats very slowly at the stagnant position in the downstream half of embayments. Only once it has retreated passed this
point of GL stagnation, a correlation between dGL and S can be seen.

330 For depressions we do not see a distinct relationship between dGL and S (Fig. SA5). This is because we measure the GL
position x_{GL} and therefore also dGL as the furthest downstream grounded point along the central flow-line of the glacier.
When the glacier ungrounds in the center of a depression, where the fjord is deepest, the dynamics of retreat are triggered
several kilometers upstream of the GL, as mentioned in sect. 3.1. Therefore, there is a correlation between fjord depth and
GL retreat in depressions. However, it is not reflected when only considering processes at the GL. Not finding a dGL over S
335 relation for depressions is hence expected by construction of our methodology, and not an actual feature.

3.5 Jakobshavn Isbræ

Given our previous results, we now aim to assess whether our principal geometric relationship Q_{GL}/S over dS can be found
for Jakobshavn Isbræ. To this end, we calculate the wetted area S along the topography of Jakobshavn Isfjord as used in Kajanto
et al. (2020) which depicts overall higher values and larger along-fjord changes in dS than our idealized settings (Fig. 8b).

340 Plotting all available data points for Q_{GL}/S over dS at Jakobshavn, we do not find the aforementioned geometric relation-
ship. This may have many reasons related to the complex dynamics of Jakobshavn Isbræ (Bondzio et al., 2017), but most
critically, there is lateral inflow of ice to the main channel from the surrounding ice sheet and tributaries (compare with Steiger




Figure 6. Relationship between grounding line discharge per wetted area Q_{GL}/S and along-fjord change in wetted area dS for all tested geometries and all instances when the GL is within a geometric perturbation (grey area in Fig. 3).




Figure 7. Relationship between grounding line retreat rate dGL and wetted area S . a): All instances when the grounding line is within a geometric perturbation for all tested geometries, except for depressions since retreat in these perturbations is governed by different dynamics than in the ones shown. b): Schematic of a typical relationship between dGL and S where dGL is constant for low S , while high S induces faster retreat. The transition between these two states occurs if the GL retreats past a point of GL stagnation. The location of these points may be controlled by either low S or high dS .

et al., 2018). This alters the stress balance at the GL compared to our experiments where the glacier is always closely confined between fjord walls. Specifically, it means that the rigid glacier-wall interface in our experiments is replaced by a changing
 345 ice-ice contact. This has implications for the lateral friction that the fast-flowing ice in the main channel experiences and for the processes transferring stabilizing back stress from the sides to the center of flow.

We thus expect our findings to be more easily transferable to settings where Jakobshavn Isbræ is enclosed by fjord walls. This is only the case in one part of the outlet channel, upstream of the present-day front (Fig. 8c). Indeed, values for Q_{GL}/S are inversely related to dS in a qualitative way here, such that an increase in dS is generally associated with a decrease in Q_{GL}/S
 350 and vice versa (Fig. 8d,e), consistent with our findings for synthetic geometries above (Fig. 6). Even though this relationship is only qualitative, meaning that one value for dS is not uniquely associated with one value for Q_{GL}/S , we find these results encouraging given the complexity of Jakobshavn Isbræ's dynamics. For settings resembling our setup more closely, such as medium-sized outlet glaciers found in, e.g. Greenland (Carr et al., 2014; Bunce et al., 2018; Catania et al., 2018), Svalbard (Schuler et al., 2020) and Novaya Zemlya (Hill et al., 2017), we expect an even stronger imprint of topography on retreat
 355 dynamics.




Figure 8. The real-world example Jakobshavn Isbræ. a) Topography of Jakobshavn Isfjord (Morlighem et al., 2017) with approximate present-day glacier front, and centerline along which profiles shown in b) of the wetted area S and its along-fjord change dS are calculated; c) zoom to area where JI is enclosed between fjord walls, with yellow circles showing all modeled grounding line positions in this section (Kajanto et al., 2020); d) dS profile in the same section of the fjord with grounding line positions indicated; e) Q_{GL}/S in this section with grounding line positions and a polynomial fit (blue dotted line). The opposing trends in dS (d) and Q_{GL}/S (e) as indicated by the arrows demonstrate qualitatively that the negative relationship Q_{GL}/S over dS can be found in this complex setting.

4 Discussion

4.1 Mechanisms behind geometric controls of glacier dynamics

The current study offers new quantitative insights into how topography influences the evolution of marine outlet glaciers, and their response to ocean warming. We demonstrate that two topographic metrics, the wetted area S and its derivative dS , jointly control the dynamics and retreat of glaciers constrained by fjord walls. Together, these metrics largely explain variations in grounding line mass flux Q_{GL} , which is important in the context of sea-level rise, and the grounding line retreat rate dGL .

Based on our stress analysis and physical principles, we propose the following physical interpretation for these results: First, a downstream narrowing or shoaling fjord (positive dS) stabilizes the glacier, as ice flow is funneled through the constriction

enhancing the glacier-fjord contact (Fig. 5a,c). This increases the basal or lateral resistance to flow, which stabilizes the glacier.
365 Conversely, a downstream widening/deepening fjord (negative dS) provides little support to the glacier as glacier-fjord contact
is reduced (Fig. 5b,d). Second, a narrow fjord (low S) stabilizes the glacier, because the distance between the lateral ice margins,
where friction with the fjord walls is high, and the center of flow is small. This means that the part of the glacier where ice flow
is largely undisturbed is reduced (Raymond, 1996; van der Veen, 2013). Third, a shallow fjord (low S) stabilizes the glacier
because the glacier is further away from flotation, and thus grounding line retreat is less likely to occur with a given amount of
370 thinning (Pfeffer, 2007; Enderlin et al., 2013). In our experiments, the area exposed to ocean melt does not have a large effect
on retreat dynamics. Even high oceanic melt rates, which could compensate for a small ice-ocean interface, do not trigger
retreat through geometric perturbations where S is low.

For a particular fjord geometry, the relative importance of S or dS in providing stability to the glacier may vary. This will
be discussed with two examples from our results: In embayments, S is larger than the reference fjord. Therefore, if S was the
375 dominant control for glacier dynamics here, retreat through embayments should occur more easily than through the reference
fjord. However, we find that the opposite is true; the grounding line stagnates at the downstream end of embayments (Fig. 3a),
while it retreats steadily through the reference fjord (Fig. SA1). This indeed confirms that S alone does not explain glacier
retreat. Rather, dS controls glacier dynamics, because the point of grounding line stagnation is where the fjord changes from
wide to narrow in the direction of ice flow. For bed bumps in our experiments, the picture is different. Our model glaciers
380 stagnate on or near the crest of the bumps, where dS is close to 0 or negative (Fig. 3d). This should not be an obstacle for retreat
if dS was the dominant control on glacier dynamics. Therefore, it must be the shallowness of the fjord at this point (indicated
by low S) which governs the dynamics here.

Given these disparities between different settings, it is all the more compelling that we find the geometric relationship
 Q_{GL}/S over dS universal to all our tested fjords. It implies that given the current grounding line mass flux Q_{GL} and the
385 upstream subglacial topography of a particular glacier, a well-founded estimate of the topographically induced future contribu-
tion to sea-level rise can be made. To the authors knowledge, this type of universal quantitative link between fjord topography
and glacier response has not been established before, going beyond the qualitative descriptions of ice-topography interaction
offered in previous studies (Enderlin et al., 2013; Carr et al., 2014; Bunce et al., 2018; Catania et al., 2018; Åkesson et al.,
2018b). For projections of future sea-level rise, this direct coupling between topography and ice discharge is highly relevant,
390 as it enables an ad-hoc assessment of the expected future sea-level contribution of a glacier on decadal to centennial time
scales. Q_{GL} is readily available for glaciers where the velocity and bathymetry is well-known. Moreover, where bathymetry is
uncertain, or for less well-studied glaciers, we demonstrate that the velocity evolution over time is also a good proxy for the
dynamic reaction of a glacier to fjord topography (Fig. SA4).

Our second quantitative relationship between dGL and S confirms the widely accepted concept that a wide or deep fjord
395 promotes fast grounding line retreat (e.g. Warren and Glasser, 1992; Enderlin et al., 2013; Carr et al., 2013; Bunce et al., 2018;
Catania et al., 2018; Åkesson et al., 2018b). However, we highlight that this relationship may not hold if the fjord is narrowing
or shoaling downstream (positive dS). In practical terms, this means that retreat from a uniformly wide and deep channel into
an upstream widening or deepening section does not automatically imply that retreat has to accelerate. Rather, we suggest that

a glacier will sit at the downstream end of the section, where S is increasing upstream, for a considerable time, or may not
400 even retreat further, because it is particularly stagnant here. Only if this position is abandoned, fast retreat will occur (Fig. 5a).
This fast retreat is in fact facilitated by the long residence time of the grounding line, since concurrent upstream thinning
preconditions the glacier for fast retreat.

In contrast to most previous studies, we emphasize the role of along-fjord change in fjord topography (dS) to explain geo-
metric controls of outlet glaciers. Along-fjord change in fjord depth is also key in the context of the marine ice sheet instability
405 (MISI) theory, according to which retrograde beds promote retreat (Schoof, 2007; Gudmundsson et al., 2012; Gudmundsson,
2013). However, even though we do have retrograde beds in our fjords with basal perturbations, we do not see any influence
of the MISI on glacier dynamics. This is simply because our tested glaciers never retreat into an area where the bed slope
is strongly negative, and where the MISI effect would be expected to occur. On bumps, the glaciers stop to retreat on the
downstream side where the bed is prograde (Fig. 3d). In depressions, the grounding line stagnates where the bed slope is
410 only slightly negative (Fig. 3c), which is not enough to trigger a MISI feedback loop. Retreat off this stagnant position occurs
through ungrounding several kilometers upstream of the grounding line. This process is not related to typical MISI dynamics.
Besides that, our quantitative relation Q_{GL}/S over dS may seem contradictory to widely accepted concepts of glacier dy-
namics, because we project high Q_{GL}/S for prograde beds (i.e., negative dS in our study). This may give the impression that
prograde beds should lead to accelerating ice discharge. However, we emphasize that we assess ice discharge *per area*, not
415 absolute values of ice discharge. A glacier retreating on a prograde bed will experience a reducing wetted area as it recedes,
and thus the ratio Q_{GL}/S may increase, but not the absolute grounding line flux. This is exemplified by our experiments with
bumps, where the glacier stagnates on a prograde bed even though Q_{GL}/S is relatively high (Fig. 6). Essentially, we thus
describe a mass conservation mechanism where a smaller flux gate requires a smaller ice flux to maintain the same grounding
line position. This is well known in the context of ice-topography interaction (Jamieson et al., 2012; Åkesson et al., 2018b).

420 Only a few studies have considered the influence of along-fjord changes, rather than absolute values, in fjord width on
glacier dynamics, and available observations are limited in time (Carr et al., 2014; Bunce et al., 2018). The main consensus is
that a fjord widening in the direction of glacier retreat promotes fast grounding line recession, while a narrowing fjord reduces
retreat rates. Furthermore, retreat onto a pinning point can stabilize the grounding line. This is related to our findings in that
we also see accelerating retreat the further the grounding line moves into a wider fjord upstream (c.f. grounding line positions
425 in the downstream half of the embayment ($55 \text{ km} < x < 65 \text{ km}$) in Fig. 3a). However, in our results, this only occurs after a
phase of grounding line stagnation at the downstream end of such fjord sections. We do not find conclusive evidence in the
observational record whether these points of grounding line stagnation are a relevant phenomenon in real-world settings or not
(Carr et al., 2014; Bunce et al., 2018; Catania et al., 2018). Further research analysing a range of fjord geometries and glacier
retreat histories is required to test this result. We do see some signs in our experiments that retreat slows down the further the
430 grounding line recedes into a narrower fjord upstream. Overall though, retreat in upstream narrowing fjords is markedly faster
than if the fjord is upstream widening (compare grounding line positions in the downstream half of bottlenecks ($55 \text{ km} < x < 65$
 km) with the ones in the upstream half ($45 \text{ km} < x < 55 \text{ km}$) in Fig. 3b). This we explain with the aforementioned enhancement
(reduction) in fjord-glacier contact for an upstream widening (upstream narrowing) fjord (Fig. 5a,b). Thus, we confirm that

retreat slows down in an upstream narrowing fjord, but in the context of a retreat cycle through both upstream widening and
435 narrowing fjord sections, overall faster retreat occurs through upstream narrowing fjords. Therefore, the observational records
of glacier retreat in Greenland may be too short and the fjord-width variations too small to testament similar dynamics as we
observe in the model (Carr et al., 2014; Bunce et al., 2018; Catania et al., 2018). However, in line with the observational record,
we can reproduce the stability that lateral pinning points offer. This is clearly demonstrated by the strong stability that narrow
bottlenecks provide in our experiments.

440 Our experiments are set-up to simulate grounding line retreat, and hence we do not offer any insights on ice-topography
interactions for advancing glaciers. Previous studies, however, suggest that fjord geometry induces hysteresis in the retreat-
advance cycle of a glacier, meaning that a reversal to colder conditions after a phase of climate warming does not allow the
grounding line to advance to the same position it occupied initially if the fjord is widening or deepening in front of the glacier
(Brinkerhoff et al., 2017; Åkesson et al., 2018b). We expect this also to hold for our experiments if we had simulated ocean
445 cooling following the warming scenarios tested.

4.2 Study limitations

As in all numerical studies, our results have limitations related to the choice of model parameters. In particular, there are three
aspects that warrant further discussion. First, the SSA used here is not a full representation of the stress regime in a glacier
which especially bears relevance near the grounding line. For weak beds and fast-flowing outlet glaciers, as we aim to mimic in
450 our synthetic setup, the SSA is a reasonable approximation and widely used in the glaciological literature. It may fail, however,
on steep bed slopes, such as the ones in our simulations of basal perturbations. In particular, the effect of basal high-friction
points may be overestimated which might explain why basal perturbations generally yield lower Q_{GL}/S values than lateral
perturbations in our experiments (Fig. 6), and why bumps were found to be excessively stabilizing. However, we do not expect
this effect to compromise our conclusions in general. For instance, Favier et al. (2012), using a setup similar to ours to simulate
455 the effect of a basal pinning point on ice dynamics with a Full-Stokes model, present stress distribution patterns that agree
favorably with the ones shown here. The influence of using the SSA as compared to a Full-Stokes model on our results is
therefore expected to be limited.

Second, choosing a Budd-type friction law which introduces an elevation-dependence of the basal friction through the
effective pressure adds complexity to the interpretation of our results. Specifically, it means that bed bumps and depressions
460 alter the basal resistance to flow purely through their elevation difference to the surroundings. This may be another reason
why no retreat over bumps was possible in our experiments. Furthermore, while the Budd friction law is one of the most
commonly used ones, previous studies have also shown that different friction laws can lead to substantial differences in transient
ice dynamics and steady-states (Brondex et al., 2017; Åkesson et al., 2021). A comparison between different friction laws
is outside the scope of this study, and thus the impact of this modeling choice is difficult to estimate. Third, the calving
465 parameterization chosen is an important control on the simulated dynamics. Other idealized studies have applied a calving
law with a prescribed ice front position or a prescribed ice shelf length (Schoof et al., 2017; Haseloff and Sergienko, 2018),
which both have the disadvantage of being unknown in general, whereas we opt for the von-Mises parameterization due to its

relatively good performance when applied to real-world glaciers (Choi et al., 2018). Again, in the absence of a universal calving law, the potential effect of this modeling choice on our results is hard to assess. Overall, it can be assumed that the geometric relationship found here is not as distinct in complex real-world settings. To what degree our results are transferable to a specific glacier will depend on the degree to which the stress regime at the glacier front and the grounding line is comparable to our setup. This is what we demonstrated with the example of Jakobshavn Isbræ. Nonetheless, since we use a state-of-the-art model with parameterizations that have been calibrated against typical values for marine outlet glaciers, we expect our findings to be applicable to a wide range of settings.

Many of the medium-sized glacier catchments in Greenland may yield a long-term contribution to sea-level rise as thinning at the outlet glaciers may propagate far upstream (Felixson et al., 2021). Here, we offer a quantitative perspective on the processes at the grounding line, and highlight the importance of assessing both the wetted area and the along-fjord change in wetted area in order to accurately describe ice-topography interactions. These two parameters together determine the geometrically induced ice discharge to the ocean, which is crucial for sea-level rise, and the expected future retreat of marine outlet glaciers.

5 Conclusions

The shape of a fjord can promote or inhibit glacier retreat in response to climate change. Here, we use a numerical model to study such ice-topography interactions in a synthetic setup under idealized conditions. We find that variable fjord topography induces gradients in lateral or basal shear stresses which then influence glacier dynamics. Increased shear at the ice-fjord interface, which stabilizes the model grounding line, is caused by converging flow towards a downstream constriction, because such flow enhances the glacier-fjord contact. Conversely, areas of reduced shear, which promote fast retreat, are found where ice flow diverges, because glacier-fjord contact is reduced. In practical terms, this means that retreat of a glacier into an upstream widening or deepening fjord does not necessarily promote retreat, but may in fact stabilize the glacier. We also confirm that rapid retreat is more likely to occur through deep and wide fjords, while slower retreat is expected for narrow and shallow topographies.

Furthermore, using the concept of the wetted area, which is the submerged cross-sectional area of a fjord, and its along-fjord change, we quantitatively link grounding line discharge and retreat rate to fjord topography. Specifically, we postulate that given the current grounding line flux and the upstream subglacial topography of a particular glacier, an ad-hoc estimate of the topographically induced component of future mass loss can be made. For less well-studied glaciers, we demonstrate that the velocity evolution over time is a promising proxy for the dynamic response of a glacier to fjord topography.

We expect that the quantitative relationships between topography and retreat dynamics are most likely to be transferable to real-world glaciers confined by fjord walls, while being less relevant for ice streams where lateral ice flow influences dynamics considerably. We demonstrate this using the example of Jakobshavn Isbræ.

Future studies should aim to verify our findings using real-world observations. Particularly valuable in this context would be long-term observations of (sub-)annual grounding line positions, calving fronts and velocity changes, combined with detailed bathymetric maps for glaciers confined by fjord walls in Greenland and the Arctic.

Code availability. ISSM 4.16 is freely available for download at <http://issm.jpl.nasa.gov/>. Key scripts needed to perform the simulations done in this study will be made publicly available on GitHub once the manuscript is published.

Author contributions. HÅ, KHN and BdF came up with the idea for the project. TF designed the model setup with significant input from HÅ, BdF and MM. TF carried out and analyzed the experiments with input from all co-authors. TF wrote the manuscript and produced the
505 figures with feedback from all co-authors.

Competing interests. The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements. HÅ was supported by the Swedish Research Council, grant no. 2016-04021. BdF is funded by the Norwegian Research Council (grant 287206). The simulations were performed on resources provided by UNINETT Sigma2 - the National Infrastructure for
510 High Performance Computing and Data Storage in Norway (nn9635k, nn4659k). We thank Karita Kajanto for providing model output on Jakobshavn Isbræ used in this study.




Figure A1. Grounding line position over time for different fjord geometries and magnitudes (for color code refer to Fig. 2). From top to bottom: linear-fjord reference run, embayments, depressions, bottlenecks, bumps




Figure A2. Retreat through *asymmetric* and *longer* perturbations. Annual grounding lines for lateral perturbations (a-d), annual profiles through glaciers for basal perturbations (e,f). Geometries shown are (with names referring to Table 2): a) ByH1800_asy, b) ByH900_lon, c) ByH-900_asy, d) ByH-675_lon, e) BuH-240_lon, f) BuH180_lon. Shaded areas indicate extent of geometric perturbation.




Figure A3. Stress fields in *asymmetric* and *longer* geometries. Lateral shear stress gradients for lateral perturbations (a,b,c,d) and longitudinal stress gradients for basal perturbations (e,f)




Figure A4. v_{GL} over dS correlation for all geometries including asymmetric and longer ones. Differences between this relationship and the relationship Q_{GL}/S over dS originate from time instances when the ice front is grounded well above flotation (and therefore $v_{GL} \not\propto Q_{GL}/S$) and because we measure v_{GL} only along the central flow-line of the glacier.




Figure A5. dGL over S correlation including depressions.

References

- Asay-Davis, X. S., Cornford, S. L., Durand, G., Galton-Fenzi, B. K., Gladstone, R. M., Gudmundsson, G. H., Hattermann, T., Holland, D. M., Holland, D., Holland, P. R., Martin, D. F., Mathiot, P., Pattyn, F., and Seroussi, H.: Experimental design for three interrelated marine ice sheet and ocean model intercomparison projects: MISIMIP v. 3 (MISIMIP +), ISOMIP v. 2 (ISOMIP +) and MISOMIP v. 1 (MISOMIP1), *Geoscientific Model Development*, 9, 2471–2497, <https://doi.org/https://doi.org/10.5194/gmd-9-2471-2016>, publisher: Copernicus GmbH, 2016.
- Bondzio, J. H., Seroussi, H., Morlighem, M., Kleiner, T., Rückamp, M., Humbert, A., and Larour, E. Y.: Modelling calving front dynamics using a level-set method: application to Jakobshavn Isbræ, West Greenland, *The Cryosphere*, 10, 497–510, <https://doi.org/10.5194/tc-10-497-2016>, 2016.
- Bondzio, J. H., Morlighem, M., Seroussi, H., Kleiner, T., Rückamp, M., Mougintot, J., Moon, T., Larour, E. Y., and Humbert, A.: The mechanisms behind Jakobshavn Isbræ’s acceleration and mass loss: A 3-D thermomechanical model study, *Geophysical Research Letters*, 44, 6252–6260, <https://doi.org/10.1002/2017GL073309>, 2017.

- Brancato, V., Rignot, E., Milillo, P., Morlighem, M., Mouginot, J., An, L., Scheuchl, B., Jeong, S., Rizzoli, P., Bueso Bello, J. L., and Prats-
525 Iraola, P.: Grounding Line Retreat of Denman Glacier, East Antarctica, Measured With COSMO-SkyMed Radar Interferometry Data, *Geophysical Research Letters*, 47, e2019GL086291, <https://doi.org/10.1029/2019GL086291>, 2020.
- Briner, J. P., Bini, A. C., and Anderson, R. S.: Rapid early Holocene retreat of a Laurentide outlet glacier through an Arctic fjord, *Nature
Geoscience*, 2, 496–499, <https://doi.org/10.1038/ngeo556>, 2009.
- Briner, J. P., Cuzzone, J. K., Badgeley, J. A., Young, N. E., Steig, E. J., Morlighem, M., Schlegel, N.-J., Hakim, G. J., Schaefer, J. M.,
530 Johnson, J. V., Lesnek, A. J., Thomas, E. K., Allan, E., Bennike, O., Cluett, A. A., Csatho, B., Vernal, A. d., Downs, J., Larour,
E., and Nowicki, S.: Rate of mass loss from the Greenland Ice Sheet will exceed Holocene values this century, *Nature*, 586, 70–74,
<https://doi.org/10.1038/s41586-020-2742-6>, 2020.
- Brinkerhoff, D., Truffer, M., and Aschwanden, A.: Sediment transport drives tidewater glacier periodicity, *Nature Communications*, 8, 1–8,
<https://doi.org/10.1038/s41467-017-00095-5>, 2017.
- 535 Brondeux, J., Gagliardini, O., Gillet-Chaulet, F., and Durand, G.: Sensitivity of grounding line dynamics to the choice of the friction law,
Journal of Glaciology, 63, 854–866, <https://doi.org/10.1017/jog.2017.51>, 2017.
- Budd, W. F., Jessen, D., and Smith, I. N.: A Three-Dimensional Time-Dependent Model of the West Antarctic Ice Sheet, *Annals of Glaciol-
ogy*, 5, 29–36, <https://doi.org/10.3189/1984AoG5-1-29-36>, publisher: Cambridge University Press, 1984.
- Bunce, C., Carr, J. R., Nienow, P. W., Ross, N., and Killick, R.: Ice front change of marine-terminating outlet glaciers in northwest and
540 southeast Greenland during the 21st century, *Journal of Glaciology*, 64, 523–535, <https://doi.org/10.1017/jog.2018.44>, 2018.
- Carr, J. R., Vieli, A., and Stokes, C.: Influence of sea ice decline, atmospheric warming, and glacier width on marine-terminating outlet glacier
behavior in northwest Greenland at seasonal to interannual timescales, *Journal of Geophysical Research: Earth Surface*, 118, 1210–1226,
<https://doi.org/10.1002/jgrf.20088>, 2013.
- Carr, J. R., Stokes, C., and Vieli, A.: Recent retreat of major outlet glaciers on Novaya Zemlya, Russian Arctic, influenced by fjord geometry
545 and sea-ice conditions, *Journal of Glaciology*, 60, 155–170, <https://doi.org/10.3189/2014JoG13J122>, 2014.
- Catania, G. A., Stearns, L. A., Sutherland, D. A., Fried, M. J., Bartholomaeus, T. C., Morlighem, M., Shroyer, E., and Nash, J.: Geometric
Controls on Tidewater Glacier Retreat in Central Western Greenland, *Journal of Geophysical Research: Earth Surface*, 123, 2024–2038,
<https://doi.org/https://doi.org/10.1029/2017JF004499>, 2018.
- Choi, Y., Morlighem, M., Wood, M., and Bondzio, J. H.: Comparison of four calving laws to model Greenland outlet glaciers, *The
550 Cryosphere*, 12, 3735–3746, <https://doi.org/https://doi.org/10.5194/tc-12-3735-2018>, 2018.
- Choi, Y., Morlighem, M., Rignot, E., and Wood, M.: Ice dynamics will remain a primary driver of Greenland ice sheet mass loss over the
next century, *Communications Earth & Environment*, 2, 1–9, <https://doi.org/10.1038/s43247-021-00092-z>, 2021.
- Courant, R., Friedrichs, K., and Lewy, H.: Über die partiellen Differenzgleichungen der mathematischen Physik, *Mathematische Annalen*,
100, 32–74, <https://doi.org/10.1007/BF01448839>, 1928.
- 555 Enderlin, E. M. and Howat, I. M.: Submarine melt rate estimates for floating termini of Greenland outlet glaciers (2000–2010), *Journal of
Glaciology*, 59, 67–75, <https://doi.org/10.3189/2013JoG12J049>, 2013.
- Enderlin, E. M., Howat, I. M., and Vieli, A.: High sensitivity of tidewater outlet glacier dynamics to shape, *The Cryosphere*, 7, 1007–1015,
<https://doi.org/10.5194/tc-7-1007-2013>, 2013.
- Favier, L., Gagliardini, O., Durand, G., and Zwinger, T.: A three-dimensional full Stokes model of the grounding line dynamics: effect of a
560 pinning point beneath the ice shelf, *The Cryosphere*, p. 12, 2012.

- Felikson, D., Bartholomaus, T. C., Catania, G. A., Korsgaard, N. J., Kjær, K. H., Morlighem, M., Noël, B., Broeke, M. v. d., Stearns, L. A., Shroyer, E. L., Sutherland, D. A., and Nash, J. D.: Inland thinning on the Greenland ice sheet controlled by outlet glacier geometry, *Nature Geoscience*, 10, 366–369, <https://doi.org/10.1038/ngeo2934>, 2017.
- Felikson, D., Catania, G. A., Bartholomaus, T. C., Morlighem, M., and Noël, B. P. Y.: Steep Glacier Bed Knickpoints Mitigate Inland Thinning in Greenland, *Geophysical Research Letters*, 48, e2020GL090112, <https://doi.org/10.1029/2020GL090112>, 2021.
- 565 Garbe, J., Albrecht, T., Levermann, A., Donges, J. F., and Winkelmann, R.: The hysteresis of the Antarctic Ice Sheet, *Nature*, 585, 538–544, <https://doi.org/10.1038/s41586-020-2727-5>, number: 7826 Publisher: Nature Publishing Group, 2020.
- Gudmundsson, G. H.: Ice-shelf buttressing and the stability of marine ice sheets, *The Cryosphere*, 7, 647–655, <https://doi.org/10.5194/tc-7-647-2013>, 2013.
- 570 Gudmundsson, G. H., Krug, J., Durand, G., Favier, L., and Gagliardini, O.: The stability of grounding lines on retrograde slopes, *The Cryosphere*, p. 10, <https://doi.org/10.5194/tc-6-1497-2012>, 2012.
- Harrison, W. D., Elsberg, D. H., Echelmeyer, K. A., and Krimmel, R. M.: On the characterization of glacier response by a single time-scale, *Journal of Glaciology*, 47, 659–664, <https://doi.org/10.3189/172756501781831837>, 2001.
- Haseloff, M. and Sergienko, O. V.: The effect of buttressing on grounding line dynamics, *Journal of Glaciology*, 64, 417–431, 2018.
- 575 Haubner, K., Box, J. E., Schlegel, N. J., Larour, E. Y., Morlighem, M., Solgaard, A. M., Kjeldsen, K. K., Larsen, S. H., Rignot, E., Dupont, T. K., and Kjær, K. H.: Simulating ice thickness and velocity evolution of Upernavik Isstrøm 1849–2012 by forcing prescribed terminus positions in ISSM, *The Cryosphere*, 12, 1511–1522, <https://doi.org/10.5194/tc-12-1511-2018>, 2018.
- Hill, E. A., Carr, J. R., and Stokes, C. R.: A Review of Recent Changes in Major Marine-Terminating Outlet Glaciers in Northern Greenland, *Frontiers in Earth Science*, 4, <https://doi.org/10.3389/feart.2016.00111>, 2017.
- 580 Jamieson, S. S. R., Vieli, A., Livingstone, S. J., Cofaigh, C., Stokes, C., Hillenbrand, C.-D., and Dowdeswell, J. A.: Ice-stream stability on a reverse bed slope, *Nature Geoscience*, 5, 799–802, <https://doi.org/10.1038/ngeo1600>, 2012.
- Jamieson, S. S. R., Vieli, A., Cofaigh, C. O., Stokes, C. R., Livingstone, S. J., and Hillenbrand, C.-D.: Understanding controls on rapid ice-stream retreat during the last deglaciation of Marguerite Bay, Antarctica, using a numerical model, *Journal of Geophysical Research: Earth Surface*, 119, 247–263, <https://doi.org/10.1002/2013JF002934>, 2014.
- 585 Kajanto, K., Seroussi, H., de Fleurian, B., and Nisancioglu, K. H.: Present day Jakobshavn Isbræ (West Greenland) close to the Holocene minimum extent, *Quaternary Science Reviews*, 246, 106492, <https://doi.org/10.1016/j.quascirev.2020.106492>, 2020.
- King, M. D., Howat, I. M., Candela, S. G., Noh, M. J., Jeong, S., Noël, B. P. Y., van den Broeke, M. R., Wouters, B., and Negrete, A.: Dynamic ice loss from the Greenland Ice Sheet driven by sustained glacier retreat, *Communications Earth & Environment*, 1, 1–7, <https://doi.org/10.1038/s43247-020-0001-2>, 2020.
- 590 Larour, E., Seroussi, H., Morlighem, M., and Rignot, E.: Continental scale, high order, high spatial resolution, ice sheet modeling using the Ice Sheet System Model (ISSM), *Journal of Geophysical Research: Earth Surface*, 117, <https://doi.org/10.1029/2011JF002140>, 2012.
- MacAyeal, D. R.: Large-scale ice flow over a viscous basal sediment: Theory and application to ice stream B, Antarctica, *Journal of Geophysical Research: Solid Earth*, 94, 4071–4087, <https://doi.org/10.1029/JB094iB04p04071>, 1989.
- Mangerud, J., Goehring, B. M., Lohne, O. S., Svendsen, J. I., and Gyllencreutz, R.: Collapse of marine-based outlet glaciers from the Scandinavian Ice Sheet, *Quaternary Science Reviews*, 67, 8–16, <https://doi.org/10.1016/j.quascirev.2013.01.024>, 2013.
- 595 McNabb, R. W. and Hock, R.: Alaska tidewater glacier terminus positions, 1948–2012, *Journal of Geophysical Research: Earth Surface*, 119, 153–167, <https://doi.org/10.1002/2013JF002915>, 2014.

- Miles, B. W. J., Stokes, C. R., Jenkins, A., Jordan, J. R., Jamieson, S. S. R., and Gudmundsson, G. H.: Intermittent structural weakening and acceleration of the Thwaites Glacier Tongue between 2000 and 2018, *Journal of Glaciology*, 66, 485–495, <https://doi.org/10.1017/jog.2020.20>, 2020.
- 600 Morland, L. W.: Unconfined ice-shelf flow, in: Van der Veen C.J., Oerlemans J. (eds) *Dynamics of the West Antarctic Ice Sheet*, vol. 4 of *Glaciology and Quaternary Geology*, pp. 99–116, Springer, Dodrecht, https://doi.org/10.1007/978-94-009-3745-1_6, 1987.
- Morlighem, M., Bondzio, J., Seroussi, H., Rignot, E., Larour, E., Humbert, A., and Rebuffi, S.: Modeling of Store Gletscher’s calving dynamics, West Greenland, in response to ocean thermal forcing, *Geophysical Research Letters*, 43, 2659–2666, <https://doi.org/10.1002/2016GL067695>, 2016.
- 605 Morlighem, M., Williams, C. N., Rignot, E., An, L., Arndt, J. E., Bamber, J. L., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L., Mouginot, J., Noël, B. P. Y., O’Cofaigh, C., Palmer, S., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., and Zinglensen, K. B.: BedMachine v3: Complete Bed Topography and Ocean Bathymetry Mapping of Greenland From Multibeam Echo Sounding Combined With Mass Conservation, *Geophysical Research Letters*, 44, 11,051–11,061, <https://doi.org/10.1002/2017GL074954>, 2017.
- 610 Morlighem, M., Wood, M., Seroussi, H., Choi, Y., and Rignot, E.: Modeling the response of northwest Greenland to enhanced ocean thermal forcing and subglacial discharge, *The Cryosphere*, 13, 723–734, <https://doi.org/10.5194/tc-13-723-2019>, 2019.
- Motyka, R. J., Hunter, L., Echelmeyer, K. A., and Connor, C.: Submarine melting at the terminus of a temperate tidewater glacier, LeConte Glacier, Alaska, U.S.A., *Annals of Glaciology*, 36, 57–65, <https://doi.org/10.3189/172756403781816374>, publisher: Cambridge University Press, 2003.
- 615 Mouginot, J., Rignot, E., Bjørk, A. A., van den Broeke, M., Millan, R., Morlighem, M., Noël, B., Scheuchl, B., and Wood, M.: Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018, *Proceedings of the National Academy of Sciences*, 116, 9239–9244, <https://doi.org/10.1073/pnas.1904242116>, 2019.
- 620 Nick, F. M., Vieli, A., Howat, I. M., and Joughin, I.: Large-scale changes in Greenland outlet glacier dynamics triggered at the terminus, *Nature Geoscience*, 2, 110–114, <https://doi.org/10.1038/ngeo394>, 2009.
- Nick, F. M., Vieli, A., Andersen, M. L., Joughin, I., Payne, A., Edwards, T. L., Pattyn, F., and van de Wal, R. S. W.: Future sea-level rise from Greenland’s main outlet glaciers in a warming climate, *Nature*, 497, 235–238, <https://doi.org/10.1038/nature12068>, 2013.
- Oerlemans, J. and Nick, F.: A minimal model of a tidewater glacier, *Annals of Glaciology*, 42, 1–6, <https://doi.org/10.3189/172756405781813023>, 2005.
- 625 Oppenheimer, M., Glavovic, B., Hinkel, J., van de Wal, R., Magnan, A., Abd-Elgawad, A., Cai, R., Cifuentes-Jara, M., DeConto, R., Ghosh, T., Hay, J., Isla, F., Marzeion, B., and Sebesvari, Z.: Sea Level Rise and Implications for Low-Lying Islands, Coasts and Communities, in: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate, 2019.
- Pattyn, F. and Morlighem, M.: The uncertain future of the Antarctic Ice Sheet, *Science*, 367, 1331–1335, <https://doi.org/10.1126/science.aaz5487>, 2020.
- 630 Pfeffer, W. T.: A simple mechanism for irreversible tidewater glacier retreat, *Journal of Geophysical Research: Earth Surface*, 112, <https://doi.org/10.1029/2006JF000590>, 2007.
- Raymond, C.: Shear margins in glaciers and ice sheets, *Journal of Glaciology*, 42, 90–102, <https://doi.org/10.3189/S0022143000030550>, 1996.

- 635 Schoof, C.: Ice sheet grounding line dynamics: Steady states, stability, and hysteresis, *Journal of Geophysical Research*, 112, <https://doi.org/10.1029/2006JF000664>, 2007.
- Schoof, C., Davis, A. D., and Popa, T. V.: Boundary layer models for calving marine outlet glaciers, *The Cryosphere*, 11, 2283–2303, 2017.
- Schuler, T. V., Kohler, J., Elagina, N., Hagen, J. O. M., Hodson, A. J., Jania, J. A., Kääb, A. M., Luks, B., Małeckı, J., Moholdt, G., Pohjola, V. A., Sobota, I., and Van Pelt, W. J. J.: Reconciling Svalbard Glacier Mass Balance, *Frontiers in Earth Science*, 8, <https://doi.org/10.3389/feart.2020.00156>, 2020.
- 640 Seroussi, H., Morlighem, M., Larour, E., Rignot, E., and Khazendar, A.: Hydrostatic grounding line parameterization in ice sheet models, *The Cryosphere*, 8, 2075–2087, <https://doi.org/10.5194/tc-8-2075-2014>, 2014.
- Steiger, N., Nisancioglu, K. H., Åkesson, H., de Fleurian, B., and Nick, F. M.: Simulated retreat of Jakobshavn Isbræ since the Little Ice Age controlled by geometry, *The Cryosphere*, 12, 2249–2266, <https://doi.org/10.5194/tc-12-2249-2018>, 2018.
- 645 Straneo, F., Heimbach, P., Sergienko, O., Hamilton, G., Catania, G., Griffies, S., Hallberg, R., Jenkins, A., Joughin, I., Motyka, R., Pfeffer, W. T., Price, S. F., Rignot, E., Scambos, T., Truffer, M., and Vieli, A.: Challenges to Understanding the Dynamic Response of Greenland’s Marine Terminating Glaciers to Oceanic and Atmospheric Forcing, *Bulletin of the American Meteorological Society*, 94, 1131–1144, <https://doi.org/10.1175/BAMS-D-12-00100.1>, 2013.
- van der Veen, C. J.: *Fundamentals of Glacier Dynamics*, CRC Press, <https://doi.org/10.1201/b14059>, 2013.
- 650 Vieli, A., Funk, M., and Blatter, H.: Flow dynamics of tidewater glaciers: a numerical modelling approach, *Journal of Glaciology*, 47, 595–606, <https://doi.org/10.3189/172756501781831747>, 2001.
- Warren, C. R. and Glasser, N. F.: Contrasting Response of South Greenland Glaciers to Recent Climatic Change, *Arctic and Alpine Research*, 24, 124–132, <https://doi.org/10.1080/00040851.1992.12002937>, 1992.
- Weertman, J.: Stability of the Junction of an Ice Sheet and an Ice Shelf, *Journal of Glaciology*, 13, 3–11, <https://doi.org/10.3189/S0022143000023327>, 1974.
- 655 Xu, Y., Rignot, E., Fenty, I., Menemenlis, D., and Flexas, M. M.: Subaqueous melting of Store Glacier, west Greenland from three-dimensional, high-resolution numerical modeling and ocean observations, *Geophysical Research Letters*, 40, 4648–4653, <https://doi.org/10.1002/grl.50825>, 2013.
- Åkesson, H., Morlighem, M., Nisancioglu, K. H., Svendsen, J. I., and Mangerud, J.: Atmosphere-driven ice sheet mass loss paced by topography: Insights from modelling the south-western Scandinavian Ice Sheet, *Quaternary Science Reviews*, 195, 32–47, <https://doi.org/10.1016/j.quascirev.2018.07.004>, 2018a.
- 660 Åkesson, H., Nisancioglu, K. H., and Nick, F. M.: Impact of Fjord Geometry on Grounding Line Stability, *Frontiers in Earth Science*, 6, <https://doi.org/10.3389/feart.2018.00071>, 2018b.
- Åkesson, H., Gyllencreutz, R., Mangerud, J., Svendsen, J. I., Nick, F. M., and Nisancioglu, K. H.: Rapid retreat of a Scandinavian marine outlet glacier in response to warming at the last glacial termination, *Quaternary Science Reviews*, 250, 106645, <https://doi.org/10.1016/j.quascirev.2020.106645>, 2020.
- 665 Åkesson, H., Morlighem, M., O’Regan, M., and Jakobsson, M.: Future projections of Petermann Glacier under ocean warming depend strongly on friction law, *Journal of Geophysical Research: Earth Surface*, 2021.