

The Holocene dynamics of Ryder Glacier and ice tongue in north Greenland

Matt O'Regan^{1,2*}, Thomas M. Cronin³, Brendan Reilly⁴, Aage Kristian Olsen Alstrup⁵, Laura Gemery³, Anna Golub³, Larry A. Mayer⁶, Mathieu Morlighem⁷, Matthias Moros⁸, Ole L. Munk⁵, Johan Nilsson^{2,9},
5 Christof Pearce¹⁰, Henrieka Detlef¹⁰, Christian Stranne^{1,2}, Flor Vermassen^{1,2}, Gabriel West^{1,2}, and
Martin Jakobsson^{1,2}

¹Department of Geological Sciences, Stockholm University, 10691, Stockholm, Sweden

²Bolin Centre for Climate Research, Stockholm University, 10691, Stockholm, Sweden

10 ³Florence Bascom Geoscience Center, U.S. Geological Survey, Reston, VA, 20192, USA

⁴Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA, 92037, USA

⁵Department of Clinical Medicine - Nuclear Medicine and PET, Aarhus University

⁶Center for Coastal and Ocean Mapping, University of New Hampshire, Durham, NH, 03824, USA

⁷Department of Earth System Science, University of California, Irvine, CA, 92697, USA

15 ⁸Leibniz Institute for Baltic Sea Research Warnemünde, D-18119, Rostock, Germany

⁹Department of Meteorology, Stockholm University, 10691, Stockholm, Sweden

¹⁰Department of Geoscience and Arctic Research Centre, Aarhus University, 8000, Aarhus, Denmark

Correspondence to: Matt O'Regan (matt.oregan@geo.su.se)

20 **Abstract.** The northern sector of the Greenland Ice Sheet is considered to be particularly susceptible to ice mass loss arising from increased glacier discharge in the coming decades. However, the past extent and dynamics of outlet glaciers in this region, and hence their vulnerability to climate change, are poorly documented. In the summer of 2019, the Swedish icebreaker *Oden* entered the previously uncharted waters of Sherard Osborn Fjord, where Ryder Glacier drains approximately 2% of Greenland's ice sheet into the Lincoln Sea. Here we reconstruct the Holocene dynamics of Ryder
25 Glacier and its ice tongue by combining radiocarbon dating with sedimentary facies analyses along a 45 km transect of marine sediment cores collected between the modern ice tongue margin and the mouth of the fjord. The results illustrate that Ryder Glacier retreated from a grounded position at the fjord mouth during the Early Holocene ($>10.7 \pm 0.4$ cal ka BP) and receded more than 120 km to the end of Sherard Osborn Fjord by the Middle Holocene (6.3 ± 0.3 cal ka BP), likely becoming completely land-based. A re-advance of Ryder Glacier occurred in the Late Holocene, becoming marine-based
30 around 3.9 ± 0.4 cal ka BP. An ice tongue, similar in extent to its current position was established in the Late Holocene (between 3.6 ± 0.4 and 2.9 ± 0.4 cal ka BP) and extended to its maximum historical position near the fjord mouth around 0.9 ± 0.3 cal ka BP. Laminated, clast-poor sediments were deposited during the entire retreat and regrowth phases, suggesting the persistence of an ice tongue that only collapsed when the glacier retreated behind a prominent topographic high at the landward end of the fjord. Sherard Osborn Fjord narrows inland, is constrained by steep-sided cliffs, contains a number of
35 bathymetric pinning points that also shield the modern ice tongue and grounding zone from warm Atlantic waters, and has a shallowing inland sub-ice topography. These features are conducive to glacier stability and can explain the persistence of

Ryder's ice tongue while the glacier remained marine-based. However, the physiography of the fjord did not halt the dramatic retreat of Ryder Glacier under the relatively mild changes in climate forcing during the Holocene. Presently, Ryder Glacier is grounded more than 40 km seaward of its inferred position during the Middle Holocene, highlighting the potential for substantial retreat in response to ongoing climate change.




Figure 1: Satellite view of north Greenland and Ellesmere Island from 07/30/2019. Velocity of outlet glaciers are from Sentinel-1, winter campaign 2019/2020 [version 1.3] (Nagler et al., 2015). Robeson Channel (RC), Hall Basin (HB), Kennedy Channel (KC), Kane Basin (KB).

1. Introduction

Mass loss from the Greenland Ice Sheet (GrIS) occurs from surface ablation (melting) and through iceberg calving (discharge) and subaqueous melt at marine terminating glaciers. It has increased six-fold since the 1980's, contributing an estimated 13.7 mm to global sea level between 1972-2019 (Mouginot et al., 2019). In north and northeast Greenland (Fig. 1), ice discharge rates from marine terminating glaciers are lower than those observed in the south and northwest (Mouginot et al., 2019). With amplified rates of high Arctic warming, and the continued loss of sea ice and buttressing ice shelves, accelerated ice discharge from the northern sector of the GrIS has been identified as a particular concern for sea-level rise in the coming decades (Moon et al., 2012; Hill et al., 2018; Mouginot et al., 2019).

Marine sediment archives provide unique insights into the past extent and dynamics of Greenland's marine-based ice margin, including the sensitivity and environmental controls on the stability of outlet glaciers and their floating ice tongues

(Jakobsson et al., 2018; 2020; Wangner et al., 2018; Reilly et al., 2019; Vermassen et al., 2020). Combined with analyses of ice cores (Vinther et al., 2009, Lecavalier et al., 2017), terrestrial and marine mapping of glacial limits (Funder et al., 2011a), and other paleoclimate time-series (Briner et al., 2016), marine sediment archives allow us to investigate the response of the
55 GrIS to natural climate variability over time scales that exceed the length of direct satellite observations and other historical records. An understanding of how marine terminating glaciers responded to past climate change, and ultimately elucidating the geologic and environmental controls on their behaviour, are critical to reduce uncertainties in future sea-level predictions (Bamber et al., 2019).

60 After the Last Glacial Maximum, the GrIS receded dramatically through the Early (11.7 – 8.2 cal ka BP) and Middle Holocene (8.2-4.2 cal ka BP) when Arctic summer air temperatures were ~1-3 °C above 20th century averages (Kaufman et al., 2004; Miller et al., 2010; Briner et al., 2016). The GrIS reached its minimum extent near the end of the Middle Holocene (8.2-4.2 cal ka BP) or sometime near the beginning of the Late Holocene (4.2-0 cal ka BP) (Young and Briner 2015). The Lincoln Sea, which surrounds much of northern Greenland hosts some of the most persistent and harsh sea-ice conditions in
65 the Arctic today. However, many terrestrial glacial (Kelly and Bennike, 1992; Landvik et al., 2001; Möller et al., 2010; Funder et al, 2011b; Larsen et al., 2019) and lacustrine paleoclimate studies (Olsen et al., 2012) show warmer climatic conditions, reduced glacial ice extent, and more open water conditions along the north Greenland coast during the Holocene Thermal Maximum (11.0-5.5 cal ka BP). Due to difficulties in accessing the Lincoln Sea, there are no marine records documenting glacier dynamics north of Petermann Glacier (Reilly et al., 2019) that can be combined with these land-based
70 studies.

In the summer of 2019, during the *Ryder 2019* expedition, the Swedish icebreaker *Oden* became the first vessel to enter the unchartered waters of Sherard Osborn Fjord, which connects Ryder Glacier with the Lincoln Sea (Fig. 1). Ryder Glacier drains about 2% of Greenland's ice sheet and is one of four major marine terminating glaciers in this sector of the GrIS (Fig.
75 1). Ryder and Petermann glaciers in the northwest and Nioghalvfjærdsfjord Glacier in the northeast are the only remaining Greenland outlet glaciers that have large, intact floating ice tongues, believed to exert an important buttressing force that slows glacier flow (Mottram et al., 2019). Ryder's ice tongue is 25 kilometers long and has been relatively stable during the last 70 years, showing a net advance of about 43 m a⁻¹ between 1948 and 2015 (Hill et al., 2018).

80 Here we combine radiocarbon dating with the analysis of lithofacies in six marine sediment cores that form a ~45 km long transect extending from the modern ice tongue margin of Ryder Glacier to the mouth of Sherard Osborn Fjord. We integrate these results with established phases of ice recession and re-growth in this sector of northern Greenland, providing the first insights into the dynamic behavior of Ryder Glacier and its ice tongue during the Holocene.

Sherard Osborn Fjord is ~17 km wide and extends ~81 km from the modern grounding zone of Ryder Glacier out towards the Lincoln Sea. Ryder Glacier is currently grounded below sea level, with an ice tongue that extends ~55 km out from the grounding zone (Fig. 2). Bathymetric mapping during the *Ryder 2019* expedition revealed two prominent sills crossing the fjord (Jakobsson et al., 2020) (Fig.2). These sills bound a basin that has a maximum depth of 890 m. The outer sill has little sedimentary cover and appears to be a bedrock feature, while the inner sill is interpreted as a sedimented former glacial grounding zone (Jakobsson et al., 2020), potentially developed on a pre-existing bedrock high.





Figure 2. Landsat image of Sherard Osborn Fjord from 30/07/2019 showing locations of cores discussed in this study. Bedrock geology is from Henriksen et al. (2009). Bathymetry of Sherard Osborn Fjord was collected during the *Ryder 2019* expedition (Jakobsson et al., 2020). Ice limits for the Kap Fulford (yellow) and Warming Land (white) Stades are redrawn from Kelly and Bennike (1992). The extent of Ryder ice tongue in 1917 (pink) is re-drawn from Koch (1928). Numbers refer to key locations of radiocarbon dates used by Kelly and Bennike (1992) to constrain Holocene ice margin positions (Table 1). Dashed cyan line (x-x') marks the location of the oceanographic profile in Fig. 3.

95 The sill depths on the outer fjord are 475 m on the east and 375 m on the west (Fig. 2). The inner sill has a ~6.2 km wide central region that ranges in depth from 193 to 300 m, with a ~1 km wide channel on the eastern side that extends to a depth of 390 m. The modern ice tongue terminus of Ryder Glacier is located approximately 5 km landward of the inner sill. Despite these bathymetric barriers, waters of Atlantic origin that circulate through the Lincoln Sea are found between the two sills at depths greater than 350 m (Fig. 3; Jakobsson et al., 2020). These relatively warm (>0.3°C) and saline (>34.7) waters are constrained by the inner sill with only a small amount of warmer water flowing across the inner sill through the <1 km wide and 390 m deep channel, to be strongly mixed with glacially derived meltwater.

100




Figure 3. Potential temperature and salinity profiles illustrating the distribution of Atlantic water in Sherard Osborn Fjord. Vertical black lines are positions of CTD casts conducted during *Ryder 19*. Yellow circles are the locations of coring sites. The position of site 10 on the outer sill is projected, as this profile does not cross its location (see supplementary for additional details).

105 Sherard Osborn Fjord cuts into the Paleozoic Franklinian Basin that extends over 2000 km from the Canadian Arctic to eastern north Greenland (Henriksen and Higgins, 2000). Along the north Greenland coast, deposition in the Franklinian Basin occurred in a passive margin environment through the latest Precambrian to Devonian, with shallow water carbonate shelf sediments found to the south and siliciclastic slope and deep-water sediments in the north (Henriksen et al., 2009). Exposed sedimentary bedrock between the present GrIS margin and the terminus of the floating ice tongue of Ryder Glacier is composed of Cambrian and Silurian carbonate shelf sediments. Further seaward, Silurian sands and siltstones deposited in a deeper water turbiditic environment are found on the lowlands of Nyeboe and Wulff Land (Henriksen and Higgins, 2000). Older, deformed Cambrian deep-water calcareous mudstones and sandy turbidites cap the outer headlands of Nyeboe and
 110 Wulff Land, and are part of the 600 km long E-W trending north Greenland fold belt that formed near the end-Devonian Ellesmerian Orogeny (Higgins et al., 1998) (Fig. 2).

115 Terrestrial mapping of glacial landforms and marine limits across Nyeboe Land, Warming Land and Wulff Land has defined three regional glacial events from the end of the Pleistocene (Late Weichselian) through the Holocene (Kelly and Bennike, 1992). The oldest marginal ice limits that are traced across north Greenland belong to the Kap Fulford Stade (Kelly and Bennike, 1992). During this time, the GrIS extended across the southern parts of many peninsulas, with outlet glaciers flowing to the outer limits of many of the fjords. Radiocarbon dates from marine macrofossils sampled by Kelly and Bennike at Kap Fulford (Station 31; 10030 ± 175 and 9390 ± 90 ^{14}C a BP) and freshwater algae from northern Wulff Land (Station 43; 10480 ± 90 ^{14}C a BP) provide minimum age estimates for the retreat of glacial ice of the Kap Fulford Stade to the latest 120 Pleistocene to Early Holocene (>10.5 cal ka BP) (Kelly and Bennike, 1992) (Table 1, Fig. 2).

125 **Table 1. Key radiocarbon dates originally used to constrain the ages of glacial stages in the vicinity of Sherard Osborn Fjord. Index numbers refer to original site numbers given in Kelly and Bennike (1992) with locations shown on Figure 2. Details of the sampling sites are from Kelly and Bennike (1985) and Bennike and Kelly (1987). Ages were re-calibrated with IntCal 20 (41, 43) or Marine20 using a ΔR 300 ± 300 , which is consistent with the new calibrated radiocarbon dates presented in this manuscript. Calibrated ages are rounded to 10 years.**

| # | Area | Relevance | Lab. ID | ^{14}C age (a) | Mean and 1- σ range (cal a BP) |
|---|--------------|--------------------|----------|-------------------------|---------------------------------------|
| 31 | Nyeboe Land | Kap Fulford Stade | K-4339 | 10030 ± 175 | 10830 ± 490 |
| Shells of <i>Portlandia arctica</i> from marine silts at an elevation of 87 m | | | | | |
| 31 | Nyeboe Land | Kap Fulford Stade | HAR-6289 | 9390 ± 90 | 9690 ± 400 |
| Shells of <i>Hiatella arctica</i> from marine silts at 92-93 m | | | | | |
| 34 | Nyeboe Land | Steensby Stade | K-4380 | 4870 ± 80 | 4560 ± 410 |
| Shells of <i>Mya truncata</i> and <i>H. arctica</i> from marine silts at 24 m | | | | | |
| 36 | Warming Land | Warming Land Stade | HAR-6290 | 8210 ± 120 | 8210 ± 370 |
| Shells of <i>M. truncata</i> and <i>H. arctica</i> from outcropping marine silts at 68-70 m, 12 km behind ice front | | | | | |
| 40 | Warming Land | Warming Land Stade | HAR-6287 | 6480 ± 100 | 6420 ± 350 |
| Shells of <i>M. truncata</i> from marine silts at 26 m that are younger than ice margin deposits | | | | | |
| 41 | Wulff Land | Steensby Stade | K-4573 | 5100 ± 130 | 5830 ± 170 |
| <i>Salix twigs</i> from a the base of sandy peat at 275 m. Station located 100 m from margin of outlet glacier | | | | | |
| 43 | Wulff Land | Kap Fulford Stade | GU-2588 | 10480 ± 90 | 12380 ± 270 |
| Filamentous green algae from a silt lens embedded in gravelly diamicton at 72 m | | | | | |
| 44 | Wulff Land | Warming Land Stade | K-4374 | 8000 ± 115 | 7990 ± 330 |
| Shells of <i>M. truncata</i> and <i>H. arctica</i> from a sand layer overlying folded silt and sand at 62 m | | | | | |

130 A second set of ice marginal deposits are found along many of the fjords, 20-60 km inland from the Kap Fulford Stade. These are assigned to the Warming Land Stade, representing a standstill of outlet glaciers draining the retreating ice front. Regionally, the age of the Warming Land Stade is bracketed between >9.5 to 8.0 cal ka BP (Kelly and Bennike, 1992). Many glacially dammed lakes formed during the Warming Land Stade, as glaciers continued to occupy the fjord systems while the

ice margin retreated on land (Kelly and Bennike, 1992). For example, an ice-dammed lake formed on Wulff Land, in the central lowlands between Ryder and C. H. Ostenfeld glaciers (Fig. 1). The most proximal minimum age constraints for the Warming Land Stade in Sherard Osborn Fjord come from western Warming Land, where marine macrofossils in sediments younger than the ice margin deposits provide an age of 8210 ± 120 ^{14}C a BP (Station 36, Kelly and Bennike, 1992). On southeastern Warming Land molluscs from a marine silt returned an age of 6480 ± 100 ^{14}C a BP (Station 40, Kelly and Bennike, 1992) implying that ice had retreated further inland than this position by this time. On the eastern edge of Wulff Land outlet glaciers built a series of deltas in the ice-dammed lake during the Warming Land Stade. Marine macrofossils in the lowest delta are dated to 8000 ± 115 ^{14}C a BP (Station 44) indicating that the retreat of C. H. Ostenfeld Glacier and drainage of the glacial lake occurred after this time (Kelly and Bennike, 1992). The ice margin continued to retreat following the Warming Land Stade. In the vicinity of Ryder Glacier, it likely reached a position equivalent to its modern one by the Middle Holocene ~ 6 cal ka BP, before receding even further inland (Kelly and Bennike, 1992).

The Steensby Stade marks the most recent re-advance of the ice margin, outlet glaciers and local ice caps to their maximum positions since the Kap Fulford Stade. Its onset is poorly dated and occurred sometime during neoglacial cooling following the Holocene Thermal Maximum. At the GrIS margin to the north of Ryder Glacier, peat deposits over which the ice margin advanced provide an age of 5100 ± 130 ^{14}C a BP (Station 41), while at Steensby Glacier, reworked marine macrofossils in lateral moraines yield an age of 4870 ± 80 ^{14}C a BP (Station 34; Kelly and Bennike, 1992). Despite uncertainty surrounding the onset of the Steensby Stade, and subsequent dynamics of Ryder Glacier through the Late Holocene, the maximum extent is believed to coincide with historical observations that place the terminus of outlet glaciers and ice tongues near the mouth of many of the major fjords. This is true for Ryder's ice tongue, which was positioned near the outer margin of Sherard Osborn Fjord in 1917 by the Danish geologist and explorer Lauge Koch (Koch, 1928) (Fig. 2). Between 1917 and 1947, Ryder ice tongue retreated to near its current position (Davies and Krinsley, 1962; Higgins, 1990), and has remained relatively stable, even exhibiting a net advance of 43 m a^{-1} between 1948-2015 (Hill et al., 2018).

155 3. Materials and methods

3.1 Marine sediment cores

Sediment cores from Sherard Osborn Fjord were collected during the 37-day *Ryder 2019* expedition (August 5-September 10, 2019). Between August 13 and 25, unusually light sea-ice conditions allowed *Oden* to occupy nine coring stations and systematically map the fjord (Fig. 2; Jakobsson et al., 2020). Coring was conducted using a 12 m long piston core (PC) and small (1-2 m) trigger weight core (TWC), and a 6 m long gravity core (GC). Piston and gravity cores were collected in liners with an inner/outer diameter of 100/110 mm, while TWCs were collected in narrower (80/88mm) liners.

Table 2. Locations, water depths and lengths of cores used in this study.

| Station | Core ID | Latitude (°N) | Longitude (°E) | Water Depth (m) | Core Length (m) |
|---------|----------------|---------------|----------------|-----------------|-----------------|
| 6 | Ryder19-6-GC1 | 82.0095 | -51.7408 | 633 | 4.93 |
| 7 | Ryder19-7-GC1 | 81.9532 | -51.5760 | 551 | 5.19 |
| | Ryder19-7-PC1 | 81.9518 | -51.5878 | 559 | 8.96 |
| | Ryder19-7-TWC1 | 81.9518 | -51.5878 | 559 | 0.96 |
| 8 | Ryder19-8-GC1 | 81.8947 | -51.1365 | 228 | 4.98 |
| | Ryder19-8-PC1 | 81.8928 | -51.1315 | 238 | 8.97 |
| 9 | Ryder19-9-GC1 | 81.8843 | -50.9848 | 271 | 5.89 |
| | Ryder19-9-PC1 | 81.8908 | -50.9682 | 274 | 8.72 |
| 10 | Ryder19-10-GC1 | 82.2713 | -52.0165 | 272 | 2.87 |
| 11 | Ryder19-11-GC1 | 82.2682 | -52.5038 | 208 | 1.34 |

165

Here we focus on cores obtained from five stations that were positioned on topographic highs, with two radiocarbon dates coming from a sixth station on the outer sill. The five main stations form a 45 km long transect extending from the edge of the modern ice tongue to the outer sill of Sherard Osborn Fjord (Table 2, Fig. 2). Sub-bottom profiles were acquired across the coring sites with *Oden's* Kongsberg SBP120 (3°x3°) chirp sonar using a 2.5-7 kHz pulse. These profiles show that a relatively thin (<10-15 m) drape of sediments exists on top of the acoustic basement (Fig. 4). This basement surface could either be sedimentary bedrock, or highly consolidated sedimentary units like till. Hogan et al. (2020) illustrated that in Petermann Fjord, lower frequency air gun seismic data was required to differentiate bedrock from more lithified sediments and till.

170




Figure 4. Sub-bottom profiles across coring stations in Sherard Osborn Fjord illustrating the amount of the generally thin sedimentary cover on top of acoustic basement penetrated by each core.

The unsplit sediment cores were allowed to equilibrate to room temperature (~20°C) and logged shipboard on a Geotek multi-sensor core logger (MSCL). The MSCL provided downcore measurements (1 cm sampling interval) of bulk density (using gamma-ray attenuation) and magnetic susceptibility. Bulk density is controlled by sediment porosity and grain composition (grain density). In predominantly lithogenic normally consolidated marine sediments, variability in bulk density is a function of grain size-driven changes to porosity (coarser sediments = lower porosity = higher bulk density). Magnetic susceptibility is a useful proxy for the input of siliciclastic versus biogenic/organic sediments, and in dominantly lithogenic sediments can be used to discriminate between provenance and grain size (Hatfield et al., 2017; Reilly et al., 2019). Gamma-ray attenuation was measured using a ¹³⁷Cs source with a 5 mm collimator and a 10 s count time. Magnetic susceptibility was measured with a 125 mm Bartington loop sensor using a 1 s acquisition time. An empirical volume correction was applied to account for the difference between the core diameter (D_C) and the loop diameter (D_L). The volume-specific magnetic susceptibility (κ, 10⁻⁵ SI) is defined as,

$$\kappa = \kappa_{\text{(uncorrected)}} / \kappa_{\text{(rel)}}$$

where

$$\kappa_{\text{(rel)}} = 4.8566 (D_C / D_L)^2 - 3.0163 (D_C / D_L) - 0.6448$$

The mass-specific magnetic susceptibility (χ) was calculated by dividing κ by the bulk density.

After logging, piston and gravity cores were split into working and archive half sections. Archive halves were described and imaged using a digital line-scanning camera on the MSCL, while the working half was sampled for shore-based analyses. Split sections were wrapped in plastic film, placed in D-tubes and stored in a refrigerated container (4°C) until they were returned to Stockholm University's refrigerated core storage facility. TWC's remained unsplit and placed directly into the refrigerated container.

Following the expedition, all archive halves were analysed on an Itrax X-ray fluorescence (XRF) core scanner at the Department of Geological Sciences, Stockholm University. Measurements were performed with a Mo tube set to 55 kV and 50 mA with a downcore measurement resolution of 2 mm and a counting time of 15 s. Here we only present the raw counts of calcium (Ca) as a proxy for detrital input from the surrounding Cambrian and Silurian carbonate bedrock terrain.

Computed tomography (CT) scanning was performed on selected archive sections and unsplit TWC's at the Department of Nuclear Medicine & PET of Aarhus University Hospital in Denmark using a Siemens Biograph Vision 600 PET/CT. The CT scan parameters were: 120 kV, 400 Eff mAs (no CARE Dose4D), 200 mm field of view, 0.6 mm slice thickness, filter kernel B60s (sharp). Images were processed using SedCT MATLAB tools to obtain a coronal slice through the central region of each core at an effective pixel resolution of 0.5 x 0.5 mm (Reilly et al., 2017). Clasts with >1 mm diameter were automatically counted from the 3-dimensional CT data, binned into 2 cm intervals, and normalized by core volume using the

algorithm of Reilly et al. (2019) which provides an objective proxy for ice rafted debris (IRD) concentration in glaciomarine lithofacies deposited beyond the limit of grounded ice.

Twenty-nine grain size measurements were made on core 6-GC to broadly characterize grain size spectra in different lithologic units. Measurements were performed using a Malvern Mastersizer 3000 laser diffraction particle size analyzer at the Department of Geological Sciences, Stockholm University. Wet samples were immersed in a dispersing agent (<10% sodium hexametaphosphate solution) and placed in an ultrasonic bath to disaggregate particles before being poured into the Malvern measurement chamber.

3.2 Composite depth scales

The first attempts at coring in Sherard Osborn Fjord illustrated that near surface sediments were very soft, and the top of the coring tools were sinking below the seafloor resulting in variable the recovery of near surface sediments between the gravity, piston and trigger weight cores. At stations 7, 8 and 9, multiple coring tools were deployed, and sediments were recovered using all of these devices. A composite depth scale was developed for each station to ensure that measurements from the different cores could be integrated. The composite depth scales were developed through correlation of the MSCL bulk density, magnetic susceptibility and XRF-scanning (mainly Ca content) data and additionally through visual correlation using the CT-images (*supplementary material Table S1*).

The piston core at each station was used as the reference depth scale. Depths of the gravity and trigger weight cores were stretched, or compressed, between tie points to the piston cores. At stations 7 and 9, the piston core recovered sediments closer to the seafloor than the gravity or trigger weight cores. At station 8, correlation of MSCL and XRF data indicate that the gravity core recovered 40 cm of near-surface sediments that were either not recovered or lost from the top of the piston core.

3.3 Radiocarbon dating

Radiocarbon measurements (^{14}C) were made on 48 samples (Table 3). Shipboard and post-cruise sampling focused on constraining the ages for lithologic units. Most radiocarbon dates were obtained on 300 specimens of the benthic foraminifera *Cassidulina neoteretis* (see Cronin et al. (2019) and Cage et al. (2021) for discussions of this foraminiferal species in the Arctic and Nordic seas respectively). Sixteen dates were obtained on mixed benthic foraminifera (including *C. neoteretis*, *C. reniforme*, *O. tener*, *E. excavatum clavata*) and one date on *Cibicides lobatulus* (Table 3). Additionally, four dates were made on specimens of the planktic foraminifera *Neogloboquadrina pachyderma*. All radiocarbon measurements were performed at the National Ocean Sciences Accelerator Mass Spectrometry (NOSAMS) facility at Woods Hole Oceanographic Institution, Massachusetts, USA. Calibration of the radiocarbon dates was performed using Oxcal v. 4.4 (Bronk Ramsey, 2009) and the Marine20 calibration curve (Heaton et al., 2020).

There is no *a priori* information on the local marine reservoir correction (ΔR) for Sherard Osborn Fjord or the Lincoln Sea. Estimates of ΔR for nearby regions vary widely and have been derived using different marine calibration curves. Results from three pre-bomb living molluscs collected in the vicinity of Thule, suggest a ΔR of 5 +/- 50 years (Mörner and Funder, 1990). Kelly and Bennike (1992) applied 150-year ΔR for marine macrofossils from the region surrounding Ryder Glacier, as suggested for areas of northernmost Greenland by Funder et al. (1982). However, Coulthard et al. (2010) found an average ΔR of 335±85 years (using Marine09; Reimer et al., 2009) based on 24 molluscs from the northwestern Canadian Arctic Archipelago, which includes the northern and western coasts of Ellesmere Island. Reilly et al. (2019) argued that a ΔR of 770 years (using Marine13; Reimer et al., 2013) provided the best fit between a stacked paleosecular variation record from Petermann Fjord sediments and a North Atlantic reference curve. This large ΔR was also consistent with offsets between radiocarbon dates on planktic and benthic foraminifera and ^{210}Pb derived chronologies for the past 100 years. In summary, estimates used in the literature range from 0–770 years. In this study we have applied the new Marine20 calibration curve (Heaton et al., 2020), which results in ages that are ~150 years younger than equivalent Holocene ^{14}C ages calibrated using Marine09 or Marine13. Taking this into consideration we have applied a ΔR of 300±300 years. This broad range provides a large uncertainty envelope. The upper bound approaches the older offset found by Reilly et al. (2019) for Petermann Fjord (770 years using Marine13, which equates to ~620 years using Marine20) and the lower bound the 150 years (which equates to ~0 years using Marine20) commonly used to reconstruct terrestrial ice margins in the area (Kelly and Bennike 1992; Young and Briner, 2015). The mean calibrated ages obtained using a ΔR of 300±300 years (equivalent to ~450 years using Marine13) provide a suitable estimate for sites that are influenced by Atlantic waters. This dependency on water mass is one of the underlying problems in determining an applicable local reservoir correction. For example, the paired benthic/planktic foraminifera samples we ran revealed offsets of 470-570 years in 6-GC and 7-PC which are currently bathed in Atlantic waters, and 0 to 200 years at 10-GC which lies closer to the mixed surface layer (Fig. 3; Table 3). Additional work is needed to resolve the issue of local marine reservoir offsets in the region, their dependency on water mass, and how they have may have changed through time.

265

Table 3. Raw and calibrated radiocarbon dates and the type of material analysed. All data were calibrated using a ΔR 300 ± 300 years. Asterix highlight ages deemed outliers. All the outliers came from samples with low microfossil abundance where mixed benthic assemblages were dated.

| # | Lab ID | Sample ID | Depth (cm) | Comp. depth (cm) | ^{14}C Age (a) | $\delta^{13}\text{C}$ (‰) | Mean (cal a BP) | 1- σ (cal a BP) | Material dated |
|---|--------|------------------|------------|------------------|-------------------------|---------------------------|-----------------|------------------------|----------------------|
| 1 | 152168 | 6-GC-2, 3-6 | 44.5 | 44.5 | 1730±75 | -0.3 | 850 | 540-1150 | <i>C. neoteretis</i> |
| 2 | 156282 | 6-GC-2, 70-75 | 112.5 | 112.5 | 4690±20 | 0.98 | 4330 | 3970-4770 | mixed benthic |
| 3 | 156283 | 6-GC-2, 93-95, A | 134 | 134 | 6870±30 | -0.13 | 6830 | 6510-7190 | mixed benthic |

| | | | | | | | | | |
|-----|--------|------------------|-------|-------|-----------|-------|-------|-------------|----------------------------|
| 4 | 156284 | 6-GC-2, 93-95, B | 134 | 134 | 6400±25 | 0.2 | 6330 | 5990-6680 | <i>N. pachyderma</i> |
| 5 | 152196 | 6-GC-3, 101-105 | 295 | 295 | 8480±40 | -0.83 | 8550 | 8180-8920 | <i>C. neoteretis</i> |
| 6 | 152197 | 6-GC-CC | 494 | 494 | 9030±35 | -0.43 | 9180 | 8760-9550 | <i>C. neoteretis</i> |
| 7* | 156272 | 7-PC-1, 65-72 | 68.5 | 68.5 | 6790±35 | -0.67 | 6750 | 6440-7120 | mixed benthics |
| 8 | 152208 | 7-PC-1, 103-109 | 106 | 106 | 2670±20 | 0.33 | 1850 | 1470-2210 | mixed benthics |
| 9 | 156294 | 7-PC-1, 103-115 | 109 | 109 | 2620±45 | -0.12 | 1790 | 1410-2130 | mixed benthics |
| 10 | 152169 | 7-GC-2, 32-34 | 101 | 140 | 2890±50 | -0.62 | 2100 | 1720-2480 | <i>C. neoteretis</i> |
| 11 | 152198 | 7-TWC-CC | 97 | 164 | 3260±25 | -0.73 | 2540 | 2160-2930 | <i>C. neoteretis</i> |
| 12 | 156309 | 7-PC-2, 23-25 | 175 | 175 | 4060±30 | -0.54 | 3540 | 3150-3940 | mixed benthics |
| 13 | 152207 | 7-GC-2, 85-87 | 154 | 197 | 5940±25 | 1.2 | 5820 | 5520-6190 | <i>Cibicides lobatulus</i> |
| 14 | 152171 | 7-PC-2, 63-65, A | 215 | 215 | 7090±70 | -0.7 | 7050 | 6750-7420 | <i>C. neoteretis</i> |
| 15 | 153807 | 7-PC-2 63-65, B | 215 | 215 | 6520±35 | -0.54 | 6470 | 6140-6830 | <i>N. pachyderma</i> |
| 16 | 156293 | 7-GC-3, 114-118 | 335 | 415 | 8060±85 | -1.62 | 8070 | 7710-8380 | mixed benthics |
| 17* | 156310 | 7-PC-3, 122-126 | 424 | 424 | 9270±45 | -1 | 9500 | 9100-9920 | mixed benthics |
| 18* | 156308 | 7-GC-3, 125-127 | 345 | 428 | 9350±45 | -1.16 | 9610 | 9250-10050 | mixed benthics |
| 19 | 152172 | 7-PC-4, 143-145 | 595.5 | 595.5 | 8410±100 | -0.91 | 8470 | 8070-8850 | <i>C. neoteretis</i> |
| 20 | 152170 | 7-GC-CC | 520 | 611 | 8900±100 | -1.06 | 9020 | 8630-9430 | <i>C. neoteretis</i> |
| 21 | 152209 | 7-PC-5, 103-105 | 706.5 | 706.5 | 9460±45 | -0.72 | 9750 | 9390-10170 | <i>C. neoteretis</i> |
| 22 | 152173 | 7-PC-5, 143-145 | 746.5 | 746.5 | 9210±95 | -0.72 | 9420 | 9010-9850 | <i>C. neoteretis</i> |
| 23* | 156295 | 7-PC-6, 76-78, A | 830.5 | 830.5 | 11000±160 | -0.96 | 11850 | 11410-12380 | mixed benthics |
| 24 | 152210 | 7-PC-6, 76-78, B | 830.5 | 830.5 | 10200±40 | -0.28 | 10730 | 10330-11160 | <i>C. neoteretis</i> |
| 25 | 154575 | 8-GC-2, 102-104 | 149 | 180 | 3290±25 | 1.46 | 2580 | 2200-2970 | <i>C. neoteretis</i> |
| 26 | 152211 | 8-GC-3, 102-104 | 301 | 341 | 4090±20 | N/A | 3580 | 3190-3980 | <i>C. neoteretis</i> |
| 27 | 156285 | 8-PC-3, 61-63, B | 362.5 | 400.5 | 4480±20 | -0.04 | 4070 | 3680-4480 | <i>C. neoteretis</i> |
| 28* | 156311 | 8-PC-3, 61-63, A | 362.5 | 400.5 | 9280±35 | -0.31 | 9520 | 9120-9930 | mixed benthics |
| 29* | 156312 | 8-PC-3, 63-65 | 364.5 | 402.5 | 6310±30 | 0.32 | 6230 | 5900-6580 | mixed benthics |
| 30 | 156288 | 8-PC-3, 73-75 | 374.5 | 412.5 | 5000±20 | -1.03 | 4720 | 4370-5170 | mixed benthics |
| 31 | 152213 | 8-PC-3, 83-85 | 384.5 | 422.5 | 5260±30 | -0.27 | 5040 | 4710-5460 | <i>C. neoteretis</i> |
| 32 | 152212 | 8-GC-4, 62-64 | 411.5 | 495 | 7220±40 | -0.37 | 7190 | 6900-7540 | <i>C. neoteretis</i> |
| 33 | 152174 | 8-GC-CC | 499 | 609 | 7690±65 | -0.85 | 7680 | 7380-8000 | <i>C. neoteretis</i> |

| | | | | | | | | | |
|----|--------|---------------------|-------|-------|-----------|-------|-------|-------------|----------------------|
| 34 | 152298 | 8-PC-5, 46-49 | 648 | 686 | 9140±40 | -0.75 | 9330 | 8940-9750 | <i>C. neoteretis</i> |
| 35 | 156313 | 8-PC-5, 68-72 | 670.5 | 708.5 | 10400±70 | -0.62 | 11010 | 10550-11450 | mixed benthics |
| 36 | 152279 | 9-TWC-CC | 101 | 201 | 3550±55 | -0.31 | 2900 | 2550-3320 | mixed benthics |
| 37 | 152300 | 9-PC-4, 103-105 | 556.5 | 556.5 | 4380±25 | -0.33 | 3940 | 3560-4350 | <i>C. neoteretis</i> |
| 38 | 152280 | 9-PC-4, 123-125 | 576.5 | 576.5 | 6620±65 | -0.47 | 6570 | 6250-6940 | <i>C. neoteretis</i> |
| 39 | 152299 | 9-GC-CC | 590 | 650 | 7780±40 | -0.5 | 7770 | 7460-8080 | <i>C. neoteretis</i> |
| 40 | 152281 | 9-PC-6, 53-55 | 810 | 810 | 8370±110 | -0.79 | 8420 | 8010-8780 | <i>C. neoteretis</i> |
| 41 | 156287 | 10-GC-1, 50-54 | 50 | 50 | 2450±20 | -0.05 | 1600 | 1250-1950 | mixed benthic |
| 42 | 152301 | 10-GC-1, 102-104, A | 103 | 103 | 6750±35 | -0.6 | 6710 | 6380-7070 | <i>C. neoteretis</i> |
| 43 | 153806 | 10-GC-1, 102-104, B | 103 | 103 | 6750±30 | 0.15 | 6710 | 6390-7070 | <i>N. pachyderma</i> |
| 44 | 152302 | 10-GC-2, 132-134 | 279.5 | 279.5 | 7880±45 | -0.34 | 7880 | 7560-8190 | <i>C. neoteretis</i> |
| 45 | 152282 | 10-GC-CC, A | 288 | 288 | 8270±90 | -0.59 | 8310 | 7910-8640 | <i>C. neoteretis</i> |
| 46 | 153808 | 10-GC-CC, B | 288 | 288 | 8060±35 | -1.61 | 8060 | 7720-8370 | <i>N. pachyderma</i> |
| 47 | 152303 | 11-GC-1, 12-14 | 13 | 13 | 7540±30 | -0.3 | 7520 | 7240-7860 | <i>C. neoteretis</i> |
| 48 | 152283 | 11-GC-1, 102-104 | 103 | 103 | 11150±250 | -0.24 | 12010 | 11580-12610 | <i>C. neoteretis</i> |

4. Results

270 4.1 Lithostratigraphic Units

Data from the MSCL, XRF-scanning data, and CT imaging are used to identify six major lithologic units that are correlated from the fjord mouth (10-GC) to the inner bathymetric sill lying seaward of the modern ice tongue (cores 8-PC and 9-PC) (Figs. 5 and 6). A sixth core, 11-GC, located on the outer sill in the shallowest water depth (208 m), did not contain the same lithostratigraphic units, and only the basal radiocarbon date is used as a constraint for Early Holocene ice retreat.

275

LU6: The lowermost lithologic unit (*LU6*) was recovered in 7-PC, 8-PC and 9-PC. It has moderately high but variable Ca contents and is a coarse-grained, poorly sorted diamicton containing abundant gravel (2-64 mm) and cobble-sized (64-256 mm) clasts with a high bulk density ($> 2 \text{ g/cm}^3$) (Figs. 5, 7). The mass-specific magnetic susceptibility is considerably higher in *LU6* compared to the overlying *LU5*, generally exceeding $80\text{-}100 \times 10^{-8} \text{ m}^3/\text{kg}$. Less than 5 cm of this unit were recovered in 9-PC, with greater recovery in 7-PC (54 cm) and 8-PC (>135 cm). The upper boundary with *LU5* is abrupt in 8-PC, gradual in 7-PC. In 9-PC, the thin *LU6* and lowermost *LU5* sediments are laminated, deformed and contain two large

280

dropstones, making it difficult to determine the nature of the boundary. The LU6/LU5 boundary is marked by a pronounced up-core decrease in bulk density and mass-specific magnetic susceptibility. Two subunits (LU6a and LU6b) are recognized in the longer sequences from 7-PC and 8-PC. These subunits are separated by a sharp boundary (Figs. 6, 7). Sediments from LU6a have a lower bulk density (2.0-2.2 g/cm³) compared to LU6b (2.2-2.5 g/cm³) and lower clast abundance. LU6b is a massive clast-supported diamicton, while LU6a contains lenses or intervals of visibly deformed fine-grained layers. Given the succession from a massive to a deformed diamicton, we interpret these units as being deposited beneath grounded ice (Figs. 6, 7).

290 *LU5*: The defining characteristic of LU5 are pronounced mm- to cm-scale laminations and absence of bioturbation. Laminae are sometimes visible in the split core sections as alternating reddish-brown and lighter tan-colored laminae (Fig. 6). The laminations are best defined by variations in Ca and bulk density. Where visible to the naked eye, the lighter tan-colored laminations are enriched in Ca and denser (lighter in CT-scanning images) (Fig. 6). Ca-enriched layers are skewed towards medium and coarse silts, while laminae with lower Ca abundance are skewed towards fine silt and clay (Fig. 8). The higher
295 bulk density of the light-colored laminae likely is caused by the coarser grain size and higher detrital carbonate content, which has a higher grain density compared to quartz and clay minerals. LU5 has a low abundance of irregularly spaced clasts >1 mm that are more prevalent near the base of the unit (Fig. 5). In cores 10-GC, 7-PC, and 6-GC, there is a notable up-core decrease in the bulk density, Ca-content and laminae thickness of LU5. Mass-specific magnetic susceptibility also increases up-core, as the relative Ca-abundance decreases, consistent with a larger contribution from siliciclastic sediments. In 6-GC
300 and 7-PC, these long-term trends are interrupted by a 0.50-1.0 m thick interval with elevated Ca contents and thicker laminae, found near the top of the unit.

At the two stations from the inner sill, 8-PC (238 mbsl) and 9-PC (274 mbsl), LU5 is considerably thinner than in 6-GC (633 mbsl) and 7-PC (559 mbsl), which lie further seaward, and in deeper water depths (Fig. 5). CT images illustrate that laminations in much of LU5 are wavy to lenticular in nature (Figs. 6, 7). Throughout LU5, the laminations are disrupted by numerous small-scale normal faults with occasional evidence of truncation suggesting erosion. This is common throughout LU5 in 8- and 9-PC and very common in the lowermost 1-2 m of 7-PC (*see example from 9-PC-6 in Fig. 7 and supplementary information for additional examples*). In 6-GC and 7-PC, faulting in LU5 is more pronounced than in the overlying laminated sequence of LU4 (Figs. 6, 7). Laminations become less convoluted and more planer towards the top of
305 LU5. On the outer sill (10-GC), LU5 is also comparatively thin, containing highly fractured and wispy laminations that are truncated or eroded in some intervals. The base of LU5 in 10-GC is dominantly composed of sand and gravel sized material.




Figure 5. Summary of stratigraphy and correlation of lithologic units between coring stations in Sherard Osborn Fjord. Locations and uncalibrated ages of radiocarbon dates are shown. Key horizons in LU5 and LU4 that can be correlated between 7-PC, 8-PC and 9-PC are illustrated by thin black lines. Depths for lithologic unit boundaries are provided for each coring station in the supplementary information.

315 The wavy and lenticular nature of laminations in much of LU5 is interpreted as evidence for traction processes under the influence of bottom water currents. We interpret this unit as a meltwater generated laminated facies that transitions from a grounding line proximal deposit at the base to distal at the top. The Ca enrichment arises from increased inputs of detrital carbonate, eroded from the surrounding Cambrian and Silurian bedrock and is more common in the grounding zone proximal base of the sequence. Evidence for truncation of laminations indicates somewhat discontinuous deposition, with more

320 frequent episodes of erosion near the base of LU5, when deposition occurred closer to the grounding zone. The thinner nature of LU5 at on the shallow inner sill (8-PC, 9-PC), and lack of relatively coarse, Ca- enriched laminae like those found near the base of LU5 in 6-GC and 7-PC, indicates a later onset for continuous sedimentation. On the outer sill, the more

condensed LU5 in 10-GC also indicates a more dynamic depositional environment, likely influenced by current activity or ice scouring.

325

LU4: Faint, planar mm-scale laminations that lack notable evidence for bioturbation define LU4. The transition between LU5 and LU4 is gradual, with the base of LU4 being identified by a notable and correlative decline in Ca-abundance that is identified at all the coring stations (Fig. 5). Laminations in LU4 are only weakly visible by the naked eye but are evident in the CT scanning data (Fig. 7). In 7-PC and 6-GC, laminations appear to transition from wavy to lenticular with frequent faulting in LU5, to planar in LU4. With the exception of 10-GC on the outer sill, coarse ice-rafted clasts remain dispersed in low numbers through LU4. LU4 is a silty clay, containing very minor amounts of sand (<1%), with some samples showing a slight elevation of medium to coarse silt (Fig. 8). The bulk density remains relatively constant (1.6-1.8 g/cm³) with mass specific magnetic susceptibility higher than in LU5, and increasing up-core towards the top of the unit, consistent with a continued up-core decrease in detrital carbonate concentrations. Occasional minor pulses of Ca are seen throughout the unit. The LU5 to LU4 transition appears correlative across the fjord, and many small-scale features in the Ca, bulk density and magnetic susceptibility logs can be traced between the sites (Fig. 5).

330

335

We interpret LU4 as laminated meltwater influenced sediments dominated by suspension settling in an ice distal environment. This is consistent with the gradual transition seen between LU5 and LU4. The apparent correlative nature of LU4 across the fjord suggests uniform and continuous deposition.

340

LU3: Throughout Sherard Osborn Fjord, LU4 is capped by LU3, a second diamicton containing large gravel to cobble-sized clasts (Figs. 6, 7). LU3 is recognized by its high bulk density (1.8-2.0 g/cm³), elevated Ca-content and abundant clast content (10 - >100 clasts/cm³) (Fig. 5). LU3 has a low mass specific magnetic susceptibility, except at 8-PC where cobble and bolder sized clasts of felsic rocks were recovered, which presumably led to the high susceptibility measurements. CT images from 6-GC and 7-PC show that the lower boundary is heavily bioturbated, while at 8-PC and 9-PC it is marked by a higher concentration of coarse clasts that tend to fine upwards (Fig. 7). At 8-PC the coarse clasts appear to have been smeared down the inside of the core liner, artificially increasing the clast content of the underlying LU4. In all cores, the lower boundary of LU3 and uppermost 5-10 cm of LU4 are bioturbated. The upper boundary of LU3 is relatively sharp at all sites. The matrix material of LU3 is clayey-silt, with a generally coarser mean grain size compared to LU4, and a larger sand-sized contribution (up to 4.5%) (Fig. 8). The only indication of deformation is seen in 8-PC, surrounding the region where large dropstones were recovered. The absence of deformation at the other sites suggests that this is an artifact of coring and not a primary depositional feature. We interpret LU3 as a glaciomarine diamicton, primarily composed of ice rafted material deposited by icebergs and sea ice.


345

350


355

360 *LU2*: A sharp but bioturbated boundary separates LU3 from LU2 at all sites except 10-GC on the outer sill. The defining characteristic of LU2 is the presence of bioturbation and lack of preserved laminations. This unit exhibits very little variation in the bulk density or Ca content (Fig. 5), is composed primarily of fine silt and clay, and contains very few scattered ice-rafted clasts (Fig. 7). The thickness of LU2 varies considerably across the fjord, with the thickest occurrences in 6-GC and 7-PC. LU2 is nearly absent in sediments from the outer sill (10-GC) where there is an abrupt and non-bioturbated contact between LU3 and LU2 (Fig. 5).


365 *LU1*: A return to a laminated facies lacking coarse ice-rafted clasts and bioturbation defines LU1. The grain size spectra of LU1 closely resemble those from LU4 (Fig. 8). The laminations in LU1 are also highly fractured and distorted along the sides of the core. This is attributed to coring disturbance and the soft to soupy nature of these near surface sediments. LU1 is divided into two subunits. LU1b is only found at stations 8 and 9 on the inner sill. It is a faintly laminated unit, with some discrete intervals where the laminations are disturbed, sometimes by bioturbation. LU1b has low abundances of Ca, similar to what is found in LU2 and LU4 (Fig. 5). LU1a is marked by the emergence of more pronounced mm- to cm-scale laminations that are only seen clearly in the CT images (Fig. 7). LU1a is marked by increased Ca concentrations, which are most apparent in 9-PC and 7-PC (Fig. 4). At stations 10, 6 and 7, LU1a sits abruptly on top of LU2 (Fig. 5), while at stations 370 8 and 9, there is a gradual transition between LU1b and LU1a.



375 **Figure 6.** CT-images and associated lithologic units for each core. White in CT-images corresponds to higher density zones, black to lower density zones. Yellow triangles represent levels of radiocarbon dates.



380 Figure 7. Examples of CT-images and color photographs representative of the six lithologic units. The vertical scale for each image is a constant 50 cm. The structures seen on the CT-images and the color photographs are not identical because the imaging planes are slightly different. The CT-images are representative 2D slices of the core interior, whereas the photos show the surface of split cores. White in CT-images corresponds to higher density zones, black to lower density zones. Composite depths for each interval are given in red.



385 Figure 8. Grain size spectra of lithologic units 1 – 5 illustrating the similar grain size distributions of sediments in LU1 and LU4, as well as the difference between the dark (less dense) and light (more dense) laminations in LU5.

4.1 Ages of unit boundaries

Radiocarbon dating is used to constrain the ages of lithostratigraphic unit and subunit boundaries. In most cases samples were collected from within a few centimeters of these boundaries in one or more cores (Fig. 5). The youngest age underlying a unit or subunit boundary is used to provide a maximum age for the boundary (Fig. 9). However, this is not possible for the LU5/LU6 and LU1b/LU2 boundaries. For LU6 the oldest age from the overlying unit is adopted as the minimum boundary age. A younger than and older than age is provided for LU1b, based upon dates obtained from two different cores (Table 4). Of the 48 radiocarbon dates, 7 were identified as outliers because the overlying ages are older and fell outside the 1σ calibrated age range. The lowermost sample in 8-PC (Sample #35, Table 3) is also considered an outlier, due to the substantially younger age returned in a sample 22.5 cm up-core (Sample #34, Table 3), and the fact that it was obtained on a mixed benthic assemblage. On the other hand, LU5 sediments from this site were not deposited continuously, implying

390

395

periods of erosion or non-deposition that could account for the large difference in ages between the lowermost dates in 8-PC (Fig. 9). In any case, due to the general uncertainty about the accuracy of this date, we have decided to treat it as an outlier.

The lithologic sequence from Sherard Osborn Fjord sediments spans nearly the entire Holocene (Table 4), with the oldest boundary (LU5/LU6) dated in 7-PC (Sample #24) to 10200 ± 40 ^{14}C years (10330 - 11160 cal. a BP) while the youngest boundary (LU1a/2) is dated in 6-GC (Sample #1) to 1730 ± 40 ^{14}C years (540 - 1150 cal. a BP). The oldest age comes from 11-GC (Sample #48, 11150 ± 250 ^{14}C years) and provides a constraint on the timing for ice retreat from the outer sill.




Figure 9. Likelihood distributions of calibrated radiocarbon dates illustrating the mean and $1\text{-}\sigma$ age ranges. Color-coding is used to identify the lithologic unit that each date was acquired from. Dates from each core are presented in their stratigraphic order. White indicates ages that are deemed to be outliers. Yellow stars indicate dates used to define the ages of unit boundaries (Table

4). On the inner sill, the age range for the LU2–LU1b transition is bracketed by an ‘older than’ and ‘less than’ age from 9-PC and 8-PC respectively. Index numbers correspond to those in Tables 3 and 4. Date #47 from 11-GC is not shown as it could not be definitively placed within a lithologic unit. *Np* indicates date obtained on *N. pachyderma*.

410

Table 4. Key radiocarbon dates and calibrated ranges defining basal ages of lithologic (sub)unit boundaries. Calibrated ages and uncertainty are rounded to 100 years, with the mean age from the calibrated age range reported. The first column refers to the radiocarbon date index number from Table 3.

| # | Sample | Unit | ¹⁴ C Age (a) | Age (cal ka BP) |
|----|----------------------|----------------|-------------------------|-----------------|
| 1 | 06-GC-2, 3-6 cm | 1a | 1730±75 | 0.9 ± 0.3 |
| 36 | 9-TWC-CC | 1b | >3550±55 | 2.9 ± 0.4 |
| 26 | 8-GC-3, 102-104 | 1b | <4090±20 | 3.6 ± 0.4 |
| 37 | 9-PC-4, 103-105 cm | 2 | 4380±25 | 3.9 ± 0.4 |
| 4 | 06-GC-2, 93-95 cm, B | 3 | 6400±25 | 6.3 ± 0.3 |
| 44 | 10-GC-2, 132-134 cm | 4 | 7880±45 | 7.9 ± 0.3 |
| 24 | 7-PC-6, 76-78 cm, B | 5 (mid fjord) | 10200±40 | 10.7 ± 0.4 |
| 34 | 8-PC-5, 46-49 cm | 5 (inner sill) | 9140±40 | 9.3 ± 0.4 |

5. Discussion

415 5.1 Overview of the Holocene lithostratigraphic succession


The succession of stratigraphic units recovered in Sherard Osborn Fjord mirrors the classic deglacial facies transitions described from ice-shelf settings of Antarctica (Smith et al., 2019) and high-latitude Northern Hemisphere fjords occupied by marine terminating outlet glaciers and floating ice tongues (O’Cofaigh et al., 2001; Reilly et al., 2019) (Fig. 10). The interpretations of the stratigraphic units are here used to reconstruct the Ryder Glacier’s dynamical history (Fig. 11). The Sherard Osborn Fjord sequence involves a basal subglacial to glaciomarine diamicton (LU6) overlain by a laminated meltwater dominated facies. The laminated facies fines upwards from a grounding zone proximal deposit with evidence for traction current activity (LU5) to a grounding zone distal facies deposited by suspension settling (LU4) beneath a floating ice tongue (Fig. 9).

420

425 The laminated grounding zone distal facies (LU4) is abruptly terminated by a clast-rich diamicton (LU3). Similar facies transitions have been described as a response to migration of the ice shelf calving front, glacier surging or advance, or ice shelf collapse (Smith et al., 2019). In Sherard Osborn Fjord, the overlying sediments (LU2) are clast poor with extensive bioturbation suggesting deposition under more productive surface waters indicative of less extensive surface ice cover. As such, the facies succession, with LU3 separating the grounding zone distal (LU4) and more open-water bioturbated sediments (LU2), is consistent with a collapse of Ryder’s ice tongue. The removal of an ice tongue marking the onset of LU3

430

435 allowed enhanced input of poorly sorted ice rafted material as debris-laden icebergs calved directly from the grounding zone were able to traverse the fjord. A return to a laminated facies (LU1) similar in character to LU4 indicates the successive re-establishment of an ice tongue that eventually extended to the outer sill in Sherard Osborn Fjord. This basic interpretation is supported by historical observations of Lauge Koch, who mapped the ice-tongue limit in proximity to the outer sill in 1917 (Koch, 1928) (Fig. 2).



440 **Figure 10.** Representative lithostratigraphic column and environmental interpretations for the sedimentary cover of the inner sill and middle fjord. Approximate linear sedimentation rates are provided for the middle fjord (station 7, Fig. 2) using the derived unit boundary ages and their uncertainty (Table 3). The interpreted lithostratigraphic column is shown alongside the Agassiz air temperature reconstruction and elevation changes of the GrIS at Camp Century (Lecavalier et al., 2017); the reconstructed extent of Petermann Glacier ice tongue (Reilly et al., 2019); the period of beach ridge formation along the north coast of Greenland

indicating the absence of perennial land fast ice (Möller et al., 2010); and driftwood delivery to northern Ellesmere Island (England et al., 2008). Radiocarbon dates from England et al. (2008) were recalibrated using Intcal20.

445 5.2 Early Holocene glacier and ice tongue dynamics


The oldest recovered sediments come from 11-GC on the outer sill where a sample of the benthic foraminifera *Cassidulina neoteretis* returned a date of 12 ± 0.5 cal ka BP. Due to the limited amount of sediment that has accumulated above the bedrock/till at 6-GC, 10-GC and 11-GC (Fig. 4), we interpret this as a minimum age for the retreat of Ryder Glacier from the outer sill, where it was grounded 80-90 km seaward from its current position (Fig. 11). Deposition of the grounding zone proximal meltwater sediments (LU5) above a likely subglacial diamicton in 7-PC commenced at 10.7 ± 0.4 cal ka BP, indicating that the glacier had retreated further inland by this time (Fig. 10). The calibrated ranges of radiocarbon dates from the base of 7-PC (10.3-11.1 cal ka BP) conform to the reported age for the Kap Fulford Stade (>10.5 cal ka BP) (Kelly and Bennike, 1992) (Fig. 11). It is less certain if the older age from 11-GC on the outer sill ($11.6-12.6$ cal ka BP) (Fig. 9, Table 3) implies that grounded ice was here during the Kap Fulford Stade or an earlier stage of deglaciation.

455 Following retreat of Kap Fulford glacial ice, the sedimentary sequence is dominated by the Ca-rich laminated meltwater facies (LU5). The coarser grained Ca-enriched laminae are associated with increased meltwater input from glacial erosion of the surrounding Cambrian and Silurian carbonate shelf bedrock (Fig. 2). The strong meltwater signal captured in the LU5 coincides with the rapid Early Holocene reduction in the height of the GrIS (Lecavalier et al., 2017) (Fig. 10).

460 Sediments from LU5 are comparatively thin at the inner sill coring sites, and difficult to directly correlate with the records from 6-GC and 7-PC. Erosional events identified in LU5 sediments from the inner sill suggest discontinuous sedimentation (see 9PC-6 Fig. 7). Sitting in much shallower water depths (228-271 mbsl; Table 1), glacial ice likely remained grounded here after it retreated from stations 6 and 7. Deposition on the inner sill would also have been influenced by bottom scouring currents and debris flows while the glacier was grounded (or near flotation) on shallow regions of the sill that rise to depths of 193 m. Therefore, we suggest that Ryder Glacier remained grounded on or near the inner sill until sometime after 9.3 ± 0.4 cal ka BP, which marks the onset of continuous sedimentation of LU5 at this location (Fig. 11; Table 4). This is consistent with the age of the Warming Land Stade, bracketed between >9.5 cal ka BP and 8.0 cal ka BP by Kelly and Bennike (1992). The shallow inner sill, first identified through bathymetric mapping during the *Ryder 2019* expedition (Jakobsson et al., 2020), is very close to the previously inferred grounding zone location for the Warming Land Stade and provides a natural pinning point for the marine-based glacier during this standstill (Fig. 2).

475 In 6-GC and 7-PC there is a broad period of elevated detrital carbonate delivery (high Ca-content) that occurs near the top of LU5 (Fig 5). This likely reflects enhanced meltwater delivery during terrestrial ice retreat at the end of the Warming Land Stade prior to the LU4/LU5 boundary at 7.6 ± 0.40 cal ka BP (Fig. 11; Table 4). Across Sherard Osborn Fjord, deposition of

the laminated LU4 sediments continued until 6.3 ± 0.3 cal ka BP (Fig. 11; Table 4), which is very similar to the age (6.0 cal ka BP) for the minimum Holocene extent reported for Ryder Glacier (Kelly and Bennike, 1992).



480 **Figure 11.** Generalised lithology and corresponding reconstructions of Ryder Glacier and ice tongue during deposition of the lithologic units. The profile location is illustrated in Fig. 11. The bathymetry was re-gridded from BedMachine v3 (Morlighem et al., 2017) after adding bathymetric data collected on *Ryder 2019*. Substantial modifications to the former digital elevation model exist around the now mapped inner sill and beneath the modern ice tongue (see supplementary material for details). Yellow dots indicate approximate positions of coring stations discussed in the text.

485 5.3 Middle Holocene inland retreat and collapse of Ryder's ice tongue

The second diamict facies (LU3) in the stratigraphic succession separates the faintly laminated sub-ice tongue sediments of LU4 and the bioturbated facies of LU2. As this unit is found from the inner sill all the way to the outer sill, and does not appear to be time-transgressive, it cannot be attributed to a slow landward migration of the ice tongue calving margin. Instead, it is interpreted as a collapse facies (Smith et al., 2019) associated with the sudden disintegration of Ryder's ice
490 tongue. While this interpretation fits with the overall facies succession, the ~ 2.4 kyr duration of LU3 (from 6.3 ± 0.3 to 3.9 ± 0.4 cal ka BP; Table 4) is too long to be a simple abrupt collapse event, as these are generally associated with increased sediment accumulation rates (Smith et al., 2019).

An alternate interpretation is that LU3 sediments represent an erosional event. However, this does not fit with the
495 overlapping ages of sediments from the top of LU4 at coring stations 6, 7, 9 and 10, and to a large degree the ascending ages found downcore through LU3 at station 8 (Fig. 9). Therefore, the widespread distribution, long duration, consistent boundary ages of LU3, and lack of evidence for pervasive ice scouring makes it hard to reconcile this facies with a glacial surge. This unit is also much younger than the inferred age for the drainage of the ice dammed lake that developed on Wulff Land between Ryder and C. H. Ostenfeld glaciers between 7.9-8.5 cal ka BP (Kelly and Bennike, 1992). These observations
500 suggest that LU3 marks a period of low sedimentation that began around 6.3 ± 0.3 cal ka BP when terrestrial studies suggest that Ryder Glacier had retreated further inland than its current position (Kelly and Bennike, 1992), similar to much of the ice margin along northwestern Greenland during the Middle and Late Holocene (Søndergaard et al., (2020) The age for the end of LU3 (3.9 ± 0.4 cal a BP, Table 4), marking the re-growth of a marine based glacier and ice tongue, is also consistent with existing dates constraining the onset of the Steensby Stade ($<4560 \pm 410$, Station 34, Table 1).

505 The explanation that best fits evidence from terrestrial field studies, and the overall facies succession, is that the condensed diamict of LU3 was deposited when Ryder Glacier retreated far enough inland to become cut-off from the main fjord. In Sherard Osborn Fjord, a relatively deep, isolated marine embayment exists behind a prominent topographic high lying 40 km inland of the modern grounding zone (Fig. 12). Here elevations increase to between 100-200 m above sea level, with
510 elevations surrounding the embayment reaching 400 – 600 m above sea level (Figs. 11, 12). Retreat of the grounding zone onto this topographic high (possibly a former grounding zone wedge), or even inland of it, would explain the apparent rapid disintegration of the ice shelf at the end of LU4. The entrainment of meltwater-derived sediments within the isolated embayment or even behind the inner sill, can account for the overall reduction in sedimentation rates in the outer fjord. This is consistent with observations from modern proglacial lakes that illustrate their efficiency at disrupting meltwater fluxes and
515 sequestering sediments (Carrivick and Tweed, 2013; Bogen et al., 2015; Piret et al., 2021). While grounded in this distal position, calving could have discharged some icebergs into Sherard Osborn Fjord, delivering coarse-grained ice rafted sediments. Elevated Ca contents within LU3 (Fig. 5) attest to sediment provenance being primarily from landward of the


inner sill (Fig. 2). Furthermore, the sedimentation rates during LU3 are low (9-10 cm/kyr; Fig. 10) and comparable with sedimentation rates of 10-30 cm/kyr reported by Dowdeswell et al. (1994) for massive diamicts derived from iceberg rafting in Scoresby Sund and the adjacent East Greenland shelf. A final indication that Ryder Glacier retreated to the innermost fjord or beyond is the lack of iceberg scouring seen in bathymetric data on the inner sill. This suggests that the calving front was grounded in water shallow enough to not produce icebergs in excess of 190-200 m thickness – a condition that can easily be met by the landward retreat of Ryder Glacier (Fig. 11).

525 A modern analogue for this configuration is seen in neighboring Victoria Fjord. Today C. H. Ostenfeld Glacier is grounded on a topographic high at the end of the fjord, in an equivalent position to the landward terminus of Sherard Osborn Fjord where the proposed Middle Holocene ice margin is located (Fig. 11). Between 2000 and 2006, the C. H. Ostenfeld ice tongue, which protruded from a narrow over-deepened marine channel, largely disintegrated (Moon and Joughin, 2008), while the grounding zone may soon become increasingly land-based and discharge grounded ice directly into Victoria Fjord (Hill et al., 2017) (Fig. 11).

The continued retreat of Ryder Glacier through the Early and Middle Holocene is consistent with many regional paleoclimate proxies from around northern Greenland. This includes beach ridge formation along the north Greenland coast, indicating more open water conditions that persisted until ~6 cal ka BP (Möller et al., 2010; Funder et al., 2011), reduced winter sea-ice conditions in Bliss Lake until 6.5 cal ka BP (Olsen et al., 2012), and peak late summer air temperatures inferred from $\delta^{18}\text{O}$ of chironomids in Secret and Deltasø lakes that were $>2^\circ\text{C}$ warmer than present until 6.2 cal ka BP (Axford et al., 2019; Lasher et al., 2017) (Fig. 1). The paleotemperature estimates from these lakes indicate a slight cooling, but persistent, stable and positive temperatures between ~6.0-4.0 cal ka BP (Lasher et al., 2017; McFarlin et al., 2018). Diminished sediment delivery due to the presumably stable inland position of Ryder Glacier lasted from 6.3 ± 0.3 to 3.9 ± 0.4 cal ka BP (Table 4). The inferred re-advance into Sherard Osborn Fjord, marked by increased sediment delivery at the onset of LU2, occurred around 3.9 ± 0.4 cal ka BP. This timing for glacier advance is consistent with cooling seen in lake based temperature reconstructions around 4 cal ka BP (Lasher et al., 2017) and the oldest estimated age (3.5 to 4.0 cal ka BP) for ice at the base of the southern dome of Hans Tausen ice cap, which had disappeared during the Middle Holocene - although northern parts of the ice cap had survived (Madsen and Thorsteinsson, 2001; Landvik et al., 2001; Zekollari et al., 2017). However, it should be emphasized that the onset of deposition of LU2 at 3.9 ± 0.4 cal ka BP, marks the advance of Ryder Glacier into Sherard Osborn Fjord, and terrestrial-based ice may have started advancing earlier.

A stable inland position for Ryder Glacier during LU3 fits with generally colder conditions that prevailed over north Greenland towards the end of the middle Holocene. This is reflected in the development of more prolonged winter ice cover in Bliss Lake after 5.9 cal ka BP (Olsen et al., 2012) and a period of low melt rates for the north GrIS inferred from elevation changes at Camp Century (Lecavalier et al., 2017) (Fig. 10). Therefore the prolonged slow deposition of coarse IRD during

LU3 is likely a combined effect of limited but persistent iceberg calving, the slow melt-out of englacial material from the disintegrated ice tongue, as well as continued input of sea-ice rafted material entrained during periods of shore-fast sea ice growth and decay.



555

Figure 12. A. Bathymetry and sub-glacial topography of north Greenland from Morlighem et al. (2017) illustrating pronounced differences between the broad submarine trough that extends inland from Petermann Glacier and the more restricted, and rapidly shoaling topography in Sherard Osborn and Victoria Fjords. B. Detail of Sherard Osborn Fjord showing the two mapped sills that correspond to the likely grounded limits of Ryder Glacier during the Kap Fulford and Warming Land Stade. At the end of the Middle Holocene, Ryder Glacier likely retreated to a more inland position becoming largely land-based and cut off from the main fjord by the restricted embayment. White dots are the coring sites discussed in the text. Red line (Y-Y') shows the position of the bathymetric profile used in Fig. 11 (additional detail in the supplementary information). Ice margin positions during the Kap Fullford Stade (yellow line) and Warming Land Stade (blue line) are from Kelly and Bennike (1992). GZ = Grounding Zone.

560

5.4 Late Holocene cooling and the re-growth of Ryder's ice tongue

565

The advance of Ryder Glacier and the re-growth of local ice caps surrounding Sherard Osborn Fjord is described by the Steensby Stade re-advance. Its onset is not well constrained by terrestrial field data, but is believed to have started sometime

after 5.1-4.7 ka (Kelly and Bennike, 1992), with outlet glaciers reaching their maximum extents by the start of the 1900's. Thus, in marine records from Sherard Osborn Fjord, the Steensby Stade encompasses deposition of LUs 2 and 1.

570 The highly bioturbated LU2 sediments appear abruptly on top of LU3, with the onset of deposition sometime after 3.9 ± 0.4
cal ka BP (Table 4). While the inland glacier may have started advancing earlier, increased sedimentation rates at the onset
of LU2 (Fig. 10) likely marks the seaward migration of Ryder Glacier to a marine-based position in the innermost fjord.
Within this setting, the re-growth of a limited ice tongue would act as a filter for coarse iceberg rafted sediments and can
explain the paucity of IRD in LU2 sediments. Extensive bioturbation in LU2 attests to warmer climatic conditions and the
575 absence of an ice tongue seaward of the mapped inner sill and likely less persistent sea-ice cover.

On the inner sill, deposition of laminated sediments lacking bioturbation (LU1b) marks the re-growth of the ice tongue out to
this location (Fig. 11) between 3.6 ± 0.4 and 2.9 ± 0.4 cal ka BP. This is roughly 300-1000 years after the glacier became
marine-based at the end of LU3. The persistence of bioturbation (LU2) at stations in the middle fjord (stations 6 and 7),
580 indicates an earlier establishment of the ice tongue over the inner sill, while more open water conditions persisted further
seaward (Fig. 9). This configuration is similar to what we found during our expedition in 2019 (Fig. 2). It appears that this
was a relatively stable configuration that persisted for another 2-2.7 kyrs (until 0.9 ± 0.3 cal ka BP; Table 4), before the ice
shelf extended towards the outer sill and laminated LU1a sediments were deposited in the middle fjord (Figs. 10, 11). The
long-term stability of the ice tongue terminus near the inner sill is consistent with the inner sill acting as a barrier for Atlantic
585 water invasion and limiting basal ice tongue melting (Jakobsson et al., 2020).

The Late Holocene re-growth of Ryder's ice tongue has some parallels with the Petermann ice tongue, which began to
reform between 1.9-2.3 cal ka BP, after being absent since its collapse around 6.9 cal ka BP (Reilly et al., 2019). Petermann
also attained a stable ice tongue with an extent similar to 20th century historical observations between 0.4-0.9 cal ka BP
590 (Reilly et al., 2019). At both Ryder and Petermann the growth of ice tongues towards the outer fjords occurred much later
than the establishment of multi-year landfast sea ice in front of Phillips Inlet and Disraeli Fjord on northern Ellesmere Island
around 5.5 cal ka BP (England et al., 2008). On the other hand, ice tongue re-growth in Petermann (1.9-2.3 cal ka BP) and
growth of Ryder's ice tongue to the inner sill in Sherard Osborn Fjord (by 2.9 ± 0.4 cal ka BP) are consistent with the
development of more extensive sea ice around the northern Greenland margin by 2.5 cal ka BP (Funder et al., 2011) and only
595 intermittent periods of sea-ice free conditions in the Lincoln Sea after 3.9 cal ka BP based on the cessation of driftwood
delivery to Clements Markham Inlet (England et al., 2008) (Fig. 10).

Uncertainty in the local reservoir correction leaves the timing for the maximum Late Holocene growth of Ryder's ice tongue
poorly constrained. However, as with the final growth of Petermann's ice tongue (Reilly et al., 2019), it appears consistent
600 with numerous regional paleoclimate records that document the development of cooler conditions close to the transition from

the Medieval Warm Period to the Little Ice Age. For example, geochemical proxies from Bliss Lake (Fig. 1) indicate the onset of cooling around 850 cal. a BP (Olsen et al., 2012), which agrees with the onset of cooler air temperatures inferred from the $\delta^{18}\text{O}$ record of the Hans Tausen ice cap around 780 cal. a BP (Olsen et al., 2012; Hammer et al., 2001).

605 The absence of a thin bioturbated facies reflecting deposition since the 20th century retreat of Ryder's ice tongue is likely due to incomplete recovery of surface sediments in the studied cores. Given the predicted range of sedimentation rates for LU1a in 7PC (51-102 cm/kyr, Fig. 10), only 4–8 cm of sediment would have accumulated since the retreat of Ryder's ice tongue to its current position ~80 years ago. This thin sedimentary cover could easily have been missed during coring, especially considering the soft nature of surface sediments encountered in the fjord.

610 **5.5 Controls on the Holocene dynamics of Ryder Glacier**

The retreat of Ryder Glacier from the fjord mouth ($>10.7\pm 0.4$ cal ka BP) is consistent with the demise of glacial ice across much of northern Greenland in the Early Holocene (Bennike and Björck, 2002; Larsen et al., 2010; Strunk et al., 2018; Larsen et al., 2018). During the early stages of deglaciation, the Lincoln Sea was covered by thick shelf ice, fed by coalescent glaciers emanating from Greenland and Ellesmere Island (England et al., 1999; Larsen et al., 2010). This eastward

615 flowing shelf ice overrode parts of the north Greenland coast (Johannes V. Jensen Land; Fig. 1) with its final break up occurring between 10.3 - 10.1 cal ka BP (Larsen et al., 2010). Reduced sea-ice conditions in the Lincoln Sea at this time are evidenced by the onset and persistence of driftwood delivery to Clements Markham Inlet (Fig. 9; England et al., 2008) after glacial ice had retreated from the northernmost Nares Strait >10.1 ka (England et al., 1999). Future work on sediment cores from *Ryder 2019* utilizing paleoenvironmental proxies for sea ice and oceanographic conditions may help resolve how
620 atmospheric warming (Cook et al., 2019) and warm Atlantic water advection (Wood et al., 2021) into the Lincoln Sea region influenced Holocene sea-ice dynamics and glacier retreat.

Ryder Glacier retreated much earlier than Petermann Glacier to the south (Jakobsson et al., 2018; Reilly et al., 2019). Petermann Glacier remained grounded at the mouth of Petermann Fjord, bordering Hall Basin, until 7.5 cal ka BP
625 (Jakobsson et al., 2018) even after Ryder had retreated from the inner sill and was >40 km from the mouth of Sherard Osborn Fjord. In this respect, it is important to acknowledge that Petermann's retreat from the outer fjord was one of the final events that occurred during deglaciation of Nares Strait and specifically of Hall Basin (Jakobsson et al., 2018). Glacial ice in Hall Basin had remained in an advanced position abutting Robeson Channel until 9.3 cal ka BP (Fig., 1; Jakobsson et al. 2018; England et al 1999). The ice margin had receded to the mouth of Petermann Fjord by 8.7 cal ka BP, prior to the
630 opening of Nares Strait between 8 and 8.5 cal ka BP (Jennings et al., 2011, 2019; Georgiadis et al., 2018). Therefore, differences in the timing of retreat between Petermann and Ryder Glaciers appears closely tied to local glacial and sea-ice conditions. These appear to have been less severe and more mobile in the Lincoln Sea in the Early Holocene compared to southern Nares Strait region.

635 The sedimentary facies from Sherard Osborn Fjord documents seemingly gradual retreat and regrowth phases for Ryder
Glacier as evidenced by the generally low abundance of coarse IRD (Fig. 10) and the absence of lenses, or pulses of IRD that
would otherwise indicate ice tongue collapse events. For example, in Petermann Fjord the collapse of the ice shelf at 6.9 cal
ka BP is marked in sedimentary records by the abrupt appearance of IRD clasts in sediments from across the fjord (Reilly et
al., 2019). The stability of Ryder Glacier's ice tongue is most likely related to the physiography of Sherard Osborn Fjord.

640

The inner sill in Sherard Osborn Fjord not only acts as a natural pinning point for glacial ice during its retreat but also
impedes the flow of warm Atlantic subsurface waters into the innermost fjord (Fig. 3; Jakobsson et al., 2020). This likely
reduces the subaqueous melting at the grounding zone and of the ice tongue today, and may have done so efficiently for
much of the Holocene. The subglacial bed topography and physiography of the inner fjord will also exert an influence on
645 glacier (and hence ice tongue) stability. There appears to be a slight retrograde slope between the inner sill and modern
grounding zone (Fig. 11), which are common in northern Greenland fjords and can be conducive to rapid glacial retreat
(Enderlin et al., 2013; Carr et al., 2015; Hill et al., 2018). However, Sherard Osborn Fjord also narrows considerably
landward of the inner sill and remains flanked by steep-sided cliffs (Fig. 12). Jamieson et al. (2012) have shown that even on
a retrograde slope, retreating ice streams slow or can become pinned where they pass narrow sections in surrounding
650 topography. The sedimentary record from seaward of the inner sill contains no evidence for massive calving events that
would otherwise suggest a collapse of the ice tongue during a rapid retreat down a retrograde slope. However, there is
tremendous uncertainty regarding the bathymetry beneath the current ice tongue as no direct measurements exist, so even the
presence of a retrograde slope is not certain. A more certain feature of the bathymetry is the steeply rising bed topography
landward of the modern grounding zone (Figs. 11, 12). This feature, which could be a former grounding zone, may have
655 promoted the gradual retreat of Ryder Glacier and the stability of its ice tongue during the Early and Middle Holocene
(Powell, 1990; Alley, 1991; Hill et al., 2018). A critical observation is that despite the apparent persistence of Ryder's ice
tongue, the physiographic setting of Sherard Osborn Fjord, which appears to be conducive to glacier stability, did not halt the
overall retreat of Ryder Glacier under the relatively mild changes in climate forcing during the Holocene.

660 From the results of this study, it is not possible to constrain the amount of inland retreat Ryder Glacier experienced during
the Middle Holocene. Based on the cessation of sedimentation in the outer fjord, we suggest that it may have retreated at
least into the restricted embayment at the end of Sherard Osborn Fjord (Figs. 11, 12). Unlike Petermann Glacier, where the
bed topography widens and extends far beneath the GrIS at elevations below modern sea level, the bed of Ryder Glacier
abruptly rises above sea level some 60 km landward of the modern grounding line (Fig. 12). This is comparable to the
665 smaller glaciers in the immediate surroundings and to the Køge Bugt glacier complex in southeast Greenland, where the
steep land-ocean boundary limited the Middle Holocene retreat of the glaciers in this area (Dyke *et al.*, 2017). This suggests
that the land-ocean interface might have been an important boundary that mediated or halted the Holocene retreat of many

glaciers that are not located in deep troughs that extend beneath the inner ice sheet. Overall, our constrains on the Holocene dynamics of Ryder Glacier suggest that it could retreat another 40-60 km inland if climatic conditions remain similar or exceed those of the Middle Holocene.

6. Conclusions

Like much of northern Greenland, Ryder Glacier responded acutely to climate variability in the Holocene. During the Early and Middle Holocene it retreated over 120 kilometers from a grounded position near the mouth of Sherard Osborn Fjord (80 km seaward of the modern grounding zone) to likely become land-based more than 40-60 km landward of its current position by 6.3 ± 0.3 cal ka BP. Throughout this long period of retreat, deposition of laminated, clast poor sediments attest to strong meltwater inputs and an overall stable ice tongue. Ryder Glacier remained land-based until the Late Holocene (3.9 ± 0.4 cal ka BP). As it again advanced into Sherard Osborn Fjord, an ice tongue developed that quickly grew out to its 21st century position near a prominent bathymetric sill, located 30 km seaward of the modern grounding zone. Today this sill reduces the incursion of Atlantic waters into the inner fjord and shields the grounding zone and ice tongue from basal melting (Jakobsson et al., 2020). This was also likely the case for 2.0-2.7 kyrs in the Late Holocene, before the ice tongue grew to its historical maximum extent around 0.9 ± 0.3 cal ka BP. Late Holocene bioturbated sediments that are nearly devoid of ice rafted debris were deposited between 3.9 ± 0.4 and 0.9 ± 0.3 cal ka BP and indicate reduced ice cover over much of the middle and outer fjord. This contrasts with the near permanent shorefast ice conditions that existed in Sherard Osborn Fjord for much of the late 20th century (Higgins, 1989), but it is similar to conditions during the summer of 2019. The physiography of Sherard Osborn Fjord appears to have had a stabilizing effect on the ice tongue of Ryder Glacier through the Holocene. Nevertheless, Ryder Glacier still retreated >40-60 km landward of its current position and remained there under the relatively mild climate forcing in the Middle Holocene, suggesting that it may again retreat completely from the fjord to an inland position if climate warming continues at its current pace.

Data availability

Upon publication, the data presented in this manuscript will be archived in the Bolin Centre Database at the Bert Bolin Centre for Climate Research (<https://bolin.su.se/data/>).

Author contributions

This manuscript was written by MO with subsequent input from all co-authors. MJ and LM led the Ryder 2019 Expedition. Cores were collected and processed shipboard by MO, TC, BR, GW, and LG. Shore-based analyses were conducted by MO,

695 BR, FV, CP, GW, AKO, OLM, AG, MM and HD. CS and JN collected, processed and interpreted the oceanographic data. MM provided the improved DEM for Sherard Osborn Fjord.

Competing interests

The authors declare that they have no conflict of interest.

700 Acknowledgments

We are grateful to Captain and crew of Icebreaker *Oden* and personnel from the Swedish Polar Research Secretariat for their support during the planning and execution of the Ryder 2019 Expedition. This expedition was endorsed as Explorer's Club Flag Expedition #51. We thank the Swedish Polar Research Secretariat, Center for Coastal and Ocean Mapping, University of New Hampshire and Stockholm University for supporting the *Ryder 2019* Expedition financially. M.O., M.J, C.S and
705 colleagues from Stockholm University were supported by grants from the Swedish Research Council (VR; grants 2016-05092, 2016-04021, 2018-04350). T.C. and L.G. were supported by the USGS Land Change Program. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government. We thank June Padman (Oregon State University) and Carina Johansson (Stockholm University) for their help in shipboard core processing.

710 References

- Alley, R. B.: Sedimentary processes may cause fluctuations of tidewater glaciers, *Annals of Glaciology*, 15, 119–124, 1991.
- Axford, Y., Lasher, G. E., Kelly, M. A., Osterberg, E. C., Landis, J., Schellinger, G. C., Pfeiffer, A., Thompson, E. and Francis, D. R.: Holocene temperature history of northwest Greenland – With new ice cap constraints and chironomid assemblages from Deltasø, *Quaternary Science Reviews*, 215, 160–172, doi:10.1016/j.quascirev.2019.05.011, 2019.
- 715 Bamber, J. L., Oppenheimer, M., Kopp, R. E., Aspinall, W. P. & Cooke, R. M.: Ice sheet contributions to future sea-level rise from structured expert judgment, *Proceedings of the National Academy of Sciences* 116, 11195, doi: 10.1073/pnas.1817205116, 2019
- Bennike, O. and Kelly, M.: Radiocarbon dating of samples collected during the 1984 expedition to North Greenland. *Rapp. Grønlands geol. Unders.* 135, 8-10, 1987.
- 720 Bennike, O. and Björck, S.: Chronology of the last recession of the Greenland ice sheet, *Journal of Quaternary Science*, 17, 211–219, 2002.

- Bogen, J., Xu, M., and Kennie, P.: The impact of pro-glacial lakes on downstream sediment delivery in Norway, *Earth Surface Processes and Landforms*, 40(7), 942-952, doi:10.1002/esp.3669, 2015.
- 725 Briner, J.P., McKay, N.P., Axford, Y., Bennike, O., Bradley, R.S., de Vernal, A., Fisher, D., Francus, P., Frechette, B., Gajewski, K., Jennings, A., Kaufman, D.S., Miller, G., Rouston, C., Wagner, B.: Holocene climate change in Arctic Canada and Greenland. *Quat. Sci. Rev.* 147, 340-364, doi:10.1016/j.quascirev.2016.02.010, 2016.
- Bronk Ramsey, C.: Bayesian analysis of radiocarbon dates, *Radiocarbon*, 51(1), 337–360, 2009,
- Carrivick J. L., Tweed F. S.: Proglacial lakes: character, behavior and geological importance, *Quaternary Science Reviews* 78, 34–52, 2013.
- 730 Carr, J. R. ,Vieli, A., Stokes, C. R., Jamieson, S. S. R., Palmer, S. J., Christoffersen, P., Dowdeswell, J. A., Nick, F. M., Blankenship, D. D., and Young, D. A.: Basal topographic controls on rapid retreat of Humboldt Glacier, Northern Greenland, *J. Glaciol.* 61,137–150, doi:10.3189/2015JoG14J128, 2015.
- Cook, A. J., Copland, L., Noel, B., Stokes, C. R., Bentley, M., Sharp, M. J., Bingham, R. G., and van den Broeke, M. R.: Atmospheric forcing of rapid marine-terminating glacier retreat in the Canadian Arctic Archipelago. *Science Advances*, 5(3), 735 eaau8507, doi: 10.1126/sciadv.aau8507, 2019.
- Coulthard, R. D., Furze, M. F. A., Pienkowski, A. J., Chantel Nixon, F., England, J. H.: New marine DR values for Arctic Canada, *Quaternary Geochronology*, 5, 419-434, doi:10.1016/j.quageo.2010.03.002, 2010.
- Cronin, T. M., Seidenstein, J., Keller, K., McDougall, K., Reufer, A., & Gemery, L.: The benthic foraminifera *cassidulina* from the Arctic Ocean: Application to paleoceanography and biostratigraphy, *Micropaleontology*, 65(2), 105–125, 2019.
- 740 Davies, V. E. and Krinsley, D. B.: The recent regimen of the ice cap margin in North Greenland. *Assoc. Internat. d'Hydrologie Sci.*, 58, 119-130, 1962.
- Dowdeswell, J. A., Whittington, R., and Marienfield, P.: The origin of massive diamicton facies by iceberg rafting and scouring, Scoresby Sund, East Greenland, *Sedimentology*, 41, 21-35, 1994.
- Dyke, L.M., Andresen, C.S., Seidenkrantz, M.-S., Hughes, A.L.C., Hiemstra, J.F., Murray, T., Bjørk, A.A., Sutherland, 745 D.A., and Vermassen, F.: Minimal Holocene retreat of large tidewater glaciers in Køge Bugt, southeast Greenland. *Sci Rep.* doi: 10.1038/s41598-017-12018-x, 2017.
- Enderlin, E. M., Howat, I. M., and Vieli, A.: High sensitivity of tidewater outlet glacier dynamics to shape, *Cryosphere* 7,1007–1015, doi:10.5194/tc-7-1007-2013, 2013.
- England, J.: Coalescent Greenland and Inuitian ice during the Last Glacial Maximum: revising the Quaternary of the 750 Canadian High Arctic, *Quaternary Science Reviews*, 18(3), 421–456, doi:10.1016/S0277-3791(98)00070-5, 1999.
- England, J. H., Lakeman, T. R., Lemmen, D. S., Bednarski, J. M., Stewart, T. G., Evans, D. J. A.: A millennial-scale record of Arctic Ocean sea ice variability and the demise of the Ellesmere Island ice shelves. *Geophysical Research Letters*, 35, L19502, doi:10.1029/2008GL034470, 2008.
- Funder, S.: Rapport Grønlands Geologiske Undersøgelse. 14C-dating of samples collected during the 1979 expedition to 755 North Greenland, 110, 9-14, 1982.

- Funder, S., Kjeldsen, K. K., Kjær, K. H., O Cofaigh, C.: The Greenland ice sheet during the past 300,000 Years, *Developments in Quaternary Sciences*, 15, 699-713, doi:10.1016/B978-0-444-53447-7.00050-7, 2011a.
- Funder, S., Goosse, H., Jepsen, H., Kaas, E., Kjaer, K.H., Korsgaard, N.J., Larsen, N.K., Linderson, H., Lysa, A., Moller, P., Olsen, J., Willerslev, E.: A 10,000-year record of Arctic Ocean sea-ice variability—view from the beach, *Science*, 333, 747-750. doi:10.1126/science.1202760, 2011b.
- Georgiadis, E., Giraudeau, J., Martinez, P., Lajeunesse, P., St-Onge, G., Schmidt, S. and Massé, G.: Deglacial to postglacial history of Nares Strait, Northwest Greenland: a marine perspective from Kane Basin, *Climate of the Past*, 14(12), 1991–2010, doi:10.5194/cp-14-1991-2018, 2018.
- Hammer, C. U., Johnsen, S. J., Clausen, H. B., Dahl-Jensen, D., Gundestrup, N., Steffensen, J. P.: The paleoclimatic record from a 345 m long ice core from the Hans Tausen Iskappe, *Meddelelser Om. Grøn. Geosci.* 39, 87-95, 2001.
- Hatfield, R.G., Stoner, J.S., Reilly, B.T., Tepley, F.J., Wheeler, B.H., Housen, B.A.: Grain size dependent magnetic discrimination of Iceland and South Greenland terrestrial sediments in the northern North Atlantic sediment record. *Earth Planet. Sci. Lett.* 474, doi: 10.1016/j.epsl.2017.06.042, 2017.
- Heaton, T., Köhler, P., Butzin, M., Bard, E., Reimer, R., Austin, W., Bronk Ramsey, C., Grootes, P. M., Hughen, K. A., Kromer, B., Reimer, P. J., Adkins, J., Burke, A., Cook, M. S., Olsen, J., and Skinner, L.: Marine20—The Marine Radiocarbon Age Calibration Curve (0–55,000 cal BP), *Radiocarbon*, 62(4), 779-820, doi:10.1017/RDC.2020.68, 2020.
- Henriksen, N. and Higgins, A. K.: 2000. Early Palaeozoic Basin Development of North Greenland - Part of the Franklinian Basin, *Polarforschung*, 68, 131-140, 2020.
- Henriksen, N., Higgins, A., Kalsbeek, F., & Pulvertaft, T. C. R.: Greenland from Archaean to Quaternary. Descriptive text to the 1995 Geological map of Greenland, 1:2 500 000. 2nd edition. *GEUS Bulletin*, 18, 1-126. doi:10.34194/geusb.v18.4993, 2009.
- Higgins, A. K.: North Greenland ice islands, *Polar Record*, 25, 207-212, 1989.
- Higgins, A. K., Soper, N. J., and Leslie, A. G.: The Ellesmerian and Caledonian Orogenic Belts of Greenland. *Polarforschung*, 68, 141–151, 1998.
- Higgins, A. K.: North Greenland glacier velocities and calf ice production, *Polarforschung*, 60, 1-23, 1990.
- Hill, E. A, Carr, J. R. and Stokes, C. R.; A Review of Recent Changes in Major Marine-Terminating Outlet Glaciers in Northern Greenland, *Frontiers in Earth Sciences*, 4, 111. doi: 10.3389/feart.2016.00111, 2017.
- Hill, E. A., Carr, J. R., Stokes, C. R. and Gudmundsson, G. H.: Dynamic changes in outlet glaciers in northern Greenland from 1948 to 2015, *The Cryosphere*, 12, 3243-3263, doi:10.5194/tc-12-3243-2018, 2018.
- Hogan, K. A., Jakobsson, M., Mayer, L., Reilly, B. T., Jennings, A. E., Stoner, J. S., Nielsen, T., Andresen, K. J., Nørmark, E., Heirman, K. A., Kamla, E., Jerram, K., Stranne, C. and Mix, A.: Glacial sedimentation, fluxes and erosion rates associated with ice retreat in Petermann Fjord and Nares Strait, north-west Greenland, *The Cryosphere*, 14(1), 261–286, doi:10.5194/tc-14-261-2020, 2020.

- Jakobsson, M., Hogan, K.A., Mayer, L.A., Mix, A., Jennings, A., Stoner, J., Eriksson, B., Jerram, K., Mohammad, R.,
790 Pearce, C., Reilly, B., Stranne, C.: The Holocene retreat dynamics and stability of Petermann Glacier in northwest
Greenland, *Nature Communications*, 9, 2104. doi:10.1038/s41467-018-04573-2, 2018.
- Jakobsson, M., Mayer, L., Nilsson, J., Stranne, C., Calder, B., O'Regan, M., Mix, A., and Ryder¹⁹ Shipboard Scientific
Party: Ryder Glacier in northwest Greenland is shielded from warm Atlantic water by a bathymetric sill, *Nature
Communications Earth & Environment*, 1, 45, doi:10.1038/s43247-020-00043-0, 2020.
- 795 Jamieson, S. S. R., Vieli, A., Livingstone, S. J., Ó Cofaigh, C., Stokes, C., Hillenbrand, C-D., and Dowdeswell, J.: Ice-stream
stability in a reverse bed slope, *Nature Geoscience*, 5, 799-802, doi: 10.1038/NGEO1600, 2012.
- Jennings, A. E., Sheldon, C., Cronin, T. M., Francus, P., Stoner, J. and Andrews, J.: The Holocene history of Nares Strait:
Transition from glacial bay to Arctic-Atlantic throughflow, *Oceanography*, 24(3), 18–33, 2011.
- Jennings, A. E., Andrews, J. T., Oliver, B., Walczak, M. and Mix, A.: Retreat of the Smith Sound Ice Stream in the Early
800 Holocene, *Boreas*, 48(4), 825-840, doi:10.1111/bor.12391, 2019.
- Kaufman, D, Ager, T, Anderson, N, Anderson, P, Andrews, J, Bartlein, P, Brubaker, L, Coats, L, Cwynar, L, Duvall, M,
Dyke, A, Edwards, M, Eisner, W, Gajewski, K., Geirsdóttir, A., Hu, F, Jennings, A, Kaplan, M, Kerwin, M, Lozhkin, A,
MacDonald, G, Miller, G, Mock, C, Oswald, W, Otto-Bliesner, B, Porinchu, D, Rühland, K., Smol, J, Steig, E. . and Wolfe,
B.: Holocene thermal maximum in the western Arctic (0–180°W), *Quaternary Science Reviews*, 23(5–6), 529–
805 560,doi:10.1016/J.QUASCIREV.2003.09.007, 2004.
- Kelly, M. and Bennike, O.: Quaternary geology of parts of central and western North Greenland: a preliminary account.
Rapp. Grønlands geol. Unders. 126, 111-116, 1985.
- Kelly, M., and O. Bennike.: Quaternary Geology of Western and Central North Greenland, *Rapport Grønlands Geologiske
Undersøgelse* 153, GGU, Copenhagen, 34 pp, 1992.
- 810 Koch, L.: Contributions to the glaciology of North Greenland, *Meddelelser om Grønland*, 65, 181-464, 1928
- Landvik, J. Y., Weidick, A., Hansen, A.: The glacial history of the Hans Tausen Iskappe and the last glaciation of Peary
Land, North Greenland, *Meddelelser Om. Grønland. Geosci.* 39, 27-44, 2001.
- Larsen, N. K., Kjaer, K. H., Funder, S., Möller, P., van der Meer, J. J. M., Schomacker, A., Linge, H., and Darby, D. A.: Late
Quaternary glaciation history of northernmost Greenland Evidence of shelf-based ice. *Quaternary Science Reviews*, 29,
815 3339-3414, doi:10.1016/j.quascirev.2010.07.027, 2010.
- Larsen, N. K., Levy, L. B., Carlson, A. E., Buizert, C., Olsen, J., Strunk, A., Bjørk, A. A. and Skov, D. S.: Instability of the
Northeast Greenland Ice Stream over the last 45,000 years, *Nature Communications*, 9(1), 1872, doi:10.1038/s41467-018-
04312-7, 2018.
- Larsen, N. K., Levy, L. B., Strunk, A., Søndergaard, A. S., Olsen, J. & Lauridsen, T. L.: Local ice caps in Finderup Land,
820 North Greenland, survived the Holocene Thermal Maximum, *Boreas*, 48, 551–562, doi:10.1111/bor.12384. ISSN 0300-
9483, 2019.

- Lasher, G. E., Axford, Y., McFarlin, J. M., Kelly, M. A., Osterberg, E. C. and Berkelhammer, M. B.: Holocene temperatures and isotopes of precipitation in Northwest Greenland recorded in lacustrine organic materials, *Quaternary Science Reviews*, 170, 45–55, doi:10.1016/j.quascirev.2017.06.016, 2017.
- 825 Lecavalier, B.S., Fisher, D.A., Milne, G.A., Vinther, B.M., Tarasov, L., Huybrechts, P., Lacelle, D., Main, B., Zheng, J., Bourgeois, J., Dyke, A.S.: High Arctic Holocene temperature record from the Agassiz ice cap and Greenland ice sheet evolution, *Proc. Natl. Acad. Sci.*, 114, 5952-5957, doi:10.1073/pnas.1616287114, 2017
- Madsen, K.N., Thorsteinsson, T.: Textures, fabrics and melt-layer stratigraphy in the Hans Tausen ice core, North Greenland - indications of late Holocene ice cap generation?, *Meddelelser Om. Grøn. Geosci.*, 39, 97-114, 2001.
- 830 McFarlin, J. M., Axford, Y., Osburn, M. R., Kelly, M. A., Osterberg, E. C. and Farnsworth, L. B.: Pronounced summer warming in northwest Greenland during the Holocene and Last Interglacial, *Proc. Natl. Acad. Sci.*, 201720420, doi:10.1073/pnas.1720420115, 2018.
- Miller, G. H., Alley, R. B., Brigham-Grette, J., Fitzpatrick, J. J., Polyak, L., Serreze, M. C., White, J. W. C.: Arctic amplification: can the past constrain the future?, *Quaternary Science Reviews*, 29, 1779-1790.
- 835 doi:10.1016/j.quascirev.2010.02.008, 2010.
- Mörner, N.-A. and Funder, S.: C-14 dating of samples collected during the NORQUA 86 expedition, and notes on the marine reservoir effect, *Meddelelser om Grønland*, 22, 57-59, 1990.
- Möller, P., Larsen, N. K., Kjær, K. H., Funder, S., Schomacker, A., Linge, H., Fabel, D.: Early to middle Holocene valley glaciations on northernmost Greenland. *Quat. Sci. Rev.* 29, 3379-3398, doi:10.1016/j.quascirev.2010.06.044, 2010.
- 840 Moon, T. and Joughin, I.: Changes in ice front position on Greenland's outlet glaciers from 1992 to 2007, *J. Geophys. Res.-Earth*, 113, 1–10, doi:10.1029/2007JF000927, 2008.
- Moon, T., Joughin, I., Sith, B., and Howat, I.: 21st-century evolution of Greenland outlet glacier velocities, *Science*, 336, 576–578. doi:10.1126/science.1219985, 2012.
- Morlighem, M., Williams, C. N., Rignot, E., An, L., Arndt, J. E., Bamber, J. L., Catania, G., Chauché, N., Dowdeswell, J.
- 845 A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L. A., Mouginot, J., Noël, B. P. Y., O'CoFaigh, C., Palmer, S., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., Zinglensen, K. B.: BedMachine v3: Complete Bed Topography and Ocean Bathymetry Mapping of Greenland From Multibeam Echo Sounding Combined With Mass Conservation, *Geophysical Research Letters*, 44 (21), 11,051-11,061, 2017.
- 850 Mottram, R., B. Simonsen, S., Høyer Svendsen, S., Barletta, V.R., Sandberg Sørensen, L., Nagler, T., Wuite, J., Groh, A., Horwath, M., Rosier, J., Solgaard, A., Hvidberg, C.S., Forsberg, R.: An Integrated View of Greenland Ice Sheet Mass Changes Based on Models and Satellite Observations, *Remote Sens.*, 11, 1407, 2019.
- Mouginot, J., Rignot, E., Bjørk, A. A., van den Broeke, M., Millan, R., Morlighem, M., Noël, B., Scheuchl, B., Wood, M.: Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018, *Proceedings of the National Academy of Sciences*,
- 855 116 (19), 9239-9244, 2019.

- Nagler, T., Rott, H., Hetzenecker, M., Wuite, J. and Potin, P.: The Sentinel-1 Mission: New Opportunities for Ice Sheet Observations, *Remote Sensing*, 7(7), 9371–9389, doi:10.3390/rs70709371, 2015, 2015.
- O’Cofaigh, C., Dowdeswell, J. A.: Laminated sediments in glacial marine environments: diagnostic criteria for their interpretation. *Quat. Sci. Rev.* 20, 1411-1436, 2001.
- 860 Olsen, J., Kjær, K. H., Funder, S., Larsen, N. K., Ludikova, A.: High-Arctic climate conditions for the last 7000 years inferred from multi-proxy analysis of the Bliss Lake record, North Greenland, *J. Quat. Sci.*, 27, 318-327. doi:10.1002/jqs.1548, 2012.
- Piret, L., Bertrand, S., Hawkings, J., Kylander, M.E., Torrejón, F., Amann, B. and Wadham, J.: High-resolution fjord sediment record of a receding glacier with growing intermediate proglacial lake (Steffen Fjord, Chilean Patagonia), *Earth Surf Process Landforms*, 46(1), doi: 10.1002/esp.5015, 2021.
- 865 Powell, R. D.: Glacial marine processes at grounding line fans and their growth to ice-contact deltas, *Geol. Soc. London, Spec. Publ.*, 53, 53–73. doi: 10.1144/GSL.SP.1990.053.01.03, 1990.
- Reilly, B.T., Stoner, J.S., Wiest, J.: SedCT: MATLAB™ tools for standardized and quantitative processing of sediment core computed tomography (CT) data collected using a medical CT scanner. *Geochem. Geophys. Geosyst.* 18, 3231-3240, doi:10.1002/2017GC006884, 2017.
- 870 Reilly, B. T., Stoner, J. S., Mix, A. C., Walczak, M. H., Jennings, A., Jakobsson, M., Dyke, L., Glueder, A., Nicholls, K., Hogan, K. A., Mayer, L. A., Hatfield, R. G., Albert, S., Marcott, S., Fallon, S. and Cheseby, M.: Holocene break-up and reestablishment of the Petermann Ice Tongue, Northwest Greenland, *Quat. Sci. Rev.*, 218, 322–342, doi:10.1016/j.quascirev.2019.06.023, 2019.
- 875 Reimer, P.J., Baillie, M.G.L., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Bronk- Ramsey, C., Buck, C.E., Burr, G.S., Edwards, R.L., Friederich, M., Grootes, P.M., Guilderson, T.P., Hajdas, I., Heaton, T.J., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B., McCormac, F.G., Manning, S., Reimer, R.W., Richards, D.A., Southon, J.R., Talamo, S., Turney, C.S.M., van der Plicht, J., and Weyhenmeyer, C.E.: IntCal09 and Marine09 radiocarbon age calibration curves, 0-50,000 years cal BP. *Radiocarbon* 51 (4), 1111-1150, 2009.
- 880 Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, C.B., Buck, C.E., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Haflidason, H., Hajdas, I., Hatte, C., Heaton, T.J., Hoffmann, D.L., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B., Manning, S.W., Niu, M., Reimer, R.W., Richards, D.A., Scott, E.M., Søndergaard, A. S., Larsen, N. K., Steinemann, O., Olsen, J., Funder, S., Egholm, D. L., and Kjær, K. H.: Glacial history of Inglefield Land, north Greenland from combined in situ 10Be and 14C exposure dating, *Clim. Past*, 16, 1999–2015, <https://doi.org/10.5194/cp-16-1999-2020>, 2020.
- 885 Southon, J.R., Staff, R.A., Turney, C.S.M., and van der Plicht, J.: IntCal13 and Marine13 radiocarbon age calibration curves 0-50,000 Years cal BP. *Radiocarbon* 55, 1869-1887. doi:10.2458/azu_js_rc.55.16947, 2013.
- Smith, J. A., Graham, A. G. C., Post, A. L., Hillenbrand, C-D., Bart, P. J., and Powell, R. D.: The marine geological imprint of Antarctic ice shelves, *Nature Communications*, 10, 5635, doi:10.1038/s41467-019-13496-5, 2019.

- 890 Strunk, A., Larsen, N. K., Nilsson, A., Seidenkrantz, M.-S., Levy, L. B., Olsen, J. and Lauridsen, T. L.: Relative sea-level changes and ice sheet history in FINDERUP Land, North Greenland, *Frontiers in Earth Science*, 6, 129, 2018.
- Vermassen, F., Bjørk, A.A., Sicre, M.A., Jaeger, J.M., Wangner, D.J., Kjeldsen, K.K., Siggaard-Andersen, M.L., Klein, V., Mouginit, J., Kjær, K.H. and Andresen, C.S.: A Major Collapse of Kangerlussuaq Glacier's Ice Tongue Between 1932 and 1933 in East Greenland, *Geophys. Res. Lett.*, 47, 1–9, doi:10.1029/2019GL085954, 2020.
- 895 Vinther, B. M., Buchardt, S. L., Clausen, H. B., Dahl-Jensen, D., Johnsen, S. J., Fisher, D. A., Koerner, R. M., Raynaud, D., Lipenkov, V., Andersen, K. K., Blunier, T., Rasmussen, S. O., Steffensen, J. P., Svensson, A. M.: Holocene thinning of the Greenland ice sheet, *Nature*, 461, 385-388, doi:10.1038/nature08355, 2009.
- Wangner, D. J., Jennings, A. E., Vermassen, F., Dyke, L. M., Hogan, K. A., Schmidt, S., Kjær, K. H., Knudsen, M. F. and Andresen, C. S.: A 2000-year record of ocean influence on Jakobshavn Isbræ calving activity, based on marine sediment cores, *Holocene*, 28, 1731–1744. doi:10.1177/0959683618788701, 2018.
- 900 Wood, M., Rignot, E., Fenty, I., An, L., Bjork, A., van den Broeke, M., Cai, C., Kane, E., Menemenlis, D., Milan, R., Morlighem, M., Mouginit, J., Noel, B., Scheuchi, B., Velicogna, I., Willis, J. K., and Zhang, H.: Ocean forcing drives glacier retreat in Greenland, *Science Advances*, 7(1), eaba7282, doi: 10.1126/sciadv.aba7282, 2021.
- Young, N. E., and Briner, J. P.: Holocene evolution of the western Greenland Ice Sheet: Assessing geophysical ice-sheet models with geological reconstructions of ice-margin change, *Quaternary Science Reviews*, 114, 1-17, doi:10.1016/j.quascirev.2015.01.018, 2015.
- 905 Zekollari, H., Lecavalier, B. S., and Huybrechts, P.: Holocene evolution of Hans Tausen Iskappe (Greenland) and implications for the palaeoclimatic evolution of the high Arctic, *Quaternary Science Reviews*, 168, 182-193, doi:10.1016/j.quascirev.2017.05.010, 2017.

910