

Understanding monsoon controls on the energy and mass balance of glaciers in the Central and Eastern Himalaya

Authors: Stefan Fugger^{1,2}, Catriona L. Fyffe^{1,3}, Simone Fatichi⁴, Evan Miles¹, Michael McCarthy^{1,5}, Thomas E. Shaw¹, Baohong Ding⁶, Wei Yang⁶, Patrick Wagnon⁷, Walter Immerzeel⁸, Qiao Liu⁹, and Francesca Pellicciotti^{1,3}

¹Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Birmensdorf, Switzerland

²Institute of Environmental Engineering, ETH Zurich, 8093 Zurich, Switzerland

³Department of Geography and Environmental Sciences, Northumbria University, Newcastle, UK

⁴Department of Civil and Environmental Engineering, National University of Singapore, Singapore

⁵British Antarctic Survey, Cambridge, UK

⁶Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing, China

⁷Univ. Grenoble Alpes, CNRS, IRD, Grenoble-INP, IGE, 38000 Grenoble, France

⁸Faculty of Geosciences, Department of Physical Geography, Utrecht University, Utrecht, The Netherlands

⁹Institute of Mountain Hazards and Environment, Chinese Academy of Sciences, Chengdu, China

Correspondence: Stefan Fugger (stefan.fugger@wsl.ch)

Abstract. The Indian and East Asian Summer Monsoons shape the melt and accumulation patterns of glaciers in High Mountain Asia in complex ways due to the interaction of persistent cloud cover, large temperature ranges, high atmospheric water content and high precipitation rates. Glacier energy and mass balance modelling using in-situ measurements offer insights into the ways in which surface processes are shaped by climatic regimes. In this study, we use a full energy- and mass-balance model and seven on-glacier automatic weather station datasets from different parts of the Central and Eastern Himalaya to investigate how monsoon conditions influence the glacier surface energy and mass balance. In particular, we look at how debris-covered and debris-free glaciers respond differently to monsoonal conditions. The radiation budget primarily controls the melt of clean-ice glaciers, but turbulent fluxes play an important role in modulating the melt energy on debris-covered glaciers. The sensible heat flux decreases during core monsoon, but the latent heat flux cools the surface due to evaporation of liquid water. This interplay of radiative and turbulent fluxes causes debris-covered glacier melt rates to stay almost constant through the different phases of the monsoon. Ice melt under thin debris, on the other hand, is amplified by both the dark surface and the turbulent fluxes, which intensify melt during monsoon through surface heating and condensation. Pre-monsoon snow cover can considerably delay melt onset and have a strong impact on the seasonal mass balance. Intermittent monsoon snow cover lowers the melt rates at high elevation. Based on these results, we expect the mass balance of thick debris-covered glaciers to react less sensitively to projected future monsoon conditions than clean-ice and glaciers with very thin debris. This work is fundamental to the understanding of the present and future Himalayan cryosphere and water budget, while informing and motivating further glacier- and catchment-scale research using process-based models.

1 Introduction

High Mountain Asia (HMA) holds the largest ice volume outside the polar regions (Farinotti et al., 2019) and due to the large elevation range and vast geographic extent, HMA glaciers are highly diverse in character and hydro-climatic context (Yao et al., 2012). Several large-scale weather patterns interact with the region's topography (Bookhagen and Burbank, 2010), causing glaciers' contrast in terms of hypsometry (Scherler et al., 2011a) and accumulation and ablation seasonality (Maussion et al., 2014). The Indian Summer Monsoon dominates the Central Himalaya and the Southeastern Tibetan Plateau during summer, and gradually loses strength moving towards the Karakoram, Pamir and Kunlun ranges in the west, where the influence of Westerlies is particularly strong. A more continental regime, influenced by both monsoon and westerlies, controls the Central Tibetan Plateau (Yao et al., 2012; Mölg et al., 2014), while the East Asia Monsoon influences the eastern slopes of the Tibetan Plateau (Yao et al., 2012; Maussion et al., 2014). These major modes of atmospheric circulation control the surface processes and runoff regimes of glaciers (e.g. Kaser et al., 2010; Mölg et al., 2012, 2014) and lead to distinct responses of glaciers to climate change (Scherler et al., 2011b; Yao et al., 2012; Sakai and Fujita, 2017; Kraaijenbrink et al., 2017). Mass losses are high throughout most of HMA, and are particularly pronounced on the South-Eastern Tibetan Plateau, while glaciers exhibit a near-neutral mass balance regime throughout the Karakoram, Pamir and Kun Lun (Gardelle et al., 2012; Brun et al., 2017; Farinotti et al., 2020; Shean et al., 2020).

Accurate glacier mass balance modelling is essential to assess glacier meltwater contribution to mountain water resources, and to predict future glacier states and catchment runoff. Physically-based models of glacier energy and mass balance represent surface and subsurface energy fluxes using physical equations to calculate the energy available for melt, and the glacier runoff. Summer-accumulation glaciers in HMA experience simultaneous accumulation and ablation. Using an energy balance model, Fujita and Ageta (2000) found that the mass balances of this type of glaciers is highly sensitive to climatic variability during the monsoon season, when warm air temperatures and high precipitation rates coincide. Using energy balance modelling for an interannual study at the Central Tibetan Zhadang glacier, Mölg et al. (2012) demonstrated that the timing of monsoon onset and the associated albedo variability can change melt rates considerably in subsequent years. At the same time, they observed a decoupling of the glacier mass balance from the Indian Summer Monsoon during the main monsoon season. Mölg et al. (2014) explain the mass balance variability of Zhadang Glacier as being controlled by both the Indian Summer Monsoon onset and mid-latitude Westerlies. Combining energy balance with weather forecast modelling, Bonekamp et al. (2019) identify the timing and quantity of snowfalls as the main source of differences in mass balance regimes between the Shimshal Valley in the Karakoram and the Langtang Valley in the Central Himalaya. Similarly, Zhu et al. (2018) attribute mass balance differences of three glaciers on the Tibetan Plateau mainly to different local rain/snow precipitation ratios and timing.

The presence of debris cover, a widespread characteristic of HMA glaciers, (e.g. Scherler et al., 2011b; Kraaijenbrink et al., 2017; Herreid and Pellicciotti, 2020), creates additional complexity to understanding and modelling the processes leading to (sub-debris) glacier melt. In recent years, much effort has gone into developing energy balance models for debris-covered glaciers, (e.g. Nicholson and Benn, 2006; Reid and Brock, 2010; Lejeune et al., 2013; Fujita et al., 2014; Collier et al., 2014; Rounce et al., 2015; Evatt et al., 2015; Steiner et al., 2018). Yang et al. (2017) compare the energy balance of a debris-covered

and a clean-ice glacier on the Southeastern Tibetan Plateau and finds the main differences, beside the differences in melt rates, is their climatic sensitivity and the important role of turbulent fluxes on debris-covered glaciers. Studies with observational data on two Indian glaciers showed that thick debris is a more important control on melt rates than elevation (Pratap et al., 2015; Shah et al., 2019) and also dampens and delays glacier melt in the diurnal cycle (Shrestha et al., 2020). Ablation is often expected to be higher on glaciers with debris around or below the critical thickness (site dependent, 1-5cm) (Nakawo and Rana, 1999) than at both clean-ice sites and at sites with thicker debris cover, as shown experimentally (Östrem, 1959; Reznichenko et al., 2010), and by means of modelling (e.g. Nakawo and Rana, 1999; Reid and Brock, 2010), with humidity being a determining factor for this enhancement (Evatt et al., 2015). Moisture in debris is an important factor under monsoonal conditions, controlling the debris' thermal properties and thus ablation (Sakai et al., 2004; Nicholson and Benn, 2006) and has been the focus of devoted modelling studies (Collier et al., 2014; Giese et al., 2020). Moreover, the representation of latent heat due to evaporation (Steiner et al., 2018; Giese et al., 2020) and atmospheric stability correction for turbulent fluxes were shown to be important to improve the simulation of sub-debris melt (Reid and Brock, 2010; Mölg et al., 2012). Previous studies explicitly dealing with the imprint of the monsoon on the surface thermal properties of glaciers remained limited to individual clean-ice glaciers in the Central Tibetan Plateau (Mölg et al., 2012, 2014).

Our main goal is to improve the understanding of monsoon controls on glaciers of various surface types in the Central and Eastern Himalaya. Applying the glacier energy and mass balance module of a land surface model suited to both debris-covered and clean-ice glaciers, and leveraging seven on-glacier automatic weather station (AWS) records from the region, we answer the following questions: 1) Which energy and mass fluxes dominate the seasonal mass balance of Himalayan glaciers? 2) How does debris cover modulate the energy balance in comparison to clean-ice surfaces? 3) How does the monsoon change the glacier surface energy balance? Answering these questions allows us to infer how these glaciers will respond to the possible future changes of the monsoons in the region. We apply the model at the point scale of individual AWSs, driven by high-quality in situ meteorological observations that guarantee accurate energy balance simulations, not affected by extrapolation of the meteorological forcing. By identifying the key surface processes of glaciers and their dynamics under monsoonal conditions, this study promotes their appropriate representation in models of glacier mass balance and the hydrology of glacierised catchments.

2 Study sites and data

In situ observations from seven on-glacier AWSs in different environments along the climatic gradient of the Himalaya were gathered and are used for forcing and evaluation of the model (Figure 1 and Table 1). Our seven study sites are located in the Central and Eastern Himalaya and cover a range of glacier types and local climates (Figure 1, 2 and Table 2). The seven sites include both spring- (24K, Parlung No.4) and summer-accumulation glaciers (all others) as indicated by the proportion of monsoon precipitation to the annual precipitation (Figure A1). Langtang, Lirung and Yala are neighbouring glaciers found in the Langtang Valley (Figure 1). The Langtang Valley is strongly influenced by the Indian Summer Monsoon (\sim June to October), during which more than 70% of the annual precipitation falls (Figure A1 and Table 2), while the period from November

85 to May is a drier season (Immerzeel et al., 2012; Collier and Immerzeel, 2015). The Valley has been a site of extensive glacio-
logical (e.g. Fujita et al., 1998; Stumm et al., 2020), meteorological (Immerzeel et al., 2014; Collier and Immerzeel, 2015;
Heynen et al., 2016; Steiner and Pellicciotti, 2016; Bonekamp et al., 2019) and hydrological (e.g. Ragettli et al., 2015) investi-
gations. On-glacier AWSs were installed during the ablation season on Lirung (2012-2015) and Langtang (2019) glaciers, and
90 tongues, but the tongue of Lirung has disconnected from the accumulation zone (Figure 2). Yala is a considerably smaller
clean-ice glacier, with most of its ice mass located at comparably high elevation. It is oriented to the southwest and has a gentle
slope (Fujita et al., 1998) (Figure 2 and Table 2).

North Changri Nup Glacier (hereafter Changri Nup Glacier) is a debris-covered valley glacier located in the Everest region in
Nepal (Figure 1). The southeast-oriented, avalanche-fed glacier discharges into the Koshi River system. The local climate is
95 similar to that of the Langtang Valley, with 70-80% of precipitation falling during monsoon (Vincent et al., 2016) (Figure 2,
A1 and Table 2).

24K and Parlung No.4 glaciers are located on the southeastern Tibetan Plateau, feeding water into the upper Parlung Tsangpo,
a major tributary to the Yarlung Tsangpo - Brahmaputra River. The summer climate is characterized by monsoonal air masses
reaching the Gangrigabu mountain range from the south through the Yarlung Tsangpo Grand Canyon. 24K Glacier is an
100 avalanche fed valley glacier with a debris-covered tongue, located 24 km from the town of Bome (Yang et al., 2017). It is small
in size, oriented to the northwest and surrounded by shrubland (Figure 1, 2 and Table 2). Parlung No.4 is a debris-free valley
glacier, which is north-east oriented, considerably larger than 24K and located 130km to the south-east from Bome (Yang et al.,
2011) (Figure 1 and Table 2). Full automatic weather stations were installed in the ablation zones of both glaciers in 2016 and
subsequent years (Table 1).

105 Hailuoguo Glacier, the second-largest of our study sites (Figure 2) is located on the eastern slope of Mt. Gongga in the east-
ernmost portion of the southeastern Tibetan Plateau (Figure 1). It is located at low elevation and large parts of its ablation zone
are continuously covered with a thin layer of fine clasts and scattered with coarser clasts, leading to high annual ablation rates
(Figure 2 and Table 2). The local climate influenced by the East Asia Monsoon with typically only 50 to 60% of the annual
precipitation arriving during the monsoon period (Figure 1 and A1). The debris-covered tongue is connected to a steep and
110 extensive accumulation zone via a large icefall, but avalanching is the primary mass supply mechanism through the icefall to
the valley tongue (Liao et al., 2020), and a dynamic disconnect is expected to occur in the near future. Weather stations were
installed at three nearby off-glacier locations and one on-glacier site during 2008, while precipitation was measured at the
Alpine Ecosystem Observation and Experiment Station of Mt. Gongga, within 1.5 km from the glacier terminus (Table 1).

115 We use the monthly averaged ERA5-Land reanalysis data (Muñoz Sabater, 2019) to evaluate the representativeness of the
AWS records in terms of seasonal variability (Figures A3 to A9), and to provide an overview of the long term climatic patterns,
e.g. the average monsoonal regime from June through September (Figure A1). We thereby focus on the qualitative aspects,
given that the absolute values from the reanalysis dataset are not representative for the AWS location at the glacier surfaces. A




Figure 1. (a) shows the context of study sites with respect to large-scale weather patterns, topography and glacier distribution (blue, source: Randolph Glacier Inventory 6.0). Blue dots indicate clean-ice study glaciers and brown dots indicate debris-covered study glaciers. (b) displays the spatial pattern of average annual precipitation from ERA5-Land (1981-2019). (c) shows the monsoonal (June-September) portion of the ERA5-Land total annual precipitation (MI). Background map source: ESRI, U.S. National Park Service.

detailed description is given in the Supplementary.




Figure 2. Characteristics of study sites, summarized (center) in terms of elevation, mean measured ice melt rate, measured debris thickness and JJAS contribution to the ERA5-Land total annual (1981-2019) precipitation (monsoon index; MI). For each site, we also show glacier (bars in aqua) and debris (bars in olive) hypsometry, with area on the x-axis [km²] and altitude on the y-axis [m.asl], and glacier and supraglacial debris extents.

Table 1. Summary of available meteorological and ablation observations at each site, as well as each site’s model period. Variables indicated with * were transferred from neighboring weather station. Variables with ** were reconstructed based on other variables measured at the same station.

	AWS Location				Variables measured			Model period	Reference
	Lat	Lon	Elevation [m.a.s.l.]	Debris thickness [cm]	AWS	Precipitation	Ablation		
Lirung	28.233	85.562	4076	30	$T, RH, W_s, W_d, SW_{\downarrow},$ $SW_{\uparrow}, LW_{\uparrow}, LW_{\downarrow}, P_{atm}^*$	Pluvio Kyangring and Yala Basecamp, 3857m.asl and 5090m.asl, hourly, partly lapsed	SR50	2014-05-05/ 2014-10-24	Ragetti et al. (2015)
Langtang	28.279	85.722	4536	50	$T, RH, W_s, W_d, SW_{\downarrow},$ $SW_{\uparrow}, LW_{\downarrow}^{**}, LW_{\uparrow}, P_{atm}^*$	Pluvio Morimoto base camp 4919m.asl, hourly	SR50	2019-05-11/ 2019-10-30	unpublished
Yala	28.233	85.612	5350	-	$T, RH, W_s, W_d, SW_{\downarrow},$ $SW_{\uparrow}, LW_{\uparrow}, LW_{\downarrow}, P_{atm}^*$	Pluvio Yala base camp 5090m.asl, hourly	SR50	2019-05-01/ 2019-10-31	ICIMOD (2021)
Changri Nup	27.993	86.780	5470	10	$T, RH, W_s, W_d, SW_{\downarrow},$ $SW_{\uparrow}, LW_{\uparrow}, LW_{\downarrow}, P_{atm}^*$	Pluvio at Pyramid meteorological station (4993 m a.s.l.), 4.9 km SE of AWS location, hourly	SR50	2016-05-01/ 2016-10-31/	Wagnon (2021)
24K	27.983	86.778	3900	20	$T, RH, W_s, W_d, SW_{\downarrow},$ $SW_{\uparrow}, LW_{\uparrow}, LW_{\downarrow}, P_{atm}^*$	On-glacier tipping bucket, hourly	stake	2016-06-01/ 2016-09-29	Yang et al. (2017)
Parlung No.4	29.761	95.720	4806	-	$T, RH, W_s, W_d, SW_{\downarrow},$ $SW_{\uparrow}, LW_{\uparrow}, LW_{\downarrow}, P_{atm}^*$	Pluvio, near terminus 4600m.asl, hourly	stake	2016-05-01 2016-10-31	Yang et al. (2017)
Hailuogou	29.558	101.969	3550	1	$T, RH, W_s, W_d, SW_{\downarrow},$ $SW_{\uparrow}, LW_{\uparrow}, LW_{\downarrow}, P_{atm}^*$	Pluvio at GAEORS station, 3000m.asl, 1.5km from terminus	stake	2008-05-15 2008-10-31	Zhang et al. (2011)

Table 2. Characteristics of the study sites. Planimetric glacier and debris surface areas, mean elevation, slope and aspect were calculated using the updated Randolph Glacier Inventory 6.0 by Herreid and Pellicciotti (2020) and the USGS GTOPO30 digital elevation model. Slope (mean) and aspect (vectorial average) for the whole glacier. MI (‘Monsoon-Index’) is the mean June-September portion of the ERA5-Land total annual precipitation (1981-2019); For Lirung, where the ablation zone has dynamically disconnected from the accumulation zone, the glacier characteristics represent both zones together.

	Area [km^2]		Elevation [$m.asl$]			Slope	Aspect	MI
	Glacier	Debris	min	max	mean	[°]	[°]	[-]
Lirung (LIR)	4.0	1.5	3990	6830	5100	27.6	151.2	0.74
Langtang (LAN)	37.0	17.8	4500	6620	5330	16.0	177.5	0.71
Yala (YAL)	1.4	-	5170	5660	5390	23.5	229.2	0.74
Changri Nup (CNU)	2.7	1.4	5270	6810	5610	15.9	189.4	0.76
24K (24K)	2.0	0.9	3910	5070	4350	18.3	302.6	0.46
Parlung No.4 (NO4)	11.0	-	4620	5950	5330	17.1	23.5	0.40
Hailuogou (HAI)	24.5	4.1	2980	7470	5360	27.0	104.3	0.56

3 Methods

3.1 Tethys-Chloris energy balance model

We use the hydrological, snow and ice modules of the Tethys-Chloris (T&C) land surface model (Fatichi et al., 2012; Paschalis et al., 2018; Mastrotheodoros et al., 2020; Botter et al., 2020) to simulate the mass and energy balance of the seven study glaciers. The T&C model simulates, in a fully distributed manner, the energy and mass budgets of a large range of possible land surfaces, including vegetated land, bare ground, water, snow and ice. Here, we apply the model at the point scale of the AWS locations to simulate the energy fluxes of the underlying surface and subsurface, which can comprise snow, ice and supraglacial debris cover layers, according to the local and dynamic conditions. The melt and accumulation of ice and snow, and the ice melt under debris are also explicitly simulated. The surface energy balances for the three different possible surfaces are for snow,

$$R_n(T_{sno}) + Q_v(T_{sno}) + Q_{fm}(T_{sno}) - H(T_{sno}) - \lambda E(T_{sno}) - G(T_{sno}) - M(T_{sno}) = 0, \quad (1)$$

for debris cover,

$$R_n(T_{deb}) + Q_v(T_{deb}) - H(T_{deb}) - \lambda E(T_{deb}) - G(T_{deb}) = 0, \quad (2)$$

and for ice,

$$R_n(T_{ice}) + Q_v(T_{ice}) - H(T_{ice}) - \lambda E(T_{ice}) - G(T_{ice}) - M(T_{ice}) = 0, \quad (3)$$

where $R_n [W m^{-2}]$ is the net radiation absorbed by the snow/debris/ice surface, $Q_v [W m^{-2}]$ is the energy advected from precipitation, $Q_{fm} [W m^{-2}]$ is the energy gained or released by melting or refreezing the frozen or liquid water that is held inside the snow pack, $H [W m^{-2}]$ is the sensible energy flux and $\lambda E [W m^{-2}]$ the latent energy flux for any of the surfaces, and $G [W m^{-2}]$ is the conductive energy flux from the surface to the subsurface. In ice, the conduction of energy is represented in the model down to a depth of 2 m after which it is assumed the ice pack is isothermal. Finally, $M [W m^{-2}]$ is the net energy input to the snow or ice pack. For debris on top of ice, and snow on top of debris or ice, the in-/outgoing fluxes towards/from the ice are adjusted according to the respective interface type. The sign convention is such that fluxes are positive when directed towards the surface. To close the energy balance, a prognostic temperature T_s is estimated for each computational element. Iterative numerical methods are used to solve the non-linear energy budget equation until convergence for the ice and snow surface, and the heat diffusion equation for the debris surface, while concurrently computing the mass fluxes resulting from snow and ice melt and sublimation. To close the energy balance, a prognostic temperature for the different surface types (T_{sno} , T_{deb} , T_{ice}) is estimated for each computational element. Iterative numerical methods are used to solve the non-linear energy budget equation until convergence for the ice and snow surface, and the heat diffusion equation for the debris surface, while concurrently computing the mass fluxes resulting from snow and ice melt and sublimation. In the case of snow, debris and ice surfaces, either of which is simulated to always fully cover a computational element, T_{sno} , T_{deb} or T_{ice} are equivalent to the element's overall surface temperature T_s . In the following, we use the surface type specific symbol for surface specific

equations, while we use T_s for equations valid for all three surface types.

155 3.1.1 Radiative fluxes

R_n is calculated as the sum of incoming and outgoing shortwave and longwave fluxes as

$$R_n = SW_{\downarrow}(1 - \alpha) + LW_{\downarrow} + LW_{\uparrow}, \quad (4)$$

where $SW_{\downarrow} [W m^{-2}]$ is the incoming shortwave radiation, $\alpha [-]$ is the surface albedo, $LW_{\downarrow} [W m^{-2}]$ and $LW_{\uparrow} [W m^{-2}]$ are the incoming atmospheric and outgoing longwave radiation components, respectively. In this study α is given as an input
160 to the model based on the AWS observations. We prescribe α for all surface types as the daily cumulated albedo, which is the 24 hour sum of SW_{\uparrow} divided by the sum of SW_{\downarrow} centred over the time of observation (van den Broeke et al., 2004).

3.1.2 Incoming energy with precipitation

For calculating the incoming energy with precipitation, rain is assumed to fall at air temperature (T_a) when positive, with a lower boundary of $0^\circ C$. Snow is assumed to fall at negative T_a with an upper boundary of $0^\circ C$. Here, Q_v is the energy
165 required to equalize the precipitation temperature with the surface temperature T_s and is therefore calculated as

$$Q_v = c_w P_{r,liq} \rho_w [\max(T_a, 0) - T_s] + c_i P_{r,sno} \rho_w [\min(T_a, 0) - T_s], \quad (5)$$

where $c_w = 4196 [J kg^{-1} K^{-1}]$ is the specific heat of water, $c_i = 2093 [J kg^{-1} K^{-1}]$ the specific heat of ice, $\rho_w = 1000 [kg m^{-3}]$ is the density of water and $P_{r,liq} [mm]$, $P_{r,sno} [mm]$ are the rain- and snowfall intensities, respectively.

3.1.3 Phase changes in the snow pack

170 The snow pack has a water holding capacity Sp_{wc} described in section 3.2.2. The energy flux gained/released by melting/refreezing the frozen/liquid water that is held inside the snow pack is calculated as:

$$Q_{fm}(t) = \begin{cases} f_{sp} \frac{\lambda_f \rho_w Sp_{wc}(t-dt)}{1000 dt}, & T_{sno}(t) < 0 \text{ and } T_{sno}(t-dt) \geq 0 \\ -f_{sp} \frac{\lambda_f \rho_w Sp_{wc}(t-dt)}{1000 dt}, & T_{sno}(t) \geq 0 \text{ and } T_{sno}(t-dt) < 0 \end{cases} \quad (6)$$

where $f_{sp} = \frac{5}{SWE} []$ with $\max(f_{sp}) = 1$ is the fraction of the snowpack water equivalent ($SWE [mm w.e.]$) involved in either melting or freezing. This choice was made in order to mimic refreezing in the upper portion of the snowpack, while the
175 snowpack is otherwise represented as a single layer. $\lambda_f = 333700 [J kg^{-1}]$ is the latent energy of melting and freezing of water, t stands for time, $dt [s]$ is the timestep, and the unit for T_{sno} is $[^\circ C]$.

3.1.4 Turbulent energy fluxes

Over snow, debris and ice surfaces, the sensible energy flux is calculated as

$$H = \rho_a C_p \frac{(T_s - T_a)}{r_{ah}}, \quad (7)$$

180 where $T_s [^{\circ}C]$ is the surface temperature (generalised term for $T_{sno}, T_{deb}, T_{ice}$), $C_p = 1005 + [(Ta + 23.15)^2] / 3364 [J kg^{-1} K^{-1}]$ is the specific heat of air at constant pressure, $\rho_a [kg m^{-3}]$ is the density of air. The aerodynamic resistance $r_{ah} [s m^{-1}]$, which is also a function of wind speed (Ws) is calculated using the simplified solution of the Monin-Obokhov similarity theory (Mascart et al., 1995; Noilhan and Mahfouf, 1996). The roughness lengths of heat ($z_{0h} [m]$) and water vapour ($z_{0w} [m]$) used in the calculation of the aerodynamic resistance are equal in the T&C model ($z_{0h} = z_{0w}$), and ($z_{0h} = z_{0w} = 0.1 z_{0m}$). The roughness
185 length of momentum (z_{0m}) is set to 0.001 m for snow and ice surfaces (Brock et al., 2000), while we optimize it against the surface temperature for debris (see section 3.3).

Correct estimates of the latent energy flux due to water phase changes at the surface are important to accurately model glacier melt, especially under moist conditions (Sakai et al., 2004). Phase changes between the water and gas phase and the resulting
190 energy fluxes are considered over all surfaces. The latent energy is limited by the availability of water in the form of ice and snow, or in the case of a debris surface, by the amount of water intercepted (interception storage capacity is set to 2mm). The latent energy flux is estimated from:

$$\lambda E = \lambda_s \frac{\rho_a (q_{sat}(T_s) - q_a)}{r_{aw}}, \quad (8)$$

where λ_s is the latent energy of sublimation defined as $\lambda_s = \lambda + \lambda_f$, with $\lambda = 1000 (2501.3 - 2.361 T_a) [J kg^{-1}]$ as the latent
195 energy of vaporisation. q_{sat} is the surface specific humidity at saturation at T_s , q_a is the specific humidity of air at the measurement height and r_{aw} the aerodynamic resistance to the vapour flux, which we assume equals r_{ah} .

Controls on turbulent fluxes

The predictive power of the temperature gradient between surface and air $\delta_T [^{\circ}C^{-1}]$ and Ws as well as their combination for
200 determining H and LE were assessed using a univariate polynomial regression model for the single predictors and a multiple polynomial regression model using both variables, and both models had 2 degrees of freedom.

3.1.5 Ground energy flux

The definition of the ground energy flux $G [W m^{-2}]$ differs based on the surface type. When there is snow, it is equal to the energy transferred from the snowpack to the underlying ice or debris surface. In the assumption of a slowly changing process,
205 G can be approximated with the temperature difference of the previous time step ($t-1$), which allows to solve for G outside the numerical iteration to find the snow surface temperature of the current time step:

$$G_{sno}(t) = k_{sno} \frac{T_{sno}(t-1) - T_{deb,ice}(t-1)}{d_{sno}} \quad (9)$$

where $k_{sno}[W K^{-1} m^{-1}]$ is the thermal conductivity of snow and $d_{sno}[m]$ is the snow depth. For ice in the absence of snow and debris, it is the energy flux from the ice pack to the underlying surface or to the ice at a depth of $2m$:

$$210 \quad G_{ice}(t) = k_{ice} \frac{T_{ice}(t-1) - T_{grd}(t-1)}{d_{ice}} \quad (10)$$

where $k_{ice}[W K^{-1} m^{-1}]$ is the thermal conductivity of ice, $T_{grd}[^{\circ}C]$ is the temperature of the underlying ice, and $d_{ice}[m]$ is the relevant ice thickness. For debris, which was discretised into eight layers at all debris-covered sites, G is the energy flux conducted into the debris layers. Its calculation is for a given time t and depth z

$$G(z, t) = -k_d \frac{\partial T_{deb}(z, t)}{\partial z_d}, \quad (11)$$

215 where $k_d[W K^{-1} m^{-1}]$ is the debris thermal conductivity (see section 3.3) and $T_{deb}(z, t)[^{\circ}C]$ is the debris temperature at time t and depth z . $G(z, t)$ can be included in the heat diffusion equation as such:

$$cv_s \frac{\partial T_{deb}(z, t)}{\partial t} = \frac{\partial}{\partial z_d} (-G(z, t)), \quad (12)$$

where cv_d is the debris heat capacity. Under the assumption of homogeneous debris layers, $\kappa[m^2 s^{-1}]$ as the debris heat diffusivity replaces the term $\frac{k_d}{cv_s}$ and equation (12) can be written as:

$$220 \quad \frac{\partial T_{deb}(z, t)}{\partial t} = \kappa \frac{\partial^2 T_{deb}(z, t)}{\partial z^2}, \quad (13)$$

The heat diffusion equation (13) is solved using iterative numerical methods. This way, the debris temperature profile $T_{deb}(z, t)$ is solved together with $G(z, t)$ at any depth and time. The conductive energy flux at the base of the debris is used to heat the ice and to calculate ice melt once above the melting point.

Note, that G can also act in the opposite direction, i.e. when energy is conducted from the snowpack/debris/ice towards the
225 surface. In our results, G sums up all types of conductive energy fluxes in the snow-debris-ice column. Part of the energy is used for heating the snow, ice or debris-ice surface layer until melt occurs (G , Table A2).

3.2 Mass balance in the T&C model

3.2.1 Precipitation partition

Precipitation is partitioned into solid Pr_{sno} and liquid Pr_{liq} precipitation, because of the differing impacts of snow and rain on
230 the energy and mass balance. For this study, the precipitation partition method described by Ding et al. (2014) was implemented into the T&C model. This parameterisation has been developed specifically for High Mountain Asia based on a large dataset of rain, sleet and snow observations, and does not require recalibration. It determines the precipitation partition based on the wet-bulb temperature, station elevation and relative humidity and allows for sleet events, as a mixture between liquid and solid precipitation. Ding et al. (2014) found the wet-bulb (T_{wb}) to be a better predictor than T_a of the precipitation type. They also
235 found that the temperature threshold between snow and rain increases at higher elevations, and that the probability of sleet is reduced in conditions of low relative humidity.

3.2.2 Water content of the snow, ice and debris layers

The water content of ice is approximated with a linear reservoir model. The liquid water outflow is proportional to the ice pack water content $I_{p_{wc}} [mm w.e.]$, which is initiated when $I_{p_{wc}}$ exceeds a threshold capacity, prescribed as 1% of the ice water equivalent ($IWE [mm w.e.]$). The $I_{p_{wc}}$ is the sum of ice melt and liquid precipitation, minus the water released from the ice pack. The water released is the sum of the ice pack excess water content plus the outflow from the linear reservoir, given as $I_{out} = I_{p_{wc}} / K_{ice}$, where K_{ice} is the reservoir constant which is proportional to the ice pack water equivalent. Unlike within snow packs, Q_{fm} is not accounted for within the ice pack, since water is presumed to be evacuated quickly from the ice due to runoff without refreezing.

The water content of the snow pack $Sp_{wc} [mm w.e.]$ is approximated using a bucket model, in which outflow of water from the snow pack occurs when the maximum holding capacity of the snow pack is exceeded. Following the method of Bélair et al. (2003), the maximum holding capacity of the snow pack is based on SWE , a holding capacity coefficient and the density of snow ρ_{sno} . Snowmelt plus liquid precipitation, minus the water released from the snow pack gives the current Sp_{wc} . If T_{sno} is greater than $0^\circ C$ then the snow pack water content is assumed to be liquid, whereas otherwise it is assumed frozen.

For supraglacial debris, both observations and methods for modelling its water content are lacking. We thus use a simplified scheme for moisture at the surface of the debris, in order to mimic the drying process of the debris surface: we assume debris to have a dynamic interception storage, which can hold a maximum of $2 mm$ water at all debris-covered sites and can be refilled by snowmelt or liquid precipitation. The evaporative flux from the debris and the latent energy flux of evaporation is therefore limited by this interception storage.

3.2.3 Snow and ice mass balance

The mass balance calculation of snow and ice is rather similar, so they will be described together here. Calculations are performed for snow if there is snow precipitation during a timestep or the modelled SWE at the surface is greater than zero. Net input of energy to the snow or ice pack will increase its temperature, and after the temperature has been raised to the melting point, additional energy inputs will result in melt. The change in the average temperature of the ice or snowpack dT is controlled using:

$$dT = \frac{M dt}{c_i \rho_w W E_b} 1000, \quad (14)$$

Where dt is the time step $[h]$ and $W E_b [mm w.e.]$ is IWE or SWE before melting and limited to a maximum of $2000 mm$, assumed to be the water equivalent mass exchanging energy with the surface. Energy inputs into an iso-thermal ice/snow pack result in melt $M [mm w.e.]$ as

$$M = \frac{M dt}{\lambda_f \rho_w} 1000 \quad (15)$$

The water equivalent mass of the snow/ice pack after melting $WE(t)$ [mm w.e.] is updated conserving the mass balance following:

$$WE(t) = WE(t - dt) + Pr_{sno}(t) - E(t) dt - M(t), \quad (16)$$

270 Here $E = \lambda E / \lambda_s$ [mm] is the sublimation from ice and snow. The snow density is assumed to be constant with depth and calculations are performed assuming one single snow pack layer. The snow density evolves over time using the method proposed by Verseghy (1991) and improved by Bélair et al. (2003). In this parameterisation the snow density increases exponentially over time due to gravitational settling and is updated when fresh snow is added to the snowpack. Two parameters are required in this scheme, ρ_{sno}^{M1} and ρ_{sno}^{M2} [$kg m^{-3}$], which represent the maximum snow density under melting and freezing conditions, respectively. The depth of the ice pack can be increased through the formation of ice from the snow pack (ice accumulation), 275 which is prescribed to occur if the snow density increases to greater than $500 kg m^{-3}$ (a density associated with the firm to ice transition) and at a rate of $0.037 mm h^{-1}$ (Cuffey and Paterson, 2010). The density of ice is assumed constant with depth and equal to $916.2 kg m^{-3}$.

3.3 Debris parameters

280 A major challenge in physically based mass-balance modelling of debris-covered glaciers is the selection of appropriate debris properties. In addition to the debris thickness, which was measured at the AWS location, values are needed for the thermal conductivity k_d , the roughness length z_0 of the debris surface, the surface emissivity ϵ_d , the debris volumetric heat capacity cv_d and the debris density ρ_d . While the latter three can be quantified using literature values, there is more uncertainty about k_d and z_0 , two highly variable quantities that are difficult to measure in the field. We thus choose to optimize them, since our 285 primary requirement is an accurate representation of the energy and mass balance: (1) in a first step, we optimize k_d simulating only the conduction of energy through the debris during snow free conditions, with the LW_{\uparrow} -derived surface temperature $T_{s,LW}$ as an input, the ice melt as the target variable and the Nash-Sutcliffe Efficiency $NSE[-]$ as performance metric. (2) Next, we run the whole energy balance model and optimize z_0 for snow-free conditions, with all required meteorological inputs, and the optimal k_d from step (1), while comparing modelled T_s against $T_{s,LW}$, using NSE as performance metric. The 290 resulting parameters are given in Table 4. All optimized values fall within the expected range based on prior energy-balance studies on debris-covered glaciers (Nicholson and Benn, 2006; Reid and Brock, 2010; Lejeune et al., 2013; Rounce et al., 2015; Collier et al., 2015; Evatt et al., 2015; Yang et al., 2017; Miles et al., 2017; Quincey et al., 2017; McCarthy, 2018; Rowan et al., 2020).

Table 3. Debris parameter values for each site derived from the two-step optimization procedure.

	Lirung	Langtang	Changri Nup	24K	Hailuoguo
k_d [$W m^{-1} K^{-1}$]	1.09	1.65	1.77	1.45	0.72
z_{0m} [m]	0.7	0.38	0.11	0.15	0.027

3.4 Uncertainty estimation

295 We calculate the uncertainty associated with all energy and mass balance components by performing 10^3 Monte Carlo simulations for each study site at the AWS location. We perturb three debris parameters (k_d, z_{0m}, ϵ_d), debris thickness h_d , as well as six measured model input variables: air temperature T_a , the vapour pressure at reference height $ea[Pa]$, SW_{\uparrow} , SW_{\downarrow} , LW_{\downarrow} , the total precipitation before partition Pr , and the wind speed Ws . Measured outgoing shortwave radiation SW_{\uparrow} was included into the Monte Carlo set, as it determines our input α , as discussed in Section 3.1.1. While the parameter uncertainty range was defined based on previously published values for debris (e.g. Yang et al., 2017; Rounce et al., 2015; Evatt et al., 2015; Reid and Brock, 2010; Nicholson and Benn, 2006; Rowan et al., 2020; Lejeune et al., 2013; Collier et al., 2015; Miles et al., 2017; Quincey et al., 2017; McCarthy, 2018), the debris thickness measurement uncertainty was given with a range of 1 cm and the range for the meteorological inputs was set based on the respective sensor uncertainties (see Table 5). All uncertainties were equally distributed around the standard parameter values and observations. Uncertainties are given as one standard deviation of the error of the Monte Carlo runs against the standard run.

Table 4. Debris parameter values for each site derived from the two-step optimization procedure.

	Lirung	Langtang	Changri Nup	24K	Hailuogou
$k_d [W m^{-1} K^{-1}]$	1.09	1.65	1.77	1.45	0.72
$z_{0m} [m]$	0.7	0.38	0.11	0.15	0.027

Table 5. Uncertainty ranges of parameters and input variables used for Monte Carlo runs. Where units are indicated with [-], the parameter or variable was perturbed by the fractional value shown.

Parameter/ Variable	Range	Parameter/ Variable	Range
k_d [-]	± 0.1	SW_{\downarrow} [-]	± 0.03
z_0 [-]	± 0.1	SW_{\uparrow} [-]	± 0.03
ϵ_d [-]	± 0.05	LW_{\downarrow} [-]	± 0.03
h_d [mm]	± 5	Pr [-]	± 0.15
Ta [$^{\circ}C$]	± 0.2	W_s [m/s]	± 0.3
ea [-]	± 0.02		

3.5 Model evaluation

The model accurately reproduces the measured surface height change (ablation and accumulation) at both debris-covered and clean-ice glaciers (Figure 3). The maximum uncertainties associated with each model output ranges from $\pm 4\%$ (Parlung No.4, Figure 3f) to $\pm 15\%$ (Yala, Figure 3c). Where Ultrasonic Depth Gauge (UDG) records are available (Lirung, Langtang, Yala, Changri Nup), the deviations of the simulations from the observations stay within the uncertainty range (Figure 3a-d). We decided to not consider the UDG record from Changri Nup after a large August snowfall, as variables describing the surface state (e.g. α , LW_{\uparrow}) following this event indicate a discontinuous snow cover at the AWS location, while the UDG, which is some meters away from the AWS, shows continuous snow cover with depths of tens of centimeters. This discrepancy was also confirmed by observation of the site from October 2016. It was thus not possible to match the UDG record with the model for the late ablation period on Changri Nup, but the model closely reproduces observed surface height change for the pre-monsoon and early monsoon (Figure 3d), when AWS and UDG observations agree in terms of surface state. The deviation to measured melt is larger than the uncertainty range at 24K and Parlung No.4 for two and one individual stake readings, respectively, but the overall agreement is very good also at these sites (Figure 3e, f). For Parlung No.4 there are no stake measurements available before July 21 due to the long-lasting snow cover.




Figure 3. (a)-(g) Observed vs. modelled surface change at all study sites, including ice melt, snow melt, sublimation, precipitation phase and snow depth. Measured melt is either from ablation stakes (black crosses) or Ultrasonic Depth Gauges (black lines). Vertical dotted lines indicate monsoon onset and end.

4 Results

4.1 Modelled mass balance

The ablation season average melt rates vary considerably across sites: the highest value of 42.7 mm d^{-1} is reached at the low-lying site with thin debris cover, Hailuogou, and the lowest value of 6 mm d^{-1} is evident at Langtang, a site at moderate elevation but with the thickest debris cover out of all study sites (Figure 4). The largest average seasonal mass loss component at all sites is ice melt, with a minimum of 65.8% of the mass losses at Changri Nup (Figure 4c) and up to 95.4% at Hailuogou, (Figure 4g). This is followed by snowmelt, accounting for only 0.1% at 24K (Figure 4e) but as much as 33.1% at Yala (Figure 4c) of the seasonal mass losses. Sublimation from ice and snow represents a very small share of the seasonal mass losses, and ranges from 0.01% (Lirung, Figure 4a) to 1.2% (Changri Nup, Figure 4d). It mostly occurs under dry conditions during pre-monsoon at the highest sites (Changri Nup, Yala).

The timing of snow cover is an important control of both amounts and patterns of ice melt, as ice melt rates are close to zero during periods of snow cover. This becomes clear in Figure 4, where ice melt rates are low during weeks when also snow melt




Figure 4. (a)-(g) Melt rates of ice and snow (stacked) as weekly averages at each site. Vertical dotted lines indicate monsoon onset and end. Error bars depict the uncertainty (standard deviation) of the estimates. Melt rates are normalized to the mean of the ice melt over the entire period (value in the upper left of each panel).

takes place. A long lasting pre-monsoonal snowpack can delay the onset of ice melt considerably, e.g. at Parlung No.4, where ice melt is delayed until the end of June (Figures 3f and 4f). Similarly, intermittent snow cover protects the ice from melting at the two highest sites (Yala and Changri Nup) during the summer months (Figure 3c-d and 4c-d).

4.2 Modelled energy balance

The largest components in the energy balance are LW_{\uparrow} , LW_{\downarrow} and SW_{\downarrow} (Figure A2). The two longwave fluxes counteract and offset each other in large parts resulting in a moderate, melt-reducing LW_{net} , which reaches its highest values during the pre- and post-monsoon (Figure 5). SW_{\downarrow} and its reflected counterpart SW_{\uparrow} result in a net shortwave flux SW_{net} , which at all sites contributes the overall largest amount of energy available for melt M (Figure 5). M is additionally increased or reduced by the turbulent fluxes H and LE , while the energy advected to the glacier surface by precipitation (Qv) remains small ($< 2 W m^{-2}$, Table A2). G links the snow/debris/ice surface to the subsurface, and is a result of all surface fluxes and the subsurface conditions. Before ice melt occurs, depending on season and site, a part dG of G between 0 and $17.8 W m^{-2}$,




Figure 5. (a)-(g) Stacked energy fluxes weekly averages at each site, depicting the components SW_{net} , LW_{net} , H , LE , Q_v , Q_{fm} , G and M . Energy fluxes are negative fluxes when directed away from the surface and positive when directed towards the surface.

is invested into warming the debris or ice pack to the melting point (Table A2). dG tends to be larger during pre-monsoon and
 345 at the higher sites (Yala, Changri Nup), where ambient temperatures frequently fall below 0°C .

4.3 Impact of debris cover

Debris cover modulates the energy balance in several ways: with the albedo of the snow free debris surface ranging between 0.05 (24K) and 0.22 (Changri Nup), a much larger amount of SW_{\downarrow} is absorbed by the surface than on clean ice glaciers, where the albedo typically ranges between 0.3 and 0.6. In contrast to clean-ice glaciers however, where the main flux re-emitting
 350 absorbed energy is LW_{\uparrow} (Figure 5c and f), a large part of the debris-absorbed energy is also re-emitted to the atmosphere by the turbulent fluxes H and LE (Figure 5a,b,d and e). As a result of this insulating effect of debris, the seasonal average melt rate of debris-covered 24K is considerably lower (19.8 mm d^{-1}) than that of clean-ice Parlung No.4 (34.4 mm d^{-1}), despite the latter site being 900 m lower in elevation than the former one (Figure 4e and f), and their geographical proximity (1). On Hailuogou, the site with very thin debris however, the turbulent fluxes act in the opposite direction, i.e. contributing energy for
 355 melt. Summed up, they can reach weekly averages of 150 W m^{-2} (Figure 5g).

4.4 Impact of the monsoon

During monsoonal conditions, increased cloudiness results in SW_{\downarrow} decreasing its melt contribution at all sites compared to pre-monsoonal conditions (Figure 6) with changes ranging between -41.8 (Hailuogou, pre-monsoon: 178.2; monsoon: 136.4) and -135 (Yala, pre-monsoon: 307.7; monsoon: 172.7) at the seven sites (all values in $W m^{-2}$, from Table A2). Note that we express fluxes in terms of the net energy absorbed by, or removed from the snow/debris/ice surface, (with positive and negative fluxes indicating energy absorbed and removed from surface, respectively). Reflected shortwave radiation SW_{\uparrow} , which removes energy from the surface, and which is controlled by the surface albedo, follows these changes (Figure 6), becoming less negative by $+5.4$ (24K, pre: -18.5 , mon: -13.8) and up to $+164.8$ (Parlung No.4, pre: -219.6 , mon: -54.8) between sites. An exception to this is Changri Nup, where SW_{\uparrow} changes by $-12.1 W m^{-2}$ (pre: -60.6 , mon: -72.7), an increase of the flux as a consequence of the high albedo of ephemeral monsoonal snow cover (Figure 3e, Table A2). On the other hand, the melt contribution of LW_{\downarrow} increases at all sites (Figure 6), by at least $+15.7$ (Hailuogou, pre: 314.6, mon: 330.3) and up to $+57.0$ (Yala, pre: 248.5, mon: 305.5) (Table A2). Its counterpart LW_{\uparrow} further reduces melt, but to a lesser extent, by -1.0 (Changri Nup, pre: -318.7 , mon: -319.7) to $-13.3 W m^{-2}$ (Langtang, pre: -339.3 , mon: -352.8) (Table A2). As a result LW_{net} plays only a minor role in cooling the glaciers at all sites during monsoon (Figure 5).

4.4.1 Impact of the monsoon on clean-ice sites

We observe opposite changes in M at the two clean-ice glaciers in the transition from pre-monsoon to monsoon: M decreases at Yala by -10.2 (pre: 74.8, mon: 64.6) and increases at Parlung No.4 by $+130.4$ (pre: 32.3, mon: 162.6) (all values in $W m^{-2}$, from Table A2). LW_{net} evolves in similar ways at the two sites and as described in Section 4.4. The difference in M is largely caused by the variability in SW_{net} , which almost entirely controls the melt of the clean-ice glaciers during monsoon. Both H and LE on average remain small energy fluxes at the clean ice sites with highest averages of $LE = -17.6$ at Parlung No.4 and of $H = -13.7$ at Yala (Table A2). Interestingly, LE changes from being an evaporative, melt-reducing energy flux during the pre-monsoon, to a small melt-contributing energy flux ($< 4 W m^{-2}$) at both clean-ice sites (Table A2).

4.4.2 Impact of the monsoon on glaciers with thick debris

Average M remains similar between the pre-monsoon and monsoon at the sites with thick debris cover, as the energy balance components adjust to monsoonal conditions: the changes in M , ranging between -1.0 (Lirung, pre: 37.5, mon: 36.5) and $+2.1$ (24K, pre: 79.5, mon: 81.6), stay below uncertainty levels (all values in $W m^{-2}$, Figure 6a, c, e, g and Table A2). Similar to the other surface types, LW_{net} reduces melt to a lesser degree during the monsoon (Section 4.4). There is a considerable reduction in the melt-contribution of SW_{net} , and the glacier-cooling H becomes a smaller flux by 49.0 (24K, pre: -99.8 , mon: -50.8) up to 68.3 (Lirung, pre: -116.7 , mon: -48.4) (Table A2). The change in LE partly offsets the changes in H , with increases in the flux ranging from -2.1 (24K, pre: -50.6 , mon: -52.7) to 24.4 (Lirung, pre: -16.0 , mon: -40.4) (Figure 6a, c, e and g, and Table A2). Therefore, the changes in the fluxes from the pre-monsoon to monsoon tend to balance each other out (although SW_{\downarrow} decreases, LW_{\downarrow} increases, and although H reduces, LE increases), so that overall melt rates remain similar.




Figure 6. (a)-(g) Differences in energy balance components from pre-monsoon to monsoon at each site including their uncertainties (error bars). The direction of change is to be considered relative to the sign of the original flux (x-axis). Due to the sign convention mentioned in Section 4.3, the presented changes reflect whether the surface receives more energy (positive change) or less energy (negative change). Background indicates the surface type of the site: gray indicates thick debris cover, light blue indicates clean-ice sites, and grey-blue indicates thin debris. (h)-(j) Alternative depiction of the changes from (a)-(f), summarizing surface types; Example Δ -flux numbers in $[W m^{-2}]$ refer to (h) Parlung No.4, (i) Lirung and (j) Hailuogou; Numbers for the remaining glaciers can be looked up in Table A2.

An interruption of the monsoon at 24K occurred in August 2016, possibly associated with an El Niño event (Kumar et al., 2006). During this interruption the energy balance returned to a pre-monsoonal regime (Figure 5e) due to clearer skies, more pronounced diurnal temperature amplitudes, low precipitation rates and lower relative humidity (Figure A7), resulting in higher melt rates during that period (Figure 4e).

4.4.3 Impact of the monsoon a glacier with thin debris

In contrast to the glaciers with thick debris, during the monsoon, the melt energy M increases considerably at Hailuogou Glacier. Although SW_{net} contributes less energy for melt during the monsoon and LW_{net} remains overall small at this site (Figure 5), M increased by 28.7 (pre: 158.1, mon: 186.8) on average (all values in $W m^{-2}$, from Table A2), and mostly during the nights (Figure A11). The increase in melt energy is mostly driven by the turbulent energy fluxes: H increases by 16.6 (pre: 9.1, mon: 25.7) and LE increases by 26.6 (pre: 5.4 mon: 31.6) (Figure 5 and Table A2), with higher increases during the nighttime than during the daytime (Figure A11). While they act to reduce melt at the glaciers with thick debris cover, here the turbulent fluxes drive additional melt during the monsoon.

4.5 Controls on the turbulent fluxes

Our results show the importance of the turbulent fluxes in the energy balance of debris-covered glaciers, their varying role as melt-enhancing or melt-reducing fluxes depending on the debris thickness, and how the monsoon modulates them.

From the regression models (Section 3.1.4), we find that H is largely controlled by the temperature gradient between surface and air (δ_T) on glaciers with thick debris: between 73 (Lirung, pre-monsoon) and 98% (24K, pre-monsoon) of the variability of H are explained by δ_T (Figure 7a), and δ_T decreases during monsoonal conditions by -0.05 (Langtang) to -1.44 $^{\circ}\text{C m}^{-1}$ (Changri Nup) (Table A3). It becomes clear that a smaller temperature gradient between surface and air during the monsoon weakens the melt-reducing effect of H . In contrast, Ws emerges as the most important control of H and LE at the glacier with thin debris, explaining up to 89 and 65% of the variability, respectively (Figure 7a). The mean magnitude of Ws increases at this site from 1.23 in pre-monsoon to 2.1 m s^{-1} in monsoon (Table A3). A cold surface in combination with a wind-enhanced turbulence and fast turnover of warm and moist air masses results in both H and LE becoming powerful drivers of melt on Hailuoguo, the glacier with thin debris cover (Figure 5).

Neither RH , g_T , or Ws on their own, nor their combination explain the variability of LE across sites with thick debris (Figure 7a). LE however increases consistently from pre-monsoon to monsoon together with an increase in the duration of moisture availability at the surface of those glaciers, with increases ranging between 22.3% at 24k and 63.1% at Changri Nup (Table A3). In fact, evaporation and its melt-reducing LE flux tend to be water-limited during the pre-monsoon, but energy-limited during the monsoon (Figure 7b). This implies that the availability of additional moisture drives the increase of LE from pre-monsoon to monsoon.

420




Figure 7. (a) Predictive power of temperature gradient between surface and air (T) and wind speed (Ws) and their combination ('all') for determining H and LE . A multiple polynomial regression model including both variables was used, otherwise a univariate polynomial regression model, and both models had 2 degrees of freedom. **(b)** Budyko-like diagram with the 5-day mean potential evaporation rate during snow-free conditions (E_{pot}) relative to the mean available intercepted water (In) on the x-axis, and the actual evaporation rate during snow-free conditions (E_{act}) relative to In on the y-axis. Only debris-covered glaciers, where LE is a glacier-cooling flux, are shown.

5 Discussion

5.1 Which mass and energy fluxes determine the seasonal mass balance of glaciers in the Central and Eastern Himalaya?

We apply our model in a systematic way to seven glaciers in a variety of environments in the Central and Eastern Himalaya. We
 425 force the model with in-situ station data and constrain and evaluate it against observations of surface height change, lending
 great confidence to the energy flux components. Previous energy balance studies in the region were limited to two (Lejeune
 et al., 2013; Yang et al., 2017; Bonekamp et al., 2019) or three (Zhu et al., 2018) study sites, and partly relied on reanalysis
 products or atmospheric modelling for forcing (Zhu et al., 2018; Bonekamp et al., 2019), without the possibility to evaluate
 the model performance. At all our study sites, ice melt is the largest mass loss component during the ablation season, followed
 430 by snow melt, while sublimation plays only a small role early and late in the season (Section 4.1). Similar to several previous
 studies (Kayastha et al., 1999; Aizen et al., 2002; Yang et al., 2011; Sun et al., 2014), we find that the largest energy source
 for snow and ice melt is SW_{net} (Section 4.2). Thus, major controls on the energy and mass balance of all glaciers are the
 snowcover dynamics (Zhu et al., 2018) and the associated variations in albedo, which in turn are modulated by the timing of
 precipitation and the partition of precipitation into rain and snow (Ding et al., 2017; Bonekamp et al., 2019). For example, in
 435 the case of Parlung No.4, the onset of glacier melt was delayed until well after monsoon onset, until all snow had disappeared
 (Section 4.1). After snow has melted out, ephemeral snowcover from monsoonal precipitation increased surface albedo and
 raised SW_{\uparrow} , protecting the ice and suppressing melt rates throughout the summer (Fujita and Ageta, 2000) (Section 4.1). This

was especially true at the highest sites (Yala, Changri Nup), highlighting the importance of observations of high-elevation surface condition for constraining seasonal glacier mass balance.

440 5.2 How does debris cover modulate the energy balance in comparison to clean-ice surfaces?

Previous energy balance studies on debris-covered glaciers were limited to one or two study sites (e.g. Lejeune et al., 2013; Collier et al., 2014; Rounce et al., 2015; Steiner et al., 2018). Applying the model at five sites with debris cover allows us to identify processes that are likely to be common for a large population of debris-covered glaciers in High Mountain Asia. At the four sites with thick debris, our work confirms that debris protects the ice by returning energy to the atmosphere in the form of
445 turbulent fluxes H and LE in addition to LW_{\uparrow} (Yang et al., 2017) and that the turbulent fluxes can be a major component in the energy balance during both dry and wet conditions (Steiner et al., 2018) (Section 4.3). We also find a melt-enhancing effect of thin debris (Östrem, 1959; Reznichenko et al., 2010; Reid and Brock, 2010) at Hailuoguo Glacier (Section 4.4.3), and that the turbulent fluxes "work against" this glacier (Section 4.5). Our analysis extends beyond most prior representations however by including a water interception storage (Section 3.2.2), which is capable of mimicking the drying process of the debris (Steiner
450 et al., 2018). Representing this process, which was often neglected in previous studies, allows for a more realistic estimation of LE , which is crucial in its role as a glacier-cooling flux at the glaciers with thick debris, and as a control of potential melt enhancement of thin debris (Evatt et al., 2015). Uncertainty remains around the size of the interception storage - for this study it was fixed to 2mm - and investigations on the water interception and holding capacity of debris are needed in order to elucidate this process. Its representation however allows us to extend the short-period results of (Steiner et al., 2018) to
455 multiple sites and across distinct meteorological conditions, emphasizing the importance of turbulent fluxes for debris-covered glacier energy balance. In contrast to the debris-covered glaciers, the turbulent fluxes play a minor role on the clean-ice glaciers (Section 4.4.1).

5.3 How does the monsoon change the glacier surface energy balance?

The ablation period occurs between April and November at all sites, and all sites are affected by the Indian and East Asian
460 summer monsoons during this period (Figures A3 to A9). A long-term average of 65 to 85% of precipitation arrives during the summer months (June-September) at the Central Himalayan sites (Lirung, Lantang, Yala and Changri Nup, Figure A1a-d and Table 2) in contrast to 40 to 56% at the eastern sites (24K, Parlung No.4 and Hailuoguo, Figure A1e-g and Table 2). SW_{\downarrow} is reduced at all glacier surfaces due to the reflection and scattering by persistent, heavy clouds (Figure 8). Overcast conditions caused by monsoon also increase LW_{\downarrow} at all sites (Figure 8). Our analysis shows that some effects of monsoon are common
465 for all surface types, while the presence or absence of debris and its thickness control how the incoming energy is absorbed and transmitted to the ice (Figure 8). We therefore provide a synthesis of the changes based on surface types:

5.3.1 Glaciers with thick debris

Overcast cloud cover, increasing air temperatures and additional moisture modify the energy balance of debris-covered glaciers, to result in a melt-equalizing effect (Section 4.3): warm clouds emit additional amounts of energy towards the glacier in the form of LW_{\downarrow} (Figure 8, Section 4.4). H reduces its cooling effect as a consequence of a smaller average temperature gradient between surface and air (Figure 8, Section 4.5). On the other hand, additional evaporative cooling in the form of LE takes place at the wet debris surface, balancing out the other, melt-enhancing changes (Figure 8, Section 4.3). The debris surface shifts from a water-limited environment during pre-monsoon to an energy-limited process during monsoon (Section 4.5 and Figure 7). We account for the debris water content through the inclusion of a simple interception storage (Section 3.2.2). This allowed us to identify the importance of the glacier-cooling LE coming from the evaporation of liquid water from the debris. Our results are derived from simulations at one location (AWS) on each glacier. To understand how representative our results are of conditions across the glacier ablation zone at each site, and across the possible range of debris thicknesses in particular, we conduct a sensitivity experiment to evaluate the transferability of our results across the glaciers' ablation areas (see detailed explanation in Section A1). This experiment shows that even accounting for the range of conditions across each glacier ablation area, the pattern of pre-monsoon to monsoon difference in flux components, and importantly the equalizing effect on M , remain similar at the glacier scale at all sites with thick debris cover.

5.3.2 Clean-ice glaciers

In contrast to debris-covered glaciers, when clean-ice glaciers are snow-free and the ice has been heated to the melting point, almost all net radiation goes into ice melt, while the turbulent flux contribution remains small (Section 4.4.1). However, when entering the monsoon period, LE tends to switch sign (Figure 8), changing from sublimation/evaporation to condensation, which adds energy to the surface instead of removing it (Section 4.4.1). This behaviour has not been indicated for the drier conditions on the Tibetan Plateau (Mölg et al., 2012; Sun et al., 2014), but has previously been observed at Himalayan sites with a 'southern influence' (Azam et al., 2014; Yang et al., 2017). In contrast to the glaciers with thick debris, the energy balance of clean-ice glaciers is highly sensitive to elevation, as shown in our sensitivity experiment (Section A1)

5.3.3 Glacier with thin debris

At the site with thin debris, we observe a melt-enhancing effect during monsoon conditions. The dark debris surface absorbs almost 90% of SW_{\downarrow} in the case of Hailuoguo (Table A2), and with a short conduction length (1 cm), the energy influx goes almost entirely to melt. Shortwave fluxes reduce during the monsoon, yet melt nonetheless increases, as higher wind speeds enhance turbulence resulting in an increase in H (Section 4.5 and Table A3). Warmer and more humid air increases LE inputs from condensation at the cold surface (Table A3 and Figure A9). Both turbulent fluxes thus become important sources of melt energy (Section 4.4.3). This adds detailed insights to prior observations and modelling inferences that debris around or below the critical thickness causes higher melt rates than at both clean-ice sites and sites with thicker debris cover (Östrem, 1959; Nakawo and Rana, 1999; Reznichenko et al., 2010; Reid and Brock, 2010; Evatt et al., 2015; Fyffe et al., 2020). Artificially




Figure 8. Symbolic representation of changes in energy balance components from pre-monsoon to monsoon. Triangles pointing down/up indicate a positive/negative flux with regards to our sign-convention, where positive/negative means a flux towards/away from the surface. Red/blue indicate an increasing/decreasing value of the flux when moving from pre-monsoon to monsoon. When signs switch, the underlying, empty triangles indicate the pre-monsoonal direction of the flux, while the overlying, colored ones indicate the monsoonal flux

applying thick debris to Hailuogou, while acknowledging the limitations of this experiment (Section A1), results in the same change pattern as the one observed on the other debris-covered glaciers: Melt rates remain almost unchanged when going from pre-monsoon to monsoon (Section A1).

5.4 Implications for Himalayan glaciers in a changing climate

Monsoon-influenced, summer-accumulation glaciers (such as Langtang, Lirung, Yala, and Changri Nup) have been previously shown to be especially vulnerable to warming due to a decrease in accumulation and an enhancement of ablation due to reduced albedo (Fujita and Ageta, 2000), and our results confirm that SW_{net} is the key control on monsoon-period melt rates for clean-ice glaciers (Section 4.4.1). Our results also emphasize that the longevity of pre-monsoon snowcover into the monsoon period is a key control on melt rates (Section 4.1), supporting past findings that the strength and timing of the monsoon onset has a profound impact on small mountain glaciers (Mölg et al., 2014, 2012) through the phase change of precipitation in the transition to monsoon conditions (Fujita and Ageta, 2000; Ding et al., 2017; Zhu et al., 2018). Importantly, our insights into the differential response of glaciers with different surface types to the monsoon and its onset offers keys to interpret their future response under a changing climate:

All future climate scenarios agree on continued warming during the 21st century over High Mountain Asia (Masson-Delmotte, 2021), together with a strengthening of elevation dependent warming (Palazzi et al., 2017) and increases in moisture availability (Masson-Delmotte, 2021). An analysis on the ensemble estimates of regionally downscaled CMIP5 projections (CORDEX) for the Himalayas (Sanjay et al., 2017) shows that total summer precipitation is projected to increase for 2036-2065 (2066-2095)

by 4.4% (10.5%) in the Central Himalaya and by 6.8% (10.4%) in the Eastern Himalaya under RCP4.5 scenarios, relative to 1976-2005. While there is broad model consensus on the increase in future precipitation, there is little consensus on the future variability, frequency and spatial distribution of precipitation across High Mountain Asia (Kadel et al., 2018; Sanjay et al., 2017). A slight shift towards an earlier monsoon onset of <5 days over the coming century together with an increasing shift
520 towards a later retreat by 5 to 10 days (mid-century) and 10 to 15 days (end-century) might increase the length of the monsoon period, with stronger lengthening in the Eastern Himalaya (Moon and Ha, 2020).

The prospect of warmer temperatures together with increased precipitation would (1) cause a shift in the precipitation partition from snow to rain in the monsoon, resulting in snow cover shifting to higher elevations and increasing total melt; (2) potentially lead to an increase in early spring snowfall, which would delay the onset of ice melt; (3) increase the likelihood of ephemeral
525 monsoonal snow cover but move it to higher elevations, thus leaving more of the lower ablation zones exposed; (4) increase the wet-bulb temperature together with humidity to result in a further reduction of the solid fraction of precipitation during monsoon. Overall it is likely that glacier ablation zones will be exposed for longer periods under future climate due to a net decrease of the snow covered duration, with a resulting increase in total ablation. A lengthening of the monsoon into autumn, on the other hand, (Moon and Ha, 2020) would somewhat offset warmer air temperatures with regards to the late-season melt
530 for all glacier types.

The expected warmer and wetter monsoonal conditions, including increased cloudiness, are likely to result in an overall increase of melt rates on clean-ice and glaciers with debris cover around or below the critical thickness. This is because (1) they are more directly controlled by net radiation (comprising all short- and long-wave fluxes), which is likely to increase in magnitude (Section 4.4.1); (2) the turbulent fluxes towards cold surfaces are also likely to increase in magnitude, and they tend
535 to 'work against' these types of glaciers (Section 4.4.1 and 4.4.3). In contrast, the turbulent fluxes 'work for' the glaciers with debris above the critical thickness, and the melt-equalizing effect of debris under monsoon (Section 4.4.2) would likely remain in place. With these components summing up to have an overall protective effect on glaciers with thick debris, they are likely to resist the projected changes in the monsoonal summer longer into the future. Previous studies suggested that the mass balance of debris-covered glaciers might be less sensitive to climate warming than clean-ice glaciers (e.g. Anderson and Mackintosh,
540 2012; Wijngaard et al., 2019; Mattson, 2000). Here we confirm this hypothesis and suggest that this is particularly so under monsoonal conditions. We also suggest that glaciers with debris under the critical thickness might be even more sensitive to future monsoons than clean-ice glaciers.

6 Conclusions

545 We model the energy and mass balance of seven glaciers in the Central and Eastern Himalaya at seven on-glacier weather stations. We find that:

1. At all sites, the largest mass loss component during the ablation season is ice melt, followed by snowmelt and sublimation, while the last only plays a role at our highest sites and outside of the core monsoon. We find that the seasonal

energy and mass balance is strongly controlled by variations of absorbed shortwave radiation, a result of the prevalence of spring snow cover and the occurrence of ephemeral monsoonal snow accumulation.

2. Debris cover above the critical thickness returns most of the energy it absorbs back to the atmosphere via longwave emission and turbulent heat fluxes. While H is primarily controlled by the temperature gradient between surface and air, LE is controlled by the availability of liquid water at the debris surface. When debris is around or under the critical thickness, the melt is more directly radiation-driven. In this case, however, melt is additionally modulated by the turbulent fluxes H and LE , for which wind speed is the primary control, and the cold surface favours condensation rather than evaporation.

3. The response of the glacier mass and energy balance to the monsoon depends on the surface type: melt-rates increase compared to the pre-monsoon at the clean-ice glaciers and the glaciers with thin debris cover, while they stay similar at the glaciers with thick debris cover. We attribute these differences to the role the turbulent fluxes play for each surface type. At the glaciers with thick debris cover, where the turbulent fluxes 'work for' the glacier, evaporation of the additionally available moisture (LE) provides extra cooling during the monsoon. The evaporation of liquid water is a moisture limited process during the pre-monsoon and turns into an energy-limited process during the monsoon. The cooling induced by H at the same time decreases, with the result of unchanged available melt-energy M during monsoon. In a sensitivity experiment, we confirm that these results are representative of the entire ablation zones of the thickly debris-covered glaciers. At the clean-ice sites, in contrast, the melt is mostly radiation controlled throughout the ablation season and varies greatly over the elevation profile of the ablation zone. The turbulent fluxes play a subordinate role there, but can switch from melt-reducing to melt-enhancing in the seasonal transition into the monsoon. At the thin debris-covered site, on the other hand, the turbulent fluxes always 'work against' the glacier and intensify during the monsoon.

Given these findings, under projected future monsoonal conditions, namely warmer and possibly longer and wetter monsoons (Sanjay et al., 2017; Moon and Ha, 2020; Masson-Delmotte, 2021), we expect the mass balance of glaciers with thick debris-cover to react less sensitively than clean-ice glaciers and glaciers with thin debris cover.

Code and data availability. Model and analysis codes as well as AWS datasets are available upon request during the discussion and review phase and will be made publicly available at a later stage.

575 *Author contributions.* SFu, FP and EM designed the study. SFu carried out the analysis with the help of CF, MM and SFa. SFu interpreted the results, created the figures and wrote the manuscript with the help of CF, EM, MM, TS and FP. SFa, PW, WI, and QL reviewed the manuscript. WY and BD facilitated field data collection and provided parameterisations for albedo and precipitation phase. WY, PW and WI also contributed data sets.

Competing interests. The authors declare that they have no conflict of interest.

580 *Acknowledgements.* This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 772751), RAVEN, "Rapid mass losses of debris-covered glaciers in High Mountain Asia". We would like to thank Jakob Steiner and ICIMOD for hosting and contributing datasets. We also thank Marin Kneib for organizing, and the Langtang and 24K field teams for helping with data collection, as well as the HRE team for their great support.

Appendix A: Climatic and meteorological conditions

585 Average mean monthly 2 m air temperatures have a similar pattern at all study sites (Figure A1a), with a slow increase from January to a peak between July and August, just after peak monsoon, and a steeper decline from post-monsoon into winter. Incoming shortwave radiation (Figure A1b) shows a clear peak before monsoon onset at all sites. A smaller secondary peak is reached just after the monsoon in October at the Central Himalayan sites, but not at the Eastern Himalayan sites. Interruptions in monsoonal overcast conditions (break periods) seem to be more common at the eastern sites, leading to occasional secondary
590 peaks in incoming shortwave radiation during monsoon. LW_{\downarrow} follows a similar regime as Ta , with highest values reached during the core monsoon (Figure A1c). The yearly cycle of wind speeds (Figure A1d) varies considerably between sites. Common characteristics for most sites (except for Changri Nup) are that wind speeds are highest around December/January and that monsoonal wind speeds are generally higher than during the shoulder seasons. There is a clear difference in the seasonal evolution of precipitation between the Central (Lirung, Lantang, Yala, Changri Nup) and the Eastern Himalayan sites (24K, 595 Parlung No.4, Hailuogou) (Figure A1e): relatively high mean monthly precipitation during the monsoon period is contrasted by comparably low precipitation outside of this period. The eastern sites have less pronounced monsoonal precipitation peaks, and more gradual changes in precipitation intensities over the annual cycle. The Parlung sites (24K and Parlung No.4) have two precipitation peaks: during spring and monsoon. Hailuogou exhibits the smoothest evolution over the annual cycle with a clear maximum in July. A simple monsoon index (MI) is calculated for each year including the study year as the ratio between
600 monsoon precipitation and annual average precipitation (Figure A1e). This value tends to be higher in the Central Himalaya compared to the sites on the South-Eastern Tibetan Plateau.

At each site, we define the onset and recession date of monsoon based on visual inspection of the AWS records, observing the seasonal shift of air temperature, relative humidity, and longwave and shortwave radiation. For this, we define the date after which clear shifts are observable in the variables' regime (Figures A3 to A9).





Figure A1. Monthly climatology derived from ERA5-Land for 1981-2019 (grey background lines), along with the monthly averages (black lines) and the study year at each glacier (colored lines). Plotted meteorological variables are **(a)** mean air temperature (Ta), **(b)** incoming shortwave radiation (SW_{\downarrow}), **(c)** incoming longwave radiation (LW_{\downarrow}), **(d)** wind speed (Ws) and **(e)** monthly precipitation sums (Pr). Black vertical lines indicate the average region-wide monsoon season. Boxplots show the monsoon index (MI) over ERA5-Land period as the fraction of monsoonal (June-September) to annual precipitation, with the colored dot indicating the value for the study year.



E

Figure A2. All energy balance components of all glaciers in comparison, split into pre-monsoon and monsoon; black bars indicate the uncertainty range;




Figure A3. Meteorological observations on Lirung during the ablation season recorded by AWS; Red vertical lines indicate monsoon onset and end; cyan indicates time steps with snow cover at the AWS location, as determined from α (>0.5)




Figure A4. Meteorological observations on Langtang during the ablation season recorded by AWS; Red vertical lines indicate monsoon onset and end; cyan indicates time steps with snow cover at the AWS location, as determined from α (>0.5)




Figure A5. Meteorological observations on Yala during the ablation season recorded by AWS; Red vertical lines indicate monsoon onset and end; cyan indicates time steps with snow cover at the AWS location, as determined from α (>0.5)




Figure A6. Meteorological observations on Changri Nup during the ablation season recorded by AWS; Red vertical lines indicate monsoon onset and end; cyan indicates time steps with snow cover at the AWS location, as determined from α (>0.5)




Figure A7. Meteorological observations on 24K during the ablation season recorded by AWS; Red vertical lines indicate monsoon onset and end; cyan indicates time steps with snow cover at the AWS location, as determined from α (>0.5)




Figure A8. Meteorological observations on Parlung No.4 during the ablation season recorded by AWS; Red vertical lines indicate monsoon onset and end; cyan indicates time steps with snow cover at the AWS location, as determined from α (>0.5)




Figure A9. Meteorological observations on Hailuogou during the ablation season recorded by AWS; Red vertical lines indicate monsoon onset and end; cyan indicates time steps with snow cover at the AWS location, as determined from α (>0.5)




Figure A11. Energy flux differences in the diurnal cycle (stacked) between pre-monsoon and monsoon; The direction of change is to be considered relative to the sign of the original flux. Positive and negative sign corresponds to energy added or removed from the glacier, respectively; Grey background indicates debris-covered site, light blue indicates clean ice sites and gray-blue indicates 1cm debris site




Figure A12. Changes in the individual fluxes when moving from premonsoon to monsoon. Color dots indicate 'standard' runs with AWS site specific conditions. Black bars indicate the uncertainty range on the standard runs. Grey indicates the sensitivity of flux changes (Δ -range) to debris thickness varied from 10 and 80 cm combined with the elevation range of debris-covered ablation zone of the individual glaciers; Positive and negative sign corresponds to energy added or removed from the glacier, respectively;

Our results are derived from simulations at one location (AWS) on each glacier. To understand how representative they are of conditions across the glacier ablation zone at each site, and across the possible range of debris thicknesses in particular, we conduct a sensitivity experiment at each site. We assume that the strongest changes in meteorological forcing with elevation would be the T_a , which in turns controls the precipitation partition and the albedo. We re-run the model synthetically varying the AWS elevation to represent the range of elevation of each glacier ablation zone by applying a T_a lapse rate of $0.6^{\circ}C/100m$ and, for the debris-covered sites, by varying also the debris thickness in the range 10-80 cm (for ranges and steps see Table A1). Using the station-measured, accumulated albedo is not appropriate during this experiment, due to changing snow conditions with varying elevation. We therefore include the parameterisation introduced by Ding et al. (2017) for modelling α . From the resulting range of EB flux outputs, we calculate the range of expected changes for the entire ablation zone when moving from pre-monsoon to monsoon (Δ -range). This allows us to place our results in the context of the changes that can be expected over the entire ablation zone, given its elevation span and debris thickness variability. Figure A12 shows that even accounting for the range of conditions across each glacier ablation area, the pattern of pre-monsoon to monsoon difference in flux components, and importantly M , remain similar for debris-covered sites: The Δ -range of M stays within the uncertainty range, with the exception of Langtang, where the unrealistic combination of relatively thin debris and low elevation causes high M Δ -range. This lends confidence to the results obtained at the individual AWS locations. Although we adjusted forcing data for elevation in this exercise, we could not represent the effects of variable debris thicknesses in modifying $2m$ meteorological variables (Steiner and Pellicciotti, 2016; Shaw et al., 2016). This comes with the assumption that surface-atmosphere interactions are negligible compared to the altitudinal patterns and temporal changes. While this might be acceptable at thicker debris sites, it is more questionable at Hailuogou, where the observations were taken above thin and cold debris. However, also at this site, the Δ -range ends up to be small ($5 W m^{-2}$) and close to zero when debris between 10 and 80 cm thickness is applied artificially.

Table A1. Ranges of elevations and debris thicknesses used for the sensitivity runs, including the glacier terminus elevation (min), the AWS elevation (AWS) and the upper debris limit on debris-covered glaciers or to the approximated ELA elevation on clean-ice glaciers (max). We also show the range of debris thicknesses h_m modelled for debris-covered glacier sites. All combinations of elevations and debris thicknesses were used.

Glacier		Lirung	Langtang	Yala	Changri Nup	24K	Parlung No.4	Hailuogou
min	[m.asl]	3990	4500	5170	5270	3910	4620	2980
AWS	[m.asl]	4076	4557	5350	5471	3900	4800	3550
max	[m.asl]	4400	5600	5400	5600	4200	5400	3700
h_d	[cm]	10, 20, 30, 40, 50, 60, 70, 80						

Table A2. Mean energy balance components at each site and for pre-monsoon (pre), monsoon (mon), and post-monsoon (post) periods, as well as flux magnitude changes from pre-monsoon to monsoon, and monsoon to post-monsoon. All values are in $W\ m^{-2}$.

<i>fluxes in W/m2</i>	SW↓			SW↑			LW↓			LW↑			LE		
	pre	mon	post	pre	mon	post	pre	mon	post	pre	mon	post	pre	mon	post
LIR															
mean	277.1	170.0	224.1	-39.8	-21.1	-41.3	293.3	341.2	264.0	-358.1	-364.8	-340.4	-16.0	-40.4	-1.9
Δ		-107.1	54.1		18.7	-20.2		47.9	-77.2		-6.7	24.4		-24.4	38.5
LAN															
mean	295.6	208.1	262.4	-55.9	-30.1	-48.6	292.3	334.7	238.6	-339.3	-352.8	-326.1	-26.4	-49.9	-5.8
Δ		-87.5	54.3		25.7	-18.4		42.3	-96.0		-13.5	26.7		-23.5	44.1
YAL															
mean	307.7	172.7	271.8	-169.4	-94.4	-195.0	248.5	305.5	212.3	-299.8	-311.6	-287.1	-13.7	0.1	-12.2
Δ		-135.0	99.1		75.0	-100.6		57.0	-93.2		-11.7	24.4		13.7	-12.3
CNU															
mean	237.1	154.7	258.1	-60.6	-72.7	-126.0	268.8	310.2	196.4	-318.7	-319.7	-300.4	-15.7	-18.1	-3.1
Δ		-82.4	103.4		-12.1	-53.3		41.5	-113.8		-1.0	19.3		-2.5	15.0
24K															
mean	296.6	219.7	140.9	-18.5	-13.0	-8.3	324.6	349.3	336.6	-369.9	-371.3	-351.0	-50.6	-52.7	-40.0
Δ		-76.9	-78.8		5.4	4.7		24.7	-12.7		-1.3	20.3		-2.1	12.7
NO4															
mean	308.5	209.1	197.3	-219.6	-54.8	-81.5	267.5	308.2	261.5	-310.3	-313.4	-310.5	-17.6	3.6	-18.5
Δ		-99.5	-11.8		164.8	-26.7		40.7	-46.7		-3.1	2.9		21.2	-22.1
HAI															
mean	178.2	136.4	105.8	-25.2	-10.8	-28.4	314.6	330.3	273.1	-324.5	-327.9	-309.1	5.4	31.6	-4.8
Δ		-41.8	-30.6		14.4	-17.6		15.7	-57.2		-3.4	18.8		26.2	-36.4

<i>fluxes in W/m2</i>	H			Qv			G			dG			M		
	pre	mon	post	pre	mon	post	pre	mon	post	pre	mon	post	pre	mon	post
LIR															
mean	-116.7	-48.4	-86.5	0.0	0.0	0.0	-33.7	-36.4	-18.7	5.6	0.0	16.6	37.5	36.5	19.0
Δ		68.3	38.0		0.0	0.0		-2.7	-17.8		-5.6	-16.6		-1.0	17.5
LAN															
mean	-126.7	-77.4	-111.4	0.0	-0.3	0.0	-22.5	-26.2	-7.8	2.2	0.0	27.3	26.9	28.7	10.4
Δ		49.3	34.1		-0.3	-0.3		-3.7	-18.4		-2.2	-27.3		1.8	18.4
YAL															
mean	-2.3	0.5	1.4	0.0	-0.2	-4.1	4.7	0.9	0.0	4.7	0.9	0.0	74.8	64.6	0.3
Δ		2.8	-0.9		-0.2	3.9		-3.8	0.9		-3.8	0.9		-10.2	64.3
CNU															
mean	-67.4	-8.7	-36.1	0.0	0.0	0.0	-39.2	-29.1	10.4	17.1	1.8	10.6	34.6	34.1	0.2
Δ		58.7	27.4		0.0	0.0		10.1	-39.5		-15.3	-8.9		-0.5	33.9
24K															
mean	-99.8	-50.8	-24.4	-0.3	0.2	0.6	-81.8	-81.4	-53.1	0.3	0.0	0.0	79.5	81.6	53.7
Δ		49.0	-26.4		0.6	-0.4		0.4	-28.4		-0.3	0.0		2.1	27.9
NO4															
mean	4.7	12.3	5.8	0.2	0.2	0.0	0.2	0.1	0.1	0.2	0.1	0.0	32.3	162.6	54.1
Δ		7.6	6.6		0.0	0.2		-0.1	0.1		-0.1	0.1		130.4	108.5
HAI															
mean	9.1	25.7	-9.9	1.2	1.9	0.1	-150.4	-186.4	-16.5	1.8	0.5	17.6	158.1	186.8	36.9
Δ		16.6	35.5		0.7	1.8		-35.9	-169.9		-1.4	-17.2		28.7	149.9

Table A3. Mean cloud-cover fraction (ccf), relative humidity (RH), temperature gradient between surface and air (δ_T), wind speed (Ws) and the percentage of time during which the debris is modelled to hold intercepted water (In) for each site and season, also indicating percent changes between the sub-seasons.

		ccf [–]			RH [%]			δ_T [$^{\circ}C\,m^{-1}$]			Ws [$m\,s^{-1}$]			In [%]		
		<i>pre</i>	<i>mon</i>	<i>post</i>	<i>pre</i>	<i>mon</i>	<i>post</i>	<i>pre</i>	<i>mon</i>	<i>post</i>	<i>pre</i>	<i>mon</i>	<i>post</i>	<i>pre</i>	<i>mon</i>	<i>post</i>
LIR	mean	0.76	0.91	0.74	68.1	90.8	67.1	1.19	0.78	0.89	0.47	0.27	0.52	40.1	74.0	26.4
	Δ		0.14	-0.16		22.6	-23.6		-0.40	0.10		-0.19	0.24		33.9	-47.6
LAN	mean	0.76	0.86	0.62	80.7	96.9	56.3	1.02	0.97	0.95	1.79	1.10	1.27	38.8	75.3	9.5
	Δ		0.10	-0.24		16.2	-40.6		-0.05	-0.02		-0.68	0.17		36.5	-65.8
YAL	mean	0.65	0.87	0.55	69.8	93.0	39.4	-0.36	-0.96	-0.89	1.74	1.00	1.68	-	-	-
	Δ		0.21	-0.32		23.2	-53.6		-0.59	0.07		-0.74	0.67		-	-
CNU	mean	0.78	0.89	0.58	71.2	89.2	39.3	1.69	0.25	0.89	1.88	1.09	2.48	16.2	79.3	5.3
	Δ		0.11	-0.31		18.1	-50.0		-1.44	0.64		-0.79	1.39		63.1	-74.0
24K	mean	0.70	0.82	0.91	73.1	80.6	81.2	1.86	0.73	0.18	1.33	1.56	1.35	56.8	79.1	84.3
	Δ		0.12	0.08		7.4	0.7		-1.13	-0.55		0.22	-0.21		22.3	5.2
NO4	mean	0.72	0.87	0.79	65.7	81.3	73.1	-0.78	-2.01	-0.62	2.96	2.67	3.23	-	-	-
	Δ		0.14	-0.07		15.7	-8.3		-1.22	1.39		-0.28	0.56		-	-
HAI	mean	0.88	0.93	0.91	81.3	92.3	90.6	-2.08	-2.61	0.38	1.23	2.15	0.93	99.8	100.0	75.6
	Δ		0.05	-0.01		10.9	-1.6		-0.53	2.99		0.91	-1.22		0.2	-24.4

References

- Aizen, V., Aizen, E., and Nikitin, S.: Glacier regime on the northern slope of the Himalaya (Xixibangma glaciers), *Quaternary International*, 97, 27–39, 2002.
- Anderson, B. and Mackintosh, A.: Controls on mass balance sensitivity of maritime glaciers in the Southern Alps, New Zealand: the role of debris cover, *Journal of Geophysical Research: Earth Surface*, 117, 2012.
- Azam, M., Wagnon, P., Vincent, C., Ramanathan, A., Favier, V., Mandal, A., and Pottakkal, J.: Processes governing the mass balance of Chhota Shigri Glacier (western Himalaya, India) assessed by point-scale surface energy balance measurements, *Cryosphere*, 8, 2195–2217, 2014.
- Bélair, S., Crevier, L.-P., Mailhot, J., Bilodeau, B., and Delage, Y.: Operational implementation of the ISBA land surface scheme in the Canadian regional weather forecast model. Part I: Warm season results, *Journal of hydrometeorology*, 4, 352–370, 2003.
- Bonekamp, P. N., de Kok, R. J., Collier, E., and Immerzeel, W. W.: Contrasting meteorological drivers of the glacier mass balance between the Karakoram and central Himalaya, *Frontiers in Earth Science*, 7, 107, 2019.
- Bookhagen, B. and Burbank, D. W.: Toward a complete Himalayan hydrological budget: Spatiotemporal distribution of snowmelt and rainfall and their impact on river discharge, *Journal of Geophysical Research: Earth Surface*, 115, 2010.
- Botter, M., Zeeman, M., Burlando, P., and Fatichi, S.: Impacts of fertilization on grassland productivity and water quality across the European Alps: insights from a mechanistic model, *Biogeosciences Discussions*, pp. 1–35, 2020.
- Brock, B. W., Willis, I. C., and Sharp, M. J.: Measurement and parameterization of albedo variations at Haut Glacier d’Arolla, Switzerland, *Journal of Glaciology*, 46, 675–688, 2000.
- Brun, F., Berthier, E., Wagnon, P., Kääb, A., and Treichler, D.: A spatially resolved estimate of High Mountain Asia glacier mass balances from 2000 to 2016, *Nature geoscience*, 10, 668–673, 2017.
- Collier, E. and Immerzeel, W. W.: High-resolution modeling of atmospheric dynamics in the Nepalese Himalaya, *Journal of Geophysical Research: Atmospheres*, 120, 9882–9896, 2015.
- Collier, E., Nicholson, L., Brock, B., Maussion, F., Essery, R., and Bush, A.: Representing moisture fluxes and phase changes in glacier debris cover using a reservoir approach, *The Cryosphere*, 8, 1429–1444, 2014.
- Collier, E., Maussion, F., Nicholson, L., Mölg, T., Immerzeel, W., and Bush, A.: Impact of debris cover on glacier ablation and atmosphere-glacier feedbacks in the Karakoram, 2015.
- Cuffey, K. M. and Paterson, W. S. B.: *The physics of glaciers*, Academic Press, 2010.
- Ding, B., Yang, K., Qin, J., Wang, L., Chen, Y., and He, X.: The dependence of precipitation types on surface elevation and meteorological conditions and its parameterization, *Journal of hydrology*, 513, 154–163, 2014.
- Ding, B., Yang, K., Yang, W., He, X., Chen, Y., Guo, X., Wang, L., Wu, H., and Yao, T.: Development of a Water and Enthalpy Budget-based Glacier mass balance Model (WEB-GM) and its preliminary validation, *Water Resources Research*, 53, 3146–3178, 2017.
- Evatt, G. W., Abrahams, I. D., Heil, M., Mayer, C., Kingslake, J., Mitchell, S. L., Fowler, A. C., and Clark, C. D.: Glacial melt under a porous debris layer, *Journal of Glaciology*, 61, 825–836, 2015.
- Farinotti, D., Huss, M., Fürst, J. J., Landmann, J., Machguth, H., Maussion, F., and Pandit, A.: A consensus estimate for the ice thickness distribution of all glaciers on Earth, *Nature Geoscience*, 12, 168–173, 2019.
- Farinotti, D., Immerzeel, W. W., de Kok, R. J., Quincey, D. J., and Dehecq, A.: Manifestations and mechanisms of the Karakoram glacier Anomaly, *Nature Geoscience*, 13, 8–16, 2020.

- Fatichi, S., Ivanov, V. Y., and Caporali, E.: A mechanistic ecohydrological model to investigate complex interactions in cold and warm water-controlled environments: 1. Theoretical framework and plot-scale analysis, *Journal of Advances in Modeling Earth Systems*, 4, 665 2012.
- Fujita, K Sakai, A. et al.: Modelling runoff from a Himalayan debris-covered glacier, *Hydrol. Earth Syst. Sci*, 18, 2679–2694, 2014.
- Fujita, K. and Ageta, Y.: Effect of summer accumulation on glacier mass balance on the Tibetan Plateau revealed by mass-balance model, *Journal of Glaciology*, 46, 244–252, 2000.
- Fujita, K., Takeuchi, N., and Seko, K.: Glaciological observations of Yala Glacier in Langtang Valley, Nepal Himalayas, 1994 and, *Bulletin of Glacier Research*, 16, 75–8, 1998.
- Fyffe, C. L., Woodget, A. S., Kirkbride, M. P., Deline, P., Westoby, M. J., and Brock, B. W.: Processes at the margins of supraglacial debris cover: quantifying dirty ice ablation and debris redistribution, *Earth Surface Processes and Landforms*, 2020.
- Gardelle, J., Berthier, E., and Arnaud, Y.: Slight mass gain of Karakoram glaciers in the early twenty-first century, *Nature geoscience*, 5, 322–325, 2012.
- 675 Giese, A., Boone, A., Wagnon, P., and Hawley, R.: Incorporating moisture content in surface energy balance modeling of a debris-covered glacier, *The Cryosphere*, 14, 1555–1577, 2020.
- Herreid, S. and Pellicciotti, F.: The state of rock debris covering Earth’s glaciers, *Nature Geoscience*, pp. 1–7, 2020.
- Heynen, M., Miles, E., Ragettli, S., Buri, P., Immerzeel, W. W., and Pellicciotti, F.: Air temperature variability in a high-elevation Himalayan catchment, *Annals of Glaciology*, 57, 212–222, 2016.
- 680 ICIMOD: AWS Yala Glacier [Data set]. ICIMOD., 2021.
- Immerzeel, W. W., Van Beek, L., Konz, M., Shrestha, A., and Bierkens, M.: Hydrological response to climate change in a glacierized catchment in the Himalayas, *Climatic change*, 110, 721–736, 2012.
- Immerzeel, W. W., Kraaijenbrink, P. D., Shea, J., Shrestha, A. B., Pellicciotti, F., Bierkens, M. F., and de Jong, S. M.: High-resolution monitoring of Himalayan glacier dynamics using unmanned aerial vehicles, *Remote Sensing of Environment*, 150, 93–103, 2014.
- 685 Kadel, I., Yamazaki, T., Iwasaki, T., and Abdillahi, M. R.: Projection of future monsoon precipitation over the central Himalayas by CMIP5 models under warming scenarios, *Climate Research*, 75, 1–21, 2018.
- Kaser, G., Großhauser, M., and Marzeion, B.: Contribution potential of glaciers to water availability in different climate regimes, *Proceedings of the National Academy of Sciences*, 107, 20 223–20 227, 2010.
- Kayastha, R. B., Ohata, T., and Ageta, Y.: Application of a mass-balance model to a Himalayan glacier, *Journal of Glaciology*, 45, 559–567, 690 1999.
- Kraaijenbrink, P., Bierkens, M., Lutz, A., and Immerzeel, W.: Impact of a global temperature rise of 1.5 degrees Celsius on Asia’s glaciers, *Nature*, 549, 257–260, 2017.
- Kumar, K. K., Rajagopalan, B., Hoerling, M., Bates, G., and Cane, M.: Unraveling the mystery of Indian monsoon failure during El Niño, *Science*, 314, 115–119, 2006.
- 695 Lejeune, Y., Bertrand, J.-M., Wagnon, P., and Morin, S.: A physically based model of the year-round surface energy and mass balance of debris-covered glaciers, *Journal of Glaciology*, 59, 327–344, 2013.
- Liao, H., Liu, Q., Zhong, Y., and Lu, X.: Landsat-based estimation of the glacier surface temperature of Hailuoguo glacier, Southeastern Tibetan Plateau, between 1990 and 2018, *Remote Sensing*, 12, 2105, 2020.
- Mascart, P., Noilhan, J., and Giordani, H.: A modified parameterization of flux-profile relationships in the surface layer using different roughness length values for heat and momentum, *Boundary-Layer Meteorology*, 72, 331–344, 1995.
- 700

- Masson-Delmotte, V., P. Z. A. P. S. L. C. C. P. S. B. N. C. Y. C. L. G. M. I. G. M. H. K. L. E. L. J. B. R. M. T. K. M. T. W. O. Y. R. Y. B. Z.: IPCC, 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press. In Press., 2021.
- Mastrotheodoros, T., Pappas, C., Molnar, P., Burlando, P., Manoli, G., Parajka, J., Rigon, R., Szeles, B., Bottazzi, M., Hadjidoukas, P., et al.: More green and less blue water in the Alps during warmer summers, *Nature Climate Change*, 10, 155–161, 2020.
- Mattson, L. E.: The influence of a debris cover on the mid-summer discharge of Dome Glacier, Canadian Rocky Mountains, *IAHS PUBLICATION*, pp. 25–34, 2000.
- Maussion, F., Scherer, D., Mölg, T., Collier, E., Curio, J., and Finkelburg, R.: Precipitation seasonality and variability over the Tibetan Plateau as resolved by the High Asia Reanalysis, *Journal of Climate*, 27, 1910–1927, 2014.
- 710 McCarthy, M. J.: Quantifying supraglacial debris thickness at local to regional scales, Ph.D. thesis, Scott Polar Research Institute and British Antarctic Survey, 2018.
- Miles, E. S., Steiner, J. F., and Brun, F.: Highly variable aerodynamic roughness length (z_0) for a hummocky debris-covered glacier, *Journal of Geophysical Research: Atmospheres*, 122, 8447–8466, 2017.
- Mölg, T., Maussion, F., Yang, W., and Scherer, D.: The footprint of Asian monsoon dynamics in the mass and energy balance of a Tibetan glacier, *The Cryosphere*, 6, 1445, 2012.
- 715 Mölg, T., Maussion, F., and Scherer, D.: Mid-latitude westerlies as a driver of glacier variability in monsoonal High Asia, *Nature Climate Change*, 4, 68–73, 2014.
- Moon, S. and Ha, K.-J.: Future changes in monsoon duration and precipitation using CMIP6, *npj Climate and Atmospheric Science*, 3, 1–7, 2020.
- 720 Muñoz Sabater, J.: ERA5-Land monthly averaged data from 1981 to present .Copernicus Climate Change Service (C3S) Climate Data Store (CDS). (Accessed on 01-10-2020), 2019.
- Nakawo, M. and Rana, B.: Estimate of ablation rate of glacier ice under a supraglacial debris layer, *Geografiska Annaler: Series A, Physical Geography*, 81, 695–701, 1999.
- Nicholson, L. and Benn, D. I.: Calculating ice melt beneath a debris layer using meteorological data, *Journal of Glaciology*, 52, 463–470, 2006.
- 725 Noilhan, J. and Mahfouf, J.-F.: The ISBA land surface parameterisation scheme, *Global and planetary Change*, 13, 145–159, 1996.
- Östrem, G.: Ice melting under a thin layer of moraine, and the existence of ice cores in moraine ridges, *Geografiska Annaler*, 41, 228–230, 1959.
- Palazzi, E., Filippi, L., and von Hardenberg, J.: Insights into elevation-dependent warming in the Tibetan Plateau-Himalayas from CMIP5 model simulations, *Climate Dynamics*, 48, 3991–4008, 2017.
- 730 Paschalis, A., Fatichi, S., Pappas, C., and Or, D.: Covariation of vegetation and climate constrains present and future T/ET variability, *Environmental Research Letters*, 13, 104012, 2018.
- Pratap, B., Dobhal, D., Mehta, M., and Bhambri, R.: Influence of debris cover and altitude on glacier surface melting: a case study on Dokriani Glacier, central Himalaya, India, *Annals of Glaciology*, 56, 9–16, 2015.
- 735 Quincey, D., Smith, M., Rounce, D., Ross, A., King, O., and Watson, C.: Evaluating morphological estimates of the aerodynamic roughness of debris covered glacier ice, *Earth Surface Processes and Landforms*, 42, 2541–2553, 2017.

- Ragettli, S., Pellicciotti, F., Immerzeel, W. W., Miles, E. S., Petersen, L., Heynen, M., Shea, J. M., Stumm, D., Joshi, S., and Shrestha, A.: Unraveling the hydrology of a Himalayan catchment through integration of high resolution in situ data and remote sensing with an advanced simulation model, *Advances in Water Resources*, 78, 94–111, 2015.
- 740 Reid, T. D. and Brock, B. W.: An energy-balance model for debris-covered glaciers including heat conduction through the debris layer, *Journal of Glaciology*, 56, 903–916, 2010.
- Reznichenko, N., Davies, T., Shulmeister, J., and McSaveney, M.: Effects of debris on ice-surface melting rates: an experimental study, *Journal of Glaciology*, 56, 384–394, 2010.
- Rounce, D., Quincey, D., and McKinney, D.: Debris-covered glacier energy balance model for Imja-Lhotse Shar Glacier in the Everest region
745 of Nepal, 2015.
- Rowan, A. V., Nicholson, L. I., Quincey, D. J., Gibson, M. J., Irvine-Fynn, T. D., Watson, C. S., Wagnon, P., Rounce, D. R., Thompson, S. S., Porter, P. R., et al.: Seasonally stable temperature gradients through supraglacial debris in the Everest region of Nepal, Central Himalaya, *Journal of Glaciology*, pp. 1–12, 2020.
- Sakai, A. and Fujita, K.: Contrasting glacier responses to recent climate change in high-mountain Asia, *Scientific reports*, 7, 1–8, 2017.
- 750 Sakai, A., Fujita, K., and Kubota, J.: Evaporation and percolation effect on melting at debris-covered Lirung Glacier, Nepal Himalayas, 1996, *Bulletin of glaciological research*, 21, 9–16, 2004.
- Sanjay, J., Krishnan, R., Shrestha, A. B., Rajbhandari, R., and Ren, G.-Y.: Downscaled climate change projections for the Hindu Kush Himalayan region using CORDEX South Asia regional climate models, *Advances in Climate Change Research*, 8, 185–198, 2017.
- Scherler, D., Bookhagen, B., and Strecker, M. R.: Hillslope-glacier coupling: The interplay of topography and glacial dynamics in High Asia,
755 *Journal of Geophysical Research: Earth Surface*, 116, 2011a.
- Scherler, D., Bookhagen, B., and Strecker, M. R.: Spatially variable response of Himalayan glaciers to climate change affected by debris cover, *Nature geoscience*, 4, 156–159, 2011b.
- Shah, S. S., Banerjee, A., Nainwal, H. C., and Shankar, R.: Estimation of the total sub-debris ablation from point-scale ablation data on a debris-covered glacier, *Journal of Glaciology*, 65, 759–769, 2019.
- 760 Shaw, T. E., Brock, B. W., Fyffe, C. L., Pellicciotti, F., Rutter, N., and Diotri, F.: Air temperature distribution and energy-balance modelling of a debris-covered glacier, *Journal of Glaciology*, 62, 185–198, 2016.
- Shean, D., Bhushan, S., Montesano, P., Rounce, D., Arendt, A., and Osmanoglu, B.: A systematic, regional assessment of High Mountain Asia Glacier mass balance. *Front, Earth Sci*, 7, 363, 2020.
- Shrestha, R., Kayastha, R., and Kayastha, R.: Effect of debris on seasonal ice melt (2016–2018) on Ponkar Glacier, Manang, Nepal. *Sciences in Cold and Arid Regions*, 12 (5): 261–271. DOI: 10.3724/SP. J. 1226.2020. 00261, Reju Shrestha et al, pp. 0261–0271, 2020.
- 765 Steiner, J. F. and Pellicciotti, F.: Variability of air temperature over a debris-covered glacier in the Nepalese Himalaya, *Annals of Glaciology*, 57, 295–307, 2016.
- Steiner, J. F., Litt, M., Stigter, E. E., Shea, J., Bierkens, M. F., and Immerzeel, W. W.: The importance of turbulent fluxes in the surface energy balance of a debris-covered glacier in the Himalayas, *Frontiers in Earth Science*, 6, 144, 2018.
- 770 Stumm, D., Joshi, S. P., Gurung, T. R., and Silwal, G.: Mass balances of Yala and Rikha Samba Glacier, Nepal from 2000 to 2017, *Earth System Science Data Discussions*, pp. 1–37, 2020.
- Sun, W., Qin, X., Du, W., Liu, W., Liu, Y., Zhang, T., Xu, Y., Zhao, Q., Wu, J., and Ren, J.: Ablation modeling and surface energy budget in the ablation zone of Laohugou glacier No. 12, western Qilian mountains, China, *Annals of glaciology*, 55, 111–120, 2014.

- van den Broeke, M., van As, D., Reijmer, C., and van de Wal, R.: Assessing and improving the quality of unattended radiation observations in Antarctica, *Journal of Atmospheric and Oceanic Technology*, 21, 1417–1431, 2004.
- 775 Versegny, D. L.: CLASS—A Canadian land surface scheme for GCMs. I. Soil model, *International Journal of Climatology*, 11, 111–133, 1991.
- Vincent, C., Wagnon, P., Shea, J., Immerzeel, W., Kraaijenbrink, P., Shrestha, D., Soruco, A., Arnaud, Y., Brun, F., Berthier, E., et al.: Reduced melt on debris-covered glaciers: investigations from Changri Nup Glacier, Nepal, 2016.
- 780 Wagnon, P.: IACS working group on debris-covered glaciers. AWS North Changri Nup Glacier [Data set]. Hosted on glacioclim.osug.fr, 2021.
- Wijngaard, R. R., Steiner, J. F., Kraaijenbrink, P. D., Klug, C., Adhikari, S., Banerjee, A., Pellicciotti, F., Van Beek, L. P., Bierkens, M. F., Lutz, A. F., et al.: Modeling the Response of the Langtang Glacier and the Hintereisferner to a Changing Climate Since the Little Ice Age, *Frontiers in Earth Science*, 7, 143, 2019.
- 785 Yang, W., Guo, X., Yao, T., Yang, K., Zhao, L., Li, S., and Zhu, M.: Summertime surface energy budget and ablation modeling in the ablation zone of a maritime Tibetan glacier, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
- Yang, W., Yao, T., Zhu, M., and Wang, Y.: Comparison of the meteorology and surface energy fluxes of debris-free and debris-covered glaciers in the southeastern Tibetan Plateau, *Journal of Glaciology*, 63, 1090–1104, 2017.
- Yao, T., Thompson, L., Yang, W., Yu, W., Gao, Y., Guo, X., Yang, X., Duan, K., Zhao, H., Xu, B., et al.: Different glacier status with atmospheric circulations in Tibetan Plateau and surroundings, *Nature climate change*, 2, 663–667, 2012.
- 790 Zhang, Y., Fujita, K., Liu, S., Liu, Q., and Nuimura, T.: Distribution of debris thickness and its effect on ice melt at Hailuoguo glacier, southeastern Tibetan Plateau, using in situ surveys and ASTER imagery, *Journal of Glaciology*, 57, 1147–1157, 2011.
- Zhu, M., Yao, T., Yang, W., Xu, B., Wu, G., and Wang, X.: Differences in mass balance behavior for three glaciers from different climatic regions on the Tibetan Plateau, *Climate Dynamics*, 50, 3457–3484, 2018.