

The stability of present-day Antarctic grounding lines — Part A: No indication of marine ice sheet instability in the current geometry

Emily A. Hill¹★, Benoît Urruty²★, Ronja Reese^{1,3}★, Julius Garbe^{3,4}, Olivier Gagliardini², Gaël Durand², Fabien Gillet-Chaulet², G. Hilmar Gudmundsson¹, Ricarda Winkelmann^{3,4}, Mondher Chekki², David Chandler⁵, and Petra M. Langebroek⁵

¹Department of Geography and Environmental Sciences, Northumbria University, Newcastle upon Tyne, UK

²Univ. Grenoble Alpes, CNRS, IRD, Grenoble INP, IGE, 38000 Grenoble, France

³Potsdam Institute for Climate Impact Research (PIK), Member of the Leibniz Association, Potsdam, Germany

⁴Institute of Physics and Astronomy, University of Potsdam, Potsdam, Germany

⁵NORCE Norwegian Research Centre, Bjerknes Centre for Climate Research, Bergen, Norway

★These authors contributed equally to this work.

Correspondence: Olivier Gagliardini (olivier.gagliardini@univ-grenoble-alpes.fr)

Abstract. Theoretical and numerical work has ~~firmly established that~~ shown that under certain circumstances grounding lines of marine-type ice sheets can enter phases of irreversible advance and retreat driven by the marine ice sheet instability (MISI). Instances of such irreversible retreat have been found in several simulations of the ~~past and future evolution of the~~ Antarctic Ice Sheet. However, it has not been assessed whether the Antarctic grounding lines are already undergoing MISI in their *current*
5 *position*. Here, we conduct a systematic numerical stability analysis ~~of all the grounding lines of the Antarctic Ice Sheet to determine if they are currently undergoing irreversible retreat through MISI. To do this, we initialise using~~ three state-of-the-art ~~ice-flow-ice-sheet~~ models, Úa, Elmer/Ice, and PISM, ~~to replicate the current geometry of the Antarctic Ice Sheet, and then apply small, but numerically significant, perturbations in ocean-induced ice-shelf melt. We find.~~ Our results show that the grounding lines around Antarctica migrate slightly away from their initial position while the perturbation is applied, and then revert to the
10 initial state once the perturbation is removed. ~~There is no indication of irreversible or self-sustaining retreat. This suggests that if the currently observed mass imbalance (external climate forcing) were to be removed, the grounding-line retreat would likely stop. However~~ This indicates that present-day retreat of Antarctic grounding lines is not irreversible or self-sustained. However, our accompanying paper (Part B, Reese et al., 2022) shows that if the grounding-lines retreat further inland, under present-day climate forcing, further grounding-line retreat is expected, and our accompanying paper (Part B, Reese et al., 2022) shows that
15 ~~this could eventually lead to a~~ it may lead to the eventual irreversible collapse of some marine regions of West Antarctica.

1 Introduction

Retreat of the Antarctic grounding lines, i.e. the zones where the grounded ice sheet becomes so thin that it floats, could destabilise large marine regions of the ice sheet (Weertman, 1974; Schoof, 2007; Mengel and Levermann, 2014; Feldmann and Levermann, 2015), thereby committing several metres of global sea-level rise over the coming centuries to millennia (e.g.,
20 DeConto and Pollard, 2016; Gollledge et al., 2015; Ritz et al., 2015; Cornford et al., 2015). Indeed, the potential for widespread

destabilisation and rapid ice discharge is one of the greatest uncertainties in future projections of ice sheet mass loss (Robel et al., 2019; Pattyn and Morlighem, 2020; IPCC, 2021). ~~Previous studies have made some suggestions that present-day retreat in regions of the West Antarctic Ice Sheet could mean that irreversible retreat has begun. However, these studies do not provide compelling evidence, and to date there has not yet been a systematic analysis to assess whether irreversible retreat of Antarctic grounding lines is already underway.~~

Marine ice sheet instability (MISI) is the proposed mechanism by which grounding lines are considered to undergo self-sustained, irreversible retreat ~~if~~ (Weertman, 1974; Schoof, 2007, 2012). The existence of MISI means that a shift in the position of the grounding line can cause it to cross a critical threshold (or ‘tipping point’), beyond which the MISI mechanism drives the system towards a different steady state. The resulting retreat is considered irreversible, because once initiated, reversing the perturbation to pre-threshold conditions is not sufficient to halt or reverse it. Instead, the forcing has to be reversed past its initial value to recover the initial state (Rosier et al., 2021). For marine, laterally uniform ice sheets with constant conditions, it has been shown that ice flux across the grounding line increases with thickness. Hence, retreat on a retrograde (inland) sloping bed into deeper water, and thus regions of greater ice thickness, promotes a positive feedback in which retreat continues, unabated, inland. ~~This self-sustaining mechanism was theoretically shown to determine the stability regime of grounding lines of marine, laterally uniform ice sheets.~~ In this case, no stable steady-state grounding lines exist on a retrograde sloping bed (Weertman, 1974; Schoof, 2007, 2012). However, in the case of laterally confined ice shelves that buttress the inland grounded ice, the MISI mechanism becomes more complex. Indeed, in the presence of buttressing ice shelves, stable steady-state grounding-line positions can exist on a retrograde bed slope (Gudmundsson et al., 2012; Pegler, 2018; Haseloff and Sergienko, 2018). Most ice shelves around Antarctica provide such buttressing, and have an important impact on inland ice dynamics (Fürst et al., 2016; Reese et al., 2018b). In addition to the ice-shelf lateral confinement, non-negligible bed topography found, for instance, under Thwaites Glacier, and very weak beds, such as under Siple Coast ice streams complicate stability conditions (Sergienko and Wingham, 2019, 2022). ~~In other words, while a retrograde sloping bed is a necessary conditions for MISI to occur, it is not a sufficient condition.~~ Crucially, this means that whether Antarctic grounding lines are undergoing irreversible retreat cannot be concluded from observed retreat and bed-slope direction ~~alone~~. Targeted ice sheet model simulations are required to answer this question.

~~The existence of MISI means that a shift in the position of the grounding line can cause it to cross a critical threshold (or ‘tipping point’), beyond which the MISI mechanism drives the system towards a different steady state. The resulting retreat is considered irreversible, because once initiated, reversing the perturbation to pre-threshold conditions is not sufficient to halt or reverse the retreat. Instead, the forcing has to be reversed past its initial value to recover the initial state. As an example term~~

MISI is often used to describe the potential for a self-reinforcing, positive feedback to be at play, i.e. the retreat of the Antarctic grounding lines is internally driven (e.g., Pattyn and Morlighem, 2020). In particular, MISI is often discussed in relation to the susceptibility of the West Antarctic Ice Sheet to collapse due to this positive feedback mechanism. Previous numerical modelling studies have shown the potential for tipping points related to MISI in the Antarctic Ice Sheet. For example Rosier et al. (2021) showed that three tipping points exists for the Pine Island Glacier. Furthermore, in their simulations of the whole Antarctic Ice Sheet, Garbe et al. (2020) ~~find~~ found that retreat of West Antarctic grounding lines could be initiated by around

1 – 2 °C of global warming above pre-industrial, while the recovery of these grounding lines to their modern positions requires temperatures that are at least –1 °C below the pre-industrial average.

Large parts of the Antarctic Ice Sheet have been identified as potentially susceptible to MISI due to their deep inland sloping topography satisfying the necessary condition for MISI (see Fig. S60 in). This encompasses in particular the marine regions of the West Antarctic Ice Sheet. Recent work on Pine Island Glacier has shown the existence of three tipping points; two smaller tipping points and a third, larger one, that could eventually induce the collapse. Previous studies have argued that sections of the West Antarctic Ice Sheet . The Amundsen-Sea Embayment (ASE) sector in West Antarctica is of particular concern as glaciers in this region have been accelerating and thinning, and the grounding lines are retreating .

Parts of the East Antarctic Ice Sheet also fulfill the necessary condition for MISI, retrograde sloping beds, in their deep marine basins, e.g. Robin subglacial basin upstream of the Filchner-Ronne ice shelf (basin 1 in Fig. 2), Recovery subglacial basin feeding the Filchner Ice Shelf (basin 2), Wilkes subglacial basin and Aurora subglacial basin upstream of Totten Ice Shelf in East Antarctica may already be undergoing self-sustained unstable retreat (Joughin et al., 2014; Favier et al., 2014; Rignot et al., 2014). However, with the exception of Totten Glacier , these regions have not yet shown substantial acceleration, thinning or retreat of the grounding line.

The aim of this paper is to determine if the currently observed grounding line retreat in any Antarctic region to date there has not been a systematic analysis to determine whether the currently observed changes in grounding line position are reversible. In this paper we use a systematic modelling approach to assess whether the current retreat of the Antarctic grounding lines is due to MISI, and therefore irreversible. To do this we perform a numerical analysis an ongoing positive feedback mechanism related to MISI. Our modelling approach is outlined at the beginning of Sect. 2. Briefly, we perform numerical experiments using three state-of-the-art ice sheet models, Elmer/Ice (Gagliardini et al., 2013), Úa (Gudmundsson, 2020) , and the Parallel Ice Sheet Model (PISM; Bueler and Brown, 2009; Winkelmann et al., 2011). We initialise Elmer/Ice and Úa to be in steady state and closely replicate the present-day geometry of the ice sheet and then perform a numerical stability analysis. To assess whether this steady state ice sheet geometry is stable, we can apply a small-amplitude perturbation to a control variable that satisfies the steady state condition, in this case the position of the grounding line. We perturb the grounding line position indirectly, by increasing the sub-shelf melt rates. This perturbation is designed to be large enough that it is numerically significant, but small enough that the state of the system is not fundamentally altered. Applying a small perturbation to a steady state solution is a common mathematical stability approach and has been previously used in numerical studies to assess the stability of grounding lines in idealised set-ups . If the steady state is stable, the system will evolve back to its initial state after the perturbation is removed. Conversely, if the steady state is unstable, a small perturbation will be enhanced and the system evolves away from its initial steady state.

Owing to the (long-term) response of ice sheets to varying climate forcing, by applying a small but numerically significant perturbation to our initial model states, that all closely replicate the current geometry and velocity of the Antarctic Ice Sheet is not in a steady state today. Indeed, observations show that mass loss and grounding line retreat are driven by external climate forcing. However, in order to assess whether MISI is playing a role in the current retreat of the Antarctic grounding lines, we need to go beyond observations, and apply a systematic modelling approach. Using the approach outlined above we can

determine whether the currently observed changes in grounding-line position are reversible or have already begun self-sustained collapse. If we can find stable steady states in the current geometry of the ice sheet in our numerical models, then we can conclude that the present-day location of Antarctic grounding lines are not inherently unstable, i.e., the positive feedback related to MISI is not automatically at play. Vice-versa, the existence of such stable steady states is rather a strong indication
95 that find the currently observed grounding lines to either revert back to their former position (if the state is steady), or stay within the vicinity (if drifting through time), then this would indicate that current retreat of Antarctic grounding lines is in response to either past, or present-day external climate forcing, without MISI being at play.

To corroborate our analysis, we repeat these perturbation experiments using a transient ice-sheet state in PISM that has been forced using ocean and atmospheric conditions from 1850 to 2015. Using this state we can additionally test whether,
100 with transient forcing included, the current grounding lines show self-sustained retreat in response to a small-amplitude perturbation. If, after removing the perturbation, the grounding lines either reverse towards their initial position or the same position obtained in a control simulation we can additionally conclude that MISI is not taking place. Note that in this manuscript for PISM simulations that replicate the transient behaviour of the ice sheet, we alternatively refer to grounding lines as ‘irreversible’ if they are engaged in MISI-driven retreat and ‘reversible’ if MISI is not at play. unlikely to be due to an ongoing
105 positive feedback mechanism, i.e. related to MISI.

While the approach outlined above answers the question of whether present-day grounding lines are or are not engaged in an ongoing MISI, the problem of their stability could also be approached from another angle. In particular, the question arises if as to whether the currently observed retreat driven by current climate conditions could commit the grounding lines to eventually undergo irreversible retreat. This is specifically addressed in our accompanying paper (Part B, Reese et al., 2022),
110 where long-term model simulations are used to assess whether present-day climate forcing has the potential to eventually lead to a collapse of major marine basins.

The paper is structured as follows: In the following section (Sect. 2.1-2) we present the common datasets and approaches used in our model initialisation. We then present details of the ice sheet model set-ups in Sect. ??, and in Sect. ?? we present the initial model states used in initialisation of our three ice sheet models and the perturbation experiments described in Sect. 2.4.
115 The results are presented for the entire ice sheet and individual drainage basins in Sect. 3, and discussed further in Sect. 4.

2 Methods

Here, we perform numerical simulations using three. To determine the stability regime of the Antarctic Ice Sheet in its current geometry we use three different ice sheet models, to explore the stability of the current positions of Antarctic grounding lines. A summary and two complementary methodologies. The ice sheet models used are Elmer/Ice, Úa and PISM. Modelling experiments were conducted independently by three modelling teams and each team tailored the details of the methodologies to the needs of their own models. The two methodologies applied are i) stability analysis of steady states, and ii) trajectory analysis of transient states.
120

125 In the *steady-state stability* analysis, we construct a steady Antarctic Ice Sheet state as close as possible to the currently observed ice sheet geometry and surface velocities. To determine whether this steady state of the ice sheet is stable or unstable, a small amplitude perturbation was applied. If the ice sheet reverted back to the steady state after the perturbation was removed, it was considered to be stable with respect to the perturbation applied. This experimental procedure is motivated by the definition of asymptotic stability, i.e. a steady state solution f_e is stable if there exists a $\delta > 0$ such that if $\|f_e - f(t)\| < \delta$, then $f(t) \rightarrow f_e$ when $t \rightarrow +\infty$. This definition requires the construction of a steady state, i.e. f_e , to which the perturbation is applied, and an analysis of the convergence, or non-convergence, over time of the perturbed state towards that steady state.

130 In the *trajectory* analysis, which is motivated by an orbital stability analysis, we perturbed a given state (steady or not) and analysed if the trajectory of the perturbed solution stayed within a given distance from the trajectory of the unperturbed solution. Formally, for the state $f(t_0)$ and the resulting perturbed state $g(t_0)$ for which $\|f(t_0) - g(t_0)\| = \delta_0$, we determined if $\|f(t) - g(t)\| \leq \delta_0$ for $t > t_0$. Hence, we observe if the grounding line of the perturbed state stays with time at the same, or smaller, initial distance from the grounding line of the unperturbed state. Note that here we are only interested in assessing the local stability of the system around its current geometry, and hence only observe the evolution of the state for a limited period after the applied perturbation.

140 An overview of our entire methodology is shown in Fig. 1. ~~We first initialise all three models using as many common aspects of our models as possible.~~ The *steady-state stability* experiments were performed using Elmer/Ice and Úa and can be summarised as follows: 1) The models are initialized using an inversion procedure to replicate the observed present-day geometry and ice velocities (Sect. 2.1) ~~to create initial states that replicate the observed geometry of the Antarctic Ice Sheet (see 2.2.1),~~ 2) A modification is made to the mass balance term, such that if held constant in time, the present-day geometry is as close as numerically possible to a steady state (Sect. ??). ~~Secondly, we use these initial states to assess the sensitivity of current grounding-line positions to perturbations in ice shelf buttressing 2.2.2),~~ 3) to test whether this state is stable or unstable the model states are perturbed (by increasing sub-shelf melt) for a short period of time to cause a small, but numerically significant, shift in the position of the grounding line (Sect. 2.4). ~~All three ice sheet models are initialised using slightly different procedures, the details of which are outlined in,~~ 4) the perturbation is reversed and we use the temporal evolution of the flux and grounding line positions to determine whether the constructed steady state is asymptotically stable (Sect. 2.2.1 and 2.3).

150 To ensure that our conclusions are not conditioned on the steady-state assumptions made in our first set of numerical experiments, we conducted a series of *trajectory* analysis experiments where the perturbation is applied to drifting grounding lines. In this case the model states are not in steady-state. In PISM we initialise the ice sheet-model as follows: 1) the model is initialised to pre-industrial conditions using a spin-up procedure, 2) the model is run using historical forcing from ISMIP6 to obtain a transient state consistent with the observed present-day trend in mass loss (see Sect. 2.3 for further details). We then perturb the model states and determine if the grounding lines of the perturbed and unperturbed transient model states remain close to each other, or if their trajectories start to diverge over time. As we are here interested in the stability of the grounding lines in their current position, we only follow the grounding-line evolution for a limited period of time. It is possible that if we had followed the grounding line evolution for extended periods of time the trajectories of the perturbed and unperturbed states

would have started to diverge. However, that would no longer be a statement about the local stability regime of the grounding lines in the current ice sheet geometry.




Figure 1. Overview of experimental setup. Schematic summarizing the experimental setup used in the stability analysis of the present-day Antarctic grounding lines. The setup comprises the different model initialisation procedures (grey box) as well as the two different types of prognostic model experiments (blue box): the numerical stability analysis presented in the current manuscript (Part A; orange box) and the committed retreat analysis presented in Reese et al. (2022) (Part B; yellow box). The three different model initialisation procedures yield three comparable initial states (“common initial state”; green box), from which all experiments are started. See text for more details.

160 2.1 Common Approach

To minimise the differences between the ice sheet initial states, we apply as many common features as possible in the model setups (Table 1), either in terms of model physics, input datasets or parameterisations. In principle, any remaining differences between the ice sheet reference states will be solely due to their individual initialisation procedures and implementation of model physics. A comparison of all three models is given in Table 1.

165 PISM is a finite difference model capable of simulating large ensembles on multi-millennial time-scales, whereas Elmer/Ice and Úa are finite element models with unstructured grids and are designed for decadal to century scale simulations. All models use ice density $\rho_i = 917$, sea water density $\rho_w = 1027$, and gravity $g = 9.81$. To use the most physics-based friction law all models impose a pressure-dependent friction law, which usually leads to more physically representative sliding close

to the grounding line as compared to the Weertman sliding law. The exact formulations of each law used are given in Appendices A1-A3.

To initialise the models to present day conditions, we require a number of observational datasets. We use We first initialise all three models using as many common aspects of our models as possible to create initial states based on the observed geometry of the Antarctic Ice Sheet. These common inputs and approaches are summarised in Table 1. Namely we use observations of bedrock topography, ice surface elevation, and ice thickness from BedMachine Antarctica ~~Version 2~~ (Version 2; Morlighem et al., 2020). To replicate the current ice flow, we take surface velocities from a recent snapshot in time (2015/16) from the MEaSUREs Annual Ice Velocity Maps ~~Version 1 dataset~~ dataset (Version 1; Mouginot et al., 2019), which has a resolution of 1 km and good coverage across the entire ice sheet. ~~In the case of Úa and Elmer/Ice these surface velocities are used directly in the inversion algorithm.~~

Across the surface of the ice sheet, all models initially apply a constant-in-time surface mass balance (later modified for Elmer/Ice and Úa), which is the output from the regional atmospheric climate model RACMO2.3p2 averaged from 01/1995 to 12/2014 (van Wessem et al., 2018). Melting at the bottom of ice shelves is a key control on the dynamics of the Antarctic Ice Sheet, and is the focus of the perturbation experiments. All three models parameterise sub-shelf melt rates using their respective implementations of the Potsdam Ice shelf Cavity mOdel (PICO; Reese et al., 2018a). We ensure that the PICO geometry is the same as that in Reese et al. (2018a). ~~While PICO is not a perfect representation of present-day melt rates, it can track the grounding line movement and provides melting rates for newly ungrounded regions.~~ PICO parameters were selected to reflect the sensitivity of sub-shelf melt rates in the ~~ASE~~ Amundsen Sea Embayment (ASE) sector and Filchner-Ronne Ice Shelf to ocean temperature changes. Further details on this tuning of PICO parameters can be found in Reese et al. (2022). This results in the parameter for vertical heat exchange $\gamma_T^* = 5 \times 10^{-5} \text{ m s}^{-1}$ and the parameter for the overturning strength $C = 2 \text{ Sv m}^3 \text{ kg}^{-1}$. ~~Temperature corrections were applied to individual PICO basins summarising the far-field ocean conditions such that~~ While PICO is not a perfect representation of present-day melt rates ~~are obtained for present-day ocean forcing from , averaged over the time period 1975 to 2012.~~, it can track the grounding line movement and provides melting rates for newly ungrounded regions.

	Elmer/Ice	PISM Úa	PISM
Numerical method	Finite element	Finite difference <u>element</u>	Finite <u>element</u> difference
Stress Balance <u>balance</u>	SSA	SSA +SIA	SSA <u>+SIA</u>
Grid Resolution <u>resolution</u>	Unstructured grid 1–50 km	8 km Unstructured grid 1– 200 km	<u>8 km</u>
Rheology	Glen’s flow law	Glen–Paterson–Budd–Liboutry–Glen <u>Paterson–Budd–Liboutry–Duval</u> flow law Glen’s flow law	<u>flow law</u>
Friction <u>Sliding</u> law	Regularized Coulomb (Joughin et al., 2019)	<u>Regularized Coulomb</u> (Asay-Davis et al., 2016)	Power-law with Mohr– Coulomb (Schoof and Hind- marsh, 2010; Bueler and van Pelt, 2015) Regularized Coulomb
Initialisation method	Data assimilation with relax- ation	Spin-up Data assimilation with relaxation	<u>Spin-up</u>

Table 1. Comparison of the models and the physics of the models which are detailed further in Sect. 2.1 and Appendices A1–A3. SSA: Shallow-shelf approximation, SIA: Shallow-ice approximation.

2.2 ~~Model Initialisation~~ Steady states in Elmer/Ice and Úa

2.2.1 ~~Elmer/Ice and Úa~~

195 ~~In this section, due to their similar initialisation procedures (see Fig. 1), we present the common approach used by~~ Elmer/Ice
and Úa ~~to create initial model states for the Antarctic Ice Sheet. Specific details related to each individual model are presented~~
~~in Appendices A1 and A2. These models have both~~ are both finite element models that solve the vertically integrated ice
dynamics equations using the Shallow Shelf Approximation (SSA; MacAyeal, 1989). Both models have been used extensively
to solve ice flow problems in Antarctica, and have participated in a number of model intercomparison experiments, e.g. Pattyn
200 et al. (2012); Cornford et al. (2020). ~~Both ice sheet codes are finite element models that solve the vertically integrated ice~~
~~dynamics equations using the Shallow Shelf Approximation. An~~ models first create an unstructured finite element mesh ~~was~~
~~created for each model~~, where element sizes were refined in fast flowing regions. ~~In both cases, mesh resolution is 1 km~~
, and close to the grounding line. ~~Observations of ice sheet topography were then linearly interpolated onto these meshes.~~
~~Throughout the forward in time experiments, the extent of the ice shelves remains unchanged; however, both models impose a~~
205 ~~minimum ice shelf thickness, which is sufficiently thin to represent ice that has been removed and provides no buttressing. The~~

~~grounding-line position is calculated using the flotation criterion~~ The initial ice sheet model states were then obtained using a two-step approach outlined in the following sections.

~~Here, we use a two step approach to initialise Elmer/Ice and Úa. First, we used a data assimilation approach in which we perform~~

210 2.2.1 Model inversions

Using present-day observations of ice sheet geometry and velocity, the models are first initialised using a model inversion to estimate ~~the basal friction and viscosity sliding-law and rheology~~ parameters using the adjoint method (MacAyeal, 1993), ~~to replicate the observed present-day ice sheet velocities.~~ Both models impose a pressure-dependent sliding law, which usually leads to more physically representative sliding close to the grounding line as compared to the Weertman sliding law (Bronde
215 et al., 2019). The optimal fields for ~~these~~ the basal slipperiness and viscosity parameters are found by minimising a cost function which is the sum of misfit and regularisation terms. The main misfit term is the difference between observed and modelled velocities. Both models also apply an additional penalty on the rates of thickness change, to reduce nonphysical ice flux divergence anomalies. We regularise the inverse solutions using Tikhonov regularisation terms that enforce smoothness of the inferred parameters (~~friction and viscosity~~ basal slipperiness and ice rate factor). The regularisation weights are determined
220 using an L -curve analysis. By design, at the end of the inversion the surface velocities are in very good agreement with observations. See Appendices ~~A1 and~~ A1 and A2 for details on the inversion.

~~Second, to be able to assess the stability of current grounding lines with respect to small-amplitude perturbations, we need to obtain an ice sheet initial state that is~~

2.2.2 Mass balance modification

225 Observations clearly show that the present-day geometry of the ice sheet is not in steady state. ~~We define steady state as a state for which ice volume, i.e. ice thickness changes through time are as close as possible to zero. While the data assimilation approach is able to replicate present-day ice velocities and grounding-line positions, once we run the models forward in time, we quickly drift away from these initial conditions, due to inconsistencies in the input datasets and uncertainties in model parameters. Similarly to and, and to overcome the challenges of model drift under control conditions, we apply a correction~~
230 ~~to the~~ $dh/dt \neq 0$, and the current mass balance is $dh/dt = \dot{m}_{pd} - \nabla \cdot (\mathbf{v}h)$, where \mathbf{v} is the ice velocity and \dot{m}_{pd} is the total present-day mass balance (surface + base). However, given that stability is strictly speaking a property of steady states, in order to test the stability of the grounding lines in their current geometry, we require a steady-state configuration of the ice sheet. Ideally, this steady state would be achieved through the inversion step, however, the penalty we apply to rates of thickness change is not sufficient to bring the model into a steady state, and when the model is run forward in time it drifts away
235 from the present-day geometry and ice velocity. Hence, we create a modified mass balance term (\dot{m}) in the form ~~This seeks to bring rates of volume change \dot{m}_{mod} such that $dh/dt = 0$, or as close as numerically possible to zero (steady state), by subtracting.~~ To do this, we follow a similar approach to others (Price et al., 2011; Goelzer et al., 2013) in which we subtract the modelled rates of ice sheet thickness change (dh/dt) thickness change $dh/dt|_{relax}$ needed to keep the model in balance :

There are subtle differences as to how \dot{m} is calculated in each model. Elmer/Ice begins with a 20-year relaxation period with
 240 $\dot{m} = \dot{b}_{\text{RACMO}} - \dot{b}_{\text{PICO}}$, where \dot{m}_{pd} as follows:

$$\dot{m}_{\text{mod}} = \dot{m}_{\text{pd}} - \left. \frac{dh}{dt} \right|_{\text{relax}}, \quad \text{where} \quad \dot{m}_{\text{pd}} = \dot{b}_{\text{RACMO}} + \dot{b}_{\text{PICO}}. \quad (1)$$

Thereby, the present-day surface mass balance field \dot{b}_{RACMO} denotes is the 1995–2014 averaged surface mass balance provided by RACMO2.3p2 (van Wessem et al., 2018) and \dot{b}_{PICO} the sub-shelf melt rates provided by PICO using 1975–2012 averaged ocean forcing from Schmidtke et al. (2014)(see Appendix A1). In a second step, rates of thickness change dh/dt
 245 were subtracted from the RACMO surface mass balance field and sub-shelf melt rates from PICO (Eq. 1). This \dot{m} field was then put back into. By construction, the modification term $dh/dt|_{\text{relax}}$ reduces the mass balance to $\nabla \cdot (vh) = \dot{m}_{\text{mod}}$ with $dh/dt \approx 0$. It is calculated in slightly different ways for the two models, details of which are in the respective Appendices A1 and A2. In summary, Elmer/Ice for a further 1-year relaxation period calculates $dh/dt|_{\text{relax}}$ after a 20-year relaxation period and then inputs this into a 1-year forward relaxation, which forms the beginning of the perturbation experiments. Úa uses a
 250 semi-iterative approach detailed in Appendix A2. This takes $dh/dt|_{\text{inv}}$ which takes $dh/dt|_{\text{inv}}$ calculated after the inversion as input to Eq. (1) for a 1-year relaxation period, t_{relax} . The \dot{m} . The \dot{m}_{mod} field is then recalculated using $dh/dt|_{t_{\text{relax}}}$ $dh/dt|_{\text{relax}}$ at 1 year. In both models \dot{m} is held constant for all remaining simulations. The correction terms applied to the mass balance $dh/dt|_{t_{\text{relax}}}$ in each model are shown in Supplementary Fig. ??, and this is the beginning of the perturbation experiments. The modified mass balance fields and terms in Eq. (1) are shown in Figs. S1 to S3.

We find that applying this modified mass balance term over the entire ice sheet (both grounded and floating portions) brings
 255 the ice sheet models to a steady state. However, by creating our modified mass balance field, we have created a surface mass balance field that deviates from climate conditions in RACMO. Broadly speaking, these fields represent the spatial pattern across the ice sheet; both models show higher positive mass balance around the coastal margins, and more precipitation around the West Antarctic coast of the ASE and Bellingshausen Sea regions (Figs. S1 and S2). Additionally, the $dh/dt|_{\text{relax}}$ fields
 260 are regionally similar, where in both models the modification needed to obtain a steady state are largest in West Antarctica, in particular in the ASE sector. This reflects the current flux imbalance in West Antarctica and ongoing retreat and mass loss from this region. While the spatial patterns are broadly the same, e.g. large corrections across the AMSE modifications across the ASE, we note that the distribution and amplitude of the mass balance correction fields vary modification to the mass balance varies greatly between the two models, likely due to subtle differences in our model initialisation. We interpret the differences
 265 between these two correction terms to show that our results are not an artefact of this field having a specific shape.

We find that applying this corrected mass balance term over the entire ice sheet (both grounded and floating portions) brings
 the ice sheet models to a steady state. As aforementioned, the Antarctic Ice Sheet is not in a steady state today. However, in
 order to determine the stability regime of the present-day grounding lines in the context of their present-day geometry, we
 require a steady state. It In Úa the modification is higher in magnitude and locally spatially heterogeneous, i.e. both positive
 270 and negative values occur in close proximity to one another (see Fig. S3). By comparison, the modification in Elmer/Ice is smoother and generally smaller in magnitude. Crucially, it is assumed that using this modifying the mass-balance correction approach in this manner has not altered the location of the critical thresholds in the grounding-line position for the real ice

sheet, and that perturbing these steady states can allow us to ~~conclude on~~ learn about the stability of the present-day grounding lines in their current position. Indeed, we interpret differences between our two modified mass balance fields, and that our results are consistent for both models (and PISM that does not modify the mass balance; Sect. 2.3), to show that our results are not an artefact of this field having a specific shape. We also conducted a series of numerical experiments where we allowed the grounding lines to drift instead of forcing them to be stationary. This was done by either applying limited or no additional modification to the surface mass balance (Figure S4). In all cases we found that once the perturbation was removed, grounding lines started to drift at the same rate as the (unperturbed) reference run. For unstable states one would expect the trajectories to diverge with time.

2.2.3 PISM

~~In contrast to Úa and Elmer~~

2.2.3 Initial states

~~The initial ice-sheet model states created by Elmer/Ice, PISM simulations use a finite difference scheme to solve the momentum balance on a regular grid of 8 km horizontal resolution. Ice flow velocities are obtained from a superposition of the SSA and the shallow ice approximation (SIA) velocity fields. PISM is thermo-mechanically coupled and solves the enthalpy evolution on a three-dimensional grid. The ice rheology / rate factor is calculated from the ice enthalpy. A pseudo-plastic power-law relationship relates the basal shear stress to the SSA basal sliding velocities. Thereby, the Mohr-Coulomb criterion is used to compute the basal yield stress from parameterised till material properties (the till friction angle, a heuristic, piece-wise linear function of bedrock topography following the assumption that subglacial material with a marine history should be weaker) and the effective pressure on the saturated till. The amount of water in the till is determined by Ice and Úa closely replicate the observed ice thickness and surface velocities, particularly in the location of fast flowing ice streams (see Figs. S5 and S6). Importantly, the positions of the grounding lines in our models closely replicates the currently observed position of the enthalpy model and a constant till water decay rate. For consistency with the other models, glacial isostatic adjustment is not considered and we only calve floating ice that extends beyond the present-day extent of Antarctic ice shelves. While report that a horizontal resolution of 2 km is required to accurately represent grounding lines (Fig. 2) from BedMachine (Morlighem et al., 2020). In addition, the good agreement of observed and modelled grounding-line dynamics, find that by using a subgrid interpolation of friction, positions for all models can be seen in our individual glacier profile figures presented in the results (Fig. 4). We calculate the error in the grounded ice area (a proxy for grounding-line reversibility in PISM is also captured at coarser ($\Delta x > 10$ km) resolutions. While a higher horizontal resolution would be wishful, we here employ this interpolation to be able to run PISM over millennial time scales. We find that PISM results are in line with results from Elmer/change), by differencing the simulated and observed grounded ice areas. To obtain a relative displacement of the grounding line itself, we normalise this area change by the simulated length of the grounding line. The resulting grounding-line position errors are 84 m for Elmer/Ice and Úa that employ finer resolution around the grounding lines Ice, 540 m for Úa.~~




Figure 2. Modelled and observed present-day Antarctic grounding-line locations. Present-day Antarctic bed topography (BedMachine; Morlighem et al., 2020) showing regions where the bed topography is below (blue shading) and above (brown shading) sea level in meters above sea level (m a.s.l.). Modelled initial grounding-line positions are shown as colored lines, observed grounding-line position is shown in black. Ice shelves are indicated by grey shading. Golden lines denote the locations of the transects shown in Fig. 4. Grey lines mark the boundaries of the IMBIE basins (Zwally et al., 2012). The inset shows a zoom into the Amundsen Sea Embayment sector of West Antarctica.

305 During a steady-state control simulation for both models there is less than 4.2 mm of change in sea-level equivalent volume over 100 years. Furthermore, there is very little drift in the position of the grounding line from the initial location, deviating by only 12.6 (Elmer/Ice) and 2.1 m (Úa) over 100 years, which is very small with respect to grid resolution (1 km close to the grounding line).

2.3 Transient state in PISM

310 To support the findings of this study further, we generate an initial state using PISM. Due to the spin-up procedure of PISM, when it arrives at a steady state, by definition that state will be stable. Hence, we would not need to additionally perturb such a state to determine if it is stable or unstable. Instead, we conduct the *trajectory* analysis described above whereby we perturb a

state including a present-day trend in mass changes. This allows us to examine if current grounding-line positions stay within the vicinity of the unperturbed control run. More details on the physics modelled in PISM and a comparison to the other models is given in Appendix A3.

2.3.1 Spin-up to pre-industrial conditions and historical run

The strategy adopted to build an initial state with PISM relies on spin-up methods, with the approach taken here detailed in Reese et al. (2022) and summarised in Fig. 1: ~~we~~ We first create a thermal equilibrium with ~~constant~~ fixed geometry, followed by an ensemble of equilibrium simulations with full dynamics run ~~for~~ under constant \sim 1850 climate conditions ~~;~~ starting from ~~these~~ (here equilibrium is defined as the integrated ice sheet mass balance being close to zero).

~~Starting from the 1850 initial configurations,~~ historic simulations are run from 1850 to 2015. We use ISMIP6 historic forcing for the atmosphere and the ocean from 1850 to 2015, ~~which is a model result and not fully representative of the past climate history.~~ 2015. Climatologies for the ~~pre-industrial~~ equilibrium state were created such that when adding the historic anomalies the atmosphere and ocean forcings between 1995 and 2014 match the present-day observations. Observed present-day velocities do not directly enter the initialisation, but instead are used, amongst other criteria, indirectly to determine optimal parameters in the initial state ~~ensemble, which is spanned over various uncertain parameters~~ related to basal sliding and ~~ice flow.~~ From this ensemble of initial states, ~~we~~ sub-shelf melting. We select the state that best replicates present-day ice thickness, grounding-line position, mass loss, and ice surface velocities ~~;~~ We do so by using a scoring scheme that tests the (root-mean-square) deviations from observations in each individual variable by normalising and multiplying them. We lay a specific focus on the Amundsen, Weddell, and Ross seas by including the regional scores in addition to the continental values. We then use the best scoring run for the ‘best’ PICO parameters ~~for the stability experiments~~ (“AIS5” in Reese et al., 2022). ~~Note that this state is slightly gaining mass over the historic period between 1992 and 2015. Further details on PISM are given in Appendix A3.~~

2.4 Ice sheet initial states

2.3.1 Initial state

~~We have generated three ice sheet model initial states which closely replicate the current geometry, surface velocity, and grounding-line positions of the Antarctic Ice Sheet. We interpolate these initial states onto a regular 2-km grid to evaluate the performance of the ice sheet initial states.~~ Despite the different initialisation procedure of PISM, the resultant ice sheet reference state with respect to ~~observations.~~ All indicators show a good agreement between each of the ice sheet models and the observations (Table 2). ~~Despite different initialisation procedures, all three models are able to closely replicate the observed ice thickness and surface velocities, particularly in the location of fast flowing ice streams (see Supplementary Figs. S5 and S6). It is not surprising that Elmer/Ice and Úa have geometry and ice velocities is in good agreement with observations, given that they impose the surface topography from BedMachine and use observed surface velocities in the inversion. However, PISM is also and the states obtained in the other models. PISM is~~ capable of locating most, if not all ice streams in their correct positions, and accurately replicating the observed ice thickness and surface velocities after the spin-up (see Figs. S5 and S6).

345 To assess the sensitivity of Antarctic grounding lines to small changes in their position, we need to closely replicate the
current observed position of the grounding line. Figure 2 shows that the Due to the spin-up procedure and the coarser grid
resolution of PISM there are some areas with greater deviation in the initial grounding-line positions of all three models
are in most parts in very good agreement with the observed position of the grounding line in the BedMachine dataset. We
calculate the position. The overall error in the grounded ice area (a proxy for grounding-line change), by differencing the
350 simulated and observed grounded ice areas. To obtain a relative displacement of the grounding line itself, we normalise this
area change by the simulated length of the grounding line. The resulting grounding-line position errors are 84 m for Elmer/Ice,
540 m for Úa and grounding line position for PISM is 11.4 km for PISM, which is below, or km, which, while higher than
the other two models, is very close to ; the minimum the horizontal grid size used in each model. In addition, the good
agreement of observed and modelled grounding-line positions can be seen in our individual glacier profile figures presented
355 in the results. As expected, there is a smaller deviation between the observed and modelled grounding lines in Elmer/Ice and
Úa, due to their inversion and relaxation initialisation procedure, in which the grounding lines have had little opportunity to
migrate away from their prescribed initial (observed) position. Due to the spin-up procedure and the coarser grid resolution of
PISM there are some areas with greater deviation in the initial the model. Overall there is good agreement between modelled
and observed grounding-line positions. Nonetheless there is overall good agreement positions (Fig. 2), including in regions of
360 particular interest, such as Thwaites Glacier.

By design, initial states created by Úa and Elmer/Ice are as close as possible to steady state. During a steady-state control
simulation for both models there is less than 4.2 mm of The model grounding-lines positions are calculated using the flotation
criterion.

During the historic simulation (1850–2015), the change in sea-level equivalent volume over 100 years. Furthermore, there is
365 very little drift in the position of the grounding line from the initial location, deviating by only 12.6 (Elmer/Ice) and 2.1 m (Úa)
over 100 years, which is very small with respect to grid resolution. To support the findings of this study further, we generate
an initial state using PISM, that includes the recent history of the ice sheet, and therefore is not in steady state (see Sect. 2.3).
Due to the spin-up procedure of PISM, when it arrives at a steady state, by definition that state will be stable. Hence, we
would not need to additionally perturb such a state to determine if it is stable or unstable. Instead, we analyse a state including
370 a present-day trend in mass changes. This allows us to examine if current is -1.88 mm. While this is lower than observed
rates of mass loss over the last decades, due to mass gains from snowfall, the pattern of thickness change is comparable to
observations (see Reese et al., 2022).

2.4 Experimental design

We have generated three ice sheet model initial states which closely replicate the current geometry, surface velocity, and
375 grounding-line positions are also reversible when transient forcing is included, and is therefore complementary to the steady-state
perturbations in the other models of the Antarctic Ice Sheet (Table 2). We account for any remaining discrepancies between
individual models by running control simulations alongside the perturbation experiments, and the results presented are then
with respect to these control runs.

Indicator	Unit	Elmer/Ice	PISM	Úa	Reference	Source
Ice extent (total)	10^6 km^2	13.48	13.58	13.57	13.52	Morlighem et al. (2020)
Ice extent (floating)	10^6 km^2	1.45	1.10	1.51	1.50	Morlighem et al. (2020)
Ice extent (grounded)	10^6 km^2	12.03	12.48	12.06	12.03	Morlighem et al. (2020)
Ice mass	10^7 Gt	2.39	2.39	2.41	2.38 ± 0.04	Morlighem et al. (2020)
Ice mass above flotation	10^7 Gt	2.05	2.05	2.03	2.09 ± 0.04	Morlighem et al. (2020)
Ice mass above flotation	m SLE	56.67	56.46	56.06	57.9 ± 0.9	Morlighem et al. (2020)
Ice flux across the grounding line	Gt yr^{-1}	1624	2124	1727	1929 ± 40	Gardner et al. (2018)
Surface mass balance (grounded)	Gt yr^{-1}	1868	2192	1874	1792	van Wessem et al. (2018)

Table 2. Comparison of the initial state indicators for the initial states created using Elmer/Ice, Úa and PISM. All variables, except for flux across the grounding line, were calculated across the same 2 km resolution grid. Reference values for ice extents (total, floating, and grounded) were calculated using the BedMachine v2-dataset (Morlighem et al., 2020), whereas ice mass values were taken from Table S3 in the dataset paper (Morlighem et al., 2020) (m SLE, metres sea-level equivalent). The total flux across the grounding line was calculated in each respective model. Observed grounding-line flux was taken from Gardner et al. (2018) and total surface mass balance is from [RACMO RACMO2.3p2](#) (van Wessem et al., 2018).

Modelled and observed present-day Antarctic grounding-line locations. Present-day Antarctic bed topography (BedMachine;) showing regions where the bed topography is below (blue shading) and above (brown shading) sea level in meters above sea level (m a.s.l.). Modelled initial grounding-line positions are shown as colored lines, observed grounding-line position is shown in black. Ice shelves are indicated by grey shading. Golden lines denote the locations of the transects shown in Fig. 4. Grey lines mark the boundaries of the IMBIE basins. The inset shows a zoom into the Amundsen Sea Embayment sector of West Antarctica.

385 2.5 Experimental design

During the control simulations the mass balance in Elmer/Ice and Úa is

$$\nabla \cdot (\mathbf{v}h) + \frac{dh}{dt} = \dot{m}_{\text{mod}} \quad (2)$$

where $\frac{dh}{dt} \approx 0$ and \dot{m}_{mod} is the same as defined for the previous models in Eq. (1). In PISM the mass balance is

$$\nabla \cdot (\mathbf{v}h) + \frac{dh}{dt} = \dot{m}_{\text{pd}} \quad (3)$$

390 and $dh/dt \neq 0$ and is broadly comparable to present-day rates of ice thickness change.

We take the initial ice sheet states presented in the previous section and apply a small-amplitude perturbation to assess the stability of the current grounding-line positions. This perturbation is designed to be as small as possible, i.e. we want to impose a small deviation in the current grounding-line position such that the signal in the We apply a perturbation to the current position

of the grounding line is clearly visible, and distinctly emerging from the noise, but we do not want to force the grounding lines significantly into a different state. In lines by increasing the far-field ocean temperature that drives the melt rates calculated by PICO. Note that in general, it would be possible to perturb the system using a number of different control parameters in our models. Given the important role that ice shelves play on inland ice sheet dynamics and grounding-line position, via buttressing forces, we here choose to perturb the system by applying a shift in ice shelf-sub-shelf melt rates.

We apply this perturbation in all three models by increasing the far-field ocean temperature that drives the melt rates calculated by PICO. We increase the input ocean temperature, which is assumed to be representative of the conditions at depth on the continental shelf, by $[+1; +3; +5] \Delta T \in \{1, 3, 5\}$ °C all around the Antarctic Ice Sheet. This perturbation in temperature is applied for 20 years to create a numerically significant grounding-line retreat. By applying different increases in ocean temperature, we are able to test the robustness of this small perturbation and found that a 5 °C perturbation over 20 years remained small, i.e. it resulted in a small but obvious deviation in the position of the grounding line. After 20 years we remove the perturbation. While +5 °C appears to be an unrealistically high magnitude change, we want to stress that this perturbation is not designed to be realistic and is only applied over a few decades. Instead, we can think of our small perturbation as a small movement of the grounding line away from its current position, on the order of a few grid cells or ice thicknesses at the grounding line. Indeed, the +5 °C perturbation leads to approximately 10 km of retreat along the profiles in the ASE region (see Results) and along other profiles (see Figs. S16 and S17). We note that a recovery for the +5 °C perturbation also implies that a smaller perturbation would also be reversible, and allow a recovery phase of the simulation for a further 80 years (Fig. 1). We extend some simulations by 400 years to test the robustness of indicates that the grounding-line evolution over longer timescales. In parallel, we run a reference simulation (control run) with no melt perturbation. In all of our control and perturbed simulations the surface mass balance remains fixed, which in Elmer/Ice and Úa is the corrected mass balance field \hat{m} described in Sect. 2.2.1, and RACMO surface mass balance in PISM. steady state is stable.

There are subtle differences in the application of the melt perturbation between models, due to their initialisation. As Elmer/Ice and Úa already include a correction to the mass balance, we apply ice shelf melt anomalies as $\Delta \dot{b}_{PICO}(t) = \dot{b}_{PICO}^{pert}(t) - \dot{b}_{PICO}^{ref}(t)$ where \dot{b}_{PICO}^{pert} and \dot{b}_{PICO}^{ref} are the perturbed and non-perturbed melt rates, respectively. Both \dot{b}_{PICO}^{pert} and \dot{b}_{PICO}^{ref} vary with time, to account for cavity geometry changes that influence the calculation of melt rates in PICO. In the control simulations and during the 480-year recovery phase, $\Delta \dot{b}_{PICO} = 0$. During Across the floating ice shelves, during the 20-year perturbation phase, $\Delta \dot{b}_{PICO}$ is then subtracted from the synthetic mass balance term \hat{m} to give a total perturbed mass balance $\hat{m}_{pert} = \hat{m} - \Delta \dot{b}_{PICO}$. In PISM, no such correction to the mass balance has been applied, instead, $\dot{m}_{\Delta T}$ is the anomaly in sub-shelf melting calculated using PICO $\delta \dot{b}_{PICO}$ applied to the modified mass balance fields in Elmer/Ice and Úa as

$$\dot{m}_{\Delta T} = \dot{m}_{mod} + \delta \dot{b}_{PICO} \quad \text{where} \quad \delta \dot{b}_{PICO} = \dot{b}_{PICO}(T + \Delta T) - \dot{b}_{PICO}(T). \quad (4)$$

Across the grounded areas no perturbation is applied, and the mass balance is unchanged (Eq. 1). In PISM the mass balance field has not been modified from present-day conditions, and unlike the other two models the initial temperatures T in PICO are perturbed directly by ΔT

$$\dot{m}_{\Delta T} = \dot{b}_{RACMO} + \dot{b}_{PICO}(T + \Delta T). \quad (5)$$

We compare the ~~basal-mass-balance-integrated sub-shelf melt~~ perturbation applied in all three models and find that it is comparable for all experiments (see Fig. S10b).

430 ~~Here, we use the integrated ice flux across the grounding line as the metric of the system state, and to determine the reversibility. After 20 years we remove the perturbation and allow a recovery phase of the simulation for a further 80 years (Fig. 1). We extend some simulations by 400 years to test the robustness of the grounding-line position. This is because the ice flux is inherently linked to the MISI hypothesis, where retreat down a sloping bed increases ice flux, leading to thinning, flotation, and further evolution over longer timescales. Throughout the forward-in-time experiments, the extent of the ice~~
435 ~~shelves remains unchanged; however, all models impose a minimum ice shelf thickness, which is sufficiently thin to represent ice that has been removed and provides no buttressing.~~

~~During the recovery phase (80–480 years after the end of the perturbation), we examine the temporal evolution of the integrated ice flux across the grounding line. We choose~~ grounding-line ~~retreat. Ice flux is also flux as our metric of the system state because it is~~ found to recover faster to a perturbation than grounded area or volume, due to the long timescales needed for
440 the ice to thicken and re-advance. In the ~~steady-state steady-state~~ initial states of Úa and Elmer/Ice, by design, the ice flux across the grounding line balances the surface accumulation upstream. Increased ~~ice-shelf melting sub-shelf melt~~ in our simulations leads to a sharp increase in ice flux across the grounding line, which is assumed to be due to a loss of buttressing as a result of ice-shelf thinning. If the flux across the grounding line returns to its initial value after the perturbation is removed, this indicates that the ice sheet reverts to a steady state with a balance between surface accumulation in the grounded regions and
445 grounding-line flux (note that surface accumulation is altered a little by the grounding-line movement). Hence, it is assumed that a return of the grounding-line flux indicates the grounding line has either reverted back to its initial position or has begun to re-advance towards its former position. When the grounding line does not retreat further, it means that it has found a new stable position very close of the previous one. If the flux were to increase away from its initial value, the grounding line is unstable. To support this we also examine the trend in ~~grounded-line grounded-line~~ position after the perturbation is removed,
450 which is calculated as the change in grounded area for a constant grounding-line length.

To further analyse the response of the grounding lines after the perturbation is removed, we calculate the e -folding relaxation time, i.e. the time taken for the flux to decrease by a factor of e (Euler's number; ≈ 2.17). To do this we fit an exponential decay function in the form

$$\Delta Q(t) = \Delta Q_{\text{pert}} e^{-t/\tau} \quad (6)$$

455 to the change in flux during the recovery period of 80 years $\Delta Q(t)$, where ΔQ_{pert} is the change in flux at the end of the 20-year perturbation relative to the initial, unperturbed flux, t is time after the perturbation, and τ is the recovery timescale. We repeat this for all three models and perturbation experiments, and these exponential curves can be seen in ~~Supplementary~~-Figs. S7 to S9.

3 Results

460 In the following sections we present the results of our perturbation experiments of present-day Antarctic grounding lines as described in Sect. 2.4. Figure 3 shows the integrated ice flux across the entire Antarctic grounding lines in each model during the perturbation experiments. We also present the results integrated across the 27 basins from IMBIE (Fig. 2; Zwally et al., 2012), excluding basins 7, 8, and 25 which contain only small ice shelves. Additional figures showing grounding-line position change and volume above flotation can be found in the Supplement ([Figs. S10 to S15](#)).

465 3.1 Antarctic Ice Sheet

On the Antarctic-wide scale, all models and all perturbations show a similar trend; a strong increase in ice flux, reaching a maximum at the end of the 20-year perturbation period, followed by an exponential decrease for the remaining 80 years (Fig. 3). The magnitude of the flux response to the melt perturbations is comparable between all ice sheet models, in particular for the 1 °C temperature experiment, in which ice flux increased by approximately 300 Gt yr⁻¹. In the higher temperature
470 scenarios (3 °C and 5 °C), the flux responses diverge slightly from one another, PISM and Úa remain similar, while Elmer/Ice shows a stronger increase. This is likely due to subtle differences in the imposed basal melt perturbation in each model, which also diverge with the magnitude of the perturbation (see Fig. S10b).

~~After the perturbation in ice shelf melt rates is removed, Antarctic-wide ice flux in all ice sheet models decreases exponentially, tending towards the initial value.~~ The calculated e -folding flux response time reveals that the Antarctic wide recovery time is in
475 good agreement for all models, ranging between 10 and 20 years, and is largely independent of the magnitude of the perturbation (Fig. 3, bottom panel). During the 80-year recovery period, the flux decreases rapidly, but is not fully recovered. To check if it is able to recover eventually, we extended the relaxation period in the 5 °C simulations in all models to 480 years and found that the Antarctic-wide flux returns to within 3.5 % of its original value (see [Supplementary](#)-Fig. S13). Alongside the ice flux evolution, the total retreat of the grounding line, and ice volume show similar trends: rapid retreat and reduction in ice volume
480 during the perturbation, after which retreat rates subside and grounding lines begin a slow recovery. ~~While they do not fully recover within 80 years, due to the aforementioned slower response timescales, in~~ [Furthermore](#), the Antarctic wide signal there is no indication of accelerated retreat (see [Supplementary](#)-Fig. S10). The recovery of the ice flux, alongside a short, two-decade e -folding time, strongly indicates that the majority of Antarctic grounding lines are reversible ~~with respect to current geometry, when perturbed by increased ice shelf melt rates. If large parts of the Antarctic grounding line were irreversible, i. e. ice flux~~
485 ~~were to increase away from the initial value, we would expect a slower or non-existent recovery for the total Antarctic ice flux. Indeed.~~ [Also](#), ice flux evolution for individual basins (in all models) appears to show an exponential decrease after the perturbation is removed (Fig. 3). However, some basins recover quicker than others. In the remainder of this section we explore the response of individual basins in more detail.




Figure 3. Reversibility experiments and recovery time scales. Change in integrated ice flux across the grounding line for perturbed simulations using three ice sheet models. All three temperature perturbation experiments (1 °C, 3 °C, 5 °C) are shown for selected individual basins. For ease we merge the results for basins flowing into the Amery, Filchner-Ronne and Ross ice shelves. See [Supplementary-Fig. S12](#) for additional basins. PISM fluxes are smoothed using a running mean filter of 5 years. Grey shading shows the perturbation period. Bar plot shows the e -folding recovery time. Each bar shows the median response time from all three experiments (1 °C, 3 °C, 5 °C) for each model, and error bars show the range. Bars are not shown for individual models for some basins (e.g. Cape Adare for PISM) where the exponential fit to the change in ice flux was deemed poor and the R^2 value is less than 0.8. There are four individual experiments for PISM where the flux results are noisy (due to coarser grid resolution) such that the exponential fit for that individual temperature perturbation has $R^2 < 0.8$ (see Fig. S8). We set these values to zero as the lower end of the error bars.




Figure 4. Profiles of the Amundsen Sea Embayment (ASE) sector for the three models at different time intervals during the 5 °C experiment. Shown are the ice-sheet geometries of the initial state (dotted line filled with light blue), at the end of the perturbation at 20 years (red line), after 80 years of recovery at 100 years (orange line), and after 500 years (cyan line). The dotted cyan line shows the location of the control simulations after 500 years for the PISM simulations. Observed (BedMachine) grounding-line positions are indicated by a black dots. The small panels show a zoom into the region marked by the black squares. The resolution of each profile depends on the model resolution. Profile locations are shown in Fig. 2.

3.2 Amundsen Sea Embayment sector

490 The Amundsen Sea Embayment (ASE) sector encompasses the Pine Island, Thwaites, and Getz drainage basins (basin numbers 20 to 22 in Fig. 2) ~~and is of particular interest as it drains ice from some of the marine regions of the West Antarctic Ice Sheet.~~ For all three of these basins, our results show a rapid exponential decay in the ice flux ~~after the 20-year perturbation. This rapid recovery is also reflected in the median~~ (e -folding times ~~for all three models and all three basins in the ASE sector, which are less than 20 years. In addition to the basin-wide flux responses shown in~~ < 20 years) ~~after the 20-year perturbation~~ (Fig. 3, ~~we~~ 495 ~~).~~ We extract profiles along four marine glaciers in the ASE sector (Fig. 4) to show the response of a vertical transect along the ice shelves at four critical points of the 5 °C experiment. These flow-lines were directly interpolated from the original model grids. We note ~~that there are~~ some differences in the bed geometry due to the different resolution and interpolation methods used by each model.

At Pine Island Glacier, the initial grounding-line position in Úa is close to observations, whereas in Elmer/Ice and PISM 500 the initial grounding-line positions are located downstream at a topographic ridge (Fig. 4a). ~~Despite their different starting locations, Grounding lines in~~ Elmer/Ice and Úa ~~grounding lines retreat (approximately 3-7 km ; retreat approximately 3 to 7 km~~ (Fig. S11) across sections of retrograde sloping bed topography during the perturbation. ~~The grounding line in PISM. In PISM the grounding line~~ retreats along a section of pro-grade slope to the top of the ridge. ~~For all models the 5 °C perturbation causes the ice shelf to disappear entirely.~~ After the perturbation is removed, ~~all models show the grounding lines to grounding~~ 505 ~~lines in all models~~ advance back to their initial positions ~~. For all three models, the 5 perturbation causes the ice shelf to progressively thin until it disappears entirely and then grows back to , at least, its~~ ~~and the ice shelves regrow to their~~ initial thickness. Grounding-line flux (Fig. 3) ~~shows a clear V-shape (increase to 20 years, and decrease after), where~~ ~~increases during the 20-year perturbation and sharply decreases thereafter. At year 50 the flux in Úa and PISM show the flux to have returned to~~ ~~has decreased~~ either below the control simulation (PISM) or within 0.7 % (Úa) of the initial flux ~~in just 30 years after the~~ 510 ~~perturbation is removed (year 50). However, the recovery is slower in.~~ In Elmer/Ice ~~, which is reflected in the longer recovery timescale (approximately 11 years) compared with PISM and Úa~~ ~~the recovery is slower (11-year e-folding time) than the other models,~~ and the flux remains 10 % higher than the initial at 100 years. ~~This is also reflected in the median response times which are 6 and 7 years respectively for Úa and PISM, and 11 years for Elmer/Ice. year 100.~~

Our results for Thwaites Glacier are similar to Pine Island (Fig. 4b). ~~During the perturbation the grounding lines in~~ In the 515 entire Thwaites basin (including Dotson and Crosson ice shelves) ~~retreat by approximately 4 to 9 km (Fig. S11) and the ice flux across the grounding line~~ ~~grounding line flux~~ increases between 50 and 170 Gt yr⁻¹ ~~. After the perturbation is removed the grounding-line fluxes are recovered in all three models and the response times are within~~ ~~during the perturbation, and then decreases rapidly (response times~~ 8-15 years (Fig. 3) ~~) in all models after the perturbation is removed.~~ At Thwaites Glacier ~~in particular,~~ the initial ~~locations of the~~ grounding lines in all three models are in close agreement with one another (Fig. 4b). For 520 Elmer/Ice and Úa this is slightly downstream of the observed grounding-line position at a topographic ridge, whereas PISM is located at the currently observed position. The retreat during the perturbation is not across a section of reverse bed slope, but instead remains downstream of a second topographic ridge (located 70 km along the profile). For PISM and Úa, the ice

shelf thins strongly during the perturbation, where as in Elmer/Ice, the ice shelf is relatively thick in the initial state and does not disappear during perturbation. After the end of the perturbation, the ice shelf grows back to its initial shape in all models. In Úa the grounding lines re-advance fully to their initial positions (within 100 years), and PISM re-advances to the position arrived at in the control simulation after 500 years. The recovery in Elmer/Ice is slower than Úa which corresponds to the longer recovery timescale for the entire Thwaites basin (Fig. 3).

Similar behaviour is also observed for Crosson and Dotson ice shelves for Elmer/Ice and Úa (Figs. 4e-dc-d). The ice shelf thins and the grounding line retreats across a prograde slope during the perturbation. The grounding line advances towards their initial position after the perturbation is removed, and in Úa returns fully to its former position after 500 years. ~~The reversing trend after 100 years indicates similar behaviour with Elmer/Ice.~~ In contrast, the Crosson and Dotson grounding lines in PISM continues to retreat and the ice surface lowers after the perturbation is removed. This additional retreat is also reflected in the longer recovery time for Thwaites basin for PISM (Fig. 3), which includes the Crosson and Dotson ice shelves show signs of ongoing retreat. Crucially, ~~however,~~ the grounding lines in PISM start slightly retreated of in slightly retreated positions compared to those of observations and the two other models. After the perturbation, ~~between 20 and 100 years of the simulation (20–100 years) the ice shelves thicken and~~ the grounding lines of both glaciers re-advance and the ice shelves thicken. However, after 100 years, the ice surface lowers and the grounding lines retreat again, to a location further inland of the perturbed position (at 20 years). Importantly, the final grounding line positions after grounding line positions at year 500 years, are very similar to their location after 500 years in the control simulation, showing that they are reversible with respect to the trend in the control simulations. Long-term (1510,000-year) simulations under present-day climate forcing also show that the Crosson and Dotson grounding lines will remain (in the absence of additional forcing) at this slightly retreated position in the future (see Fig. 4 in Reese et al., 2022).

For additional basins in this region of West Antarctica, the Getz (Fig. 3) and those in the Bellingshausen Sea sector (basins 23 and 24 in Supplementary Fig. S12), the results are similar, i.e. the ice flux tends toward its initial value, the grounding lines re-advance to their former positions (Fig. S11), and the response times are less than 20 years (Fig. 3). In summary, despite strong thinning of the buttressing ice shelves and associated ice shelf thinning and retreat of the grounding lines in the ASE sector of the ice sheet during our perturbation experiments, we find that the grounding lines return to their initial positions after the perturbation is removed and the buttressing is reapplied.

3.3 East Antarctica

Alongside the strong signal of recovery seen in West Antarctica, a number of Our experiments for basins in East Antarctica show a similar response to melt rate perturbations similarly strong signal of recovery to those in the ASE region. In particular, Wilkes, Totten and Amery basins all display a rapid exponential decay in flux (e -folding times < 20 years) after the perturbation is removed, which is reflected in their short response timescales, in most cases < 20 years (Fig. 3). Within 100 years, the flux in these basins appears to be reversible in Úa and PISM, whereas within 100 years, and within 500 years in Elmer/Ice the flux is reversible within 500 years (Fig. S15). Alongside this, these basins Grounding lines in these basins also show a strong signal of re-advance of the grounding line after the perturbation is removed (Fig. S11). This is also evident in selected

glacier profiles at Totten and Cook ~~Glaciers~~ glaciers (Fig. S17**b-eb-c**). At both glaciers, the ice shelves disappear under the strongest perturbation (5 °C) ~~results in the entire loss of the ice shelves. Despite this, after~~ but at year 100 years the ice shelves have almost recovered their former ice thickness and grounding-line positions. At Cook Glacier the grounding lines have fully
560 returned to their initial positions after 500 years in Úa and PISM and in Elmer/Ice is tending towards the initial position. ~~We note that PISM shows retreat and subsequent re-advance along a section of reverse bed slope at Cook Glacier, but this retreat takes the grounding line to the same starting location as that in Úa and Elmer/Ice.~~ At Totten Glacier, all models start close to the observed grounding-line position, and retreat approx~~3-6~~ 3 to 6 km inland (Fig. S11). After the perturbation is removed Úa fully recovers by 500 years and Elmer/Ice has regrounded at the observed position (Fig. S17). ~~The ice shelf in PISM has thickened, but~~ In PISM the grounding line has not reverted to its initial position after 500 years. ~~However, the grounding line has, but has instead~~ approached the same location ~~after 500 years~~ in the control simulation and can therefore still be considered reversible.

At Amery ~~ice shelf~~ Ice Shelf, all models show that the ice shelves thicken, and the ice flux has almost fully recovered within 100 years, but the grounding line has not returned to its former position. However, it appears to have found a new stable
570 grounding-line position within a short (approx. < 10 km) distance inland, and does not experience accelerated retreat thereafter (Fig. S17a). We note that a number of smaller basins surrounding Totten, Wilkes and Amery (basins 12, 15, and 16) show a similar recovery of the ice flux within approximately < 20 years and re-advance of the grounding lines.

In ~~the~~ Dronning Maud Land ~~region of East Antarctica~~ (basins 4, 5 and 6), the signal of recovery is similar; all three ice sheet models show an increase in ice flux and retreat of the grounding line during the perturbation, and a decreasing trend in ice flux
575 after the perturbation is removed (~~Supplementary~~ Figs. S12 and S11). In general, the recovery of the grounding-line position is slower in these basins ~~, as reflected in their recovery timescales which are often~~ (e-folding times > 15 years, ~~in particular in Úa~~ (Fig. 3). However, the extended relaxation period shows that the flux in the 5 °C experiment recovers to within 3 % of its initial value for all models and all three basins ~~(4-6). At the same time~~ 4-6. Concurrently, all grounding lines ~~are slowly re-advancing~~ slowly re-advance to their former positions, and show no signs of accelerated retreat. ~~We note that these basins are not considered to be vulnerable to MISI, and currently show mass gain rather than loss.~~

580

3.4 Filchner-Ronne and Ross Ice Shelf sectors

Elsewhere, the large basins draining into the Filchner-Ronne and Ross ice shelves show more complicated behaviour. In general, we do not see any strong signs of instability, i.e. the flux decays towards its initial value after the perturbation is removed, similar to the previously discussed basins. However, for the Filchner-Ronne basin in particular, the ice flux in all
585 models remains 20 – 40 % away from its initial value after the 80-year relaxation period in the 5 °C experiment, and all models show response times of approximately 20 years (Fig. 3). In addition, all models show some signs of further retreat after the perturbation is removed (between ~~20-100~~ 20-100 years in Fig. S11), but at a reduced rate. However, we find that in our extended simulations (for 5 °C) the grounding lines either (1) tend to re-advance towards their initial positions, (2) show a re-advance of a few to several kilometers before appearing to reach a new steady position (Fig. S14), or (3) settle on a slightly
590 retreated position a short distance inland. In no cases do we see further retreat inland by the end of the simulations after 500

years. For both the Filchner-Ronne and Ross basins the flux decreases to within 10 % (Elmer/Ice), or 6 % (PISM and Úa) of its initial value after 500 years.

We extract profiles at major outlets feeding into the large Filchner-Ronne and Ross ice shelves (see Fig. S16). For both profiles feeding the Ross Ice Shelf and the Ronne Ice Shelf, all models show signs of thinning and some additional retreat after the perturbation is removed (between 20 and 100 years). However, during the extended relaxation period (100 to 500 years) the ice shelves have begun to thicken again and the grounding lines do not retreat further inland. In the Filchner profile we find similar behaviour, with some additional retreat after the end of the perturbation (20 years). The grounding lines in PISM and Elmer/Ice do not retreat further after 100 years, but remain in the same (slightly inland) position. In Úa the grounding line begins to re-advance and has almost returned to its initial (and full) ice shelf extent after 500 years. Overall, there are no signs of accelerated retreat in these basins, and instead the grounding lines of the larger ice shelves may need longer to recover from a perturbation.

4 Discussion

~~Here, using three ice sheet models, we have applied small-amplitude perturbations to the current position of the Antarctic grounding lines to assess whether their retreat is due to the MISI. Our results show that most grounding lines, including those of major marine basins in West Antarctica, are able to recover their former positions when the perturbation in ocean-induced melt is reversed, and therefore can be considered stable in the case of Elmer/Ice and Úa, as well as recoverable in the case of the transient PISM state. This is clearly shown by the rapid exponential decay (e -folding time of < 20 years) of the ice flux (our metric of system state) towards its initial value after the perturbation was removed, both for the entire ice sheet and for most individual basins. Our results have also shown that individual grounding lines can return towards their former positions (Fig. 4) within 100 years, particularly at the smaller basins/ice shelves, whereas the larger basins re-advance at a slower rate. Importantly, there are no signs of accelerated retreat with respect to a small perturbation in the position of the grounding line from its current location. This conclusion is further supported by the PISM simulations, which also show reversible grounding lines after removing the perturbation. report that grounding lines can persist at bed topography peaks and ‘initiate retreat potentially long after the onset of a change in local climate.’ In our experiments, we capture the centennial time scale over which such a response could occur for the entire Antarctic grounding lines in our 480 year relaxation period. Since we find no indication of further retreat during these simulations, we are relatively confident that this non-linear response is not at play in our simulations. We further test the millennial timescales in PISM and find no indication of this process over 10,000 years.~~

~~Overall, we have shown that the present-day grounding-line positions of The perturbation experiments described above all showed that with time the grounding lines reverted back towards the state prior to the application of the perturbation. Where we ensured that the initial state was steady, as done in experiments using Elmer/Ice and Úa are stable in their current geometry; in the absence of any present day external forcing. However, it is imperative that this finding is understood in the context of our steady ice sheet model states.~~

The current state of the Antarctic Ice Sheet is not in steady state; areas in the ASE sector in particular are thinning at a rate of several metres per year. To assess the stability of the grounding lines in their current geometry, we required a steady state model set-up. As aforementioned, this is to exclude the role of external forcing (i.e. climate/ocean warming) and allow us to evaluate whether present-day grounding lines have already crossed an internal instability threshold. To bring Elmer/Ice and Úa into steady state, a small correction to the mass balance imposed in these models has to be made (see Fig. ??). This amounts to 76 in Elmer/Ice and 81 in Úa, which represents 4.2% and 4.5% of the total surface mass balance respectively (Table 2). In both models the correction terms needed to obtain a steady state are largest in West Antarctica, in particular in the ASE sector, when compared to East Antarctica. This reflects the current flux imbalance in West Antarctica, that is driving present-day migrated back to their original steady-state locations. Hence, these steady states are stable. In other experiments, where the grounding lines were allowed to migrate freely following the initialisation procedure, the grounding-line retreat. Compared to migration rates reverted back to those of the reference (i.e., unperturbed) runs. In no experiments conducted with the three ice-sheet models did we see any indication of self-enhanced irreversible retreat from the currently observed locations of the impact the perturbation has on the flux, this imbalance only accounts for 3–5% in the 5 °C experiments (Fig. 3). For the smaller perturbations the ratio is larger, but lower than 15–30%. Given that the amplitude and spatial distribution of the corrected mass balance fields are highly variable between the two models, and neither model shows signs of instability, we assume that our results are independent of this correction. In addition, we do not find any signs of self-sustained retreat in the simulations conducted by PISM (that does not correct the surface mass balance), which is a further indication that the results are independent of individual model choices.

While correcting the mass balance in this manner allows our modelled grounding lines and geometry to remain close to observations, the compromise is an ice sheet that is less representative of the actual balance between ice flux across the grounding line and the integrated surface mass balance upstream. This imbalance in the present-day ice fluxes (i.e. the integrated accumulation over the grounded area is lower than the integrated flux across the grounded line) clearly highlights that present-day grounding lines are, and likely will continue, to retreat over the coming decades grounding-lines of the Antarctic Ice Sheet. Our results do not suggest the contrary, but only that present-day retreat is externally driven rather than due to an internal ice sheet instability. Note that to draw this conclusion about no ongoing internally-driven retreat, our methodology requires that the critical thresholds for the real ice sheet (1) are correctly represented in our initial configurations and (2) have not been altered by modifications to any of our model parameters to obtain a steady state. We have achieved both of these by first creating initial states as close as possible to present-day observations, and second, only applying a small correction to the surface mass balance in order to bring Úa and Elmer/Ice into steady state.

Given the ongoing impact of changes in external climate forcing on the Antarctic Ice Sheet, it is important to address whether present-day climatic conditions are sufficient to force grounding lines to eventually cross a critical system state and embark on irreversible retreat. We address this question in our accompanying paper, by conducting multi-millennia scale simulations using an ensemble of present-day configurations in PISM (including the state used here) to assess whether present-day Antarctic grounding lines are committed to enter irreversible, self-sustaining retreat under current climate forcing. Indeed, we show that when present-day external climatic drivers are present, current grounding lines can eventually be forced into regions of the

bed topography in which the grounding-line retreat then becomes irreversible. Hence, some grounding lines are committed to enter into MISI driven retreat unless present-day climate forcing were to be reversed. This supports previous suggestions that present-day climate conditions may be sufficient to trigger rapid grounding-line retreat in West Antarctica in the long term therefore suggest that all grounding lines of Antarctica are not likely to be undergoing retreat due to MISI at present, including the grounding lines which are currently observed to be retreating.

It is perhaps surprising that the results of this paper ~~reveal~~ suggest that the ASE sector of West Antarctica has reversible grounding-line positions in response to a small deviation in their position, ~~and are therefore stable in their current geometry.~~ This is clearly replicated in all three ice flow models. Alongside this, Pine Island and Thwaites glaciers have fast response timescales, consistent with the results of Levermann and Feldmann (2019). This reversibility is evident even under the strongest perturbation in which given that a number of the smaller ice shelves disappear entirely during the perturbation period. A number of previous studies have suggested that the retrograde bed slopes of Pine Island and Thwaites glaciers may mean they have already begun entered internally-driven (MISI) retreat. Our experiments show that, under steady state conditions, i.e., the absence of an externally driven imbalance in the ice flux (albeit an unlikely scenario in present day), the Pine Island and Thwaites Glacier grounding lines are stable with respect to their current geometrical positions. Previous work has (e.g., Joughin et al., 2010; Rignot et al., 2014; Joughin et al., 2014; Favier et al., 2014; Mouginot et al., 2014; Seroussi et al., 2017; Milillo et al., 2022). Our results instead show no indication of ongoing retreat in this sector of the ice sheet due to MISI. While previous work has indeed shown that the bed topography could cause tipping points to be crossed at Pine Island Glacier for example (Rosier et al., 2021). ~~Importantly, our steady state simulations do not contradict this finding, but show that, our results show that it is unlikely that~~ the current retreat of the grounding line at Pine Island is not (yet) driven by MISI, but is instead externally driven. This is also supported by observations that show grounding-line retreat to have recently stagnated, suggesting the current position is indeed stable. ~~When due to MISI. Importantly, once~~ the grounding line retreats further towards the critical regions identified in Rosier et al. (2021), internal instability is likely to ~~dominate over external forcing~~ be found.

At Thwaites Glacier, all models have shown that a small perturbation in the current grounding-line position (in the absence of external forcing) is not sufficient to initiate MISI retreat. We note that this still holds when present-day climate forcing is included (PISM experiments); the grounding line and ice shelf thickness return close to observations as shown by the steady state simulations of the other models (see Fig. 4). Crucially, We find similar results for Thwaites Glacier where an entire removal of the ice shelf ~~in our strongest perturbation~~ (in the strongest perturbation) is not sufficient to destabilise the current grounding-line position, as has been suggested, ~~due to the limited buttressing it provides.~~ Thus, present-day observed retreat at Thwaites is unlikely to be due to MISI. However, under present-day forcing all ensemble members in our accompanying paper show that the Thwaites grounding line retreats sufficiently far inland to enter irreversible retreat. A number of by previous studies (e.g., Wild et al., 2022). Several modelling studies have also shown ~~accelerated retreat under future climate forcing~~ is possible when the grounding line retreats further inland that once the grounding lines retreat further inland under future increases in ocean forcing (and in the absence of any changes in surface mass balance from present day), it is possible that they will enter phases of accelerated retreat (Joughin et al., 2014; Seroussi et al., 2017), ~~which.~~ This accelerated retreat could be an

indication of the onset of MISI. Note however that the projected grounding-line evolution of Thwaites Glacier in response to future ocean changes is associated with large uncertainties (Nias et al., 2019; Robel et al., 2019).

695 ~~Elsewhere, in George V Coast and Wilkes Land, regions with deep marine subglacial basins (see Basins 13 and 14 in Fig. 2) that are considered prone to MISI. To assess whether present-day Antarctic grounding lines are committed to enter irreversible retreat under current climate forcing, we conduct multi-millennial scale simulations using an ensemble of present-day configurations in PISM (including the state used here), see our accompanying paper (Reese et al., 2022). Indeed, we find no signs of unstable that current climate conditions can force grounding lines in their current position. Instead, these basins show a rapid recovery of the ice flux to its initial value, and re-advance of the grounding line to its former position within 500 years (see Basins 13 and 14 in Fig. S14). Cook and Ninnis glaciers in particular, have their current grounding-line positions seaward of the ‘ice plug’ region identified in . However, once the grounding lines retreat beyond this stabilising region, there is potential for rapid MISI-driven retreat into the deep Wilkes subglacial basin. A deep subglacial trough also exists at Totten Glacier, but again we find the grounding line to be stable in its current location. Hence, present day and near future retreat of Totten Glacier’s grounding~~
700 ~~line is likely to be driven by external drivers rather than MISI. the ASE sector to eventually enter irreversible retreat, but not within 300 years of constant present-day climate. This supports previous suggestions that present-day climate conditions may be sufficient to trigger rapid grounding-line retreat in West Antarctica in the long term (Golledge et al., 2021; Garbe et al., 2020; Joughin et al., 2014).~~

~~By repeating our perturbation~~ Here, by repeating our experiments using three different ice sheet models we have captured structural uncertainties associated with individual ice sheet models. ~~Despite differences in initialisation procedure, grid resolution, and in some cases the initial position of the grounding lines, no model shows any signs of irreversible retreat of grounding lines.~~ However, we note several potential sources of uncertainty. First, we ~~chose to perturb the grounding lines by increasing~~ parameterise the sub-shelf melt rates using the PICO sub-shelf melt model. While any parameterisation of sub-shelf melt in the PICO box model. Any parameterisation of ice shelf melting has associated uncertainties, and may not capture
715 ~~some important~~ certain physical processes in sub-shelf cavities. ~~However, the ice shelf, the sub-shelf~~ melt distribution is not considered to greatly influence our results for two main reasons: ~~–~~ First, our results are consistent across a wide range of temperature perturbations, including the ~~unrealistic scenario~~ strongest perturbation (+5 °C) in which several ice shelves ~~are removed entirely~~ disappear. Second, ~~our results are consistent across all three models despite the~~ different implementations of PICO ~~in each model will, that~~ lead to subtly different melt rate distributions. ~~However, our results are consistent despite this.~~
720 In addition, we could have perturbed the grounding line using different model parameters, e.g. relating to ice viscosity or basal sliding. Indeed, we modified the basal sliding using \dot{U}_a and the results were consistent with those presented here; grounding lines re-advanced after the perturbation was removed.

~~The goal of our experiments is~~ As our experiments aimed to assess the stability of the observed grounding lines in their current geometry. Hence, the results are dependent ~~not only on replicating current geometry in our ice flow models, but also upon observations of the~~ on accurate representations of the ice thickness and bedrock topography. ~~While our understanding of the subglacial topography of the Antarctic Ice Sheet has improved greatly in the last two decades, uncertainties remain, particularly in poorly surveyed regions of the ice sheet.~~ One approach to ~~quantify these uncertainties~~ assess the impact of ice

thickness and bedrock topography on our results would be to conduct additional perturbation experiments, with modifications to the topography within some bounds. In part, we have achieved this by ~~conducting simulations using using three~~ different ice sheet models ~~and their respective~~ with different grid resolutions, where in particular the bed topography around the grounding lines in PISM is smoother than the other models (see Fig. 4). Grid resolution itself is a source of uncertainty in ice sheet model simulations, in particular around the grounding line (Durand et al., 2009; Pattyn et al., 2013). ~~Again, we have partly captured uncertainties in grid resolution by conducting experiments using different models, but again this is captured by our experiments with different models and grids.~~ In addition, we repeated some experiments in Úa in which we modified the mesh resolution around the grounding line, and found that our results were not affected by a finer resolution (500 m) at the grounding line.

~~While our models replicate the current geometry of~~ In a previous study it was reported that grounding lines can persist at bed topography peaks and 'initiate retreat potentially long after the onset of a change in local climate' (Robel et al., 2022). We further test the millennial timescales in PISM (Reese et al., 2022) where we find long-term retreat in the ASE sector. The present-day state in the model could hence be such a "persistence". However, Robel et al. (2022) do not discuss the link between persistence and reversibility. In our case, we find that the ice sheet, it remains challenging to accurately represent all the processes and dynamics of the ice sheet. One particular physical process that is currently challenging to model is iceberg calving present-day retreat is not irreversible yet.

An additional source of uncertainty in our results is the absence of dynamic calving front positions in our models. In Elmer/Ice and Úa the ice front is fixed and the ice shelves maintain a numerically required minimum thickness, ~~even in the case of complete melting of the ice shelf. This is a common approach in a number of ice sheet models~~ due to the numerical challenges of removing ice entirely from the domain. It is possible that this thin layer of ice promotes ice shelf regrowth at a quicker rate than a true re-advance of the ice front, due to the external forcing, surface and basal mass balance, that are still applied across areas that have reached the minimum ice thickness. However, we note that the experiments conducted with PISM show similar recovery time scales, where in this case ice shelf ~~'cells'~~ cells are converted to ocean ~~'cells'~~ cells when the ice thickness decreases to zero. In addition, as the calving fronts are kept fixed, they are unable to advance beyond our prescribed initial position, which could also alter the stability of the grounding lines. We do not expect this to be relevant, because previous studies have shown that downstream regions (close to the calving fronts) of ice shelves provide little to no buttressing to upstream flow (Fürst et al., 2016; Reese et al., 2018b). We acknowledge that not imposing calving during our experiments is a limitation, especially as recent work has shown that (in the presence of buttressing) the calving law could impact the stability regime of grounding lines (Haseloff and Sergienko, 2022). Future studies should seek to include calving in such stability analyses. However, ~~in particular~~ for smaller ice shelves, the perturbations we apply effectively remove the entire ice shelves. As long as the ice shelves are allowed to regrow to their current extent after a calving event, our results can also be interpreted for such ~~calving styles~~ a perturbation. Alongside calving, ice shelf damage can have an important impact on the magnitude of grounding-line retreat due to weakening in the shear zones (Lhermitte et al., 2020). Hence, it is possible that with damage included, the retreat of the grounding line in our perturbation experiments would be larger. However, damage is not

yet sufficiently well parameterised for inclusion in our models, and implementing time-dependent damage in particular, is an ongoing (computational) challenge.

5 Conclusions

765 ~~Here we show~~ Our key finding is that the grounding lines of Antarctica, including Pine Island and Thwaites glaciers, ~~exhibit no~~
~~indication that~~ show no indication of marine ice sheet instability (MISI) ~~is already underway in the~~ in their ~~current~~ geometry
~~of the ice sheet. We initialised Úa and Elmer/Ice to be in steady state close to the present-day geometry of state. We arrive at~~
~~this conclusion by performing a large number of perturbation experiments to model states that replicate the current observed~~
~~geometry and velocity of the Antarctic Ice Sheet. In those experiments where we eliminated model drift to ensure that the~~
770 ~~initial state was steady,~~ the ice sheet, ~~and then performed a numerical stability analysis in which we applied a small-amplitude~~
~~perturbation in ocean-induced sub-shelf melt. In all cases we found that the grounding lines retreated slightly away from their~~
~~initial position while the perturbation was applied, and then returned towards their initial position after the perturbation was~~
~~removed. We repeated these experiments using an ice sheet state in PISM which included the present-day trend in mass loss and~~
~~found that this does not affect the reversibility of the grounding lines in the perturbation experiments~~ grounding line reverted
775 back towards to its original location, and in those experiments where model drift was allowed, the grounding line reverted
to the prior rate of migration. Taken together, this indicates that the Antarctic grounding lines are not currently engaged in
irreversible retreat caused by internal (MISI) dynamics.

~~While the experiments presented in this paper do not~~

There is a general consensus within the ice-sheet modelling community that the West Antarctic Ice Sheet is susceptible
780 to MISI. Here, we have argued that the currently observed changes of the Antarctic Ice Sheet are not a manifestation of an
ongoing positive-feedback related to MISI. While our experiments suggest that an internal instability threshold has ~~already not~~
~~yet~~ been crossed in Antarctica, ~~crucially, these experiments do not replicate the present-day mass imbalance of the ice sheet.~~
~~Given that it appears highly unlikely that present-day climatic changes can be reversed, future retreat of the grounding lines~~
~~future retreat~~ driven by changes in atmospheric/oceanic climate conditions could force the grounding lines to ~~reach regions of~~
785 ~~the bed topography at which the ice sheet crosses~~ cross a tipping point, ~~and after which~~ retreat becomes driven by ~~the MISI.~~
~~Future MISI. Further~~ work is needed ~~on an Antaretic-wide scale~~ to quantify the amplitude and duration of forcing required for
~~present-day grounding lines to transgress into an irreversible retreat that would cause grounding lines to the Antarctic Ice Sheet~~
~~to enter a phase of a large-scale irreversible collapse involving grounding-line~~ retreat over hundreds of kilometers, ~~thereby~~
~~causing several metres of~~ and a concomitant sea-level rise of potentially several meters.

A1 Elmer/Ice**A1.1 Model description**

Elmer/Ice (<https://elmerice.elmerfem.org/>) is an open-source finite-element software for ice-sheets, glaciers and ice flow modelling built on the multi-physics finite-element-model suite Elmer (<http://www.elmerfem.org/blog/>). Elmer/Ice is a very general and flexible tool and as such has been used for a large diversity of applications (181 publications since 2004). The main features and capabilities of Elmer/Ice have been described in Gagliardini et al. (2013) and in the associated publications (<https://elmerice.elmerfem.org/publications>). Here, only the main characteristics relevant for our analysis will be presented. Regarding the physics included in Elmer/Ice, the ice flow velocity is computed solving the Shallow Shelf Approximation (SSA) assuming an isotropic rheology following Glen’s flow law. The initial viscosity field is computed using the 3D ice temperature field given by Lieffering and Pattyn (2013) and using the values given in Cuffey and Paterson (2010) for the activation energies and prefactors used to compute the temperature-dependent Glen’s rate-factor A . This initial viscosity is then modified using inverse methods. Regarding the boundary conditions, the ice front of ice-shelves is assumed to be fix (i.e., the sum of calving flux and frontal melt flux equals the ice flux at the front). For the grounded part, the ~~friction-sliding-law~~ parameter field is ~~first~~ inferred using inverse methods ~~and a regularized friction law is then used for the relaxation phase~~ (see Sect. 2.2.1). For the floating part of the basal boundary, basal melt from the ocean is applied using the PICO ~~box~~-model (Reese et al., 2018a) and no melt is applied to partially floating mesh elements. The grounding line is determined using a flotation criterion and a sub-grid scheme is applied for the ~~friction-in-basal-drag-over~~ partially floating elements; SEP3 in Seroussi et al. (2014). Regarding the mesh, we use an anisotropic mesh adaptation scheme that uses the observed ice velocities and thickness. The mesh is preferentially refined along the directions of highest curvature of these two fields with an additional criterion function of the distance to the grounding line. The resulting mesh contains 545,837 nodes and 1,070,444 linear elements and the size varies from 1 km close to the grounding line to 50 km in the interior. The mesh is held constant during the transient simulations.

Details concerning the model initialisation, relaxation and the ~~friction-basal sliding~~ law used in the transient simulations are given below.

815 A1.2 Model inversion

To initialise the model, we use an inverse method to estimate model parameters that control the ~~friction-and-ice-viscosity-basal slipperiness and the ice rate factor~~ by minimising the misfit between observed (\mathbf{u}_{obs}) and modelled (\mathbf{u}_{mod}) velocities. ~~For the friction, we~~ We optimise β in a linear ~~friction-sliding~~ law that relates the basal shear stress, \mathbf{t}_b , to the basal sliding velocity, \mathbf{v}_b :

$$\mathbf{t}_b = -10^\beta \mathbf{v}_b = -C_{\text{eff}} \mathbf{v}_b, \quad (\text{A1})$$

820 This ensures that ~~the effective friction coefficient $C_{eff} = 10^\beta$ remains~~ $C_{eff} = 10^\beta$ is positive. For the ice rheology, the vertically averaged effective viscosity used in the SSA is written as

$$\bar{\mu} = \eta^2 \bar{\mu}_{ini} I_D^{(1-n)/n}, \quad \text{with} \quad \bar{\mu}_{ini} = 1/H \int_{z_b}^{z_s} (2A)^{-1/n} dz, \quad (\text{A2})$$

where n is the Glen exponent, I_D is the second invariant of the strain-rate tensor \mathbf{D} defined as $I_D^2 = 2\mathbf{D} : \mathbf{D}$, $\bar{\mu}_{ini}$ is the initial vertically averaged ice rigidity computed using the 3D ice temperature field given by Lieffering and Pattyn (2013) as
 825 explained above, and, to ensure that the viscosity remains positive, we optimise the parameter η starting from an initial guess of one (i.e. no modification from the viscosity initially predicted from the temperature field).

We apply a standard inverse methodology in which a cost function $J(\beta, \eta)$, which is the sum of misfit (I) and regularisation (R) terms, is minimized. The gradients of J with respect to β and η are determined in a computationally efficient way using the adjoint method. The misfit (~~I~~) ~~and regularisation~~ (~~and regularisation~~ R) terms are defined as:

$$830 \quad I = \frac{1}{2} \sum_{i=1}^{N_{\text{obs}}} \|\mathbf{u}_{\text{mod}}^i - \mathbf{u}_{\text{obs}}^i\|^2 \quad (\text{A3})$$

$$R = \lambda_1 \frac{1}{2} \int_{\Omega} (\nabla \cdot (\mathbf{u}_{\text{mod}} h) - \dot{a})^2 d\Omega + \lambda_2 \frac{1}{2} \int_{\Omega} \|\nabla \beta\|^2 d\Omega + \lambda_3 \frac{1}{2} \int_{\Omega} \|\nabla \eta\|^2 d\Omega \quad (\text{A4})$$

As the velocity observation grids might be incomplete or have a better spatial resolution than the finite-element mesh in the ice-sheet interior, the difference between the model and the observations in I (Eq. A3) is evaluated at the N_{obs} locations where
 835 observations are present. The model velocities are interpolated using the natural finite-element interpolation functions. For the regularisation term R (Eq. A4), the first term computes the misfit between the modelled flux divergence $\nabla \cdot (\mathbf{u}_{\text{mod}} h)$ and the apparent point mass balance \dot{a} integrated over the model domain Ω . Due to uncertainties in other model parameters that are not controlled during the inversion (e.g. the bed elevation), it is not possible in general to match both the velocities and the apparent point mass balance. So this term acts more as a regularisation terms that penalizes the highest ice flux divergence anomalies.
 840 The remaining anomalies are then dampened during a relaxation period, see next section. Here, for the apparent point mass balance we use the ~~parametrization~~ ~~parameterization~~ by DeConto and Pollard (2016) for the basal mass balance and RACMO for the surface mass balance and neglect the thickness rate of changes. The two other terms impose a smoothness constraint for the two inferred parameters β and η .

The regularisation weights used in this study are $\lambda_1 = 3.162 \times 10^{-5}$, $\lambda_2 = 1.259 \times 10^2$ and $\lambda_3 = 7.943 \times 10^4$. Following the
 845 principle of the L -curve analysis, they have been empirically chosen from a large set of inversions, as those that give a good compromise with a low misfit and small regularisation terms.

As the velocity data set used for the common initial state is spatially incomplete, the inversion is first performed with a mosaic that aggregates observations from 2007 to 2018 (Mouginot et al., 2019) and thus has a nearly complete spatial coverage but comes at the expense of the accuracy in areas where velocities have largely changed. The minimisation is then continued

850 for 100 iterations using the 2015-2016 dataset to get closer to those observations while staying close to the initially inverted values in areas where observations are missing.

A1.3 Relaxation

The role of the relaxation is to reduce the inconsistency between input data and inverted data when we switch from a diagnostic to a prognostic simulation (Gillet-Chaulet et al., 2012), which results in unreasonably high surface elevation rate of change.

855 The model relaxation in Elmer/Ice is divided into three different steps. During the first 5 years, the relaxation is applied with the linear sliding law used for the inversion and the inverted friction coefficient. In floating areas, the friction parameter cannot be inverted, but is set to a fixed value of $C = 1 \text{ Pa m}^{-1} \text{ a}$ to allow some friction if the grounding line was to advance. Then, to use a more realistic friction law, the regularized law from Joughin et al. (2019) (see section below) is applied for the following 15 years of relaxation. The conversion from the linear friction law to the regularized one is also described in Sect. A1.4. The
 860 last part of the relaxation is applying the mass balance [correction-modification](#) as explained in Sect. 2.2.1. It consists of 1 year of balanced flux relaxation, in which the mass balance term is defined from Eq. (1) and the [correction-term](#) $\frac{dh}{dt}|_{t_{\text{relax}}}$ [term](#) $\frac{dh}{dt}|_{\text{relax}}$ is defined as the thickness rate of change from the last time step of the previous relaxation step. This step permits to reach a near-steady state.

A1.4 Regularized Coulomb sliding law

865 For basal sliding, we adopt the regularized Coulomb sliding law proposed by Joughin et al. (2019):

$$\mathbf{t}_b = -\lambda C_{s,m} \left(\frac{\|\mathbf{v}_b\|}{\|\mathbf{v}_b\| + u_0} \right)^{1/m} \frac{\mathbf{v}_b}{\|\mathbf{v}_b\|}, \quad (\text{A5})$$

that depends on the two parameters $C_{s,m}$ and u_0 and where

$$\lambda = \begin{cases} 1, & \text{for } h_{af} \geq h_T \\ \frac{h_{af}}{h_T}, & \text{otherwise} \end{cases} \quad (\text{A6})$$

with h_{af} the height of ice above flotation and h_T a threshold height. Following Joughin et al. (2019), we adopt $m = 3$,
 870 $u_0 = 300 \text{ m a}^{-1}$ and $h_T = 75 \text{ m}$.

This [friction-sliding](#) law (Eq. A5) exhibits two asymptotic behaviours, a Weertman regime $\mathbf{t}_b = -C_{s,m}/u_0^{1/m} \|\mathbf{v}_b\|^{1/m-1} \mathbf{v}_b$ for $\|\mathbf{v}_b\| \ll u_0$ and a Coulomb regime $\mathbf{t}_b = -\lambda C_{s,m} \mathbf{v}_b / \|\mathbf{v}_b\|$ for $\|\mathbf{v}_b\| \gg u_0$. It does not include a direct dependency on effective pressure N which role is subsumed in the model parameters. In pressure-dependant [friction-sliding](#) laws like that used in Úa (Eq. A10), u_0 depends on the effective pressure. However, assuming a perfect hydrological connection between
 875 the sub-glacial drainage system and the ocean to compute N usually restricts the Coulomb regime to a small area close to the grounding line where the ice is close to flotation. Note that in this particular case, we have $N = \rho g h_{af} = \rho g h_T \lambda$, such that both [Eqs. A5 and A10](#) [Eq. \(A5\) and \(A10\)](#) have the same dependency to water pressure in the vicinity of the grounding line. As the [friction-sliding-law](#) parameter $C_{s,m}$ is determined through an inversion, it should include the dependency to N ; so that

keeping it constant through time implicitly assumes that N does not change. Because this assumption is certainly not valid as
 880 the ice column approach flotation, the factor λ imposes a linear correction to the friction when h_{af} drops below the threshold
 h_T , so that the friction decreases smoothly toward zero at the grounding line.

The inversion being done assuming a linear Weertman sliding law (Eq. A1), $C_{s,m}$ is inferred from the inverted effective
~~friction coefficient~~ C_{eff} ~~sliding-law coefficient~~ C_{eff} , such that

$$C_{s,m} = C_{eff} \|\mathbf{v}_b\| \left(\frac{\|\mathbf{v}_b\| + u_0}{\|\mathbf{v}_b\|} \right)^{1/m} / \lambda, \quad (\text{A7})$$

885 In floating areas, where ~~the friction was~~ $C_{s,m}$ ~~is~~ not constrained by the inversion, we set a constant value of $C_{s,m} = 10$ kPa.

A2 Úa

A2.1 Model description

Úa is an open source finite-element ice flow model (Gudmundsson et al., 2012; Gudmundsson, 2020). The model is based on
 a vertically integrated formulation of the momentum equations and can be used to simulate the flow of large ice sheets such
 890 as the Antarctic and Greenland ice sheets, ice caps and mountain glaciers. Úa solves the ice dynamics equations using the
 shallow-shelf approximation (SSA) (MacAyeal, 1989) and Glen's flow law (Glen, 1955). The location of the grounding line is
 determined using the flotation condition. During forward transient experiments Úa allows for fully implicit time integration,
 and the non-linear system is solved using the Newton-Raphson method. A minimum ice thickness constraint is used to ensure
 that ice thicknesses remain positive.

895 To initialise the Antarctic-wide model we take the ice extent from BedMachine Antarctica ~~v2~~ (Morlighem et al., 2020) and
 within this boundary we generated a finite-element mesh with 194,193 nodes and 385,097 linear elements using the Mesh2D
 Delaunay-based unstructured mesh generator (Engwirda, 2015). Element sizes were refined based on effective strain rates and
 distance to the grounding line, and have a maximum size of 226 km in the very interior of the ice sheet, a mean size of 3.8 km,
 a median size of 1.57 km, and a minimum size of 0.68 km. Element sizes close to the grounding line are 1 km in size. We then
 900 linearly interpolated ice surface, thickness and bed topography from BedMachine Antarctica ~~v2~~ (Morlighem et al., 2020) onto
 the model mesh. The surface boundary condition is stress-free, allowing the surface to respond freely to changes in surface
 velocity and surface mass balance. Surface mass balance is initially prescribed using output from ~~RACMO~~ [RACMO2.3p2](#)
 (van Wessem et al., 2018), and sub-shelf melt is parameterised using an implementation of the PICO ~~box~~ model (Reese et al.,
 2018a). Parameters of the basal slipperiness coefficient C in the sliding law and ice rate factor A in the flow law are determined
 905 using an inverse approach described in the following section.

A2.2 Model inversion

To initialise the model we use a data assimilation approach in which we estimate unknown parameters C and A by minimising
 the misfit between observed (u_{obs}) and modelled (u_{mod}) velocities. Observed ice velocities (Mouginot et al., 2019) were
 linearly interpolated onto the model mesh. Úa uses a standard inverse methodology in which a cost function J , which is the

910 sum of a misfit (I) and regularisation (R) term, is minimized. The gradients of J with respect to A and C are determined in a computationally efficient way using the adjoint method and Tikhonov-type regularisation. The misfit (I) and regularisation (R) terms are defined as:

$$I = \frac{1}{2\mathcal{A}} \int (u_{\text{mod}} - u_{\text{obs}})^2 / \epsilon_{\text{obs}}^2 d\mathcal{A} \quad (\text{A8})$$

915
$$R = \frac{1}{2\mathcal{A}} \int (\gamma_s^2 (\nabla \log_{10}(p/\hat{p}))^2 + \gamma_a^2 (\nabla \log_{10}(p/\hat{p}))^2) d\mathcal{A} \quad (\text{A9})$$

where $\mathcal{A} = \int d\mathcal{A}$ is the area of the model domain, ϵ_{obs} are measurement errors, and \hat{p} are the prior values for model parameters (\hat{A} and \hat{C}). We use a uniform prior $\hat{A} \approx 1.15 \times 10^{-8} \text{ kPa}^{-3} \text{ yr}^{-1}$ equivalent to an ice temperature of approx. -10°C using an Arrhenius temperature relation. For \hat{C} we estimate the prior as $\hat{C} = u_{\text{obs}}/\tau_b^m$ with $\tau_b = 80 \text{ kPa}$ and $m = 3$. The value of C beneath the ice shelves do not deviate from this prior value. Tikhonov regularisation parameters γ_s and γ_a control the slope and amplitude of the gradients in A and C . Optimum values were determined using L -curve analysis and are equal to $\gamma_s = 10000$ and $\gamma_a = 1$. The inversion was run for 10,000 iterations after which the cost function had converged.

925 Here we invert the model to estimate C using a commonly-used sliding law (Eq. A10) that relates the basal traction t_b to the horizontal components of the bed tangential basal sliding velocity \mathbf{v}_b . This was proposed by Asay-Davis et al. (2016) (Eq. 11 in that paper) and used in a recent inter-comparison experiment (Cornford et al., 2020). In our notation it reads

$$\mathbf{t}_b = - \frac{\mu_k N \mathcal{G} \beta^2 \|\mathbf{v}_b\|}{[(\mu_k N)^m + (\mathcal{G} \beta^2 \|\mathbf{v}_b\|)^m]^{1/m}} \frac{\mathbf{v}_b}{\|\mathbf{v}_b\|} \quad (\text{A10})$$

where μ_k is a coefficient of kinetic friction, and \mathcal{G} is the grounding/floating mask, with $\mathcal{G} = 1$ where the ice is grounded and $\mathcal{G} = 0$ otherwise. Here β^2 is defined as,

$$\beta^2 = C^{-1/m} \|\mathbf{v}_b\|^{1/m-1}, \quad (\text{A11})$$

930 where C is the slipperiness coefficient and we set $m = 3$. When calculating the effective pressure, N , we assume a perfect hydrological connection with the ocean, i.e. $N = \mathcal{G} \rho g (h - h_f) = \mathcal{G} \rho g h_{\text{af}}$ where h_f the flotation ice thickness (maximum ice thickness possible for a given ocean water column thickness, H , where $H = S - B$ and S is the ocean surface and B is the bedrock) or $h_f = H \rho_w / \rho$. This ensures that the basal drag approaches zeros as the grounding line is approached from above, i.e., $\mathbf{t}_b = \mathbf{0}$ at the grounding line.

935 The sliding law (Eq. A10) combines basal drag as calculated separately by the Coulomb and Weertman sliding laws through reciprocal weighting and, thus, represents a combination of those two sliding laws. Equation (A10) gives the limits,

$$\|\mathbf{t}_b\| = \begin{cases} -\mathcal{G} \beta^2 \|\mathbf{v}_b\|, & \text{for } N \rightarrow +\infty \text{ or } \|\mathbf{v}_b\| \rightarrow 0 & (\text{Weertman}) \\ -\mu_k N, & \text{for } N \rightarrow 0 \text{ or } \|\mathbf{v}_b\| \rightarrow +\infty & (\text{Coulomb}) \end{cases} \quad (\text{A12})$$

A2.3 Relaxation

Similar to Elmer/Ice as described above (Sect. A1.3), Úa requires a period of relaxation after the inversion to dampen ice flux anomalies. Here, we use a two-step, semi-iterative approach to apply a correction-modification to the mass balance term in Eq. (1). First, we take rates of thickness change that are calculated at the end of the inversion $\frac{dh}{dt}|_{inv} \sim \frac{dh}{dt}|_{inv}$ and use these as input to Eq. (1). Then, we relax the model forward in time for 1 year using this initial correction-modification to \dot{m} , after which we take rates of thickness change $\frac{dh}{dt}|_{relax} \sim \frac{dh}{dt}|_{relax}$ and use these as second correction-modification to the mass balance. This correction-term-modified mass balance field is sufficient to bring the model into a steady state, in which ice volume changes are approximately zero. This correction-term-modified mass balance field then remains fixed for all of the remaining control and perturbation simulations.

A3 PISM

We here extend on the model description of PISM. For more details on the spin-up procedure, please see Reese et al. (2022).

A3.1 Model description

The Parallel Ice Sheet Model (PISM; <https://www.pism.io>; Bueler and Brown, 2009; Winkelmann et al., 2011) is an open-source ice dynamics model developed at the University of Alaska, Fairbanks, and the Potsdam Institute for Climate Impact Research. In PISM, ice rheology Ice flow velocities are obtained from a superposition of the SSA and the shallow ice approximation (SIA) velocity fields. PISM is thermo-mecanically coupled and solves the enthalpy evolution on a three-dimensional grid. The ice rheology / rate factor is calculated taking into account the ice enthalpy and following the Glen–Paterson–Budd–Liboutry–Duval flow law (Liboutry and Duval, 1985), assuming a Glen exponent of $n = 3$ and a SSA flow enhancement factor of $E_{SSA} = 1$. The SIA flow enhancement factor is varied in the parameter ensemble. For the run presented in this manuscript, we use a value of $E_{SIA} = 2$.

Basal sliding is parameterized in the form of a generalized power-law formulation (Schoof and Hindmarsh, 2010),

$$\mathbf{t}_b = -\tau_c \frac{\mathbf{v}_b}{u_0^q \|\mathbf{v}_b\|^{1-q}}, \quad (\text{A13})$$

where \mathbf{t}_b is the basal shear stress, τ_c is the yield stress for basal till (see below), \mathbf{v}_b is the SSA basal sliding velocity, and u_0 is a threshold velocity. The sliding-law exponent $q = 1/m$ can vary between 0 (purely-plastic Coulomb sliding) and 1 (linear relationship between basal velocity and shear stress with coefficient τ_c/u_0). For the experiments presented here we adopt values of $u_0 = 100$ m/a and $q = 0.75$, respectively.

Basal shear stress in the vicinity of the grounding line is linearly interpolated on a sub-grid scale between adjacent grounded and floating grid cells according to the height above buoyancy (Feldmann et al., 2014), which allows the grounding line to evolve freely. Note that sub-shelf melt is not interpolated across the grounding line and not applied in partially floating grid cells, as usually done in PISM. The yield stress τ_c is a function of parameterized till material properties (heuristic till friction

angle ϕ) and effective till pressure N_{till} (“Mohr-Coulomb criterion”; Bueler and van Pelt, 2015):

$$\tau_c = \tan(\phi) N_{\text{till}}, \quad (\text{A14})$$

970 where N_{till} is a function of the ice overburden pressure and the modelled effective amount of water in the till layer. No connection to the ocean is assumed in the calculation of the till water content, however, the till is assumed to be fully saturated when in contact with the ocean (in grid cells with floating ice or ice-free ocean) which means that freshly grounded grid cells are usually slippery.

975 In the simulations presented here, for consistency with the other models, glacial isostatic adjustment is not considered and we only calve floating ice that extends beyond the present-day extent of Antarctic ice shelves.

980 In contrast to Úa and Elmer/Ice, PISM simulations use a finite difference scheme to solve the momentum balance on a regular grid of 8 km horizontal resolution. While Seroussi et al. (2014) report that a horizontal resolution of 2 km is required to accurately represent grounding-line dynamics, Feldmann et al. (2014) find that by using a subgrid interpolation of friction, grounding-line reversibility in PISM is also captured at coarser ($\Delta x > 10$ km) resolutions. While a higher horizontal resolution is desirable, we here employ this interpolation to be able to run PISM over millennial time scales at 8 km horizontal resolution. Despite the above mentioned differences, we find that PISM results are in line with results from Elmer/ice and Úa that employ finer resolution around the grounding lines.

Code availability. Elmer/Ice code is publicly available through GitHub (<https://github.com/ElmerCSC/elmerfem>; Gagliardini et al., 2013). All the simulations were performed with version 9.0 (Rev: 242e4bb) of Elmer/Ice. PISM code is publicly available at <https://github.com/pism/pism>. The Úa code is publicly available through Github (<https://github.com/GHilmarG/UaSource/>), and an archived version of the model can be found at <https://doi.org/10.5281/zenodo.3706623> (Gudmundsson, 2020).

Data availability. Model output data will be made publicly available on Zenodo. DOI links to the repositories will be provided upon publication.

990 *Author contributions.* The experiments presented in this paper were collectively designed by members of the TiPACCs work package 2. BU, RR and EH set up and ran the simulations for Elmer/Ice, PISM and Úa, respectively, and wrote the paper. JG created Figs. 1, 2 and 4. All authors contributed to the writing and discussion of ideas.

Competing interests. All authors declare that they have no competing interests.

Acknowledgements. This work is part of the TiPACCs project, which receives funding from the European Union's Horizon 2020 research and innovation programme under grant agreement no. 820575. The Elmer/Ice computations presented in this paper were performed using HPC resources of CINES under the allocation 2021-A0060106066 made by GENCI. We further acknowledge the European Regional Development Fund (ERDF), the German Federal Ministry of Education and Research (BMBF) and the Land Brandenburg for supporting this project by providing resources on the high-performance computer system at the Potsdam Institute for Climate Impact Research. The development of PISM is supported by NSF grants PLR-1644277 and PLR-1914668 and NASA grants NNX17AG65G and 20-CRYO2020-0052. We acknowledge the use of the Northumbria University HPC facility Oswald used to perform the Úa simulations. We like to thank Christian Schoof for providing helpful input on stability analysis.

References

- Asay-Davis, X. S., Cornford, S. L., Durand, G., Galton-Fenzi, B. K., Gladstone, R. M., Gudmundsson, G. H., Hattermann, T., Holland, D. M., Holland, D., Holland, P. R., et al.: Experimental design for three interrelated marine ice sheet and ocean model intercomparison projects: MISMIP v. 3 (MISMIP+), ISOMIP v. 2 (ISOMIP+) and MISOMIP v. 1 (MISOMIP1), *Geoscientific Model Development*, 9, 2471–2497, <https://doi.org/https://doi.org/10.5194/gmd-9-2471-2016>, 2016.
- 1005 Brondex, J., Gillet-Chaulet, F., and Gagliardini, O.: Sensitivity of centennial mass loss projections of the Amundsen basin to the friction law, *The Cryosphere*, 13, 177–195, <https://doi.org/https://doi.org/10.5194/tc-13-177-2019>, 2019.
- Bueler, E. and Brown, J.: Shallow shelf approximation as a "sliding law" in a thermomechanically coupled ice sheet model, *Journal of Geophysical Research: Solid Earth*, 114, 1–21, <https://doi.org/10.1029/2008JF001179>, 2009.
- 1010 Bueler, E. and van Pelt, W.: Mass-conserving subglacial hydrology in the Parallel Ice Sheet Model version 0.6, *Geoscientific Model Development*, 8, 1613–1635, <https://doi.org/https://doi.org/10.5194/gmd-8-1613-2015>, 2015.
- Cornford, S. L., Martin, D. F., Payne, A. J., Ng, E. G., Le Brocq, A. M., Gladstone, R. M., Edwards, T. L., Shannon, S. R., Agosta, C., Van Den Broeke, M. R., Hellmer, H. H., Krinner, G., Ligtenberg, S. R., Timmermann, R., and Vaughan, D. G.: Century-scale simulations of the response of the West Antarctic Ice Sheet to a warming climate, *Cryosphere*, 9, 1579–1600, <https://doi.org/10.5194/tc-9-1579-2015>, 1015 2015.
- Cornford, S. L., Seroussi, H., Asay-Davis, X. S., Gudmundsson, G. H., Arthern, R., Borstad, C., Christmann, J., Dias dos Santos, T., Feldmann, J., Goldberg, D., Hoffman, M. J., Humbert, A., Kleiner, T., Leguy, G., Lipscomb, W. H., Merino, N., Durand, G., Morlighem, M., Pollard, D., Rückamp, M., Williams, C. R., and Yu, H.: Results of the third Marine Ice Sheet Model Intercomparison Project (MISMIP+), *The Cryosphere*, 14, 2283–2301, <https://doi.org/10.5194/tc-14-2283-2020>, 2020.
- 1020 Cuffey, K. M. and Paterson, W. S. B.: *The physics of glaciers*, Academic Press, 2010.
- DeConto, R. M. and Pollard, D.: Contribution of Antarctica to past and future sea-level rise, *Nature*, 531, 591–597, <https://doi.org/https://doi.org/10.1038/nature17145>, 2016.
- Durand, G., Gagliardini, O., De Fleurian, B., Zwinger, T., and Le Meur, E.: Marine ice sheet dynamics: Hysteresis and neutral equilibrium, *Journal of Geophysical Research: Earth Surface*, 114, <https://doi.org/https://doi.org/10.1029/2008JF001170>, 2009.
- 1025 Engwirda, D.: Locally optimal Delaunay-refinement and optimisation-based mesh generation, in: Ph.D Thesis, School of Mathematics and Statistics, University of Sydney, <https://ses.library.usyd.edu.au/handle/2123/13148>, 2015.
- Favier, L., Durand, G., Cornford, S., Gudmundsson, G., Gagliardini, O., Gillet-Chaulet, F., Zwinger, T., Payne, A., and Le Brocq, A.: Retreat of Pine Island Glacier controlled by marine ice-sheet instability, *Nature Climate Change*, 4, 117–121, <https://doi.org/doi:10.1038/nclimate2094>, 2014.
- 1030 Feldmann, J. and Levermann, A.: Collapse of the West Antarctic Ice Sheet after local destabilization of the Amundsen Basin, *Proceedings of the national academy of sciences*, 112, 14 191–14 196, <https://doi.org/https://doi.org/10.1073/pnas.1512482112>, 2015.
- Feldmann, J., Albrecht, T., Khroulev, C., Pattyn, F., and Levermann, A.: Resolution-dependent performance of grounding line motion in a shallow model compared with a full-Stokes model according to the MISMIP3d intercomparison, *Journal of Glaciology*, 60, 353–360, <https://doi.org/https://doi.org/10.3189/2014JoG13J093>, 2014.
- 1035 Fürst, J. J., Durand, G., Gillet-Chaulet, F., Tavard, L., Rankl, M., Braun, M., and Gagliardini, O.: The safety band of Antarctic ice shelves, *Nature Climate Change*, 6, 479–482, <https://doi.org/https://doi.org/10.1038/nclimate2912>, 2016.

- Gagliardini, O., Zwinger, T., Gillet-Chaulet, F., Durand, G., Favier, L., Fleurian, B. d., Greve, R., Malinen, M., Martín, C., Råback, P., et al.: Capabilities and performance of Elmer/Ice, a new-generation ice sheet model, *Geoscientific Model Development*, 6, 1299–1318, <https://doi.org/https://doi.org/10.5194/gmd-6-1299-2013>, 2013.
- 1040 Garbe, J., Albrecht, T., Levermann, A., Donges, J. F., and Winkelmann, R.: The hysteresis of the Antarctic ice sheet, *Nature*, 585, 538–544, <https://doi.org/https://doi.org/10.1038/s41586-020-2727-5>, 2020.
- Gardner, A. S., Moholdt, G., Scambos, T., Fahnestock, M., Ligtenberg, S., Van Den Broeke, M., and Nilsson, J.: Increased West Antarctic and unchanged East Antarctic ice discharge over the last 7 years, *The Cryosphere*, 12, 521–547, <https://doi.org/https://doi.org/10.5194/tc-12-521-2018>, 2018.
- 1045 Gillet-Chaulet, F., Gagliardini, O., Seddik, H., Nodet, M., Durand, G., Ritz, C., Zwinger, T., Greve, R., and Vaughan, D. G.: Greenland ice sheet contribution to sea-level rise from a new-generation ice-sheet model, *Cryosphere*, 6, 1561–1576, <https://doi.org/10.5194/tc-6-1561-2012>, 2012.
- Glen, J. W.: The creep of polycrystalline ice, *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences*, 228, 519–538, <https://doi.org/10.1098/rspa.1955.0066>, 1955.
- 1050 Goelzer, H., Huybrechts, P., Fürst, J. J., Nick, F. M., Andersen, M. L., Edwards, T. L., Fettweis, X., Payne, A. J., and Shannon, S.: Sensitivity of Greenland ice sheet projections to model formulations, *Journal of Glaciology*, 59, 733–749, <https://doi.org/https://doi.org/10.3189/2013JoG12J182>, 2013.
- Golledge, N. R., Kowalewski, D. E., Naish, T. R., Levy, R. H., Fogwill, C. J., and Gasson, E. G.: The multi-millennial Antarctic commitment to future sea-level rise, *Nature*, 526, 421–425, 2015.
- 1055 Golledge, N. R., Clark, P. U., He, F., Dutton, A., Turney, C., Fogwill, C., Naish, T., Levy, R. H., McKay, R. M., Lowry, D. P., et al.: Retreat of the Antarctic Ice Sheet during the Last Interglaciation and implications for future change, *Geophysical Research Letters*, 48, e2021GL094513, <https://doi.org/https://doi.org/10.1038/nature15706>, 2021.
- Gudmundsson, G. H., Krug, J., Durand, G., Favier, L., and Gagliardini, O.: The stability of grounding lines on retrograde slopes, *The Cryosphere*, 6, 1497–1505, <https://doi.org/10.5194/tc-6-1497-2012>, 2012.
- 1060 Gudmundsson, H.: GHilmarG/UaSource: Ua2019b, <https://doi.org/10.5281/zenodo.3706624>, 2020.
- Haseloff, M. and Sergienko, O. V.: The effect of buttressing on grounding line dynamics, *Journal of Glaciology*, 64, 417–431, <https://doi.org/https://doi.org/10.1017/jog.2018.30>, 2018.
- Haseloff, M. and Sergienko, O. V.: Effects of calving and submarine melting on steady states and stability of buttressed marine ice sheets, *Journal of Glaciology*, pp. 1–18, 2022.
- 1065 IPCC: Summary for Policymakers, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by MassonDelmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., Cambridge University Press, 2021.
- Joughin, I., Smith, B. E., and Holland, D. M.: Sensitivity of 21st century sea level to ocean-induced thinning of Pine Island Glacier, Antarctica, *Geophysical Research Letters*, 37, <https://doi.org/https://doi.org/10.1029/2010GL044819>, 2010.
- 1070 Joughin, I., Smith, B. E., and Medley, B.: Marine ice sheet collapse potentially under way for the Thwaites Glacier Basin, West Antarctica, *Science*, 344, 735–738, <https://doi.org/10.1126/science.1249055>, 2014.
- Joughin, I., Smith, B. E., and Schoof, C. G.: Regularized Coulomb friction laws for ice sheet sliding: Application to Pine Island Glacier, Antarctica, *Geophysical research letters*, 46, 4764–4771, <https://doi.org/https://doi.org/10.1029/2019GL082526>, 2019.

- 1075 Levermann, A. and Feldmann, J.: Scaling of instability timescales of Antarctic outlet glaciers based on one-dimensional similitude analysis, *The Cryosphere*, 13, 1621–1633, <https://doi.org/https://doi.org/10.5194/tc-13-1621-2019>, 2019.
- Lhermitte, S., Sun, S., Shuman, C., Wouters, B., Pattyn, F., Wuite, J., Berthier, E., and Nagler, T.: Damage accelerates ice shelf instability and mass loss in Amundsen Sea Embayment, *Proceedings of the National Academy of Sciences*, 117, 24735–24741, <https://doi.org/https://doi.org/10.1073/pnas.1912890117>, 2020.
- 1080 Liefvering, B. V. and Pattyn, F.: Using ice-flow models to evaluate potential sites of million year-old ice in Antarctica, *Clim. Past Discuss*, 9, 2859–2887, <https://doi.org/10.5194/cpd-9-2859-2013>, 2013.
- Liboutry, L. and Duval, P.: Various isotropic and anisotropic ices found in glaciers and polar ice caps and their corresponding rheologies, *Annales geophysicae*, 3, 207 – 224, <https://doi.org/https://doi.org/10.1016/0148-9062%2885%2990267-0>, 1985.
- MacAyeal, D. R.: Large-scale ice flow over a viscous basal sediment: theory and application to ice stream B, Antarctica, *Journal of Geophysical Research*, 94, 4071–4087, <https://doi.org/10.1029/jb094ib04p04071>, 1989.
- 1085 MacAyeal, D. R.: A tutorial on the use of control methods in ice-sheet modeling, *Journal of Glaciology*, 39, 91–98, <https://doi.org/10.3189/S0022143000015744>, 1993.
- Mengel, M. and Levermann, A.: Ice plug prevents irreversible discharge from East Antarctica, *Nature Climate Change*, 4, 451–455, <https://doi.org/https://doi.org/10.1038/nclimate2226>, 2014.
- 1090 Milillo, P., Rignot, E., Rizzoli, P., Scheuchl, B., Mouginit, J., Bueso-Bello, J., Prats-Iraola, P., and Dini, L.: Rapid glacier retreat rates observed in West Antarctica, *Nature Geoscience*, pp. 1–6, <https://doi.org/https://doi.org/10.1038/s41561-021-00877-z>, 2022.
- Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., Eisen, O., Ferraccioli, F., Forsberg, R., Fretwell, P., et al.: Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet, *Nature Geoscience*, 13, 132–137, <https://doi.org/https://doi.org/10.1038/s41561-019-0510-8>, 2020.
- 1095 Mouginit, J., Rignot, E., and Scheuchl, B.: Sustained increase in ice discharge from the Amundsen Sea Embayment, West Antarctica, from 1973 to 2013, *Geophysical Research Letters*, 41, 1576–1584, <https://doi.org/https://doi.org/10.1002/2013GL059069>, 2014.
- Mouginit, J., Rignot, E., and Scheuchl, B.: Continent-wide, interferometric SAR phase, mapping of Antarctic ice velocity, *Geophysical Research Letters*, 46, 9710–9718, <https://doi.org/https://doi.org/10.1029/2019GL083826>, 2019.
- Nias, I. J., Cornford, S. L., Edwards, T. L., Gourmelen, N., and Payne, A. J.: Assessing uncertainty in the dynamical ice response to ocean warming in the Amundsen Sea Embayment, West Antarctica, *Geophysical Research Letters*, 46, 11253–11260, <https://doi.org/https://doi.org/10.1029/2019GL084941>, 2019.
- 1100 Pattyn, F. and Morlighem, M.: The uncertain future of the Antarctic Ice Sheet, *Science*, 367, 1331–1335, <https://doi.org/https://doi.org/10.1126/science.aaz5487>, 2020.
- Pattyn, F., Schoof, C., Perichon, L., Hindmarsh, R., Bueller, E., Fleurian, B. d., Durand, G., Gagliardini, O., Gladstone, R., Goldberg, D., et al.: Results of the marine ice sheet model intercomparison project, MISMIP, *The Cryosphere*, 6, 573–588, <https://doi.org/https://doi.org/10.5194/tc-6-573-2012>, 2012.
- 1105 Pattyn, F., Perichon, L., Durand, G., Favier, L., Gagliardini, O., Hindmarsh, R. C., Zwinger, T., Albrecht, T., Cornford, S., Docquier, D., et al.: Grounding-line migration in plan-view marine ice-sheet models: results of the ice2sea MISMIP3d intercomparison, *Journal of Glaciology*, 59, 410–422, <https://doi.org/https://doi.org/10.3189/2013JoG12J129>, 2013.
- 1110 Pegler, S. S.: Suppression of marine ice sheet instability, *Journal of Fluid Mechanics*, 857, 648–680, <https://doi.org/https://doi.org/10.1017/jfm.2018.742>, 2018.

- Price, S. F., Payne, A. J., Howat, I. M., and Smith, B. E.: Committed sea-level rise for the next century from Greenland ice sheet dynamics during the past decade, *Proceedings of the National Academy of Sciences*, 108, 8978–8983, <https://doi.org/https://doi.org/10.1073/pnas.1017313108>, 2011.
- 1115 Reese, R., Albrecht, T., Mengel, M., Asay-Davis, X., and Winkelmann, R.: Antarctic sub-shelf melt rates via PICO, *The Cryosphere*, 12, 1969–1985, <https://doi.org/https://doi.org/10.5194/tc-12-1969-2018>, 2018a.
- Reese, R., Gudmundsson, G. H., Levermann, A., and Winkelmann, R.: The far reach of ice-shelf thinning in Antarctica, *Nature Climate Change*, 8, 53–57, <https://doi.org/https://doi.org/10.1038/s41558-017-0020-x>, 2018b.
- Reese, R., Garbe, J., Hill, E. A., Urruty, B., Naughten, K. A., Gagliardini, O., Durand, G., Gillet-Chaulet, F., Chandler, D., Langebroek, P. M., and Winkelmann, R.: The stability of present-day Antarctic grounding lines – Part B: Possible commitment of regional collapse under current climate, *The Cryosphere Discussions*, 2022, 1–33, <https://doi.org/10.5194/tc-2022-105>, 2022.
- 1120 Rignot, E., Mouginot, J., Morlighem, M., Seroussi, H., and Scheuchl, B.: Widespread, rapid grounding line retreat of Pine Island, Thwaites, Smith, and Kohler glaciers, West Antarctica, from 1992 to 2011, *Geophysical Research Letters*, 41, 3502–3509, <https://doi.org/https://doi.org/10.1002/2014GL060140>, 2014.
- 1125 Ritz, C., Edwards, T. L., Durand, G., Payne, A. J., Peyaud, V., and Hindmarsh, R. C.: Potential sea-level rise from Antarctic ice-sheet instability constrained by observations, *Nature* 2015 528:7580, 528, 115–118, <https://doi.org/10.1038/nature16147>, 2015.
- Robel, A. A., Seroussi, H., and Roe, G. H.: Marine ice sheet instability amplifies and skews uncertainty in projections of future sea-level rise, *Proceedings of the National Academy of Sciences*, 116, 14 887–14 892, <https://doi.org/https://doi.org/10.1073/pnas.1904822116>, 2019.
- Robel, A. A., Pegler, S. S., Catania, G., Felikson, D., and Simkins, L. M.: Ambiguous stability of glaciers at bed peaks, *Journal of Glaciology*, pp. 1–8, 2022.
- 1130 Rosier, S. H., Reese, R., Donges, J. F., De Rydt, J., Gudmundsson, G. H., and Winkelmann, R.: The tipping points and early warning indicators for Pine Island Glacier, West Antarctica, *The Cryosphere*, 15, 1501–1516, <https://doi.org/https://doi.org/10.5194/tc-15-1501-2021>, 2021.
- Schmidtko, S., Heywood, K. J., Thompson, A. F., and Aoki, S.: Multidecadal warming of Antarctic waters, *Science*, 346, 1227–1231, <https://doi.org/https://doi.org/10.1126/science.1256117>, 2014.
- 1135 Schoof, C.: Ice sheet grounding line dynamics: Steady states, stability, and hysteresis, *Journal of Geophysical Research: Earth Surface*, 112, <https://doi.org/https://doi.org/10.1029/2006JF000664>, 2007.
- Schoof, C.: Marine ice sheet stability, *Journal of Fluid Mechanics*, 698, 62, <https://doi.org/https://doi.org/10.1017/jfm.2012.43>, 2012.
- Schoof, C. and Hindmarsh, R. C.: Thin-film flows with wall slip: an asymptotic analysis of higher order glacier flow models, *Quarterly journal of mechanics and applied mathematics*, 63, 73–114, <https://doi.org/https://doi.org/10.1093/qjmam/hbp025>, 2010.
- 1140 Sergienko, O. and Wingham, D.: Grounding line stability in a regime of low driving and basal stresses, *Journal of Glaciology*, 65, 833–849, 2019.
- Sergienko, O. V. and Wingham, D. J.: Bed topography and marine ice-sheet stability, *Journal of Glaciology*, 68, 124–138, 2022.
- Seroussi, H., Morlighem, M., Larour, E., Rignot, E., and Khazendar, A.: Hydrostatic grounding line parameterization in ice sheet models, *The Cryosphere*, 8, 2075–2087, <https://doi.org/https://doi.org/10.5194/tc-8-2075-2014>, 2014.
- 1145 Seroussi, H., Nakayama, Y., Larour, E., Menemenlis, D., Morlighem, M., Rignot, E., and Khazendar, A.: Continued retreat of Thwaites Glacier, West Antarctica, controlled by bed topography and ocean circulation, *Geophysical Research Letters*, 44, 6191–6199, 2017.
- van Wessem, J. M., van de Berg, W. J., Noël, B. P. Y., van Meijgaard, E., Amory, C., Birnbaum, G., Jakobs, C. L., Krüger, K., Lenaerts, J. T. M., Lhermitte, S., Ligtenberg, S. R. M., Medley, B., Reijmer, C. H., van Tricht, K., Trusel, L. D., van Ulf, L. H., Wouters, B., Wuite,

- J., and van den Broeke, M. R.: Modelling the climate and surface mass balance of polar ice sheets using RACMO2 – Part 2: Antarctica (1979–2016), *Cryosphere*, 12, 1479–1498, <https://doi.org/10.5194/tc-12-1479-2018>, 2018.
- 1150 Weertman, J.: Stability of the junction of an ice sheet and an ice shelf, *Journal of Glaciology*, 13, 3–11, <https://doi.org/https://doi.org/10.3189/S0022143000023327>, 1974.
- Wild, C. T., Alley, K. E., Muto, A., Truffer, M., Scambos, T. A., and Pettit, E. C.: Weakening of the pinning point buttressing Thwaites Glacier, West Antarctica, *The Cryosphere*, 16, 397–417, <https://doi.org/https://doi.org/10.5194/tc-16-397-2022>, 2022.
- 1155 Winkelmann, R., Martin, M. A., Haseloff, M., Albrecht, T., Bueler, E., Khroulev, C., and Levermann, A.: The Potsdam parallel ice sheet model (PISM-PIK)–Part 1: Model description, *The Cryosphere*, 5, 715–726, <https://doi.org/https://doi.org/10.5194/tc-5-715-2011>, 2011.
- Zwally, H. J., Giovinetto, M. B., Beckley, M. A., and Saba, J. L.: Antarctic and Greenland drainage systems, GSFC cryospheric sciences laboratory, http://icesat4.gsfc.nasa.gov/cryo_data/ant_grn_drainage_systems.php, 2012.