

Does combining **Combining modelled snowpack stability with machine learning** help with predicting **to predict avalanche activity**?

Léo Viallon-Galinier^{1,2,3}, Pascal Hagenmuller¹, and Nicolas Eckert²

¹Univ. Grenoble Alpes, Université de Toulouse, Météo-France, CNRS, CNRM, Centre d'Études de la Neige, Grenoble, France

²Univ. Grenoble Alpes, INRAE, CNRS, IRD, Grenoble INP, IGE, 38000 Grenoble, France

³École des Ponts, Champs-sur-Marne, France

Correspondence: Léo Viallon-Galinier (leo.viallon@meteo.fr)

Abstract. Predicting avalanche activity from meteorological and snow cover simulations is critical in mountainous areas to support operational forecasting. Several numerical and statistical methods have tried to address this issue. However, it remains unclear how combining snow physics, mechanical analysis of snow profiles and observed avalanche data improves avalanche activity prediction. This study combines extensive snow cover and snow stability simulations with observed avalanche occurrences within a Random Forest approach to predict avalanche situations at a spatial resolution corresponding to elevations and aspects of avalanche paths in a given mountain range. We develop a rigorous leave-one-out evaluation procedure including an independent **test evaluation** set, confusion matrices, and receiver operating characteristic curves. In a region of the French Alps (Haute-Maurienne) and over the period 1960-2018, we show the added value within the **statistical machine learning** model of considering advanced snow cover modelling and mechanical stability indices instead of using only simple meteorological and bulk information. Specifically, using mechanically-based stability indices and their time-derivatives in addition to simple snow and meteorological variables increases the probability of avalanche situation detection from around 65% to 76%. However, due to the scarcity of avalanche events and the possible misclassification of non-avalanche situations in the training data set, the predicted avalanche situations that are really observed remains low, around 3.4%. These scores illustrate the difficulty of predicting avalanche occurrence with a high spatio-temporal resolution, even with the current **cutting-edge** data and modelling tools. Yet, our study opens perspectives to improve modelling tools supporting operational avalanche forecasting.

Keywords: snow avalanche, machine learning, mechanical stability indices, snow cover modelling, cross-validation, avalanche forecasting

1 Introduction

Avalanches are a significant issue in mountain areas where they threaten recreationists and infrastructures (Wilhelm et al., 2000; Stethem et al., 2003). The **long-term** (Keylock et al., 1999; Eckert et al., 2010b) and **short-term** (Schweizer et al., 2020) assessments of avalanche hazards and the **mapping** (Keylock et al., 1999; Eckert et al., 2010b) and **forecasting** (Schweizer et al., 2020) of avalanche hazard and related risks are therefore important challenges for local authorities (Bründl and Margreth, 2021; Eckert and Giacona, 2022). Most of the countries facing such hazards rely on operational services for avalanche hazard forecasting (LaChapelle, 1977;

Morin et al., 2019) and hazard mapping (Eckert et al., 2018). In this work, we focus on the issue of [short-term prediction forecasting](#) (estimation of the outcomes of unseen data) of daily avalanche activity from simulated meteorological and snow data. Indeed, inferring the relation between avalanche activity and given weather and snow conditions is one of the essential components of operational avalanche hazard forecasting (prediction in the future based on predicted snow and weather conditions).

Prediction of avalanche activity is mainly based on the knowledge of the snowpack evolution and of the mechanical processes leading to avalanches (e.g. LaChapelle, 1977; Morin et al., 2019). Information on the snowpack evolution can be collected through field observations and measurements (e.g. Coléou and Morin, 2018), and numerical simulations (e.g. Bartelt and Lehning, 2002; Vionnet et al., 2012). These data typically include a detailed description of the snowpack stratigraphy with vertical profiles of snow properties (Fierz et al., 2009). Several methods allow for identifying avalanche-prone situations from these profiles. Detection of weak layers based on mechanical and expert rules, such as the so-called lemons technique ([Jamieson and Schweizer, 2005](#))([Schweizer and Jamieson, 2007](#)), comprises one qualitative approach. Numerical computation of stability indices based on mechanical theories constitutes an automated method to quantify the snowpack stability (Roch, 1966; Föhn, 1987; Lehning et al., 2004; Schweizer et al., 2006; Viallon-Galinier et al., 2021). These approaches rely on the knowledge of mechanical processes involved in avalanche release (Schweizer, 2017; Viallon-Galinier et al., 2021). Numerical models, which are currently used as an aid to decision-making for avalanche forecasters, generally combine mechanical stability indices and expert rules to provide information on snowpack stability (Morin et al., 2019; Schweizer et al., 2006; Giraud et al., 2002; Viallon-Galinier et al., 2021).

Machine learning techniques can approach the complex link between simple snow cover variables and avalanche occurrence. These methods allow taking advantage of the knowledge of past avalanche activity to determine objective delimitation of avalanche-prone conditions within the space defined by their potential drivers. The first attempt to use machine learning techniques in the avalanche community was performed using linear methods by Bois et al. (1974). In the next decade, several attempts were made to use nearest neighbors for local avalanche danger forecasting (e.g. Navarre et al., 1987; Buser, 1989). Classification trees quickly become another common choice, as it is conceptually close to decision processes used by forecasters (e.g. Kronholm et al., 2006; Hendrikx et al., 2014). The first use of random forests was performed by Mitterer and Schweizer (2013). This method become then popular in the community (e.g. Sielenou et al., 2021; Pérez-Guillén et al., 2021; Sielenou et al., 2021). Other techniques have also been tested, such as support vector machine (e.g. Pozdnoukhov et al., 2011; Choubin et al., 2019; Sielenou et al., 2021) and more advanced techniques appeared in the last years such as convolutional neural networks (e.g. Singh and Ganju, 2008; Dekanova et al., 2018).

Most existing studies use meteorological variables as input or simple bulk variables such as snow depth to feed the [statistical model](#). [These variables machine learning model](#). The first machine learning models (Navarre et al., 1987; Buser, 1989) mainly rely on meteorological observations, simple snow observations and avalanche records. The use of modelled snow information was therefore developed to complement or replace observations (e.g. Schirmer et al., 2009; Sielenou et al., 2021) and expert analyses were introduced to provide appropriate variables (Schweizer and Föhn, 1996). However, most of the commonly used variables are only surrogates for the true drivers of avalanche processes. By contrast, studies using mechanically-based variables closely related to the processes involved in avalanche formation (e.g. Viallon-Galinier et al.,

2021) are **seldom used less frequent** in machine learning approaches (e.g. Schweizer and Föhn, 1996; Mayer et al., 2022).
60 However, these variables could increase the interpretability of the algorithm results and bring complementary non-linear information readily oriented toward the prediction of avalanche activity. Hence, they may reduce the complexity of statistical tools to implement (simpler statistical relations and a smaller number of variables to consider) compared to a model that directly uses the snow model output, and improve the overall predictive power.

Existing statistical prediction approaches are difficult to compare. Different spatial extensions are considered from large
65 mountain ranges (e.g. Kronholm et al., 2006; Sielenou et al., 2021) to avalanche paths (e.g. Choubin et al., 2019). In the literature, different measures of avalanche activity are also considered from binary classes (e.g. Kronholm et al., 2006; Hendrikx et al., 2014) to ordinal multi-classes (e.g. Mosavi et al., 2020; Sielenou et al., 2021). Yet, the most important difficulty for the comparison is that existing studies do not share a common evaluation process which includes a relevant segmentation of the training and **testing evaluation** datasets and common performance metrics. This absence of a homogeneous methodology for
70 evaluating machine learning approaches within the snow and avalanche community limits the comparison between studies.

On this basis, this paper aims to determine whether combining machine learning on avalanche data and mechanical stability analysis of snow profiles helps predict avalanche activity. **In particular, we compare the prediction score of the model trained either only on meteorological and simple snow variables as input or also on variables related to the snowpack stability and derived from the full snowpack stratigraphy.** We use random forest techniques to relate meteorological, **snow conditions**
75 **modelled snowpack information** and mechanically-based stability indices to observed avalanche occurrences. We also employ time-derivatives of mechanical indices to account for short-time persistence of avalanche-prone conditions in certain cases. We eventually present a rigorous leave-one-out evaluation procedure of broad interest for evaluating avalanche prediction efficiency that includes an independent **test evaluation** set, confusion matrices, receiver operating characteristic (ROC) curves and additional scores derived from the confusion matrix. The study area is located in Haute-Maurienne in the French Alps
80 where extensive avalanche data and snow cover reanalyses over 58 years (1960-2018) are available.

2 Material and methods

2.1 Study area

We selected an area belonging to the Haute-Maurienne massif in the Northern French Alps, consisting of the three district municipalities of Bessans, Bonneval-sur-Arc and Lanslevillard (Figure 1). This area is frequently studied for avalanche-related
85 issues (e.g. Ancey et al., 2004; Eckert et al., 2009; Favier et al., 2014; Kern et al., 2021; Zgheib et al., 2020) because it is prone to intense avalanche activity **and local observations of avalanche events are particularly reliable (see below), especially over the last decades.** The area is characterized by a relatively high elevation ranging from 1400 to 3700 m, and its avalanche activity does not yet seem to be reduced by adverse climate warming effects (Lavigne et al., 2015; Zgheib et al., 2022). Located in the eastern French Alps next to the Italian border, the area experiences extreme snowfall events known as "easterly return", which drive most of the
90 avalanche activity (Eckert et al., 2010a; Le Roux et al., 2021). We considered data on the winters between 1960 and 2018. When referring to the winter season, we consider days between the 15th of October and the 15th of May. These dates are




Figure 1. Studied area. General situation on the left and contour of avalanche paths surveyed each day (EPA) in blue for the three district municipalities of our studied area (delimited with orange lines). Only the avalanches that flow below a certain threshold (blue line at the bottom of each avalanche path) are systematically reported.

consistent with the dates of production of avalanche bulletins in France and were already selected as suitable bounds in other studies (e.g. Sielenou et al., 2021).

2.2 Avalanche observations

95 Our proxy of avalanche activity relies on the *Enquête Permanente sur les Avalanches* (EPA). The EPA reports all avalanches in approximately 3,000 pre-defined paths over French mountain ranges (Bourova et al., 2016). About 110 of these are located in the studied area and are shown in Figure 1. Each avalanche record indicates the period during which the avalanche is likely to have released and some additional information, such as the elevation and the aspect of the starting zone. EPA was initially designed to capture large natural avalanche events in exposed areas and was extensively used for hazard mapping (Bourova et al., 2016). Hence, only avalanches whose run-out reaches a certain threshold (pre-identified run out threshold (defined for each avalanche path, with a threshold elevation, e.g., reaching : a road, or the valley floor, see Figure 1) are systematically recorded. The avalanche activity derived from the EPA depends on this specific sampling procedure. Moreover, it relies on human-based observations and inevitably contains some uncertainties. However, EPA remains one of the longest avalanche activity records. The selected area is characterized by a dense observation network covering a large variety of avalanche paths. Besides, the steep topography of Haute-Maurienne reduces the effect of the observation threshold as most of the avalanches reach avalanches flow far downslope, close to the valley floor, providing a representative screenshot of the overall avalanche activity in the area. Further discussion on the EPA strengths and weaknesses is out of the scope of the paper and can be found in Jomelli et al. (2007) or Eckert et al. (2013).

100

105

One of the drawbacks of this data for the current study is the uncertainty of the date of some avalanche events, which can be large for remote sites paths or during low visibility periods (28.6% of the reports have an uncertainty above one day and 23.6% above 3 days, as estimated by the observers). To associate meteorological and snow conditions to each observed avalanche, we remove observations with an uncertainty of (length of the period on which the avalanche can have occurred) of more than three days on the release date, from the dataset. When the uncertainty is larger than one day, the last day of the period was defined as the day of the avalanche event. For instance, if an observer reports that an avalanche has occurred between the 21st and 23rd of January in a given path, we consider that the uncertainty of the report is 3 days (≤ 3 days) and we arbitrarily consider that the avalanche occurred on the 23rd of January. Moreover, the aspect and the elevation of the starting zone were not reported in a few cases (representing less than 5% of the total number of events) because the starting point was not visible from the observation point or due to a lack of time for the observation. In these cases, the starting zone was defined by the average elevation and aspect of the typical release area defined for each avalanche path. We applied this definition of release day and zone to the 2779 avalanche events observed avalanches in the studied spatio-temporal domain. i

We grouped these observations into eight aspect sectors (from North to North-West) and three elevation bands (centered at 1800, 2400 and 3000 m). This choice defines the spatial resolution of our model. All observations are represented in this geometry in Figure 2. When considering all avalanche and non-avalanche situations, the avalanche situations represent 1.1% of the overall dataset. This is called the base rate and acts as a reference for further comparisons.

2.3 Simulated snowpack

The SAFRAN-SURFEX/ISBA-Crocus model chain (Durand et al., 1999; Lafaysse et al., 2013) was used to simulate the snow and meteorological conditions in the Haute-Maurienne massif. SAFRAN provides meteorological information adapting numerical weather prediction on a gridded domain to the area of interest and assimilates observed meteorological data (Durand et al., 2009). We used the publicly available reanalysis (Vernay et al., 2020). This modelling scheme assumes that meteorological conditions depend only on elevation and aspect. The SURFEX/ISBA-Crocus model is a one-dimensional snowpack model representing snowpack evolution with a multi-layered scheme based on physical evolution laws (Brun et al., 1989; Vionnet et al., 2012). It uses as an input the meteorological data from SAFRAN model, and it is coupled to the soil scheme ISBA-DIF (Decharme et al., 2011) to represent energy and mass exchange at the bottom of the snowpack. Accordingly to the spatial resolution of the avalanche observations, snow conditions are computed for eight aspects and three elevation levels (1800, 2400 and 3000 m). The temporal resolutions of the meteorological and snow conditions considered here were 1 h and 3 h, respectively.

These simulations retrieve meteorological and bulk snow conditions but also the full snowpack stratigraphy. Hence an additional step is required to take advantage of this information, which is here done through the computation of stability indices as presented right after.

2.4 Stability indices

Nine stability indices have been selected based on their applicability with our snow cover model: five for dry snow avalanches and four for wet snow avalanches. In addition, we also computed time-derivatives of these indices.

2.4.1 Dry snow indices

For dry snow, three indices are related to failure initiation, namely natural strength-stress ratio (S_n , (Föhn, 1987)), skier strength-stress ratio (S_a , (Föhn, 1987)) and external strength-stress ratio (S_r , (Reuter et al., 2015)). These indices compare shear strength to shear stress, [for a given layer interface](#), where the stress originates from the weight of the overlying layers (S_n and S_a) and/or of an external load (skier, for instance) at the top of the snowpack (S_a and S_r) (Viallon-Galinier et al., 2021). Moreover, we selected two formulations of critical crack length for representing crack propagation (Viallon-Galinier et al., 2021): the original formulation by (Heierli et al., 2008; van Herwijnen et al., 2016) and the alternative approach by Gaume et al. (2017). Both approaches require a slab modulus, determined from density according to Scapozza (2004), and fracture energy estimated from strength. Details on these indices are available in Viallon-Galinier et al. (2022).

145 These indices were computed for each layer. For each time step, based on the values of each index, we identified five weak layers (one per index). We defined a weak layer as a layer characterized by a local minimum of the considered stability index (excluding the top and the bottom layers). This approach allows identifying the five weakest layers, with five complementary ways of estimating the weakness (five stability indices). It has the advantage of providing a constant number of variables (25 variables: 5 stability indices on 5 weak layers) for further statistical analysis.

155 2.4.2 Wet snow indices

To characterize the conditions prone to wet snow avalanches, we used the mean liquid water content in the whole snowpack (Mitterer et al., 2013, 2016) and the thicknesses of humid snow layers. For the latter index, we considered a snow layer as humid as soon as its liquid water content exceeds either 0, 1 and 3% in volume. These three indices are denoted I_{h0} , I_{h1} and I_{h3} . [We also introduce the snow depth as an indicator of the amount of snow that can be involved in a potential avalanche.](#)

160

2.4.3 Time-derivatives

Stability indices values at a given time may not be sufficient to represent the avalanche activity. The time evolution of snow properties is supposed to be represented by snow cover models. However, considering snowpack properties only at a given date and disregarding its past evolution does not indicate whether the snowpack is becoming more prone to avalanches or is in a stabilization phase. For instance, low values of a stability index may indicate an avalanche-prone situation. However, if these values are preceded by even lower ones, the possible avalanche should already have occurred when the stability was minimal, or even before, but not after. Yet, few stability indices include the time dimension in the literature. To our knowledge, only Conway and Wilbour (1999) ([also used by Reuter et al. \(2022\)](#)) and MEPRA natural risk (Giraud et al., 2002; Viallon-Galinier et al., 2021) include explicit time dependence. Here, we used time derivatives of the previously defined stability indices. We defined the time-derivative of stability index f on a given weak layer as $(f(t) - f(t - dt))/dt$ with several time intervals dt (6, 24, 48, 72, 120 and 240 h). The derivatives represent 150 variables for dry snow indices and 24 variables for wet snow indices.

170




Figure 2. Number of avalanche situations recorded in our study area over the full time period at the presented spatial resolution, i.e. per elevation band (1800, 2400 and 3000 m) and aspects (8 from N to NW).

Time derivatives on snow depth were also used as a straightforward indicator of stability for dry snow conditions (accumulation of new snow) and wet snow conditions (settling and melting).

2.5 Learning procedure

175 Random forests are used to relate snow and meteorological conditions to avalanche activity in the presented spatial resolution.

2.5.1 Avalanche activity

Avalanche activity is based on EPA records in the selected area. For each day, aspect and elevation band, we classify avalanche and non-avalanche situations. A given day on given aspect sector and elevation band is considered as an avalanche situation if at least one avalanche is reported for this day (after filtering of observations and attribution of dates). All other situations are non-avalanche ones. The model resolution defined by classes of elevation and aspect is more demanding than more common approaches applied to whole mountain ranges (e.g. Hendrikx et al., 2014; Kronholm et al., 2006; Sielenou et al., 2021) but provides more detailed information, closer to the spatial resolution used in avalanche operational forecasting (Morin et al., 2019). The number of avalanche situations observed by elevation and aspect range is shown in Figure 2.

2.5.2 General overview of input variables

185 For each elevation and aspect selected, input variables used are summarized in Table 1.

These variables gather information from the meteorological SAFRAN model (Meteo), SURFEX-ISBA/Crocus (BulkSimple snow), stability indices (Stability) computed on the basis of modelled snowpack and derivatives of these variables (Derivatives)

Table 1. Variables used to predict avalanche activity using machine learning

Category	Sub-category	Name	time intervals	Number of variables
Meteo	Snowfall	Snowfall accumulation (mm)	24 and 72h	2
	Rainfall	Rainfall accumulation (mm)	24 and 72h	2
	Temperature	Min, max, mean values (K)	24 and 72h	6
	Wind	Max and mean wind speed (km/h)	24 and 72h	4
		Projected mean direction on N-S axis and E-W axis	24 and 72h	4
Bulk Simple snow	Snow depth	Snow depth (m)	—	1
	Depth of new snow	Depth (m) of snow fallen since (see intervals)	24, 72, 120h	3
Stability	Dry snow	Stability indices (S_n, S_a, S_k, a_c, a_g) for the 5 identified weak layers and depths of each weak layer	—	25
		Depth of the corresponding weak layers (m)	—	5
	Wet snow	Maximum mean liquid water content	24h	1
		Maximum height of wet snow with thresholds of 0, 1, 3% of liquid water to consider layer as wet (m)	24h	3
	Snow depth variation	Snow depth variation (m)	24, 72, 120h —	3 1
Derivatives	Dry snow indices	All dry snow indices	6, 24, 48, 72, 120, 240h	150
	Wet snow indices	All wet snow indices	6, 24, 48, 72, 120, 240h	24
	Snow depth variation	Snow depth variation (m)	24, 72, 120h	3

as described in Section 2.4. Hereafter, if no special mention is added, all these variables (All) are used but for studying variable importance, subsets of this list are also used.

190 2.5.3 Machine learning algorithm

To relate snow and meteorological conditions to avalanche activity as defined above, we used Random Forest (RF) techniques (Breiman, 2001; Hastie et al., 2009). Random Forest is an ensemble method used for classification. Each decision tree in the ensemble is built from a random subset of the data. This technique allows going beyond the limitations of single decision trees but without dramatically increasing the algorithm complexity and with similar introspection capabilities. Once trained, each
195 tree of the Random Forest predicts a class for the input data. Aggregating all trees allow to define a probability for each class as the portion of trees predicting the given class.

Random Forest classifiers require two hyper-parameters: the number of trees and the tree depth. Here, we let the trees fully grow until there is only one element in each leaf, as usually done (Hastie et al., 2009). An optimization on our full dataset showed that 3000 trees were sufficient (more trees did not improve the results), so [that](#) this value was selected for the whole
200 study.

We use two classes, namely avalanche and non-avalanche situations, that are highly unbalanced (mean of 1.1% of avalanche situations in the winter season depending on elevation and aspect, see Figure 2). Machine learning techniques, if not handled with care, do not perform well on unbalanced data (e.g. Hastie et al., 2009; Sielenou et al., 2021). They are designed to optimize the overall classification accuracy or a similar score. Their results thus tend to be biased towards the majority class (Chawla et al., 2004; Sielenou et al., 2021), here the non-avalanche situations. The most common techniques to limit this effect are
205 oversampling of the minority classes, undersampling of the majority classes or dedicated learning algorithms. We here used a combination of these techniques. We only considered situations of the winter season characterized by a simulated snow depth larger than 10 cm. This first selection led to the undersampling of the majority class. Note that we chose this conservative threshold [as the goal is not to substitute to the to remove very obvious non-avalanche situations from the dataset \(no snow in the starting zone means no avalanche\)](#). We do not expect this threshold to be optimal as this is the goal of the training phase of the machine learning algorithm, [but only to remove situations for which we are sure they were not prone to avalanche](#). However, this first step was not sufficient to fully balance the dataset. We therefore used an adaptation of RF classifier to deal with unbalanced data (Chen et al., 2004): each tree of the forest is trained on a subset of the data randomly drawn; the probability law for drawing is adapted so that the probability of drawing non-avalanche or avalanche situations are identical. This second step acts as an oversampling
215 of the minority class.

2.6 Evaluation methods

2.6.1 Evaluation process

We evaluated the model performance with a leave one year out approach (LOYO). The snowpack completely melts in summer, and new snowfall in autumn occurs on bare ground. Therefore, there is no memory between winter seasons and they are

220 exchangeable. This is not the case between successive days during the winter season, with highly correlated snowpack characteristics. A simple leave one out (i.e., leave one day out) would yield better scores but would be less relevant. For each of the 58 seasons between 1960 and 2018, a **test an evaluation** set is composed of one winter season and a learning set of the remaining 57 seasons. This leads to 58 sets of trained random forests, each one being evaluated on one year. On a single winter season, there are not enough avalanche situations to be statistically relevant. Therefore, the confusion matrix of 58 **test evaluation** years
225 were aggregated to compute scores with all information available. This leave one year out approach is used for all evaluations presented.

We also quantified the statistical uncertainty related to the sample size. As we used 58 years of evaluation data computed separately, we were able to define an uncertainty by bootstrapping evaluation years used to compute the considered score. In practice, 1000 independent draws of 58 years (with replacement) were randomly produced and the scores were computed on
230 each draw. The 20th and 80th percentiles were used to quantify the uncertainty of the produced scores.

2.6.2 Scores

The Random Forest model produces the probability of being an avalanche situation, defined as a situation with at least one avalanche event, given the snow and meteorological conditions. We selected a threshold (t) on this probability to discriminate avalanche and non-avalanche situations. It is possible to construct a confusion matrix (as presented in Table 2) based on this
235 threshold. We derived three scores from the confusion matrix. The true positive rate (TPR) or recall is the ratio between correctly predicted avalanche situations divided by the number of observed avalanche situations. **This score is also called probability of detection (POD)**. It quantifies how many avalanche situations have been correctly predicted. The false positive rate (FPR), also called false alarm ratio (FAR), is the ratio between the number of false positives (non-avalanche situations that are identified as avalanche situations) and the total number of non-avalanche situations. It corresponds to the probability that a
240 false alarm will be raised. These two complementary indicators are interesting but do not fully characterize the performance of a binary classifier in case the two classes are unbalanced (which is the case here). We used a third score to represent how many predicted avalanche situations are really observed as such. This score is called precision and is defined as the ratio between correctly predicted avalanche situations and the number of predicted avalanche situations. The definition of these scores is summarized in Table 3. **We also mention the specificity ($1 - \text{FPR}$), to be compared with the true positive rate. Finally, we
245 also compute the balanced precision, which is the precision we would have considering balanced positive and negative classes (avalanche and non avalanche situations), computed as**
$$\frac{AA}{AA+NA*(AA+AN)/(NA+NN)}$$
.

2.6.3 Scores presentation

These scores can be computed for any threshold t on the avalanche situation probability. The impact of this threshold on the overall scores can be represented with two graphs: the ROC (Receiver Operating Characteristic) curve and the precision-recall
250 graph. The ROC curve shows the true positive rate as a function of the false positive rate for all possible thresholds between 0 and 1. When the threshold is equal to 0, all situations are considered avalanche situations (true positive rate is 1, false positive rate is close to 1). When the threshold is 1, all situations are considered non-avalanche situations (true positive rate is 0 and

Table 2. Confusion matrix: observed and predicted avalanche situations ("Avalanche") and non-avalanche ones ("Non-avalanche").

		Predicted	
		Avalanche	Non-avalanche
Observed	Avalanche	AA	AN
	Non-avalanche	NA	NN

Table 3. Scores derived from the confusion matrix.

Name	expression
True positive rate or recall	$\frac{AA}{AA+AN}$
False positive rate	$\frac{NA}{NA+NN}$
Precision	$\frac{AA}{AA+NA}$

false positive rate is close to zero). A perfect classifier would have a threshold value for which the true positive rate is 1, and the false positive rate is 0. A random classifier would have values of the true positive rate close to values of the false positive rate (first bisector) **Random classification is usually associated with the diagonal in the ROC diagram**. A standard measure derived from this curve is the area between the first bisector and the ROC curve (the area under curve or AUC) (Bradley, 1997). The AUC quantifies how good the model is compared to a random classifier. We used the AUC value to compare different classifier configurations. In addition, recall is also plotted as a function of precision to capture the model capacity to identify avalanche situations (precision) while limiting the number of false positives (recall). In this graph, the optimal point would be (1,1) i.e., a 100% precision and a 100% recall.

260 2.6.4 Importance of variables

The importance of variables was estimated through the separative power of each variable in the trees by computing the normalized mean decrease of impurity (also called Gini importance) on nodes where the given variable is used to separate the data in two groups (Breiman, 2001). A variable importance of zero means that the variable could be removed without reducing model performance and a high value denotes a high separative capacity (between avalanche and non-avalanche situations) of the variable. If two variables contain similar information, each variable will be picked randomly in the tree construction and these variables will consequently share out the importance of the common information (Breiman, 2001). This first approach is commonly used with random forest but only provides a first rough insight into variable importance. We thus use a more robust discrimination of the importance of variables by using different subsets of variables (see subsets in Table 1). The performance difference between more independent groups gives an idea of the importance of the variables present in each group.




Figure 3. Random forest model output (trained with all variables) for winter 1998-1999 at 2400 m for aspects (a) NW and (b) SE. The grey bars represent the days for which avalanches were observed in the selected aspect and elevation range. The base rate of avalanche situations in the full dataset is 0.011. Dates represent the beginning of months.

270 3 Results

3.1 Overview of random forest output

The trained Random Forest model provides the probability of being an avalanche situation for each day, aspect and elevation. Figure 3 provides an overview of the output for a specific season (1998–1999), altitude elevation (2400 m) and two aspects (NW and SE). We observe a high variability of the output between days. The time series differs between aspects, which gives a rough
 275 idea of the interest of the selected spatial scale. When considering the observations, most peaks of the random forest output correspond to observed avalanche activity. The random forest thus provides, in this example, a relevant image of the expected avalanche activity. There are also false positives (such as late March in NW aspect) or false negatives (such as early February in SE). This first overview is insufficient for an evaluation of the performance of the model that must be conducted over longer periods, all aspects and elevations.

280 3.2 Model performance

The ROC curve of the model trained with all input variables at our spatial resolution and evaluated independently on each winter season since 1958 is shown in Figure 4a. Fortunately, the model is far better than a random classifier (ROC curve above the first diagonal) but it also remains far from an optimal classifier (no points close to (0, 1)). The uncertainty around the ROC curve is very low, which indicates that a sufficient amount of data is available to constrain the model and that the evaluation is




Figure 4. (a) ROC curve for of the test dataset. model trained with all input variables at our spatial resolution and evaluated independently on each winter season since 1958. The optimal point (threshold value of 0.01) is represented by a red dot. (b) Precision and recall (Table 3) curve. Shading represents the uncertainty based on the 20th and 80th percentile of the bootstrap on test evaluation years (see methods section).

285 not sensitive to the choice of the winter season. The optimal threshold, defined as the threshold which leads to the ROC point closest to (0, 1), is here 0.01. In other words, a situation is considered an avalanche situation when the model probability is larger than 0.01. For this threshold, we provide the corresponding confusion matrix in Table 4, classifying situations between observed and predicted avalanche and non-avalanche situations in all elevation and aspect bands. The corresponding scores are 75.3% for the true positive rate or recall, 23.6% for the false positive rate and 3.4% for precision. The balanced precision is 76.2%. These scores mean that more than three-quarter of the observed avalanche situations are correctly identified but avalanches were actually observed only on 3.4% of the situations when avalanches were predicted. The recall (75.3%) and sensitivity (complementary of the false positive rate, here 76.4%) are similar, indicating similar performances on observed avalanche and non-avalanche situations. An alternative point of view is to consider precision and recall rather than true and false positive rates (Figure 4b). The maximal precision that can be reached with our model is around 30% but with a very low value of recall (below 5%). With higher values of recall, the precision ranges between 2 and 10%.

290

295

3.3 Variable importance

As described in Section 2.6.4, the predictive power of the input variables can be estimated in two ways.

First, we computed the feature importance of all variables and aggregated (summed) them by groups (as defined in Table 1) (Figure 5). The most important variables are related to snow depth (Figure 5) and, in particular, the new snow amounts or snow depth variations for a time step of 72 and 120. Variables related to dry snow stability appear to also be of large importance (13.6%) but with much more variables in the corresponding group: 30 25 dry stability indices, whereas there are only four variables

300

Table 4. Confusion matrix for the **test evaluation** dataset: observed and predicted avalanche situations ("Avalanche") and non-avalanche ones ("Non-avalanche") summed over elevation and aspect ranges. A threshold value of 0.01 is used, i.e., predicted probabilities over 0.01 are considered to identify avalanche situations. The corresponding recall is 75.3%, the false positive rate is 23.6% and the precision is 3.4%.

		Predicted	
		Avalanche	Non-avalanche
Observed	Avalanche	1 895	623
	Non-avalanche	55 005	178 357




Figure 5. Feature importance (Gini importance) on train dataset, aggregated (summed) by groups of variables. The number of variables in each group is reported on the right.

for in the new snow depth group. The depths of weak layers is also of importance (3.6%). Derivatives of dry snow indices decrease in importance with time step, whereas for wet snow indices, the importance is more pronounced for a time step of 72 h. Temperature and wind are also important, even described with few involved variables. By contrast, snowfall and rainfall (on 24 h) are variables with low importance. The variability between years is limited (not shown), giving confidence in the robustness of these results. However, absolute values have to be taken with care as this analysis method is strictly valid only when the different variables are independent, which is far from the case we have here.

Second, we studied the importance of variable groups by removing the data related to different groups of variables before learning and observing changes in evaluation scores. Specifically, we selected six subsets of the presented variables (see Table 1): the meteorological variables only (Meteo), bulk variables only (Bulk), both together (Meteo+Bulk), Simple snow, stability variables without derivatives (Stability), stability variables and derivatives (Stability+Derivatives) and all variables (All). The ROC curves for all these subsets are presented in Figure 6. The associated scores for the optimal threshold are reported in




Figure 6. ROC curves of the model trained with different sets of variables. Shading represents the uncertainty by bootstrap on test evaluation years (see methods section). Labels of subsets of variables correspond to those of Table 1. Scores associated to the optimal points (nearest to (0, 1)) are reported in Table 5

Table 5. **These thresholds are coherent with the base rate of our dataset.** The ROC curve of the model trained only on meteorological variables is very close to the first bisector (Area Under the Curve AUC=0.09, Figure 6). In other words, this model is almost not much better than a random classifier. Using the **bulk simple snow** variables (snow depth and **derivatives new snow depth**) allows for a first improvement in scores with an AUC of 0.19. **As expected, adding meteorological variables to the bulk set (Meteo+Bulk) does not change the model scores significantly. Using only the 37 stability variables increases the AUC to 0.27** Using the stability variables also allows for an **AUC of 0.19** and combining it with the associated 174 time derivatives increases the AUC to 0.32, **which**. **This result highlights the importance of time dimension in avalanche activity forecasting. The AUC of 0.32 for stability and derivatives** is close to the value (AUC=0.33) obtained by using all variables. **However** **Moreover**, the uncertainty linked to inter-annual variability is larger than the difference between the two latter approaches. This means that using all stability indices and their derivatives contains all relevant information available (in the context of the variables tested in this study) for discriminating avalanche and non-avalanche situations. The other scores (false positive rate FPR, recall, precision, see Table 3) present similar trends between groups compared to AUC **except for the stability group that has a higher FPR and a lower recall compared to other groups** (. **Some differences are nevertheless observed, with for instance a higher recall but a higher FPR for stability and derivatives or all variables.** **This compared to all variables, which** highlights that the selection of an optimal classifier is always a question of compromise between these two scores.

Table 5. Predictive performance of the model trained with different sets of variables. The scores include area under ROC curve (AUC), false positive rate (FPR), recall and precision. We also report the associated optimal threshold used to compute these scores (associated to the point of the ROC curve nearest to the optimal one). Subsets of variables correspond to those of Table 1

Subset	AUC	FPR (%)	recall (%)	precision (%)	threshold
Meteo	0.009	49.9	51.7	1.1	0.025
Bulk Simple snow	0.195	32.9	65.3	2.1	0.001
Meteo and Bulk 0.211 27.4 60.0 2.3 0.008 Stability	0.273 0.188	30.9 38.1	71.8 65.9	2.4 1.8	0.01
Stability and derivatives	0.319 0.321	26.6 26.5	76.3 76.7	3.0	0.01
All	0.334	23.6	75.3	3.4	0.01

4 Discussion

4.1 Machine learning for predicting avalanche activity

330 The model performance in the studied area decomposed into eight aspects and three elevation bands, is summarized with the confusion matrix shown in Table 4. Values of recall (75.3%), false positive rate (23.6%) or precision (3.4%) may seem quite low compared to current literature. Hendrixx et al. (2005) or Kronholm et al. (2006) obtained accuracy for separation around 85% with regression trees and meteorological variables or simple **bulk snow** variables (snow depth or simple melting model). The accuracy of our model is 76.5% but this metric may not be the most informative when classes are highly unbalanced, as in our problem because it mainly gathers information on non-avalanche situations. Sielenou et al. (2021) reported scores above 95% for accuracy but did not exploit other metrics. Hendrixx et al. (2014) reported a recall (focusing on observed avalanche situations) of 76 to 79%, close to our value of 75.3%. Some studies, such as Pérez-Guillén et al. (e.g. 2021); Mayer et al. (e.g. 2022) Pérez-Guillén et al. (2021) or Mayer et al. (2022), did similar work using different targets (manually predicted avalanche hazard or measured stability) with accuracy also in comparable ranges (72 to 88%). Precision is highly influenced by the **balance between avalanche base rate (proportion of avalanche situations). Here, avalanche and non-avalanche situations, that are here highly unbalanced, but are highly unbalanced. We nevertheless consider that the balance is** representative of avalanche activity in the considered area. Moreover, low values of precision (around 3% for our model) are not uncommon for such difficult problems in related but different contexts (e.g. Rubin et al., 2012). **Eventually, to compare our results to some studies with balanced dataset, the balanced precision should be considered, which is 76.1%.**

345 However, it remains difficult to compare scores to other studies due to differences in evaluation methods and reported scores. All studies used different methods for defining a training and **a testing an evaluation** dataset. In this study, we used a robust and conservative method, consisting in isolating winter seasons for evaluation. Indeed, with the snow melting between seasons, we get rid of the snowpack memory and provide a robust separation between training and **testing evaluation** datasets, leading

to trustworthy evaluation results with our method. Moreover, we discard all the situations where the snow depth is less than
350 10 cm in the release zone, and situations outside the winter period where avalanche release is very unlikely. Consequently, our
evaluation does not include the most obvious non-avalanche situations. It is thus more strict than Sielenou et al. (2021), for
instance, who used the random forest out of bag method with oversampling of the minority class. It resembles the methodology
of Hendrikx et al. (2014) who selected two independent years for evaluation. Our method may be used for future benchmarks
to compare competing methods on a robust and homogeneous basis. In addition, the scores reported are not homogeneous
355 between studies either. Some of them focus on global accuracy (e.g. Kronholm et al., 2006; Pérez-Guillén et al., 2021), others
on accuracy per class (e.g. Sielenou et al., 2021) and a few propose other metrics such as recall, precision or F1 score (harmonic
mean of precision and recall) (e.g. Hendrikx et al., 2014). The choice of the score depends on the goal of each study and must
be adapted to it. However, limiting to a few values for summarizing the model performances limits the information available.
These differences in the evaluation processes - both separation between **test evaluation** and train sets and computed scores -
360 limit the possibility of model comparison.

Our model predicts the probability that at least one avalanche occurs on a given day within a spatial unit corresponding to one
elevation band (centred at 1800, 2400 and 3000 m) and one aspect (among 8 aspects). This spatial resolution **represents the information
currently used for avalanche forecasting in many countries enables to capture the spatial distribution of the expected avalanche activity in
one region. This latter information is crucial to evaluate and describe the avalanche danger at regional scale** (Morin et al.,
365 2019). This prediction goal is more demanding than a prediction at larger scales, as generally used in previous studies. **Indeed,
prediction at aspect-elevation resolution implies to correctly predict the avalanche activity for each aspect and elevation
band and not globally at a larger scale. For instance, if one avalanche occurs one day, it implies to identify that we have
one avalanche situation but also in which aspect and elevation sector to be considered a success. An avalanche predicted
in an other elevation or aspect will be considered as one false negative (in the elevation-aspect it really occurred) and
370 one false positive (in the elevation-aspect it was predicted).** It inevitably leads to lower performances for similar models but
provides more precise information about the spatial distribution of the avalanche hazard (Statham et al., 2018). Indeed most
studies considered avalanche activity at the scale of mountain ranges, of some thousands of km² (e.g. Kronholm et al., 2006;
Hendrikx et al., 2014; Sielenou et al., 2021; Pérez-Guillén et al., 2021). These approaches have the advantage of using machine
learning to also aggregate information at larger scales but provide a less geographically precise indicator of avalanche activity.
375 More local approaches have the advantage of providing a relation between snow and meteorological conditions and observed
or expected avalanche activity.

4.2 Added value of physical modelling of snow coverand**, stability analysis and time-derivatives for predicting avalanche activity**

We tested different input variables to train our model: meteorological variables, **bulk simple snow** variables (mainly snow
380 depth), stability indices and derivatives**of bulk variables and stability indices**. We evaluated the added value of the different groups of
variables with two different methods (described in Section 2.6.4). Meteorological information only was insufficient **to predict
avalanche activity with our method** (Figure 6). Contrarily to many other studies (e.g. Buser, 1989; Mayer et al., 2022),

we did not use observed meteorological information but large-scale modelled information (Durand et al., 2009). Thus, the meteorological information is uncertain and nearly identical for all aspects and elevations, while underlying snowpacks are generally significantly different. Therefore, we did not expect a good prediction at high spatio-temporal resolution with only meteorological information.

Most of the developed models used, at least, some basic output of snow cover models or observed snow evolution such as snow depth (e.g. Hendrikx et al., 2014). In our study, snow depth and new snow depth appeared as an essential variable in both methods used to estimate variable importance: its Gini importance is high (Figure 5) and adding it to the input variables improves a lot the model performance (Figure 6). This result is consistent with current literature identifying snow depth as the first statistical predictor for avalanche activity (e.g. Schweizer et al., 2003; Castebrunet et al., 2012; Sielenou et al., 2021). Some studies used more advanced diagnostics from snow cover models (e.g. Gassner and Brabec, 2002; Pérez-Guillén et al., 2021; Mayer et al., 2022) or computed expert aggregated variables similarly to what snow cover models do from temperature and precipitations (e.g. Kronholm et al., 2006). Snow modelling with physical models for taking into account snowpack history thus appears of high interest for automatic avalanche activity prediction.

The novelty of our model is to add a wide range of stability indices to reduce the complex information of snow cover models with the help of knowledge of physical processes. The stability indices are and combine it with a time-dependent analysis with the use of time-derivatives of stability indices. The combination of stability indices and derivatives is crucial in our random forest model (Figure 5). The time dimension have been identified as a critical information. Since the first statistical forecasts, differences between time steps, for instance on temperature (e.g. Obled and Good, 1980; Navarre et al., 1987), have been used. Conway and Wilbour (1999) also have developed a stability index that explicitly uses time derivatives. We here show that the use of time derivatives, especially in a statistical system that is not able to treat simultaneously different time steps, allows for an improvement of the prediction of avalanche activity. More generally, we showed that the introduction of stability indices and time-derivatives could help identify avalanche-prone situations with statistical machine learning models. This group of variables also gathers a great deal of information as it nearly replaces the information from other variables. This is shown by our results, which indeed, our results are quite similar when using only stability indices and their derivatives versus using all variables (Figure 6). This result indicates that stability indices combined with time-derivatives are a relevant way to summarize the information of meteorological and snow cover models for avalanche-prone conditions prediction, which is a new way of validating the interest of such stability indices.

Computing feature importance can drive the selection of relevant input variables but correlations between variables can affect the computed importance. Re-training the full model with a subset of input variables provides a robust estimation of their effective added value. In particular, the analysis of feature importance allows for selecting the right time steps for derivative computations in the wide range of possibilities included (last column of Table 1). The most important derivatives are the short-time ones (6 to 72 h) for dry snow and 72 h for wet snow (Figure 5). This result is consistent with the knowledge of involved processes (van Herwijnen et al., 2018), whereas it was never demonstrated so far from a statistical approach. The spontaneous release in dry snow occurs during or immediately after snowfall whereas wet snow problems are more linked to the progressive wetting of the snowpack with solar radiations (time scales of one to several days) or rain (e.g. Reuter et al., 2022). Variable

importance allows for selecting the most relevant variables which may be kept for further work, especially on stability variables and derivatives, which our results prove to be of interest (Figure 6).

420 4.3 Other advantages and disadvantages of our approach

We used the EPA as the ground truth of avalanche activity. This dataset is unique in its spatial and temporal extension but mainly focuses on large avalanches often reaching valley floors. In consequence, the high-elevation avalanche activity and smaller avalanches are not reported, which leads to a limited number of avalanche situations in the dataset. Yet, for the spatio-temporal domain selected in this study (Haute-Maurienne, 1960-2018), the number of avalanche events reported in the EPA
425 remains large enough (2518 avalanche situations). The local topography with steep slopes and the repartition of the recorded avalanche path allow for a **representative reasonable** screenshot of the **overall** avalanche activity. However, the scarcity of reported avalanche events might become a problem in other regions **as our balancing methods may become insufficient**. Observation may not be possible every day (e.g., poor visibility or remote sites), and only avalanches are reported (i.e., no information on the observation that no avalanche occurred). This means that the dataset does not allow to clearly define non-avalanche
430 situations: some situations may be identified as non-avalanche situations, while an avalanche occurred but was not reported. The data also suffers from uncertainty on the dates of avalanches. This may reduce the obtained score. Other data sources may be used to complement avalanche observation dataset, such as observations from ski resorts (e.g. Giard et al., 2018) or satellite avalanche detection (e.g. Karas et al., 2022), but no other data source has the temporal extension of EPA, except archival data that require in-depth investigations which cannot always be undertaken (Giacona et al., 2017, 2021).

435 Moreover, we here trained the model with the Haute-Maurienne data. Some climatological or terrain features may lead to a predicted avalanche activity specific to the Haute-Maurienne area, especially with a higher sensitivity of certain aspects or elevations (**eastern crests e.g.**, during easterly returns). Hence, the model may not be transferable directly to other areas without a new calibration. Finally, this study presents a binary classification as there is rarely more than one avalanche per day and spatial unit (aspect-elevation), which limits the definition of several classes of avalanche activity. In the future, such machine learning
440 techniques may benefit from the use of other sources of data to complement EPA data and identify more avalanches, such as remote sensing (e.g. Karas et al., 2022), infrasound (Mayer et al., 2020) or seismic detection (van Herwijnen and Schweizer, 2011).

In this study, we chose to treat all avalanche types in a single learning process, including dry and wet avalanches. Some previous studies separate different avalanche activities on pre-defined time periods (e.g. Obléd and Good, 1980), or by type
445 of avalanches, restricting to dry or wet avalanches (e.g. Mayer et al., 2022; Pérez-Guillén et al., 2022). If we assume that decision trees (or here, Random Forest) can capture dry avalanche activity on one hand and wet avalanche activity on the other hand, and if we provide information to discriminate between situations, such as liquid water content or height of wet snow in this study, then a decision tree (or an ensemble of them) will be able to be optimized on the overall avalanche activity by introducing a split in the overall tree to distinguish between dry and wet situations, if relevant. Some other studies also mix
450 dry and wet avalanches, such as the MEPRA French operational avalanche activity indicator (Giraud et al., 2002). Moreover, the observation dataset does not always allow to infer the processes that led to the avalanche and some situations may remain

uncertain in case of a mix of dry and wet snow in the snowpack. For the forecasters, complementary information may be provided with additional tools to identify the processes or situations involved, such as Reuter et al. (2022).

455 The **interest of physics to predict avalanche activity impact of using physically-based indices of snow stability as predictors of avalanche activity instead of simpler variables** was studied through a specific statistical tool, namely random forests. This method is popular due to its simple background (decision trees (Breiman et al., 1984)) which allows for in-depth analysis and interpretation to some extent and its capacity to represent non-linear phenomena (e.g. Sielenou et al., 2021; Pérez-Guillén et al., 2021; Mayer et al., 2022). Many other statistical methods are available but random forests have been shown to be as relevant as other ones (e.g. Sielenou et al., 2021). We introduced time-derivatives and cumulative values to represent the importance of history for
460 snowpack-related processes. Methods in the range of recurrent neural networks are specifically designed to cope with processes having a memory of previous states (Hochreiter and Schmidhuber, 1997). **This kind of statistical method could be These alternative statistical methods could be further** compared to our random forest approach **to . It may provide improvements in the prediction scores or** strengthen our results on the effectiveness of combining snow physics and machine learning for predicting avalanche activity.

465 Our results were obtained with a reanalysis of meteorological and snow conditions, that is to say, input data that have been retroactively corrected with all available observations. This may not be completely representative of operational forecasting (prediction in the future) situation in which models are corrected by observations of the past but run unconstrained for the forecast. This transposition to the forecasting context would be the next step in terms of complexity for machine learning methods. However, the use of the reanalysis allows for a better evaluation of the capabilities of the machine learning model
470 with fewer input errors, which was the goal of this paper.

5 Conclusion and outlooks

This paper combines snow cover modelling, mechanical stability indices and observational data through machine learning for avalanche **activity** prediction. In particular, we considered numerous stability indices and their time-derivatives. To evaluate the random forest model, we defined a robust method adapted to the specific behaviour of the snowpack (long-term memory).
475 This evaluation was conducted on three district municipalities of the French Alps with 58 years of a comprehensive dataset of avalanche observations, with a high spatial resolution (8 aspects and 3 elevation ranges) and an extended set of variables describing both meteorological, mechanical stability variables and their time evolution.

The combination of snow physics through snowpack modelling, stability indices and their derivatives, and random forest proves to be **validuseful for avalanche activity prediction**. The snow depth **remains and new snow depth remain** the most important **predictor predictors** but this study highlights the interest in using mechanical stability indices and their derivatives. This is the primary finding of our research as this had never been demonstrated with such a large variety of indices and their derivatives in previous studies (Kronholm et al., 2006; Hendrixx et al., 2014; Sielenou et al., 2021, e.g.)**(e.g. Zeidler and Jamieson, 2004; Kronholm et al., 2006; Hendrixx et al., 2014; Sielenou et al., 2021)**, even the rare ones using simple stability indices within **statistical machine**

learning models (Mayer et al., 2022). Our results also more generally underline the interest of physically-based snow cover models
485 and stability indices for identifying avalanche-prone conditions.

Obtained scores of recall (75.3%), false positive rate (23.6%) and recall precision (3.4%) are consistent with current literature
with similar goals and methods. These scores illustrate the difficulty to predict avalanche occurrence with high spatio-temporal
resolution, even with the cutting-edge data and modelling tools currently available. Moreover, we used a rather strict evaluation
method leading to lower but more representative scores compared robust and conservative scores, which are not directly comparable to
490 other studies (e.g. Sielenou et al., 2021). Hence, this method may be seen as the first step for future formal comparison between
approaches. More widely, with its high spatio-temporal resolution and use of cutting-edge physical and mechanical models, our
study opens the perspective to improve modelling tools supporting operational avalanche forecasting.

We here focus on the avalanche activity reported by EPA. The method may be extended in the future to other target variables
describing more precisely avalanche hazard such as release volumes or typical situations (Schweizer et al., 2003; Statham et al.,
495 2018; Reuter et al., 2022; Mayer et al., 2022). Similarly, we used meteorological reanalysis for snow modelling for the quality
of the data but this may not be completely representative of forecast conditions and tests have to be conducted with re-forecasts
rather than reanalysis.

References

- 500 Ancey, C., Gervasoni, C., and Meunier, M.: Computing extreme avalanches, *Cold Regions Science and Technology*, 39, 161–180, <https://doi.org/10.1016/j.coldregions.2004.04.004>, 2004.
- Bartelt, P. and Lehning, M.: A physical SNOWPACK model for the Swiss avalanche warning: Part I: numerical model, *Cold Regions Science and Technology*, 35, 123 – 145, [https://doi.org/10.1016/S0165-232X\(02\)00074-5](https://doi.org/10.1016/S0165-232X(02)00074-5), 2002.
- Bois, P., Obléd, C., and Good, W.: Multivariate data analysis as a tool for day-by-day avalanche forecast, in: *Snow Mechanics Symposium*, 1974.
- 505 Bourova, E., Maldonado, E., Leroy, J., Alouani, R., Eckert, N., Bonnefoy-Demongeot, M., and Deschatres, M.: A new web-based system to improve the monitoring of snow avalanche hazard in France, *Natural Hazards and Earth System Sciences*, 16, 1205–1216, <https://doi.org/10.5194/nhess-16-1205-2016>, 2016.
- Bradley, A. P.: The use of the area under the ROC curve in the evaluation of machine learning algorithms, *Pattern Recognition*, 30, 1145–1159, [https://doi.org/10.1016/s0031-3203\(96\)00142-2](https://doi.org/10.1016/s0031-3203(96)00142-2), 1997.
- 510 Breiman, L.: Random Forests, *Machine Learning*, 45, 5–32, <https://doi.org/10.1023/a:1010933404324>, 2001.
- Breiman, L., Friedman, J. H., Olshen, R. A., and Stone, C. J.: *Classification and regression trees*, Cole Advanced Books and Software, 1984.
- Brun, E., Martin, E., Simon, V., Gendre, C., and Coleou, C.: An Energy and Mass Model of Snow Cover Suitable for Operational Avalanche Forecasting, *Journal of Glaciology*, 35, 333–342, <https://doi.org/10.3189/S0022143000009254>, 1989.
- Bründl, M. and Margreth, S.: Integrative risk management: The example of snow avalanches, in: *Snow and Ice-Related Hazards, Risks, and Disasters*, pp. 259–296, Elsevier, <https://doi.org/10.1016/b978-0-12-817129-5.00002-0>, 2021.
- 515 Buser, O.: Two Years Experience of Operational Avalanche Forecasting using the Nearest Neighbours Method, *Annals of Glaciology*, 13, 31–34, <https://doi.org/10.3189/s026030550000759x>, 1989.
- Castebrunet, H., Eckert, N., and Giraud, G.: Snow and weather climatic control on snow avalanche occurrence fluctuations over 50 yr in the French Alps, *Climate of the Past*, 8, 855–875, <https://doi.org/10.5194/cp-8-855-2012>, 2012.
- 520 Chawla, N. V., Japkowicz, N., and Kotcz, A.: Editorial, *ACM SIGKDD Explorations Newsletter*, 6, 1–6, <https://doi.org/10.1145/1007730.1007733>, 2004.
- Chen, C., Liaw, A., Breiman, L., et al.: Using random forest to learn imbalanced data, University of California, Berkeley, 110, 24, <https://statistics.berkeley.edu/sites/default/files/tech-reports/666.pdf>, 2004.
- Choubin, B., Borji, M., Mosavi, A., Sajedi-Hosseini, F., Singh, V. P., and Shamshirband, S.: Snow avalanche hazard prediction using machine learning methods, *Journal of Hydrology*, 577, 123 929, <https://doi.org/10.1016/j.jhydrol.2019.123929>, 2019.
- 525 Coléou, C. and Morin, S.: Vingt-cinq ans de prévision du risque d’avalanche à Météo-France, *La Météorologie*, 100, 79–84, <https://doi.org/10.4267/2042/65147>, 2018.
- Conway, H. and Wilbour, C.: Evolution of snow slope stability during storms, *Cold Regions Science and Technology*, 30, 67 – 77, [https://doi.org/10.1016/S0165-232X\(99\)00009-9](https://doi.org/10.1016/S0165-232X(99)00009-9), 1999.
- 530 Decharme, B., Boone, A., Delire, C., and Noilhan, J.: Local evaluation of the Interaction between Soil Biosphere Atmosphere soil multilayer diffusion scheme using four pedotransfer functions, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2011JD016002>, 2011.
- Dekanova, M., Duchon, F., Dekan, M., Kyzek, F., and Biskupic, M.: Avalanche forecasting using neural network, in: *2018 ELEKTRO, IEEE*, <https://doi.org/10.1109/elektro.2018.8398359>, 2018.

- 535 Durand, Y., Giraud, G., Brun, E., Mérindol, L., and Martin, E.: A computer-based system simulating snowpack structures as a tool for regional avalanche forecasting, *Journal of Glaciology*, 45, 469–484, <https://doi.org/10.3189/S002214300001337>, 1999.
- Durand, Y., Laternser, M., Giraud, G., Etchevers, P., Lesaffre, B., and Mérindol, L.: Reanalysis of 44 Yr of Climate in the French Alps (1958–2002): Methodology, Model Validation, Climatology, and Trends for Air Temperature and Precipitation, *Journal of Applied Meteorology and Climatology*, 48, 429–449, <https://doi.org/10.1175/2008JAMC1808.1>, 2009.
- 540 Eckert, N. and Giacona, F.: Towards a holistic paradigm for long-term snow avalanche risk assessment and mitigation, *Ambio*, pp. 1–22, 2022.
- Eckert, N., Parent, E., Faug, T., and Naaim, M.: Bayesian optimal design of an avalanche dam using a multivariate numerical avalanche model, *Stochastic Environmental Research and Risk Assessment*, 23, 1123–1141, <https://doi.org/10.1007/s00477-008-0287-6>, 2009.
- Eckert, N., Coleou, C., Castebrunet, H., Deschatres, M., Giraud, G., and Gaume, J.: Cross-comparison of meteorological and avalanche data
545 for characterising avalanche cycles: The example of December 2008 in the eastern part of the French Alps, *Cold Regions Science and Technology*, 64, 119–136, <https://doi.org/10.1016/j.coldregions.2010.08.009>, 2010a.
- Eckert, N., Naaim, M., and Parent, E.: Long-term avalanche hazard assessment with a Bayesian depth-averaged propagation model, *Journal of Glaciology*, 56, 563–586, <https://doi.org/10.3189/002214310793146331>, 2010b.
- Eckert, N., Keylock, C., Castebrunet, H., Lavigne, A., and Naaim, M.: Temporal trends in avalanche activity in the French
550 Alps and subregions: from occurrences and runout altitudes to unsteady return periods., *Journal of Glaciology*, 59, 93 – 114, <https://doi.org/10.3189/2013JoG12J091>, 2013.
- Eckert, N., Naaim, M., Giacona, F., Favier, P., Lavigne, A., Richard, D., Bourrier, F., and Parent, E.: Repenser les fondements du zonage réglementaire des risques en montagne « récurrents », *La Houille Blanche*, 104, 38–67, <https://doi.org/10.1051/lhb/2018019>, 2018.
- Favier, P., Eckert, N., Bertrand, D., and Naaim, M.: Sensitivity of avalanche risk to vulnerability relations, *Cold Regions Science and
555 Technology*, 108, 163–177, <https://doi.org/10.1016/j.coldregions.2014.08.009>, 2014.
- Fierz, C., R.L. A., Y. D., P. E., Greene, E., Mcclung, D., Nishimura, K., Satyawali, P., and Sokratov, S.: The International Classification for Seasonal Snow on the Ground, *IHP-VII Technical Documents in Hydrology*, 83, <https://unesdoc.unesco.org/ark:/48223/pf0000186462>, 2009.
- Föhn, P. M.: The stability index and various triggering mechanisms, *IAHS publication*, 162, 195–214, https://iahs.info/uploads/dms/iahs_162_0195.pdf, 1987.
560
- Gassner, M. and Brabec, B.: Nearest neighbour models for local and regional avalanche forecasting, *Natural Hazards and Earth System Science*, 2, 247–253, <https://doi.org/10.5194/nhess-2-247-2002>, 2002.
- Gaume, J., Van Herwijnen, A., Chambon, G., Wever, N., and Schweizer, J.: Snow fracture in relation to slab avalanche release: critical state for the onset of crack propagation, *The Cryosphere*, 11, 217–228, <https://doi.org/10.5194/tc-11-217-2017>, 2017.
- 565 Giacona, F., Eckert, N., and Martin, B.: A 240-year history of avalanche risk in the Vosges Mountains based on non-conventional (re) sources, *Natural Hazards and Earth System Sciences*, 17, 887–904, 2017.
- Giacona, F., Eckert, N., Corona, C., Mainieri, R., Morin, S., Stoffel, M., Martin, B., and Naaim, M.: Upslope migration of snow avalanches in a warming climate, *Proceedings of the National Academy of Sciences*, 118, e2107306 118, 2021.
- Giard, D., Poli, P., Morin, S., Cohuet, J.-B., Marin, F., Souverain, C., Coléou, C., Regimbeau, A., and Créau, M.: L'approche participative au
570 service de l'observation météorologique, *La Météorologie*, p. 105, <https://doi.org/10.4267/2042/65152>, 2018.
- Giraud, G., Navarre, J.-P., and Coléou, C.: Estimation du risque avalancheux dans le système expert MEPRa, *Tech. rep.*, CNRM - Centre national de recherches météorologiques, <https://hal.archives-ouvertes.fr/hal-02638748>, unpublished work, 2002.

- Hastie, T., Tibshirani, R., and Friedman, J.: *The Elements of Statistical Learning*, Springer New York, <https://doi.org/10.1007/978-0-387-84858-7>, 2009.
- 575 Heierli, J., Gumbsch, P., and Zaiser, M.: Anticrack Nucleation as Triggering Mechanism for Snow Slab Avalanches, *Science*, 321, 240–243, <https://doi.org/10.1126/science.1153948>, 2008.
- Hendrikx, J., Owens, I., Carran, W., and Carran, A.: Avalanche activity in an extreme maritime climate: The application of classification trees for forecasting, *Cold Regions Science and Technology*, 43, 104–116, <https://doi.org/10.1016/j.coldregions.2005.05.006>, 2005.
- Hendrikx, J., Murphy, M., and Onslow, T.: Classification trees as a tool for operational avalanche forecasting on the Seward Highway, Alaska, *Cold Regions Science and Technology*, 97, 113 – 120, <https://doi.org/10.1016/j.coldregions.2013.08.009>, 2014.
- 580 Hochreiter, S. and Schmidhuber, J.: Long short-term memory, *Neural computation*, 9, 1735–1780, <https://doi.org/10.1162/neco.1997.9.8.1735>, 1997.
- Jamieson, B. and Schweizer, J.: *Using a checklist to assess manual snow profiles*, *Avalanche News*, 72, 57–61, 2005.
- Jomelli, V., Delval, C., Grancher, D., Escande, S., Brunstein, D., Hetu, B., Filion, L., and Pech, P.: Probabilistic analysis of recent snow avalanche activity and weather in the French Alps, *Cold Regions Science and Technology*, 47, 180–192, <https://doi.org/10.1016/j.coldregions.2006.08.003>, 2007.
- 585 Karas, A., Karbou, F., Giffard-Roisin, S., Durand, P., and Eckert, N.: Automatic Color Detection-Based Method Applied to Sentinel-1 SAR Images for Snow Avalanche Debris Monitoring, *IEEE Transactions on Geoscience and Remote Sensing*, 60, 1–17, <https://doi.org/10.1109/tgrs.2021.3131853>, 2022.
- 590 Kern, H., Eckert, N., Jomelli, V., Grancher, D., Deschatres, M., and Arnaud-Fassetta, G.: Brief communication: Weak control of snow avalanche deposit volumes by avalanche path morphology, *The Cryosphere*, 15, 4845–4852, <https://doi.org/10.5194/tc-15-4845-2021>, 2021.
- Keylock, C. J., McClung, D. M., and Magnússon, M. M.: Avalanche risk mapping by simulation, *Journal of Glaciology*, 45, 303–314, <https://doi.org/10.3189/S0022143000001805>, 1999.
- 595 Kronholm, K., Vikhamar-Schuler, D., Jaedicke, C., Isaksen, K., Sorteberg, A., and Kristensen, K.: Forecasting snow avalanche days from meteorological data using classification trees; Grasdalen, Western Norway, in: *Proceedings of the International Snow Science Workshop*, Telluride, Colorado, pp. 1–6, Citeseer, <https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.855.5195&rep=rep1&type=pdf>, 2006.
- LaChapelle, E. R.: *Snow Avalanches: A review of Current Research and Applications*, *Journal of Glaciology*, 19, 313–324, <https://doi.org/10.3189/s0022143000215633>, 1977.
- 600 Lafaysse, M., Morin, S., Coléou, C., Vernay, M., Serça, D., Besson, F., Willemet, J.-M., Giraud, G., Durand, Y., and Météo-France, D.: Towards a new chain of models for avalanche hazard forecasting in French mountain ranges, including low altitude mountains, in: *Proceedings of International Snow Science Workshop Grenoble–Chamonix Mont-Blanc*, vol. 7, pp. 162–166, http://arc.lib.montana.edu/snow-science/objects/ISSW13_paper_O1-02.pdf, 2013.
- Lavigne, A., Eckert, N., Bel, L., and Parent, E.: Adding expert contributions to the spatiotemporal modelling of avalanche activity under different climatic influences, *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 64, 651–671, <https://doi.org/10.1111/rssc.12095>, 2015.
- 605 Le Roux, E., Evin, G., Eckert, N., Blanchet, J., and Morin, S.: Elevation-dependent trends in extreme snowfall in the French Alps from 1959 to 2019, *The Cryosphere*, 15, 4335–4356, <https://doi.org/10.5194/tc-15-4335-2021>, 2021.
- Lehning, M., Fierz, C., Brown, B., and Jamieson, B.: Modeling snow instability with the snow-cover model SNOWPACK, *Annals of Glaciology*, 38, 331–338, <https://doi.org/10.3189/172756404781815220>, 2004.
- 610

- Mayer, S., van Herwijnen, A., Olivieri, G., and Schweizer, J.: Evaluating the performance of an operational infrasound avalanche detection system at three locations in the Swiss Alps during two winter seasons, *Cold Regions Science and Technology*, 173, 102962, <https://doi.org/10.1016/j.coldregions.2019.102962>, 2020.
- 615 Mayer, S., van Herwijnen, A., Techel, F., and Schweizer, J.: A random forest model to assess snow instability from simulated snow stratigraphy, *The Cryosphere Discussions*, <https://doi.org/10.5194/tc-16-4593-2022>, 2022.
- Mitterer, C. and Schweizer, J.: Analysis of the snow-atmosphere energy balance during wet-snow instabilities and implications for avalanche prediction, *The Cryosphere*, 7, 205–216, <https://doi.org/10.5194/tc-7-205-2013>, 2013.
- Mitterer, C., Techel, F., Fierz, C., and Schweizer, J.: An operational supporting tool for assessing wet-snow avalanche danger, in: *Proceedings ISSW*, vol. 33, https://www.researchgate.net/profile/Frank_Techel/publication/261672459_An_operational_supporting_tool_for_assessing_wet-snow_avalanche_danger/links/00b49534f63d14b4ae000000/An-operational-supporting-tool-for-assessing-wet-snow-avalanche-danger.pdf, 2013.
- 620 An-operational-supporting-tool-for-assessing-wet-snow-avalanche-danger.pdf, 2013.
- Mitterer, C., Heilig, A., Schmid, L., van Herwijnen, A., Eisen, O., and Schweizer, J.: Comparison of measured and modelled snow cover liquid water content to improve local wet-snow avalanche prediction, in: *International Snow Science Workshop Proceedings*, https://arc.lib.montana.edu/snow-science/objects/ISSW16_O6.01.pdf, 2016.
- 625 Morin, S., Horton, S., Techel, F., Bavay, M., Coléou, C., Fierz, C., Gobiet, A., Hagenmuller, P., Lafaysse, M., Ližar, M., Mitterer, C., Monti, F., Müller, K., Olefs, M., Snook, J. S., van Herwijnen, A., and Vionnet, V.: Application of physical snowpack models in support of operational avalanche hazard forecasting: A status report on current implementations and prospects for the future, *Cold Regions Science and Technology*, p. 102910, <https://doi.org/https://doi.org/10.1016/j.coldregions.2019.102910>, 2019.
- Mosavi, A., Shirzadi, A., Choubin, B., Taramideh, F., Hosseini, F. S., Borji, M., Shahabi, H., Salvati, A., and Dineva, A. A.: Towards an Ensemble Machine Learning Model of Random Subspace Based Functional Tree Classifier for Snow Avalanche Susceptibility Mapping, *IEEE Access*, 8, 145 968–145 983, <https://doi.org/10.1109/access.2020.3014816>, 2020.
- 630 Ensemble Machine Learning Model of Random Subspace Based Functional Tree Classifier for Snow Avalanche Susceptibility Mapping, *IEEE Access*, 8, 145 968–145 983, <https://doi.org/10.1109/access.2020.3014816>, 2020.
- Navarre, J., Guyomar'ch, G., and Giraud, G.: Un modèle statistique pour la prévision locale des avalanches, *IAHS-Publ*, 162, 571–580, *Unmod{è}lestatistiquepourlapr{é}visionlocaledesavalanches*, 1987.
- Obled, C. and Good, W.: Recent Developments of Avalanche Forecasting by Discriminant Analysis Techniques: A Methodological Review and Some Applications to the Parsenn Area (Davos, Switzerland), *Journal of Glaciology*, 25, 315–346, <https://doi.org/10.3189/S0022143000010522>, 1980.
- 635 Recent Developments of Avalanche Forecasting by Discriminant Analysis Techniques: A Methodological Review and Some Applications to the Parsenn Area (Davos, Switzerland), *Journal of Glaciology*, 25, 315–346, <https://doi.org/10.3189/S0022143000010522>, 1980.
- Pérez-Guillén, C., Hendrick, M., Techel, F., van Herwijnen, A., Volpi, M., Tasko, O., Pérez-Cruz, F., Obozinski, G., and Schweizer, J.: Data-driven automatic predictions of avalanche danger in Switzerland, in: *EGU General Assembly*, Copernicus GmbH, <https://doi.org/10.5194/egusphere-egu21-6154>, 2021.
- 640 Data-driven automatic predictions of avalanche danger in Switzerland, in: *EGU General Assembly*, Copernicus GmbH, <https://doi.org/10.5194/egusphere-egu21-6154>, 2021.
- Pozdnoukhov, A., Matasci, G., Kanevski, M., and Purves, R. S.: Spatio-temporal avalanche forecasting with Support Vector Machines, *Natural Hazards and Earth System Sciences*, 11, 367–382, <https://doi.org/10.5194/nhess-11-367-2011>, 2011.
- Pérez-Guillén, C., Techel, F., Hendrick, M., Volpi, M., van Herwijnen, A., Olevski, T., Obozinski, G., Pérez-Cruz, F., and Schweizer, J.: Evaluation of an automated data-driven decision support tool for operational avalanche forecasting in Switzerland, in: *IGS Symposium on Snow*, 2022.
- 645 Evaluation of an automated data-driven decision support tool for operational avalanche forecasting in Switzerland, in: *IGS Symposium on Snow*, 2022.
- Reuter, B., Schweizer, J., and van Herwijnen, A.: A process-based approach to estimate point snow instability, *The Cryosphere*, 9, 837–847, <https://doi.org/10.5194/tc-9-837-2015>, 2015.

- Reuter, B., Viallon-Galinier, L., Horton, S., van Herwijnen, A., Mayer, S., Hagenmuller, P., and Morin, S.: Characterizing snow instability with avalanche problem types derived from snow cover simulations, *Cold Regions Science and Technology*, 194, 103462, <https://doi.org/10.1016/j.coldregions.2021.103462>, 2022.
- 650 Roch, A.: Les déclenchements d'avalanche, *IAHS Publ.*, 69, 86–99, <https://iahs.info/uploads/dms/069021.pdf>, 1966.
- Rubin, M. J., Camp, T., Herwijnen, A. V., and Schweizer, J.: Automatically Detecting Avalanche Events in Passive Seismic Data, in: 2012 11th International Conference on Machine Learning and Applications, IEEE, <https://doi.org/10.1109/icmla.2012.12>, 2012.
- Scapozza, C.: Entwicklung eines dichte-und temperaturabhängigen Stoffgesetzes zur Beschreibung des visko-elastischen Verhaltens von Schnee, Ph.D. thesis, ETH Zurich, <https://doi.org/10.3929/ethz-a-004680249>, 2004.
- 655 Schirmer, M., Lehning, M., and Schweizer, J.: Statistical forecasting of regional avalanche danger using simulated snow-cover data, *Journal of Glaciology*, 55, 761–768, <https://doi.org/10.3189/002214309790152429>, 2009.
- Schweizer, J.: On recent advances in avalanche research, *Cold Regions Science and Technology*, 144, 1 – 5, <https://doi.org/10.1016/j.coldregions.2017.10.014>, international Snow Science Workshop 2016 Breckenridge, 2017.
- Schweizer, J. and Föhn, P. M. B.: Avalanche forecasting — an expert system approach, *Journal of Glaciology*, 42, 318–332, <https://doi.org/10.3189/s0022143000004172>, 1996.
- 660 Schweizer, J. and Jamieson, J. B.: A threshold sum approach to stability evaluation of manual snow profiles, *Cold Regions Science and Technology*, 47, 50–59, <https://doi.org/10.1016/j.coldregions.2006.08.011>, 2007.
- Schweizer, J., Bruce Jamieson, J., and Schneebeli, M.: Snow avalanche formation, *Reviews of Geophysics*, 41, <https://doi.org/10.1029/2002RG000123>, 2003.
- 665 Schweizer, J., Bellaire, S., Fierz, C., Lehning, M., and Pielmeier, C.: Evaluating and improving the stability predictions of the snow cover model SNOWPACK, *Cold Regions Science and Technology*, 46, 52 – 59, <https://doi.org/10.1016/j.coldregions.2006.05.007>, 2006.
- Schweizer, J., Mitterer, C., Techel, F., Stoffel, A., and Reuter, B.: On the relation between avalanche occurrence and avalanche danger level, *The Cryosphere*, 14, 737–750, <https://doi.org/10.5194/tc-14-737-2020>, 2020.
- Sielenou, P. D., Viallon-Galinier, L., Hagenmuller, P., Naveau, P., Morin, S., Dumont, M., Verfaillie, D., and Eckert, N.: Combining random forests and class-balancing to discriminate between three classes of avalanche activity in the French Alps, *Cold Regions Science and Technology*, p. 103276, <https://doi.org/10.1016/j.coldregions.2021.103276>, 2021.
- 670 Singh, A. and Ganju, A.: Artificial Neural Networks for snow avalanche forecasting in Indian Himalaya, in: Proceedings of 12th International Conference of International Association for Computer Methods and Advances in Geomechanics, IACMAG, vol. 16, 2008.
- Statham, G., Haegeli, P., Greene, E., Birkeland, K., Israelson, C., Tremper, B., Stethem, C., McMahon, B., White, B., and Kelly, J.: A conceptual model of avalanche hazard, *Natural hazards*, 90, 663–691, <https://doi.org/10.1007/s11069-017-3070-5>, 2018.
- 675 Stethem, C., Jamieson, B., Schaerer, P., Liverman, D., Germain, D., and Walker, S.: Snow Avalanche Hazard in Canada – a Review, *Natural Hazards*, 28, 487–515, <https://doi.org/10.1023/a:1022998512227>, 2003.
- van Herwijnen, A. and Schweizer, J.: Monitoring avalanche activity using a seismic sensor, *Cold Regions Science and Technology*, 69, 165–176, <https://doi.org/10.1016/j.coldregions.2011.06.008>, 2011.
- 680 van Herwijnen, A., Gaume, J., Bair, E. H., Reuter, B., Birkeland, K. W., and Schweizer, J.: Estimating the effective elastic modulus and specific fracture energy of snowpack layers from field experiments, *Journal of Glaciology*, 62, 997–1007, <https://doi.org/10.1017/jog.2016.90>, 2016.
- van Herwijnen, A., Heck, M., Richter, B., Sovilla, B., and Techel, F.: When Do Avalanches Release: Investigating Time Scales in Avalanche Formation, in: Proceedings, International Snow Science Workshop, 2018.

- 685 Vernay, M., Lafaysse, M., Hagenmuller, P., Nheili, R., Verfaillie, D., and Morin, S.: The S2M meteorological and snow cover reanalysis in the French mountainous areas (1958 - present), version 2020.2, <https://doi.org/10.25326/37>, 2020.
- Viallon-Galinier, L., Hagenmuller, P., and Lafaysse, M.: Forcing and evaluating detailed snow cover models with stratigraphy observations, *Cold Regions Science and Technology*, 180, 103 163, <https://doi.org/10.1016/j.coldregions.2020.103163>, 2021.
- Viallon-Galinier, L., Hagenmuller, P., Reuter, B., and Eckert, N.: Modelling snowpack stability from simulated snow stratigraphy: Summary and implementation examples, *Cold Regions Science and Technology*, 201, 103 596, <https://doi.org/10.1016/j.coldregions.2022.103596>, 2022.
- 690 Vionnet, V., Brun, E., Morin, S., Boone, A., Faroux, S., Le Moigne, P., Martin, E., and Willemet, J.-M.: The detailed snowpack scheme Crocus and its implementation in SURFEX v7.2, *Geoscientific Model Development*, 5, 773–791, <https://doi.org/10.5194/gmd-5-773-2012>, 2012.
- Wilhelm, C., Wiesinger, T., and Ammann, W.: The avalanche winter 1999 in Switzerland-an overview, Tech. rep., WSL/SLF Davos, 2000.
- 695 Zeidler, A. and Jamieson, B.: A nearest-neighbour model for forecasting skier-triggered dry-slab avalanches on persistent weak layers in the Columbia Mountains, Canada, *Annals of Glaciology*, 38, 166–172, <https://doi.org/10.3189/172756404781815194>, 2004.
- Zgheib, T., Giacona, F., Granet-Abisset, A.-M., Morin, S., and Eckert, N.: One and a half century of avalanche risk to settlements in the upper Maurienne valley inferred from land cover and socio-environmental changes., *Global Environmental Change*, 65, 102 149, <https://doi.org/10.1016/j.gloenvcha.2020.102149>, 2020.
- 700 Zgheib, T., Giacona, F., Granet-Abisset, A.-M., Morin, S., Lavigne, A., and Eckert, N.: Spatio-temporal variability of avalanche risk in the French Alps, *Regional Environmental Change*, 22, <https://doi.org/10.1007/s10113-021-01838-3>, 2022.

Code and data availability. The meteorological and snow cover reanalysis used in this study is freely available at <http://dx.doi.org/10.25326/37#v2020.2>. The whole EPA avalanche database is freely available at <https://www.avalanches.fr/>.

Author contributions. Léo Viallon-Galinier developed the model code and performed the simulations. Léo Viallon-Galinier prepared the manuscript with contributions from all co-authors.

705

Competing interests. The authors declare no competing interests.

Acknowledgements. IGE/INRAE and CNRM/CEN are members of LabEx OSUG. Authors are grateful to the numerous people from ONF-RTM that contributed to the EPA data collection.