

## Response to reviewers

We thank the reviewers for the time and effort taken to read through and comment upon this manuscript, and for their helpful comments which have helped us to make improvements. We attach a document showing changes made to the manuscript in response.

### Reviewer 1

This paper presents a suite of synthetic simulations of the Amundsen sea sector of the Antarctic Ice-Sheet. The objective is to study the sensitivity of centennial scale ice-sheet simulations to the basal friction law formulation.

The results show that the predicted volumes and rates of volume change are sensitive to the friction law, but also to the exponent  $m$ , with complex patterns at the regional scale. However they found that all simulations follow the same trajectories (i.e. mass loss or gain) for a given perturbation, giving some confidence that ice-flow models have some predictive skills at these time-scales.

The paper is well structured and written and the results are convincing. I have only few minor comments.

First, effective pressure dependent friction laws, by definition, depends on the basal water pressure, and thus this requires a specific model. However, it's only in the discussion section that we learn the assumption that has been made, i.e. perfect connectivity. This assumption should already be presented in the model set-up section.

A good point! This assumption is now mentioned in the model setup.

Second, for the *Decreased Melt Branch* experiments, I suspect that this perturbation might lead to an advance of the grounding lines? However, the friction parameters are not constrained by observations in initially floating areas, so the differences between the simulations might more depend on the assumptions for the different priors than on the friction law or  $m$ ?

Decreasing the melt does indeed cause advance, but we do not believe this is a problem for our experiments. One advantage of allowing the model to run for 20 years before introducing the perturbations is that in most places the grounding line retreats during this time, and the initial response to the perturbation occurs at a point where the sliding parameter is constrained. A new paragraph has been added to the discussion to address this.

Minor comments :

- Page 8, line 171 : « *a slight change had to be made in order* ». Please explain here what are these slight changes ; I think we read later that this is changes in the prior ?

The changes to the prior are now explained as part of the overview of experiments.

- Page 13, last paragraph : I don't understand the meaning of the first two sentences.

The sentences have been edited to hopefully make the meaning clearer.

### Reviewer 2

In this manuscript, the authors present the results of sensitivity study in which they vary the choice of sliding parameterization along with the numerical values of the associated parameters and evaluate the different solutions that result for a pair of important glaciers in the Amundsen Sea embayment under several idealized geometric perturbations. They find that there are broad similarities in the trajectories of models using these different parameterizations, with the overall sign remaining the same for all cases, but with some qualitative differences as well as some regional peculiarities.

I found the paper to address an important question in ice sheet modelling and to be reasonably easy to follow. The results are interesting and provide some degree of comfort that, even though as modellers we are generally ignorant of the appropriate way to model sliding, there is some hope in West Antarctica that the models are still producing useful results. I have a handful of specific comments and concerns - some more serious than others - which I have included below.

Comments:

L21 Define effective pressure here rather than on line 33.

The definition has been moved.

L58 The initialism 'ASE' is first defined on line 88, below where it appears here.

The text has been changed to avoid use of the acronym before definition.

L65 It doesn't really make sense to say that a higher  $m$  produces higher velocities unless it's qualified by a description of what's going on with the friction coefficient, which is, in this work and others, treated as an unconstrained spatially-varying free parameter. Can this notion of velocity increasing with  $m$  be made a bit more rigorous?

A sentence has been added to explain that the sliding parameter  $C$  is set using initialisation methods based on matching to observations, and a clarification made that the higher velocities referred to are in forward simulation, rather than the initial velocity field. Hopefully this makes the meaning clearer.

L97–99 There are a few too many commas in this sentence.

This has been re-written.

Eqs. 1–3 These equations are defined again later, albeit in slightly different form, which is a bit confusing. I understand why they are here for exposition, but maybe

they could either be described qualitatively, or not numbered, or something else to make clear that they are not necessarily the equations actually implemented.

The numbering has been removed from these equations, to distinguish them from those used in the model which are introduced later.

L103 While most readers of this paper will likely already understand the assumptions of the SSA, for completeness they should be briefly mentioned here.

An additional descriptor of “vertically-integrated” has been added to SSA in this sentence, and a reference added to MacAyeal(1989).

Eq. 4 Probably not necessary to include this, unless you are going to state the full set of governing equations. I think it’s okay just having it in the Ua reference.

The equation has been removed.

L108 What flux and stress conditions are imposed at the inland boundary?

L109 It is not very clearly described how calving is treated in this model. Is the calving front held fixed or evolved? How?

Descriptions of the boundary conditions (inland and calving front) have been added to the model setup.

L112 We’re not given any context into how these densities are used (and are not mentioned again in the text). These should either be omitted or the full model needs to be described.

Clarification has been made that these densities are consistent with the BedMachine geometry. We believe they must be included in the setup as important physical parameters of the system.

Eqs. 6–10 It’s not clear to me what the practical difference is between  $C$  and  $\mu$ . While they may have different physical interpretations, both would appear to be friction coefficients. However, in this paper, one is treated as a spatially varying field to be inferred from velocity observations while the other is a scalar parameter. It would be helpful to try to clarify this in the text. This also implies that different sliding laws have substantially different degrees of flexibility and, as a result presumably exhibit different degrees of fidelity to observed surface velocities. It would be worthwhile to comment on this issue as well: are the differences in evolution really the result of the different sliding laws, or is it because some laws match observed velocities better than others?

We believe that calling  $C$  a “friction parameter” (as often the case in the literature) is misleading. We refer to  $C$  as a sliding parameter, or sometimes basal slipperiness. The  $C$  distribution incorporates behaviour due to the thermal and mechanical properties of ice as well as bed roughness. Since all of our sliding laws include the sliding parameter  $C$ , which is now explicitly noted in the text, the degrees of flexibility are comparable, whereas using a spatially varying friction coefficient would indeed add extra degrees of freedom to some laws (in particular the regularised Coulomb law) but

not others. The approach we use is common among modelling studies. There is not a large variation in how well velocities are matched in the inversion process by different sliding laws, and this is now mentioned in the model setup section.

Eq. 7–10 I think that the redefining of  $\beta_2$  is a confusing choice of notation, given that  $\beta_2$  usually means what you call C. Furthermore, this substitution makes the dependence of the various laws on the sliding exponent  $m$  much less transparent. I suggest writing the laws in a way that makes this dependence more plain (perhaps by just substituting the given expression for  $\beta_2$  back into the equations).

The description of the sliding laws has been simplified by putting  $\beta_2$  back into the equations as suggested.

L130–131 As written, the Budd law only becomes similar to the Coulomb law for high  $m$  when  $q = m$ , which these experiments don't explore.

The statement was referring to the Weertman law, which was not sufficiently clear. This has been rectified.

Eq. 12 The choices made for  $p_k$ , prior need to be shown here. Is it a spatially varying field or is it a scalar? how is it selected? Furthermore, how well the resulting velocity solutions correspond to observations (and to one another between sliding laws) needs to be quantified here.

More detail about the priors has been added to the model setup section.

L143 Total area of what? Perhaps introduce a symbol ( $\Omega$  is frequently used) to represent the model domain.

The total area of the domain. This has been clarified in the text.

L148 I don't know what it means for the system to 'settle', nor is it clear what it means for a strongly non-linear system as this to 'evolve steadily'. The manuscript needs to include a more detailed description of the criteria used for assessing whether the so-called 'transients' associated with data-model incompatibility in time-dependent simulations have reached a sufficiently small degree of influence so as to render the resulting simulations plausible.

This part has been rewritten. From experience across the userbase of the model we know that the initial transients in  $\dot{U}_a$  are only present for the first year or so of a run, and our 20 year relaxation period allows plenty of time for these to become irrelevant.

L152–157 I would suggest using the 'description' LaTeX environment (or equivalent) here rather than 'itemize'; the two different hyphens look weird together.

A good suggestion. This has been done.

L165 Median might be a better choice than mean here, since the sample contains some outliers (e.g. Budd  $q = 1$ ,  $m = 7$ ).

A good point. The median value has been added, although in this case it was not much different from the mean.

L170–174 It would be better to be specific about what change was made to convince the solver to converge so that a reader can properly assess the impacts that such a change would have when compared to the other simulations.

The details of this have been added to the model setup section.

L186 and elsewhere Perhaps two significant digits is sufficient when reporting percentage changes.

The reported values in the text have been reduced to two decimal places.

Fig. 5 This may be a comment on  $U_a$  more than the present manuscript, but Fig. 5 shows clear indication of numerical artifacts in the thickness field (presumably the result of an unstable continuity discretization). If this is indeed the case, this needs to be mentioned in the text and the potential implications mentioned (which might be significant: after all, the flux across the grounding line depends quite sensitively on the thickness there).

There is indeed evidence of numerical artifacts in the thickness field upstream where the resolution is coarse. When compared to a finer mesh, this choice of resolution did not change our findings regarding the patterns of behaviour due to sliding law choices, and so does not affect the relevant outcomes within the context of this paper. However, it is important to note that the numerical artifacts are among the reasons that the experiments in this study should not be treated as predictive, something which has been made clear. This has been addressed both in the model setup and the discussion sections.

L245–246 It is not immediately clear why the oceanic pressure condition shouldn't apply away from the grounding line. In any case, a reference for this claim is needed.

The point has been expanded slightly, and a reference added.

L254 typo: 'deivation'

Corrected.

L316–324 It might be worth mentioning that this effect of faster transport of upstream ice to the grounding line is probably a strictly transient phenomenon.

It is unclear to us what is meant by this, so no change has been made to this paragraph.

L358–362 The conclusion that the sliding law choice doesn't matter much over centennial time scales is reasonably well supported by the results of this paper, but this might not be true in non-WAIS scenarios; it's worth remembering that this is a system that is already undergoing a strong and non-linear transient response and may

be much less sensitive to the types of choices explored here than other places. I think that the language of this section needs to be tempered a bit to reflect this.

A fair point. This conclusion has been qualified to recognise that it may not be true everywhere.

# Compare Results

Old File:

**Paper\_SlidingLaws.pdf**

21 pages (2.07 MB)

04/08/2022 16:03:50

versus

New File:

**PIG\_Thwaites\_Sliding\_Laws.pdf**

22 pages (2.54 MB)

05/09/2022 11:01:11

## Total Changes

328

## Content

102 Replacements

112 Insertions

89 Deletions

## Styling and Annotations

4 Styling

21 Annotations

[Go to First Change \(page 1\)](#)

# The predictive power of ice sheet models and the regional sensitivity of ice loss to basal sliding parameterisations: A case study of Pine Island and Thwaites Glaciers, West Antarctica

Jowan M. Barnes<sup>1</sup> and G. Hilmar Gudmundsson<sup>1</sup>

<sup>1</sup>Department of Geography and Environmental Sciences, Northumbria University, Newcastle upon Tyne, UK

**Correspondence:** Jowan M. Barnes (jowan.barnes@northumbria.ac.uk)

**Abstract.** Ice sheet models use a wide range of sliding laws to define a relationship between ice velocity and basal drag, generally comprising some combination of a Weertman-style power law and Coulomb friction. The exact nature of basal sliding is not known from observational data, making assessment of the suitability of different sliding laws difficult. The question of how much this choice could affect predictions of future ice sheet evolution is an important one. Here we conduct a model study of a large sector of the West Antarctic Ice Sheet (WAIS), a particularly critical component of the cryosphere, using a range of sliding parameterisations, and provide an assessment of the sensitivity of ice loss to the choice of sliding law. We show that, after initialisation, various sliding laws result in broadly similar ranges of sea-level contribution over 100 years, with the range primarily dependent on exact parameter values used in each sliding law. Comparing mass loss from Thwaites and Pine Island glaciers, and the neighbouring regions, reveals significant qualitative geographical differences in the relationship between sliding parameters and the modelled response to changes in forcing. We show that the responses do not necessarily follow universal systematic patterns and, in particular, higher values of the sliding exponent  $m$  do not necessarily imply larger rates of mass loss. Despite differences in the magnitudes of ice loss and rates of change in the system, all our experiments display broad similarities in behaviour which serve to reinforce the decade-to-century-scale predictive power of ice sheet models, regardless of the choice of basal sliding.

15 *Copyright statement.* ©2022, all rights reserved

## 1 Introduction

For fast flowing areas of the Antarctic Ice Sheet, a significant portion of the forward motion is due to sliding along the bedrock beneath them. Ice sheet models must mathematically represent this sliding in order to simulate the physical system, but the exact nature of the relationship between a glacier and its bedrock cannot be determined from observational data, and is not fully understood. This process enters models in the form of a sliding law describing the form of basal drag. In general, these laws express basal drag in terms of the basal velocity and/or effective pressure, the difference between the hydrostatic overburden of ice and the basal water pressure. Sliding laws contain some parameters whose values are unknown, and are prescribed as part

of setting up a model. As a result, they can vary between models, and between simulations, giving different results in future predictions. While there are some common choices for these parameters, it is unknown which are the most representative of the physics of the ice sheet system.

An early sliding law, and one which is still widely used in modelling, is that of Weertman (1957). This is a power law of the form

$$v_b = C\tau_b^m,$$

where  $\tau_b$  is the basal drag,  $C$  is a sliding parameter,  $v_b$  is the basal velocity and  $m$  is a sliding exponent. We refer to this as the Weertman sliding law.

Throughout its history, the Weertman law has been criticised for neglecting the effects of cavitation, in which water under the ice can affect the basal drag. Lliboutry (1968) proposed the idea that a sliding law should depend on effective pressure. This can be done using Coulomb's law of solid friction, giving the relationship

$$\tau_b = \mu_k N,$$

where  $\mu_k$  is a coefficient of friction and  $N$  is the effective pressure.

The idea of including effective pressure in the sliding law has been explored further. Budd et al. (1979) proposed a law of the form

$$v_b = N^{-q} C_B \tau_b^m,$$

where  $C_B$  is a sliding parameter and  $q$  is a positive constant. Fowler (1987) produced a sliding law with broadly similar behaviour to that of Budd's, but which approached Coulomb friction at high velocities. Schoof (2005) asserts that laws taking the form of the Budd law are unphysical as they allow arbitrarily large stresses at the bed. Motivated by the work of Iken (1981), Schoof (2005) derived a regularised Coulomb sliding law, stated to be "essentially the same" as that of Fowler (1987). Variations of this have been used in several models since (e.g., Gagliardini et al., 2007; Leguy et al., 2014; Cornford et al., 2020).

Another approach, suggested by Tsai et al. (2015), is to compute the basal drag using both a Weertman sliding law and a Coulomb friction law, and take the minimum value of the two at each point. We refer to this as the Tsai sliding law. The Coulomb friction is applicable to softer beds, and is more representative of sliding near the grounding line. Taking the minimum value of this and the Weertman law effectively applies Coulomb friction near the grounding line, and Weertman sliding elsewhere.

With several sliding laws available, and parameter options within them, it is useful for modellers to know more about the possible implications of their choice. Previous studies have compared sliding laws in different ways. Brondex et al. (2017)

compares the Weertman, Budd, regularised Coulomb and Tsai sliding laws in an idealised setting, finding different levels of transient response in each case. This was followed up in Brondex et al. (2019) by simulations on a West Antarctic domain using linear and nonlinear versions of the Weertman and Budd laws, and two friction coefficients in the regularised Coulomb law. In both studies, the Weertman law was found to produce the smallest changes in volume above flotation (VAF), while the Budd law caused the largest changes. Simulations using the non-linear versions of the Budd and Weertman sliding laws lost ice faster than those using the linear versions.

The effect of the value of  $m$  in the Weertman sliding law has been investigated in a few cases. In these studies, the sliding parameter  $C$  is set for each value of  $m$ , usually by using inversion methods, in order to match an initial velocity field. Gillet-Chaulet et al. (2016) tests different values, and finds that higher values of  $m$  generally produce higher velocities in forward simulations and a closer match to an observed surface velocity field of Pine Island Glacier (PIG), with  $m = 20$  suggested as the best value in this case. Joughin et al. (2019) compares experiments using the Weertman law with different values of  $m$  to a regularised Coulomb law, focusing on a profile along PIG. The regularised Coulomb law was found to reproduce observations with greater success, and higher values of  $m$  in the Weertman law approached a better result. De Rydt et al. (2021) explores the idea that  $m$  in the Weertman sliding law does not necessarily need to take a uniform value, presenting a map of "optimal"  $m$  values over PIG which come closest to reproducing observed velocity changes. In general, this involves higher values of  $m$  on the fastest flowing areas. Sun et al. (2020) compares the response of several models to ice shelf collapse. It is shown that there is a general tendency for models using higher values of  $m$  to respond faster to the changes in forcing. This is in agreement with results from within single models in Brondex et al. (2019), Joughin et al. (2019) and De Rydt et al. (2021).

75

Some examples can be found of the opposite being true, indicating a more complex relationship between the value of  $m$  and the rates of change to a system. For Pine Island Glacier in particular, Nias et al. (2018) shows that higher values of  $m$  can produce higher or lower changes in VAF depending on the bathymetry product being used. This sensitivity to topography is further demonstrated by Wernecke et al. (2022).


80

The focus of the work presented in this paper is on the effects of parameter choices in sliding laws on the forward evolution of an ice sheet system and the response to changes in ice shelf forcing. The values of  $m$ ,  $\mu_k$  and  $q$  are varied, and we seek to quantify the difference these choices can make to the modelled response when different changes in the buttressing of an ice sheet system are applied, as well as whether there are universal systematic patterns when varying sliding parameters. We also set out to assess the degree of confidence with which predictions can be made by an ice sheet model, despite the lack of empirical knowledge of basal conditions to inform a sliding law.

The chosen domain for the experiments presented in this paper is the Amundsen Sea Embayment (ASE) in West Antarctica (Figure 1), the region of the continent which is experiencing the fastest ice mass loss (Shepherd et al., 2018; Rignot et al., 2019), at a rate which is accelerating (Sutterley et al., 2014). Understanding how the West Antarctic Ice Sheet (WAIS) is evolving

ing is seen as one of the most important questions in Antarctic research (National Academies of Sciences, Engineering, and Medicine, 2015). A lot of past work has been centred around PIG, including the previously mentioned studies. Recently, there has also been a focus on Thwaites Glacier due to the rapid changes observed in recent years (e.g., Mouginot et al., 2014; Milillo et al., 2019), and it is the subject of a large multidisciplinary investigation, the International Thwaites Glacier Collaboration (Scambos et al., 2017). Many models have been used to make predictions of the future evolution of Thwaites Glacier and the ASE region, and they generally agree that the trend of rapid retreat in the ASE will continue into the future (e.g., Favier et al., 2014; Joughin et al., 2014; DeConto and Pollard, 2016; Yu et al., 2018). Models tend to differ on the rates of change they calculate for a variety of reasons (e.g., Cornford et al., 2020), and as such efforts are ongoing to improve the understanding and functionality of all aspects of ice sheet models, and to reduce uncertainty in their predictions.

100



**Figure 1.** The catchment basins of the Amundsen Sea Embayment (ASE) using MEASUREs Antarctic Boundaries (Rignot et al., 2013) and, inset, the location of the ASE within Antarctica. The different colours indicate the four regions we focus on. The thicker black outline is the boundary of our model domain.

## 2 Experimental design

### 2.1 Model setup and forcing

We use the ice sheet model Úa (Gudmundsson, 2020), which implements the vertically-integrated Shallow Shelf Approximation (SSA) (MacAyeal, 1989), and Glen’s flow law (Glen, 1958) with exponent  $n = 3$ . Our domain covers the ASE region including PIG, Thwaites Glacier and the glaciers flowing into the Dotson and Crosson ice shelves. The inland boundary was created based on the MEaSUREs Antarctic Boundaries (Rignot et al., 2013), with smoothing applied. We use the geometry from BedMachine Antarctica v2 (Morlighem et al., 2020), from which we also derive the calving front boundary for our domain. A Dirichlet condition is used to set all velocities along the inland boundary to zero, since the domain boundary generally follows the edges of drainage basins. The ice-front stress boundary condition is applied along the seaward boundaries, which are fixed at the calving front. The mesh resolution is 1.5 km at the grounding line, and becomes coarser further upstream. The resolution was chosen for computational efficiency, and test cases were compared to finer meshes to ensure this resolution is sufficient to demonstrate the relative behaviours of sliding laws required for this study.

The densities are given the values of  $917 \text{ kg m}^{-3}$  for ice and  $1027 \text{ kg m}^{-3}$  for ocean water, consistent with the BedMachine geometry. The effective pressure is calculated by assuming a perfect hydrological connection to the ocean. The surface mass balance is taken from RACMO2.3p2 (Van Wessem et al., 2018). Where the ice is afloat, we use a simple depth-based melt parameterisation of ocean-induced melt

$$M = \begin{cases} 0 & \text{if } z \geq 0 \\ -\frac{M_{\max}}{500} z & \text{if } 0 > z > -500 \\ M_{\max} & \text{if } z \leq -500, \end{cases} \quad (1)$$

where  $M$  is the basal melt rate in  $\text{m a}^{-1}$ ,  $M_{\max}$  is a prescribed maximum melt rate and  $z$  is the vertical coordinate in metres, positive upwards with zero at sea level.

Our experiments test four different sliding laws, and vary the parameters within them. We use the Weertman, Budd and Tsai sliding laws introduced in section 1, and a regularised Coulomb law used in Cornford et al. (2020), which is a power-weighted reciprocal sum of the Weertman and Coulomb sliding laws (see Eq. 6). We present these sliding laws below in a form

125 which makes clear how they relate to each other.

$$\text{Coulomb friction:} \quad \tau_C = \mu_k N \frac{\mathbf{v}_b}{\|\mathbf{v}_b\|} \quad (2)$$

$$\text{Weertman sliding:} \quad \tau_W = C^{-\frac{1}{m}} \|\mathbf{v}_b\|^{\frac{1}{m}-1} \mathbf{v}_b \quad (3)$$

$$\text{Budd sliding:} \quad \tau_B = N^{\frac{q}{m}} C_B^{-\frac{1}{m}} \|\mathbf{v}_b\|^{\frac{1}{m}-1} \mathbf{v}_b \quad (4)$$

$$\text{Tsai sliding:} \quad \tau_T = \min(\tau_C, \tau_W) \quad (5)$$

$$130 \text{ Regularised Coulomb sliding:} \quad \tau_R^m = \frac{1}{\tau_W^m} + \frac{1}{\tau_T^m} \quad (6)$$

Note that Coulomb friction alone is not used in any of our model simulations, meaning that all of our sliding laws contain the sliding parameter  $C$  which incorporates the effects of various thermal and mechanical properties of ice as well as basal roughness (e.g., Cuffey and Paterson, 2010). The parameters are given a range of values, exploring a range around common values used in modelling studies. We use  $m = 1, 3, 5, 7$  and  $9$ . Commonly,  $m = 3$  is used by ice sheet models, but higher values of  $m$  in the Weertman law could produce behaviour closer to that of regularised Coulomb laws. We use  $q = 1$  and  $2$ . The value  $q = 2$  was originally suggested by Budd et al. (1984), but  $q = 1$  is a more common choice in recent studies (e.g., Brondex et al., 2019). A range of friction coefficients have been used in modelling, and mostly fall within the range of our chosen values,  $\mu_k = 0.25, 0.5, 0.75$ .

## 140 2.2 Overview of experiments

We use an adjoint inversion method to calculate fields of values for  $A$  and  $C$  independently for each set of parameters in each sliding law. The method involves referring to observed velocities  $u_{\text{obs}}$  and  $v_{\text{obs}}$  with observational errors  $u_{\text{err}}$  and  $v_{\text{err}}$ , from the 2016-17 year of the dataset presented by Mouginot et al. (2017), to minimise a cost function  $\mathcal{J} = I + R$ , consisting of a misfit function

$$145 \quad I = \frac{1}{2\mathcal{A}} \int ((u - u_{\text{obs}})/u_{\text{err}})^2 d\mathcal{A} + \frac{1}{2\mathcal{A}} \int ((v - v_{\text{obs}})/v_{\text{err}})^2 d\mathcal{A}, \quad (7)$$

and a regularisation term

$$R = \sum_{k=1,2} \frac{1}{2\mathcal{A}} \int \left( \gamma_{s,k}^2 (\nabla(p_k - p_{k,\text{prior}}))^2 + \gamma_{a,k}^2 (p_k - p_{k,\text{prior}})^2 \right) d\mathcal{A}, \quad (8)$$

where  $\mathcal{A} = \int d\mathcal{A}$  is the total area of the domain,  $u$  and  $v$  are the modelled horizontal  $x$  and  $y$  velocity components, respectively,  $\gamma_{s,k}$  and  $\gamma_{a,k}$  are slope and amplitude regularisation parameters,  $p_1 = \log_{10} A$ ,  $p_2 = \log_{10} C$  and  $p_{k,\text{prior}}$  are prior values, or initial estimates, for the parameters  $p_k$ . For the inversions in this study,  $\gamma_{s,1} = \gamma_{s,2} = 1 \times 10^4$  m,  $\gamma_{a,1} = 100$  and  $\gamma_{a,2} = 1$ . Further details of the inversion method can be found in Barnes et al. (2021). In most cases, the priors used in these experiments are the outputs of a previous inversion for the Weertman law with  $m = 3$ , adjusted using an analytical conversion to account for the value of  $m$ . The original Weertman inversion used the priors named 'Priors2' in Barnes et al. (2021), which

are spatially varying based on observed velocities and a temperature field. A change had to be made to the value of  $p_{2,prior}$  in Equation 8 in order for the inversion process to run successfully in the  $m = 9$  cases for the Regularised Coulomb law with  $\mu_k = 0.5$  and  $0.75$ , and for the Tsai law with  $\mu_k = 0.75$ . In these cases, the inverted  $C$  field for the Tsai law with  $m = 9$  and  $\mu_k = 0.5$  is used as the prior. The mean absolute speed misfit over the domain is within the range  $27\text{-}52 \text{ ma}^{-1}$  for all inversions.

Each simulation is run for 20 years using  $M_{max} = 100$ . This relaxation period allows ample time to move past any initial transient effects, which are fairly common in ice flow models (e.g., Pattyn et al., 2012) and in Úa only have a significant influence the output during the first few years of model evolution. We do not introduce different forcing from the beginning of the run, as we wish to assess the effects of a change introduced to a system which is already in motion. Changes are introduced after 20 years of simulation, and the experiments are split into four branches from this point:

- Control Branch - the simulation continues with the same forcing.
- Increased Melt Branch - there is an immediate step change increase in the melt rate, with the value of  $M_{max}$  raised to 200.
- Decreased Melt Branch - melting stops immediately, with  $M_{max}$  lowered to 0.
- Ice Shelf Removed Branch - all the floating ice is immediately removed and no more is allowed to form for the remainder of the run.


These four branches are run for a further 100 years. These are not realistic perturbations, but are designed to test the most extreme changes to the ice shelf buttressing, as well as one more moderate case, in order to clearly see the differences caused by altering the sliding parameterisation.

### 3 Results

The experiments in the *Control Branch* all display loss of ice from the system, with the loss of volume above flotation (VAF) over the 120 years of simulation being between 37 and 104 mm sea level equivalent, depending on the sliding law and parameter choices (Figure 2 and Table 1). The mean ice loss is 59.5 mm sea level equivalent, and the median is 59.3 mm. There is a clear pattern of lower values of  $m$  causing smaller VAF changes in all sliding laws other than Budd. The Regularised Coulomb and Tsai laws produce very similar curves, becoming more similar with higher values of  $m$ . When using the same values of  $m$ , both produce larger mass loss than the Weertman law even for the highest friction coefficient in our ensemble ( $\mu_k = 0.75$ ), although the gap between the two becomes significantly smaller with higher values of  $m$ . There is also a systematic pattern of higher values of  $\mu_k$  (i.e. higher friction) causing less change in volume above flotation. In a few cases, the curves do not fit the pattern, due to the use of a different prior during initialisation. These particular cases should perhaps be discounted when assessing patterns in the results, but are presented here to demonstrate a dependence of results on the initialisation.

185 The Budd sliding law shows less systematic behaviour with respect to the value of  $m$ , but a higher value of  $q$  produces a larger change in VAF, for values of  $m$  up to 5. When  $m$  becomes too high, the Budd sliding law produces very high ice loss, as demonstrated when  $m = 7$  and  $q = 1$  in Figure 2. Raising  $m$  further soon makes the sliding law unusable. For both values of  $q$  in our ensemble, simulations using  $m = 9$  ran into convergence issues early on, and could not be completed.

190 Turning our attention to the responses to changes in ice shelf forcing, the differences in VAF loss for the entire ensemble of experiments are displayed in Figure 3 and Table 2. The results are presented with reference to the *Control Branch* in each case. Since the *Control Branch* is run for 20 years before any changes are made to the forcing, the system state at the point of perturbation will be slightly different in each case. It is therefore useful to look at VAF differences in terms of a percentage change




**Figure 2.** VAF loss in the *Control Branch* for every set of parameter choices in the sliding laws.

Sliding law	Value of $m$				
	1	3	5	7	9
Weertman	37.11	44.13	52.47	59.30	62.39
Budd, $q = 1$	61.27	57.76	60.95	103.82	-
Budd, $q = 2$	67.97	75.24	71.34	63.34	-
Reg. Coulomb, $\mu_k = 0.25$	57.76	61.01	66.01	70.14	71.33
Reg. Coulomb, $\mu_k = 0.5$	49.00	52.97	58.71	64.24	58.61*
Reg. Coulomb, $\mu_k = 0.75$	45.50	49.80	55.85	61.68	58.20*
Tsai, $\mu_k = 0.25$	51.06	60.41	65.56	69.74	71.42
Tsai, $\mu_k = 0.5$	44.09	51.78	57.36	62.11	65.18
Tsai, $\mu_k = 0.75$	41.02	48.29	54.79	60.17	57.68*

**Table 1.** The VAF loss (mm sea level rise equivalent) at the end of the *Control Branch* simulations, for all experiments. Values marked with \* indicate where a different prior had to be used in initialisation.

Sliding law	<i>Decreased Melt Branch</i>					<i>Increased Melt Branch</i>					<i>Ice Shelf Removed Branch</i>					
	Value of $m$	1	3	5	7	9	1	3	5	7	9	1	3	5	7	9
Weertman		-23.35	-32.68	-34.07	-34.52	-34.39	3.28	3.72	3.10	3.05	4.82	23.70	22.33	12.90	11.42	11.79
Budd, $q = 1$		-44.58	-41.61	-40.77	-34.70	-	14.66	10.74	7.18	15.69	-	52.86	31.72	23.97	32.74	-
Budd, $q = 2$		-56.32	-47.89	-45.59	-44.57	-	13.14	14.65	11.11	14.90	-	56.98	43.30	30.87	33.40	-
Reg. Coulomb, $\mu_k = 0.25$		-50.33	-52.32	-51.77	-49.65	-48.53	20.40	13.35	13.46	14.87	14.96	69.95	43.44	34.75	34.26	37.68
Reg. Coulomb, $\mu_k = 0.5$		-40.00	-43.36	-43.30	-42.62	-41.71*	10.98	9.33	8.39	8.98	8.89*	63.20	35.38	28.00	27.88	21.96*
Reg. Coulomb, $\mu_k = 0.75$		-35.69	-39.88	-40.53	-39.98	-38.44*	9.37	7.66	7.37	7.70	6.85*	59.89	29.75	25.80	24.21	14.33*
Tsai, $\mu_k = 0.25$		-47.12	-51.820	-52.13	-50.35	-50.71	21.15	13.50	13.35	14.20	14.54	103.01	48.04	34.80	33.88	35.27
Tsai, $\mu_k = 0.5$		-34.57	-42.43	-43.45	-43.26	-42.93	9.58	9.75	9.17	9.74	10.25	69.53	39.97	28.19	28.67	29.75
Tsai, $\mu_k = 0.75$		-29.87	-38.87	-40.93	-41.01	-39.25*	8.85	8.26	7.56	7.38	7.09*	62.10	31.34	25.81	23.26	15.89*

**Table 2.** The percentage differences in VAF at the end of simulation compared to the *Control Branch*, for all experiments. Values marked with \* indicate where a different prior had to be used in initialisation.



**Figure 3.** Additional VAF loss compared to the *Control Branch* after the changes in forcing are introduced, for the entire ensemble of sliding laws and parameter choices.

195 compared to the *Control Branch*, rather than as a volume in  $\text{km}^2$ . The Weertman sliding law displays the smallest response to all changes, while at a glance the other three sliding laws appear to respond quite similarly, when using the same values of  $m$ .


For the *Decreased Melt Branch*, it is notable in most panels of Figure 3 that a larger difference is seen between linear sliding ( $m = 1$ ) and the non-linear versions ( $m > 1$ ), than between any two non-linear cases, which all respond very similarly. For the Weertman law, the VAF loss difference at the end of the simulation compared to the *Control Branch* is  $-23.4\%$  for  $m = 1$  and within one percentage point of  $33.6\%$  for all other values of  $m$ . For the Tsai and regularised Coulomb laws, this gap between

the  $m = 1$  response and other values of  $m$  is larger for higher values of  $\mu_k$ , but the difference from the *Control Branch* is larger for lower values of  $\mu_k$ . The difference in response with varying friction coefficient is not linear. As an example, when  $m = 3$ , the VAF loss difference for the Regularised Coulomb law is -52.3 % with  $\mu_k = 0.25$ , -43.4 % for  $\mu_k = 0.5$  and -39.9 % for  $\mu_k = 0.75$ . Taking  $\mu_k = 0.5$  as a midpoint, lowering the friction causes a significantly larger difference in the response than raising it by the same amount. For the Budd law, there is more variability and a slightly larger percentage difference with  $q = 2$  compared to  $q = 1$ . The effect of  $m$  is also reversed compared to the other laws, with  $m = 1$  responding more than higher values.

The *Increased Melt Branch*, as the less extreme of the three forcing changes, generally causes the lowest differences in the responses. The VAF loss differences using the Weertman law range from 3.1 % to 4.8 %. The obvious outliers are in the Regularised Coulomb and Tsai laws with low friction ( $\mu_k = 0.25$ ), where the simulations using  $m = 1$  start to display an increased response towards the end of the run, and the Budd law for which there is more variability in the responses for different values of  $m$ , particularly when  $q = 1$ . Using Regularised Coulomb with  $m = 3$  as an example again, the response changes with  $\mu_k$  in a similar way to that seen in the *Decreased Melt Branch*. For  $\mu_k = 0.25$ , the VAF loss is 13.3 % greater than in the *Control Branch*, and increasing the friction to  $\mu_k = 0.5$  and  $\mu_k = 0.75$  decreases the difference to 9.3 % and 7.7 %, respectively.

The largest variability is found in the *Ice Shelf Removed Branch*. The Budd sliding law shows similar behaviour here as in the other two branches, with lower values of  $m$  and higher values of  $q$  causing larger responses. The other sliding laws all display a clearly new behaviour. In all cases, the general trend seen in other branches holds initially, with higher values of  $m$  causing greater changes to the VAF. However, the response for lower values of  $m$  in this branch becomes much larger at some point during each simulation. This point is reached earlier with lower values of  $\mu_k$  in the Tsai and Regularised Coulomb laws, and earliest when using the Tsai law, which produces just over twice the VAF loss of the *Control Branch* when  $m = 1$ . Within these two laws, the difference is far more evident with  $m = 1$ . Using the Weertman law, the difference in response is lower overall and very similar for  $m = 1$  and  $m = 3$ , which have VAF loss 23.7 % and 22.3 % higher than the *Control Branch*, respectively, while the other values of  $m$  are in the range 11.4-12.9 %. The effects of altering  $\mu_k$  are similar to those in the other branches, with the difference from the *Control Branch* higher with  $\mu_k = 0.25$  than when  $\mu_k = 0.5$ , and the difference when  $\mu_k = 0.75$  a little lower.

The VAF values are just one metric by which one can assess the system. Another option is to look at the positions of the grounding lines at the end of simulations. Figure 4 shows the grounding lines for all sliding laws using the values  $m = 3$ ,  $q = 1$  and  $\mu_k = 0.5$ , and for all values of  $m$  within the Weertman sliding law, in the *Ice Shelf Removed Branch*. Since this shows the response to the most extreme forcing change, this is the case for which differences are most evident. When comparing the sliding laws using the same parameter values, it is clear that the Weertman law causes the least retreat, and that the Tsai and Regularised Coulomb laws cause the most, ending up in very similar positions. For the most part, the grounding line produced by the Budd law lies somewhere between the two. When looking at the grounding lines for the Weertman law using different parameters, it can be seen that the grounding line retreats less far for higher values of  $m$ . This trend is much more noticeable



**Figure 4.** Grounding lines at 120 years in the *Ice Shelf Removed Branch* for (a) all sliding laws using  $m = 3$ ,  $q = 1$  and  $\mu_k = 0.5$  and (b) different values of  $m$  in the Weertman sliding law. The initial grounding line ( $t=0$ ) is also indicated.


around the Pine Island and Kohler Glaciers, while the grounding lines are located in very similar positions around Thwaites Glacier and the Crosson Ice Shelf.

We can also look directly at model outputs of velocities and ice thickness. Figure 5 displays the differences in speed and thickness at the end of the *Ice Shelf Removed Branch* for each non-linear version of the Weertman law, compared to that of the linear case ( $m = 1$ ). Across much of the domain, the ice is thinner and moves faster when higher values of  $m$  are used in the sliding law. However in some areas, specifically Pine Island and Kohler Glaciers, the opposite is true. The differences between experiments using successive values of  $m$  become less as  $m$  increases, so the largest difference is between the linear case and  $m = 3$ , while the difference between the  $m = 7$  and  $m = 9$  cases is much smaller. We note that there is evidence of numerical artifacts in the thickness field upstream where the resolution is coarse. When compared to a finer mesh, this choice of resolution does not change our findings regarding the patterns of behaviour due to sliding law choices, but the numerical artifacts do indicate that the experiments in this study should not be treated as predictive.

## 4 Discussion

### 4.1 The whole domain

As outlined in the previous section, the results display fairly systematic behaviour in many respects. Some of the patterns are simple, such as higher friction coefficients  $\mu_k$  resulting in smaller changes to the system. Other trends are more complicated.



**Figure 5.** Differences in final speed (panels a-d) and ice thickness (panels e-h) in the *Ice Shelf Removed Branch*, for the Weertman sliding law. All other values of  $m$  are compared to the case where  $m = 1$ . Areas which are not grounded in either case are not shaded as part of each comparison.

In particular, there is an apparent inconsistency in the effects of the value of  $m$ . When the melt is decreased, it is clear that higher values of  $m$  respond more to the change, with the system gaining more ice when  $m > 1$ . However, with an increased melt rate, and especially with the removal of the entire ice shelf, low values of  $m$  cause a larger response.

255

The Budd sliding law responds in ways which set it apart as an outlier among those tested. This is most likely because it relies on the value of effective pressure throughout the domain, which is calculated assuming a perfect hydrological connection with the ocean. This is a reasonable assumption close to the grounding line but becomes unlikely further upstream (e.g., Cuffey and Paterson, 2010), especially where the bedrock is above sea level and subglacial water pressure is set to zero. With more accurate information about the effective pressure at the base of the ice sheet, the Budd law would likely produce results more consistent with the other sliding laws. However, in our case the assumption causes erratic behaviour, and the Budd law is not considered in detail during our discussion. The Tsai and Regularised Coulomb sliding laws also include  $N$ , but in these laws the Coulomb component of the equation is only relevant in close proximity to the grounding line, with a transition into

260

Weertman-style sliding upstream.

265

Excluding the Budd sliding law, all of the *Control Branch* experiments (Figure 2) show a similar evolution in the volume of ice contained within the domain. After 120 years of transient simulation, the loss of volume above flotation ranges from about 37 mm to 71 mm sea level rise equivalent, with a mean value of 57.05 mm. The standard deviation is 8.68 mm, giving a coefficient of variation 0.15 which implies moderate variation in rates of change in the system. But no matter which sliding law or parameters are used, the direction of evolution is always the same, suggesting that the choice of sliding parameterisation does not introduce a level of uncertainty which would affect predictions of the broader patterns of future ice movement in a system forced by ice shelf melt rates in a realistic range. Indeed, it has been shown by Hill et al. (2021) that far larger uncertainties are generally introduced by oceanic forcing than by parameters of the ice flow dynamics.

270

275

The difference in rates of change when varying basal sliding conditions can be seen more clearly when perturbations are introduced. Again, all sliding laws react in broadly the same way, but with different rates and magnitudes of change. From this, we can say that the choice of sliding law does not impact predictions of how an ice sheet system will change in terms of the direction of evolution, but it does affect how fast the changes occur. It is also worth noting that the large differences in behaviour for low values of  $m$  in the *Ice Shelf Removed Branch* are due to an unrealistic extreme forcing accelerating the rate of change in the system, and that the results in the more moderate *Increased Melt Branch* do not differ so much.

280

In the *Decreased Melt Branch*, the grounding line advances beyond its initial position, where  $C$  was not inverted for and the prior value could have an influence on the results. However, due to the 20 years of model evolution before the perturbation is introduced, the initial responses to the perturbation are unaffected by this. With the exception of the Budd law, none of our experiments show a change in behaviour midway through the simulation.

285

The significantly larger percentage rises in VAF resulting from using lower values of  $m$  cannot be accounted for by differences in the system states after the first 20 years of simulation. They demonstrate differences between the rates at which the forward simulations respond after the perturbations are introduced. It is likely that the overall trend is driven by the large speed reductions of Pine Island as the value of  $m$  increases. The grounding line positions (Figure 4) and differences in ice thickness and speed (Figure 5) clearly suggest that the modelled response to different  $m$  values, and the sensitivity to basal sliding representation in general, is affected by regional differences.

290

## 4.2 Regional Variation


To investigate differences within the ASE, we divide the domain into four regions, as indicated by the colours on Figure 1. These regions are the catchment basins of three glaciers - Pine Island, Thwaites and Kohler - and the combined catchments of the small glaciers which flow out to the Crosson Ice Shelf. Figure 6 shows the magnitude of differences in VAF for these four regions. Pine Island Glacier has the largest overall ice loss, and thus contributes heavily to the behaviour seen for the whole

295


domain in the previous section.

300 Figure 7 shows the percentage differences in VAF loss compared to the *Control Branch* for all other branches using the Weertman sliding law. Figure 8 displays the same information in the case of using the Tsai sliding law with  $\mu_k = 0.5$ . The results from the Regularised Coulomb law are very similar to those from the Tsai law. We can see a variety of responses across the domain, showing that regional factors can play an important role in how sliding laws react to changes. In terms of **percent-**  
**age** increases, Kohler Glacier responds the most by far to changes involving a loss of buttressing, with up to 150% greater  
 305 VAF using the Tsai law. The differences in VAF for Thwaites and Crosson at the end of the simulations is very small, less than 10% for the Weertman law, and 30% for the Tsai law.


The large changes when the ice shelf is removed and  $m$  has a low value are not universal. They occur in the Pine Island and Kohler Glaciers, the same locations in which the largest grounding line retreat is seen in Figure 4. Thwaites Glacier,  
 310 and those flowing into the Crosson Ice Shelf, do not display the same behaviour. Looking at flowlines through Pine Island



**Figure 6.** Additional VAF loss compared to the *Control Branch* for the Weertman sliding law with different values of  $m$ . The contributions of each region within the ASE domain are shown.



**Figure 7.** Additional VAF loss compared to the *Control Branch* for the Weertman sliding law with different values of  $m$ . The contributions of each region within the ASE domain are shown. Note that the  $y$ -axes for each region use different scales.



**Figure 8.** Additional VAF loss compared to the *Control Branch* for the Tsai sliding law with  $\mu_k = 0.5$  and different values of  $m$ . The contributions of each region within the ASE domain are shown. Note that the  $y$ -axes for each region use different scales.

315 Glacier, we discover that 50 years into the simulations, the glacier is grounded on a ridge in the bathymetry. With low values of  $m$ , the ice unpins from this point earlier, producing the sudden rise in VAF differences. The region-specific effect at play here could be the bathymetry itself. Nias et al. (2018) found that for Pine Island Glacier, using different bathymetry datasets can effect whether more or less ice is lost with increasing values of  $m$ . They conclude that the buttressing force of a bump in one bathymetry dataset is amplified by larger values of  $m$ , slowing down the flow. Our bathymetry is a different product to either tested in that study, but it is likely that there is a similar cause for the pattern observed in our results and that such bathymetric features do not exist around Thwaites Glacier and Crosson Ice Shelf, so the same effect is not seen in those regions.

320 The relationship between the value of  $m$  and the rates of retreat and ice loss is an issue which could be explored further. Nias et al. (2018) identifies local bathymetry as a likely source of differences in behaviour, but there could be other factors to consider, such as variation in the basal sliding parameter  $C$  or flow rate factor  $A$ . We note that even a small change required for successful inversion in three of our cases caused the responses in those cases to fall out of line with the patterns seen in the rest, suggesting a high sensitivity to changes in the inverted fields. Whatever the cause, we can confidently say that the relationship contains complexities and can be affected by regional differences, so there is no simple statement to be made about what the general effect of increasing  $m$  will be on a system. We believe the regional variance in the effect of  $m$  supports the idea that  $m$  should perhaps be treated not as a universal constant, but as a distributed field of values, as suggested by De Rydt et al. (2021). At the very least, different glaciers may need to use different values in order to reproduce their observed rates of change.

330 In general, whatever differences are seen when increasing the value of  $m$ , they are smaller between higher values. This can be seen in the control experiments (Figure 2), in which the  $m = 7$  and  $m = 9$  cases are closer together than those of lower sequential values in our ensemble. It is also seen clearly in most panels of Figure 3 for the *Decreased Melt Branch* that the  $m = 1$  case is more different from the other cases, and in the differences in final speed and thickness presented in Figure 5. This demonstrates that when increasing the value of  $m$  in sliding laws, the response converges. With large values of  $m$ , the Weertman law is often said to approach the behaviour of a Coulomb friction law in this way.

335

An observation which appears strange at first is that when a change in forcing increases the melt rate or removes the buttressing ice shelf entirely, higher values of  $m$  can cause larger VAF losses but at the same time less retreat of the grounding line. We see this around Thwaites glacier when using the Weertman sliding law, for which the  $m = 9$  experiment produces the largest VAF loss in the *Control Branch* (Figure 2a), the largest percentage response in the *Ice Shelf Removed Branch* (Figure 7c), and the greatest ice speed and thinning (Figure 5), yet the least grounding line retreat (Figure 4b). We suggest that when more ice is lost from upstream areas, it crosses the grounding line at a rate faster than the ice shelf is melting away, thus causing a buildup of ice in this region. Because the ice around the grounding line is thicker, the glaciers stay grounded. In this way, it is possible for higher values of  $m$  to cause glaciers to lose a greater volume of ice while at the same time retaining a larger grounded area.

345 In most cases, we see that the responses in each region follow the same patterns using the Weertman and Tsai laws. There are some differences, most notably in the response of Thwaites, for which a significantly different pattern can be seen. However, this is another effect of the greater speed of response using the Tsai law. As can be seen in the grounding lines of Figure 4, for the Tsai and Regularised Coulomb laws, the ice retreats far enough for the Thwaites and Pine Island ice shelves to merge. The two glaciers become much more closely linked, and this has an effect on the evolution of Thwaites. Even before the ice sheets merge, the retreat of the small western section of Pine Island starts cutting into the Thwaites region, as we have defined the regions by present-day boundaries. The same behaviour would be seen in the Weertman case if the model was run for longer. This is a significant example of the importance of differing rates of change caused by the choice of basal sliding parameterisation.

## 5 Conclusions

355 Parameter choices in sliding laws have an important effect on the speed and magnitude of responses to changes in the forcing of the ice system, although the exact nature of the effect can be highly region-dependent. The Weertman, Tsai and Regularised Coulomb laws all display similar behaviours in the effects of changing the values of  $m$  and  $\mu_k$ . The Budd sliding law responds differently to the other laws, due to the simplistic hydrological condition used in our model, but in general raising the value of  $q$  increases the magnitude of the response.

360

Changing the value of the friction coefficient  $\mu_k$  produces a very structured response in all cases. Simulations with lower values of  $\mu_k$  respond more to changes in forcing. The relationship is not linear, with the differences in response compared to the *Control Branch* converging as  $\mu_k$  increases. This is to be expected, as there will be a convergence to the limit at which friction is sufficiently high to allow no basal sliding at all.

365

Changing the value of the sliding exponent  $m$  produces a more varied range of responses, and is affected by regional differences in bathymetry as well as the inverted fields  $A$  and  $C$ . In the case of lowering the ice shelf melt rate and allowing increased buttressing,  $m = 1$  is a notable outlier, producing less difference in VAF compared to the *Control Branch*, while higher values of  $m$  respond very similarly to the change. The response to increased melting or removal of the ice shelf also systematically shows higher values of  $m$  responding more in terms of VAF loss, up to a point at which the trend switches. This is due to regional changes at Pine Island and Kohler Glaciers, where lower values of  $m$  cause more rapid retreat along the main glacier trunks. Thwaites Glacier, and those flowing out to the Crosson Ice Shelf, do not display this behaviour.

375 Despite the differences in our results when using different parameterisations of basal sliding, with the highest VAF loss at the end of the *Control Branch* being almost twice that of the lowest, there is also a degree of similarity between them. In broad terms, the system always reacts in the same way to a particular change in forcing, no matter which sliding law or parameter values are used. The magnitude and speed of these reactions vary, but the response is clearly bounded. Our results suggest that ice sheet models, if initialized by inverting for sliding-law parameters using present-day geometry and surface velocities and driven by forcing in a realistic range, follow approximately the same trajectory for at least several decades and up to a century, irrespective of the choice of sliding law. In this sense, the predictions of ice sheet models about near-future ice loss are reasonably well constrained despite our incomplete knowledge of basal sliding processes. The implication is that initialized ice sheet models therefore have predictive power over shorter times scales of several decades, at least for fast-flowing glaciers such as those in West Antarctica. Further work is required to determine if this holds in other areas, or when other poorly constrained processes such as calving and ice damage are included.

385 *Code availability.* Version 2019b of the source code for *Úa*, sufficient to run the experiments in this project, is available at <http://doi.org/10.5281/zenodo.3706623> (Gudmundsson, 2020). The code is under continuous development, and the latest version is available at <https://github.com/GHilmarG/UaSource>.

*Author contributions.* JMB ran the model simulations and led the writing. Both authors were involved in devising the experiments and editing the manuscript.

390 *Competing interests.* The authors declare no competing interests.

*Acknowledgements.* This work is from the PROPHET project, a component of the International Thwaites Glacier Collaboration (ITGC). Support from Natural Environment Research Council (NERC: Grant NE/S006745/1). This is ITGC Contribution No. 074.

The authors would like to thank Douglas Brinkerhoff and one anonymous reviewer for their insightful feedback which helped to improve the manuscript, and Alexander Robinson for handling the editing of the paper.

395

## References

- Barnes, J. M., Dias dos Santos, T., Goldberg, D., Gudmundsson, G. H., Morlighem, M., and De Rydt, J.: The transferability of adjoint inversion products between different ice flow models, *The Cryosphere*, 15, 1975–2000, 2021.
- 400 Brondex, J., Gagliardini, O., Gillet-Chaulet, F., and Durand, G.: Sensitivity of grounding line dynamics to the choice of the friction law, *Journal of Glaciology*, 63, 854–866, 2017.
- Brondex, J., Gillet-Chaulet, F., and Gagliardini, O.: Sensitivity of centennial mass loss projections of the Amundsen basin to the friction law, *The Cryosphere*, 13, 177–195, 2019.
- Budd, W., Keage, P., and Blundy, N.: Empirical studies of ice sliding, *Journal of glaciology*, 23, 157–170, 1979.
- Budd, W., Janssen, D., and Smith, I.: A three-dimensional time-dependent model of the West Antarctic ice sheet, *Annals of Glaciology*, 5, 405 29–36, 1984.
- Cornford, S. L., Seroussi, H., Asay-Davis, X. S., Gudmundsson, G. H., Arthern, R., Borstad, C., Christmann, J., Dias dos Santos, T., Feldmann, J., Goldberg, D., Hoffman, M. J., Humbert, A., Kleiner, T., Leguy, G., Lipscomb, W. H., Merino, N., Durand, G., Morlighem, M., Pollard, D., Rückamp, M., Williams, C. R., and Yu, H.: Results of the third Marine Ice Sheet Model Intercomparison Project (MISMIP+), *The Cryosphere*, 14, 2283–2301, <https://doi.org/10.5194/tc-14-2283-2020>, <https://tc.copernicus.org/articles/14/2283/2020/>, 2020.
- 410 Cuffey, K. M. and Paterson, W. S. B.: *The physics of glaciers*, Elsevier, 2010.
- De Rydt, J., Reese, R., Paolo, F. S., and Gudmundsson, G. H.: Drivers of Pine Island Glacier speed-up between 1996 and 2016, *The Cryosphere*, 15, 113–132, 2021.
- DeConto, R. M. and Pollard, D.: Contribution of Antarctica to past and future sea-level rise, *Nature*, 531, 591–597, 2016.
- Favier, L., Durand, G., Cornford, S. L., Gudmundsson, G. H., Gagliardini, O., Gillet-Chaulet, F., Zwinger, T., Payne, A., and Le Brocq, 415 A. M.: Retreat of Pine Island Glacier controlled by marine ice-sheet instability, *Nature Climate Change*, 4, 117–121, 2014.
- Fowler, A.: Sliding with cavity formation, *Journal of Glaciology*, 33, 255–267, 1987.
- Gagliardini, O., Cohen, D., Råback, P., and Zwinger, T.: Finite-element modeling of subglacial cavities and related friction law, *Journal of Geophysical Research: Earth Surface*, 112, 2007.
- Gillet-Chaulet, F., Durand, G., Gagliardini, O., Mosbeux, C., Mouginit, J., Rémy, F., and Ritz, C.: Assimilation of surface velocities acquired 420 between 1996 and 2010 to constrain the form of the basal friction law under Pine Island Glacier, *Geophysical Research Letters*, 43, 10–311, 2016.
- Glen, J.: The flow law of ice: A discussion of the assumptions made in glacier theory, their experimental foundations and consequences, *IASH Publ*, 47, 171–183, 1958.
- Gudmundsson, G. H.: GHilmarG/UaSource: Ua2019b (Version v2019b), <http://doi.org/10.5281/zenodo.3706623>, 2020.
- 425 Hill, E. A., Rosier, S. H., Gudmundsson, G. H., and Collins, M.: Quantifying the potential future contribution to global mean sea level from the Filchner–Ronne basin, Antarctica, *The Cryosphere*, 15, 4675–4702, 2021.
- Iken, A.: The effect of the subglacial water pressure on the sliding velocity of a glacier in an idealized numerical model, *Journal of Glaciology*, 27, 407–421, 1981.
- Joughin, I., Smith, B. E., and Medley, B.: Marine ice sheet collapse potentially under way for the Thwaites Glacier Basin, West Antarctica, 430 *Science*, 344, 735–738, 2014.
- Joughin, I., Smith, B. E., and Schoof, C. G.: Regularized Coulomb friction laws for ice sheet sliding: application to Pine Island Glacier, Antarctica, *Geophysical research letters*, 46, 4764–4771, 2019.

- Leguy, G., Asay-Davis, X., and Lipscomb, W.: Parameterization of basal friction near grounding lines in a one-dimensional ice sheet model, *The Cryosphere*, 8, 1239–1259, 2014.
- 435 Lliboutry, L.: General theory of subglacial cavitation and sliding of temperate glaciers, *Journal of Glaciology*, 7, 21–58, 1968.
- MacAyeal, D. R.: Large-scale ice flow over a viscous basal sediment: Theory and application to ice stream B, Antarctica, *Journal of Geophysical Research: Solid Earth*, 94, 4071–4087, 1989.
- Milillo, P., Rignot, E., Rizzoli, P., Scheuchl, B., Mouginot, J., Bueso-Bello, J., and Prats-Iraola, P.: Heterogeneous retreat and ice melt of Thwaites Glacier, West Antarctica, *Science advances*, 5, eaau3433, 2019.
- 440 Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., Eisen, O., Ferraccioli, F., Forsberg, R., Fretwell, P., et al.: Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet, *Nature Geoscience*, 13, 132–137, 2020.
- Mouginot, J., Rignot, E., and Scheuchl, B.: Sustained increase in ice discharge from the Amundsen Sea Embayment, West Antarctica, from 1973 to 2013, *Geophysical Research Letters*, 41, 1576–1584, 2014.
- Mouginot, J., Rignot, E., Scheuchl, B., and Millan, R.: Comprehensive annual ice sheet velocity mapping using Landsat-8, Sentinel-1, and 445 RADARSAT-2 data, *Remote Sensing*, 9, 364, 2017.
- National Academies of Sciences, Engineering, and Medicine: A Strategic vision for NSF investments in Antarctic and southern ocean research, National Academies Press, Washington, DC, 2015.
- Nias, I., Cornford, S., and Payne, A.: New mass-conserving bedrock topography for Pine Island Glacier impacts simulated decadal rates of mass loss, *Geophysical Research Letters*, 45, 3173–3181, 2018.
- 450 Pattyn, F., Schoof, C., Perichon, L., Hindmarsh, R., Bueler, E., De Fleurian, B., Durand, G., Gagliardini, O., Gladstone, R., Goldberg, D., et al.: Results of the marine ice sheet model intercomparison project, MISMIP, *The Cryosphere*, 6, 573–588, 2012.
- Rignot, E., Jacobs, S., Mouginot, J., and Scheuchl, B.: Ice-shelf melting around Antarctica, *Science*, 341, 266–270, 2013.
- Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessem, M. J., and Morlighem, M.: Four decades of Antarctic Ice Sheet mass balance from 1979–2017, *Proceedings of the National Academy of Sciences*, 116, 1095–1103, 2019.
- 455 Scambos, T. A., Bell, R. E., Alley, R. B., Anandakrishnan, S., Bromwich, D., Brunt, K., Christianson, K., Creyts, T., Das, S., DeConto, R., et al.: How much, how fast?: A science review and outlook for research on the instability of Antarctica’s Thwaites Glacier in the 21st century, *Global and Planetary Change*, 153, 16–34, 2017.
- Schoof, C.: The effect of cavitation on glacier sliding, *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 461, 609–627, 2005.
- 460 Shepherd, A., Ivins, E., Rignot, E., Smith, B., Van Den Broeke, M., Velicogna, I., Whitehouse, P., Briggs, K., Joughin, I., Krinner, G., et al.: Mass balance of the Antarctic Ice Sheet from 1992 to 2017, *Nature*, 558, 219–222, 2018.
- Sun, S., Pattyn, F., Simon, E. G., Albrecht, T., Cornford, S., Calov, R., Dumas, C., Gillet-Chaulet, F., Goelzer, H., Gollledge, N. R., et al.: Antarctic ice sheet response to sudden and sustained ice-shelf collapse (ABUMIP), *Journal of Glaciology*, 66, 891–904, 2020.
- Sutterley, T. C., Velicogna, I., Rignot, E., Mouginot, J., Flament, T., Van Den Broeke, M. R., Van Wessem, J. M., and Reijmer, C. H.: 465 Mass loss of the Amundsen Sea Embayment of West Antarctica from four independent techniques, *Geophysical Research Letters*, 41, 8421–8428, 2014.
- Tsai, V. C., Stewart, A. L., and Thompson, A. F.: Marine ice-sheet profiles and stability under Coulomb basal conditions, *Journal of Glaciology*, 61, 205–215, 2015.

- Van Wessem, J. M., Van De Berg, W. J., Noël, B. P., Van Meijgaard, E., Amory, C., Birnbaum, G., Jakobs, C. L., Krüger, K., Lenaerts, J.,  
470 Lhermitte, S., et al.: Modelling the climate and surface mass balance of polar ice sheets using RACMO2–Part 2: Antarctica (1979–2016),  
The Cryosphere, 12, 1479–1498, 2018.
- Weertman, J.: On the sliding of glaciers, *Journal of glaciology*, 3, 33–38, 1957.
- Wernecke, A., Edwards, T. L., Holden, P. B., Edwards, N. R., and Cornford, S. L.: Quantifying the impact of bedrock topography uncertainty  
in Pine Island Glacier projections for this century, *Geophysical Research Letters*, p. e2021GL096589, 2022.
- 475 Yu, H., Rignot, E., Seroussi, H., and Morlighem, M.: Retreat of Thwaites Glacier, West Antarctica, over the next 100 years using various ice  
flow models, ice shelf melt scenarios and basal friction laws, *The Cryosphere*, 12, 3861–3876, 2018.