

# Characteristics of the contemporary Antarctic firn layer simulated with IMAU-FDM v1.2A (1979-2020)

Sanne B.M. Veldhuijsen<sup>1</sup>, Willem Jan van de Berg<sup>1</sup>, Max Brils<sup>1</sup>, Peter Kuipers Munneke<sup>1</sup>, and Michiel R. van den Broeke<sup>1</sup>

<sup>1</sup>Institute for Marine and Atmospheric Research Utrecht, Utrecht University, Utrecht, The Netherlands

**Correspondence:** Sanne B.M. Veldhuijsen (s.b.m.veldhuijsen@uu.nl)

**Abstract.** Firn simulations are essential for understanding Antarctic ice sheet mass change as they enable us to convert satellite altimetry observed volume changes to mass changes, column thickness to ice thickness, and to quantify the meltwater buffering capacity of firn. Here, we present and evaluate a simulation of the contemporary Antarctic firn layer using the updated semi-empirical firn model IMAU-FDM for the period 1979-2020. We have improved previous fresh snow density and firn compaction parameterizations, and used updated atmospheric forcing. In addition, the model has been calibrated and evaluated using 112 firn core density observations across the ice sheet. We found that 63 to 68 % of the seasonal and decadal surface height variability is due to variations in firn air content rather than firn mass. Comparison of simulated surface elevation change with a previously published multi-mission altimetry product for the period 2003-2015 shows that performance of the updated model has improved, notably in Dronning Maud Land and Wilkes Land. However, a substantial trend difference ( $>10 \text{ cm yr}^{-1}$ ) remains in the Antarctic Peninsula and Ellsworth Land, mainly caused by uncertainties in the spin-up forcing. By estimating previous climatic conditions from ice core data, these trend differences can be reduced by 38 %.

## 1 Introduction

The Antarctic ice sheet (AIS) is the largest freshwater reservoir on Earth, and is losing mass since at least 2002 (Shepherd et al., 2018; Rignot et al., 2019), thereby contributing to around 10 % of global average sea level rise since 1993 (Oppenheimer et al., 2019). About 99 % of the AIS is covered by a firn layer (Winther et al., 2001), which represents the transition of fresh snow to glacier ice. The firn layer thickness typically ranges from 40 m in the coastal zone and can exceed 100 m in the interior (van den Broeke, 2008), and also fluctuates in time due to changes in accumulation, melt and firn compaction. A detailed understanding of the firn layer and its variability is important for constraining current and future AIS mass change for three key reasons. Firstly, firn depth and density estimates are required to convert altimetry observed volume-to-mass changes, which remains a major source of uncertainty in mass balance studies (Morris and Wingham, 2015; Verjans et al., 2021). Secondly, firn depth and density estimates are required to convert column thickness to ice thickness, to calculate solid ice discharge over the grounding line (Rignot et al., 2019). Thirdly, firn provides pore space in which nearly all ( $>94 \%$ ) the surface meltwater refreezes or is retained, thereby buffering surface meltwater and preventing mass loss by runoff of surface meltwater (Medley et al., 2022).

25 From both observational and modelling studies, firn layer depth and density are known to vary greatly across the AIS, resulting from the wide range of (near) surface climatic conditions (e.g., Keenan et al., 2021), i.e. temperature, wind speed, accumulation and melt. Low temperatures in the interior (Fig. 1) cause slow densification rates resulting in a firn layer thickness (defined here as the depth of the  $830 \text{ kg m}^{-3}$  density level) greater than 100 m. On the other hand, on warm and dry ice shelves, densification and melt is enhanced, resulting in denser firn layers with a depth less than 50 m (van den Broeke, 2008).

30 Changes of the AIS surface elevation are an expression of changes in the firn layer and of dynamical change of the underlying ice and bedrock. The difficulty is that these seasonal cycles and decadal changes in elevation are not solely caused by fluctuations in mass, but also for a large part by fluctuations in the firn density (Arthern and Wingham, 1998; Ligtenberg et al., 2012; Medley et al., 2022). In altimetry studies, it is therefore important to accurately separate these changes in a mass- and density change component. The mass component can subsequently be separated in a ice-dynamical, bedrock and surface mass  
35 balance component (Willen et al., 2021).

Another important characteristic of the firn layer is its capability to retain meltwater in pore spaces by capillary forces and refreezing. In a warmer future climate enhanced firn compaction, melt, and refreezing can all potentially lead to firn air depletion, thereby limiting the meltwater storage capacity of the firn. This is especially important over the ice shelves, where meltwater accumulation can lead to hydrofracturing-induced ice shelf collapse (Munneke et al., 2014; Datta et al., 2019),  
40 thereby accelerating future Antarctic mass loss and sea level rise (Gilbert and Kittel, 2021). Accurate information about firn conditions is thus essential both for understanding the current AIS mass balance, and to predict its future.

Firn models can fill spatial and temporal gaps between the relatively sparse observations from firn cores and snow pits. While remote sensing products provide high temporal and spatial resolution information about surface height changes, and indirectly, firn properties of the top few meters (e.g., Davis and Poznyak, 1993; Medley et al., 2015; Schröder et al., 2019; Shepherd  
45 et al., 2019), they do not provide information about the deeper firn column. Firn models are, therefore, also important to obtain vertical variations in density, and can be used to perform future simulations of firn depth and density in response to climate change scenarios (Ligtenberg et al., 2012; Munneke et al., 2014).

Firn models can roughly be divided in two classes: physically based and semi-empirical models. Physics-based models describe densification using a constitutive relationship between stress and strain for snow, while semi-empirical models use  
50 physics based densification equations in combination with parameters that are calibrated with observational density profiles. Semi-empirical models neglect some processes such as drifting-snow compaction (e.g., Sommer et al., 2018) and are calibrated with observations representative of the past or present climate, while these conditions might be not representative of a future climate. Nevertheless, semi-empirical models require a smaller number of poorly constrained parameters, such as roughness length, snow grain shape and size, and computational demands are lower (Keenan et al., 2021). This is especially advantageous  
55 for data-scarce regions and large-scale studies. Therefore, semi-empirical firn models have been used widely for correction of satellite altimetry products (e.g., Adusumilli et al., 2018; Smith et al., 2020; Willen et al., 2021) and to assess the meltwater buffering capacity of the Antarctic firn layer (Ligtenberg et al., 2014; Munneke et al., 2014). Several studies have found that the performance of semi-empirical models is comparable to that of physically based models (Steger et al., 2017; Vandecrux et al., 2020; Keenan et al., 2021).

60 Differences in the formulations of, e.g., firn densification and fresh snow density can lead to a substantial spread in modelled  
firn thickness and air content (Lundin et al., 2017; Verjans et al., 2021). Using statistical emulation of firn densification models,  
Verjans et al. (2021) quantified how much different sources of uncertainty contribute to the total uncertainty in modelled firn  
thickness change of the East Antarctic ice sheet. This study shows that, at the basin scale, the ensemble uncertainty in firn  
thickness changes ranges from 0.2 to 1.0 cm yr<sup>-1</sup>, which amounts to a 15-300 % relative uncertainty. While differences in  
65 climate forcing have the largest influence on the spread, in basins with high snowfall and a large spatial variability of climatic  
conditions, firn densification formulation also has a substantial contribution to the spread (up to 46 %). In basins with recent  
increases in snowfall rates the contribution of fresh snow density to the total uncertainty is also substantial (up to 28 %).

In this study, we have improved the semi-empirical IMAU Firn Densification model (IMAU-FDM) (Ligtenberg et al., 2011)  
by updating (i) the fresh snow density, (ii) the firn densification parameterizations, and (iii) by using updated atmospheric  
70 forcing fields. In addition, the model has been calibrated and evaluated using 112 in situ firn core density measurements across  
the AIS. This version 1.2A of IMAU-FDM supersedes the previous version (Ligtenberg et al., 2011). Here, we present the  
simulated contemporary characteristic of the Antarctic firn layer for the period 1979-2020. We focus on spatial, seasonal and  
decadal variability in firn thickness and firn air content (FAC). To evaluate the decadal surface elevation changes and seasonal  
cycles we compare the simulations to a multi-mission remote sensing surface height change product for the period 1992-2015  
75 (Schröder et al., 2019). Finally, we test how sensitive the modelled firn layer is to uncertainties in model variables and spin-up  
procedure.

## 2 Materials and methods

### 2.1 IMAU-FDM

IMAU-FDM is a semi-empirical 1D firn densification model that simulates the time evolution of firn density, temperature,  
80 liquid water content and surface height changes due to firn and SMB processes. The model employs up to 3000 layers of 3 to  
15 cm thickness, which represent the firn properties in a Lagrangian way. Abbreviations of all versions of IMAU-FDM used in  
this study are listed in Table 1. The model was originally developed by Helsen et al. (2008), and updated by Ligtenberg et al.  
(2011) to a version abbreviated here to FDM v1.1. Brils et al. (2022) recently improved IMAU-FDM applied to the Greenland  
ice sheet by including a refined parameterization of the thermal conductivity to version FDM V1.2G. The various versions  
85 of IMAU-FDM have been extensively evaluated against firn density and temperature observations from both Greenland and  
Antarctica (Ligtenberg et al., 2011; Kuipers Munneke et al., 2015; Brils et al., 2022). We further improved the model to FDM  
v1.2A by updating (i) the fresh snow density parameterization and (ii) the firn densification, as discussed below.

#### 2.1.1 Fresh snow density

The fresh snow density ( $\rho_s$ ; kg m<sup>-3</sup>) is an important, but poorly constrained upper boundary condition of firn models. FDM  
90 v1.1 used a fresh snow density parameterization based on Kaspers et al. (2004) with a correction by Helsen et al. (2008),

**Table 1.** Abbreviations and characteristics of IMAU-FDM versions used in this study.

Abbreviation	IMAU-FDM version	Forcing	Fresh snow density	MO <sub>830*</sub> fit
FDM v1.2A	FDM v1.2 Antarctica	RACMO2.3p2, ERA-5	Eq. (2); this study	Power (Eq. 5)
FDM FS-K	FDM v1.2 Antarctica	RACMO2.3p2, ERA-5	Eq. (1); Kaspers et al. (2004)	Power (Eq. 5)
FDM FS-L	FDM v1.2 Antarctica	RACMO2.3p2, ERA-5	Eq. (2); Lenaerts et al. (2012)	Power (Eq. 5)
FDM v1.2A-log	FDM v1.2 Antarctica	RACMO2.3p2, ERA-5	Eq. (2); this study	Log (Eq. 4)
FDM v1.1p1	FDM v1.1	RACMO2.3p1, ERA-Interim	Eq. (1); Kaspers et al. (2004)	Log (Eq. 4)
FDM v1.1p2	FDM v1.1	RACMO2.3p2, ERA-5	Eq. (1); Kaspers et al. (2004)	Log (Eq. 4)

which has been calibrated over Antarctica and yields density values that typically represent the first 0.5 m of the snowpack. It varies as a function of annual average surface temperature ( $\bar{T}_s$ ; K), 10-m wind speed ( $\bar{V}_{10}$ ; m s<sup>-1</sup>) and accumulation ( $\dot{b}$ ; mm w.e. yr<sup>-1</sup>):

$$\rho_s = A + B\bar{T}_s + C\bar{V}_{10} + E\dot{b} \quad (1)$$

95 in which  $A$ ,  $B$ ,  $C$  and  $E$  are fit coefficients. Snow crystal type and degree of riming are indeed temperature dependent, and previous work shows that fresh snow density increases with increasing temperature (Judson and Doesken, 2000). Also, strong winds during snowfall cause crystal breaking, thereby reducing the snow effective grain size (Sato et al., 2008), which results in more efficient packing and increased fresh snow densities. However, at least two concerns can be raised against Eq. (1). First, due to the partly similar spatial patterns of accumulation and temperature in Antarctica, a statistical relationship between fresh  
100 snow density and accumulation may exist, however this relationship has no physical grounds. Other studies have therefore not found a dependency between annual accumulation and fresh snow density in Antarctica, Greenland and the Alps (Lehning et al., 2002; Fausto et al., 2018; Lenaerts et al., 2012). Second, Eq. (1) neglects the impact of the meteorological conditions at the time of deposition, which can be highly variable in time. Therefore, we tested the more recent parameterization of Lenaerts et al. (2012), which has been calibrated over Antarctica, in which  $\rho_s$  is a function of instantaneous surface temperature ( $T_s$ ; K)  
105 and 10-m wind speed ( $V_{10}$ ; m s<sup>-1</sup>):

$$\rho_s = A + BT_s + CV_{10} \quad (2)$$

in which  $A$ ,  $B$  and  $C$  are fit coefficients. The resulting density values are representative of the fresh skin layer density, which typically represents the upper few cm of the snowpack. Within days after depositing, the crystal structure of freshly deposited snow breaks down due to wind and redistribution of drifting snow, which increases the surface snow density (Groot Zwaafink et al., 2013). IMAU-FDM does not calculate this densification by wind and redistribution of drifting snow. The fit coefficients  
110  $A$ ,  $B$  and  $C$  in Eq. (2) for the fresh snow density are recalibrated to improve the fit of the simulated with observed surface snow densities, defined as the top 0.5 m of the firn column, as this best matches the thickness of the sampled layer. These fit coefficients are then used to simulate the fresh snow density. This recalibration is described in Section 3.1. As we only

use 10 additional measurements in the evaluation dataset compared to the calibration dataset, we also include a 10-fold cross  
 115 evaluation.

### 2.1.2 Dry snow densification rate

Over time, the freshly deposited snow becomes denser. The rate of the dry firm densification ( $d\rho/dt$ ) is calculated using the semi-empirical equation of Arthern et al. (2010):

$$\frac{d\rho}{dt} = D\dot{b}g(\rho_i - \rho)e^{\left(\frac{E_c}{RT} - \frac{E_g}{RT_{ave}}\right)} \quad (3)$$

120 where  $D$  is a constant,  $\dot{b}$  is the long-term average accumulation rate ( $\text{kg m}^{-2} \text{y}^{-1}$ ),  $g$  is the gravitational acceleration,  $\rho_i$  is the density of bubble free ice ( $917 \text{ kg m}^{-3}$ ),  $R$  is the gas constant,  $\rho$  is the layer density ( $\text{kg m}^{-3}$ ),  $T$  is the instantaneous layer temperature (K),  $T_{ave}$  is the long-term average surface skin temperature (K), and  $E_c$  and  $E_g$  are the activation energies for creep ( $60 \text{ kJ mol}^{-1}$ ) and grain-growth ( $42.4 \text{ kJ mol}^{-1}$ ), respectively.  $\dot{b}$  is used as a measure of the overburden pressure, thereby assuming that the accumulation rate is constant over time. Using FDM v1.2G for Greenland, Brils et al. (2022) show  
 125 that assuming a constant  $\dot{b}$  introduces a minor error in the load experienced by a layer of firm (e.g.  $<3.2\%$  at Summit and  $<1.9\%$  at Dye-2). The constant  $D$  has different values above ( $0.03$ ) and below ( $0.07$ ) the critical density level  $\rho = 550 \text{ kg m}^{-3}$  to represent the distinct densification mechanisms. For  $\rho < 550 \text{ kg m}^{-3}$ , densification mainly occurs by settling and sliding of grains, and for  $\rho > 550 \text{ kg m}^{-3}$ , it mainly occurs by deformation, sublimation and diffusion (Herron and Langway, 1980).

By comparing simulated and observed depths of the  $550$  and  $830 \text{ kg m}^{-3}$  density levels ( $z_{550}$  and  $z_{830}$ , respectively),  
 130 Ligtenberg et al. (2011) found that Eq. (3) requires correction terms  $MO_{550}$  for  $\rho < 550 \text{ kg m}^{-3}$  and  $MO_{830*}$  for  $550 < \rho < 830 \text{ kg m}^{-3}$ , which are defined as the ratio of modelled and observed values of  $z_{550}$  and  $z_{830*}$ , where  $z_{830*} = z_{830} - z_{550}$ . The correction terms  $MO_{550}$  and  $MO_{830*}$  are added as a multiplier to Eq. (3);  $MO$  values below one reduce the densification rate, and values above one enhance the densification rate.  $MO_{550}$  and  $MO_{830*}$  are chosen as functions of the long-term mean accumulation rate. Ligtenberg et al. (2011) and Brils et al. (2022) used logarithmic correction functions, thus:

$$135 \quad MO = \alpha - \beta \ln(\dot{b}) \quad (4)$$

in which  $MO$  is either  $MO_{550}$  or  $MO_{830*}$ , and  $\alpha$  and  $\beta$  are fit coefficients which differ for  $MO_{550}$  and  $MO_{830*}$ . In FDM v1.1 a minimum value of  $0.25$  is imposed on  $MO_{550}$  and  $MO_{830*}$ . For Greenland and Antarctica, different values for  $\alpha$  and  $\beta$  have been used to optimize densification. Here, we also tested a power-law function for  $MO_{830*}$ :

$$MO_{830*} = \delta \dot{b}^{-\epsilon} + \phi \quad (5)$$

140 in which  $\delta$ , and  $\epsilon$  and  $\phi$  are fit coefficients. The fit coefficients in Eq. (4) and (5) are calibrated, which is described in Section 3.2. As we only use 10 additional measurements in the evaluation dataset compared to the calibration dataset, we also include a 10-fold cross evaluation.

### 2.1.3 Heat conduction, meltwater percolation and refreezing

The conduction of heat is simulated by using a one-dimensional heat transfer equation, which couples vertical heat conduction to temperature gradients through the thermal conductivity of firn. In FDM v1.1 the thermal conductivity is a function of firn density only (Anderson, 1976), and in FDM v1.2G this has been extended to a function of density and temperature (Calonne et al., 2019), which is also included in FDM v1.2A. The lower boundary condition assumes a constant heat flux across the lowest layer, where the deep temperature is allowed to change along with long-term changes in surface temperature or internal heat release. Meltwater percolation is simulated using the bucket method, whereby each firn layer has a maximum irreducible water content that decreases with increasing density (Coléou et al., 1999). The meltwater can percolate through all layers in a single timestep. The retained meltwater refreezes when it reaches a layer with a temperature below the freezing point. To capture the percolation and refreezing of meltwater more accurately, the model timestep is reduced from 3600 to 300 s for locations with melt. Further details and evaluation of heat conduction, percolation and refreezing in IMAU-FDM are provided by Brils et al. (2022).

## 2.2 Model strategy

An initial firn layer is obtained by spinning up the model over a reference period until the firn layer is in equilibrium with the surface climate. The equilibrium is roughly reached when the entire firn column, defined here as the depth where the density  $\rho$  reaches  $830 \text{ kg m}^{-3}$ , is fully refreshed by accumulation. The required spin-up time in years is therefore calculated as the total mass of the modelled firn layer divided by the annual accumulation in w.e. (Kuipers Munneke et al., 2015). As no obvious strong long-term trends have been detected in Antarctica's surface climate and SMB during the period 1979-2020 (e.g., Favier et al., 2017; Mottram et al., 2021), and no reliable surface climate fields are available prior to 1979, the spin-up forcing is obtained by looping over the 1979-2020 forcing data. Thereby, we make the rather strong assumption that this 42-year period is representative for the past 100-1000 years and that the firn layer is in equilibrium with that climatic period. In the actual simulation after the spin-up, a minor trend ( $<0.6 \text{ mm yr}^{-1}$  averaged over the AIS) in total FAC remains, because at the bottom of the column, ice with a density between  $830 - 917 \text{ kg m}^{-3}$  slowly replaces bubble free ice ( $\rho = 917 \text{ kg m}^{-3}$ ), which originates from the initialization of the firn column prior to the spin-up. This trend is removed before further analysis of the results.

### 2.3 Surface elevation change

IMAU-FDM quantifies the impact of firn and SMB processes on the depth of the firn layer. The resulting vertical velocity of the firn top surface ( $v_{tot}$ ) is defined as the sum of these components:

$$v_{tot} = v_{snow} + v_{sub} + v_{snd} + v_{melt} + v_{fc} + v_{by} + v_{ice} \quad (6)$$

where  $v_{snow}$  represents the vertical velocity component as a result of snowfall,  $v_{sub}$  of sublimation/riming,  $v_{snd}$  of snowdrift erosion/deposition,  $v_{melt}$  of snowmelt,  $v_{fc}$  of firn compaction and  $v_{by}$  represents the vertical motion associated with the changing ice shelf draft when the mass of the firn layer changes.  $v_{ice}$  represents the downward/upward movement of the surface by

the local divergence/convergence of the ice flow, driven by the long-term vertical mass flux through the lower boundary of the  
175 firn column. In a steady-state firn layer, this equals the mass flux at the upper boundary.  $v_{ice}$  is therefore equal to the long-term  
average annual SMB ( $\text{kg m}^{-2} \text{yr}^{-1}$ ) over the reference period divided by  $\rho_i$ , but of opposite sign. All components are derived  
from run-time output of IMAU-FDM. The minor residual trend in  $dH/dt$  due to increasing air content in the ice below the firn  
layer is added to  $v_{ice}$ .  $v_{by}$ , only relevant over ice shelves, is equal to the negative change in firn mass divided by the density of  
sea water.


## 180 2.4 RACMO2.3p2 forcing

IMAU-FDM is forced at the upper boundary with 3-hourly fields of snowfall, sublimation, snowdrift erosion, 10-m wind  
speed, surface temperature, snowmelt and rainfall. In this study, FDM v1.2A is driven with ERA-5 reanalysis data (Hersbach  
et al., 2020), dynamically downscaled with the regional atmospheric climate model RACMO2.3p2 (Van Wessem et al., 2018).  
RACMO2.3p2 is run with a horizontal resolution of 27x27 km and the simulation covers the period 1979-2020. RACMO2.3p2  
185 has been thoroughly evaluated for the Antarctic and Greenland ice sheets by Van Wessem et al. (2018) and Noël et al. (2018),  
respectively. In comparison to RACMO2.3p1, RACMO2.3p2 employs upper-air relaxation, has updated topography, and a  
retuned cloud scheme and snow properties. Upper-air relaxation is the indiscriminate nudging applied to the upper part of the  
atmosphere only. The most important effects of these changes are (i) increased snowfall in the ice sheet interior, (ii) reduced  
snowdrift sublimation and (iii) increased snowmelt (Van Wessem et al., 2018). Several Antarctic studies use output from FDM  
190 v1.1 forced with RACMO2.3p2 (e.g., Schröder et al., 2019; Keenan et al., 2021), but the impact of the improved forcing has  
not yet been described. As snowfall and surface melt are defining drivers of the state of the firn, we also assess the impact of  
the updated forcing in this paper. In addition to the updates of RACMO2.3p2, previous FDM simulations for Antarctica used  
dynamically downscaled ERA-Interim reanalysis data, which has been replaced by ERA-5 in 2019. This results in an improved  
forcing. For example, surface mass balance anomalies in Dronning Maud Land from 2006 to 2018 are better captured in ERA-5  
195 compared to ERA-Interim (Gossart et al., 2019). We expect that forcing differences in surface mass balance have a larger impact  
on the temporal evolution of the surface height compared to adjustments in the fresh snow density and dry snow densification  
parameterizations (Verjans et al., 2021).

## 2.5 Observational data

To calibrate and evaluate the firn model, we collected published firn density profiles from widely varying locations across  
200 the AIS (Fig. 1). We used 112 density profiles from firn cores and 8 density profiles from neutron density probe measure-  
ments, combining multiple datasets (van den Broeke, 2008; Schwanck et al., 2016; Bréant et al., 2017; Fernandez et al., 2010;  
Montgomery et al., 2018; Fourteau et al., 2019; Olmi et al., 2021; Winstrup et al., 2019). The firn cores and neutron density  
probe measurements are mostly obtained in summer months between 1980 and 2020. For calibrating the fresh snow density  
parameterization, observations of the density of surface snow, defined as the top 0.5 m, from 61 firn cores could be used. Eight  
205 additional surface snow density values from neutron density probe measurements and two from firn cores (Montgomery et al.,  
2018) were added after calibration, and are thus only used for evaluation. We used densities of the top 0.5 m, as this depth

is representative of many in situ surface snow density measurements, it generally consists of snow from multiple snowfall events, and it captures the densification by wind and redistribution of drifting snow (Brun et al., 1997). For calibrating the MO fits, only dry firn cores are used, as Eq. (3) describes dry snow densification only. A firn core is considered dry if its location experiences on average less melt than 5 % of the average annual accumulation. For the MO fits, 93 dry firn cores could be used, which improves upon Ligtenberg et al. (2011), who used data from 48 dry firn cores. In addition to the 93 dry cores used for calibration of the MO ratios, another 10 wet cores and five additional dry cores were used to evaluate the  $z_{550}$  and  $z_{830}$  model output. For evaluating FAC output, 31 firn cores could be used. Table S1 in the Supplement lists all measurements that have been used, the corresponding coordinates, method and citation.



**Figure 1.** Map of Antarctica showing annual average surface temperature and the locations of the in situ density observations. The yellow circles indicate observational locations of the depths of the critical density level of  $\rho = 550 \text{ kg m}^{-3}$  ( $z_{550}$ ), the white circles of the depths of the critical density levels of both  $\rho = 550 \text{ kg m}^{-3}$  ( $z_{550}$ ) and  $\rho = 830 \text{ kg m}^{-3}$  ( $z_{830}$ ), the purple circles of the depths of the critical density levels of both  $\rho = 550 \text{ kg m}^{-3}$  ( $z_{550}$ ) and  $\rho = 830 \text{ kg m}^{-3}$  ( $z_{830}$ ), and of the firn air content (FAC), and the black crosses of the surface snow densities (top 0.5 m of the firn column). The numbers indicate regions mentioned throughout the manuscript.

In order to verify whether the modelled surface elevation trends and seasonal variability align with observations, we used a multi-mission satellite altimetry product covering the period January 1992 to December 2015 as presented by Schröder et al. (2019). In this product, data from the ERS-1, ERS-2, Envisat, ICESat, and CryoSat-2 missions are combined into a monthly time series on a  $10 \times 10 \text{ km}$  grid, with the polar gap, floating ice shelves and steep terrain areas being excluded. Validation of this dataset with in situ and airborne observations showed that the products are successfully combined (Schröder et al., 2019). The altimetry product has been resampled to the FDM grid using bilinear resampling. The measurement precision over flat terrain



ranges from 5 to 15 cm depending on the satellite. However, as the precision decreases with increasing surface slope, the uncertainty can reach several meters in the coastal regions with steep topography. Offsets due to varying penetration depth of the radar signals (thus relevant for ERS-1, ERS-2, Envisat and CryoSat-2) caused by variations in electromagnetic properties of the ice sheet are not included in this uncertainty estimate. These additional offsets can reach up to several centimeters per year, and can for instance increase the seasonal amplitude at continent-wide scales by up to several centimeters (Nilsson et al., 2022). While corrections are applied to account for variations in penetration depths, these corrections are in many cases unable to fully correct the artificial signals, which can e.g. be seen from the intermission variation in seasonal amplitude (Schröder et al., 2019; Nilsson et al., 2022). On the other hand, signal penetration is negligible for laser measurements (such as ICESat, ICESat-2, Operation IceBridge). However, ICESat and Operation IceBridge datasets have poor spatial and temporal coverage, and we believe that the record of ICESat-2 is currently is too short (only two years and two months overlap with FDM v1.2A, and no overlap with FDM v1.1p1) to provide a viable evaluation. Therefore, we used the multi-mission satellite altimetry product as presented by Schröder et al. (2019), which has a good spatial and temporal coverage. Our firm simulations only provide elevation change due to anomalies in firn thickness caused by variations in e.g. snow accumulation, surface temperature, firn densification and melt. The difference between the observed and simulated elevation change therefore represents a combination of elevation changes due to ice-dynamical imbalance, bedrock motion, errors in the firn model and its climate forcing and errors in the altimetry observations. For this comparison, we therefore exclude regions with ice-dynamical imbalance, by using the ice-dynamical imbalance mask of Shepherd et al. (2019), which mainly consists of Pine Island, Thwaites, Totten and Getz glacier basins and the Kamb ice stream.

## 2.6 Sensitivity analysis

To study the model sensitivity to input and parameter uncertainty we performed a sensitivity analysis for the 106 locations shown in Figure S1a. These locations include 95 locations from firn core sites (the firn cores used are located at 95 unique grid points, excluding five sites that were later added for evaluation). To improve the representativeness of the climatic conditions over the AIS, we used 11 additional locations in this analysis that are located in high-accumulation or low-temperature regions, as these areas are underrepresented in the 95 firn core sites (Figs. S1b,c).

All the performed experiments are listed in Table 2. Verjans et al. (2021) and Lundin et al. (2017) showed that differences in accumulation, temperature, firn densification formulation and fresh snow density can lead to a substantial spread in modelled firn thickness and air content. For each of these components, we separately performed a model sensitivity test, which includes the spin-up as well as the main run. The fresh snow density was varied with the RMSE from the evaluation with in situ measurements ( $30 \text{ kg m}^{-3}$ ; Section 3.1). To test the sensitivity to uncertainty of the dry snow densification rate, we perform a simulation where we use the 95 % confidence interval boundaries of the MO fits (Eqs. 4 and 5, Fig. 3). To test the sensitivity to uncertainty in accumulation and temperature forcing, we use the standard deviation of snowfall and temperature among regional climate models and reanalysis products (Carter et al., 2022) (Fig. S1). This spread varies spatially and we vary the accumulation with the ratio of the ensemble standard deviation to the ensemble mean of each location, and vary the temperature with the ensemble standard deviation.

255 Another source of forcing uncertainty is the climate forcing during the spin-up period. As explained in Section 2.2, the  
spin-up forcing is obtained by looping over the 1979-2020 forcing data. However, firm core (Thomas et al., 2015, 2017; Medley  
and Thomas, 2019) and isotopes (Stenni et al., 2017) studies show that in the Antarctic Peninsula and Ellsworth Land, the  
accumulation and temperature were typically about 10 % and 1 K lower during the last centuries. In addition, over the remainder  
of the AIS, the accumulation and temperature were about typically 5 % and 0.5 K higher or lower during the last centuries. To  
260 investigate the typical effect this has on the IMAU-FDM results, we include schematic sensitivity tests in which we adjusted  
the accumulation and temperature forcing with these values only during the spin-up period. To mimic a gradual increase in  
precipitation over the Antarctic Peninsula and Ellsworth Land, we performed a spin-up experiment in which the precipitation  
up until the third-to-last loop (the 42 year long reference period) is decreased by 10 %, in the second-to-last by 6.66 %, and in  
the last by 3.33 % with respect to the mean precipitation of 1979-2020. To mimic a gradual increase in temperature over the  
265 Antarctic Peninsula and Ellsworth Land, we create a spin-up experiment in which the temperature up until the third-to-last is  
decreased by 1 K, in the second-to-last by 0.66 K, and in the last by 0.33 K. Over the remainder of the AIS accumulation and  
temperature were both decreased and increased with 5 % and 0.5 K during the spin-up period. As the uncertainties between  
temperature and accumulation during the spin-up period are dependent, we also performed tests in which we simultaneously  
adjusted the temperature and precipitation.

270 For all our experiments, except for the MO fits, we re-ran our model calibration procedure to get the optimal MO fits. We  
then ran the model with these updated MO fits and compared the outputs to our reference FDM v1.2A run. We investigate the  
impact of the uncertainties on the FAC and surface elevation change. As the model set up of FDM v1.2A assumes a steady-state  
firm layer, there is no net surface elevation trend in this simulation over the period 1979-2020, except for sensitivity tests in  
which the accumulation and temperature during the spin-up period are adjusted. In the remainder of the sensitivity experiments,  
275 we therefore compare surface elevation change over the randomly selected period 2015-2020, instead of over 1979-2020. By  
relating the sensitivities of FAC and surface elevation change of each experiment to climatic variables at the sample locations,  
we can expand the uncertainty estimates over the entire ice sheet. The empirical relations for each sensitivity experiment are  
included in Figure S3. The uncertainties of each test were combined quadratically to obtain a value for the total uncertainty,  
where for the spin-up experiments only the combined temperature and accumulation experiments are included.

## 280 **3 Calibration and model performance**

### **3.1 Fresh snow density**

To calibrate and evaluate the fresh snow density parametrization, we performed simulations with FDM v1.2A using the pa-  
rameterizations of Kaspers et al. (2004) (Eq. 1), Lenaerts et al. (2012) (Eq. 2), the latter with updated constants, and of FDM  
v1.1p1. These simulations are abbreviated as FDM FS-K, FDM FS-L, FDM v1.2A and FDM v1.1p1, respectively. The fit  
285 coefficients and statistics of the simulations are listed in Table 3. Figure 2a shows the modelled against observed surface snow  
densities. FDM FS-K somewhat overestimates the surface snow densities (bias = 17.6 kg m<sup>-3</sup>), especially for high surface den-  
sity, and FDM v1.1p1 to a lesser extent (bias = 10.3 kg m<sup>-3</sup>), where the update in the atmospheric forcing results in a poorer

**Table 2.** Overview of the sensitivity experiments. The inputs and parameters are adjusted by subtracting and adding the uncertainties either during the entire run or only during the spin-up. The values used are based on the references provided or on analysis within this work.

Experiment name	Variable	Variation	When	Reference
MO fits	MO fits	+ and -95 % confidence interval	entire run	Figs. 3a,b
$\rho_s$	$\rho_s$	+ and -30 kg m <sup>-3</sup>	entire run	RMSE evaluation Section 3.1
$T_s$	$T_s$	+ and - spatial variable K	entire run	$\sigma$ from Carter et al. (2022)
b	b	+ and - spatial variable %	entire run	$\sigma$ from Carter et al. (2022)
spin-up $T_s$	$T_s$	+0/+1 K and -1/-0.5 K <sup>a</sup>	spin-up	Stenni et al. (2017)
spin-up b	b	+0/+10% and -10/-5 % <sup>a</sup>	spin-up	Thomas et al. (2017)
spin-up $T_s$ , b	b and $T_s$	+0/+5 %, +0/+0.5 K and -10/-5 %, -1/-0.5 K <sup>a</sup>	spin-up	Stenni et al. (2017); Thomas et al. (2017)

<sup>a</sup>The values on the left side of the slash indicate the variation for the Antarctic Peninsula and Ellsworth Land, and the values on the right side of the slash for the remainder of the AIS. Temperature and accumulation over the Antarctic Peninsula and Ellsworth Land are the second-to-last loop of spin-up reduced by 6.66 % and 0.66 K and in the last loop with 3.33 % and 0.33 K.

performance of FDM FS-K compared to FDM v1.1p1. FDM FS-L substantially underestimates surface snow densities (bias = -45.1 kg m<sup>-3</sup>), but has a higher correlation with the observations than FDM FS-K. The degree of underestimation increases  
290 with increasing wind speed, which suggests that a higher value of fit coefficient C is needed to properly capture the impact of wind compaction. This aligns with the fact that IMAU-FDM does not include densification by wind packing. In the optimized fit coefficients (FDM v1.2A, Table 3) the value of C is 2.6 times higher than the value reported by Lenaerts et al. (2012). The wind dependency is also 2.4 times stronger than proposed by Kaspers et al. (2004), which is in agreement with Sugiyama et al. (2012) who found a factor 2.8. The RMSE of the surface snow density of FDM v1.2A is respectively 28 % and 23 % lower  
295 compared to FDM FS-K and FDM v1.1p1, due to both minimizing the bias and improving the representation of the variability. Compared to FDM FS-K, the surface snow density is reduced by 18 kg m<sup>-3</sup>, especially in the high-accumulation margins, and to a lesser extent in windy escarpment regions (Fig. 2b). With the 10-fold cross evaluation for FDM v1.2A, we found that the RMSE for the surface snow density slightly increases to 30.4 kg m<sup>-3</sup> (compared to 28.8 kg m<sup>-3</sup> for the reference FDM v1.2A). The fresh snow density parameterization for Greenland used in FDM v1.2G, is only a function of annual average temperature  
300 (Brils et al., 2022), owing to a lack of co-located surface snow density, temperature and wind speed observations on the GrIS. This contrasts to the expressions used here, which includes a dependency on instantaneous wind speed and temperature. For all sensitivity tests presented here, we calculated the associated MO fits. However, the impact of the different MO fits on the surface snow densities is small (<0.1 %).


### 3.2 Dry snow densification rate

305 To calibrate the dry snow densification rate we first performed a simulation of FDM v1.2A without MO corrections. The resulting MO fits and statistics are listed in Table 4 and shown in Figures 3a,b. The optimal MO<sub>550</sub> fit is less steep with respect to accumulation ( $\beta = 0.117$  vs 0.151) than the one in FDM v1.1p1. The fresh snow density is now independent of accumulation

**Table 3.** Fit coefficients and evaluation of the fresh snow density parameterizations (Eqs. 1,2). The fit coefficients of FDM FS-K and FDM FS-L are taken from Kaspers et al. (2004) and Lenaerts et al. (2012), respectively.

Version	A ( $\text{kg m}^{-3}$ )	B ( $\text{kg m}^{-3} \text{K}^{-1}$ )	C ( $\text{kg s m}^{-4}$ )	E ( $\text{kg yr m}^{-3} \text{mm w.e.}^{-1}$ )	$R^2$	RMSE ( $\text{kg m}^{-3}$ )	Bias ( $\text{kg m}^{-3}$ )
FDM v1.2A <sup>b</sup>	83	0.77	11.67		0.46	28.8	-1.3
FDM FS-K <sup>a</sup>	-77	1.5	6.8	0.075	0.39	40.2	17.6
FDM FS-L <sup>b</sup>	97.5	0.77	4.49		0.41	55.3	-45.1
FDM v1.1p1 <sup>a</sup>	-77	1.5	6.8	0.075	0.37	37.3	10.3

<sup>a</sup>using annual average accumulation, temperature and wind speed. <sup>b</sup>using instantaneous temperature and wind speed.



**Figure 2.** (a) Scatter plot of simulated against observed surface snow density using FDM v1.2A, FDM FS-K (Kaspers et al., 2004), FDM FS-L (Lenaerts et al., 2012) and FDM v1.1p1. (b) Map of average difference between the simulated surface snow densities of FDM v1.2A and FDM FS-K for the locations with observations.

and generally lower, and hence more densification is needed at high-accumulation sites to match the density measurements. This might explain the slightly deteriorated correlation of  $\text{MO}_{550}$ . For the  $\text{MO}_{830*}$  regression, we found that a power fit (FDM v1.2A) compared to a logarithmic fit (FDM v1.2A-log) improves correlation ( $R^2 = 0.88$  compared to 0.83). Furthermore, a minimum value is now explicitly derived with the power fit, so that a rather arbitrary minimum value no longer needs to be prescribed. When the locations are rerun using the new MO values with the power fit for  $\text{MO}_{830*}$ , the resulting RMSE of the modelled  $z_{830*}$ ,  $z_{830}$  (firm thickness) and FAC are respectively 39, 36 and 49 % lower compared to the logarithmic fit (Figs. 3c,d). The update in atmospheric forcing without updated MO calibration does not result in an improvement in FAC (shown by the similar RMSE of FDM v1.1p1 and FDM v1.1p2). Also, the RMSE of FAC simulated with FDM v1.2A is respectively


**Table 4.** Fit coefficients and statistics of the MO fits (Eqs. 4,5). The fit coefficients and statistics of FDM v1.1p1 are taken from Ligtenberg et al. (2011). The FDM v1.2A-log uses the settings of FDM v1.2A; only Eq. (3) is optimized using a logarithmic  $MO_{830^*}$  fit.

Version	Fit	$\alpha$	$\beta$	$\sigma$	$\epsilon$	$\phi$	$R^2$	RMSE (m)
$MO_{550}$ FDM v1.1p1	Log (Eq. 4)	1.435	0.151				0.43	
$MO_{550}$ FDM v1.2A	Log (Eq. 4)	1.288	0.117				0.37	
$MO_{830^*}$ FDM v1.1p1	Log (Eq. 4)	2.366	0.292				0.71	
$MO_{830^*}$ FDM v1.2A-log	Log (Eq. 4)	2.504	0.334				0.83	0.13
$MO_{830^*}$ FDM v1.2A	Power (Eq. 5)			6.387	0.477	0.195	0.88	0.11

21 and 23 % lower compared to FDM v1.1p1 and FDM v1.1p2, which indicates smaller improvement compared to FDM v1.2A-log (49 %). This is caused by the difference in MO fit between FDM v1.2A-log and FDM v1.1 (Fig. 3b), which results from a combination of the larger firn core dataset and updated atmospheric forcing.

Using a power compared to logarithmic fit results in higher  $MO_{830^*}$  values for low ( $<400 \text{ mm yr}^{-1}$ ) and high ( $>1,400$   
320  $\text{mm yr}^{-1}$ ) accumulation regions, and lower  $MO_{830^*}$  values for intermediate accumulation regions ( $400\text{-}1,400 \text{ mm yr}^{-1}$ ) (Fig. 3b). Therefore, the densification rate is increased in high and low-accumulation regions and reduced in intermediate accumulation regions. This results in decreased firn thickness in high and low-accumulation regions, and in increased firn thickness in intermediate accumulation regions (Fig. 3d). Especially large differences are found in high-accumulation areas where the MO fit curve decreases asymptotically towards zero. With the 10-fold cross evaluation for FDM v1.2A, we found that the  
325 RMSE for  $z_{550}$  slightly increases to 2.33 m (compared to 2.26 m for the reference FDM v1.2A), and the RMSE of  $z_{830}$  to 9.14 m (compared to 8.87 m for the reference FDM v1.2A).

In FDM v1.2G, the version for the Greenland ice sheet, a logarithmic  $MO_{830}$  fit was applied. Dry firn cores from the GrIS cover a smaller range of average annual accumulation,  $80\text{-}680 \text{ mm yr}^{-1}$  compared to  $20\text{-}960 \text{ mm yr}^{-1}$  in Antarctica. If we only include cores from the  $80\text{-}680 \text{ mm yr}^{-1}$  accumulation range, we find a similar  $R^2$  value for the logarithmic vs power  
330 fit (0.67 vs 68). Another difference is that in FDM v1.2G, an almost constant value of 0.67 for the  $MO_{550}$  fit was found, which implies a linear correlation between the densification rate and the accumulation rate. In FDM v1.2A we do find a moderate ( $R^2=0.37$ ,  $\beta = 0.12$ ) correlation of  $MO_{550}$  with accumulation, i.e. AIS densification rate depends more strongly on accumulation than temperature compared to the GrIS. Again, if we only include cores from the  $80\text{-}680 \text{ mm yr}^{-1}$  accumulation range this correlation weakens ( $R^2=0.08$ ,  $\beta = 0.07$ ).




**Figure 3.** Scatter plots defining the MO ratios for (a)  $z_{550}$  and (b)  $z_{830}$  as a function of the annual average accumulation including various MO fits. The blue shaded areas indicate the 95 % confidence intervals. (c) Scatter plots of the simulated against observed  $z_{830}^*$  ( $Z_{830}-Z_{550}$ ) for FDM v1.2A and FDM v1.2A-log. (d) Scatter plots of the simulated against observed FAC for FDM v1.2A, FDM v1.2A-log, FDM v1.1p1 and FDM v1.1p2. (e) Map of the difference between simulated firn thickness ( $z_{830}$ ) of the two tests (FDM v1.2A – FDM v1.2A-log) for locations with observations. The statistics and fits of the MO fits are listed in Table 4.

## 335 4 Spatial and temporal variability of AIS firn depth and density

### 4.1 Spatial density variability

Using the improved firn model, we simulate the spatial and temporal patterns of the AIS firn characteristics at 27 km horizontal resolution. The spatial patterns of the densities, including observed values, are shown in Figure 4. Figure 4a shows the surface snow density (top 0.5 m), which is on average  $376 \text{ kg m}^{-3}$ . The patterns are primarily a reflection of the variation in fresh snow density (eq. 2), with the lowest values on the cold and calm East Antarctic plateau and higher values in the windy escarpment regions and along the warm coastal margins. In addition to this, the surface snow density further increases over the ice shelves,

340




**Figure 4.** Maps from FDM v1.2A of the average (a) surface snow density (defined as the top 0.5 m), depths of critical density levels of (b)  $\rho = 550 \text{ kg m}^{-3}$  ( $z_{550}$ ) and of (c)  $\rho = 830 \text{ kg m}^{-3}$  ( $z_{830}$ ) (the firn thickness), and (d) firn age at the critical density level of (c)  $\rho = 830 \text{ kg m}^{-3}$ . Circles indicate the values found by field measurements.

where the average surface snow density amounts to  $414 \text{ kg m}^{-3}$ , and over the coastal margins, where intermittent melting and refreezing of retained meltwater increases the near-surface density.

In line with observations (Winther et al., 2001), nearly all (>99 %) of the AIS is covered by a layer of firn. This represents the ice sheet fraction where average SMB is positive. The spatial pattern of the depths of the critical density levels  $z_{550}$  and  $z_{830}$  are shown in Figures 4b and c, which are on average 14.1 and 73.8 m. The patterns are roughly the inversed pattern of the surface snow density, which constitutes the upper boundary of the depth-density profile. For simplicity, henceforth we define the firn thickness as  $z_{830}$ . The patterns vary spatially across climatic regions with temperature as a primary driver and accumulation and surface melt as secondary drivers. The highest  $z_{550}$  and  $z_{830}$  values (30.6 and 114.2 m, respectively) are found in the interior, where the densification rate is low due to low temperatures, and the surface snow density is low. Generally,  $z_{550}$  and  $z_{830}$  are small along the warm coastal margins due to faster densification, higher surface snow densities and melt. However, especially the  $z_{830}$  values can be relatively high in high-accumulation regions along the coast, particularly in West Antarctica and the Antarctic Peninsula. The high accumulation rates in these regions result in less densified snow with depth and thus thick firn layers despite relatively high temperatures. The  $z_{550}$  and  $z_{830}$  values are lowest in relatively warm and dry regions on and around Ross, Filchner Ronne and Amery ice shelf, where fresh snow is buried more slowly, and the densification rates are high due to relatively high temperatures.

Figure 4d shows the age of firn at the pore close off depth, here assumed to equal  $z_{830}$ , which is on average 754 yr. The combination of strongly increasing firn depth and decreasing accumulation towards the interior leads to firn ages at close-off depth of up to 3240 yr in central East Antarctica, whereas the pore close-off firn age in warm and wet coastal margins can be as low as 20 yr. If we compare this to observations, we find an  $\text{RMSE} = 231 \text{ yr}$ ,  $R^2 = 0.99$ . On average, the RMSE is 25 % of the observed values. In comparison, RMSE of  $z_{830}$  is 15 % of the observed values and  $R^2 = 0.73$ . These simulations add to our understanding of the large spread in timescales of the firn.



**Figure 5.** Maps from FDM v1.2A of average (a) firn air content (FAC), (b) the difference in FAC (FDM v1.2A - FDM v1.1p1), (c) peak-to-peak FAC difference, (d) relative peak-to-peak FAC difference compared to the average FAC from (a). The peak-to-peak is defined as the difference between the highest and the lowest FAC value occurring during the study period (1979-2020).

## 4.2 Firn air content

Firn air content (FAC), the total (vertically integrated) pore space, is an important output of firn models as it is used to convert  
 365 observed volume changes to mass changes, column thickness to ice thickness, and is an indicator of the meltwater buffering  
 capacity of the firn. FAC is expressed in m and represents the change in depth that would occur if the firn layer would be  
 compressed to the density of glacier ice. The average FAC of the AIS according to IMAU-FDM is 22.8 m, and it varies  
 spatially between 0 to 40 m across climatic regions (Fig. 5a). The pattern resembles the  $z_{830}$  pattern (Fig. 4c); thick firn layers  
 contain the most air. High values are found in the cold interior and along parts of the coast, particularly in West Antarctica, and  
 370 in areas with high accumulation. Low values are found along the warmer coastal margins, with ice shelves having an average  
 FAC of 15.9 m. These average FAC estimates are 5 % and 7 % lower compared to values simulated with the CFM forced with  
 MERRA-2 climate data of 24.0 and 17.0 m respectively (Medley et al., 2022).

The average FAC over the entire AIS (including the ice shelves) simulated with FDM v1.2A is 0.6 m higher than simulated  
 with FDM v1.1. The spatial FAC differences between FDM v1.2A and FDM v1.1 are shown Figure 5b. In the highly elevated  
 375 and low-accumulation regions the new FAC is lower, whereas in the intermediate accumulation regions the new FAC is higher.  
 This implies that the range of FAC is smaller and less spatial variable than previously modelled. The differences in FAC  
 pattern can be explained by differences in (i) the  $MO_{830^*}$  fit, (ii) the fresh snow density, (iii) snowfall and (iv) snowmelt. FDM  
 v1.2A has higher  $MO_{830^*}$  values for low- and high-accumulation regions compared to FDM v1.2A, which results in a faster  
 densification and therefore lower FAC values. In intermediate accumulation regions the  $MO_{830^*}$  values are lower, resulting in  
 380 a slower densification and higher FAC values. The fresh snow densities of FDM v1.2A are on average lower ( $19.1 \text{ kg m}^{-3}$ ,  
 Fig. 2), which contributes to FAC values being higher on average. On top of that, increased accumulation in the interior and  
 increased snowmelt along the coastal margins in RACMO2.3p2 compared to RACMO2.3p1 (Van Wessem et al., 2018), also  
 contributes to the general pattern of higher FAC in the interior and lower values along the coastal margins. However, in the



highly elevated and lowest accumulation regions, we find a reduction in FAC, which implies that the  $MO_{830^*}$  update outweighs  
385 the effects of higher accumulation and lower fresh snow density in this region.


Figures 5c and d focus on temporal variability in FAC. Figure 5c shows the peak-to-peak FAC difference, which represents  
the difference between the highest and the lowest FAC value occurring during the study period (1979-2020). The peak-to-peak  
FAC difference is low in the interior, less than 0.1 m, despite the high FAC. Along the coastal margins, the peak-to-peak FAC  
difference is higher, up to 5 m, even though the FAC is generally lower. Therefore, the relative peak-to-peak FAC difference  
390 is low in the interior (<1 %) and high near the ice sheet margins (>5 %) (Fig. 5d). The average relative peak-to-peak FAC  
difference for the ice shelves, excluding the two largest ones, the Ross and Ronne-Filchner ice shelves, amounts to 13 %. This  
high peak-to-peak variability is caused by a combination of substantial temporal variability in accumulation and melt. The  
temporal FAC variability and the driving processes are discussed further in Sections 4.3 and 4.4 below.

### 4.3 Seasonal firn air content and firn depth variability

395 In order to better understand the FAC temporal variability and its drivers, we decompose the signal into seasonal and interannual  
contributions. Figure 6a shows the average seasonal cycle of the vertical firn surface velocity ( $v_{tot}$ ) including its components  
from Eq. (6). Four main processes drive the seasonality of the firn thickness: snowfall ( $v_{snow}$ ), firn densification ( $v_{fc}$ ), melt  
( $v_{melt}$ ) and sublimation ( $v_{sub}$ ). Averaged over the ice sheet, snowfall is highest in winter, while firn densification, melt and  
sublimation peak in summer. The net result is that the firn thickness and FAC steadily increase over the winter, due to enhanced  
400 accumulation and reduced firn densification, melt and sublimation (Fig. 6b). This is followed by a sharp decrease over the  
summer due to reduced accumulation and increased firn densification and melt. The firn height change by ice flux divergence  
( $v_{ice}$ ) is constant by definition, while the AIS average impacts of snowdrift ( $v_{snd}$ ) and buoyancy ( $v_{by}$ ) are small (<1 %). The  
vertical bars in Figure 6a indicate the interannual variability, which is driven solely by snowfall except for the two summer  
months, which also have contributions from sublimation and melt.

405 The seasonal amplitudes of the ice-sheet-wide firn thickness and FAC are 3.1 and 2.1 cm, which amounts to Antarctic wide  
integrated volumes of 428 and 307 km<sup>3</sup>, respectively. This agrees with the modelled firn thickness amplitude of 3 cm found  
by Medley et al. (2022). The average of all local seasonal amplitudes are considerably higher, 6.8 and 4.2 cm, amounting to  
volume changes of 892 and 562 km<sup>3</sup>, respectively (Fig. 6c). The reason is that there is a spatial variation in timing of the  
seasonal maximum and minimum (Fig. 6d). The difference between the firn thickness and FAC seasonal cycles (6.8 vs 4.2 cm)  
410 suggests that around 62 % of the seasonal surface elevation fluctuations are caused by a change in air content rather than actual  
mass change. This result implies that not including the seasonal cycles of FAC in altimetry studies can generate significant  
errors, especially when investigating changes in subannual periods and in regions with large seasonal SMB cycles.

Figure 6c shows that the seasonal firn thickness amplitude can be up to 3 m in the western Antarctic Peninsula, caused by a  
rare combination of high snowfall and melt. In regions with little melt, such as coastal West Antarctica, the values are typically  
415 1 m. In interior East Antarctica, values of less than 2.5 cm are found, as the annual average accumulation is low and melt  
absent. Figure 6d shows the average day of the year of the maximum firn height, which generally occurs during late winter or




**Figure 6.** Ice sheet averaged seasonal cycle from FDM v1.2A of the (a) vertical firm surface velocity including its separate components from Eq. 6, and of (b) surface elevation and firm air content (FAC). The bars and colored areas indicate the interannual standard deviations. Maps from FDM v1.2A (c) of the average seasonal firm thickness peak-to-peak difference and (d) of average day of the year of maximum firm height.

spring, before the summer decrease in accumulation and increase in temperature. In some regions accumulation and firm depth peak in late summer or early winter, e.g., along the Weddell Sea coast, in Enderby Land, Adélie Land and parts of the interior.

#### 4.4 Interannual firm air content and firm depth variability

420 Figure 7 shows time series of AIS and above/below 2,000 m a.s.l. integrated FAC and the cumulative surface temperature and vertical firm surface velocity anomalies (against the long-term mean) including its components (Eq. 6). The figure confirms that the seasonal firm thickness and FAC variability are driven by snowfall, densification, melt and sublimation resulting in a firm thickness peak in late winter. The figure also reveals decadal firm thickness and FAC variability mainly resulting from fluctuations in snowfall (46 % of the total cumulative  $dH$  anomaly), and that firm densification, despite the long time scale, 425 reduces these snowfall-induced fluctuations by about 20 % (calculated from difference in standard deviation (sd) between the cumulative anomalies of  $v_{snow}$  and  $v_{snow+v_{fc}}$ ). We also find interannual variability in melt, but snowfall variability dominates. Below 2,000 m a.s.l. the variability in firm thickness and FAC are substantially larger, which can be attributed to the higher accumulation (70 % of the total accumulation falls on only 40 % of the ice sheet), in combination with larger temporal



**Figure 7.** Time series from FDM v1.2A of FAC, of the cumulative anomalies (against long-term mean) of surface temperature, of the vertical firn surface velocity, and of its separate components from Eq. (6). Time series are shown for (a) the entire ice sheet, (b) the part of the ice sheet situated above 2,000 m a.s.l. and (c) the part of the ice sheet situated below 2,000 m a.s.l.

variability in snowmelt and firn densification. Above 2,000 m a.s.l. the relative contribution of firn densification to the total cumulative dH anomaly is larger (32 %) than below 2,000 m a.s.l., where this is only 14 %. This difference can partly be explained by the larger interannual temperature variability above 2,000 m a.s.l. (sd in annual means = 0.78 K compared to 0.48 K) and the absence of melt. The interannual variability in snowfall and temperature are related to low and mid-latitude climate oscillations such as the El Niño-Southern Oscillation and the Southern Annular Mode (Kwok and Comiso, 2002; Marshall et al., 2017). Figure 7 shows that the magnitude of the decadal FAC variability (sd = 1.8 cm) constitutes a large fraction (67 %) of the total surface elevation change (sd = 2.7 cm). This reiterates the necessity of the common practice to remove decadal variability in FAC from altimetry signals to avoid errors in mass change estimates.

## 5 Comparison with altimetry and FDM v1.1 surface elevation change


In sections 3.1, 3.2 and 4 we presented and evaluated the spatial variation in firn thickness and density. To also evaluate the temporal variation, we compare our simulated elevation trends and seasonal variability to multi-mission satellite altimetry observations reported by Schröder et al. (2019) for the period 1992-2015.

### 5.1 Seasonal amplitude

The altimetry observations prior to 2003 exhibit a larger short-term variability as a result of the lower measurement precision of the satellites during that period (Schröder et al., 2019; Nilsson et al., 2022), and are excluded from the seasonal amplitude comparison here. To enable comparison with the FDM v1.1p1 simulation, we focus on the period 2003-2015. To reduce the impact of the long-term trends we remove a linear trend from all datasets. After doing so, we find that the seasonal amplitude averaged over the ice sheet of the altimetry observations amounts to 5.2 cm, of the FDM v1.2A simulations to 3.1 cm (60 % of the observed amplitude) and of the FDM v1.1p1 simulations to 2.5 cm (48 % of the observed amplitude). The higher seasonal amplitude of FDM v1.2A results from increased snowmelt along the coastal margins in RACMO2.3p2 compared to RACMO2.3p1 in combination with a lower fresh snow density (Fig. 2). Using updated corrections and a normalization of seasonal amplitudes from different missions, Nilsson et al. (2022) found that seasonal amplitudes from 1992-2016 are 47 % smaller than the ones from Schröder et al. (2019), which compares well with the FDM seasonal amplitudes.

### 5.2 Decadal variability


In Figure 8 we compare observed (Fig. 8a) and simulated 2003-2015 surface elevation trends. To obtain these numbers, a linear regression is fitted to monthly values for the period 2003-2015. The FDM v1.2A surface elevation trend over this period is positive in Dronning Maud Land, Enderby Land, and negative in most other regions (Fig. 8b). FDM v1.1p1 (Fig. 8c) has roughly similar patterns as FDM v1.2A, but the regional magnitudes differ. Figure 8d shows the residual surface elevation trend of FDM v1.2A, which is calculated by subtracting the FDM v1.2A elevation trend from the altimetry trend, and thus combines unresolved trends and the errors in the firn model and altimetry. In Figure 8e we show the difference in altimetry agreement between FDM v1.2A and FDM v1.1p1, which is calculated by subtracting the absolute residual surface elevation trend compared to altimetry (Fig. 8d) of FDM v1.1p1 from the similarly derived absolute residual of FDM v1.2A. The blue areas indicate an improvement in altimetry agreement of FDM v1.2A compared to FDM v1.1p1. The average absolute residual trend of FDM v1.2A compared to altimetry has been reduced by 17 % compared to FDM v1.1p1 (from 2.6 to 2.1  $\text{cm yr}^{-1}$ ). The improvement is most notable in Dronning Maud Land, Wilkes Land and Adélie Land. In Dronning Maud Land the FDM v1.2A trend has become more positive, and in Wilkes Land and Adélie Land the trend has become less negative (Fig. 8b,c, for region names see Fig. 1). In the Antarctic Peninsula and Ellsworth Land the altimetry agreement of FDM v1.2A has reduced compared to FDM v1.1p1. In these regions, a substantial residual positive trend remains ( $>10 \text{ cm yr}^{-1}$ ) (Fig. 8d), which could be due to a long-term increase in accumulation and/or ice-dynamical thickening. Smith et al. (2020) used ICESat data to show that the mass gain is mainly located along the steep slopes of the Antarctic Peninsula and decreases with distance from the



**Figure 8.** Maps of trends in surface elevation change over the period 2003-2015 from (a) multi-mission altimetry (Schröder et al., 2019), (b) FDM v1.2A and (c) FDM v1.1p1 obtained using linear regression. (d) Residual trend of altimetry minus FDM v1.2A. (e) Difference in altimetry agreement between FDM v1.2A and FDM v1.1p1. (absolute residual of FDM v1.1p1 compared to altimetry minus absolute residual of FDM v1.2A compared to altimetry). Regions within the ice-dynamical imbalance mask reported by Shepherd et al. (2019) are excluded in (d) and (e) and indicated by the grey shaded area.

ocean, which is indicative of an increase in snow accumulation. The increase in accumulation on centennial time scales in these regions (e.g., Thomas et al., 2015, 2017; Medley and Thomas, 2019) implies that the actual firn column is not in balance with the 1979-2020 climate, as assumed in the FDM spin-up procedure. As a result, the vertical downward ice flow (Eq. 6) is overestimated, which results in underestimated surface elevation change. This is further discussed in the next paragraph.

In Figure 9 we compare time series of altimetry observed elevation change over the period 1992-2015 to simulated elevation change for eleven locations across the AIS. These locations were selected as they cover the main distinct patterns from Figure 8 and have continuous observations. The time series have been smoothed using a 6-monthly moving average window, in order to reduce the impact of seasonal variations in radar penetration depth. The red shaded area is the aggregated uncertainty range from the sensitivity analysis, which is primarily determined by uncertainty in the climate forcing during the spin-up period (see next section for details of the sensitivity analysis). The agreement of FDM v1.2A compared to FDM v1.1p1 has improved ( $R^2 = 0.58$  compared to 0.36), which is mainly due to changes in the RACMO2.3p2 forcing, given the strong similarity between elevation change of FDM v1.2A and FDM v1.1p2. The altimetry observations prior to 2003 exhibit relatively stronger short-term variability (+9 % sd compared to after 2003), in line with the smaller measurement precision (Schröder et al., 2019). The agreement with FDM v1.2A increases after 2003 ( $R^2 = 0.93$  compared to 0.64, RMSE recued by 49 %), however the altimetry variability remains higher than the simulated variability (+13 % sd compared to FDM v1.2A), as these more recent observations still contain noise (<11 cm over flat terrain) and have variations in radar penetration depth (Schröder et al., 2019), while on the other hand, it can also partly be related to errors in the firn model. The uncertainty bands in Figure 9 show that a large part (73 %) of the difference between FDM v1.2A and altimetry falls within the uncertainty range. As in Figure 8d, we find a substantial positive residual altimetry trend ( $\sim 14$  and  $5 \text{ cm yr}^{-1}$ ) at location 5 (Antarctic Peninsula) and 6 (Ellsworth Land). The upper bound of the uncertainty for those two locations is determined by the sensitivity tests in which the climate



**Figure 9.** Time series of the altimetry observations and modelled elevation change for eleven locations over the period 1992-2015. The time series have been smoothed using a 6-monthly moving average window. The modelled elevation change includes FDM v1.2A, FDM v1.1p1 and FDM v1.1p2. The number of each panel corresponds to the numbered locations on the maps. The red shaded areas indicate the total uncertainty from the sensitivity analysis. The red dotted lines in panels (5) and (6) indicate the surface elevation change from the spin-up  $T_{s, b}$  experiment.

during the spin-up period was colder and drier (spin-up  $T_{s, b}$  experiment), indicated by the red dotted lines in Figure 9. With this presumption, the trend difference for location (5) greatly reduces and largely disappears for location (6).

## 6 Sensitivity analysis

We investigated the impacts of uncertainties in the model densification rate expression, fresh snow density, climate forcing, and climate forcing during the spin-up period, on the final FAC and surface elevation change results for (i) the 106 sample locations and (ii) expanded to the entire ice sheet (Table 5). For the sample locations, the sensitivities with updated MO fits are lower than without updated MO fits, especially the FAC uncertainties, which are 41 % lower. The results for the sample locations with updated MO fits are then expanded to the entire ice sheet. Some differences between the ice-sheet-wide and sample locations exist, which can be explained by the sample locations being biased towards high accumulation, which is needed to obtain optimal empirical relations for expanding the sample locations uncertainties. The average simulated FAC, for the sample locations and entire ice sheet (both with updated MO fits), is most sensitive to the uncertainties in the MO fits (5.1

500 and 5.2 %), temperature forcing (1.8 and 2.3 %) and accumulation forcing (4 and 3.7 %), and least sensitive to uncertainties in fresh snow density (0.7 and 1.2 %), and the spin-up forcing (ranging from 0.5 to 0.7 %). The simulated surface elevation change, for the sample locations and the entire ice sheet, is most sensitive to uncertainties in the accumulation during the spin-up (22.4 and 19.0 mm yr<sup>-1</sup>), followed by the accumulation forcing (8.0 and 5.6 mm yr<sup>-1</sup>), and temperature during the spin-up forcing (5.2 and 4.4 mm yr<sup>-1</sup>). As the uncertainties in temperature and accumulation during the spin-up period  
 505 have opposite effects, the tests in which accumulation and temperature were adjusted simultaneously yield an surface elevation trends of of 17.4 and 14.9 mm yr<sup>-1</sup>. The simulated surface elevation change is least sensitive to uncertainties in the MO fits (0.7/0.8 and 0.5 mm yr<sup>-1</sup>), fresh snow density (2.1 and 1.5 mm yr<sup>-1</sup>) and temperature forcing (1.5 and 0.7 mm yr<sup>-1</sup>).


**Table 5.** Overview of the relative change in annual average firm air content (FAC) and surface elevation change (dH/dt) of the ice-sheet-wide estimates and sample locations experiments. The sample location experiments also include the experiments without adjusted MO fits. An overview of the experiments including the prescribed uncertainties are listed in Table 2.

Experiment name	ice-sheet-wide estimates		Sample locations (without adjusted MO fits) <sup>a</sup>	
	FAC (%)	dH/dt (mm yr <sup>-1</sup> )	FAC (%)	dH/dt (mm yr <sup>-1</sup> )
MO fits	5.1	0.5	5.2	0.7
$\rho_s$	1.2	1.5	0.7 (5.8)	2.1 (2.4)
$T_s$	1.8	0.7	2.3 (4.9)	0.7 (1.5)
<b>b</b>	4.0	5.6	3.7 (6.8)	7.8 (8.0)
spin-up $T_s$	0.5	4.4	0.5 (1.5)	5.2 (5.2)
spin-up <b>b</b>	0.7	19.0	0.7 (1.1)	22.4 (22.4)
spin-up $T_s$ , <b>b</b>	0.7	14.9	0.6 (0.8)	17.4 (17.4)
total <sup>b</sup>	6.9	16.0	6.8 (11.5)	19.2 (19.4)

<sup>a</sup>The values between brackets indicate the uncertainties of the experiments without updated MO fits. <sup>b</sup>For the spin-up experiments only the combined temperature and accumulation experiments are included.

The spatial patterns of the total ice-sheet-wide FAC and surface elevation change uncertainties are shown in Figure 10, and for all experiments individually in Figures S4 and S5. The total FAC uncertainty, dominated by the MO fits, accumulation and  
 510 temperature uncertainties, is high (>10 %) along the high-accumulation coastal margins and in part of the interior, especially near the Transantarctic Mountains. The surface elevation change uncertainty is dominated by the spin-up and accumulation uncertainty, and mainly follows the accumulation pattern, with high values (>25 mm yr<sup>-1</sup>) along the high-accumulation coastal margins, especially in the Antarctic Peninsula, and in the Transantarctic Mountains.

We show time series of the uncertainty in surface elevation change and FAC at the sample locations in Figure S7a for all  
 515 experiments, and in Figure S7b for the non-spin-up experiments. First, we discuss uncertainties in surface elevation change for non-spin-up and spin-up experiments. Next, we look at uncertainties in FAC. In non-spin-up experiments, the uncertainty is only 4 % of the total surface elevation change. This uncertainty is smaller than the 40 % that can be derived from Table 3. This is because the imposed uncertainties have an amplifying effect. For example, lowering the fresh-snow density increases



**Figure 10.** Maps from FDM v1.2A of the estimated ice-sheet-wide total uncertainty in (a) firn air content (FAC) and (b) surface elevation change (dH/dt) from the sensitivity analysis.

surface elevation change rates of both signs. However, in Table 3 the uncertainty in surface elevation change is calculated over a short 5-year period for each location separately. Spatial and temporal variability in sign of elevation change tend to cancel out the uncertainties. In contrast, the uncertainty in the spin-up experiments gives a spatially and temporally uniform response. This paves the way to expand ice-sheet-wide estimates of uncertainty into time series, like in Figure S7c.

For FAC, its ice-sheet-wide change is only 19 % of the uncertainty in surface elevation change, similar to the 23 % for the suite of sample locations. It indicates that imposing an uncertainty in the experiments mostly has an impact on the high-density, low-FAC part of the firn column. The spatial pattern in FAC change uncertainty for the spin-up experiments is different from that of surface elevation uncertainty (Fig. S6), with a decreasing uncertainty for high-accumulation regions ( $>600$  mm w.e.  $\text{yr}^{-1}$ ). We attribute this to the fact that in high-accumulation regions, the imposed uncertainty has an effect on the entire firn layer much more quickly. In contrast, in the non-spin-up experiments, the magnitude of the FAC uncertainty is similar ( $\sim 85$  %) to the uncertainty of surface elevation change. The finding that imposing an uncertainty has its main impact on deeper, higher-density firn, means that the impact on a volume to mass conversion is limited. On the other hand, the large uncertainties in the surface elevation change compromise our ability to separate ice-dynamical from SMB and firn-related surface elevation changes.

In addition, in most places except the Antarctic Peninsula and Ellsworth Land, the long-term trends in accumulation and temperature are often weak, not significant, and variable in time and space. It suggests that the ice-sheet-wide uncertainty due to uncertainty in the spin-up climate is lower than suggested in Figure S7c. For the Antarctic Peninsula and Ellsworth Land, accumulation and temperature has increased strongly during the 20th century. To evaluate our results from the spin-up sensitivity tests in these two regions, we used surface elevation trends from altimetry observations (Schröder et al., 2019) (Fig. S8). We found that absolute residual trend altimetry minus the spin-up sensitivity test was reduced by 25 % compared to the reference FDM v1.2A (from 6.1 to 4.5  $\text{cm yr}^{-1}$ ). When we leave out pixels with a  $z_{830}$  age  $< 42$  years (6% of the pixels),



where we assume the underlying ice to have responded to the changing climatic conditions, the residual trend was reduced by  
540 38 % (from 5.6 to 3.5 cm yr<sup>-1</sup>).

## 7 Implications, remaining limitations, and outlook

In this study, the fresh snow density and firn compaction parameterizations of IMAU-FDM have been improved, and an updated atmospheric forcing has been used. The resulting difference in firn changes over time of FDM v1.2A compared to FDM v1.1p1 has an impact on conversion of volume-to-mass changes in altimetry studies. Over the period 2003-2019, (Smith et al., 2020)  
545 found that the total mass change of East Antarctica amounts to 195 Gt yr<sup>-1</sup>, and of West Antarctica to -245 Gt yr<sup>-1</sup>. When FDM v1.2A is used instead of FDM v1.1p1, the mass change over the period 2003-2015 for East Antarctica is 7.3 Gt yr<sup>-1</sup> lower, and for West Antarctica this is 1.2 Gt yr<sup>-1</sup> higher. These temporal firn differences are predominantly caused by the updated forcing as is shown by Figure 9. In addition, Figures 2 and 3 show that updates in the fresh snow density and firn densification parameterizations have substantial impact on the time averaged firn density profiles. These outputs are required  
550 to assess the meltwater buffering capacity of the firn layer, and to convert column thickness to ice thickness, to calculate solid ice discharge over the grounding line (Rignot et al., 2019)

The comparison with satellite altimetry measurements are used to assess the performance of FDM. However, especially in the interior, both the long-term trend and seasonal variability are small compared to the error range of the altimetry observations, which explains part of the remaining discrepancy that exists there, but also hampers the comparison (Verjans et al., 2021).  
555 Accurate in situ measurements of surface height evolution and firn densification are sparse. Future evaluation of firn models will therefore likely continue to depend on satellite altimetry comparison. The ICESat-2 ATL15 (Smith et al., 2021) is a precise Antarctic-wide 3-month height change laser product that is available from May 2018 onwards, thereby providing an opportunity to assess seasonal and decadal variations, especially in the coming years when more data will become available. Smith et al. (2022) already used ICESat-2 measurements to evaluate regional climate and firn densification models in Greenland.

The sensitivity analysis shows that updating the MO fits reduces the average FAC sensitivity to uncertainties in fresh snow density and climatic forcing by 41 % (Table 5), which implies that the MO factors, partly correct for biases in these boundary conditions. Thus, application of FDM v1.2A calibrated MO fits when using a different fresh snow density or climate forcing is not recommended. The fresh snow density parameterization is also dependent on the climate conditions derived from RACMO2.3p2, so any biases will impact the updates of the coefficients. In addition, the simulated fresh snow density accounts  
565 for the wind-driven compaction, and the updated coefficients are thus not recommended to use in combination with a firn model that includes wind-driven densification of the upper firn layers.

Another limitation, which is evident from both the remote sensing comparison and the sensitivity analysis, is the assumption of a steady-state firn layer over the period with reliable climate forcing (1979-present). In our sensitivity analysis we show that correcting for this by estimating previous climatic conditions from ice core data, reduces the difference between simulated and  
570 altimetry observed surface elevation trends by 38 % over the Antarctic Peninsula and Ellsworth Land. Future work could focus on further accounting for pre-1979 climatic conditions.

## 8 Summary and conclusions

In this study, we present and evaluate a simulation of the contemporary characteristics of the Antarctic firn layer using the updated IMAU-FDM (Firn Densification Model), version v1.2A, for the period 1979-2020. In this new version, the fresh  
575 snow density and firn compaction parameterizations have been improved, and an updated atmospheric forcing has been used. The RMSE when compared to observations of the fresh snow density and firn thickness have been reduced by 28 and 36 %, respectively, compared to the previous parameterizations (FDM FS-K and FDM v1.2A-log). The RMSE of FAC is 21 % lower compared to FDM v1.1p1. In general, the modelled fresh snow densities have decreased, especially in high-accumulation areas and to a lesser extent in windy escarpment regions. Firn thickness and FAC have decreased in low- and high-accumulation  
580 areas, whereas they have increased in intermediate accumulation areas.

In accordance with observations, the firn thickness and density patterns vary spatially across climatic regions with temperature as a primary and accumulation and surface melt as a secondary drivers. Along the coastal margins, we found that the firn thickness and FAC vary strongly in time, whereas in the interior the variations are small. The temporal variations can be split into a rather stable seasonal cycle driven by snowfall, compaction, summer melt and sublimation, and more irregular decadal  
585 variations mainly driven by slow snowfall variations. As variations in FAC and firn thickness have a similar phase, 63 to 68 % of the seasonal and decadal surface height variability is due to variations in FAC rather than firn mass, which emphasizes the importance of correcting for seasonal and decadal FAC variations in altimetry studies.

Comparison of simulated surface elevation change with altimetry confirms that the performance of the updated model has improved (a 17 % reduction in error), notably in Dronning Maud Land and Wilkes Land, which is predominantly due to updated  
590 climate forcing. However, a substantial trend difference ( $>10 \text{ cm yr}^{-1}$ ) remains in the Antarctic Peninsula and Ellsworth Land, which is likely caused by increasing accumulation over the past centuries, which violates the model assumption of a steady state over the simulated period (1979-2020). Correcting for this over the Antarctic Peninsula and Ellsworth Land, by estimating previous climatic conditions from ice core data, substantially improves this (an error reduction of 38 %). In general, uncertainties in the model formulation, forcing or forcing during the spin-up cause rather small changes in simulated FAC  
595 ( $<5.2 \%$ ), while uncertainties in the accumulation forcing, and in the accumulation and temperature forcing during the spin-up period can lead to substantial differences in simulated surface elevation change (4 to  $19 \text{ mm yr}^{-1}$ ).

*Code availability.* The code of IMAU-FDM v1.2A is available on GitHub at <https://github.com/brils001/IMAU-FDM> (last access: 23 November 2022).

*Data availability.* A list of all firn cores used and corresponding references can be found in the Table S1 in the Supplement. IMAU-FDM  
600 data can be obtained from the authors without conditions.

*Author contributions.* SV, WJvdB, MB, PKM and MRvdB defined the research goals and designed the study. SV improved the model, performed the simulations and analyzed the results. All authors contributed to discussions on the manuscript.

*Competing interests.* MRvdB is a member of the editorial board of journal The Cryosphere

*Acknowledgements.* This work was funded by the Netherlands Organization for Scientific Research (grant no. OCENW.GROOT.2019.091).  
605 MRvdB acknowledges support from the Netherlands Earth System Science Centre (NESCC). We acknowledge ECMWF for computational time on their supercomputers.

## References

- Adusumilli, S., Fricker, H. A., Siegfried, M. R., Padman, L., Paolo, F. S., and Ligtenberg, S. R.: Variable basal melt rates of Antarctic Peninsula ice shelves, 1994–2016, *Geophysical Research Letters*, 45, 4086–4095, 2018.
- 610 Anderson, E. A.: A point energy and mass balance model of a snow cover., Stanford University, 1976.
- Arthern, R. and Wingham, D.: The natural fluctuations of firn densification and their effect on the geodetic determination of ice sheet mass balance, *Climatic Change*, 40, 605–624, 1998.
- Arthern, R. J., Vaughan, D. G., Rankin, A. M., Mulvaney, R., and Thomas, E. R.: In situ measurements of Antarctic snow compaction compared with predictions of models, *Journal of Geophysical Research: Earth Surface*, 115, <https://doi.org/10.1029/2009JF001306>, 2010.
- 615 Bréant, C., Martinerie, P., Orsi, A., Arnaud, L., and Landais, A.: Modelling firn thickness evolution during the last deglaciation: constraints on sensitivity to temperature and impurities, *Climate of the Past*, 13, 833–853, <https://doi.org/10.5194/cp-13-833-2017>, 2017.
- Brils, M., Kuipers Munneke, P., van de Berg, W. J., and van den Broeke, M.: Improved representation of the contemporary Greenland ice sheet firn layer by IMAU-FDM v1. 2G, *Geoscientific Model Development*, 15, 7121–7138, 2022.
- Brun, E., Martin, E., and Spiridonov, V.: Coupling a multi-layered snow model with a GCM, *Annals of Glaciology*, 25, 66–72, <https://doi.org/10.3189/S0260305500013811>, 1997.
- 620 Calonne, N., Milliancourt, L., Burr, A., Philip, A., Martin, C. L., Flin, F., and Geindreau, C.: Thermal Conductivity of Snow, Firn, and Porous Ice From 3-D Image-Based Computations, *Geophysical Research Letters*, 46, 13 079–13 089, <https://doi.org/10.1029/2019GL085228>, 2019.
- Carter, J., Leeson, A., Orr, A., Kittel, C., and van Wessem, J. M.: Variability in Antarctic surface climatology across regional climate models and reanalysis datasets, *The Cryosphere*, 16, 3815–3841, 2022.
- 625 Coléou, C., Xu, K., Lesaffre, B., and Brzoska, J.-B.: Capillary rise in snow, *Hydrological processes*, 13, 1721–1732, 1999.
- Datta, R. T., Tedesco, M., Fettweis, X., Agosta, C., Lhermitte, S., Lenaerts, J. T., and Wever, N.: The effect of Foehn-induced surface melt on firn evolution over the northeast Antarctic peninsula, *Geophysical Research Letters*, 46, 3822–3831, <https://doi.org/10.1029/2018GL080845>, 2019.
- 630 Davis, C. H. and Poznyak, V. I.: The depth of penetration in Antarctic firn at 10 GHz, *IEEE Transactions on Geoscience and Remote Sensing*, 31, 1107–1111, <https://doi.org/10.1109/36.263784>, 1993.
- Fausto, R. S., Box, J. E., Vandecrux, B., Van As, D., Steffen, K., MacFerrin, M. J., Machguth, H., Colgan, W., Koenig, L. S., McGrath, D., et al.: A snow density dataset for improving surface boundary conditions in Greenland ice sheet firn modeling, *Frontiers in Earth Science*, 6, 51, <https://doi.org/10.3389/feart.2018.00051>, 2018.
- 635 Favier, V., Krinner, G., Amory, C., Gallée, H., Beaumet, J., and Agosta, C.: Antarctica-regional climate and surface mass budget, *Current Climate Change Reports*, 3, 303–315, <https://doi.org/10.1007/s40641-017-0072-z>, 2017.
- Fernandoy, F., Meyer, H., Oerter, H., Wilhelms, F., Graf, W., and Schwander, J.: Temporal and spatial variation of stable-isotope ratios and accumulation rates in the hinterland of Neumayer station, East Antarctica, *Journal of glaciology*, 56, 673–687, <https://doi.org/10.3189/002214310793146296>, 2010.
- 640 Fourteau, K., Martinerie, P., Faïn, X., Schaller, C. F., Tuckwell, R. J., Löwe, H., Arnaud, L., Magand, O., Thomas, E. R., Freitag, J., et al.: Multi-tracer study of gas trapping in an East Antarctic ice core, *The Cryosphere*, 13, 3383–3403, <https://doi.org/10.5194/tc-13-3383-2019>, 2019.

- Gilbert, E. and Kittel, C.: Surface melt and runoff on Antarctic ice shelves at 1.5 C, 2 C, and 4 C of future warming, *Geophysical Research Letters*, 48, e2020GL091733, <https://doi.org/10.1029/2020GL091733>, 2021.
- 645 Gossart, A., Helsen, S., Lenaerts, J., Broucke, S. V., Van Lipzig, N., and Souverijns, N.: An evaluation of surface climatology in state-of-the-art reanalyses over the Antarctic Ice Sheet, *Journal of Climate*, 32, 6899–6915, 2019.
- Groot Zwaaftink, C., Cagnati, A., Crepaz, A., Fierz, C., Macelloni, G., Valt, M., and Lehning, M.: Event-driven deposition of snow on the Antarctic Plateau: analyzing field measurements with SNOWPACK, *The Cryosphere*, 7, 333–347, 2013.
- Helsen, M. M., Van Den Broeke, M. R., Van De Wal, R. S., Van De Berg, W. J., Van Meijgaard, E., Davis, C. H., Li, Y., and Goodwin, I.:  
650 Elevation changes in Antarctica mainly determined by accumulation variability, *science*, 320, 1626–1629, 2008.
- Herron, M. M. and Langway, C. C.: Firn densification: an empirical model, *Journal of Glaciology*, 25, 373–385, 1980.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., et al.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Judson, A. and Doesken, N.: Density of freshly fallen snow in the central Rocky Mountains, *Bulletin of the American Meteorological Society*,  
655 81, 1577–1588, 2000.
- Kaspers, K., Van de Wal, R., Van den Broeke, M., Schwander, J., Van Lipzig, N., and Brenninkmeijer, C.: Model calculations of the age of firn air across the Antarctic continent, *Atmospheric Chemistry and Physics*, 4, 1365–1380, <https://doi.org/10.5194/acp-4-1365-2004>, 2004.
- Keenan, E., Wever, N., Dattler, M., Lenaerts, J., Medley, B., Kuipers Munneke, P., and Reijmer, C.: Physics-based SNOWPACK model  
660 improves representation of near-surface Antarctic snow and firn density, *The Cryosphere*, 15, 1065–1085, <https://doi.org/10.5194/tc-15-1065-2021>, 2021.
- Kuipers Munneke, P., Ligtenberg, S., Noël, B., Howat, I., Box, J., Mosley-Thompson, E., McConnell, J., Steffen, K., Harper, J., Das, S., et al.: Elevation change of the Greenland Ice Sheet due to surface mass balance and firn processes, 1960–2014, *The Cryosphere*, 9, 2009–2025, <https://doi.org/10.5194/tc-9-2009-2015>, 2015.
- 665 Kwok, R. and Comiso, J. C.: Spatial patterns of variability in Antarctic surface temperature: Connections to the Southern Hemisphere Annular Mode and the Southern Oscillation, *Geophysical Research Letters*, 29, 50–1, <https://doi.org/10.1029/2002GL015415>, 2002.
- Lehning, M., Bartelt, P., Brown, B., and Fierz, C.: A physical SNOWPACK model for the Swiss avalanche warning: Part III: Meteorological forcing, thin layer formation and evaluation, *Cold Regions Science and Technology*, 35, 169–184, [https://doi.org/10.1016/S0165-232X\(02\)00072-1](https://doi.org/10.1016/S0165-232X(02)00072-1), 2002.
- 670 Lenaerts, J. T., Van den Broeke, M., Déry, S., Van Meijgaard, E., Van de Berg, W., Palm, S. P., and Sanz Rodrigo, J.: Modeling drifting snow in Antarctica with a regional climate model: 1. Methods and model evaluation, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2011JD016145>, 2012.
- Ligtenberg, S., Helsen, M., and Van den Broeke, M.: An improved semi-empirical model for the densification of Antarctic firn, *The Cryosphere*, 5, 809–819, 2011.
- 675 Ligtenberg, S., Horwath, M., Van den Broeke, M., and Legrésy, B.: Quantifying the seasonal “breathing” of the Antarctic ice sheet, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL053628>, 2012.
- Ligtenberg, S., Kuipers Munneke, P., and Van den Broeke, M.: Present and future variations in Antarctic firn air content, *The Cryosphere*, 8, 1711–1723, <https://doi.org/10.5194/tc-8-1711-2014>, 2014.
- Lundin, J. M., Stevens, C. M., Arthern, R., Buizert, C., Orsi, A., Ligtenberg, S. R., Simonsen, S. B., Cummings, E., Essery, R., Leahy, W.,  
680 et al.: Firn model intercomparison experiment (FirnMICE), *Journal of Glaciology*, 63, 401–422, 2017.

- Marshall, G. J., Thompson, D. W., and van den Broeke, M. R.: The signature of Southern Hemisphere atmospheric circulation patterns in Antarctic precipitation, *Geophysical Research Letters*, 44, 11–580, <https://doi.org/10.1002/2017GL075998>, 2017.
- Medley, B. and Thomas, E.: Increased snowfall over the Antarctic Ice Sheet mitigated twentieth-century sea-level rise, *Nature Climate Change*, 9, 34–39, <https://doi.org/10.1038/s41558-018-0356-x>, 2019.
- 685 Medley, B., Ligtenberg, S., Joughin, I., Van den Broeke, M. R., Gogineni, S., and Nowicki, S.: Antarctic firn compaction rates from repeat-track airborne radar data: I. Methods, *Annals of Glaciology*, 56, 155–166, 2015.
- Medley, B., Neumann, T. A., Zwally, H. J., Smith, B. E., and Stevens, C. M.: Simulations of firn processes over the Greenland and Antarctic ice sheets: 1980–2021, *The Cryosphere*, 16, 3971–4011, 2022.
- Montgomery, L., Koenig, L., and Alexander, P.: The SUMup dataset: compiled measurements of surface mass balance components over ice  
690 sheets and sea ice with analysis over Greenland, *Earth System Science Data*, 10, 1959–1985, <https://doi.org/10.5194/essd-10-1959-2018>, 2018.
- Morris, E. M. and Wingham, D. J.: Uncertainty in mass-balance trends derived from altimetry: a case study along the EGIG line, central Greenland, *Journal of Glaciology*, 61, 345–356, 2015.
- Mottram, R., Hansen, N., Kittel, C., van Wessem, J. M., Agosta, C., Amory, C., Boberg, F., van de Berg, W. J., Fettweis, X., Gossart, A.,  
695 et al.: What is the surface mass balance of Antarctica? An intercomparison of regional climate model estimates, *The Cryosphere*, 15, 3751–3784, <https://doi.org/10.5194/tc-15-3751-2021>, 2021.
- Munneke, P. K., Ligtenberg, S. R., Van Den Broeke, M. R., and Vaughan, D. G.: Firn air depletion as a precursor of Antarctic ice-shelf collapse, *Journal of Glaciology*, 60, 205–214, <https://doi.org/10.3189/2014JoG13J183>, 2014.
- Nilsson, J., Gardner, A. S., and Paolo, F. S.: Elevation change of the Antarctic Ice Sheet: 1985 to 2020, *Earth System Science Data*, 14,  
700 3573–3598, 2022.
- Noël, B., van de Berg, W. J., Van Wessem, J. M., Van Meijgaard, E., Van As, D., Lenaerts, J., Lhermitte, S., Kuipers Munneke, P., Smeets, C., Van Uft, L. H., et al.: Modelling the climate and surface mass balance of polar ice sheets using RACMO2–Part 1: Greenland (1958–2016), *The Cryosphere*, 12, 811–831, <https://doi.org/10.5194/tc-12-811-2018>, 2018.
- Olmi, R., Bittelli, M., Picard, G., Arnaud, L., Mialon, A., and Priori, S.: Investigating the influence of the grain size and distribution on  
705 the macroscopic dielectric properties of Antarctic firn, *Cold Regions Science and Technology*, 185, 103 254, <https://doi.org/10.1016/j.coldregions.2021.103254>, 2021.
- Oppenheimer, M., Glavovic, B., Hinkel, J., van de Wal, R., Magnan, A. K., Abd-Elgawad, A., Cai, R., Cifuentes-Jara, M., Deconto, R. M., Ghosh, T., et al.: Sea level rise and implications for low lying islands, coasts and communities, 2019.
- Rignot, E., Mouginot, J., Scheuchl, B., Van Den Broeke, M., Van Wessem, M. J., and Morlighem, M.: Four decades of Antarctic Ice  
710 Sheet mass balance from 1979–2017, *Proceedings of the National Academy of Sciences*, 116, 1095–1103, <https://doi.org/10.1073/pnas.1812883116>, 2019.
- Sato, T., Kosugi, K., Mochizuki, S., and Nemoto, M.: Wind speed dependences of fracture and accumulation of snowflakes on snow surface, *Cold Regions Science and Technology*, 51, 229–239, <https://doi.org/10.1016/j.coldregions.2007.05.004>, 2008.
- Schröder, L., Horwath, M., Dietrich, R., Helm, V., Van Den Broeke, M. R., and Ligtenberg, S. R.: Four decades of Antarctic surface elevation  
715 changes from multi-mission satellite altimetry, *The Cryosphere*, 13, 427–449, <https://doi.org/10.5194/tc-13-427-2019>, 2019.
- Schwanck, F., Simões, J. C., Handley, M., Mayewski, P. A., Bernardo, R. T., and Aquino, F. E.: Drilling, processing and first results for Mount Johns ice core in West Antarctica Ice Sheet, *Brazilian Journal of Geology*, 46, 29–40, <https://doi.org/10.1590/2317-4889201620150035>, 2016.

- 720 Shepherd, A., Ivins, E., Rignot, E., Smith, B., Van Den Broeke, M., Velicogna, I., Whitehouse, P., Briggs, K., Joughin, I., Krinner, G., et al.: Mass balance of the Antarctic Ice Sheet from 1992 to 2017, *Nature*, 558, 219–222, <https://doi.org/10.1038/s41586-018-0179-y>, 2018.
- Shepherd, A., Gilbert, L., Muir, A. S., Konrad, H., McMillan, M., Slater, T., Briggs, K. H., Sundal, A. V., Hogg, A. E., and Engdahl, M. E.: Trends in Antarctic Ice Sheet elevation and mass, *Geophysical Research Letters*, 46, 8174–8183, <https://doi.org/10.1029/2019GL082182>, 2019.
- 725 Smith, B., Fricker, H. A., Gardner, A. S., Medley, B., Nilsson, J., Paolo, F. S., Holschuh, N., Adusumilli, S., Brunt, K., Csatho, B., et al.: Pervasive ice sheet mass loss reflects competing ocean and atmosphere processes, *Science*, 368, 1239–1242, <https://doi.org/10.5194/tc-5-809-2011>, 2020.
- Smith, B. E., Medley, B., Fettweis, X., Sutterley, T., Alexander, P., Porter, D., and Tedesco, M.: Evaluating Greenland Surface-Mass-Balance and Firn-Densification Data Using ICESat-2 Altimetry, *The Cryosphere Discussions*, pp. 1–24, <https://doi.org/10.5194/tc-2022-44>, 2022.
- 730 Smith, B., B. P. J. S. D. T. S. T. A. N., Harbeck, K., et al.: ATLAS/ICESat-2 L3B Gridded Antarctic and Arctic Land Ice Height Change, Version 1, 2021.
- Sommer, C. G., Wever, N., Fierz, C., and Lehning, M.: Investigation of a wind-packing event in Queen Maud Land, Antarctica, *The Cryosphere*, 12, 2923–2939, <https://doi.org/10.5194/tc-12-2923-2018>, 2018.
- 735 Steger, C. R., Reijmer, C. H., Van Den Broeke, M. R., Wever, N., Forster, R. R., Koenig, L. S., Kuipers Munneke, P., Lehning, M., Lhermitte, S., Ligtenberg, S. R., et al.: Firn meltwater retention on the Greenland ice sheet: A model comparison, *Frontiers in Earth Science*, 5, 3, <https://doi.org/10.3389/feart.2017.00003>, 2017.
- Stenni, B., Curran, M. A., Abram, N. J., Orsi, A., Goursaud, S., Masson-Delmotte, V., Neukom, R., Goosse, H., Divine, D., Van Ommen, T., et al.: Antarctic climate variability on regional and continental scales over the last 2000 years, *Climate of the Past*, 13, 1609–1634, <https://doi.org/10.5194/cp-13-1609-2017>, 2017.
- 740 Sugiyama, S., Enomoto, H., Fujita, S., Fukui, K., Nakazawa, F., Holmlund, P., and Surdyk, S.: Snow density along the route traversed by the Japanese-Swedish Antarctic Expedition 2007/08, *Journal of Glaciology*, 58, 529–539, <https://doi.org/10.3189/2012JoG11J201>, 2012.
- Thomas, E. R., Hosking, J. S., Tuckwell, R. R., Warren, R., and Ludlow, E.: Twentieth century increase in snowfall in coastal West Antarctica, *Geophysical Research Letters*, 42, 9387–9393, <https://doi.org/10.1002/2015GL065750>, 2015.
- 745 Thomas, E. R., Van Wessem, J. M., Roberts, J., Isaksson, E., Schlosser, E., Fudge, T. J., Vallelonga, P., Medley, B., Lenaerts, J., Bertler, N., et al.: Regional Antarctic snow accumulation over the past 1000 years, *Climate of the Past*, 13, 1491–1513, <https://doi.org/10.5194/cp-13-1491-2017>, 2017.
- van den Broeke, M.: Depth and density of the Antarctic firn layer, *Arctic, Antarctic, and Alpine Research*, 40, 432–438, [https://doi.org/10.1657/1523-0430\(07-021\)\[BROEKE\]2.0.CO;2](https://doi.org/10.1657/1523-0430(07-021)[BROEKE]2.0.CO;2), 2008.
- 750 Van Wessem, J. M., Van De Berg, W. J., Noël, B. P., Van Meijgaard, E., Amory, C., Birnbaum, G., Jakobs, C. L., Krüger, K., Lenaerts, J., Lhermitte, S., et al.: Modelling the climate and surface mass balance of polar ice sheets using RACMO2–Part 2: Antarctica (1979–2016), *The Cryosphere*, 12, 1479–1498, <https://doi.org/10.5194/tc-12-1479-2018>, 2018.
- Vandercrux, B., Mottram, R., Langen, P. L., Fausto, R. S., Olesen, M., Stevens, C. M., Verjans, V., Leeson, A., Ligtenberg, S., Kuipers Munneke, P., et al.: The firn meltwater Retention Model Intercomparison Project (RetMIP): evaluation of nine firn models at four weather station sites on the Greenland ice sheet, *The Cryosphere*, 14, 3785–3810, <https://doi.org/10.5194/tc-14-3785-2020>, 2020.
- 755 Verjans, V., Leeson, A., McMillan, M., Stevens, C., van Wessem, J. M., van de Berg, W. J., van den Broeke, M., Kittel, C., Amory, C., Fettweis, X., et al.: Uncertainty in East Antarctic firn thickness constrained using a model ensemble approach, *Geophysical Research Letters*, 48, e2020GL092060, <https://doi.org/10.1029/2020GL092060>, 2021.

- Willen, M., Broerse, T., Groh, A., Wouters, B., Kuipers Munneke, P., Horwath, M., van den Broeke, M., and Schröder, L.: Separating Long-Term and Short-Term Mass Changes of Antarctic Ice Drainage Basins: A Coupled State Space Analysis of Satellite Observations and Model Products, *Journal of Geophysical Research: Earth Surface*, 126, e2020JF005966, <https://doi.org/10.1029/2020JF005966>, 2021.
- 760 Winstrup, M., Vallelonga, P., Kjær, H. A., Fudge, T. J., Lee, J. E., Riis, M. H., Edwards, R., Bertler, N. A., Blunier, T., Brook, E. J., et al.: A 2700-year annual timescale and accumulation history for an ice core from Roosevelt Island, West Antarctica, *Climate of the Past*, 15, 751–779, <https://doi.org/10.5194/cp-15-751-2019>, 2019.
- Winther, J.-G., Jespersen, M. N., and Liston, G. E.: Blue-ice areas in Antarctica derived from NOAA AVHRR satellite data, *Journal of Glaciology*, 47, 325–334, <https://doi.org/10.3189/172756501781832386>, 2001.