

Simulating the current and future northern limit of permafrost on the Qinghai-Tibet Plateau

Jianting Zhao¹, Lin Zhao^{1,2,3}, Zhe Sun^{1,4}, Fujun Niu⁵, Guojie Hu³, Defu Zou³,
5 Guangyue Liu³, Erji Du³, Chong Wang¹, Lingxiao Wang¹, Yongping Qiao³,
Jianzong Shi³, Yuxin Zhang¹, Junqiang Gao⁶, Yuanwei Wang¹, Yan Li¹, Wenjun
Yu⁷, Huayun Zhou^{2,3}, Zanpin Xing^{2,3}, Minxuan Xiao¹, Luhui Yin¹, Shengfeng
10 Wang¹

¹ School of Geographical Sciences, Nanjing University of Information Science & Technology, Nanjing 210044,
10 China

² University of Chinese Academy of Sciences, Beijing 101408, China

³ Cryosphere Research Station on Qinghai-Xizang Plateau, State Key Laboratory of Cryospheric Sciences,
Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences, Lanzhou 730000, China

⁴ Key Laboratory of Environment Change and Resources Use in Beibu Gulf, Ministry of Education, Nanning
15 Normal University, Nanning 530001, China

⁵ State Key Laboratory of Frozen Soil Engineering, Northwest Institute of Eco-Environment and Resources,
Chinese Academy of Sciences, Lanzhou 730000, China

⁶ School of Mathematics and Statistics, Nanjing University of Information Science & Technology, Nanjing 210044,
China

⁷ School of Hydrology and Water Resources, Nanjing University of Information Science & Technology, Nanjing
20 210044, China

Correspondence to : Lin Zhao (lzhao@nuist.edu.cn/lzhao@lzb.ac.cn)

Abstract: Permafrost has been warming and thawing globally, with subsequent effects on the climate, hydrological, and ecosystem. However, the permafrost thermal state variation in the northern lower limit of the permafrost zone (Xidatan) on the Qinghai-Tibet Plateau (QTP) is unclear. This study attempts to explore the changes and variability of this permafrost using historical (1970–2019) and future projection datasets from remote sensing-based Land Surface Temperature product (LST) and climate projections from Earth System Model (ESM) outputs of Coupled Model Intercomparison Project Phase 5 and 6 (CMIP5, CMIP6). Our model considers phase change processes of soil pore water, thermal properties difference between frozen and unfrozen soil, geothermal flux flow, and ground ice effect. Our model can consistently reproduce the vertical ground temperature profiles and active layer thickness (ALT), recognizing permafrost boundaries, and capture the evolution of the permafrost thermal regime. The spatial distribution of permafrost

and its thermal conditions over the study area were controlled by elevation with a strong influence
35 of slope orientation. From 1970 to 2019, the mean annual ground temperature (MAGT) in the region
warmed by 0.49 °C in the continuous permafrost zone and 0.40 °C in the discontinuous permafrost
zone. The lowest elevation of the permafrost boundary (on the north-facing slopes) rose
approximately 47 m, and the northern boundary of discontinuous permafrost has retreated
southwards approximately 1~2 km, while the lowest elevation of the permafrost boundary remains
40 unchanged for the continuous permafrost zone. The warming rate in MAGT is projected to be more
pronounced under Shared Socioeconomic Pathways (SSPs) than that of Representative
Concentration Pathways (RCPs), but no distinct discrepancies in the areal extent of the continuous,
discontinuous permafrost and seasonally frozen ground among SSP and RCP scenarios. This study
highlights the slow delaying process in the response of permafrost in QTP to a warming climate,
45 especially in terms of the areal extent of permafrost distribution.

1 Introduction

Permafrost is one of the crucial components of the cryosphere that is largely sensitive to climate
change (Li et al., 2008; Nitze et al., 2018; Smith et al., 2022). Owing to its high elevation (mean
elevation above 4000 m above sea level (a.s.l.)) and extreme cold climate, the QTP is considered
50 the largest and highest elevational permafrost region (occupies a permafrost area of $1.06 \times 10^6 \text{ km}^2$
or 40 % of the total area of the QTP) located in the mid- to low- latitude regions (Zhou et al., 2000;
Yang et al., 2010; Zou et al., 2017; Zhao et al., 2020). Since the 20th century, climate warming has
been evident on the QTP, particularly in the permafrost regions, which has significantly impacted
the permafrost, manifested by rising ground temperatures, increase in ALT, thinning of permafrost,
55 melting of ground ice, and disappearing of permafrost ultimately (Wang et al., 2000; Cheng and Wu,
2007; Wu et al., 2008; Jin et al., 2011; Li et al., 2012; Zhao et al., 2020; Zhang et al., 2022). Changes
in the permafrost have substantial impacts on the hydrological process (Cheng et al., 2013; Zhao et
al., 2019), the energy exchange between land and atmosphere (Xiao et al., 2013; Hu et al., 2017),
natural hazards (Hjort et al., 2022), carbon budget (Schädel et al., 2016; Miner et al., 2022; Hjort et
60 al., 2022; Fewster et al., 2022), and ecological environment (Yi et al., 2014; Jin et al., 2021).
Therefore, it has become a pressing issue for research to diagnose how and at what rate permafrost

responds to global warming. It has prompted a great concern among geocryologists, cold regions engineers, and international society (Schuur et al., 2011; IPCC, 2019).

The northern fringe of the continuous permafrost zone of QTP is exceptionally vulnerable to climate variability, as characterized by permafrost and seasonally frozen ground coexistence, and a thicker active layer, much thinner and warmer permafrost in this region compared with the interior of the QTP (Wu et al., 2005; Liu et al., 2020). Considering the location of the northern lower limit of the continuous permafrost zone of the QTP, detailed permafrost environmental investigation and monitoring have been started systematically since 1987 (Zhao et al., 2021). The latest information from high resolution remote sensing products (e.g., Zou et al., 2014, 2017; Li et al., 2015b) is readily available. The Xidatan constitutes an ideal region to assess the response of marginal permafrost to a warming climate. Multiple field investigations and borehole monitoring were started in the late 1960s to aid in infrastructure construction of the Qinghai-Tibet Highway (QTH), documenting that the warming and thawing of permafrost have been striking in the region (Jin et al., 2000, 2006; Cheng et al., 2007). Less is known about the spatial variations, as the logistics of borehole installation is highly expensive and challenging in remote areas (e.g., remote alpine mountain areas with steep and complex topography). The higher degree of spatial heterogeneity (e.g., permafrost and seasonally frozen ground coexist) strongly influence permafrost distribution (Cheng et al., 2004), and a simple point observation representing regional conditions is problematic. Therefore, it is difficult to accurately delineate the permafrost distribution margin by traditional cartographic techniques from the limited field survey data, aerial photographs, satellite images, and topographic features dataset (Ran et al., 2012; Zou et al., 2017). It highlights the demand for a spatial study approach to achieve a realistic picture of permafrost distribution for further study of thermal state and dynamics in response to climate variability.

Models have the potential to overcome the shortage of in-situ data and field surveying in mapping permafrost conditions and change studies (Riseborough et al., 2008). A variety of models can be applied for the quantitative assessment of the response of marginal permafrost to the warming climate (Cheng et al., 1984; Li et al., 2008; Lawrence et al., 2012; Guo et al., 2016, 2012; Lu et al., 2017; Chang et al., 2018; Wang et al., 2019; Ni et al., 2021). However, most models are poor at

90 interpreting marginal permafrost, especially true in the region of northern or southern permafrost boundaries, such as the Xidatan. Such challenges, in part, are attributed to the effect of local factors (e.g., topography, vegetation, snow cover, thermal properties of the surface soil, etc.). Near the lower limit of permafrost, the permafrost and seasonally frozen ground coexist. High spatial heterogeneities of the land surface make it a challenging area for permafrost modeling (Cheng et al.,
95 2004; Zou et al., 2017, Luo et al., 2018; Yin et al., 2021). Due to the lack of detailed field observations, most existing simulation results have not considered the effects of water phase change and ground ice and the thermal state of deep permafrost. Hence, there is a considerable discrepancy among their results on the timing, rate, and magnitude of permafrost degradation (Zhao et al., 2020; Smith et al., 2022). Thus, it hardly makes us agree on a quantitative assessment of the response of
100 marginal permafrost to a warming climate. To address these issues, Sun et al. (2019) proposed a transient numerical heat conduction permafrost model and successfully simulating the evolution and dynamics of the permafrost thermal regime from 1962 until the end of this century at a monitoring borehole (QT09) located in the Xidatan comprehensive observation site (COS).

105 In this work, we attempt to upscale our model for the whole region, aiming to accurately simulate mountain permafrost spatial distribution and dynamics. The objective includes the production of high-resolution (1 km×1 km) data for the period of 1970–2019 and anticipate possible changes by 2100 under different climate change scenarios, forced by improved remote sensing-based spatial product (LST) and CMIP5 (under RCP 2.6, RCP4.5 and RCP8.5) and CMIP6 (under SSP1–2.6, SSP2–4.5, and SSP5–8.5) projections. Our model fully considers the thermal properties
110 difference between frozen and thawed soil, the phase variations of the unfrozen water in frozen soil, the distribution of the ground ice, and geothermal heat flow. We aim that this study will simulate the distribution of marginal permafrost on the QTP, quantitatively assess the thermal regime spatiotemporal dynamics under climate change, and anticipate changes for future climate scenarios.

2 Study area, materials, and methods

115 2.1 Study area

The study focuses on the Xidatan of QTP, situated in a narrow down-faulted basin at the

northern foot of the Eastern Kunlun Mountains within the northern limit of the permafrost on the QTP (Fig. 1a). The region encompasses a land area of ~ 220 km 2 and is characterized by discontinuous permafrost (Wu et al., 2005; Liu et al., 2020; Yin et al., 2021). Some periglacial landforms, such as block fields, stripes, and stone rings, have developed in the mountainous terrain (Luo et al., 2018). Several glaciers extend from the peaks of the East Kunlun Mountains downwards along the valley in the southern area (Fig. 1b). The elevation varies from 4100 m a.s.l. in the east to 5700 m a.s.l. in the west. Topographic relief in the majority (~90 %) is minimal (slopes lower than 5°), with some exceptions in mountainous areas. The plant community composition is mainly dominated by sparse alpine steppe, and the alpine desert consists of >10 m thick soil layer of gravel, fluvial sand, and silt (Wang et al., 2000; Jin et al., 2000; Yue et al., 2013; Yin et al., 2021) (Fig. 1b–g). According to the COS (Fig. 1b), from 2004 to 2018, the mean annual air temperature and mean annual precipitation were -3.6 °C and 384.5 mm, respectively. In 2017, permafrost thickness was approximately 26 m, with the MAGT at zero annual amplitude (ZAA, where the annual difference in ground temperature is less than 0.1 °C) was approximately -0.66 °C and ALT is about 1.60 m (Zhao et al., 2021).




Figure 1. The geographical location of the Xidatan on the QTP, its topography and the location of 24 borehole sites (a). Surface conditions at monitoring borehole sites (b–g): view over the Xidatan COS (b), QT09, view towards the south (c), QT09, view towards the northeast (d),

view from the vicinity of QT09 towards the east (e), XD2-1-2-7, view towards the south (f), XD1-1-1-6, view towards the east (g) (the spatial distribution of frozen ground types are derived from Zou et al. (2017); topography was generated by the Digital Elevation Model constructed (DEM) from the Shuttle Radar Topography Missions (SRTM) with a 1-arcsecond (~30 m) (Jarvis et al., 2008), Tibet Plateau boundary was taken from National Tibet Plateau Data Center (Zhang et al., 2019). All photographs were taken during the field investigation from 23 July 2021 to 2 August 2021).

2.2 Materials

2.2.1 Field monitoring and borehole observation datasets

There are fifteen monitoring boreholes with long-term observations (for the last 10 years) established in the Xidatan (Fig. 1a). A COS is in the central part of the Xidatan, where the ground surface is composed of sparse dry alpine meadows, and the soil layer is made of fluvial sand and gravel (Fig. 1b). A monitoring borehole QT09 (30 m in deep at 4538 m a.s.l.) and an automatic weather station (AWS) automatically recorded long-term observed basic meteorological data, including the soil moisture content in the active layer (October 2009 to December 2018), and soil temperature at multiple depths (January 2005 to December 2017). Approximately 4 km from the COS, another 30 m deep borehole BT01 (4530 m a.s.l.) was drilled in sparse dry steppe with considerable coarse sand and gravel, where continuous soil temperature measurements were taken continuously at depths of 0.5 to 30 m span from 2004 to 2017. In these two sites, the soil moisture content in shallow layers (<1.1 m) ranged from 15 to 39 % and from 4 to 15 %, respectively, and the organic matter content of 4.2 % and 1.68 %, respectively (Liu et al., 2020).

In addition, during August 2012, thirteen boreholes from 8 to 15 m depth (XD1-1-XD1-6, XD2-1-XD2-7) were drilled along parallel altitudinal transects at the east (3.15 km length) and west (3.86 km length) part of the Xidatan (Luo et al., 2018). The soil temperature records are available at these borehole locations covering November 2012 to September 2017. The six boreholes (XD1-1-XD1-6) are located in dry and sparse grassland on the eastern altitudinal transect between 4368 m a.s.l. and 4380 m a.s.l. (Fig. 1g), among these, the XD1-1-XD1-4 boreholes are all 15 m deep, and the two other boreholes (XD1-5-XD1-6) are 8 m depth. A frozen layer has been observed

in the five uppermost boreholes (XD1-1~XD1-5), while it was absent in the lowermost borehole
165 XD1-6 (Luo et al., 2018). Similarly, seven boreholes were drilled at the western side of the Xidatan,
resulting in an altitudinal transect from 4490 m a.s.l. (on the north) down to 4507 m a.s.l. (on the
south). The first three boreholes (XD2-1 to XD2-3) and XD2-6 are 15 m deep in sparse grassland.
Similarly, boreholes XD2-4, XD2-5, and XD2-7, are 15 m, 15 m, and 8 m deep and located in river
erosion-induced sand-rich sediment (Fig. 1e). The ground temperature monitoring results showed
170 that permafrost existed in boreholes XD2-1 to XD2-3 and XD2-6, but there is no permafrost in
boreholes XD2-4 to XD2-5 and XD2-7 (Luo et al., 2018; Yin et al., 2021).

The air temperature (height of 2, 5, and 10 m) and the volumetric unfroze soil water content in
the active layer were recorded by a CR1000/CR3000 data acquisition instrument (Campbell
Scientific Inc., USA, with ± 0.5 °C accuracy), and by a hydra-soil moisture sensor connecting to a
175 CR1000 datalogger (Campbell Scientific, USA, with an accuracy of ± 2.5 %). A cable equipped
with 20 to 30 high accuracy (± 0.1 °C) thermistors (SKLFSE, CAREERI, CAS) chain is connected
to a CR3000/CR1000 (Campbell Scientific, Logan, UT, USA) data loggers and vertically arranged
at depths from 0 to 30 m (the depths are not the same for all sites, details are given Table A1). The
ground temperature has been recorded automatically every 1 or 4 h at different depths. A more
180 detailed description of the dataset, as well as the thermistor set up and installations, can be found in
Luo et al. (2018) and Zhao et al. (2021). Before further proceeding, errors in the sensor were
identified and fixed, and the outliers were replaced with values generated by the data before and
185 after (see Zhao et al. (2021) for more details on the quality control procedures). Then, the data were
re-sampled for the daily average, used to calibrate, and validate the model performance. The spatial
distribution of these borehole sites is displayed in Fig. 1a, and the crucial information about these
boreholes employed for model calibration and validation is summarized in Tab. A1.

2.2.2 Meteorological observations from the Chinese Meteorological Administration

The observed temperature dataset from China Meteorological Administration (CMA) ground-
based meteorological stations were used to extend the land surface temperature (LST) series since
190 the 1970s. For that, observed daily mean air temperature data for the 1970 to 2019 period at two
AWS of CMA nearby (Wudaoliang:35°13' N, 93°05' E and Golmud:36°25' N, 94°55' E) was

downloaded from China Meteorological Data Sharing Service System (<http://data.cma.cn/>).

2.2.3 Remotely sensed land surface temperature datasets

Modified Moderate Resolution Imaging Spectroradiometers Land Surface Temperature (MODIS LST) product is used to force transient heat flow model for spatial modeling of alpine permafrost distribution. The MODIS onboard the Terra and Aqua satellites have provided LST measurements at a spatial resolution of $1\text{ km} \times 1\text{ km}$ since 2003 (<https://modis.gsfc.nasa.gov/>). Here, we employ clear-sky MOD11A2 (Terra MODIS) and MYD11A2 (Aqua MODIS) products (processing version 6), which contain two observations (day time and night-time) per day for the same pixel (Zou et al., 2017). Before proceeding, time series of irregularly spaced observations owing to clouds or other factors were identified, and gaps were filled by the Harmonic Analysis Time-Series (HANTS) algorithm (Obu et al., 2019). An empirical model (Zou et al., 2014, 2017) was subsequently established to get mean daily values from Aqua and Terra daytime and nighttime transient LST. Notably the model validation was quite well over the Xidatan, with the square of the correlation coefficients (R^2) above 0.9, $P < 0.01$. Details of these algorithms can be found in Xu et al. (2013) and Zou et al. (2014).

2.2.4 Additional validation datasets

The comprehensive investigation of permafrost and its environments in the Xidatan was conducted in 1975 and 2012, respectively (Nan et al., 2003; Luo et al., 2018). The lowest elevation of permafrost boundary in 1975 and 2012 was approximately 4360 m a.s.l and 4388 m a.s.l., respectively, by ground-penetrating radar (GPR) profiles combined with drilling boreholes. Subsequently, permafrost distribution in this region was delineated on a topographic map at a scale of 1:50000 by hand empirically using contour elevations line based on the field survey data, aerial photographs, and satellite images (Fig. 2a–b). In addition, one benchmark map of permafrost distribution in 2016 was accomplished by Zou et al. (2017), simulated by the temperature at the top of the permafrost (TTOP) model (Fig. 2c). The abovementioned three maps were used as the validation data to evaluate model performance in permafrost distribution. Furthermore, the long-term continuous ALT observed dataset in the BT01, QT09, XD-1, XD2-4, and XD2-6 interpolated from the in-situ soil temperature profile (Liu et al., 2020; Yin et al., 2021) were also used to evaluate

220 the model performance. Moreover, the observed permafrost distribution of boreholes (CRSQTP, JXG, XD1, XD2, XD3, XD4, XDT1, XDT2, CN13) was used to assist in determining whether permafrost exists or not.

2.3 Methods

2.3.1 Model description

225 We simulated the subsurface temperature dynamics along the soil column by numerically solving the one-dimensional transient Fourier's law heat conduction equation. The model physical basis and operational details are documented in Sun et al. (2019), and only a brief overview of the model properties for a single grid cell is given here. Ground temperature T changes over time t and depth Z through heat conduction, as described by:

$$C_{eff}(z, T) \frac{\partial T}{\partial t} - \frac{\partial}{\partial z} \left(k(z, T) \frac{\partial T}{\partial z} \right) = 0 \quad (1)$$

230 A constant geothermal heat flow of $Q_{geo}=0.08 \text{ W m}^{-2}$ as the lower boundary condition (Wu et al., 2010), and LST as the upper boundary condition. The thermal properties of the ground are described in terms of heat capacity C , thermal conductivity k and total volumetric water/ice content VWC . The latent heat effects of the water–ice phase transition is accounted for in terms of an effective heat capacity $C_{eff}(z, T)$. The heat transfer equation (Eq.1) was discretized along with a soil 235 domain to 100 m depth using finite differences. Subsequently, the trapezoidal rule was applied to numerically solve moderately stiff ordinary differential equations (Schiesser, 1991; Westermann et al., 2013). With comprehensive consideration of the modeling precision and computation cost, we choose the calculated time step to be one day and set a total of 282 vertical levels for each soil column, with the vertical resolution configurations of 0.05m (the upper 4 m) and 0.5m (remaining 240 soil layer to 100 m).

2.3.2 Model calibration and validation

We selected four borehole sets (Fig. 1a), which represented different soil type classes with various thermal properties for the initial model calibration, and the remaining sites for cross-

validation. The sites were selected based on surface deposits, vegetation coverage, and soil types at 245 a 1 km×1 km spatial resolution (Li et al., 2015b, Luo et al., 2018). Thermophysical properties (e.g., stratigraphies, texture, ground ice content, organic matter content, dry bulk density) of distinct soil layers were measured or assessed from field surveys, laboratory and on-site measurements of soil samples obtained from fifteen borehole cores (depths between 8~30 m). These boreholes were specific for each soil class and geographical location. The detailed information for the bulk density 250 and moisture content measurements of soil samples, is referred to Zhao et al. (2015). Furthermore, a time series of observed soil water content dataset in the active layer (Sun et al., 2019, 2022; Zhao et al., 2021) vicinity of the site (QT09) and the ground ice distribution maps accomplished by Zhao et al. (2010) is used for water content estimates of each soil type. And then, we pre-selected narrow 255 ranges of plausible values of typical soil thermophysical parameters (thermal conductivity and heat capacity, details see Table A2); and fine-adjusted during model calibration. The manual stepwise optimization procedures were used to adjust parameters based on the suggestions by Hipp et al. (2012). Specifically, calibration was performed by systematically changing k over the given plausible ranges to improve the agreement between the simulated and observed ground temperature at different depth levels. Subsequently, minor adjustments were made to C to promote the model's 260 performance.

The model was initialized by cyclical forcing of the first year LST data until the soil temperature profile reached a steady state to estimate an initial temperature profile. The number of spin-up cycles was between 2000 to 3300, and the criterion of soil temperature profile reached equilibrium under the upper and lower boundary condition was set at less than 0.0001 °C/cycle. The 265 last-day ground temperature profile was subsequently used as the initial condition for subsequent modeling. The agreement between the model grid and borehole monitoring site was quantified at each depth in terms of the mean absolute error (MAE) and root mean square error (RMSE) (Willmott and Matsuura, 2005; Jafarov et al., 2012):

$$MAE = \frac{1}{n} \sum_{i=1}^n |Ob_i - Sm_i| \quad (2)$$

$$RMSE = \sqrt{\frac{\sum_{i=1}^n (Ob_i - Sm_i)^2}{n}} \quad (3)$$

where Ob_i , Sm_i is observation and simulation value, respectively. And n is the total amount of data. The MAE shows an overall error between observing and simulating when the $RMSE$ emphasizes an error variation.

2.3.3 Historical and future long-term LST series

We extended LST by establishing statistical relationships between local LST and air temperature (AT) from nearby AWS to derive historical and future LST series for each grid from historical (1970–2019) AT observation and the multi-model ensemble AT projection by 2100 under different climate change scenarios.

The AT_cma, AT, and LST denote the air temperature from the CMA, air temperature at 2 m from our COS and ground surface temperature derived from modified MODIS LST, respectively. Firstly, we established a linear regression between LST and AT from the measured period of 2004 to 2018, where the temperature variability was highly correlated between LST and AT with $R^2 = 0.83$, $P < 0.01$; secondly, the daily AT series from 1970 to 2019 were generated utilizing a stepwise linear regression between measured AT from 2004 to 2018 and those extracted from CMA meteorological stations (AT_cma) nearby which worked well with $R^2 = 0.88$ $P < 0.01$; In the third, we generated a time series of LST staring from 1970 based on the AT–LST linear regression model induced in step 1 and extending series AT in step 2.

For future AT projections, the Sixth Assessment Report of the Intergovernmental Panel on Climate Change Work Group1 (IPCC WG 1 AR6) (Iturbide et al., 2020; IPCC, 2021) has evaluated and projected climate change over the QTP during the 21st century (<https://interactive-atlas.ipcc.ch>). The model estimated warming between 1995–2014 and 2081–2100 in mean annual AT in QTP under three RCPs scenarios is $0.013 \text{ }^{\circ}\text{C a}^{-1}$ (RCP2.6, low concentration of emissions), $0.028 \text{ }^{\circ}\text{C a}^{-1}$ (RCP4.5, stable concentration of emissions) and $0.060 \text{ }^{\circ}\text{C a}^{-1}$ (RCP8.5, high concentration of emission), respectively, calculated from multi-model ensemble median (21–29 model outputs) of CMIP5. The

mean warming rate is $0.017^{\circ}\text{C a}^{-1}$ (SSP1–2.6, strong climate change mitigation), $0.032^{\circ}\text{C a}^{-1}$ (SSP2–4.5, moderate mitigation), and $0.064^{\circ}\text{C a}^{-1}$ (SSP5–8.5, no mitigation), estimated from CMIP6 ensemble median of 31–34 model outputs. Using the AT–LST linear regression relationship model, we obtained a mean LST warming rate of $0.012^{\circ}\text{C a}^{-1}$ (RCP2.6), $0.025^{\circ}\text{C a}^{-1}$ (RCP4.5), and $0.050^{\circ}\text{C a}^{-1}$ (RCP8), and a mean LST increase rate of $0.015^{\circ}\text{C a}^{-1}$ (SSP1–2.6), $0.030^{\circ}\text{C a}^{-1}$ (SSP4–4.5) and $0.057^{\circ}\text{C a}^{-1}$ (SSP5–8.5).

2.3.4 Spatial modeling

The extended and projected LST were used to force our calibrated model for simulating the spatial distribution of permafrost in the Xidatan. The ground thermal regime was simulated for a specific ground stratigraphy under boundary conditions from a one-dimensional multilayer soil profile down to 100 m depth at each grid point. The thermophysical parameters of multilayer soil columns were specified and assigned for each soil type based on the soil type map at $1\text{km} \times 1\text{km}$ spatial resolution (Li et al., 2015b). If the maximum temperature of any soil layer in the grid point was $\leq 0^{\circ}\text{C}$ for two consecutive years, the model cells were identified as permafrost. In contrast, the seasonally frozen ground was defined from the not-yet-assigned cells, in which the minimum soil temperature of any layer in the same two years was $\leq 0^{\circ}\text{C}$. The remaining cells were unfrozen ground (Wu et al., 2018). The continuous permafrost zone is defined as the region where the area coverage of permafrost is more than 90 % (of the total accounts area). Otherwise, it was demarcated as a discontinuous permafrost zone (Qin et al., 2014). The simulation domain comprises about 280 km^2 with a horizontal resolution of $1\text{ km} \times 1\text{ km}$, corresponding to 280 independent runs.

3 Result

3.1 Model evaluation

Simulated ground temperatures results demonstrate relatively larger bias (with the MAE ranging from 0.69 to 2.02°C and RMSE ranging from 0.87 to 2.46°C) for surface soil layer to 1 m in depth at all calibration sites (Fig. Tab. A1). These could be explained by frequent fluctuation and complex variation pattern of ground temperature itself at shallow depth greatly affected by local factors (e.g., terrain, water bodies, snow cover, vegetation, etc.). However, these discrepancies

320 between simulated and observed ground temperature gradually reduce with the increase of soil depth. Most calibration boreholes showed a good correspondence between modelled and measured ground temperature at the intermediate (3 m, 8 m) and deep (15 m, 30 m) layers (Fig. A2), with an MAE of 0.05–0.52 °C, 0.04–0.38 °C, as well as an RMSE of 0.06–0.58 °C, 0.04–0.38 °C, respectively (Tab.1). The same pattern appeared at validation sites (Fig. A3–4). Ground temperatures in 325 validation sites were equally well reproduced by the calibrated model, yielding an MAE of 0.86–1.27 °C (RMSE of 1.15–1.63 °C) in the 0.5 and 1 m and 0.01–0.52 °C (RMSE of 0.08–0.80 °C) in 3 and 15 m (Tab.1). Generally, the consistent daily fluctuations of the simulated and observed soil temperature at all observational depths for most calibration and validation sites indicated the satisfactory simulation by our calibrated model.


330 Site XD2–6 has relatively poor performance in the deep layer (8 m and 15 m) compared with the shallow layers (Fig. A2). The deviation between measured and simulated soil temperature in this special case might be caused by micro-scale heterogeneity in surface cover, topography, and soil stratigraphy at the sub-grid scale, which led to more difficulty in accurate modeling. Nevertheless, the deviation between this site-modelled results and measured values is within 0.38 °C 335 at the deep layer (15 m). Furthermore, the permafrost at this site was simulated to disappear in the mid–late 2010s, which was in line with the observation (Yin et al., 2021).

Table 1. Error metrics for assessing daily average ground temperature at different depths derived from the observed with simulated for individual calibration and validation site (good criteria values<0.20 °C are displayed in italics).

Criteria	Site	0.5 m	1 m	3 m	8 m	15 m	30 m
MAE	BT01	1.04	1.04	0.52	0.41	<i>0.19</i>	0.09
(°C)	XD2-7	2.02	1.46	0.38	<i>0.05</i>		
	QT09	1.06	0.89	0.23	<i>0.16</i>	<i>0.18</i>	<i>0.04</i>
	XD2-6	1.42	0.69	0.23	0.22	0.38	
	XD2-1	1.05	0.95	0.41	<i>0.13</i>	<i>0.19</i>	
	XD2-4	1.01	0.86	0.21	<i>0.14</i>	<i>0.01</i>	
	XD1-1	1.27	1.18	0.52	0.25	0.19	
	XD1-4	1.11	0.92	0.44	<i>0.19</i>	<i>0.08</i>	
RMSE	BT01	1.36	1.38	0.72	0.41	<i>0.19</i>	0.09
(°C)	XD2-7	2.46	1.79	0.58	<i>0.06</i>		
	QT09	1.40	1.48	0.40	<i>0.17</i>	<i>0.18</i>	<i>0.04</i>
	XD2-6	1.78	0.87	0.30	0.23	0.38	
	XD2-1	1.36	1.20	0.54	0.24	<i>0.19</i>	

XD2-4	1.31	1.15	0.35	0.14	0.02
XD1-1	1.63	1.48	0.80	0.25	0.19
XD1-4	1.41	1.19	0.62	0.20	0.08

340 To better estimate the model performance in spatial modeling, we compared our simulations with three permafrost maps investigated in 1975, 2012, and 2016. Based on the validation of the various maps against the permafrost and seasonally frozen ground observation at 24 boreholes (Fig. 2), we found that both the 1975 and 2012 maps can well interpret the continuous permafrost zone at the central-western Xidatan. However, there are many erroneous (12.5 % for 1975, and 16.6 % for 345 2012) in recognitions of seasonally frozen ground at the discontinuous permafrost zone. It indicated that these two permafrost maps could not well represent the historical permafrost distribution status in the study region permafrost and seasonally frozen ground coexist zones. In addition, these two maps are strongly inconsistent with the 2016 map and our simulations (Fig. 2a–b). The 2016 map and our simulations showed a consistent permafrost distribution pattern and correctly identified 350 almost all continuous permafrost locations (Fig. 2c–f). However, a slight discrepancy existed between the 2016 map and our simulation in permafrost (8.3 %) and seasonally frozen ground (8.3 %) locations over the margins of the discontinuous continuous permafrost zone. Our simulated results were consistent well the investigated results; and indicated its good recognition of the seasonally frozen ground in this region.




355

Figure 2. Geographic distribution of permafrost and seasonally frozen ground across the Xidatan for three permafrost maps accomplished in 1975, 2012 and 2016 (left panels 1975 (a), 2012 (b), 2016 (c), published in Nan et al. (2003), Luo et al. (2018) and Zou et al.(2017)) compared to corresponding modelled outputs (right

panels, 1975 (d), 2012 (e), 2016 (f)).

360 Continuous multi-year ALTs derived from five monitoring sites were compared with those
from the model-simulated (Fig. 3). The results indicated that there is a strong positive correlation
between the simulated and observed ALT ($R^2=0.98$, $P<0.01$), and the simulation bias in the ALT
from these sites are within ± 0.25 m. In terms of geographical structure, the spatial characteristics of
ALT across the study area are well captured by our model. Both observed and simulated ALT in
365 XD2–6 varied from 4.15 to 4.31 m, which is higher than other sites (BT01 of 2.55 to 2.85 m; the
QT09 of 1.45 m to 1.60 m; the XD2–1 of 2.30 to 2.48 m; the XD2–3 of 2.95 to 3.05 m).



370 **Figure 3. Comparison between annually observed ALT and simulated at different sites (the TB01 and QT09**
(Liu et al. (2019), Zhao et al. (2021) observed from period 2005 to 2017, 2005 to 2018 are available, respectively,
observation period at the XD2-1, XD2-3, and XD2-6 (Yin et al. (2021)) are from 2013 to 2019, 2013 to 2017,
respectively. The solid line is a 1:1 line and the dashed line shows biases within ±0.25 m, dots are colored to
represent the different sites).

3.2 Historical permafrost evolution

375 Our simulation outputs were combined with topographic data (elevation and slopes) derived

from 30 m-DEM to analyse the permafrost distribution and its dynamics. MAGT at the depths of ZAA, permafrost table, permafrost base, and permafrost thickness are defined from vertical temperature profiles, as critical parameters to describe the permafrost thermal regime, which was also chosen for analysis and discussion. Areas with t seasonally frozen ground were excluded from

380 the subsequent studies.

3.2.1 Initial situation of permafrost distribution

The simulation results (Tab.2, Fig. 4) showed the initial situation in 1970. The lower limit of continuous permafrost modelled was ca.4525 and 4732 m a.s.l., respectively, on the north-and south-facing slopes. While the lowest elevation of the permafrost boundary simulation was 4138 m

385 a.s.l (on the north-facing slopes) and 4357 m a.s.l (on the south-facing slopes). Approximately 80 % of the total counting area was underlain by permafrost (33.93 % was continuous, and 46.07 % was discontinued) in the Xidatan. Regionally, the distribution characteristics of permafrost conditions are predominantly controlled by elevations. With altitude ascending westward gradually, permafrost temperature and permafrost table show a decreasing trend, whereas the position of permafrost base and permafrost thickness increase. Furthermore, local topographic factors in slope also govern permafrost distribution in the study area. Permafrost temperature on the north-facing slopes were far colder than that of on south-facing slopes within the same elevations (Fig. 4a). On the south-facing slopes (with high altitudes above 4500 m a.s.l.) and north-facing slopes modelled shows a comparatively cold permafrost temperature (MAGT ranges from -0.5 to -4.5 °C). Simulated 390 permafrost table was less than 2.5 m, permafrost base of 20 to 48 m, and permafrost of up to 46 m at the maximum. Whereas on the south-facing slopes with low altitudes below 4500 m a.s.l., MAGT modelled is higher than -0.5 °C, the position of the permafrost table modelled varies from 2.5 to 4.5 m, permafrost base is at a depth of fewer than 20 m, and permafrost thickness of approximately 4 m at the thinnest.

400




Figure 4. Spatial distributive features of MAGT (a), permafrost table (b), permafrost base (c), and permafrost thickness (d) for the initial simulation of the 1970s over the Xidatan (grey areas with the seasonally frozen ground were excluded).

3.2.2 Changes in permafrost conditions

405 From 1970 to 2019, the simulation results indicate that the lower limit of the continuous permafrost zone remained unchanged over the study areas. The lowest elevation of the permafrost boundary has a remarkable rise of 47 m on the north-facing slopes, while that remained unchanged on the south-facing slopes. Correspondingly, around 12.86 % of the discontinuous permafrost zone has transformed into seasonally frozen ground (Tab. 2), which caused the northern boundary of the 410 discontinuous permafrost zone to have approximately retreated southwards 1~2 km, but that is unchanged for the continuous permafrost zone (Fig. 5).




Figure 5. Spatial distributive changes of continuous and discontinuous permafrost; and seasonally frozen ground zone over the Xidatan from 1970 to 2019.

415 **Table 2. Variations of the permafrost boundary and areal extent of frozen ground type over the Xidatan for 1970-2019, and that of projected variations by 2100 under different climate change scenarios.**

	The lower limit or the lowest elevation of permafrost boundary (m a.s.l.)		Areal extent (%)		
	North-facing	South-facing	Con.	Disc.	Seas.
1970	4525(4138)	4732 (4357)	33.93	46.07	20.00
2019	4525 (4185)	4732 (4357)	33.93	33.21	32.86
SSP1–2.6 (2100)	4567 (4308)	4732 (4516)	28.57	30.36	41.07
SSP2–4.5 (2100)	4567 (4308)	4732 (4516)	28.57	28.57	42.86
SSP5–8.5 (2100)	4567 (4309)	4754 (4570)	27.14	21.79	51.07
RCP2.6 (2100)	4567 (4308)	4732 (4416)	28.57	30.36	41.07
RCP4.5 (2100)	4567 (4308)	4732 (4516)	29.29	27.50	43.21
RCP8.5 (2100)	4567 (4309)	4737 (4558)	28.93	22.50	48.57

Note: Outside brackets are the lower limit of the permafrost, while in brackets are the lowest elevation of the permafrost boundary. Con., Disc., and Seas. were indicate continuous permafrost, discontinuous permafrost, and seasonally frozen ground.

420 The regional-average MAGT has increased by 0.44 °C over the past 50 years for permafrost characteristics. With temperature warming, we found a gradual decline with a mean amplitude of 0.36 m in the position of the permafrost table whereas a drastic moved-up permafrost base is 1.12 m. Correspondingly, permafrost had thawed an average of nearly 1.54 m in thickness. Spatially, the mean MAGT warmed up to 0.49 °C, and the average permafrost table declined by 0.37 m for the 425 continuous permafrost zone, but its permafrost base (around -0.80 m) and thickness (around -1.18 m) variations were comparatively slight. By comparison, comparatively low variations in MAGT (0.40 °C) and in the permafrost table (average declined by 0.76 m), but dramatic changes of -4.23 m occurred in the discontinuous permafrost zone, which is roughly twice compared to changes on the continuous permafrost area. Correspondingly, an average of about -1.96 m in thick permafrost 430 had quickly thawed, owing to a remarkably rising effect of the permafrost base.

Table 3. Changes in characteristics of frozen ground type over the Xidatan for 1970 to 2019, and projected changes by the 2090s, relative to the 2010s, under different climate change scenarios.

Types	1970–2019	SSP1–2.6	SSP2–4.5	SSP5–8.5	RCP2.6	RCP4.5	RCP8.5	
MAGT (°C)	Con.	0.49	0.73	0.94	1.03	0.65	0.91	1.06
	Disc.	0.40	0.53	0.66	0.96	0.48	0.65	0.86
Permafrost table (m)	Con.	0.37	0.56	1.76	6.24	0.44	1.23	4.95
	Disc.	0.35	0.87	3.13	7.02	0.64	2.26	6.13
Permafrost base (m)	Con.	-0.80	-3.52	-3.87	-3.99	-3.41	-3.81	-4.13
	Disc.	-1.60	-4.87	-5.09	-5.17	-4.80	-5.08	-5.17


Permafrost thickness (m)	Con.	-1.18	-4.11	-5.23	-10.38	-3.87	-5.11	-9.42
	Disc.	-1.96	-5.78	-7.94	-12.76	-5.46	-7.44	-11.65

Note: Con., Disc., and Seas. indicate continuous permafrost, discontinuous permafrost, and seasonally frozen ground.

3.3. Projection of permafrost condition

435 The projected changes in the lower limit of permafrost, the lowest elevation of permafrost boundary, and the spatial distribution of continuous, discontinuous permafrost, and seasonally frozen ground as well as its characteristics (MAGT, permafrost table, permafrost base, and permafrost thickness), are presented in Tab.2-3 and Fig.6.

440 The result indicates that the lower limit of permafrost on the north-facing slopes is projected to increase by 42 m until 2100, relative to 2019, under all RCPs or SSPs. On the south-facing slopes, this value is about 22 m under very high emission scenarios (SSP5-8.5 or RCP8.5), which is far smaller than the changes in the lowest elevation of the permafrost boundary. The lowest elevation of the permafrost boundary on the north-facing slopes is projected to increase by 123 m by 2100, relative to 2019, under both low and medium emission scenarios, and by 124 m under very high 445 emission scenarios SSP5-8.5 or RCP8.5. South-facing slopes are projected to increase by 159 m by 2100, compared to 2019, under both low and medium emissions of RCP or SSP scenarios. Still, a more pronounced increase of around 213 and 201 m is projected under SSP5-8.5 and RCP8.5. Relative to 2019, the areal extent of continuous permafrost zone is projected to decrease by 5.36 (5.36), 5.36 (4.64), and 6.79 (5.00) %, respectively, by 2100, under SSP1-2.6 (RCP2.6), SSP2-4.5 (RCP4.5), and SSP5-8.5 (RCP8.5), respectively. Compared with the decrease of 3.57 (2.85), 4.64 (5.71), and 11.42 (10.71) % for discontinuous permafrost zone. In contrast, the areal extent of 450 seasonally frozen ground is projected to increase by 8.93 (8.21), 10.00 (10.36), and 18.21 (15.71) %, respectively, by 2100, relative to 2019, under SSP1-2.6 (RCP2.6), SSP2-4.5 (RCP4.5), and SSP5-8.5 (RCP8.5). Northern limit of the continuous permafrost zone is projected to retreat southwards around 1~2 km under SSP1-2.6 (RCP2.6) or SSP2-4.5 (RCP4.5) or RCP8.5, and about 1~3 km under SSP5-8.5. By comparison, the northern boundary of the discontinuous permafrost zone is anticipated to shift southward around 1 km under SSP1-2.6 (RCP2.6) or SSP2-4.5 (RCP4.5), and around 1~2 km under SSP5-8.5 (RCP8.5).



460 **Figure 6. Projected spatial distributive changes of frozen ground type over the Xidatan by 2100 under RCPs and SSPs scenarios (left column, from top to bottom, each row shows under SSP1–2.6, SSP2–4.5, and SSP5–8.5 scenarios, right column, from top to bottom, each row shows under RCP2.6, RCP4.5, and RCP8.5 scenarios).**

465 Under global climate warming scenarios, the permafrost temperature is anticipated to increase further, but its variation lags substantially behind the changes in air temperature. Relative to 2010s, the regional-average MAGT is projected to warm by 0.63, 0.81, and 0.99 °C, respectively, by 2090s, under SSP1–2.6, SSP2–4.5, SSP5–8.5, which is slightly higher than that of RCP scenarios (0.56, 0.78, and 0.98 °C, respectively). Along with MAGT rising, relative to the 2010s, the permafrost table is projected to further decline by 0.72 to 6.70 m under SSP scenarios (0.72 m, 470 2.48 m, and 6.70 m, respectively, under SSP1–2.6, SSP2–4.5, and SSP5–8.5), and decline by 0.54 to 5.47 m under RCP scenarios (0.54 m, 1.73 m, and 5.47 m, respectively, under RCP2.6, RCP4.5, and RCP8.5), at the end–century (the 2090s). The average permafrost base is projected to rise by 4.22, 4.54, and 4.56 m, respectively, by the 2090s, compared to the 2010s, under SSP1–2.6, SSP2–4.5, and SSP5–8.5. Meanwhile, a relative decrease in permafrost base of 4.14, 4.43, and 4.60 m; 475 are estimated under RCP2.6, RCP4.5, and RCP8.5. An average thinning in the permafrost thickness is projected to be 4.97, 6.66, and 11.74 m, respectively, under SSP1–2.6, SSP2–4.5, and SSP5–8.5, and that would be 4.71, 6.26, and 10.43 m, respectively, under RCP2.6, RCP4.5, and RCP8.5.

Spatially, the average MAGT is projected to rise by 0.73 (0.65), 0.94 (0.91), and 1.03 (1.06) °C,

respectively, for continuous permafrost zone, under SSP1–2.6 (RCP2.6), SSP2–4.5 (RCP4.5), and
480 SSP5–8.5 (RCP8.5). Compared with the rising of 0.53 (0.48), 0.66 (0.65), and 0.96 (0.86) °C,
respectively, for discontinuous permafrost zone. As for the permafrost table, both continuous and
discontinuous permafrost zone is projected to gradual decline under SSP1–2.6 (0.56 and 0.87 m),
and RCP2.6 (0.44 and 0.64 m), but a remarkable decline is projected under medium, and very high
emission scenarios, and a more pronounced decline are anticipated under SSPs scenarios than that
485 projection under RCPs scenarios. The average permafrost table in continuous permafrost zone is
projected to decline by 1.76 (1.23) and 6.24 (4.95) m, respectively, under SSP2–4.5 (RCP4.5) and
SSP5–8.5 (RCP8.5). Compared with the decline of 3.13 (2.26) and 7.02 (6.13) m, respectively,
under SSP2–4.5 (RCP4.5), and SSP5–8.5 (RCP8.5) for continuous permafrost zone. Permafrost
base is projected to move remarkably up under all scenarios. For continuous permafrost zone,
490 permafrost base is projected to rise by 3.52 (3.41), 3.87 (3.81), and 4.13 (3.99) m, respectively,
under SSP1–2.6 (RCP2.6), SSP2–4.5 (RCP4.5), and SSP5–8.5 (RCP8.5), which is slightly smaller
than that projected for discontinuous permafrost zone (4.87 (4.80), 5.09 (5.08), and 5.17 (5.17) m,
respectively, under SSP1–2.6 (RCP2.6), SSP2–4.5 (RCP4.5), and SSP5–8.5 (RCP8.5)). The
average permafrost thickness of continuous and discontinuous permafrost zone is projected to thing
495 4.11 (3.87) and 5.78 (5.46) m, respectively, under SSP1–2.6 (RCP2.6) as the main effect of
permafrost base move up. Whereas a more prominent decrease of 5.23 (5.11) and 7.94 (7.44) m,
respectively, under SSP2–4.5 (RCP4.5), 10.38 (12.76), and 9.42 (11.65) m, respectively, under
SSP5–8.5 (RCP8.5), owing to both effect of permafrost table declining and the permafrost base
rising.

500 4 Discussion

4.1 Comparison with previous studies

505 In this work, our simulated distribution of continuous permafrost zone had a substantial
agreement with three permafrost maps investigated in 1975, 2012, and 2016. Still, a remarkable
difference in the discontinuous permafrost zone where permafrost and seasonally frozen ground
coexist (Fig. 2). Compared with the 2016 map and our simulated results, the 1975 and 2012 maps
underestimated the permafrost area in the discontinuous permafrost zone. This contradiction might

be due to differences using data, methods, study periods, spatial resolutions, etc. (Yang et al., 2010; Ran et al., 2012; Zou et al., 2017). The 1975 and 2012 maps were plotted on a topographic map at a 1:50000 scale based on field investigations, aerial photographs, and satellite images (Nan et al., 510 2003; Luo et al., 2018). These coarse-resolution maps cannot accurately consider the effect of local factors since they cannot describe variations in ground conditions over a short distance (Zhang et al., 2013). Moreover, comparing them with field observations makes the result difficult to validate. Although the 1975 and 2012 maps may represent the corresponding permafrost status in that year, they are limited by field investigations, and there is not a clear understanding of whether permafrost 515 existed in the northeaster high-altitude areas or not. In the 2012 map, these isolated top mountain areas are uniformly considered as the seasonally frozen ground when permafrost mapping (Luo et al., 2018), which is unreasonable and underestimated in the areal coverage of the permafrost in that area. Furthermore, the artefactual errors were hard to control when mapping permafrost distribution by conventional cartographic techniques that manually delineated the permafrost boundaries on the 520 topographic maps (Zou et al., 2017; Ran et al., 2012). These factors inevitably led to existing uncertainties in the 1975 and 2012 maps.

By comparison, the 2016 map and our simulation results have a much higher spatial resolution (1 km×1 km) than field-investigated-based ones (e.g., 1975 and 2012 maps) by improved MODIS LST application. In addition, it showed higher and more accuracy in identifying both permafrost 525 and seasonally frozen ground boreholes and performed better recognizing the seasonally frozen ground in regions with complex terrain. This finding highlights the potential advantage of remote sensing-based data in improving the spatial modeling of marginal permafrost simulations on the QTP. Overall, our simulated distribution of continuous permafrost and discontinuous permafrost zone were similar to that of the 2016 map. Differences mainly due to the TTOP model did not 530 consider the thermal state of the deep permafrost. Therefore, the areal extent of permafrost distribution in the 2016 map likely was slightly underestimated compared with our simulation results. Moreover, the 2016 map assumes that permafrost is in equilibrium with the long-time climate. However, the ground temperature observations of permafrost on the QTP have increased 535 during the past several decades (Zou et al., 2017; Zhao et al., 2010, 2020; Yao et al., 2019; Ehlers et al., 2022), and this means a disequilibrium of permafrost under ongoing global warming. So, a map

based on a contemporary climate forcing will likely underestimate permafrost extent (Zou et al., 2017). By contrast, in our study, we used a transient numerical heat conduction permafrost model, which integrated climate and ground condition variables to quantify the change in permafrost. Our model performed well in modeling the disappeared evolution of two permafrost islands since 1975 540 and shifting the northern boundary of discontinuous permafrost (Fig. 2d–f), which can be confirmed by direct observation (Jin et al., 2006; Luo et al., 2018; Yin et al., 2021). These phenomena implied our model could accurately capture marginal permafrost thermal state dynamics under a warming climate.

Furthermore, using our model, we quantified the spatial distribution of permafrost over the 545 study area. We simulated a striking elevation dependence in permafrost distribution. Specifically, permafrost temperature decreases, decreases in thickness, and the permafrost table becomes thinner with the increased elevation, which is consistent with previous observation-based studies (Cheng et al., 1984; Wu et al., 2010; Zhao et al., 2010, 2019, 2020; Li et al., 2012; Jin et al., 2011; Luo et al., 2018). Moreover, Cheng et al. (2019) further indicated that the MAGT varied from -5 to 0.5°C, and 550 the average permafrost thickness was approximately 26 m, as deduced by considerable monitoring and field investigation dataset. The monitoring network of ALT along the Qinghai–Tibet Highway (QTH) (Li et al., 2012) has demonstrated that the mean ALT was 218 cm, ranging from 100 to 320 cm from 1981 to 2010. This evidence strongly corroborates our simulation results accuracy, giving us more confidence in upscaling our model to the study area to investigate spatiotemporal dynamics 555 and anticipate possible changes in permafrost.

4.2 Process of permafrost degradation

In this paper, we simulated a slow response of the permafrost thermal state to a warming 560 climate in the northern lower limit of the permafrost zone (Xidatan) on the QTP. As shown in our simulation, from 1970 to 2019, we simulated that roughly 12.86 % of the discontinuous permafrost zone over the study area has ultimately converted into seasonally frozen ground, which is very close to observed data (13.8 %) here in 2012 (Luo et al., 2018). Permafrost distribution and its thermal conditions over the study area were spatially controlled by elevation. In addition, due to the orientation of slope influenced the amount of solar radiation received by the ground surface (Cheng

et al., 2004). Specifically, much thicker, colder permafrost and a thinner ALT on the north-facing slopes than on the south-facing slopes within the same elevation. So, a distinct spatial discrepancy of permafrost thermal regimes in response to a warming climate as different thermal states. Over the past 50 years, the rising rate of MAGT for the continuous permafrost zone was relatively faster (regional-average warmed by 0.49 °C) due to more energy being available to heat the ground. By contrast, as permafrost temperature is close to the thawing point (about 0 °C), accumulated energy is enormously consumed by melting ground ice, and MAGT for the discontinuous permafrost zone slowly rises (regional-average warmed by 0.40 °C). Meanwhile, both the continuous (regional-average declined by 0.37 m) and discontinuous permafrost zone (regional-average declined by 0.35 m) displayed a gradual decline in the position of the permafrost table. But we simulated a drastically risen permafrost base, especially in the discontinuous permafrost zone, due to heat transfer in strata from top to bottom, leading the geothermal gradients in permafrost to keep dropping. When the geothermal gradient in permafrost temperature drops to less than that of the underlying thawed soil layers, the geothermal heat flux from the deep stratum is completely used to thaw the permafrost base. Hence, permafrost thaws from bottom to top and moves upward. As permafrost was relatively warm and thin and geothermal flow relatively high over the Xidatan (Wu et al., 2010, Sun et al., 2019), the main degradation mode of permafrost over this region is simulated to be upward thawing from the permafrost base. This degradation mode is also confirmed by several monitoring boreholes across this region (Jin et al., 2006, 2011; Cheng and Wu. 2007; Liu et al., 2020). In general, the pattern of permafrost degradation over the Xidatan from 1970 to 2019 can be summarized like as: the continuous permafrost zone has gradually converted to warm permafrost, whereas the discontinuous permafrost zone has been upward thawing remarkably. Notably, the margin of the discontinuous permafrost zone has converted to seasonally frozen ground.

As for the projections under different climate change scenarios, the latest generation of ESMs from CMIP6 projected a substantially warmer climate by 2100 than the previous generation, for instance, CMIP5 (Fewster et al., 2022). In our study, MAGT is anticipated to increase further, and the warming rate is projected to be slightly higher under SSPs than RCPs, but very small discrepancies exist among SSP and RCP scenarios in projecting changes of permafrost distribution extent. This further verify that the response of permafrost to climate warming is a slow and nonlinear

process, and its variation lags substantially behind the changes in air temperature. But contrary findings are reported by some previous studies. Based on the empirical equilibrium model, Lu et al. 595 (2017) predicted that extensive reduction of permafrost area on the QTP by the end of the 21st century under RCP2.6 (22.44 %) and RCP8.5 (64.31 %) and permafrost would retreat into the Qiangtang Plateau hinterland. Likewise, Chang et al. (2018) suggested that in the next 20 years permafrost area on the QTP are projected to shrink by 9.7 %, and 10.7 %, under RCP4.5 and RCP8.5, respectively, with projected shrinkage of 26.6 % and 32.7 % in the next 50 years. Guo and Wang 600 (2016) projected almost no permafrost on the QTP by 2080 to 2099 under RCP8.5. In addition, Yin et al. (2021) projected around 26.9 %, 59.9 %, and 80.1 % of permafrost on the QTP is likely to disappear, by the end of the 21st century under SSP1–2.6, SSP2–4.5, and SSP5–8.5 scenarios. From transient numerical modeling, Guo et al. (2012) using the Community Land Model4 (CLM4) projected an approximately 81% reduction in near-surface (<4.5 m) permafrost area on the QTP by 605 the end of the 21st century under the A1B emission scenario. Additionally, the deep permafrost of 10 and 30 m depths would be largely degraded by 2030-2050. Zhang et al. (2022) applied Noah LSM to project much of $44 \pm 4\%$, $59 \pm 5\%$, and $71 \pm 7\%$ permafrost is likely to degrade in the late 21st century, under SSP2-4.5, SSP3-7.0, and SSP5-8.5 scenarios, respectively.

610 The abovementioned results and our projections unanimously projected further degradation trend in permafrost on the QTP under warming climate scenarios, but a considerable discrepancy among results on the magnitude of permafrost degradation exists. This discrepancy can partly be attributed to those approaches that established a simple statistical relationship between the current permafrost distribution and air temperature based on the surface energy balance theory. However, permafrost in the QTP formed over a long period of cold paleoclimate and developed an energy 615 state characterized by low ground temperature and ground ice in permafrost (Buteau et al., 2004; Jin et al., 2011; Sun et al., 2019; Zhao et al., 2020). The present state of permafrost is a response to historical climatic changes and impacts (Wu et al., 2010; Cao et al., 2014). The current project of permafrost degradation from abovementioned does not consider the historical energy accumulation in permafrost and the impact of ground ice conditions buried below 1 m depth (Zhao et al., 2020; 620 Smith et al., 2022). For example, most of LSMs studies mainly focused on optimizing parametrization schemes for shallow soil layers (<4 m) and simply extending the soil column

simulation depth. Its performance in assessing the ground ice existence was poor considering the thermal state of deep permafrost (Lee et al., 2014; Sun et al., 2019). Furthermore, ignoring the geothermal heat flux by setting zero flux or constant temperature as the bottom boundary condition
625 (Wu et al., 2010; Xiao et al., 2013). These factors play a crucial role in the long-term evolution of permafrost (Zhao et al., 2020). Thus, the relationship between the decrease in the areal extent of permafrost and the warming air temperature over the present-day permafrost region is approximately linear simulated by these empirical statistics or LSMs. Such high rates of permafrost loss are not observed, indicating high sensitivity for those models predicting such losses (Zhao et
630 al., 2020).

In comparison, our model considers the thermal properties difference between frozen and thawed soil, the phase variations of the unfrozen water in frozen soil, the distribution of the ground ice, and geothermal heat flow. Thereby, it describes the heat transfer process in permafrost very well and reasonably capture the attenuation and time lag of heat transfer in deep permafrost as water or
635 ice content and ground is a poor conductor of heat. Our model is characterized by vertical modeling domains of one hundred meters with a vertical resolution of 0.05 m within the active layer (the upper 4 m) and provides sufficient accuracy to resolve the annual dynamics of active layer thawing and refreezing, as well as the evolution of ground temperatures in deeper layers. The model results were carefully validated against considerable long-term continuous monitoring of soil temperatures at
640 various depths, ALT, and observed permafrost distribution of boreholes as well as three existing permafrost distribution maps investigated in 1975, 2012, and 2016—our simulation results are in compliance with the observed facts. And the magnitude and evolution of permafrost degradation projections on the QTP derived from our transient simulations agree well with that of the heat conduction permafrost model, accounting for the thermal state of deep ground ice (Li et al., 1996; Li et al., 2008; Sun et al., 2019 and 2022). It can be noted that existing studies largely ignore the thermal properties of deeper permafrost. Our findings highlight that the initial permafrost thermal state is influenced by historical climate, stratigraphic thermal properties, ground ice distribution, geothermal heat flow, and propagation of the phase-transition interface plays a critical role in permafrost degradation.

This study may have uncertainties, including the extended MODIS LST series used as the model inputs, soil parameters heterogeneity at sub-grid in terms of surface cover, topography, and soil stratigraphy, and the permafrost model's physics. Due to a significant linear relationship between LST and AT over the study area, moreover, in this work, we mainly focus on the long-trend permafrost temperature over the foreseeable future. The biases of the estimated LST by simple regression relationship of AT–LST cannot affect the long-term mean change trend in LST. Furthermore, the one-dimensional approach of the model is another limitation, which assumes each grid cell to be uniform without lateral exchange. Our simulations, therefore, are considered as conservative changes in the ground temperature in areas with lateral water fluxes, such as flood land in the valley (Bense et al., 2012; Westermann et al., 2016; Sjöberg et al., 2016). The representation of the horizontal fluxes exchange of heat and water deserves increased attention in future modeling approaches, and coupling the current model with this physical process of heat transfer could be an important step toward better simulation in the next generation permafrost models. We projected the possible fate of permafrost over Xidatan till 2100 under an area-average warming rate scenario of QTP. The anticipated permafrost degradation in this study, may not be the basic overview, as it does not consider the regional-level or small-scale-based future climate change. We believe that our simulation results can provide a relatively reasonable projection of permafrost degradation magnitude on the QTP under the different climate change scenarios in the foreseeable future. Meanwhile, high-resolution climate models and improved numerical representations of atmospheric circulation systems and land-atmosphere interactions over the heterogeneous QTP region could be crucial in improving the climate model performance, which will improve the accuracy in the projection of permafrost degradation in the future.

5 Conclusions

This study applied a new transient numerical permafrost model to simulate permafrost distribution and its thermal dynamics at 1 km×1 km resolutions near the northern limit of permafrost on the QTP for current (1970–2019) and future (2020–2100) climatic conditions. Overall, we simulated vertical ground temperature profiles and ALT closely matching the long-term continuous

field observations over the study area. Our model well-describe permafrost heat transfer process and reasonably captures heat attenuation and time lag in deep permafrost. We accurately identified
680 permafrost boundaries, and can realistically capture the evolution of the permafrost thermal regime. According to the simulations, permafrost distribution and its thermal conditions over the study area were controlled by elevational with a strong influence of slope aspects. From 1970 to 2019, the lowest elevation of permafrost (north-facing slope aspect) rose approximately 47 m, and the northern boundary of discontinuous permafrost retreated southwards, approximately 1~2 km. But
685 that remains unchanged for the continuous permafrost area. The regional average MAGT warmed by 0.44 °C, and 0.49 °C on continuous and discontinuous permafrost zones. In general, over the past 50 years, the continuous permafrost zone over the study area has gradually warmed, whereas the discontinuous permafrost zone has remarkably upward thawing, and the margin of the discontinuous permafrost zone reduced by about 12.86%. Under gradual warming climate scenarios, the MAGT
690 is anticipated to rise further, and the warming rate is projected to be slighter higher under SSP than RCP. There are no distinct discrepancies in projection changes in the areal extent of permafrost among SSP and RCP scenarios. These findings highlight the slow process and delays in the response of permafrost in QTP to a warming climate. The projected rate of change in the permafrost extent is far lower than those that do not account for the effects of water phase change, historical climate
695 change, and the thermal state of deep permafrost. In summary, our study provides improved simulations for permafrost distribution and thermal regime dynamics in marginal permafrost on the QTP at decadal to centennial time scales. More importantly, these results may give a better understanding of degradation processes and mechanisms of marginal permafrost on the QTP, and it is fundamental prerequisites for guidelines for the further accurate evaluation of changes in the areal extent of the permafrost on a hinterland the QTP or global scale. Thus, supporting policy-makers
700 and researchers to develop strategies for the cold regions in environmental management, hazard mitigation, adaptation, stability of engineering foundations design, and conservation of land and water resources.

Appendix A

705 **Table A1. A list of monitoring boreholes in the study area and a summary of the ground properties are shown.**

Borehole (altitude/ m a.s.l.)	Coordinat es	Sensor depths (m)	Frozen ground type	Soil stratigraphy
QT09 (4538)	35°43'02" 94°07'05"	0.5–5 m (0.5 m intervals) 5–20 m (1 m intervals) 20–30 m (2 m intervals)	Permafrost	Loam (0–0.2 m) Sandy loam (0.2–1.4 m) Sandy loam with gravel (1.4–2.4 m) Sandy with gravel (2.4–10 m) Rock (10–21 m) Sandy loam (0–1.2 m) Sand (1.3–3 m) Sand with gravel (3–10 m) Weathered mudstone (>10 m)
TB01 (4530)	35°43'00" 94°04'09"	Same as QT09	Permafrost	
XD1–1 (4379)	35°41'55" 94°12'05"	0.5–10 m (0.5 m intervals) 10–15 m (1 m intervals)	Permafrost	Sandy cobble (0–4.5 m) Fluvial sand (4.5–15 m)
XD1–2 (4377)	35°41'59" 94°12'07"	Same as XD1–1	Permafrost	Sandy cobble (0–4.5 m) Fluvial sand (4.5–15 m)
XD1–3 (4576)	35°42'04" 94°12'07"	Same as XD1–1	Permafrost	Sandy cobble (0–5 m) Fluvial sand (5–15 m)
XD1–4 (4374)	35°42'10" 94°12'07"	Same as XD1–1	Permafrost	Sandy cobble (0–5.5 m) Fluvial sand (5.5–15 m)
XD1–5 (4370)	35°42'16" 94°12'08"	0.5–8 m (0.5 m intervals)	Permafrost	Sandy cobble (0–5.5 m) Fluvial sand (5.5–10 m)
XD1–6 (4368)	35°42'24" 94°12'09"	Same as XD1–5	Seasonally frozen ground	Sandy cobble (0–4.5 m) Fluvial sand (4.5–8 m) Sand (0–2.5 m)
XD2–1 (4508)	35°41'56" 94°05'08"	Same as XD1–1	Permafrost	Sand with massive ground ice (2.5– 7 m) Clay (7–9 m) Weathered mudstone (9–15 m) Sand (0–2.8 m)
XD2–2 (4503)	35°42'01" 94°05'09"	Same as XD1–1	Permafrost	Sand with massive ground ice (2.5– 6 m) Weathered mudstone (6–15 m)
XD2–3 (4500)	35°42'10" 94°05'09"	Same as XD1–1	Permafrost	Sand cobble (0–4 m) Fluvial sand (4–15 m)
XD2–4 (4498)	35°42'18" 94°05'09"	Same as XD1–1	Seasonally frozen ground	Sandy cobble (0–4 m) Fluvial sand (4–15 m)
XD2–5 (4493)	35°42'26" 94°05'10"	Same as XD1–1	Seasonally frozen ground	Sandy cobble (0–4 m) Fluvial sand (4–15 m)
XD2–6 (4490)	35°42'36" 94°05'11"	Same as XD1–1	Permafrost	Sandy cobble (0–4 m) Fluvial sand (4–15 m)
XD2–7 (4492)	35°43'00" 94°05'05"	Same as XD1–5	Seasonally frozen ground	Sand (0–4.5 m) Sandstone (4–8 m)
JXG (4530)	35°43'12" 94°04'01	1–10 m (1 m intervals) 10–30 m (2 m intervals) 0.4 m 1.6 m	Permafrost	--
CRSQTP (4530)	35°43' 94°05'	4–10 m (2 m intervals) 10–18 m (4 m intervals) 18–20 m (2 m intervals) 20–29 m (3 m intervals)	Permafrost	--
XD1 (4427)	35°43'12" 94°08'24"	--	Permafrost	--
XD2 (4530)	35°43'12" 94°04'14"	--	Permafrost	--
XD3 (4480)	35°43'12" 94°05'24"	--	Permafrost	--


XD4 (4427)	35°42'00" 94°08'24"	--	Permafrost	--
XDT1 (4602)	35°42'36" 94°02'24"	--	Permafrost	--
XDT2 (4530)	35°42'36" 94°05'24"	--	Permafrost	--
CN13 (4448)	35°42'12" 94°07'48"	--	Permafrost	--

Note: The symbol “--” is field-observed frozen ground types collected from previously published literature (Wang et al., 2000; Jin et al., 2000, 2006; Cheng et al., 2007).

Table A2. Calibration thermophysical parameters of different soil layers used for soil temperature modeling

Texture	K (W m ⁻¹ °C ⁻¹)		C (kJ m ⁻³ °C ⁻¹)		VWC (%)
	Frozen	Thawed	Frozen	Thawed	
Loam	1.25–1.57	0.85–1.28	1639–1879	2208–2475	15–20
Clay	0.83–1.30	0.61–1.03	1756–1907	1881–2191	15–20
Sandy loam	1.31–1.93	1.17–1.71	1844–2107	2258–2634	10–20
Loamy sand	1.02–1.38	1.11–1.24	2040–2208	2541–2676	15–20
Sand cobble	1.0–1.29	0.89–1.10	1639–1739	2007–2208	13–15
Fluvial sand	1.32–1.60	1.09–1.30	1288–1413	1568–1819	6–10
Sand	1.86–2.15	1.48–1.64	1505–1639	1940–2208.1	10–14
Sandstone	0.94–1.91	0.77–1.47	1317–1459	1493–1777	2–6
Sand with Gravel	1.91–2.20	1.47–1.68	1459–1601	1777–2061	6–10
Weathered mudstone	2.27	1.71	1543	1881	6
Rock	0.33	0.33	1940	1940	2

Note: K is the thermal conductivity; C is the volumetric heat capacity; as well as VWC represents total volumetric water/ice content. Soil texture information was collected from Luo et al. (2018) and Liu et al. (2020), and the values of thermal conductivity and heat capacity were from the Construction of the Ministry of PRC. (2011) and Yershov. (2016), and fine-adjusted during the calibration, water content was determined by the soil samples of the borehole cores combined with the observation dataset vicinity of QT09 and the ground ice distribution maps from Zhao et al. (2010).



715

Figure A1. Comparison of the simulated (red lines) to observed (blue lines) daily mean ground temperature at 0.5 m, 1.0 m, 3.0 m depth in four calibration boreholes (BT01, XD2-7, QT09, and XD2-6) during the observation period (There were some data gaps due to temperature probe failure in some years, at the BT01, the data gaps in the record mainly occurred at 0.5-15 m in 2007-2008, and at 15-30 m during 2005-2007 and 720 2011-2018, at the QT09, observations at 15-30 m of 2006-2008, 2011-2013, and 2015-2018 are not available, at the XD2-6, the data gap in the record in 2016-2017).





Figure A2. Same as Figure 2, but for the daily mean ground temperature at 8 m, 15 m, and 30 m.



725 **Figure A3. Comparison of the simulated (red lines) to observed (blue lines) daily mean ground temperature**
 at 0.5 m, 1.0 m, and 3.0 m depth in four validation boreholes (XD2-1, XD2-4, XD1-1, and XD1-4) during the
 observation period from 2013 to 2018 (There were some data gaps due to temperature probe failure in some
 years, at the XD2-1, the data gaps in the record mainly occurred at 0.5-3.0 m in the first half of 2015, at the
 XD1-1, the data gap in the record at 0.5-3.0 in 2014-2015, at 8-15 m during 2013-2015, at the XD1-4, the data
 730 gap in the record in the first half of 2015).




Figure A4. Same as Figure 4, but for daily mean ground temperature at 8 m, and 15 m

Code and data availability. Monitoring data *in-situ* from the field observation sites provided by the Cryosphere Research Station on Qinghai-Xizang Plateau of the Chinese Academy of Sciences (CAS), is available online <https://data.tpdc.ac.cn/en/disallow/789e838e-16ac-4539-bb7e-906217305a1d/> (Zhao et al., 2021), and <https://doi.org/10.1007/s11629-017-4731-2>, (Luo et al., 2018), respectively. Improved MODIS LST data were provided by (Zou et al., 2017) (<https://doi.org/10.5194/tc-11-2527-2017>). Meteorological observation from the Chinese

Meteorological Administration (CMA) are available from the China Meteorological Data Sharing
740 Service System (<http://data.cma.cn/data/cdcdetail/dataCode/A.0012.0001.html>). Climate
projections of CMIP5 and CMIP6 data are freely available online at (<https://interactive-atlas.ipcc.ch>)
(Iturbide et al., 2020). The Shuttle Radar Topography Mission (SRTM) with a 1-arcsecond (~30 m)
DEM data were from Hole-filled seamless SRTM data V4, International Centre for Tropical
Agriculture (CIAT), available at: <http://srtm.csi.cgiar.org> (Jarvis et al., 2008). Integration dataset of
745 Tibet boundary was provided by National Tibetan Plateau Data Center (Zhang et al., 2019), are
freely available online at (<http://data.tpdc.ac.cn/zh-hans/>). Three existing permafrost distribution
maps investigated in 1975, 2012, and 2016 were available by Nan et al. (2003)
(<https://doi.org/10.1007/s11629-017-4731-2>), Luo et al. (2018) (<https://doi.org/10.1007/s11629-017-4731-2>), and Zou et al. (2017) (<https://doi.org/10.5194/tc-11-2527-2017>). The new permafrost
750 model source code is available on request from the first authors or corresponding authors or co-
authors of this study: Jianting Zhao, jt.zhao@nuist.edu.cn, Lin Zhao, lzhao@nuist.edu.cn, Zhe Sun,
sunzhe@lzb.ac.cn.

Author contribution. LZ conceived and conceptualized the idea; JZ and ZS developed the
methodology; LZ, ZS, and GH supervised the study; JZ, MX, LY, and SW performed data
755 processing and analyses. LZ and FN acquired the funding and provided the resources; FN, DZ, GL,
DE, CW, YQ, JS, and HZ participated in the fieldwork and maintained the observation sites; JZ
wrote the draft version, and ZS, FN, GH, LW, YZ, JG, YW, YL, WY, and ZX reviewed and edited
the writing.

Competing interests. The author has declared that neither they nor their co-authors have any conflict
760 of interest.

Acknowledgments. A warm thanks to all the scientists, engineers, and students who participated in
the field measurement and helped to maintain the observation network for data acquisition. We
would like to express our gratitude to Dr. Waheed Ullah, Reading Academy, Nanjing University of
Information Science & Technology, for Language editing.

765 *Financial support.* This study was jointly supported by the National Natural Science Foundation of

China (grant no. 41931180) and the Second Tibet Plateau Scientific Expedition and Research (STEP) program (grant no. 2019QZKK0201, 2019QZKK0905).

References

770 Bense, V., Kooi, H., Ferguson, G., and Read, T.: Permafrost degradation as a control on hydrogeological regime shifts in a warming climate, *J Geophys Res Earth Surf* 117, <https://doi.org/10.1029/2011JF002143>, 2012.

Buteau, S., Fortier, R., Delisle, G., and Allard, M.: Numerical simulation of the impacts of climate warming on a permafrost mound, *Permafrost and Periglac. Process.*, 15, 41-57, <https://doi.org/10.1002/ppp.474>, 2004.

775 Cao, Y., Sheng, Y., Wu, J., Li, J., Ning, Z., Hu, X., Feng, Z., and Wang, S.: Influence of upper boundary conditions on simulated ground temperature field in permafrost regions, *J. Glaciol Geocryol.*, 36, 802-810, doi: 10.7522 / j. issn.1000-0240.2014.0096, 2014.

Chang, Y., Lyu, S., Luo, S., Li, Z., Fang, X., Chen, B., Chen, S., Li, R., and Chen, S.: Estimation of permafrost on the Tibetan Plateau under current and future climate conditions using the CMIP5 data, *Int. J. Climatol.*, 38: 5659–5676. <https://doi.org/10.1002/joc.5770>, 2018.

780 Cheng, G. and Jin, H.: Permafrost and groundwater on the Qinghai Tibet Plateau and in Northeast China, *Hydrogeol. J.*, 21, 5–23, <https://doi.org/10.1007/s10040-012-0927-2>, 2013.

Cheng, G. and Wu, T.: Responses of permafrost to climate change and their environmental significance, *Qinghai-Tibet Plateau, J. Geophys. Res.*, 112, 1–10, <https://doi.org/10.1029/2006JF000631>, 2007.

785 Cheng, G. D.: Problems on zonation of high-altitude permafrost, *Ac. Geogr. Sin.*, 39, 185–193, 1984.

Cheng, G.: Influences of local factors on permafrost occurrence and their implications for Qinghai-Xizang Railway design, *Sci China Ser D: Earth Sci.*, 47, 704–709, <https://doi.org/10.1007/BF02893300>, 2004.

790 Construction Ministry of PRC.: *Code for design of soil and foundation of building in frozen soil region* (in Chinese), China Architecture and Building Press, Beijing, China, 2011.

Ehlers, T., Chen, D., Appel, E., Bolch, T., Chen, F., Diekmann, B., Dippold, M., Giese, M., Guggenberger, G., Lai, H., Li, X., Liu, J., Liu, Y., Ma, Y., Miehe, G., Mosbrugger, V., Mulch, A., Piao, S., Schwalb, A., Thompson, L., Su, Z., Sun, H., Yao, T., Yang, X., Yang, K., and Zhu L.:
795 Past, present, and future geo-biosphere interactions on the Tibetan Plateau and implications for permafrost, *Earth Science Reviews*, <https://doi.org/10.1016/j.earscirev.2022.104197>, 2022.

Fewster, R., Morris, P., Ivanovic, R., Swindles, G., Peregon, A., and Smith, C.: Imminent loss of climate space for permafrost peatlands in Europe and Western Siberia, *Nat. Clim. Chang*, 12,

373–379, <https://doi.org/10.1038/s41558-022-01296-7>, 2022.

800 Guo, D., and Wang, H.: CMIP5 permafrost degradation projection: a comparison among different regions. *J. Geophys. Res. Atmos.*, 121, 4499–4517, <https://doi.org/10.1002/2015JD024108>, 2016.

Guo, D., Wang, H., and Li, D. 2012. A projection of permafrost degradation on the Tibetan Plateau during the 21st century. 117, D05106. *Journal of Geophysical Research: Atmospheres*.
805 <https://doi.org/10.1029/2011JD016545>.

Hanna Lee, H., Swenson, S., Slater, A., and Lawrence, D.: Effects of excess ground ice on projections of permafrost in a warming climate, *Environ. Res. Lett.* 9 124006, doi:10.1088/1748-9326/9/12/124006, 2014.

810 Hipp, T., Etzelmüller, B., Farbrot, H., Schuler, T., and Westermann, S.: Modelling borehole temperatures in Southern Norway—insights into permafrost dynamics during the 20th and 21st century, *The Cryosphere*, 6, 553–571, <https://doi.org/10.5194/tc-6-553-2012>, 2012.

Hjort, J., Streletskiy, D., Doré, G., Wu, Q., Bjella, K., and Luoto, M.: Impacts of permafrost degradation on infrastructure, *Nat. Rev. Earth Environ.* 3, 24–38, <https://doi.org/10.1038/s43017-021-00247-8>, 2022.

815 Hu, G., Zhao, L., Wu, X., Li, R., Wu, T., Xie, C., Pang, Q., and Zou, D.: Comparison of the thermal conductivity parameterizations for a freeze-thaw algorithm with a multi-layered soil in permafrost regions, *Catena*, 156, 244–251, <http://dx.doi.org/10.1016/j.catena.2017.04.011>, 2017.

IPCC. Climate change 2021: the physical science basis, 820 https://www.ipcc.ch/report/ar6/wg1/downloads/report/IPCC_AR6_WGI_Full_Report.pdf, 2021.

IPCC.: Special report on the ocean and cryosphere in a changing climate, <https://archive.ipcc.ch/srocc/>, 2019.

825 Iturbide, M., Gutiérrez, J., Alves, L., Bedia, J., Cerezo-Mota, R., Cimadevilla, E., Cofiño, A., Di, L., Faria, S., Gorodetskaya, I., Hauser, M., Herrera, S., Hennessy, K., Hewitt, H., Jones, R., Krakovska, S., Manzanas, R., Martínez-Castro, D., Narisma, G., Nurhati, I., Pinto, I., Seneviratne, S., van den Hurk, B., and Vera, C.: An update of IPCC climate reference regions for subcontinental analysis of climate model data: definition and aggregated datasets, *Earth Syst. Sci. Data*, 12, 2959–2970, <https://doi.org/10.5194/essd-12-2959-2020>.

830 Jafarov, E., Marchenko, S., and Romanovsky, V.: Numerical modeling of permafrost dynamics in Alaska using a high spatial resolution dataset., *The Cryosphere*, 6, 613–624, <https://tc.copernicus.org/articles/6/613/2012/>, 2012.

Jarvis, A., Reuter, H., Nelson, A., and Edith, G.: Hole-filled seamless SRTM data V4, Tech. rep., International Centre for Tropical Agriculture (CIAT), Cali, Columbia, available at: <http://srtm.csi.cgiar.org>, 2008.

835 Jin, H., Li, S., Cheng, G., Wang, S., and Li, X.: Permafrost and climatic change in China, *Global Planet. Change.*, 26, 387-404, [https://doi.org/10.1016/S0921-8181\(00\)00051-5](https://doi.org/10.1016/S0921-8181(00)00051-5), 2000.

Jin, H., Luo, D., Wang, S., Lü, L., and Wu, J.: Spatiotemporal variability of permafrost degradation on the Qinghai-Tibet Plateau, *Sci. Cold Arid Reg.*, 3, 281–305, DOI: 10.3724/SP.J.1226.2011.00281, 2011.

840 Jin, H., Wu, Q., and Romanovsky, V.: Degrading permafrost and its impacts, *Adv. Clim. Change Res.*, 12, <https://doi.org/10.1016/j.accre.2021.01.007>, 2021.

Jin, H., Zhao, L., Wang, S., and Jin, R.: Thermal regimes and degradation modes of permafrost along the Qinghai-Tibet Highway, *Sci. China Ser. D: Earth Sci.*, 49, 1170–1183, <https://doi.org/10.1007/s11430-006-2003-z>, 2006.

845 Kane, D., Hinkel, K., Goering, D., Hinzman, L., and Outcalt, S.: Non-conductive heat transfer associated with frozen soils, *Global Planet. Change.*, 29, 275-292, [https://doi.org/10.1016/S0921-8181\(01\)00095-9](https://doi.org/10.1016/S0921-8181(01)00095-9), 2001.

Lawrence, D., Slater, A., and Swenson, S: Simulation of Present-Day and Future Permafrost and Seasonally Frozen Ground Conditions in CCSM4, *J. Clim.*, 25, 2207-2225, 850 <https://doi.org/10.1175/JCLI-D-11-00334.1>, 2012.

Li, D., Chen, J., Meng, Q., Liu, D., Fang, J., and Liu, J.: Numeric simulation of permafrost degradation in the eastern Tibetan Plateau, *Permafrost and Periglac. Process.*, 19, 93-99, <https://doi.org/10.1002/ppp.611>, 2008.

855 Li, R., Zhao, L., Ding, Y., Wu, T., Xiao, Y., Du, E., Liu, G., and Qiao, Y.: Temporal and spatial variations of the active layer along the Qinghai-Tibet Highway in a permafrost region, *Chinese Sci. Bull.*, 57, 4609–4616, <https://doi.org/10.1007/s11434-012-5323-8>, 2012

Li, S., Cheng, G., and Guo, D.: The future thermal regime of numerical simulating permafrost on the Qinghai-Xizang (Tibet) Plateau, China, under a warming climate. *Science in China, Ser. D*, 434-441, 1996.

860 Li, W., Zhao, L., Wu, X., Zhao, Y., Fang, H., and Shi, W.: Distribution of soils and landform relationships in the permafrost regions of Qinghai-Xizang (Tibetan) Plateau, *Chinese Sci. Bull.*, 60, 2216–2226, <https://doi.org/10.1360/N972014-01206>, 2015b.

Li, X., Cheng, G., Jin, H., Kang, E., Che, T., Jin, R., Wu, L., Nan, Z., Wang, J., and Shen, Y.: Cryospheric Change in China, *Global Planet. Change.*, 62, 210–218, 865 <https://doi.org/10.1016/j.gloplacha.2008.02.001>, 2008.

Liu, G., Xie, C., Zhao, L., Xiao, Y., Wu, T., Wang, W., and Liu, W.: Permafrost warming near the northern limit of permafrost on the Qinghai-Tibetan Plateau during the period from 2005 to 2017, A case study in the Xidatan area, *Permafrost and Periglac. Process.*, 32: 323– 334, <https://doi.org/10.1002/ppp.2089>, 2020;

870 Lu, Q., Zhao, D. and Wu, S.: Simulated responses of permafrost distribution to climate change on the Qinghai–Tibet Plateau. *Sci. Rep.*, 7, 3845. <https://doi.org/10.1038/s41598-017-04140-7>, 2017.

875 Luo, J., Niu, F., Lin, Z., Liu, M., and Yin, G.: Variations in the northern permafrost boundary over the last four decades in the Xidatan region, Qinghai–Tibet Plateau. *J. Mt. Sci.* 15, 765–778, <https://doi.org/10.1007/s11629-017-4731-2>, 2018.

Miner, K., Turetsky, M., Malina, E., Bartsch, A., Tamminen, J., McGuire, A., Fix, A., Sweeney, C., Elder, C., and Miller, C.: Permafrost carbon emissions in a changing Arctic, *Nat. Rev. Earth Environ.* 3, 55–67, <https://doi.org/10.1038/s43017-021-00230-3>, 2022.

880 Nan, Z., Gao, Z., Li, S., and Wu, T.: Permafrost changes in the northern limit of permafrost on the Qinghai-Tibet plateau in the last 30 years, *J. Geogr. Sci.*, 58, 817-823, 2003.

Ni, J., Wu, T., Zhu, X., Hu, G., Zou, D., Wu, X., Li, R., Xie, C., Qiao, Y., Pang, Q., Hao, J., and Yang, C.: Simulation of the present and future projection of permafrost on the Qinghai-Tibet Plateau with statistical and machine learning models, *J. Geophys. Res. Atmos.*, 126, e2020JD033402, <https://doi.org/10.1029/2020JD033402>, 2021.

885 Nitze, I., Grosse, G., Jones, B. M., Romanovsky, V. E., and Boike, J.: Remote sensing quantifies widespread abundance of permafrost region disturbances across the Arctic and Subarctic, *Nat. Commun.* 9, 5423, <https://doi.org/10.1038/s41467-018-07663-3>, 2018.

890 Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H., Dashtseren, A., Delaloye, R., Elberling, B., Etzelmüller, B., Kholodov, A., Khomutov, A., Kääb, A., Leibman, M., Lewkowicz, A., Panda, S., Romanovsky, V., Way, R., Westergaard-Nielsen, A., Wu, T., Yamkhin, J., and Zou, D.: Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale, *Earth-Sci. Rev.*, 193, 299-316, <https://doi.org/10.1016/j.earscirev.2019.04.023>, 2019.

Qin, D. *Glossary of cryosphere science*. Meteorological Press (in Chinses), Beijing, China, 2014.

895 Ran, Y., Li, X., Cheng, G., Zhang, T., Wu, Q., Jin, H., and Jin, R.: Distribution of permafrost in China: an overview of existing permafrost maps, *Permafrost and Periglac. Process.*, 23: 322-333. <https://doi.org/10.1002/ppp.1756>, 2012.

900 Riseborough, D., Shiklomanov, N., Etzelmüller B, Gruber, S., and Marchenko, S.: Recent advances in permafrost modelling, *Permafrost and Periglac. Process.*, 19, 137–156, <https://doi.org/10.1002/ppp.615>, 2008.

905 Schädel, C., Bader, M., Schuur, E., Biasi, C., Bracho, R., Čapek, P., Baets, S., Baets, S., Diáková, K., Ernakovich, J., Aragones, C., Graham, D., Hartley, I., Iversen, C., Kane, E., Knoblauch, C., Lupascu, M., Martikainen, P., Natali, S., Norby, R., O'Donnell, J., Chowdhury, T., Šantrůčková, H., Shaver, G., Sloan, V., Treat, C., Turetsky, M., Waldrop, M., and Wickland, K.: Potential carbon emissions dominated by carbon dioxide from thawed permafrost soils, *Nature Clim. Change*, 6, 950–953, <https://doi.org/10.1038/nclimate3054>, 2016.

Schiesser, W.: The Numerical Method of Lines: Integration of Partial Differential Equations, vol. 212, Academic Press, San Diego, USA, 1991.

Schuur, E., and Abbott, B.: High risk of permafrost thaw. *Nature*, 480, 32–33, 910 <https://doi.org/10.1038/480032a>, 2011.

Sjöberg, Y., Coon, E., Sannel, A., Pannetier, R., Harp, D., Frampton, A., Painter, S., and Lyon SW.: Thermal effects of groundwater flow through subarctic fens: a case study based on field observations and numerical modeling, *Water Resour Res* 52:1591–1606. <https://doi.org/10.1002/2015WR017571>, 2016.

915 Smith, S., O'Neill, H., Isaksen, K., Noetzli, J., and Romanovsky, V.: The changing thermal state of permafrost, *Nat. Rev. Earth Environ.*, 3, 10–23 <https://doi.org/10.1038/s43017-021-00240-1>, 2022.

920 Sun, Z., Zhao, L., Hu, G., Qiao, Y., Du, E., Zou, D., and Xie, C.: Modeling permafrost changes on the Qinghai-Tibetan plateau from 1966 to 2100: a case study from two boreholes along the Qinghai-Tibet engineering corridor. *Permafrost and Periglac. Process.*, 32:156-171, <https://doi.org/10.1002/ppp.2022>, 2019.

925 Sun, Z., Zhao, L., Hu, G., Zhou, H., Liu, S., Qiao, Y., Du, E., Zou, D., and Xie, C.: Numerical simulation of thaw settlement and permafrost changes at three sites along the Qinghai-Tibet Engineering Corridor in a warming climate, *Geophysical Research Letters*, 49, e2021GL097334, <https://doi.org/10.1029/2021GL097334>, 2022.

Wang, C., Wang, Z., Kong, Y., Zhang, F., Yang, K., and Zhang, T.: Most of the Northern Hemisphere Permafrost Remains under Climate Change, *Sci. Rep.*, 9, 3295, <https://doi.org/10.1038/s41598-019-39942-4>, 2019.

930 Wang, S., Jin, H., Li, S., and Zhao, L.: Permafrost degradation on the Qinghai-Xizang (Tibet) Plateau and its environmental impacts, *Permafrost and Periglac. Process.*, 11: 43-53, [https://doi.org/10.1002/\(SICI\)1099-1530\(200001/03\)11:1<43::AID-PPP332>3.0.CO;2-H](https://doi.org/10.1002/(SICI)1099-1530(200001/03)11:1<43::AID-PPP332>3.0.CO;2-H), 2000.

935 Westermann, S., Langer, M., Boike, J., Heikenfeld, M., Peter, M., Etzelmüller, B., and Krinner, G.: Simulating the thermal regime and thaw processes of ice-rich permafrost ground with the land-surface model CryoGrid 3, *Geosci. Model Dev.*, 9, 523–546, <https://doi.org/10.5194/gmd-9-523-2016>, 2016.

Westermann, S., Schuler, T. V., Gisnås, K., and Etzelmüller, B.: Transient thermal modeling of permafrost conditions in Southern Norway, *The Cryosphere*, 7, 719-739, <https://tc.copernicus.org/articles/7/719/2013/>, 2013.

940 Willmott, C., and Matsuura, K.: Advantages of the mean absolute error (MAE) over the root mean square error (RMSE) in assessing average model performance, *Clim. Res.*, 30, 79–82, doi:10.3354/cr030079, 2005.

Wu, J., Sheng, Y., Wu, Q., and Wen, Z.: Processes and modes of permafrost degradation on the Qinghai-Tibet Plateau, *Sci. China Ser. D-Earth Sci.*, 53, 150–158, <https://doi.org/10.1007/s11430-009-0198-5>, 2010.

945

Wu, Q. and Zhang, T.: Recent permafrost warming on the Qinghai-Tibetan Plateau, *J. Geophys. Res.*, 113, D13108, doi:10.1029/2007JD009539, 2008.

Wu, Q., Zhang, T., and Liu, Y.: Permafrost temperatures and thickness on the Qinghai-Tibet Plateau, *Global Planet. Change.*, 72, 32-38, <https://doi.org/10.1016/j.gloplacha.2010.03.001>, 2010.

950

Wu, T., Li, S., Cheng, G., and Nan, Z.: Using ground-penetrating radar to detect permafrost degradation in the northern limit of permafrost on the Tibetan plateau, *Cold Reg. Sci. Technol.*, 41, 211-219, <https://doi.org/10.1016/j.coldregions.2004.10.006>, 2005.

Wu, X., Nan, Z., Zhao, S., Zhao, L., and Cheng, G.: Spatial modeling of permafrost distribution and properties on the Qinghai-Tibet Plateau, *Permafrost and Periglac. Process.*, 29, 86-99, <https://doi.org/10.1002/ppp.1971>, 2018.

955

Xiao, Y., Zhao, L., Dai, Y., Li, R., Pang, Q., and Yao, J.: Representing permafrost properties in CoLM for the Qinghai-Xizang (Tibetan) plateau, *Cold Reg. Sci. Technol.*, 87, 68-77, <http://dx.doi.org/10.1016/j.coldregions.2012.12.004>, 2013.

960

Xu, Y., Shen, Y., and Wu, Z.: Spatial and Temporal Variations of Land Surface Temperature Over the Tibetan Plateau Based on Harmonic Analysis, *Mt. Res. Dev.*, 33, 85–94, <https://doi.org/10.1659/MRD-JOURNAL-D-12-00090.1>, 2013.

Yang, M., Nelson, F. E., Shiklomanov, N. I., Guo, D., and Wan, G.: Permafrost degradation and its environmental effects on the Tibetan Plateau: A review of recent research, *Earth-Sci. Rev.*, 103, 31–44, 2010.

965

970

Yao, T., Xue, Y., Chen, D., Chen, F., Thompson, L., Cui, P., Koike, T., Lau, W. K., Lettenmaier, D., Mosbrugger, V., Zhang, R., Xu, B., Dozier, J., Gillespie, T., Gu, Y., Kang, S., Piao, S., Sugimoto, S., Ueno, K., Wang, L., Wang, W., Zhang, F., Sheng, Y., Guo, W., Yang, X., Ma, Y., Shen, S. S. P., Su, Z., Chen, F., Liang, S., Liu, Y., Singh, V. P., Yang, K., Yang, D., Zhao, X., Qian, Y., Zhang, Y., and Li, Q.: Recent Third Pole's Rapid Warming Accompanies Cryospheric Melt and Water Cycle Intensification and Interactions between Monsoon and Environment: Multidisciplinary Approach with Observations, Modeling, and Analysis, *B. Am. Meteorol. Soc.*, 100, 423-444, <https://doi.org/10.1175/BAMS-D-17-0057.1>, 2019.

Yershov, E.: *Principles of Geocryology*, Lanzhou University Press (in Chinse), Lanzhou, China, 2016.

975

Yi, S., Wang, X., Qin, Y., Xiang, B., and Ding, Y.: Responses of alpine grassland on Qinghai-Tibetan plateau to climate warming and permafrost degradation: a modeling perspective, *Environ. Res. Lett.*, 9, 074014, <https://doi.org/10.1088/1748-9326/9/7/074014>, 2014.

Yin, G., Luo, J., Niu, F., Lin, Z., and Liu, M.: Thermal regime and variations in the island permafrost
980 near the northern permafrost boundary in Xidatan, Qinghai–Tibet Plateau, *Front. Earth Sci.*, 560,
<https://doi.org/10.3389/feart.2021.708630>, 2021.

Yue, G., Zhao, L., Zhao, Y., Du, E., Wang, Q., Wang, Z., and Qiao, Y.: Relationship between soil
properties in permafrost active layer and surface vegetation in Xidatan on the Qinghai-Tibetan
Plateau, *J. Geogr. Sci.*, 35, 565-573, doi: 10.7522/j. issn. 1000-0240.2013.0065, 2013.

985 Zhang, G., Nan, Z., Hu, N., et al. 2022. Qinghai-Tibet Plateau permafrost at risk in the late 21st
Century. *Earth's Future*. 10, e2022EF002652. <https://doi.org/10.1029/2022EF002652>.

Zhang, Y., Wang, X., Fraser, R., Olthof, I., Chen, W., McLennan, D., Ponomarenko, and S., Wu, W.:
Modelling and mapping climate change impacts on permafrost at high spatial resolution for an
Arctic region with complex terrain, *The Cryosphere*, 7, 1121–1137, <https://doi.org/10.5194/tc-7-1121-2013>, 2013.

990 Zhang, Y.: Integration dataset of Tibet Plateau boundary. National Tibetan Plateau Data Center, DOI:
10.11888/Geogra.tpdc.270099. CSTR: 18406.11. Geogra.tpdc.270099, 2019.

Zhao L, Ding Y, Liu G, Wang S, and Jin H.: Estimates of the reserves of ground ice in permafrost
regions on the Tibetan plateau, *J. Glaciol Geocryol.*, 32:1-9, 2010.

995 Zhao, L. and Sheng, Y.: Permafrost survey manual (in Chinses), Science Press, Beijing, 13–14, 2015.

Zhao, L., Hu, G., Zou, D., Wu, X., Ma, L., Sun, Z., Yuan, L., Zhou, H., and Liu, S.: Permafrost
Changes and Its Effects on Hydrological Processes on Qinghai-Tibet Plateau, *Bull. Chin. Acad.
Sci.*, 34, 1233–1246, DOI: 10.16418/j.issn.1000-3045.2019.11.006, 2019.

1000 Zhao, L., Wu, Q., Marchenko, S. S., and Sharkhuu, N.: Thermal state of permafrost and active layer
in Central Asia during the international polar year, *Permafrost and Periglac. Process.*, 21, 198–
207, <https://doi.org/10.1002/ppp.688>, 2010.

1005 Zhao, L., Zou, D., Hu, G., Du, E., Pang, Q., Xiao, Y., Li, R., Sheng, Y., Wu, X., Sun, Z., Wang, L.,
Wang, C., Ma, L., Zhou, H., and Liu, S.: Changing climate and the permafrost environment on
the Qinghai–Tibet (Xizang) Plateau, *Permafrost and Periglac.*, 31, 396–405,
<https://doi.org/10.1002/ppp.2056>, 2020.

Zhao, L., Zou, D., Hu, G., Wu, T., Du, E., Liu, G., Xiao, Y., Li, R., Pang, Q., Qiao, Y., Wu, X., Sun,
Z., Xing, Z., Sheng, Y., Zhao, Y., Shi, J., Xie, C., Wang, L., Wang, C., and Cheng, G.: A synthesis
dataset of permafrost thermal state for the Qinghai–Tibet (Xizang) Plateau, China, *Earth Syst.
Sci. Data*, 13, 4207–4218, <https://doi.org/10.5194/essd-13-4207-2021>, 2021.

1010 Zhou, Y., Guo, D., Qiu, G., Cheng, G., and Li, S.: China Permafrost, Science Press, Beijing, 145–
151, 2000.

Zou, D., Zhao, L., Sheng, Y., Chen, J., Hu, G., Wu, T., Wu, J., Xie, C., Wu, X., Pang, Q., Wang, W.,
Du, E., Li, W., Liu, G., Li, J., Qin, Y., Qiao, Y., Wang, Z., Shi, J., and Cheng, G.: A new map of

1015 permafrost distribution on the Tibetan Plateau, *The Cryosphere*, 11, 2527–2542,
<https://doi.org/10.5194/tc-11-2527-2017>, 2017.

1020 Zou, D., Zhao, L., Wu, T., Wu, X., Pang, Q., and Wang, Z.: Modeling ground surface temperature
by means of remote sensing data in high-altitude areas: test in the central Tibetan Plateau with
application of moderate-resolution imaging spectroradiometer Terra/Aqua land surface temperature
and ground based infrared radiometer, *J. Appl. Remote Sens.*, 8, 083516,
<https://doi.org/10.1117/1.JRS.8.083516>, 2014.