

Permafrost saline water and Early to Mid-Holocene permafrost aggradation in Svalbard

Dotan Rotem^{1,2}, Vladimir Lyakhovsky³, Hanne Hvidtfeldt Christiansen², Yehudit Harlavan³, Yishai Weinstein¹

¹Department of Geography and Environment, Bar-Ilan University, Ramat-Gan, 52900, Israel.

5 ²Arctic geophysics Department, the University Centre in Svalbard, UNIS, Longyearbyen 9170, Norway.

³Geological Survey of Israel, 32 Yesha'yahu Leibowitz, Jerusalem 9692100, Israel.

Corresponding author: Dotan Rotem (dotanrotem1969@gmail.com)

Abstract. Deglaciation in Svalbard was followed by seawater ingress and deposition of marine (deltaic) sediments in fjord valleys, while elastic rebound resulted in fast land uplift and the exposure of these sediment to the atmosphere, in which the formation of epigenetic permafrost formed. This was then followed by the accumulation of aeolian sediments, with syngenetic permafrost formation. Permafrost was studied in the eastern Adventdalen Valley, Svalbard, 3-4 km from the maximum up-valley reach of post-deglaciation seawater ingress, and its ground ice was analyzed for its chemistry. While ground ice in the syngenetic part is basically fresh, the epigenetic part has a frozen fresh-saline water interface (FSI), with chloride concentrations increasing from the top of the epigenetic part (at 5.5 m depth) to about 15% that of seawater at 11 m depth. We applied a one-dimensional freezing model to examine the rate of top-down permafrost formation, which could accommodate with the observed frozen FSI. The model examined permafrost development under different scenarios of mean average air temperature, water-freezing temperature and the degree of pore-water freezing. We found that even at the relatively high air temperatures of the Early to mid-Holocene, permafrost could aggrade quite fast down to 20 to 37 m (the whole sediment fill of 25 m at this location) within 200 years. This, in turn, allowed freezing and preservation of the fresh-saline water interface despite of the relatively fast rebound rate, which apparently resulted in an increase in topographic gradients toward the sea. The permafrost aggradation rate could also be enhanced due to non-complete pore water freezing. We conclude that freezing must have started immediately after the exposure of the marine sediment to atmospheric conditions.

1. Introduction

30 Cycles of global warming and cooling are well documented in the geological history (e.g., Imbrie et al., 1993; Benn and Evans, 2014; Arnscheidt and Rothman, 2020). During the Pleistocene, these cycles followed Northern Hemisphere glaciation and deglaciation, which influenced both marine and land temperatures (Park et al., 2019). This also affected the extent of cryotic conditions in the periglacial environment (e.g. Murton, 2021), i.e., the distribution of permafrost, which currently covers 22% of the
35 Northern Hemisphere land areas (Obu et al., 2019). While temperatures during the Holocene were significantly higher than during the Last Glacial period, the retreat of glaciers and the follow-up elastic rebound and exposure of new land in the Arctic and the sub-Arctic environment allowed freezing and the aggradation of permafrost (e.g. Landvik et al., 1988). Nevertheless, the relatively high temperatures during the Early and the mid-Holocene warm period raise questions about the timing of initiation and the
40 extent of this process (e.g. Landvik et al., 1988; Humlum, 2005).

In Svalbard (Fig. 1), the fast retreat of glaciers during the end of Late-Pleistocene into the beginning of the Holocene resulted in the ingression of seawater in fjord valleys, which was followed by gradual uplifting and exposure due to elastic rebound. This resulted in epigenetic permafrost aggradation followed by the deposition of fluvial and aeolian sediments, and the formation of syngenetic permafrost
45 during the last ca. 4 ka (Gilbert et al., 2018). In the present study, we use the presence of saline water in the epigenetic permafrost to constrain the timing of freezing.

Permafrost is a soil or rock, which has been below 0°C for at least two consecutive years (French, 2017). While winter freezing of the ground is common in a large extent of land areas, the existence of permafrost and its aggradation depends on the annual energy balance between the atmosphere and the land (Black,
50 1954). Accordingly, permafrost develops when the land heat loss during winter exceeds the gain during the summer for long enough time. This is controlled by both seasonal solar radiation and the soil/rock thermal properties. Heat exchange between soil and the atmosphere is also strongly affected by land cover, whereby permafrost is usually not developed neither under the sea nor beneath warm-based glaciers (Waller et al., 2012). Nevertheless, permafrost can occur beneath lagoons in association with
55 taliks, as well as beneath bottom-fast ice in shallow water (Solomon et al., 2008). The extent and depth of permafrost can be significantly reduced by thick vegetation or snow cover (e.g. Grünberg et al., 2020). During the last glacial cycle, the Barents Sea and the Svalbard area (Fig. 1) were covered by one to three ice caps (Mangerud et al., 2002; Patton et al., 2017). Glacier retreat has been followed since by

elastic rebound, which is well documented in Svalbard (Bondevik et al., 1995; Lønne and Nemeč, 2004; 60 Sessford et al., 2015), with a land rise of up to 130 m in eastern Svalbard and 65 m in the western part of the archipelago (Forman, 2004). In western Svalbard, the focus of this study, research indicates a fast land rise of 19-15 mm y⁻¹ during Early to the mid-Holocene (11.7 – 8.2 ka BP), which decreased to 5 - 4 mm y⁻¹ toward the end of mid-Holocene (Salvigsen, 1984; Sessford et al., 2015) and ca. 1 mm y⁻¹ during the late Holocene (last 4 ka, e.g. Forman et al., 2004).

65 Land uplift and exposure is accompanied by the establishment of a surficial drainage system, as well as the development of a groundwater flow network, which strongly depends on the rate of permafrost deepening (Edmunds et al., 2001). The permeability of frozen soils is greatly reduced (Burt and Williams, 1976; Cochand et al., 2019), such that extensive permafrost prevents penetrating of surface water and recharging of groundwater (McEwen and de Marsily, 1991). While in sporadic and discontinuous 70 permafrost, groundwater flow is possible through non-frozen sections or taliks, flow is practically impossible through continuous permafrost land areas (Lemieux et al., 2008; Walvoord and Kurylyk, 2016), while it may be active in sub-permafrost zones or along faults in relation with pingos (Hornum et al., 2020). Flow may also provide the necessary conditions for the formation of cryopegs which holds overcooled liquid brines (Ahonen 2001). Although cryopegs were identified in Adventdalen (Tavakoli et 75 al., 2021), water paths were not yet described.

According to the Ghyben-Hertzberg approximation (Bear and Dagan 1964; Verruijt 1968), depth from seawater level to the fresh-saline water interface should be about 1:40 of the groundwater head above sea level. This ratio increases with decreasing salinity of saline water. With typical Early to mid-Holocene rebound rates of 15 to 4.5 mm y⁻¹ (Sessford et al., 2015), and assuming that the groundwater table 80 (saturated conditions) followed the topography, the fresh-saline interface is expected to be pushed downwards to as deep as 120 to 36 m respectively, within 200 years after exposure. A groundwater table of 1 m below the surface would result in a delay of 100 to 200 years, but also in this case a sediment section of tens of meters will be completely flushed within several hundred years. This is unless sediment freezing practically halts flow in the subsurface.

85 The objective of this paper is to test the presented hypothesis by studying the ground ice geochemistry of a permafrost core from Adventdalen, Svalbard, and by using a 1-D numerical heat transfer model to simulate permafrost aggradation under various surface temperature conditions and degrees of freezing.

2. Study site


Adventdalen is a U-shaped glacially eroded valley located in western Spitsbergen, Svalbard, centred on 78.110N, 16.180E (Fig. 1a). During the last glacial cycle, the valley was eroded to the basement, which was then covered by glacial deposits (Elverhøi et al., 1995; Gilbert et al., 2018). This was followed by deglaciation, which was completed ca. 10.5 ka BP (Mangerud et al., 1992; Svendsen and Mangerud, 1997; Lønne and Lyså, 2005; Farnsworth et al., 2020). Deglaciation was followed by up-valley seawater ingression, up to 13.5 km from the current end of the fjord (Cable et al., 2018; Lønne and Nemec, 2004) and subsequent valley infilling with sediment and a delta front prograding downvalley. Elastic rebound resulted in the exposure of the eastern part of the valley before 9.5 ka BP, which progressed down-valley, arriving at the current coastline location at about 4 ka (Gilbert et al., 2018). The exposed surface was first covered by fluvial sediments, followed by aeolian deposits during 4-2 ka (Gilbert et al., 2018). Permafrost in Svalbard, both epigenetic and syngenetic, is continuous and is estimated to be >100 m thick in the valleys (Humlum, 2005). The active-layer thickness is commonly 60–100 cm in the valley bottom sediments (Christiansen and Humlum, 2003; Gilbert et al., 2018; Weinstein et al., 2019; Strand et al., 2020).

Meteorological data is measured continuously for more than 100 years at the Svalbard Airport (e.g. Nordli et al., 2020), located on the Adventfjorden coast, ca. 12 km northwest of our study site. Mean annual air temperature (hereafter MAAT) was -5.9°C from 1971 to 2000, although in 2018 it was merely -1.8°C . Mean annual precipitation during 1971-2000 was 196 mm, while in 2018 it was 177 mm (Strand et al., 2021). MAAT and sea surface temperature during Early to the mid-Holocene was $2-4^{\circ}\text{C}$ higher than today, as suggested by marine mollusc's shells (Mangerud and Svendsen, 2018), lacustrine alkenons (van der Bilt et al., 2018), flora DNA (Alsos et al., 2016) and models incorporating physical and biological considerations (e.g. Park et al., 2019). Since the mid-Holocene, a continuous decline in MAAT is recorded, which changed into a fast temperature rise during the last several decades (e.g. Christiansen et al., 2013).

The study site, Adventdalen East (ADE), is located on a river terrace (78.1722°N 16.0613°E), 9.8 km upvalley from the Adventfjorden at 23 m a.s.l (Fig. 1a and b). We drilled at the same S1 location as Gilbert et al., (2018). The permafrost section (valley-fill sediments) at the ADE site (ca. 20 m) consists of a syngenetic part from 1.0- 5.5 m depth, which includes a shallow 1.5 m of fine-grained aeolian deposits and thick ice bodies of segregation ice intruded above fluvial gravel channel fill (Gilbert et al., 2018).

Below 5.5 m the epigenic permafrost consists of 3-4 m of fluvial sediments (mud and pebbles, ice-rich). This is underlain by back-delta, deltaic and fore-delta sediments (5.5-17.5 m), which cover glacial sand deposits (17.5-20 m) (Gilbert et al., 2018). The study site was deglaciated by 11.3 ka and emerged above seawater at 9.2 ka (Gilbert et al., 2018), exposing it to atmospheric conditions then, which allowed the development of a groundwater system on one hand and possibly the aggradation of permafrost on the other hand.

125





130 Figure 1. (a) Study site ADE is located at the Adventdalen, Svalbard, on a river terrace. Map provided with courtesy of Norwegian Polar Institute. (b) Drilling at ADE, spring 2017 (top); core samples before processing at UNIS cold room lab, (bottom). Photos: Dotan Rotem and Yishai Weinstein.

3. Methods

135 Two cores, 0.5 m apart, one 13 m and the other 9 m long, were retrieved at ADE in March 2017, using the UNIS permafrost drill-rig (Gilbert et al., 2015), which has core barrels of 43 mm diameter (ID). Core length, borehole depth, core condition and gravel content were recorded in the field. Retrieved core sections were placed in plastic bags and marked with a serial number and an arrow pointing towards the core top. Cores were stored at -18°C in a freezer at UNIS until processing. Cores were sectioned in a

cold room (-5°C) to 0.5 m depth intervals (Fig 1b). Intervals of the same depth in the two cores were combined to gain enough ground-ice per section for Ra isotope measurements. Samples were first scraped and then crashed to small chips, which were placed in 250 ml centrifuge tubes. Ra-free water (up to 40 ml) was added to some of the tubes, to facilitate the extraction of pore fluid. Samples were then thawed in a microwave set to 600 W for 2 min, followed by centrifuging for 8 min (11,000 RPM, high g) to separate thawed water from the soil. Extracted water was run through 3 μm , followed by 0.45 μm filters. Most of the water was used for Ra isotopes analysis (see Weinstein et al., 2019), while 30-60 ml was used for chemistry analyses. Water of the added Ra-free water was analysed to correct for element concentrations. Major elements were analysed in the Geological Survey of Israel (GSI) by ICP-AES (Optima 3000), where Sc was added as an internal standard, whereas Cl^- and SO_4^{2-} was determined by potentiometer titration, using Metrohm 702 SM Titrino Titrator connected to a chlorine electrode. The error for all majors is considered less than 5%.

4. Ground ice chemistry

Major elements of thawed ground ice are presented in Table 1 and concentration profiles of Cl^- , Na^{2+} and SO_4^{2-} are shown in Fig. 2 a-c. While the salinity of ground ice in the syngenetic permafrost is that of fresh water (e.g. Cl^- : 10-74 mg L^{-1} , Na^{2+} : 10-33 mg L^{-1} and SO_4^{2-} : 9-31 mg L^{-1}), epigenetic permafrost ground ice demonstrates a trend of increasing concentrations down to 9-12 m depth: 440-3600, 80-2700 and 150-740 mg L^{-1} of Cl^- , Na^{2+} and SO_4^{2-} , respectively. Between 9-12 m, concentrations are quite scattered, and the increasing pattern is less clear. While the Cl^- content (Fig. 2a) of the ground ice is no more than 15% seawater salinity, and the salinity of a deeper-seated saline water end member could be significantly higher, the observed increased salinity clearly presents a fresh-saline water interface.

The ionic ratio of Na^{2+} to Cl^- in both epigenetic and the syngenetic permafrost mostly exceeds 1 (Fig. 2d), significantly higher than in seawater (0.86), which is probably the result of sediment dissolution (e.g. of micas), since ion exchange should result in either a conservative behaviour (during freshening, as is the case in the ADE marine section) or in Na^+ depletion (in the case of salinization, e.g. Russak and Sivan, 2010). On the other hand, SO_4/Cl in the epigenetic permafrost is close to that of seawater (Fig. 2f), implying a relative conservative behaviour. Nevertheless, SO_4/Cl in the syngenetic part is very variable (Fig. 2f), reaching ratios as high as 2, which could be due to shale dissolution (Hindshaw et al., 2016; Cable et al., 2018). High concentration of SO_4 was also recorded in sub-permafrost (Pingo) water,

170 which was attributed to gypsum dissolution (Hodson et al., 2020). Ca/Cl decreases with depth in the syngenetic permafrost and is very low (<0.01) in the epigenetic permafrost (Fig. 2e), which is in agreement with freshening experiments in fresh-saline water zones (Russak and Sivan 2010). The high ratio of Ca/Cl and SO₄/Cl at -5.45 m depth is enigmatic and may present paleo active layer - permafrost table zone were major and minor elements concentrate (e.g. Cary and Mayland, 1972; Kokelj et al., 2002) but it should be further studied.


Table 1. Major elements (mg/L) of grownd-ice samples from the ADE core.

Sample name	Depth (m)	Permafrost Type ¹	Cl ⁻	Br	SO ₄ ²⁻	SiO ₂	Na ⁺	K ⁺	Sr	Ca ²⁺	Mg ²⁺	Freezing state of sample ₃
Sea water conc. ₂			19,354	67.3	2,712		10,760	399	7.9	412	1,290	
DR-AD-55	-1.3	Syngenetic	27.1	0.5	19.0	13.0	10.7	17.0	0.2	5.4	15.1	Frozen
DR-AD-58	-2.1	Syngenetic	12.8	0.6	30.7	14.0	19.8	4.1	0.4	8.8	27.8	Frozen
DR-AD-52	-2.9	Syngenetic	74.1	0.7	8.9	9.0	13.6	44.7	0.6	12.7	35.2	Frozen-icy
DR-AD-57	-3.3	Syngenetic	13.9	0.5	27.9	5.9	32.6	4.4	0.1	3.6	7.6	Frozen-icy
DR-AD-56	-3.5	Syngenetic	10.0	0.3	12.3	5.3	13.3	4.5	0.1	1.5	2.6	Frozen-icy
DR-AD-61	-4.0	Syngenetic	14.5	0.7	13.0	11.6	9.5	5.2	0.2	3.6	8.8	Frozen
DR-AD-63	-4.7	Syngenetic	26.8		11.6	30.0	12.8	13.6	0.1	1.7	2.6	Frozen
DR-AD-59	-5.5	Epigenetic	67.4	0.6	146.6	20.9	83.3	16.0	0.5	13.6	27.0	Partly frozen
DR-AD-53	-6.3	Epigenetic	438.1	1.9	116.1	19.4	422.0	6.7	0.5	14.2	6.1	Frozen
DR-AD-64	-7.1	Epigenetic	847.0		369.5	38.1	828.7	22.7	0.1	11.6		Frozen
DR-AD-65	-8.3	Epigenetic	1678.4		465.1	21.2	1296.3	29.1	0.1	0.5		Frozen
DR-AD-67	-8.8	Epigenetic	3677.6		570.4	25.0	1680.1	55.6	0.1	2.2		Partly frozen
DR-AD-54	-9.4	Epigenetic	2145.1	7.9	508.2	9.1	2497.5	194.6	0.3	17.0		Partly frozen
DR-AD-66	-10.4	Epigenetic	3500.4		738.9	24.8	2705.8	42.4	0.4	9.5		Partly frozen
DR-AD-60	-11.8	Epigenetic	2412.6	7.0	408.1	20.5	2319.8	98.0	0.6	18.3		Partly frozen

¹ after Gilbert et al., 2018

175 ² after de Baar et al., 2017, Salinity of 35%

³ Freezing state as recorded in the field



180

Figure 2. Major element concentrations in ground ice from Adventdalen-East (ADE) deep drillhole: (a) Cl⁻, (b) Na⁺ and (c) SO₄²⁻ in mg/L. Figures (d) - (f) present selected equivalent ratios along the profile. Dashed line separates the syngenetic and the epigenetic permafrost (Gilbert et al., 2018).

185 5. Model of permafrost formation

5.1 Conceptual model

To study the rate of permafrost formation, we developed a numerical model that solves the temperature distribution in space and time and freezing front progression, i.e. Stefan solution (e.g., Šarler, 1995). Considering low horizontal temperature variations, the problem was reduced to one dimensional depth-
190 dependent heat transfer with moving internal phase transition boundary. Various analytical and numerical methods have been developed to obtain stable solution of the Stefan problem (e.g., Crank, 1984). However, unlike the simple and clean single component systems, many natural systems, including water saturated porous rocks, change their phases under a specified temperature range rather than isothermally (Lunardini, 1988; Růhaak et al., 2015). In this case, an evolving “mushy zone” emerges,
195 which separates between the solid and liquid regions, where the thawing or freezing begins and proceeds, accompanied by latent heat absorbance or release (e.g., Crank, 1984; Yang et al., 2020). Following this approach, instead of the mathematical boundary, we use a narrow transition mushy zone (shaded area in Fig. 3), with pore space consisting of a mixture of ice and water (Rubinstein, 1982). The local enthalpy in the mushy zone takes values in the range between those of the pure solid and liquid,
200 and the temperature is approximated by a constant value, $T=T_{\text{wit}}$, equal to the phase change temperature (Crank, 1984).

Heat exchange in the sub-surface is controlled by the ground temperature gradient, as well as by the soil thermal properties (i.e. thermal conductivity, Burn, 2011). Above ground, the main factor is the air temperature, which is measured and reported as MAAT (Luo et al., 2018; Szafraniec and Dobiński, 2020)
205 and was taken as representing the MAGST (Mean Annual Ground Surface Temperature). Initial surface temperature was defined according to the temperature of the shallow seawater during mid-Holocene (2°C, Rasmussen et al., 2012), while the initial temperature profile (black solid line in Fig. 3) was defined using the regional average geothermal gradient of $0.033^{\circ}\text{C m}^{-1}$, as discussed by Olausen et al., (2019) and Betlem et al., (2018). The lower boundary of the model was set at 300 m depth, with temperature of
210 12°C according to the initial temperature distribution. Throughout the simulation, we searched for the depth and time-dependent temperature distribution $T(z,t)$, schematically shown as a dashed line in Fig. 3.

Several MAAT values were used for the modelling: (1) the current -5.8°C (measured at the Adventdalen 'Polygons' site, (Christiansen, 2005), 7 km from the fjord; (2) -4°C , which was taken from climate

215 simulation models for the mid-Holocene (Park et al., 2019; see also Mangerud and Svendsen 2017; van
der Bilt et al., 2019); (3) -3°C and 0°C assumed by Humlum (2005) for the mid-Holocene. we present
simulation for -4°C and with 0°C as an extreme high value. While snow may cause differences between
MAAT and MAGST due to thermal insulation (Zhang, 2005), it was found that in western Svalbard,
specifically in the flat landforms at Adventdalen, the differences between MAAT and MAGST are less
220 than 0.5°C (Christiansen, 2005; Lüthi 2010; Etzelmüller et al., 2011; Farnsworth 2013). Therefore, MAAT
was taken as representing MAGST. Amplitude of seasonal temperature oscillation at the surface was
set to 12°C , similar to the current fluctuation (Nordli et al., 2014; Christiansen, 2005; Osuch and
Wawrzyniak, 2017).

To distinguish between a frozen and water saturated cell, we defined a time and depth-dependent
225 freezing ratio, $B(z,t)$, shown by dotted line in Fig. 3. $B=1$ means that the soil is water-saturated, while in
the case of $B=0$ pore space is fully ice-saturated. In the mushy zone (shaded area in Fig. 3), the B -value
changes between 0 and 1, and the rate of its change defines the amount of energy or latent heat
associated with water-ice phase transition. Since it is now well-established that permafrost is not
necessarily fully frozen (e.g. Keating et al., 2018; Oldenborger and LeBlanc, 2018; see Table 1), we also
230 investigated permafrost aggradation under “partial freezing” conditions of 25% and 50%. Note that our
model assumes fully-saturated pore-water conditions, since freezing starts at sea level, soon after
exposure, therefore groundwater level is expected to be at the surface.




Figure 3. Schematic freezing profile during the top-down (epigenetic) freezing process. Initial and the
 235 developing geothermal gradients are also shown; the mushy region with constant temperature is the
 depth zone where phase transition occurs. Dotted line schematically represents the freezing condition,
 where $B=0$ stands for fully frozen and $B=1$ for liquid only. Active layer seasonality is neglected.

Another factor affecting the rate of permafrost formation is the water-freezing temperature (hereafter:
 240 WFT), which varies with salinity (Morgenstern and Anderson, 1973), as well as due to other
 environmental factors (Farouki, 1981; Morgenstern and Anderson 1973). Since salinities in the ADE site,
 down to 12 m, do not exceed 15% that of seawater, fresh water freezing temperature is the most
 appropriate for our simulations. Nevertheless, we also conducted simulations with WFT of -2°C , which
 is close to that of seawater ($T_m=-1.9^{\circ}\text{C}$, Bodnar, 1993; Marion et al., 1999), as well as with -4 , -5 and $-$
 245 6°C , following the reports of Gilbert et al., (2019) and Tavakoli et al., (2021) about high salinities (up to
 73 ppt) in cryopegs in Adventdalen.

For simplicity, thermal conductivity was calculated considering three major components, namely water,
 ice and dry sediments. While the thermal conductivity of water and ice are well defined (0.569 and 2.24
 $\text{W m}^{-1} \text{K}^{-1}$, respectively, e.g. Williams and Smith, 1989), the range of conductivity values for dry sediment

250 in the literature is large, from 0.25 for peat to 7.6 W m⁻¹ K⁻¹ for sand depending mainly on quarts content, water content and other factors (Farouki, 1981; Williams and Smith, 1989; Cosenza et al., 2003; Balland and Arp, 2005; Zhang and Wang, 2017; He et al., 2021). We decided to use the value of 3 W m⁻¹ K⁻¹ for the mineral fraction of the sediment (e.g. Overduin et al., 2019).

Porosity is an important factor in the aggradation or thawing of permafrost (Hornum et al., 2020). It
255 determines both the amount of heat released or required during freezing/thawing, and the thermal characteristics of the soil. Porosity of sediments at ADE was mainly set to its average value, 0.3 (Gilbert et al., 2018), although we also tested the effect of other porosity values (Appendix 1, DOI: [10.6084/m9.figshare.23684877](https://doi.org/10.6084/m9.figshare.23684877)). Bedrock (≥ 25 m depth) porosity was taken as 0.1, the value suggested by Hornum et al. (2020), assuming that the bedrock is mainly composed of fractured shales
260 (Benn and Evans, 2014; Grundvåg et al., 2019).

Freezing temperature and degree of freezing were kept uniform in each of the simulations, although as pore water freezes, the remaining fluid becomes saltier, further lowering the freezing temperature of the remaining solute (Herut et al., 1990), such that fully frozen pore space (i.e. 100% freezing) can only be
265 reached at extremely low temperatures, which are not relevant to our study sites, as well as to most other permafrost areas (Homshaw, 1980; Dobinski, 2011).

The 1-D freezing model provides a good approximation of freezing rate and permafrost aggradation as shown by many studies (e.g., Harada and Yoshikawa, 1996; Kukkonen and Šafanda, 2001; Farbrot et al., 2007; Etzelmüller et al., 2011; Hornum et al., 2020). Such models actually provide the maximum rates of freezing propagation, as lateral heat transport by groundwater flow is neglected. Neglecting
270 lateral heat transfer is quite justified, considering that (1) upstream shallow groundwater arrives from areas that were exposed earlier, therefore should not be warmer than the ADE groundwater, (2) experimental data suggest that a temperature drop by a tenth of a centigrade below 0°C reduces the hydraulic conductivity by several orders of magnitude (Burt and Williams, 1976; Rūhaak et al., 2015). Nevertheless, as stressed above, soon after freezing initiates, the hydraulic conductivity dramatically
275 decreases and the impact of lateral flow can be neglected.

5.2 Mathematical formulation

The heat conduction equation for time and depth-dependent temperature profile, $T(z,t)$, is (Crank, 1984):

$$280 \quad \rho C_p \frac{\partial T}{\partial t} = \frac{\partial}{\partial z} \left(\kappa \frac{\partial T}{\partial z} \right) + Q \quad (1)$$

where Q is the energy sink or source, representing the latent heat associated with water-ice phase transition, ρ is the soil density, C_p is the specific heat capacity and K is heat conductivity. The depth and time-dependent material properties were calculated assuming linear superposition of the soil, water, and ice properties (e.g., Lunardini, 1988). Thus, the depth-dependent density, heat capacity, and thermal
 285 conductivity were calculated using the porosity, θ , and freezing ratio, $B(z,t)$:

$$\begin{aligned} \rho &= (1 - \theta) \rho_{soil} + \theta ((1 - B)\rho_{ice} + B\rho_{water}) \\ \kappa &= (1 - \theta) \kappa_{soil} + \theta ((1 - B)\kappa_{ice} + B\kappa_{water}) \\ C_p &= (1 - \theta) C_{p_{soil}} + \theta ((1 - B)C_{p_{ice}} + BC_{p_{water}}) \end{aligned} \quad (2)$$

290

The thermal properties and density used for all system components (soil, ice and water) are listed in Table 2.

Out of the mushy zone, for either $B=0$ or $B=1$ (unfrozen sediments), the heat exchange leads to its cooling below or heating above the freezing temperature. When no latent heat is involved, assuming
 295 homogeneous heat conductivity, the heat conduction equation (1) is reduced to:

$$\frac{\partial T}{\partial t} = D \frac{\partial^2 T}{\partial z^2} \quad (3)$$

where D (diffusivity) [$m^2 s^{-1}$] defined as:

$$D = \frac{\kappa}{\rho C_p}$$

In the mushy zone, where the water-ice phase transition occurs, both the B value and the thermal
 300 properties (C_p and K) are depth-dependent. Accordingly, the complete heat conduction equation (1) is solved, including the latent heat term. The heat source/sink is equal to the mass of the freezing/thawing water per unit time multiplied by the latent heat, L . The water mass is equal to the rate of the B -value change times porosity and density. Finally, the source term is:

$$Q = -L \theta \rho \frac{\partial B}{\partial t} \quad (4)$$

305 We neglected the kinetics of the phase transition and assumed that thermodynamic equilibrium is established instantaneously in the mushy zone. This means that the rate of freezing/thawing is defined

by the heat flux to and from the mushy zone with $0 < B < 1$. Substituting (4) into (1) and using $\frac{\partial T}{\partial t} = 0$ leads to:

$$\theta L \rho \frac{\partial B}{\partial t} = \frac{\partial}{\partial z} \left(\kappa \frac{\partial T}{\partial z} \right) \quad (5)$$

310 The above equations are solved numerically for two functions $T(z, t)$ and $B(z, t)$, using the explicit-in-time finite difference scheme. These functions were approximated using the constant grid steps in depth Δz and in time Δt :

$$T_{n,m} = T(n\Delta z, m\Delta t)$$

$$B_{n,m} = B(n\Delta z, m\Delta t)$$

315 where n is the grid point number ($z = n\Delta z$) and m is the time step number ($t = m\Delta t$). With this notation, the finite difference form of the heat conduction equation (1) is:

$$\begin{aligned} \rho_n C p_n \frac{T_{n,m+1} - T_{n,m}}{\Delta t} = \\ = \frac{1}{\Delta z^2} \left[\frac{\kappa_{m+1} + \kappa_m}{2} (T_{n+1,m} - T_{n,m}) - \frac{\kappa_m + \kappa_{m-1}}{2} (T_{n,m} - T_{n,m-1}) \right] + L \rho \theta \frac{B_{n,m+1} - B_{n,m}}{\Delta t} \end{aligned} \quad (6)$$

where density and the thermal properties are calculated using equation (2).

320 Equation (6) is solved using time step $\Delta t = 10,800$ s (i.e. 3 hours) and a depth spacing $\Delta z = 0.25$ m. The solution was obtained for the model size down to 300 m depth, summing up to 1,200 grid points. These numerical parameters satisfy the von Neumann stability condition for explicit-in-time numerical scheme (e.g., Ames, 1977) for the material properties of Table 2. The numerical code was written with Python (Wang and Oliphant, 2012). It allows simulating the permafrost dynamics and sub-surface sediments
325 freezing under various scenarios of MAAT, water freezing temperatures (WFT) and freezing extent of pore space water.

330

Table 2: 1-D heat transfer model physical parameters of water, ice and sediment.

	Thermal conductivity	Heat capacity	Density	Diffusivity	Latent heat
	κ [W m ⁻¹ K ⁻¹]	Cp [J K ⁻¹ Kg ⁻¹]	ρ [Kg m ⁻³]	D [m ² s ⁻¹]*	L [J Kg ⁻¹]
Ice	2.24	2100	916.2	1.17X10 ⁻⁶	334000
Water	0.569	4192	999.85	1.36X10 ⁻⁷	
Mineral fraction	3	837**	2400	1.74x10 ⁻⁷	

* $D = \kappa / \rho * C_p$

** Value refers to saturated pore space without air.

5.3 Model results

We present the results of model runs with various combinations of surface temperature, water freezing temperature and sediments porosities and thermal conductivities. Complementary modelling results are presented in the Appendix (DOI: [10.6084/m9.figshare.23684877](https://doi.org/10.6084/m9.figshare.23684877)). In all cases, simulations started in the spring (May) and followed an amplitude of 12°C around the chosen (fixed) MAAT.

We first present simulations with MAAT of $-4 \pm 12^\circ\text{C}$, WFT 0°C , and complete (100%) freezing. Figure 4a presents results of a one-year simulation, with temperature profiles shown every second month. The model suggests that freezing under these conditions can reach down to three meters within the first year. The freezing depth increases to 6 m within 6 years with a slight, but significant deepening of the inflection point (Fig. 4b). After 50 years, freezing arrives at 16-17 m, and within 1000 years the freezing front is already at 73 m (Fig. 4c), within basement rocks (considering that sediment cover at ADE is ca. 25 m). The depth affected by cooling also progresses with time (<50 m in 50 years, >150 m in 1000 years) and T profile approaches linearity.




Figure 4. 1-D freezing model results for: (a) 1 year; the model starts (and finishes) at spring, defined as mid-time between minimum and maximum surface T (e.g. May), followed by summer increase in temperature (solid and dashed lines), cooling and start of freezing during fall (dotted line, e.g. November), followed by colder winter months (two-points dashed line and wide dashed line) and concluding in the spring (one point dashed line); (b) 6 years (starting in spring). (c) 50 to 1000 years. MAAT $-4 \pm 12^{\circ}\text{C}$, WFT set at 0°C , 100% freezing. Full black circles denote -4°C . Note the different scale between (a) (b) and (c).

Lowering the WFT results in a decrease of the freezing rate. For example, with WFT of 0°C the freezing front will reach 23 and 37 m after 100 and 200 years while with WFT of -2°C and -4°C it will reach 15 and 20 m and 5.5 and 6.5 m during the same periods, respectively (Fig. 5). A thin layer of permafrost may aggrade even under WFT of -5°C which is lower than the MAAT of -4°C . The freezing front will reach 2.5 m resulting in 0.5 m of permafrost, considering hypothetical, conservative, active layer of 2 m thick for the high temperatures of the mid-Holocene, which is deeper than present (Humlum, 2003; Weinstein et al., 2019). This is because the thermal conductivity of ice is higher than that of water (Farouki, 1981), which results in a deeper advance of the winter freezing front (through ice) than the advance of summer thawing (through water). Last, with WFT of -6°C (i.e. significantly lower than the MAAT), there is no apparent permafrost aggradation (not shown).




Figure 5. Results of 1-D freezing simulations with mid-Holocene MAAT of $-4 \pm 12^\circ\text{C}$ (Park et al., 2019);
 380 pore-water freezing temperature (WFT) taken as 0°C , -2°C and -4°C . Simulations were run for 100 and
 200 years.

In Figure 6, we examine the effect of partial freezing. Partial freezing (25% and 50% in our scenarios),
 result in deepening of the freezing depth, but differences are relatively small for WFT lower than 0°C .
 385 With MAAT of -4°C and WFT of 0°C , after 200 years freezing depth will reach 37, 42 and 45 m with 100%
 50% and 25% freezing, respectively. Lowering WFT to -2°C , freezing depth is indifferent to the freezing
 degree (Fig. 6). This is because of the trade-of between reducing latent heat and the lower thermal
 conductivity of the partially frozen pore space. With a WFT of -4°C , under 100% freezing 6.5 m of
 permafrost will aggrade after 200 years while with partial freezing scenarios it will aggrade to 3.25 and
 390 4.25 m for 50 and 25% respectively (Fig. 6). With WFT of -5°C a thin layer of permafrost may aggrade

under the 100% freezing scenario (0.5 m, not shown) while with partial freezing of 25-50%, it will not aggrade. , i.e. no permafrost will develop (assuming active layer depth of 1-2 m), which further suggests that when WFT is lower than the average MAAT (-4°C), aggradation is controlled by the ice thermal conductivity rather than by latent heat.

395





Figure 6. Mid-Holocene permafrost aggradation with variable freezing degrees. MAAT was set to -4°C (Park et al., 2019), and freezing proportions were taken as 100%, 50% and 25%. The shaded/grayish zone represents a hypothetical, conservative, active layer, which is taken as 2 m thick. We note that active layer in Adventdalen is usually ≤ 1 m, and 2 m was chosen due to the higher temperature during the mid-Holocene.

In general, higher MAAT results in a lower aggradation rate e.g. with MAAT of 0°C and WFT of 0°C 100% freezing, freezing depth will reach 9.5 m within 200 years. For the same setting, but with MAAT of -4°C

the freezing front will result in significant deepening and will reach 37 m (Figure 7a). Setting the system to 25% freezing; MAAT = -4°C and WFT of 0 and -2°C , freezing depth will reach 47 and 20 m respectively (Fig. 7b). Replacing MAAT to 0°C , freezing depth will reach 4.25 m for WFT of 0°C and 2.75 m for WFT of -2°C in both scenarios freezing depth is constant for 200 years (Fig. 7b). Moreover, we show that even
410 with MAAT > WFT (e.g., 0°C and -2°C , respectively) freezing will arrive at 2.75 m after 10 years and may create a thin layer of permafrost for at least 200 years. . As mentioned above, this is due to the asymmetry in the seasonal freezing/thawing process. Higher conductivity during winter (frozen pore space) enhances the loss of heat, while the lower conductivity during summer (warming front goes through thawed pore space) slows the thawing process (Kukkonen and Šafanda, 2001). We note that lower
415 proportions of freezing (e.g., 25%) will have the effect of reducing this asymmetry due to the higher proportions of liquid water in the cryotic pore-space, therefore lower thermal conductivity during freezing. Accordingly, permafrost deepening is hardly observed in the scenario of MAAT= 0°C (in particular with WFT= -2°C).

420



425 Figure 7. Early and mid-Holocene freezing depth in the first 50 and 200 years with MAAT -4°C and 0°C ; WFT of 0 and -2°C . (a) 100% freezing, (b) 25% freezing. Also shown are curves of two scenarios of corresponding depths of the fresh-saline water interface, using a 1:40 Ghyben-Hertzberg (G-H) approximation for isostasy of 4.5 and 15 mm y^{-1} . See text for more details. (c) and (d) are zoomed in of (a) and (b) for the first 50 years. The fast deepening in figures a, and b is due to change in porosity as the freezing front reaches the bedrock (25 m). The legend is the same for all figures.

6. Discussion

6.1 Ground ice salinity and the frozen interface

430 When the freezing front propagates downwards in a recently emerged land and epigenetic permafrost is formed, it might freeze old subsurface brines (Cascoyne, 2000). As the freezing process proceeds, solute concentrations in the non-frozen residual water commonly increase (e.g., Cocks and Brower, 1974; Herut et al., 1990; El Kadi and Janajreh, 2017). This results in a pore space with high salt concentrations. These brines may then migrate away from the freezing surface, driven by density and capillary forces, and coalesce to form separate saline water lenses ('cryopegs'; Cascoyne, 2000). The level of salinity and water composition will depend on the initial water composition and the extent of freezing.

Complete permafrost freezing can hardly be obtained, since the eutectic point of seawater freezing is at -36°C to -54°C (Gitterman 1937; Ringer 1905; Nelson and Thompson 1954; Marion et al., 1999), while in Adventdalen permafrost temperatures does not usually get below -6°C, below depth of zero annual amplitude (ZAA) (Christiansen et al., 2010), and are never lower than -12°C even in the shallow permafrost (Christiansen et al., 2020; Isaksen et al., 2007). Although the eutectic point is well below the expected temperature values, the freezing-salt expulsion process still prevails. Under these conditions, the permafrost pore space should hold a small fraction of residual brine solution, which contains most of the solutes originally dissolved in the bulk pore-space water. Partly frozen permafrost has been often observed in the study area during drilling, in particular deeper than a few meters. This was also found in both geophysical and geochemical observations (e.g., Keating et al., 2018; Weinstein et al., 2019). Nevertheless, when ground ice is thawed, the extracted fluid from the relatively large segments used in this study should roughly indicate the salinity of the original in situ pore fluid, assuming no major brine migration had occurred. Pore-water composition may be significantly altered from the original fluid that circulated in the sediments (e.g. seawater) due to ion exchange or dissolution prior to or even after cryotic conditions occurred, which is reflected in the Na/Cl and SO₄/Cl ratios in the thawed ground ice (Fig. 2). Nevertheless, Cl⁻ concentration is probably close to and represents the salinity of the original pore fluid. We note that while in certain cases permafrost contains lenses or pockets of unfrozen brine-containing cryotic soils ('cryopegs', e.g. Van Everdingen, 1998), which are commonly attributed to the segregation and migration of fluids (i.e. non in situ). The relatively low salinity (Fig. 2) and the evident mixing profile (Fig. 2a, b and C), suggest that this is not the case in the ADE site, and that the observed fresh-seawater

interface is an in situ observation. We relate to the chemistry of the extracted fluid as 'ground ice chemistry', although it could as well be that some of it was not actually frozen.

460 Cable et al., (2018) presented ground ice chemistry of cores from the Adventdalen, albeit closer to the current fjord (<4 km), west of the ADE site. In these cores, chloride, sodium and sulphate concentrations at depths of 3-11 m were up to 50% that of seawater. At ADE, farther away from the sea, ground ice in the epigenetic permafrost, 5.5 m from the surface and deeper, shows a gradual increase in salinity (i.e., fresh-saline interface), with Cl⁻ concentrations reaching 15% that of seawater at 9 m below the surface. Although salinities do not change much between 9-12 m, it is likely that more saline water, close to 465 seawater salinity, either exists today or existed in the past (prior to freezing) at deeper permafrost levels. The existence of a fresh-saline interface in the very shallow permafrost, in the top of the epigenetic permafrost, suggests that freezing at ADE occurred straightaway after emergence above seawater. This is further discussed below.

6.2 Rebound, exposure and fresh-saline interface deepening

470 Assuming that Early Holocene (11-8 ka BP) precipitation was slightly higher than present (200 mm per year, McFarlin et al., 2018; Kjellman et al., 2020), and using a conservative infiltration factor of 0.2 (whether thawing snow or direct rain) and the porosity used in our simulations (0.3), this amounts to an effective annual infiltration of ca. 130 mm per year. This could easily keep-up with the Early Holocene rebound rates of 15 mm y⁻¹ (established for the nearby Sassendalen Valley, Salvigsen, 1984; Sessford 475 et al., 2015), therefore preserving the groundwater table close to the surface of the emerging land. Using a Ghyben-Herzberg approximation (Bear and Dagan 1964), this would result in an Early Holocene fresh-saline interface deepening of ca. 60 m in 100 years (Fig. 7), assuming the saline water body had a common seawater density of 1,025 kg m⁻³. FSI deepening could even be faster, if the deep-water body is less saline than seawater. Even if sub-aerial exposure occurred later, during the mid-Holocene, when 480 the rebound rate decreased to 4.5 mm y⁻¹ (Forman et al., 2004), the fresh-saline interface would still deepen at a rate of 180 mm y⁻¹, i.e. 18 m in 100 years (Fig. 7).

The existence of a mixing zone at the top of the epigenetic permafrost (from 5.5 m below the current terrain surface), with Cl⁻ content 15% that of seawater at 3.5 m below the Early Holocene surface, suggests that the marine sediment section at ADE was hardly flushed with meteoric water. This further 485 suggests that permafrost aggradation commenced shortly after emergence above the sea (e.g., Kasprzak et al., 2020). Indeed, some of the simulated freezing scenarios can clearly cope with the above

fresh-saline interface deepening rates (e.g., MAAT of -4°C and WFT of 0°C , Fig. 7). Moreover, assuming that partial freezing (e.g., 50-25%) can also block flushing, this results in even faster permafrost aggradation (Fig. 7b). However, as permafrost deepens, the freezing rate slows down (e.g., Fig. 7), and none of the scenarios can cope with the assumed deepening of the fresh-saline interface, which should result in flushing of deeper zones.

It seems likely that the key factor in fresh-saline interface fossilization in a continuous permafrost landscape is the permanent freezing of the very shallow permafrost, which hydraulically disconnects the sub-permafrost zone from the surface and prevents recharge of this zone with meteoric water. As shown (Fig. 6 and 7), freezing of the top 3-5 m can occur within several years even with the relatively high temperatures of the Early to mid-Holocene, e.g., MAAT of -4°C or even warmer and WFT 0°C or -2°C will result in 5 m of frozen sediments within 5 to 35 years (Fig. 7), therefore, the fresh-saline interface could effectively be preserved.

We did not include salt diffusion in our model (e.g. Angelopoulos et al., 2019), a process that will reduce WFT as freezing progresses. It can explain the reason for partially frozen samples extracted from the epigenetic section (table 1). Including salt diffusion, we assume the freezing front may have advanced somewhat more slowly than suggested by our model.

6.3 Permafrost aggradation during the Holocene

Gilbert et al., (2018), suggested that our drilling site at ADE emerged from the sea at 10 to 9 Ka BP and that the delta front advanced westwards at a rate of 4.4 m y^{-1} prior to 9.2 Ka, which decreased to 0.9 m y^{-1} during the rest of the Holocene. Considering the relatively high rebound rates during 9 to 8 ka (e.g., $15\text{-}19\text{ mm y}^{-1}$), this suggests that the land surface at ADE reached 3-4 meters above sea level and a topographic gradient of 1-2% towards the sea within 200 years. Assuming the groundwater table was close to the surface, this should further result in a good flushing of the subsurface, unless freezing took control (Fig. 8). The observed mixing zone, which reaches the very top of the pre-Late Holocene surface, suggests that freezing started within just a few years after exposure.

Our simulations suggest that both cryotic conditions (i.e., $<0^{\circ}\text{C}$) and actual ground ice formation were started very soon after exposure to the atmosphere (Fig. 5, 6 and 7), and that significant freezing depths of 20-37 m could be achieved with 200 years (Fig. 6 and 7). This is true for both the Early to mid-Holocene warmer period (Kutzbach and Guetter, 1986; McFarlin et al., 2018; Mangerud and Svendsen, 2017; Park

et al., 2019; Kjellman et al., 2020) and for any sub-zero MAAT, regardless the size of the annual fluctuations. We even tested MAAT of +1°C and found that some freezing could occur (not shown), which is a seasonal effect, derived from the different thermal conductivities of ice and water. We also tested
520 our model with an order of magnitude lower thermal conductivity, following recently published research in Adventdalen (Hornum et al., 2020) and found that permafrost aggradation, although shallower, will result with freezing of the FSI under the mid-Holocene conditions. Our simulations are in good agreement with Harada and Yoshikawa (1996), who used 1-D model with a MAGST of -5.7°C but not completely saturated sediments and found that 533 years are needed to freeze 31.7 m of sediments in
525 Moskuslagoon, slightly to the west of our site, on the Adventfjorden shore.


The ADE site was free of sea water in the Early Holocene, prior to 9.2 ka BP (Gilbert et al. 2018). At that time, an abrupt cooling has been described in Svalbard (Mangerud and Svendsen, 2017; van der Bilt et al., 2018, 2019). The presented model results show that the initiation of permafrost and its gradual aggradation is possible under relatively high temperatures (yet MAAT \leq 0°C) of the mid-Holocene.
530 Christiansen et al. (2013), pointed out that local topographic conditions and winds in Adventdalen can induce lower temperatures at low altitude depressions, which could enhance the permafrost aggradation during the mid-Holocene.

Our simulations also show that lower percentage of freezing (e.g., 25%) may enhance the permafrost aggradation rate (e.g., Fig. 7b); however, this is not true for lower WFT (e.g., -2°C) or for relatively high
535 MAAT (e.g. 0°C), which as mentioned above is due to the trade-off between latent heat and thermal conductivity differences between ice and liquid water. Nevertheless, MAAT of 0°C seems unlikely (e.g., van der Bilt et al., 2019).


In summary, the simulations suggest that permafrost aggradation could and did occur immediately following exposure also during the relatively warm period of early to mid-Holocene (8-10 ka BP). This is
540 in disagreement with Hornum et al., 2020, who suggested that while freezing did occur during the Early Holocene the ground thawed during the mid-Holocene and refroze at about 6.5 Ka BP.

Pleistocene permafrost dynamics were studied in other permafrost regions. While in some areas there are records of permafrost degradation and peatland expansion already in the Early Holocene (post deglaciation, e.g., Lenz et al., 2015; Kaufman et al., 2015; Grinter et al., 2018; Li et al., 2021), cumulated
545 evidence indicates that air temperature during this period was highly variable, sometimes higher and sometimes lower than presently (Kaufman et al., 2015). Nevertheless, it is a common observation that during the Holocene Thermal Maximum (mid-Holocene, 8.2-4.2 ka BP) permafrost has been degrading


and thermokarst peaked (e.g., Lenz et al., 2015; Ulrich et al., 2017; Anderson et al., 2019). Permafrost aggradation resumed post- 6 ka, and mainly during the past 4-3 ka (e.g., Grinter et al., 2018; Treat and Jones, 2018). As shown above, we believe that this was not the case in Svalbard, and exposed lands continued freezing throughout the Pleistocene.




Stage 1: 22K Ka BP: Last glacial cycle, the glacier advances, and the valley bottom is eroded down to the bedrock (Elverhøi et al., 1995). The FSI is probably located at the meeting point of the sea and land.



Stage 2: 10-10.5 Ka BP: Maximal sea ingression (Lønne & Nemeč, 2004) and deltaic sediments deposition (Gilbert et al., 2018). The FSI migrates eastward to the new location where sea and land meet.



Stage 3: 9.5 Ka BP: The FSI migrates westward and freezes when temperatures drop sharply. Freezing front (from top down) exceeds fresh water lateral flow. Epigenetic permafrost aggrades.



Stage 4: ADE site at present. Fluvial and aeolian deposition freeze syngenetically (Gilbert et al., 2018).

555 Figure 8. Conceptual presentation of the type of ground water freezing processes in Adventdalen since the last glacial cycle with a focus on the fresh-saline water interface (FSI) at the ADE site.

7. Summary and conclusions

560 Arctic landscapes, including western Svalbard, were rising relatively fast in the Early to mid-Holocene due to glacial isostatic rebound. Accordingly, the preservation of a frozen saline water (mixing zone) at a very shallow depth 5.5 to 12 m is taken as evidence for fast permafrost aggradation, which could halt the infiltration of fresh meteoric water and the flushing of saline water toward the sea. This is despite of the prevailing relatively high temperatures during this period.

565 Our modelling confirms that freezing could progress relatively fast down the exposed Adventdalen sediments, i.e. to 20-35 m within 200 years, even under the reconstructed relatively high mid-Holocene air temperature of -4°C used in the 1-D model.

The modelling further suggests that permafrost may aggrade even when water freezing temperature (WFT) is slightly lower than mean annual air temperature (MAAT). This is attributed to the differences in thermal properties between ice and liquid water, where the higher conductivity of ice allows a faster 570 propagation of the freezing front during winter than the propagation of thawing during the summer.

Non-complete freezing of the cryogenic pore space (which is often the case) could result in faster deepening of the freezing front when MAAT is smaller than WFT or even when it is higher, but in the latter case the difference is not large (e.g., $\text{MAAT} < 0^{\circ}\text{C}$ and $0 < \text{WFT} > -2^{\circ}\text{C}$). However, when $\text{MAAT} \gg \text{WFT}$ (e.g., $\text{MAAT} = 0^{\circ}\text{C}$ and $\text{WFT} \leq -2^{\circ}\text{C}$), the presence of liquid water in the pore space and its lower 575 thermal conductivity would result in a halt of permafrost aggradation.

This concept of fast freezing under relatively high air temperatures may suggest that recently exposed areas may still go through permafrost aggradation even under the current global warming. Also, it could imply that a short (years to decades) cooling period could significantly slow down permafrost thawing.

8. Appendix

580 Appendix 1: Porosity analyses

Selected simulation results demonstrating the effects of the porosity values on the rate of permafrost formation are presented in figures 9 - 12. In general, higher porosity (i.e., more pore water to freeze) will result in slower permafrost aggradation. due to the higher latent heat involved. DOI: [10.6084/m9.figshare.23684877](https://doi.org/10.6084/m9.figshare.23684877).




Figure 9. Simulations of freezing front progress with different porosities for MAAT of -4°C , WFT of 0°C and 100% freezing. Inset present results for 1000 years. The fast deepening at depth > 25 m is due to
 590 change in porosity as the freezing front reaches the bedrock.




Figure 10. Simulations as in Fig. 9, but with WFT of -2°C.

595





Figure 11. Simulations as in Fig. 9 (MAAT= -4°C, WFT= 0°C), but with 25% freezing.



600 Figure 12. Simulations as in Fig. 9 (MAAT = -4°C and WFT = -2°C) but with 25% freezing.

9. Code availability

DOI: [10.6084/m9.figshare.23684493](https://doi.org/10.6084/m9.figshare.23684493)

1-D freeze/thaw model code – copy and paste code rows into Python (Spider-Anaconda).

605 # This Python-script describes the 1-D Heat transfer model code developed for the research first Rotem et al., (2023). The 1-D model is a transient one-dimensional heat transfer model suitable for simulating permafrost dynamics. The core of the model is an explicit forward-difference time approximation of the one-dimensional heat transfer equation. This script is tailored to simulate the Holocene ground temperature development in Adventdalen, Svalbard, but may be modified to fit other purposes.

610 #Usage must be cited by reference to Rotem et al. (2023).

#For references cited below see: Rotem, D., Lyakhovsky, V., Christiansen, H. H., Harlavan, Y., & Weinstein, Y. (2023). Permafrost saline water and Early to Mid-Holocene permafrost aggradation in Svalbard. *The Cryosphere*.

```

615 #Importing relevant packages for Python
import numpy as np
import matplotlib.pyplot as plt
import sys
import pandas as pd

620
n =1201          # number of point grid
m1 =20          #number of snapshots – i.e. No. of curves as display in fig. 4 and 5, or No. of sub
sets of data to be exported.
m2 = 365*50     # number of steps for every snapshot
625 m = m1*m2    # number of time step

dt = 3600.0*24.0/8    #1/8 of a day 10800 sec
t_final = dt*m        #defines the final time step
t_days=t_final/3600./24. #converts time steps to days
630 t_snap= dt*m2/3600./24.
print ('t_snap')
z=300.0          #meters depth
dz = (z/(n-1))   #meter - defines the cell width if cell width (dz).
Tfr=0           #Freezing temperature define as WFT in article

635
#define depth-porosity profile
Depth = np.zeros(n)
por = np.zeros(n)
for i in range (0,n):
640   Depth [i] = -i*dz
      por[i] = 0.3          # sediment's porosity
      if Depth[i] < -25.0: # change from sediment's porosity to rock porosity
          por[i] = 0.1

```



```

645 L = 334000.0          # J/kg water and ice Latent heat

#Density of materials
p_ice = 916.2          # Kg/m^3
p_water = 999.85      # Kg/m^3
650 p_soil = 2400.0    # Kg/m^3

#Defines Heat conductivity (K) and Heat Capacity (Cp) as function of porosity.
K_dufsoil = 3          #K_dufsoil = k of dry soil at temperatures ca. 5°C centigrade.
K_dfsoil = 3          #K_dfsoil = k of dry soil at temperatures lower than 0°C
655 Cp_dufsoil = 837    #heat capacity of dry soil (silt) in ca. 10°C Cp_dufsoil = Heat capacity of dry soil
above 0°C.
Cp_dfsoil = 712       #heat capacity of dry soil (silt) in ca. -10°C Cp_dfsoil = Heat capacity of dry soil
below 0°C.
K_ice = np.zeros(n)
660 K_water = np.zeros(n)
Cp_ice = np.zeros(n)
Cp_water = np.zeros(n)

for i in range (0,n):
665     K_ice[i] = 2.24 *por[i] + K_dfsoil *(1-por[i])          # W/(m*K) regression with 0<B<1 -1.651x
+ 2.22
     K_water[i] = 0.569*por[i] + K_dufsoil*(1-por[i])          # W/(m*K)
     Cp_ice[i] = 2100.0*por[i] + Cp_dfsoil *(1-por[i])          # j/(K*Kg)
     Cp_water[i] = 4192.0*por[i] + Cp_dufsoil*(1-por[i])        # j/(K*Kg) regression with 0<B<1 2.192x
670 + 2. considering 90% soil and 10% water.

#control on model stability condition for explicit-in-time numerical scheme
dd2=K_ice[i]/(p_soil*Cp_ice[i])*dt/dz/dz
dd3=K_water[i]/(p_soil*Cp_water[i])*dt/dz/dz
675 if dd2>0.25 or dd3>0.25:

```

```

        print(' Values must be < 0.25 ')
        print(dd2,dd3)
        print(i,Depth[i],por[i])
        sys.exit(' Decrease time step ')

680
x=np.linspace(0,z,n)

#creating time field for data export
time=np.zeros(n)
685 for i in range (0,n):
    time[i]=0+dt*i

#initial conditions
B=np.ones(n)      # B is a variable between 0 and 1 considering the ratio of ice or water in a cell.
690 B=1=water, B=0=ice creates an array of 1
T=np.ones(n)     # Initial Temperature 2 Centigrade creates array of 2 centigrade across the soil profile
for i in range(0,n-1):
    T[i]= 2.0 + 0.033*i*dz #Thermal gradient 0.033 centigrade per m

695 #Define working arrays
Tn=np.zeros(n)      # creates an array for each one of the variables
Bnew=np.zeros(n)
K=np.zeros(n)
Cp=np.zeros(n)
700 p=np.zeros(n)
dE2=np.zeros(n)

for j1 in range(0,m1):    #a loop on the snapshots
705 for j2 in range(0,m2) :
    #Boundary conditions

```

```

Time=(j1*m2+j2)*dt/3600./24/365 #years
T[0]=12*np.sin(2.*np.pi*Time)-4 #surface temperature for seasonal variation
B[0]=0. #B=1=water, B=0=ice
710 if T[0]>0:
    B[0]=1
    T[n-1]= 2 + 0.033*z #bottom of profile temperature. when sediments exposed to air it still has the sea
water temperature.
    B[n-1]=1. #B=1=water, B=0=ice
715
    for i in range(0,n): # calculate mixture properties each variable is calculated with linear ratio to B
(ice to water ratio in a cell).
        K[i]=K_ice[i]*(1.0-B[i])+K_water[i]*B[i]
        Cp[i]=Cp_ice[i]*(1.0-B[i])+Cp_water[i]*B[i]
720 p[i]=p_ice*(1.0-B[i])+p_water*B[i]

    for i in range(1,n-1): # loop over internal points. the dE2 (energy equation) equation is split for more
convenient calculations
        dE2[i]=(K[i-1]+K[i])/2.0 * (T[i-1]-T[i])/dz
725 dE2[i]=dE2[i]-(K[i]+K[i+1])/2.0 * (T[i]-T[i+1])/dz
        dE2[i]=dE2[i]*dt # Total energy flux into cell "i"
        Tn[i]=T[i]+dE2[i]/(p_soil*Cp[i]*dz) # conduction calculating the new temperature in the next
cell.
        Bnew[i]=B[i] # calculating the new ice/water ratio in the next cell
730 if dE2[i]<0.0 and Tn[i]<Tfr: #condition that verify the amount of energy and the new temperature.
if the energy gets less than 0 value it means that energy is escaping the cell and it will cool down or
freezes.
        if B[i]>0.0001: # Freezing
            Bnew[i]=B[i]+(dE2[i]-(T[i]-Tfr)*(p[i]*Cp[i]*dz))/(p[i]*L*dz*por[i]) # the Bnew depends on the
735 amount of energy that has been used to freeze the previous cell - the rest of the energy
            if Bnew[i]>0.0: # if the condition is true the new temperature equals freezing temperature.
                Tn[i]=Tfr

```

```

else:
    Tn[i]=Tfr+Bnew[i]*(L*por[i])/Cp[i] # if the condition is false (Bnew <0.0) the new temp. equals
740 the freezing temp.+Bnew.
    Bnew[i]=0.

if dE2[i]>0.0 and Tn[i]>Tfr:

745     if B[i]<0.9999:      # Thawing
        Bnew[i]=B[i]+(dE2[i]-(T[i]-Tfr)*(p[i]*Cp[i]*dz))/(p[i]*L*dz*por[i])
        if Bnew[i]<1.0:
            Tn[i]=Tfr
        else:
750         Tn[i]=Tfr+(Bnew[i]-1.0)*(L*por[i])/Cp[i]
            Bnew[i]=1.

for i in range(1,n-1):
    T[i]=Tn[i]
755    B[i]=Bnew[i]

plt.plot(T,-x, 'r',label="Temperature",linewidth=0.5)

#exporting data to csv file. Graphs was created with Microsoft Excel.
760 name_dict} =
'    Temperature': T,[ :0]
'    time':time,[ :0]
'    depth': Depth,[ :0]
'    B': B,[ :0]
765 '    K': K,[ :0]
'    Cp': Cp,[ :0]
'    p': p,[ :0]
'    dE2': dE2[ :0]

```

```

{
770 df = pd.DataFrame(name_dict)
      df.to_csv(r'C:\Users\ADMIN\Desktop\Python      Dotan\1Dmodel_1.csv',mode='a',      header=True,
float_format='%.3f') # defines the location of the data exported
pd.read_csv('trial2.csv').count()

775 #Commands for graph in python script. Graphs for article was created with Microsoft Excel.
axes1 = plt.gca()
plt.ylabel('Depth')
plt.legend(loc = "lower left")
plt.grid()
780 axes2 = axes1.twinx()
axes1.set_xlabel("Temperature")
plt.plot(B,-x, 'g',linestyle = '--',linewidth=0.7, label="B")
axes2.set_xticks([0., .2, .4, .6, .8, 1.0])
axes2.set_xlabel("B")
785 plt.legend(loc = "lower left")
plt.show()

```

10. Data availability

All raw data can be provided by the corresponding authors upon request.

790 **11. Executable research compendium (ERC)**

12. Sample availability

13. Supplement link: the link to the supplement will be included by Copernicus, if applicable.

14. Author contribution:

DR, YW, and HHC planned the drilling campaign; DR and YW, processed the cores and lab work in
795 UNIS Svalbard; DR and YH performed the water chemistry analysis at GSI; VL and DR developed the
1-D model; DR, YW and VL wrote the manuscript. All authors commented on the manuscript.

15. Competing interests:

The authors declare that they have no conflict of interest.

16. Disclaimer

800 **17. Acknowledgments**

We would like acknowledge Ullrich (Ulli) Neuman for leading the 2017 drilling campaign in Adventdalen.
Andreas Alexander and Graham L. Gilbert for field assistance. Danni Rohdent for lab assistance. Gerd-
Irena and UNIS logistics for their assistance with field and laboratory gear. GSI geochemical lab
members Olga Berlin, Galit Sharabi and Dina Siber for their assistance with chemistry analysis. Yosi
805 Yechieli for consulting about various issues of the article. The drilling campaign was supported by an
Arctic Field Grant from the Norwegian Research Council, Project Number: 269988 RiS ID: 10664.

References

- Ahonen, L.: Permafrost: occurrence and physiochemical processes (POSIVA--01-05). Finland, [ISBN](#)
[951-652-106-1](#), 2001.
- 810 Alsos, I. G., Sjögren, P., Edwards, M. E., Landvik, J. Y., Gielly, L., Forwick, M., Coissac E., Brown A.
G., Jakobsen L. V., Føreid M. K., and Pedersen, M. W.: Sedimentary ancient DNA from Lake

- Skartjørna, Svalbard: Assessing the resilience of arctic flora to Holocene climate change, *The Holocene*, 26(4), 627-642, <https://doi.org/10.1177%2F0959683615612563>, 2016.
- Ames, W. F.: Numerical methods for partial differential equations, Second edition, Academic press
815 INC, 1977.
- Anderson L, Edwards, M, Shapley, M. D., Finney, B. P. and Langdon, C.: Holocene Thermokarst Lake Dynamics in Northern Interior Alaska: The Interplay of Climate, Fire, and Subsurface Hydrology, *Front. Earth Sci.* 7:53, <https://doi.org/10.3389/feart.2019.00053>, 2019.
- Angelopoulos, M., Westermann, S., Overduin, P., Faguet, A., Olenchenko, V., Grosse, G., and
820 Grigoriev, M. N.: Heat and salt flow in subsea permafrost modeled with CryoGRID2. *Journal of Geophysical Research: Earth Surface*, 124(4), 920-937, <https://doi.org/10.1029/2018JF004823>, 2019.
- Arnscheidt, C. W. and Rothman, D. H., Routes to global glaciation. *Proceedings of the Royal Society A*, 476(2239), <https://doi.org/10.1098/rspa.2020.0303>, 2020.
- Balland, V., and Arp, P. A.: Modeling soil thermal conductivities over a wide range of
825 conditions. *Journal of Environmental Engineering and Science*, 4(6), 549-558, <https://doi.org/10.1139/s05-007>, 2005.
- Bear, J., and Dagan, G.: Some exact solutions of interface problems by means of the hodograph method, *JGR*, 69(8), 1563-1572, <https://doi.org/10.1029/JZ069i008p01563>, 1964.
- Benn, D., and Evans, D. J.: *Glaciers and glaciation*, Second edition, pp 707, Routledge, 2014.
- 830 Betlem, P., Midttømme, K., Jochmann, M., Senger, K., and Olausen, S.: Geothermal Gradients on Svalbard, Arctic Norway. In First EAGE/IGA/DGMK Joint Workshop on Deep Geothermal Energy (pp. cp-577). European Association of Geoscientists and Engineers, <https://doi.org/10.3997/2214-4609.201802945>, 2018.
- <https://doi.org/10.1177/095968369100100303>, 1991.
- 835 Black, R. F.: Permafrost: a review, *GSA Bulletin*, 65(9), 839-856, [https://doi.org/10.1130/0016-7606\(1954\)65\[839:PR\]2.0.CO;2](https://doi.org/10.1130/0016-7606(1954)65[839:PR]2.0.CO;2), 1954.
- Bodnar, R. J.: Revised equation and table for determining the freezing point depression of H₂O-NaCl solutions. *GCA*, 57(3), 683-684, <http://www.osti.gov/scitech/biblio/6951353>, 1993.
- Burn, C. R.: Permafrost distribution and stability, edited by: French, H., and Slaymaker, O., *Changing*
840 *Cold Environments: A Canadian Perspective*, John Wiley and Sons, Ltd, 126-143, <https://doi.org/10.1002/9781119950172.ch7>, 2011.

- Burt, T. P., and Williams, P. J.: Hydraulic conductivity in frozen soils, *Earth Surface Processes*, 1(4), 349-360, <https://doi.org/10.1002/esp.3290010404>, 1976.
- Cable, S., Elberling, B., and Kroon, A.: Holocene permafrost history and cryostratigraphy in the High-
845 Arctic Adventdalen Valley, central Svalbard. *Boreas*, 47(2), 423-442, <https://doi.org/10.1111/bor.12286>, 2018.
- Cary, J. W., and Mayland, H. F.: Salt and water movement in unsaturated frozen soil. *Soil Science Society of America Journal*, 36(4), 549-555, <https://doi.org/10.2136/sssaj1972.03615995003600040019x>, 1972.
- 850 Cascoyne, M.: A review of published literature on the effects of permafrost on the hydrogeochemistry of bedrock, Technical Report, Posiva Oy, Helsinki Finland, 2000.
- Christiansen, H. H.: Thermal regime of ice-wedge cracking in Adventdalen, Svalbard. *PERMAFROST PERIGLAC*, 16(1), 87-98, <https://doi.org/10.1002/ppp.523>, 2005.
- Christiansen, H. H., French, H. M., and Humlum, O.: Permafrost in the Gruve-7 mine, Adventdalen,
855 Svalbard. *NORSK GEOGR TIDSSKR*, 59(2), 109-115, <https://doi.org/10.1080/00291950510020592>, 2005.
- Christiansen, H. H., Etzelmüller, B., Isaksen, K., Juliussen, H., Farbrot, H., Humlum, O., Johansson, M., Ingeman-Nielsen, T., Kristensen, L., Hjort, J., Holmlund, P. Sannel, A. B. K. Sigsgaard, C. Åkerman, H. J. Foged, N. Blikra, L. H. Pernosky, M. A. and Ødegård, R. S.: The thermal state of
860 permafrost in the Nordic area during the International Polar Year 2007–2009. *PERMAFROST PERIGLAC*, 21(2), 156-181, <https://doi.org/10.1002/ppp.687>, 2010.
- Christiansen, H. H., Humlum, O., and Eckerstorfer, M.: Central Svalbard 2000–2011 meteorological dynamics and periglacial landscape response, *ARCT ANTARCT ALP RES*, 45(1), 6-18,
865 <https://doi.org/10.1657/1938-4246-45.16>, 2013.
- Christiansen, H.H., Gilbert, G.L., Demidov, N., Guglielmin, M., Isaksen, K., Osuch, M. and Boike, J.: Permafrost temperatures and active layer thickness in Svalbard 2017-2018. *State of Environmental Science in Svalbard*, Van den Heuvel F, Hübner C, Błaszczuk M, Heimann M, Lihavainen H (eds) 2020: SESS report 2019, Longyearbyen, Svalbard Integrated Arctic Earth Observing System, Report
870 card, p. 236-249, [10013/epic.b4472816-40ba-4089-9ce7-d7539e10e0a3](https://doi.org/10.10013/epic.b4472816-40ba-4089-9ce7-d7539e10e0a3), 2020.
- Cocks, F. H., and Brower, W. E.: Phase diagram relationships in cryobiology, *CRYOBIOLOGY*, 11(4), 340-358, [https://doi.org/10.1016/0011-2240\(74\)90011-X](https://doi.org/10.1016/0011-2240(74)90011-X), 1974.

- Cochand, M., Molson, J., and Lemieux, J. M.: Groundwater hydrogeochemistry in permafrost regions, *PERMAFROST PERIGLAC*, 30(2), 90-103, <https://doi.org/10.1002/ppp.1998>, 2019.
- 875 Crank, J.: Free and moving boundary problems. Oxford University Press, USA, 1984. Cosenza, P., Guerin, R., and Tabbagh, A.: Relationship between thermal conductivity and water content of soils using numerical modelling. *European Journal of Soil Science*, 54(3), 581-588, <https://doi.org/10.1046/j.1365-2389.2003.00539.x>, 2003.
- 880 Dobinski, W.: Permafrost. *EARTH-SCI REV*, 108(3-4), 158-169, <https://doi.org/10.1016/j.earscirev.2011.06.007>, 2011.
- de Baar, H.J.W., van Heuven, S.M.A.C., Middag, R. : Ocean Salinity, Major Elements, and Thermohaline Circulation. In: White, W. (eds) *Encyclopedia of Geochemistry*. Encyclopedia of Earth Sciences Series. Springer, Cham. https://doi.org/10.1007/978-3-319-39193-9_120-1, 2017.
- 885 Edmunds, W. M., Hinsby, K., Marlin, C., de Melo, M. C., Manzano, M., Vaikmaa, R., and Travi, Y.: Evolution of groundwater systems at the European coastline. *GEOL SOC SP*, London, 189(1), 289-311, <https://doi.org/10.1144/GSL.SP.2001.189.01.17>, 2001.
- El Kadi, K., and Janajreh, I.: Desalination by freeze crystallization: an overview, *Int. J. Therm. Environ. Eng*, 15(2), 103-110, Doi: 10.5383/ijtee.15.02.004, 2017.
- 890 Elverhøi, A., Svendsen, J. I., Solheim, A., Andersen, E. S., Milliman, J., Mangerud, J., and Hooke, R. L.: Late Quaternary sediment yield from the high Arctic Svalbard area. *The Journal of Geology*, 103(1), 1-17, <https://www.jstor.org/stable/30071132>, 1995.
- Etzelmüller, B., Schuler, T. V., Isaksen, K., Christiansen, H. H., Farbrot, H., and Benestad, R.: Modeling the temperature evolution of Svalbard permafrost during the 20th and 21st century, *The Cryosphere*, 5, 67–79, <https://doi.org/10.5194/tc-5-67-2011>, 2011.
- 895 Farbrot, H., Etzelmüller, B., Schuler, T. V., Guðmundsson, Á., Eiken, T., Humlum, O., and Björnsson, H.: Thermal characteristics and impact of climate change on mountain permafrost in Iceland. *J GEOPHYS RES-EARTH*, 112(F3), <https://doi.org/10.1029/2006JF000541>, 2007.
- Farnsworth, W. R. The Topographical and Meteorological Influence on Snow Distribution in Central Spitsbergen: How the spatial variability of snow influences slope-scale stability, permafrost landform dynamics and regional distribution trends The Topographical and Meteorological Influence on Snow Distribution in Central Svalbard. Master Thesis, department of geosciences faculty of mathematics and natural sciences university of Oslo, 2013.
- 900

- Farnsworth, W. R., Ingólfsson, Ó., Alexanderson, H., Allaart, L., Forwick, M., Noormets, R., Retelle, M.,
905 and Schomacker, A.: Holocene glacial history of Svalbard: Status, perspectives and
challenges, *EARTH-SCI REV*, 103249, <https://doi.org/10.1016/j.earscirev.2020.103249>, 2020.
- Farouki, O. T.: Thermal properties of soils. Cold Regions Research and Engineering Lab Hanover NH.,
1981.
- Forman, S. L., Lubinski, D. J., Ingólfsson, Ó., Zeeberg, J. J., Snyder, J. A., Siegert, M. J., and
910 Matishov, G. G.: A review of postglacial emergence on Svalbard, Franz Josef Land and Novaya
Zemlya, northern Eurasia. *QUATERNARY SCI REV*, 23(11-13), 1391-1434,
<https://doi.org/10.1016/j.quascirev.2003.12.007>, 2004.
- French, H. M.: The periglacial environment, fourth edition, John Wiley and Sons LTD, 2017.
- Gilbert, G. L., Christiansen, H. H., and Neumann, U.: Coring of unconsolidated permafrost deposits:
915 methodological successes and challenges, In Proceedings GeoQuébec 2015 – 68th Canadian
Geotechnical Conference and 7th Canadian Permafrost Conference, 20–23, Québec, Canada. Paper 6
pp. <https://hdl.handle.net/1956/17626>, 2015.
- Gilbert, G. L., O'Neill, H. B., Nemeč, W., Thiel, C., Christiansen, H. H., and Buylaert, J. P.: Late
Quaternary sedimentation and permafrost development in a Svalbard fjord-valley, Norwegian high
920 Arctic. *Sedimentology*, 65(7), 2531-2558, <https://doi.org/10.1111/sed.12476>, 2018.
- Gilbert, G., Instanes, A., Sinitsyn, A., and Aalberg, A.: Characterization of two sites for geotechnical
testing in permafrost: Longyearbyen, Svalbard. <http://hdl.handle.net/11250/2632119>, 2019.
- Gitterman, K. E.: Thermal analysis of seawater. *CRREL TL*, 287, 1937.
- Grinter, M., Lacelle, D., Baranova, N., Murseli, S., and Clark, I. D.: Late Pleistocene and Holocene ice-
925 wedge activity on the Blackstone Plateau, central Yukon, Canada. *QUATERNARY RES*, 1–15.
<https://doi.org/10.1017/qua.2018.65>, 2018.
- Grünberg, I., Wilcox, E. J., Zwieback, S., Marsh, P., and Boike, J.: Linking tundra vegetation, snow, soil
temperature, and permafrost, *Biogeosciences*, 17, 4261–4279, [https://doi.org/10.5194/bg-17-4261-](https://doi.org/10.5194/bg-17-4261-2020)
[2020](https://doi.org/10.5194/bg-17-4261-2020), 2020.
- 930 Grundvåg, S. A., Jelby, M. E., Śliwińska, K. K., Nøhr-Hansen, H., Aadland, T., Sandvik, S. E.,
Tennvassås, I., Engen, T., and Olausen, S.: Sedimentology and palynology of the Lower Cretaceous
succession of central Spitsbergen: integration of subsurface and outcrop data. *NORW J GEOL*,
99(2):253-284, <https://dx.doi.org/10.17850/njg99-2-02>, 2019.

- Harada, K., and Yoshikawa, K.: Permafrost age and thickness near Adventfjorden, Spitsbergen. *Polar Geography*, 20(4), 267-281, <https://doi.org/10.1080/10889379609377607>, 1996.
- 935 He, H., Flerchinger, G. N., Kojima, Y., Dyck, M., and Lv, J.: A review and evaluation of 39 thermal conductivity models for frozen soils. *Geoderma*, 382, 114694, <https://doi.org/10.1016/j.geoderma.2020.114694>, 2021.
- Herut, B., Starinsky, A., Katz, A., and Bein, A.: The role of seawater freezing in the formation of subsurface brines. *GEOCHIM COSMOCHIM AC*, 54(1), 13-21, [https://doi.org/10.1016/0016-7037\(90\)90190-V](https://doi.org/10.1016/0016-7037(90)90190-V), 1990.
- 940 Hodson, A. J., Nowak, A., Hornum, M. T., Senger, K., Redeker, K., Christiansen, H. H., ... & Marca, A.: Sub-permafrost methane seepage from open-system pingos in Svalbard. *The Cryosphere*, 14(11), 3829-3842, <https://doi.org/10.5194/tc-14-3829-2020>, 2020.
- 945 Homshaw, L. G.: Freezing and melting temperature hysteresis of water in porous materials: Application to the study of pore form, *J SOIL SCI*, 31(3), 399-414, <https://doi.org/10.1111/j.1365-2389.1980.tb02090.x>, 1980.
- Hornum, M. T., Hodson, A. J., Jessen, S., Bense, V., and Senger, K.: Numerical modelling of permafrost spring discharge and open-system pingo formation induced by basal permafrost aggradation, *The Cryosphere*, 14, 4627–4651, <https://doi.org/10.5194/tc-14-4627-2020>, 2020.
- 950 Hornum, M. T., Betlem, P., and Hodson, A.: Groundwater flow through continuous permafrost along geological boundary revealed by electrical resistivity tomography. *Geophysical Research Letters*, 48(14), <https://doi.org/10.1029/2021GL092757>, 2021.
- Humlum, O., Instanes, A., and Sollid, J. L.: Permafrost in Svalbard: a review of research history, climatic background and engineering challenges. *POLAR RES*, 22(2), 191-215, <https://doi.org/10.1111/j.1751-8369.2003.tb00107.x>, 2003.
- 955 Humlum, O.: Holocene permafrost aggradation in Svalbard. *GEOL SOC SPEC PUBL*, London, 242(1), 119-129, <https://doi.org/10.1144/GSL.SP.2005.242.01.11>, 2005.
- Imbrie, J., Berger, A., Boyle, E., Clemens, S. C., Duffy, A., Howard, W. R., Kukla, G., Kutzbach, J., Martinson, D. G., McIntyre, A., Mix, A. C., Molino, B., Morley, J. J., Peterson, L. C., Pisias, N. G., Prell, M. E., Raymo, W. L., Shackleton, N. J. and Toggweiler, J. R.: On the structure and origin of major glaciation cycles 2. The 100,000-year cycle. *Paleoceanography*, 8(6), 699-735, <https://doi.org/10.1029/93PA02751>, 1993.
- 960

- Isaksen, K., Benestad, R. E., Harris, C., and Sollid, J. L.: Recent extreme near-surface permafrost
965 temperatures on Svalbard in relation to future climate scenarios. *Geophys Res Lett*, 34(17),
<https://doi.org/10.1029/2007GL031002>, 2007.
- Kasprzak, M., Łopuch, M., Głowacki, T., and Milczarek, W., Evolution of near-shore outwash fans and
permafrost spreading under their surface: A case study from Svalbard, *Remote Sens-Basel*, 12(3),
482, <https://doi.org/10.3390/rs12030482>, 2020.
- 970 Kaufman, D. S., Axford, Y. L., Henderson, A. C., McKay, N. P., Oswald, W. W., Saenger, C.,
Anderson, R. S., Bailey, H. L., Clegg, B., Gajewski, K., Hu, F. S., Jones, M. C., Massa, C., Routson, C.
C., Werner, A., Wooller, M. J., and Yu, Z.: Holocene climate changes in eastern Beringia (NW North
America)—A systematic review of multi-proxy evidence, *Quaternary Sci Rev*, 147, 312-339,
<https://doi.org/10.1016/j.quascirev.2015.10.021>, 2015.
- 975 Keating, K., Binley, A., Bense, V., Van Dam, R. L., and Christiansen, H. H.: Combined geophysical
measurements provide evidence for unfrozen water in permafrost in the Adventdalen valley in
Svalbard. *Geophys Res Lett*, 45(15), 7606-7614, <https://doi.org/10.1029/2017GL076508>, 2018.
- Kjellman, S. E., Schomacker, A., Thomas, E. K., Håkansson, L., Duboscq, S., Cluett, A. A.,
Farnsworth, W. R., Allaart L., Cowling, O. C., McKay, N. P., Brynjólfsson, S., and Ingólfsson, Ó.:
980 Holocene precipitation seasonality in northern Svalbard: influence of sea ice and regional ocean
surface conditions. *Quaternary Sci Rev*, 240, 106388, <https://doi.org/10.1016/j.quascirev.2020.106388>,
2020.
- Kokelj, S. V., Smith, C. A. S., and Burn, C. R.: Physical and chemical characteristics of the active layer
and permafrost, Herschel Island, western Arctic Coast, Canada. *Permafrost and Periglacial*
985 *Processes*, 13(2), 171-185, <https://doi.org/10.1002/ppp.417>, 2002.
- Kukkonen, I. T., and Šafanda, J.: Numerical modelling of permafrost in bedrock in northern
Fennoscandia during the Holocene. *Global and Planet Change*, 29(3-4), 259-273,
[https://doi.org/10.1016/S0921-8181\(01\)00094-7](https://doi.org/10.1016/S0921-8181(01)00094-7), 2001.
- Kutzbach, J. E., and Guetter, P. J.: The influence of changing orbital parameters and surface boundary
990 conditions on climate simulations for the past 18 000 years. *Journal of atmospheric sciences*, 43(16),
1726-1759, [https://doi.org/10.1175/1520-0469\(1986\)043<1726:TIOCOP>2.0.CO;2](https://doi.org/10.1175/1520-0469(1986)043<1726:TIOCOP>2.0.CO;2), 1986.

- Landvik, J. Y., Mangerud, J., and Salvigsen, O.: Glacial history and permafrost in the Svalbard area. In Proceedings of the 5th International Conference on Permafrost (Vol. 1, pp. 194-198), Trondheim, Tapir Publishers., 1988.
- 995
- Lemieux, J. M., Sudicky, E. A., Peltier, W. R., and Tarasov, L.: Simulating the impact of glaciations on continental groundwater flow systems: 1. Relevant processes and model formulation. *J Geophys Res-Earth*, 113(F3), <https://doi.org/10.1029/2007JF000928>, 2008.
- Lenz, J., Grosse, G., Jones, B. M., Walter Anthony, K. M., Bobrov, A., Wulf, S., and Wetterich, S.: Mid-1000 Wisconsin to Holocene Permafrost and Landscape Dynamics based on a Drained Lake Basin Core from the Northern Seward Peninsula, Northwest Alaska. *Permafrost Periglac*, 27(1), 56–75. <https://doi.org/10.1002/ppp.1848>, 2015.
- Lønne, I., and Lyså, A.: Deglaciation dynamics following the Little Ice Age on Svalbard: implications for shaping of landscapes at high latitudes. *Geomorphology*, 72(1-4), 300-319, <https://doi.org/10.1016/j.geomorph.2005.06.003>, 2005.
- 1005
- Lønne, I., and Nemeč, W.: High-arctic fan delta recording deglaciation and environment disequilibrium. *Sedimentology*, 51(3), 553-589, <https://doi.org/10.1111/j.1365-3091.2004.00636.x>, 2004.
- Lunardini, V. J.: Freezing of soil with an unfrozen water content and variable thermal properties (Vol. 1010 88, No. 2). US Army Corps of Engineers, Cold Regions Research and Engineering Laboratory, 1988.
- Luo, D., Jin, H., Marchenko, S. S., and Romanovsky, V. E.: Difference between near-surface air, land surface and ground surface temperatures and their influences on the frozen ground on the Qinghai-Tibet Plateau. *Geoderma*, 312, 74-85., <https://doi.org/10.1016/j.geoderma.2017.09.037>, 2018.
- Lüthi, Z. L.: Thermal State of Permafrost in Central and Western Spitsbergen 2008-2009, Master's 1015 Thesis Faculty of Science University of Bern, 2010.
- Mangerud, J., Bolstad, M., Elgersma, A., Helliksen, D., Landvik, J. Y., Lønne, I., Lycke, A. K., Salvigsen, O., Sandahl, T., and Svendsen, J. I.: The last glacial maximum on Spitsbergen, Svalbard. *Quaternary Res*, 38(1), 1-31, [https://doi.org/10.1016/0033-5894\(92\)90027-G](https://doi.org/10.1016/0033-5894(92)90027-G), 1992.
- 1020
- Mangerud, J., Astakhov, V., and Svendsen, J. I.: The extent of the Barents–Kara ice sheet during the Last Glacial Maximum, *Quaternary Sci Rev*, 21(1-3), 111-119, [https://doi.org/10.1016/S0277-3791\(01\)00088-9](https://doi.org/10.1016/S0277-3791(01)00088-9), 2002.

- Mangerud, J., and Svendsen, J. I.: The Holocene thermal maximum around Svalbard, Arctic North Atlantic; molluscs show early and exceptional warmth, *The Holocene*, 28(1), 65-83, <https://doi.org/10.1177/0959683617715701>, 2017.
- 1025 Marion, G. M., Farren, R. E., and Komrowski, A. J.: Alternative pathways for seawater freezing. *Cold Reg Sci Technol*, 29(3), 259-266, [https://doi.org/10.1016/S0165-232X\(99\)00033-6](https://doi.org/10.1016/S0165-232X(99)00033-6), 1999.
- McEwen, T., and Marsily, G.de.: The potential significance of permafrost to the behaviour of a deep radioactive waste repository, (SKI-TR--91-8). Sweden, 1991.
- 1030 McFarlin, J. M., Axford, Y., Osburn, M. R., Kelly, M. A., Osterberg, E. C., and Farnsworth, L. B.: Pronounced summer warming in northwest Greenland during the Holocene and Last Interglacial. *PNAS*, 115(25), 6357-6362, <https://doi.org/10.1073/pnas.1720420115>, 2018.
- Morgenstern, N. R., and Anderson, D. M.: Physics, chemistry, and mechanics of frozen ground: a review. In *Permafrost: North American Contribution [to The] Second International Conference (Vol. 2, p. 257)*. National Academies, 1973.
- 1035 Murton, J. B.: What and where are periglacial landscapes?. *Permafrost Periglac*, 32(2), 186-212, <https://doi.org/10.1002/ppp.2102>, 2021.
- Nelson, K. H., and Thompson, T. G.: Deposition of salts from sea water by frigid concentration, 1954.
- Nordli, Ø., Przybylak, R., Ogilvie, A. E., and Isaksen, K.: Long-term temperature trends and variability on Spitsbergen: the extended Svalbard Airport temperature series, 1898–2012. *Polar res*, 33(1), 21349, <https://doi.org/10.3402/polar.v33.21349>, 2014.
- 1040 Nordli, Ø., Wyszynski, P., Gjeltén, H., Isaksen, K., Łupikasza, E., Niedźwiedz, T., and Przybylak, R.: Revisiting the extended Svalbard Airport monthly temperature series, and the compiled corresponding daily series 1898–2018, <http://repozytorium.umk.pl/handle/item/6323>, 2020.
- 1045 Major, H., and Nagy, J.: *Geology of the Adventdalen map area: with a geological map, Svalbard C9G 1: 100 000*, Norsk Polarinsitutt, Oslo, 1972.
- Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H. H., Dashtseren, A., Delaloye, R., Elberling, B., Etzelmüller, B., Kholodov, A., Khomutov, A., Kääh, A., Leibman, M. O., Lewkowicz, A. G., Panda, S. K., Romanovsky, V., Way, R. G. Westergaard-Nielsen, A., Wu, T., Yamkhin, J. and Zou, D.: Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale. *Earth-Sci Rev*, 193, 299-316, <https://doi.org/10.1016/j.earscirev.2019.04.023>, 2019.
- 1050

- Olaussen, S., Senger, K., Braathen, A., Grundvåg, S. A., and Mørk, A.: You learn as long as you drill; research synthesis from the Longyearbyen CO2 Laboratory, Svalbard, Norway. *Norw J Geol*, 99(2), 157-187, <https://doi.org/10.17850/njg008>, 2019.
- 1055 Oldenborger, G. A., and LeBlanc, A. M.: Monitoring changes in unfrozen water content with electrical resistivity surveys in cold continuous permafrost. *Geophys J Int*, 215(2), 965-977, <https://doi.org/10.1093/gji/ggy321>, 2018.
- Osuch, M., and Wawrzyniak, T.: Inter-and intra-annual changes in air temperature and precipitation in western Spitsbergen. *Int J Climatol*, 37(7), 3082-3097, <https://doi.org/10.1002/joc.4901>, 2017.
- 1060 Overduin, P. P., Schneider von Deimling, T., Miesner, F., Grigoriev, M. N., Ruppel, C. D., Vasiliev, A., et al.: Submarine permafrost map in the Arctic modeled using 1-D transient heat flux (SuPerMAP) *Journal of Geophysical Research: Oceans*, 124, 3490– 3507, <https://doi.org/10.1029/2018JC014675>, 2019.
- Park, H. S., Kim, S. J., Stewart, A. L., Son, S. W., and Seo, K. H.: Mid-Holocene Northern Hemisphere warming driven by Arctic amplification. *Science advances*, 5(12), eaax8203, <https://www.science.org/doi/abs/10.1126/sciadv.aax8203>, 2019.
- Patton, H., Hubbard, A., Andreassen, K., Auriac, A., Whitehouse, P. L., Stroeven, A. P., Shackleton, C., Winsborrow, M., Heyman, J., and Hall, A. M.: Deglaciation of the Eurasian ice sheet complex. *Quaternary Sci Rev*, 169, 148-172, <https://doi.org/10.1016/j.quascirev.2017.05.019>, 2017.
- 1070 Rasmussen, T. L., Forwick, M., and Mackensen, A.: Reconstruction of inflow of Atlantic Water to Isfjorden, Svalbard during the Holocene: Correlation to climate and seasonality, *Mar Micropaleontol*, 94, 80-90, <https://doi.org/10.1016/j.marmicro.2012.06.008>, 2012.
- Ringer, W. E. *Über die Veränderungen in der Zusammensetzung des Meerwassersalzes beim Ausfrieren*. O. Verlag, 1905.
- 1075 Rubinstein, L., Geiman, H., and Shachaf, M.: Heat transfer with a free boundary moving within a concentrated thermal capacity. *IMA J Appl Math*, 28(2), 131-147, <https://doi.org/10.1093/imamat/28.2.131>, 1982.
- Rühaak, W., Anbergen, H., Grenier, C., McKenzie, J., Kurylyk, B. L., Molson, J., Roux, N., and Sass, I. Benchmarking numerical freeze thaw models. *Energy Procedia*, 1080 <https://doi.org/10.1016/j.egypro.2015.07.866>, 2015.

- Russak, A., and Sivan, O.: Hydrogeochemical tool to identify salinization or freshening of coastal aquifers determined from combined field work, experiments, and modeling. *Environmental science and technology*, 44(11), 4096-4102, <https://doi.org/10.1021/es1003439>, 2010.
- 1085 Salvigsen, O.: Occurrence of pumice on raised beaches and Holocene shoreline displacement in the inner Isfjorden area, Svalbard, *Polar Res*, 2(1), 107-113, <https://doi.org/10.1111/j.1751-8369.1984.tb00488.x>, 1984.
- Šarler, B.: Stefan's work on solid-liquid phase changes. *Eng Anal Bound Elem*, 16(2), 83-92, [https://doi.org/10.1016/0955-7997\(95\)00047-X](https://doi.org/10.1016/0955-7997(95)00047-X), 1995.
- 1090 Sessford, E. G., Strzelecki, M. C., and Hormes, A.: Reconstruction of Holocene patterns of change in a High Arctic coastal landscape, Southern Sassenfjorden, Svalbard. *Geomorphology*, 234, 98-107, <https://doi.org/10.1016/j.geomorph.2014.12.046>, 2015.
- Solomon, S. M., Taylor, A. E., and Stevens, C. W.: Nearshore ground temperatures, seasonal ice bonding, and permafrost formation within the bottom-fast ice zone, Mackenzie Delta, NWT. In *Proceedings of the Ninth International Conference on Permafrost*, Fairbanks, Alaska (Vol. 29, pp. 1095 1675-1680). Fairbanks: Institute of Northern Engineering, University of Alaska Fairbanks. 2018
- Strand, S. M., Christiansen, H. H., Johansson, M., Åkerman, J., and Humlum, O.: Active layer thickening and controls on interannual variability in the Nordic Arctic compared to the circum-Arctic. *Permafrost and Periglacial Processes*, 32(1), 47-58, <https://doi.org/10.1002/ppp.2088>, 2021.
- 1100 Svendsen, J. I., and Mangerud, J.: Holocene glacial and climatic variations on Spitsbergen, Svalbard. *The Holocene*, 7(1), 45-57, <https://doi.org/10.1177/095968369700700105>, 1997.
- Szafraniec, J. E., and Dobiński, W.: Deglaciation rate of selected Nunataks in Spitsbergen, Svalbard—Potential for permafrost expansion above the glacial environment, *Geosciences*, 10(5), 202. <https://doi.org/10.3390/geosciences10050202>, 2020.
- 1105 Tavakoli, S., Gilbert, G., Lysdahl, A. O. K., Frauenfelder, R., and Forsberg, C. S.: Geoelectrical properties of saline permafrost soil in the Adventdalen valley of Svalbard (Norway), constrained with in-situ well data. *Journal of Applied Geophysics*, 195, 104497, <https://doi.org/10.1016/j.jappgeo.2021.104497>, 2021.
- Treat, C. C., and Jones, M. C.: Near-surface permafrost aggradation in Northern Hemisphere peatlands shows regional and global trends during the past 6000 years. *The Holocene*, 28(6), 998-1110 1010, <https://doi.org/10.1177/0959683617752858>, 2018.

- Ulrich, M., Wetterich, S., Rudaya, N., Frolova, L., Schmidt, J., Siegert, C., Fedorov A. N., and Zielhofer, C.: Rapid thermokarst evolution during the mid-Holocene in Central Yakutia, Russia, *The Holocene*, 27(12), 1899–1913, <https://doi.org/10.1177/0959683617708454>, 2017.
- van der Bilt, W. G., D'Andrea, W. J., Werner, J. P., and Bakke, J.: Early Holocene temperature oscillations exceed amplitude of observed and projected warming in Svalbard lakes, *Geophys Res Lett*, 46(24), 14732-14741, <https://doi.org/10.1029/2019GL084384>, 2019.
- van der Bilt, W. G., D'Andrea, W. J., Bakke, J., Balascio, N. L., Werner, J. P., Gjerde, M., and Bradley, R. S.: Alkenone-based reconstructions reveal four-phase Holocene temperature evolution for High Arctic Svalbard, *Quaternary Sci Rev*, 183, 204-213, <https://doi.org/10.1016/j.quascirev.2016.10.006>, 2018.
- Van Everdingen R. V.: Multi-language glossary of permafrost and related ground-ice terms, Boulder, CO: National Snow and Ice DataCenter/World Data Center for Glaciology; Revised January 2005:98, 1998.
- Verruijt, A.: A note on the Ghyben-Herzberg formula, *Hydrolog Sci J*, 13(4), 43-46, <https://doi.org/10.1080/02626666809493624>, 1968.
- Waller, R. I., Murton, J. B., and Kristensen, L.: Glacier–permafrost interactions: Processes, products and glaciological implications. *Sediment Geol*, 255, 1-28, <https://doi.org/10.1016/j.sedgeo.2012.02.005>, 2012.
- Walvoord, M. A., and Kurylyk, B. L.: Hydrologic impacts of thawing permafrost—A review. *Vadose Zone J*, 15(6), <https://doi.org/10.2136/vzi2016.01.0010>, 2016.
- Wohlfarth, B., Lemdahl, G., Olsson, S., Persson, T., Snowball, I., Ising, J., and Jones, V.: Early Holocene environment on Bjørnøya (Svalbard) inferred from multidisciplinary lake sediment studies, *Polar Res*, 14(2), 253-275, <https://doi.org/10.1111/j.1751-8369.1995.tb00693.x>, 1995.
- Weinstein, Y., Rotem, D., Kooi, H., Yechieli, Y., Sültenfuß, J., Kiro, Y., Harlavan, Y., Feldman, M., and Christiansen, H. H.: Radium isotope fingerprinting of permafrost-applications to thawing and intra-permafrost processes. *Permafrost Periglac*, 30(2), 104-112, <https://doi.org/10.1002/ppp.1999>, 2019.
- Williams, P. J., and Smith, M. W.: *The frozen earth: fundamentals of geocryology* (Vol. 306). Cambridge: Cambridge University Press., 1989.
- Yang, B., Bai, F., Wang, Y., and Wang, Z.: How mushy zone evolves and affects the thermal behaviours in latent heat storage and recovery: A numerical study, *Int J Energ Res*, 44(6), 4279-4297, <https://doi.org/10.1002/er.5191>, 2020.

Zhang, T. Influence of the seasonal snow cover on the ground thermal regime: An overview. *Reviews of Geophysics*, 43(4), <https://doi.org/10.1029/2004RG000157>, 2005.

1145 Zhang, N., and Wang, Z.: Review of soil thermal conductivity and predictive models. *International Journal of Thermal Sciences*, 117, 172-183, <https://doi.org/10.1016/j.ijthermalsci.2017.03.013>, 2017.