

Impact of icebergs on the seasonal submarine melt of Sermeq Kujalleq

Karita Kajanto¹, Fiammetta Straneo^{2,3}, and Kerim Nisancioglu¹

¹Department of Earth Science, University of Bergen and Bjerknes Centre for Climate Research, Bergen, Norway

²Scripps Institution of Oceanography, UCSD, CA, USA

³Geophysical Institute, University of Bergen and Bjerknes Centre for Climate Research, Bergen, Norway

Correspondence: Karita Kajanto (karita.kajanto@uib.no)

Abstract. The role of icebergs in narrow fjords hosting ~~marine-terminating~~marine-terminating glaciers in Greenland is poorly understood, even though ~~icebergs provide~~iceberg melt results in a substantial freshwater flux that can exceed the subglacial discharge. ~~Iceberg melt is distributed at depth, contributing to fjord stratification, thus impacting melt and dynamics of the~~
5 ~~Furthermore, the melting of deep-keeled icebergs modifies the vertical stratification of the fjord and, as such, can impact~~
~~ice-ocean exchanges at the~~ glacier front. We model an idealized representation of the high-silled Ilulissat Icefjord in ~~Western~~
~~West~~ Greenland with the MITgcm ocean circulation model, using the IceBerg package to study the effect of ~~icebergs on~~
~~fjord properties~~submarine iceberg melt on fjord water properties over a runoff season, and compare our results with available
observations from 2014. We find the subglacial discharge plume to be the primary driver of the seasonality of circulation,
glacier melt and iceberg melt. ~~Icebergs are necessary to include to correctly understand the properties of Ilulissat Icefjord,~~
10 ~~since they modify~~Furthermore, we find that melting of icebergs modifies the fjord in three main ways: First, icebergs cool and
freshen the water column ~~within over~~within over their vertical extent; Second, ~~icebergs depress~~iceberg melt causes the neutral buoyancy
depth of the plume and the ~~outflow route~~export of glacially modified ~~water~~waters to be deeper; Third, icebergs modify the
deep basin, below their vertical extent, ~~due to both increased entrainment of glacially modified water into the fjord, and iceberg~~
~~modification of~~by driving mixing of the glacially modified waters with the deep basin waters and by modifying the incoming
15 ambient ~~water~~. ~~Furthermore, the depressed neutral buoyancy depth of the plume limits melt to the deep section of the front~~
~~of Sermeq Kujalleq (Jakobshavn Isbræ) even during peak summer, and thus promotes undercutting.~~We ~~waters~~. Through the
combination of cooling and causing the subglacial-discharge-driven plume to equilibrate deeper, icebergs suppress glacier
melting in the upper layer, resulting in undercutting of the glacier face. Finally, we postulate that the impact of ~~icebergs~~
~~submarine iceberg melt~~ on the neutral buoyancy depth of the plume is a key mechanism ~~connecting iceberg mélange and~~
20 ~~linking the presence of an iceberg mélange with~~ glacier calving, ~~independent of mechanical support~~without needing to invoke
mechanical effects.

1 Introduction

Marine-terminating outlet glaciers contribute to approximately half of the mass loss from the Greenland Ice Sheet ~~, and~~
(Goelzer et al., 2020; Mankoff et al., 2020). These glaciers are sensitive to ~~changes in external forcing~~ (Slater et al., 2019; Straneo et al., 20

25 ~~Glaciers terminating in a narrow fjord are controlled by the fjord geometry (Åkesson et al., 2018; Frank et al., 2021) and fjord stratification (Slater et al., 2016; De Andrés et al., 2020). The~~ the conditions in the glacial fjords where they terminate (Straneo et al., 2013; Slater et al., 2019), and the ice-ocean interface remains the main source of uncertainty in the ~~sea-level future sea-level~~ contribution estimates from the Greenland ~~ice-sheet (Goelzer et al., 2020), despite significant effort through modelling, observations and satellite monitoring (Straneo et al., 2019; Catania et al., 2020). The key uncertainties of the field,~~
 30 ~~such as subglacial discharge, shape of the plume and calving have been thoroughly discussed in Straneo et al. (2019). Despite the uncertainties, there is accumulating evidence of the ocean driving retreat of marine-terminating glaciers~~ Ice Sheet (Goelzer et al., 2020). Increased ice loss from the ice sheet, although individual glaciers in Greenland can have very different responses due to local features (Slater et al., 2019; Catania et al., 2020). ~~in turn, leads to increased freshwater discharge into the North Atlantic, with the potential of altering the ocean circulation (Böning et al., 2016) and local marine ecosystems (Meire et al., 2017). Key to~~
 35 ~~understanding the drivers of ice sheet change and the impact on the ocean is understanding water mass transformation and circulation in glacial fjords (Straneo and Cenedese, 2015). Unfortunately, these processes tend to be under-observed because of the challenges in making measurements in these remote and ice-covered regions (Straneo et al., 2019). Thus, fjord models parameterizing the ice-ocean processes have been instrumental in understanding both ocean-driven melting of the glaciers and the export of meltwater (Jenkins, 2011; Sciascia et al., 2013; Carroll et al., 2017).~~
 40 ~~Many~~ Until recently, models have ignored the impact of the many icebergs present in these fjords on the fjord dynamics, meltwater export and glacier melting (Gladish et al., 2015; Carroll et al., 2017). However, many of the fast-flowing, marine-terminating glaciers in Greenland discharge the majority of their ice through the calving of icebergs (Mouginot et al., 2019; Wood et al., 2021). ~~Observations indicate a close connection between dense iceberg melange in front of the glacier and glacier calving. This connection has been interpreted to be due to mechanical support provided by a rigid iceberg melange that~~
 45 ~~suppresses calving (Joughin et al., 2008; Amundson et al., 2010; Burton et al., 2018; Joughin et al., 2020). Meanwhile, there is accumulating observational evidence that icebergs are an important contributor to fjord stratification, and that much of the transformation of ocean waters due to,~~ and recent studies show that iceberg melt can be a dominant freshwater source to the fjord most of the year (Enderlin et al., 2016; Moon et al., 2018). A large fraction of the icebergs melt within the fjord (Mortensen et al., 2020), releasing freshwater below the surface (Moon et al., 2018), with potential impacts both on the vertical
 50 stratification and on the circulation (Hughes, 2022). Subglacial discharge driven buoyant plumes are considered as the key drivers of circulation within glacial fjords (Sciascia et al., 2013; Carroll et al., 2017), and the accompanying high glacier melt rate is significant for the dynamics at the ice-ocean ~~exchanges takes place along the fjord — due to~~ interface (Slater et al., 2016, 2017b, c, 2020). Plumes are sensitive to the water column properties within the fjord, for example, icebergs — rather than directly at which directly impact the plume melt rate and the neutral buoyancy depth of the plume (Jenkins, 2011; Cowton et al., 2015). This
 55 implies that iceberg-induced changes in the hydrography of the fjord have the potential to impact both the direct melt of the glacier front as well as the properties of the plume. Moreover, the height reached by the plume along the glacier terminus (Moon et al., 2018; Mortensen et al., 2020; Muilwijk et al., 2021). Estimates for rapidly calving glaciers indicate that icebergs can be a larger freshwater source than subglacial discharge (Enderlin et al., 2016; Moon et al., 2018), and take up the majority of the heat used for melting in a such fjord (Davison et al., 2020). Thus, irrespective of a mechanical connection,

iceberg-modification-of-the-water-column-provides-a-mechanism-that-enables-icebergs-to-potentially-impact-the-glacier-front has the potential to increase undercutting and thus impact the shape of the terminus, that again can cause further changes in the calving of the glacier front (Slater et al., 2017b, 2021). For marine-terminating glaciers, rigid iceberg mélange is interpreted to provide mechanical support, or buttressing, that suppresses calving (Joughin et al., 2008; Amundson et al., 2010; Burton et al., 2018; Joughin et al., 2020). However, due to the lack of a comprehensive understanding of both calving and iceberg mélange, the dynamics controlling the iceberg mélange and its impact on buttressing and iceberg calving remain speculative.

Recent advances in introducing icebergs in models (Davison et al., 2020, 2022) show that buoyant meltwater from icebergs can drive a circulation within the fjord, and that iceberg melt can take up almost all available heat for melting and significantly freshen the upper layer of the fjord. Davison et al. (2022) find that the response of a fjord basin to iceberg-induced modification is reversed from warming to cooling, when the sill is shallower than the deepest iceberg keels. However, they do not describe the processes causing this. Furthermore, Davison et al. (2022) find that icebergs reduce the overall melt of the glacier front but find little impact of icebergs to a single point-source plume. We find that the dynamics and the interactions between icebergs and the plume merits further study due to the potential significance for the response of marine terminating glaciers to changes in calving.

We-In this study, we investigate how icebergs modify the seasonal stratification and circulation in the fjord, and the implications of this modification to a glacial fjord with a shallow sill. In particular, we are interested in how the presence of icebergs in the fjord impact the melt of the glacier front. We construct a-an idealized model of the shallow-silled Ilulissat Icefjord, which-is-a-high-silled,-rapidly-calving-and-an iceberg-congested fjord in Western-West Greenland. We use-apply the Iceberg toolbox (Davison et al., 2020) within the MITgcm ocean model-to-include-the-impact-of-icebergs,-and-to-investigate and-quantify-their-circulation model (Marshall et al., 1997) to include iceberg melt and quantify its impact on fjord stratification and circulation. We prescribe the seasonal evolution of the subglacial discharge from winter to peak-summer,-the-end-of-summer and characterize the role icebergs play at each phase of the discharge-runoff season. We study the sensitivity of the model to the distribution and draft of the icebergs, and the configuration of the subglacial discharge plume. The results are compared with available observational data from the fjord. Finally, we summarize and discuss the impact the icebergs have on the stability-of-the-glacier-discharge of Sermeq Kujalleq (Jakobshavn Isbræ), and consider the implications for the future response of Sermeq Kujalleq-the glacier in a warming climate.

2 Ilulissat Icefjord

Ilulissat Icefjord (also known as Kangia) is a 50 km long East–West oriented fjord at the eastern edge of Disko Bay, Western-West Greenland (also known as Qeqertarsuup tunua) (Fig. 1). The terminus of the fastest-fastest flowing glacier of Greenland, Sermeq Kujalleq (also known as Jakobshavn Isbræ), is located in the eastern end of the Ilulissat Icefjord. Sermeq Kujalleq is the most rapidly calving glacier of the Greenland Ice Sheet: during the high-discharge years of 2004–2014 its calving rate was estimated to reach over 70-55 Gt/a in the height of summer (Bondzio et al., 2017), leaving-the-fjord-clogged-with

icebergs (Mankoff et al., 2020). The sill at the mouth of Ilulissat Icefjord is relatively high — appr. 250 m — compared to the fjord depth, which is 700–800 m (Morlighem et al., 2017). Iceberg drafts can reach down to 400–500 m (Enderlin et al., 2016), causing these large icebergs to spend substantial amounts of time on the sill, before melting enough to exit over the sill into Disko Bay. ~~This dense melange makes~~ The high calving rate in combination with the high sill leaves the fjord clogged with icebergs, making the fjord difficult to study, as it is inaccessible by boat most of the year, ~~however expendable CTD-campaigns~~ . However, expendable conductivity-temperature-depth-instruments (XCTD) and instrumented seals provide observational data for peak and late summer (Gladish et al., 2015; Mernild et al., 2015; Fenty, I. et al., 2016; Beaird et al., 2017).

Observations indicate that the fjord can be described with three different layers: surface layer (0–50 m), intermediate layer (50–300 m) and the deep basin (300–800 m), see Fig. 2b. The dominant characteristic of the surface layer is that it is cold and fresh, due to melt of the iceberg ~~melange~~ mélange and a smaller contribution from surface runoff (Gladish et al., 2015; Mernild et al., 2015; Beaird et al., 2017; Mojica et al., 2021). The intermediate layer is a weakly stratified layer of glacially modified water (GMW) ~~;~~ — a mixture of ambient water, subglacial discharge and submarine meltwater — where the lower portion of the large icebergs resides (Beaird et al., 2017; Mojica et al., 2021). The deep basin is below the extent of most icebergs, and contains the warmest and most saline water in the fjord (Gladish et al., 2015; Mernild et al., 2015; Beaird et al., 2017; Mojica et al., 2021), with water temperature varying interannually within 1.5–3°C (Gladish et al., 2015; Khazendar et al., 2019). There is little information on the seasonality ~~in of water properties or circulation within~~ the fjord (Mernild et al., 2015), but the conventional idea is that basin water renewal by warm water over the sill takes place only during summer and is driven by the subglacial discharge plume ~~(Gladish et al., 2015; Carroll et al., 2017)~~ (Gladish et al., 2015; Carroll et al., 2017; Khazendar et al., 2019).

The high sill of the Ilulissat Icefjord acts as a barrier in both directions, isolating the basin from the warmest and most saline water in Disko Bay, while blocking large icebergs from leaving the fjord. Both Gladish et al. (2015) and Beaird et al. (2017) find a sharp gradient in the surface layer properties at the sill, as the cold and fresh surface conditions in the fjord switch to the relatively warm summer conditions of Disko Bay. Seasonal profiles from Disko Bay, close to Qeqertarsuaq in Disko Island, roughly 100 km west from the Ilulissat Icefjord, show a strong seasonal signal, reaching down to 300 m depth, with significant warming and freshening during the summer, and slow cooling during winter (Fig. S1) (Greenland Ecosystem Monitoring, 2020). Summer profiles obtained ~~in-front-west~~ of the sill demonstrate a similar strong summer surface warming (Fig. S1) (Beaird et al., 2017).

3 Methods

We use the Massachusetts Institute of Technology general circulation model, MITgcm, which solves the incompressible Navier-Stokes equations with finite volume methods (Marshall et al., 1997). We use an idealized, hydrostatic, high-silled fjord setup in MITgcm (Marshall et al., 1997), illustrated in Fig. 2, based on the Ilulissat Icefjord. The domain is a rectangular East–West oriented fjord that has a high sill at the fjord entrance and a vertical glacier front at the eastern end. The domain is 50 km ~~’s~~ long, 8.5 km wide, and the sill is located at 5 km from the western boundary. The fjord is 700 m deep, the sill 250 m deep and the area ~~in-front-west~~ of the sill representing Disko Bay 400 m. Grid resolution is 312.5*~~400~~*10 m. ~~Even though the~~




Figure 1. Map of Ilulissat Icefjord, indicating the sill (black line) and ~~present-day~~ Sermeq Kujalleq front location in 2014 (blue line). XCTD profile locations marked with red stars and CTD profile location used for boundary condition marked with a ~~green-pink~~ diamond, all obtained in August 2014 (Beaird et al., 2017). Next-to-glacier, mid-fjord and close-to-sill profiles used in result plots are marked with a white outline. Bathymetry and topography are from BedMachine v3 (Morlighem et al., 2017).

$\times 400 \times 10 \text{ m}$ ($\Delta x, \Delta y, \Delta z$). The model is three dimensional, ~~we do not consider Coriolis force, and use the third~~; however, we focus on the along-fjord evolution and do not include rotation. We use the lateral dimension primarily for icebergs and plume width considerations. We run the model for three months with winter conditions, ~~and then run the model for 30 days with each monthly conditions~~ (See Fig. followed by a forward run with varying seasonal forcings (Figs. 2 and S2).

130 The vertical glacier front is represented with the MITgcm IcePlume package (Cowton et al., 2015) that computes melt from the glacier front both with and in the absence of a subglacial discharge plume. The subglacial discharge outlet width — hereafter called plume width — is a key parameter of the model, since it determines the width of the plume and contributes to the melt rate and neutral buoyancy depth (Jenkins, 2011). It is also a key uncertainty of our model, which we will discuss further in the sensitivity experiments in Section 4.4. In 1985, the base of the floating tongue of Sermeq Kujalleq of the time, showed a
 135 single wide channel ~~Motyka et al. (2011)~~ (Motyka et al., 2011). The front has retreated more than 10 km since then, and is now vertical, with two branches, although the southern branch is likely a much larger contributor to subglacial discharge. A large volume of surface runoff, as in the Sermeq Kujalleq catchment, does suggest a high degree of subglacial ~~channeling~~ channelling, and thus a narrow plume (Cuffey and Paterson, 2010). However, Jackson et al. (2017) suggest that plumes with a width of ~~several hundred meters~~ 200 m are a better match ~~to the observations~~ than a point-source plume based on their observations
 140 from Kangerlussuup Sermia, West Greenland. Cavanagh et al. (2017) find that large portions of the surface runoff of Sermeq Kujalleq drain through both shear margins relatively close to the front, which suggests that there could also be subglacial




Figure 2. Model domain and forcings: a) Disko Bay temperature boundary condition from March to August. b) 2D-section of the model domain, grey-blue shading indicates iceberg extent, dotted lines separate vertical layers used in describing the results, light grey shading indicates the OBCS sponge layer where the western boundary is restored to the boundary conditions and the vertical pale-blue-turquoise block to the east indicates the vertical front of Sermeq Kujalleq ('SK'). c) Number of icebergs of each depth at 10 m intervals. d) Monthly subglacial discharge forcing.

discharge ~~beneath whole of the fast-flowing southern branch of the glacier~~ outlets at the lateral margins. The model study by Cook et al. (2021) of the subglacial drainage and discharge from Store Glacier, West Greenland, show that the degree of channelisation can vary significantly within the year, due to the evolution of the subglacial drainage system. The absence of observations of the plume of Sermeq Kujalleq itself, points toward a laterally distributed discharge (Slater et al., 2017a). In the absence of direct information, we assume that there is likely a wider subglacial discharge outlet at the hundred-meter-scale, but potentially also side outlets and laterally distributed sheet discharge during the season.

Here, we choose a 1.2-km-wide sheet plume (Jenkins, 2011) as our default (Indicated with 'P' in experiments *IBP*, *NoIBP*, *IB200P* and *IB400P* in Table 1). ~~For, and for~~ simplicity, the plume width is kept constant during each experiment, ~~although~~ Cook et al. (2021) show that for Store glacier, slightly further north along the west coast of Greenland, plume width can vary significantly during the year due to the evolution of. This is a middle-ground assumption, taking into account the likely variation of the ~~subglacial drainage system~~ degree of channelisation during the season, and potential side outlets, which are all described with a single, wider plume. To account for uncertainty in the plume width, we run sensitivity experiments with two additional plume widths: a wide plume of 4 km (*IB200WP*, *IBWP* and *IB400WP* in Table 1) and a narrow plume of 400 m (*IB200NP*, *IBNP* and *IB400NP* in Table 1). The wide plume corresponds to a situation where there is subglacial discharge along the whole southern branch of the glacier's calving front, while the narrow plume assumes that all runoff is routed into

a single subglacial channel. ~~All plumes are seasonal, with a bell-shaped discharge rate commencing in May and~~ We assume idealized, symmetric Gaussian seasonality of the subglacial discharge from May to November, peaking in August. ~~We assume peak subglacial discharge to take place in August, since surface runoff tends to peak around in Mid-July (?)~~ (Fig. 2d). We acknowledge that a small amount of discharge as is plausible also in winter (Cook et al., 2021); however, for simplicity, we assume zero subglacial discharge during the winter. The maximum volume flux ~~of is set to~~ $1200 \text{ m}^3 \text{ s}^{-1}$ ~~might be slightly underestimated for Sermeq Kujalleq, as Enderlin et al. (2016) estimate, based on estimates by~~ Enderlin et al. (2016) that the peak might reach up to $1200\text{--}1300 \text{ m}^3 \text{ s}^{-1}$. ~~This uncertainty is nevertheless in Sermeq Kujalleq. Nevertheless, uncertainty in the subglacial discharge volume is~~ small compared to the uncertainty ~~of in~~ the plume width.

We choose to impose no ~~separate surface forcing in the fjord, similar air-sea fluxes, similar~~ to Gladish et al. (2015) and Davison et al. (2020), ~~and assume that iceberg melt provides the dominant surface forcing and somewhat disconnects the fjord from atmospheric forcing as our primary focus is on subsurface processes. However, the seasonal surface forcing is partially accounted for by the Disko Bay boundary conditions.~~ We also ignore the contribution from surface runoff, which we expect to mainly have an effect on the surface salinity in the fjord (Mernild et al., 2015). ~~However, the summer warming and freshening of the fjord surface due to atmospheric forcing and runoff is accounted for by the Disko Bay boundary forcing; in the absence of icebergs, surface conditions in the fjord will follow those of Disko Bay.~~ In the model, the western Disko Bay boundary is restored to idealized temperature and salinity profiles representing Disko Bay, using a 3.2 km long sponge layer of the OBCS package (Fig. 2ab). The restoration timescale within the sponge layer ranges from is 17 hours on the outer boundary up to 1 month on the inner boundary, ~~to ensure a smooth change of the ambient conditions~~ (See Fig. S2). ~~The winter boundary condition replicates observed profiles from Disko Bay (Greenland Ecosystem Monitoring, 2020) (see Fig. S1), while the summer profile replicates~~ A volume equal to the subglacial discharge is allowed to exit from the western boundary.

The summer boundary conditions replicate a CTD profile taken in Disko Bay on August 2014, outside of the sill (Beaird et al., 2017) (green pink diamond in Fig. 1). ~~Monthly boundary condition profiles are created by interpolating linearly between~~ The winter boundary condition replicates observed profiles close to Qeqertarsuaq, further out in Disko Bay in 2018, available down to 150 m depth (Greenland Ecosystem Monitoring, 2020) (Fig. S1), below which we assume that the winter and summer profiles conditions merge. As a consequence, seasonality is applied only above sill depth, and water properties at the sill depth are constant throughout the experiments (Figs. 2a). ~~However, S1). In addition, we adjust temperature and salinity at sill depth to values the highest value observed at depth in the fjord (Beaird et al., 2017), and assume seasonality applies only above sill depth to ensure our boundary conditions allow for a sufficiently saline inflow to the fjord. Monthly boundary conditions are~~ created by interpolating linearly between the winter and summer profiles (Fig. S12a). We initialize the fjord above sill depth with the winter profile, and constant temperature and salinity below sill depth, leading to no density gradient over the sill (Fig. S2). We do not include sea ice formation, since we do not expect it to play a ~~big role in spring and summer~~ major role during the melt season.

We implement the iceberg forcing with the IceBerg package of MITgcm, presented in Davison et al. (2020). In this package, the fjord is filled with a randomly created block iceberg distribution following a power law of exponent -2.1, derived for Ilulissat Icefjord icebergs (Enderlin et al., 2016). We define the minimum and maximum iceberg depth and the surface area coverage.

~~Melt and Heat uptake due to iceberg melt and the accompanying~~ negative salinity flux are then computed from each iceberg face, similar to the sub-grid approach of Cowton et al. (2015), resulting in cell-averaged fluxes that force the ocean model. The effect of ~~drift-velocity~~ on melt rate is taken into account by considering the flow field in each computational cell with respect to the iceberg face orientation (Davison et al., 2020). ~~The effect of iceberg drift velocity — the average water velocity from the fjord surface to the iceberg keel depth — on iceberg melt is included, noting that there is little rigid mélange observed in 2014~~ (Joughin et al., 2020). The iceberg distribution remains fixed during the calculation, and the volume they uptake is ~~reduced from the volume of accounted for in~~ each computational cell occupying icebergs. Other physical aspects, such as iceberg size decrease due to melt, or the sub-grid scale flow ~~network-icebergs-create~~, icebergs create are not included. Surface melt of icebergs above sea level is not included, nor other deterioration mechanisms than submarine melt.

Enderlin et al. (2016) find for Ilulissat Icefjord that there often is a significant number of icebergs with estimated draft deeper than 200 m, but rarely many icebergs with drafts deeper than 400 m. Our default iceberg distribution covers uniformly 60% of the surface area of the fjord from the highest point of the sill to the glacier (grey area in Fig. 2b), with an iceberg-depth distribution from 10–300 m (Fig. 2c, experiments *IBP*, *NoIBP*, *IBWP* and *IBNP* in Table 1). To account for uncertainty in the iceberg distribution, we run two sets of sensitivity experiments with deeper and shallower icebergs (Fig. 10c). The deeper iceberg distribution has a maximum depth of 400 m and a high surface coverage of 90%, which is needed to fit the power law because deeper icebergs mean also more small icebergs (*IB400WP*, *IB400P* and *IB400NP* in Table 1). The shallow distribution has a maximum depth of 200 m and a surface coverage of 60%, leading to a larger number of small icebergs than in the default distribution (*IB200WP*, *IB200P* and *IB200NP* in Table 1). For a ~~comperhensive~~ comprehensive list of all model parameters, see Table S1.

Table 1. Experiment naming and key parameters

Model name	IB max depth (m)	IB surface coverage (%)	Plume width (m)
Control without icebergs			
<i>NoIBP</i>	-		1200
Ilulissat Icefjord with icebergs			
<i>IBP</i>	300	60	1200
Sensitivity experiments			
<i>IB200P</i>	200	60	1200
<i>IB400P</i>	400	90	1200
<i>IBNP</i>	300	60	400
<i>IBWP</i>	300	60	4000
<i>IB200NP</i>	200	60	400
<i>IB200WP</i>	200	60	4000
<i>IB400NP</i>	400	90	400
<i>IB400WP</i>	400	90	4000

4 Results

We run our idealized Ilulissat Icefjord model, *IBP*, with monthly varying Disko Bay temperature and salinity, ~~monthly-increasing subglacial-discharge~~, subglacial discharge and a constant iceberg distribution (See Fig. 2) over a synthetic runoff season described in Sect. 3. We compare ~~this-the results~~ to a control experiment with no icebergs, *NoIBP* — otherwise identical to *IBP* — to investigate the impact icebergs have on fjord properties, circulation and glacier melt rate. Our experiments start from the prescribed winter conditions of the fjord (Fig. 2a), before the onset of Disko Bay surface warming and the increase in subglacial discharge, ~~and we present results at the end of each month~~. We present results to early season (May–June), peak (July–August–September) and late season (October–November).

4.1 Seasonality without icebergs

4.1.1 Winter and early season

We consider first the seasonality in Ilulissat Icefjord in the control experiment, *NoIBP*, without icebergs but including the plume. Above the sill, winter conditions in the fjord are dominated by the cold Disko Bay boundary condition. ~~Below~~, while below the sill, properties remain virtually constant since there is ~~no~~ little circulation in the deep basin (Figs. 3a and 4a, 4a and S2). In the model, spring starts in April with a warming and freshening in Disko Bay (Figs. 2a and S1). ~~Warming in Disko Bay progresses throughout spring~~, increasing the surface temperature in the fjord (Fig. 3a–eb). In May, the runoff season starts with a small subglacial discharge (Fig. 2d). ~~This subglacial~~ Subglacial discharge creates a buoyant plume that ascends vertically along the glacier front, until it reaches neutral buoyancy relative to the fjord water properties (red lines in Fig. 4b–e, Fig S4). ~~After reaching the neutral buoyancy depth, the high vertical velocity transports the plume higher, further mixing with the fjord water, until the plume~~), and transforms to horizontal outflow of GMW (black arrows in Figs. 3 and 4). ~~In May, the GMW outflow takes place at 230 m depth, partly exiting which is compensated by inflow over the sill (Fig. 4b). The neutral buoyancy depth is at 225 m in May and at 105 m in June. Due to the deep neutral buoyancy depth in the early season, the GMW outflow originating from the plume partly exits over the sill, partly mixing back into mixes and returns with the inflow to the basin (Fig Figs. 4ab and S7), slightly cooling and freshening the deep basin (Fig. 8). This mixing back into the deep basin at the sill is defined as sill-driven reflux in Hager et al. (2022).~~

~~As the subglacial discharge volume increases through June and July, the plume rises higher, and more of the GMW outflow is able to exit over the sill. The plume melt rate is low during the early season due to the small volume of subglacial discharge, and plume melt is vertically limited due to the relatively deep neutral buoyancy depth (Fig. 4c–d). However, there is still some sill-driven reflux of the GMW outflow in June, and to a lesser extent in July. The subglacial discharge is largest in August, resulting in the shallowest GMW outflow at 0–150 m (Fig. 5a,b).~~

4.1.2 Peak season

During the peak season, subglacial discharge volume is high, the plume rises up to 75 m depth (Fig. 4e). This, and the GMW outflow is able to exit over the sill, with a compensating inflow bringing Disko Bay water into the deep basin (Fig. 4e). The evolution of the melt rate at the glacier front follows the evolution of the plume: a weak plume is associated with a low melt rate and a limited vertical extent of the plume, while in peak summer the plume melt rate reaches 5.1 m d^{-1} , and melt extends up to 40 m depth (Fig. 5). Melt rate is small outside of the plume, resulting in a significantly smaller horizontally-averaged melt rate (Fig. 5).

c). Since the GMW outflow exits over the sill in August during peak season, modification in the deep basin, below the 26.9 kg m^{-3} isopycnal is the result of early season entrainment sill-driven reflux (Figs. 3e-c and 7). This modification causes the deep basin to be early season refluxing causes a cooling of $0.2 \text{ }^{\circ}\text{C}$ colder than Disko Bay water at sill depth in the deep basin by peak summer (Figs. 3a-d and Fig. c and 8). In the surface and intermediate layer, cooling due to the GMW reaches Meanwhile, the surface layer is $2 \text{ }^{\circ}\text{C}$ in August, cooler compared to the Disko Bay temperature surface layer, due to mixing with plume water. Overall, glacial modification leads the entire fjord to be almost uniformly $3 \text{ }^{\circ}\text{C}$ by August in peak summer (Fig. 3e-c). The freshwater flux into the fjord consists of 96% of subglacial discharge, and only 4% out of melt from the glacier front (Fig. 6). Plume melt rate reaches up to 5.1 m d^{-1} in peak summer and extends vertically close to the surface (Fig. 5c-e).

4.1.3 Late season

During the late season the subglacial discharge decreases again, and the surface of Disko Bay starts to cool down (Fig. 2a,d). The deep basin has freshened due to the early-season refluxing of GMW, and density-driven renewal of deep basin water begins to dominate (Fig. 4d). Due to smaller subglacial discharge, the plume reaches neutral buoyancy deeper, and the correspondingly deeper GMW outflow again causes refluxing at the sill (Figs. 4d, S7). In spite of the refluxing, the basin starts to return to more saline conditions, (Figs. 3d and 8). Plume melt rate also decreases as the subglacial discharge decreases, both in magnitude and vertical extent. However, the early and late seasons are not symmetrical, since the changed stratification of the fjord slightly decreases the maximum melt rate, and the neutral buoyancy depth of the plume, from 105 m in early season (June) to 155 m in late season (October) (Figs. 3d and 4d). However, the cold surface and intermediate layer conditions in the early season compared to the warm conditions in the late season balance out the differences in melt rate (Fig. 5).

4.2 Impact of icebergs on the seasonality

4.2.1 Winter and early season

Introducing icebergs to the upper 300 m within the fjord leads to significant changes in both water properties and circulation. Since icebergs extend to the warm water layer throughout the year, there is a substantial freshwater flux from the icebergs, also during winter (Fig. 6). This fresh meltwater drives a slight weak mixing within the intermediate layer and surface during winter (Fig. 4f-e), which leads to no distinct thermocline, and a notable freshening within the extent of the icebergs (Figs. 3f and 7). The stronger wintertime circulation also increases the frontal melt of the glacier to a total flux of $4.5 \text{ m d}^{-3} \text{ s}^{-1}$, compared

to $0.7 \text{ m-d}^3 \text{ s}^{-1}$ in *NoIBP*, although the freshwater flux from glacier melt is small compared to iceberg melt (Fig. 6). ~~Spring surface warming in Disko Bay drives iceberg melt in-~~

275 ~~Outflow of iceberg meltwater within the upper 100 m prevents warming within the surface layer, which keeps the surface cold and causes slight freshening of the fjord, which remains cold and increasingly fresh (Figs. 3f-j, 4f and 8), while also creating outflow in the upper 100 m (Fig. 4e-f). This additional outflow increases the~~ is compensated by increased inflow over the sill compared to *NoIBP* (Fig. 4b,g).

~~As the subglacial discharge season starts in May, the wintertime cooling and freshening caused by icebergs affects f). During the early season, wintertime changes in the stratification due to icebergs impact the neutral buoyancy depth of the plume: The~~
280 ~~plume equilibrates 120 m-80-120 m (May/June) deeper in the water column than in NoIBP (Fig. 5e Figs 4b,f and 5a,b). The correspondingly deeper GMW outflow does not exit over the sill, but instead GMW outflow starts out correspondingly deeper, but gets further modified by icebergs along the fjord, and reaches the sill at a comparable depth to NoIBP, and refluxes into the deep basin. These dynamics affect (Figs. 4f, S7). The changes in stratification and the neutral buoyancy depth of the plume are reflected in the plume melt rate, which reaches the same peak; the peak melt rate value at depth, but is limited to below~~
285 ~~300 m is the same as in NoIBP, but vertical extent is limited (Fig. 5a). In June and July when b).~~

4.2.2 Peak season

~~During peak season, once subglacial discharge increases, the plume extends higher in the water column (4h,i). However, the plume remains 50 m deeper, and the GMW outflow close to the glacier is 50-100 m deeper than in, although remaining 50 m deeper than in NoIBP (Fig. 4c,g). The modelled peak GMW outflow takes place within the 26.3 kg m^{-3} and 26.5 kg m^{-3}~~
290 ~~density layers, while the surface above 26.3 kg m^{-3} is dominated by iceberg melt (Figs. 7, and 3g). The relatively warm and rapid GMW outflow promotes iceberg melt, peaking at $1540 \text{ m}^3 \text{ s}^{-1}$ (Figs. 6 and S3), causing the surface and intermediate layers to cool significantly compared to NoIBP. Furthermore, icebergs melt along (Figs. 3c,g and 8). Meanwhile, the GMW outflow route, due to both the temperature and velocity associated with the outflowing plume (Figs. 6 and S3), leading to a colder intermediate layer than in gets further modified by iceberg melt along the fjord and rises up to exit the fjord at a depth~~
295 ~~similar to NoIBP (Figs. 3h-i and 8 Fig. 4c,g and S7). The additional outflow due to iceberg melt is compensated by stronger inflow over the sill, increasing the peak value of up-fjord volume flux over the sill by 50%, from $1 * 10^5 \text{ m}^3 \text{ s}^{-1}$ in NoIBP to $1.5 * 10^5 \text{ m}^3 \text{ s}^{-1}$ in IBP. The buoyancy-driven circulation due to iceberg melt draws a fraction of the inflowing water towards the base of the icebergs to compensate for the outflow, rather than deeper into the basin (Fig. 4g).~~

In addition to the ~~modification of~~ surface and intermediate layers ~~modification~~, icebergs modify the deep basin through two
300 separate processes: by ~~entrainment of mixing, or refluxing, of the~~ GMW outflow into the inflowing water to the basin, and by iceberg-modification of inflowing ambient water. While the GMW is also iceberg-modified, we call the incoming ambient water modified by iceberg melt **iceberg-modified ambient water, IMAW**, to separate these two processes of ambient water modification: IMAW has not been in contact with the glacier terminus or the plume, as is the case for GMW. IMAW is ~~always~~
305 ~~present in the deep basin always~~ when there is inflow over the sill and icebergs along the inflow route, whereas GMW will mix into the deep basin when outflow is too deep to exit over the sill. Both processes contribute to cooling and freshening of the

deep basin throughout the season (Fig. 3g–j,g and 8), which is also reflected in a slight decrease of the plume melt rate at depth. However, during peak summer, GMW exits the fjord completely, and thus all modification of the incoming ambient water at this time is due to iceberg melt (Fig. 5b–e).

The plume reaches higher up the watercolumn in August — still 50 m deeper than in *NoIBP* — and the GMW outflow takes place at 100 m depth, with some slight entrainment into the inflowing waters (S7). The inflow region is located below approximately 200 m depth and below isopycnal 26.7 kg m^{-3} close to the sill and 26.5 kg m^{-3} next to the glacier, and is near-parallel to the melt line, since it contains only IMAW (Fig. 4h). The fjord has a cold surface, cool intermediate layer and slightly cooled deep basin water (Fig. 3j). Meanwhile, relatively warmer plume water reaching the icebergs drives iceberg melt, with a peak melt value of $1540 \text{ m}^3 \text{ s}^{-1}$ (Fig. 6). The increase in iceberg melt from July to August is primarily due to an increase in drift-induced melt, as temperature does not increase (Figs. 3i–j and S3). The additional outflow due to iceberg melt is compensated by a stronger inflow over the sill, increasing the volumetric flow rate over the sill by 50%, from $1 * 10^5 \text{ m}^3 \text{ s}^{-1}$. Below 26.8 kg m^{-3} the basin contains IMAW mixed with early season GMW.

The melt rate experienced by the glacier front reflects the iceberg-induced changes in the fjord properties: The deeper neutral buoyancy depth of the plume limits the vertical extent of plume melt, while the iceberg-melt induced cooling of the surface and intermediate layers decrease melt also outside of the plume. The modification of the basin water is reflected in the plume melt rate, as cooler and fresher deep basin water entrains into the plume. This is seen as the deviation of plume melt rate of *IBP* from *NoIBP* upwards from the depth where iceberg modification extends each month (Fig. 5). This results to an overall reduction of the freshwater flux from direct glacier melt in August from $66 \text{ m}^3 \text{ s}^{-1}$ in *NoIBP* to $1.5 * 10^5 \text{ m}^3 \text{ s}^{-1}$ in *IBP*. The iceberg-melt driven circulation directs more of the inflowing water towards the base of the icebergs, rather than deeper into the basin.

4.2.3 Late season

Once the subglacial discharge decreases, density-driven inflow starts to dominate in *IBP*, as in *NoIBP* (Fig. 4jh). The modelled August GMW outflow takes place within the 26.3 kg m^{-3} and 26.5 kg m^{-3} density layers, while the surface above 26.3 kg m^{-3} is dominated by iceberg melt (Fig. 7 and Fig. 3j) neutral buoyancy depth of the plume is deep, 235 m, but GMW outflow rises along the fjord due to iceberg modification to mostly exit the sill, with some refluxing into the deep basin (Figs. 4h and S7). The inflow region is located below approximately 200 m depth and isopycnal 26.5 kg m^{-3} , and the inflow can be further divided into three vertical layers: 26.5 – 26.7 kg m^{-3} is the rapid inflow region at the base of the icebergs, containing IMAW mixed with entrained GMW outflow water, 26.7 – 26.8 kg m^{-3} surface and intermediate layers are dominated by outflow of glacial meltwater (Fig. 4h). The freshwater flux from the icebergs does not decrease symmetrically compared to the increase in spring (Fig. 6), as it is $1100 \text{ m}^3 \text{ s}^{-1}$ contains only IMAW, and water below 26.8 kg m^{-3} in October compared to $940 \text{ m}^3 \text{ s}^{-1}$ contains IMAW mixed with early season GMW.

To sum up, icebergs modify the fjord properties and circulation in three main ways: Firstly, by cooling and freshening the surface and intermediate layers through iceberg melt; Secondly, by causing the plume and GMW outflow to be deeper; Thirdly, by modifying the deep basin water through iceberg modification of inflowing ambient water (IMAW), and entrainment of

340 ~~GMW into the inflow into s^{-1} in June with the same subglacial discharge. This is due to higher temperatures in the surface layer, where most of the icebergs are located (Fig. S3). As with *NoIBP*, the change in the deep basin~~ ~~These changes impact the melt rate by reducing the melt rate at depth, due to the cooling of the deep basin, and by the limited vertical extent due to the plume being deeper~~ properties causes a decrease in the October and November maximum plume melt rates, compared to May and June in the early season (Fig. 5). ~~Thus, iceberg-modification of the fjord will cause less melt of the glacier, and the~~
 345 ~~melt will be limited to the deep part of the glacier front (Fig. 11a,b,f,g).~~

4.3 Comparison to observations

~~The observed~~ The observed peak summer temperature in the fjord shows a cold and fresh surface layer, cooled intermediate layer water and a slightly modified deep basin (Figs. 3m-i and 8) — which is consistent with the results of experiment *IBP* (Fig. 3j-h). In the observations, the ~~GMW outflow is mostly located~~ contribution of runoff is most significant between isopycnals
 350 26.3 kg m^{-3} and 26.9 kg m^{-3} , corresponding to a relatively narrow depth range of approximately 100–200 m (Figs. 3m-i and 7). ~~Even though the density of the GMW outflow is different in the observations and in *IBP*, the deepest extent of a significant contribution of GMW outflow is approximately 200 m for both in the mid-fjord location (Fig. 8).~~ Below 26.9 kg m^{-3} , the
~~observed deep basin is~~ vertical changes in the observed properties are dominated by melt, following the melt line almost perfectly with negligible contribution of surface runoff, except for the profile directly next to the glacier (Fig. 7). We interpret
 355 this to indicate a large contribution of IMAW rather than GMW, ~~due to the similarity to the modelled melt-dominated inflow region of $26.5\text{--}26.7 \text{ kg m}^{-3}$.~~ IMAW is, by definition, formed along the inflow route into the fjord. ~~Thus, and thus,~~ a layer in the fjord filled with purely IMAW would be a layer of inflow towards the glacier. ~~This suggests that~~ In *IBP*, isopycnal 26.5 kg m^{-3}
~~separates in- and outflow regions in fjord, away from the sill. Although the observed properties are denser than in the model, we interpret that the shift from runoff-dominated to melt-dominated properties at 26.9 kg m^{-3} in the observations could~~
 360 distinguish between inflow and outflow in the fjord. ~~Furthermore, following a similar reasoning, we interpret $26.3\text{--}26.9 \text{ kg m}^{-3}$ in the observed profile as the GMW outflow. Despite the density difference between *IBP* and the observations, the separation between inflow and outflow take place at approximately 200 m depth for both (dashed horizontal line in Fig. 8).~~

To sum up, icebergs modify the fjord properties and circulation in three main ways: Firstly, by cooling and freshening the surface and intermediate layers through iceberg melt; Secondly, by causing the neutral buoyancy depth of the plume and the
 365 ~~GMW outflow to be deeper; Thirdly, by modifying the deep basin water through iceberg modification of inflowing ambient water (IMAW) and refluxing of GMW into the inflow to the deep basin. These changes to the water column properties reduce the maximum submarine melt rate of the glacier front by up to 10% in the deep basin and limit the vertical extent by 40–100 m, depending on the season (Fig. 5). Thus, iceberg-modification of the fjord will cause less submarine melt of the glacier, and the melt will be limited to the deep part of the glacier front (Fig. 11).~~

4.4 The relative impact of plume width and iceberg distribution on fjord properties

We run ~~three different plumes and three different iceberg distributions~~ simulations using three plume/outlet widths and three
 iceberg distributions (Table 1) from winter to peak summer in order to investigate the sensitivity of our results to plume width




Figure 3. Along-fjord temperature sections for Ilulissat Icefjord with (f-j-h), *IBP*) and without icebergs (a-e-a-d), *NoIBP*), from onset of spring for winter (April/March) to, early-season (June), peak summer (August) and late season (October), as a snapshot at along the end centreline of each month the fjord. ki) observed temperature section from XCTD data in August 2014 (Beaird et al., 2017), interpolated from XCTD locations are-marked in Fig 1. Red contours mark the isopycnals of 26.3 kg m^{-3} , 26.5 kg m^{-3} , 26.7 kg m^{-3} and 26.9 kg m^{-3} . Black arrows indicate the centerline-centreline of the outflow of glacially modified water from the plume (See also Fig. 4).




Figure 4. Snapshots of the horizontal flow rate along the fjord from the start of the subglacial discharge season in May to for winter (March), early (June), peak summer in (August, a–d) and late season (October), without icebergs (NoIBP), and e–h) and including icebergs (IBP, e–h). Black arrow indicates the centerline-centreline of the GMW outflow from the plume, and grey contours streamlines of the horizontal flow rate at $0.02 \text{ m}^3 \text{ d}^{-1}$ intervals. Red horizontal line marked on the pale blue glacier front indicates the neutral buoyancy depth of the plume.




Figure 5. Seasonality of ~~the~~ frontal melt of the glacier from May to August as both plume melt rate for *NoIBP* (dashed line), and *IBP* (solid line), and ~~also~~ width-averaged frontal melt rate throughout the runoff season for both *NoIBP* (dashed line, dotted line), and *IBP* (solid line, shaded area).




Figure 6. Modelled seasonality of ~~the~~ freshwater volume flux-fluxes into Ilulissat Icefjord from three different sources: icebergs (*IBP*, purple), subglacial discharge forcing (blue line) and glacier melt for *NoIBP* (dashed black line) and *IBP* (pale blue). Shadings indicate the range covered by the sensitivity experiments (see Table 1 and Sect. 4.4).




Figure 7. August potential temperature versus salinity next to the glacier (brown), mid-fjord (orange) and close to the sill (yellow), for observations (circles) (Beaird et al., 2017), *IBP* (solid lines) and *NoIBP* (dashed lines), with runoff and melt lines (grey lines) and isopycnals (dotted grey lines). Grey dotted contours indicate isopycnals at 0.1 kg m^{-3} intervals, with 26.3 kg m^{-3} , 26.5 kg m^{-3} , 26.7 kg m^{-3} and 26.9 kg m^{-3} indicated in black (Also plotted in Fig. 3).

and iceberg distribution (Table 1). Comparisons of temperature, circulation and melt rate demonstrate the separate contributions of the plume and iceberg distribution, and also the separate impact GMW and IMAW have on the deep basin properties (Figs. 9 and 10). Plume width is the primary controlling factor of the vertical extent of the plume. ~~Changes in the depth of the GMW outflow determine the circulation of the fjord, which are reflected in the fjord temperature and maximum melt rate of the plume, which impacts fjord circulation and water properties.~~ Decreasing plume width ~~creates a more concentrated plume that entrains less~~ reduces the volume of deep basin water entrained into the plume, and ~~The plume~~ thus rises higher in the water column, ~~transporting heat efficiently up and exports GMW closer to the fjord surface~~ (Fig. 9). ~~This concentrated plume leads to a higher~~ The shallower and more concentrated outflow of GMW, ~~which is less prone to entrain into inflowing water leads to reduced refluxing~~ at the sill, leading to a less modified deep basin (Fig. 9c,f,i). For the wide plume, on the other hand, ~~the total entrainment of the~~ deep basin water ~~entrainment~~ into the plume is ~~more efficient~~ larger, because of the greater surface area




Figure 8. Mid-fjord temperature and salinity profiles for ~~April~~ March (winter), June ~~and~~ (early season), August (peak season) and October (late season) for both IBP (solid lines), NoIBP (dashed lines). Grey circles indicate the XCTD profiles in August 2014 (Beaird et al., 2017). The dotted horizontal line marks 195 m depth, which coincides in IBP with isopycnal 26.5 kg m^{-3} and the inflow/outflow border in August, and in the observations with isopycnal 26.9 kg m^{-3} . Mid-fjord location is indicated with a red star and vertical dotted lines in Fig. 3 and in Fig. 1, and as an outlined red star in Fig. 1.

o the plume, causing the plume to remain deep, leading to entrainment of GMW reach neutral buoyancy deeper in the water column. This leads to refluxing of the GMW outflow at the sill — even during peak summer — and significant modification of the deep basin properties (Fig. 9a,d,g). While the peak plume melt rate for the wide plume is less than half that of the narrow plume, the horizontally averaged melt rate is significantly higher doubled (Fig. 10).

Iceberg depth impacts directly the properties of the intermediate layer and the deep basin, as the extent of the cooling and freshening increases with increasing iceberg depth (Fig. 9). These changes are This modification of the deep basin is a result of the combined effect of increased contribution of IMAW with iceberg depth, and the entrainment-refluxing of increasingly iceberg-modified GMW. Icebergs depress the neutral buoyancy depth of the plume, the more the wider Furthermore, changes in the properties of the deep basin water are reflected in the properties of the plume, as more iceberg-modified ambient water is entrained into the plume the plume becomes correspondingly cooler and fresher. This is reflected in a lower plume melt rate (Fig. 10a, grey shaded areas compared to lines) and in a cooler GMW outflow and thus intermediate layer (Fig. 9).

The neutral buoyancy depth of the plume is slightly impacted by the change in the deep basin properties, but more so of the iceberg-induced modification of the intermediate layer. Iceberg distributions *IB* and *IB400* have similar number of icebergs within the top 300 m (Fig. 10c), and thus the differences in the neutral buoyancy depth are small between these two iceberg distributions. *IB200*, however, has approximately twice as many icebergs within the top 200 m as the other two distributions, leading to the neutral buoyancy depth being depressed by 20 m more than for the default and narrow plumes. The other two iceberg distributions in *IB200P* and *IB200NP*. On the other hand, *IB200WP* has less iceberg modification of the plume than the other two distributions, since the neutral buoyancy depth is below the maximum iceberg extent ('WP' lines in Fig. 10a). Overall, this indicates that the number of icebergs extending down to the expected neutral buoyancy depth imposes a stronger control than the maximum iceberg depth. Iceberg-induced deep basin cooling decreases the maximum plume melt rate at depth, but this effect is small causes only a small difference between the iceberg distributions (Fig. 10a).

The sensitivity experiments demonstrate the separate contributions of IMAW and GMW, as IMAW contribution in the deep basin increases with iceberg depth, and GWM-GMW contribution decreases when decreasing the plume width. *IB400NP* has a narrow plume, causing GMW to exit right below the surface layer, leaving the deep basin unmodified by the GMW, except for a small early-season contribution. However, since *IB400NP* has deep icebergs, the resulting deep basin has a significant IMAW contribution (Fig. 9i). Conversely, *IB200NP* has icebergs shallower than the sill depth, leading to no IMAW and very little early-season GMW in the deep basin (Fig. 9c). *IB200WP*, meanwhile, has similarly negligible IMAW contribution and iceberg-modification of the entraining GMW (Fig. 9a); however, the overall entrainment-mixing of the GMW to the basin is significant due to the deep neutral buoyancy depth of the wide plume, leading which leads to significant modification of the deep basin with little contribution from the icebergs. The significance of early-season processes to the summer temperature profile is clearly demonstrated by *IBNP*, where the early-season iceberg-modified water is located at 400–600 m depth, below less modified water inflowing during peak summer (Fig. 9f). These results show that — while the plume is a determining factor for the circulation in the fjord — increasing iceberg depth increases modification in the deep basin.

5 Discussion

5.1 Iceberg modification in Ilulissat Icefjord

Our results show that icebergs must be included in order reproduce the realistic water column properties of Ilulissat Icefjord (Figs. 8, 7). Icebergs cool and freshen the entire water column of the fjord, most intensely in the top 50 m, which hosts a large number of small icebergs. Our simulation including icebergs matches well with the observed temperature profile in Ilulissat Icefjord in August 2014 (Fig. 3j-mg.i), which is typical for Ilulissat Icefjord in peak summer (Gladish et al., 2015; Fenty, I. et al., 2016; Beaird et al., 2017; Mojica et al., 2021). We find that iceberg melt-modification is the main source of cooling in the surface and intermediate layers of the fjord, while the combined effect of subglacial discharge and frontal melt of the glacier only account for approximately 25% of the simulated cooling (Fig. 8a); The iceberg-induced cooling and freshening reaches up to 4 °C and 0.7 PSU in peak summer (Fig. 8). This matches well with the cooling and freshening of 5 °C and 0.7 PSU simulated for Sermilik Fjord in southeastern Greenland by Davison et al. (2020) with a similar iceberg




Figure 9. August along-fjord sections of horizontally averaged temperature for all sensitivity experiments with three different plumes (columns), and three different iceberg distributions (rows). See Table 1 for experiment naming. Black arrow indicates the centerline-centreline of the GMW outflow from the plume, and grey contours a snapshot of streamlines of the horizontal flow rate at $0.02 \text{ m}^3 \text{ d}^{-1}$ intervals.

~~distribution~~ similarly dense iceberg distribution, as well as Davison et al. (2022) studies in an idealized fjord. Comparison of *IBP* and *NoIBP* to observations in TS-space further highlights the contribution of iceberg melt (Fig. 7). The small contribution of subglacial discharge and direct glacier melt ~~is to the cooling of the fjord~~ supported by findings from other West Greenland fjords, which show that the majority of glacial modification takes place along the fjord — through, for example, iceberg melt — while direct melt from the glacier terminus and subglacial discharge are ~~a small contributor~~ small contributors (Mortensen et al., 2020; Muilwijk et al., 2021). Our model slightly underestimates the surface cooling and freshening, which we assume to be due to the exclusion of runoff to the surface of the fjord (Mernild et al., 2015), or possibly due to underestimated contribution from small icebergs by the power law size distribution (Rezvanbehbahani et al., 2020).

We find that the iceberg-induced cooling and freshening in the intermediate layer depresses the neutral buoyancy depth of the plume and the outflow route of the GMW (~~illustrated~~ illustrated in Fig. 11). This is supported by Beaird et al. (2017) who find




Figure 10. August vertical melt rate profiles for a) plume melt and b) horizontally averaged frontal melt for the nine iceberg experiments with three different iceberg distributions (colours, see panel c) for colour labels), and wide (WP), default (P) and narrow plumes (NP) (semi-dashed, solid, and dotted lines respectively). See Table 1 for experiment naming. The three different grey shadings in panel a) indicate the vertical melt rate profile of each of the three plumes without icebergs. c) shows the number of icebergs extending to a certain depth at 10 m intervals in the three used iceberg distributions.

the water above 250 m close to the glacier to be anomalously cold compared to what is predicted by plume theory, indicating a depressed plume. They speculate that this mismatch is due to the impact of icebergs, which our model results confirm. Previous estimates of the neutral buoyancy depth of the plume of Sermeq Kujalleq in peak summer range from 350 m depth to

440 the surface (Jenkins, 2011; Gladish et al., 2015; Carroll et al., 2016), depending on the subglacial discharge flux, ambient water properties and ice front shape. Truffer and Motyka (2016) briefly speculate on the possibility that the ice ~~melange-mélange~~ could mechanically prevent the plume to reach the surface in the Ilulissat Icefjord. However, since the ~~melange-mélange~~ is not rigid during peak summer (Joughin et al., 2020), we argue that the plume is deep due to the ~~iceberg-duced-iceberg-induced~~ stratification, rather than a mechanical effect. A similar phenomenon but at much smaller scale was observed in the nearby
445 Saqqarlep fjord, where extreme melt caused surface freshening and thus prevented the plume from reaching the surface, as it normally does in this location (De Andrés et al., 2020).

~~Davison et al. (2022) study the impact of icebergs on plume melt in an idealized fjord setup, but do not find significant changes in plume temperature and melt rate. They study a 500 m fjord with a single point-source plume, which has limited entrainment of ambient water into the plume, and is thus relatively unaffected by iceberg-modification of the fjord water (Fig. 9 in (Davison et al., 2022)). Their plume reaches the surface in all their different iceberg distributions, but since they do not discuss stratification or buoyancy, it is not clear if the neutral buoyancy depth changes in their different iceberg distributions. However, since Jackson et al. (2017) suggest that wide plumes match better with observations, and observations by Beaird et al. (2017) support our model results, we conclude that a plume with a width on the hundred-meter-scale is more realistic.~~

455 The depressed plume leads to enhanced basin water modification due to increased ~~entrainment of~~ refluxing of the GMW outflow into inflowing Disko Bay water, as not all outflow is able to exit over the sill (FigFigs. 11b, S7). However, icebergs also modify directly the inflowing ambient water from Disko Bay, creating iceberg-modified ambient water (IMAW) that has not had any contact with the glacier front or the plume. Both GMW and IMAW are typically present in a high-silled fjord with a ~~marine-terminating-marine-terminating~~ glacier, such as Ilulissat Icefjord; however, their ~~origin-origins~~ are different. Thus,
460 even if the GMW outflow is able to completely exit the fjord, the deep basin water will still be modified by IMAW, as long as there are icebergs present along the inflow route. Modification of the deep basin of Ilulissat Icefjord is supported by Beaird et al. (2017), who find the fjord to be glacially modified to at least 600 m depth. We specify that in the deep basin the modified water is mostly IMAW, rather than GMW (Sec. 4.3). This interpretation is different from Gladish et al. (2015), who describe the basin to be filled with Disko Bay water at sill depth, with no significant glacial modification. Since their model lacks both
465 icebergs and seasonal considerations, all GMW will exit the fjord, leading to no modification in the deep basin. Our results highlight the importance of including both icebergs and seasonal variations of subglacial discharge. ~~Davison et al. (2020) find that icebergs increase heat transfer towards the glacier due to iceberg-melt-driven estuarine circulation in Sermilik Fjord and for an idealized fjord without a sill. However, Davison et al. (2022) do observe~~

Davison et al. (2022) observe in a modelling study of an idealized steady-state fjord in summer conditions that when icebergs
470 extend below sill depth, the deep basin is cooled, as it does in our model. On the other hand, without a sill the deep basin is warmed, due to increased up-fjord heat transport (Davison et al., 2020, 2022). Our analysis explains the dynamics of GMW depression and IMAW formation at the high sill that leads to this ~~iceberg-duced-iceberg-induced~~ deep basin modification. Hager et al. (2022) also discover a similar mixing of GMW outflow down into the deep basin in a high-silled-high-silled glacial fjord in LeConte Bay in Alaska. Our results together with Davison et al. (2022) and Hager et al. (2022) show that high sills

475 in glacial fjords are significant not only as barriers preventing warm ambient water inflow ,but contribute to deep basin water modification.

Davison et al. (2022) study the impact of icebergs on plume melt in an idealized fjord setup but find only small changes in plume temperature and melt rate. They study a 500 m deep fjord with a single point-source plume, which has limited entrainment of ambient water into the plume and is thus relatively unaffected by iceberg-modification of the fjord water (Fig. 9 in (Davison et al., 2022)). The plume reaches the surface with all their different iceberg distributions, but since they do not discuss stratification or buoyancy, it is not clear if the neutral buoyancy depth changes with their different iceberg distributions. The fjord remains relatively warm at the surface in all of their scenarios (Figs. 4 and 6 in (Davison et al., 2022)), likely due to relatively lower average iceberg concentrations along the fjord (Fig. 2 in (Davison et al., 2022)). Comparison to our results further highlights the significance of iceberg concentration to changes both within the fjord and at the glacier terminus. However, they do also find that the cooling caused by icebergs reduces submarine melt both within and outside of the plume.




Figure 11. Illustration of iceberg modification of the plume and GMW outflow. Icebergs depress the neutral buoyancy depth of the plume and the subsequent GMW outflow. The GMW outflow gets further modified by icebergs, and mixes partially back into the fjord, when reaching a high sill.

5.2 Subglacial ~~dischare~~ discharge as a driver of seasonality

In our model, as in previous studies of high-silled fjords (Gladish et al., 2015; Carroll et al., 2017), the plume drives the circulation in the fjord .~~We find subglacial discharge to be the primary driver of seasonality of the circulation, as the high sill prevents the outflow of a weak early-season plume during peak summer. However, we find distinct circulation patterns for each season, due to interaction with the high sill: Subglacial discharge drives the early-season circulation in the fjord, with a significant amount of refluxing at the sill due to relatively deep GMW outflow, leading to basin water modification . Hager et al. (2022) discover a similar strong dynamics (Fig. 4b,f, S7). During peak summer, a strong GMW outflow is located~~

high enough to fully exit the fjord, and the compensating inflow over the sill reaches buoyancy at the upper part of the deep basin (Fig. 4c,g). In the late-season, GMW outflow is again refluxed at the sill (Fig. S7), but now the basin has freshened sufficiently that density-driven renewal of deep water starts to dominate the circulation, as the subglacial discharge wanes (Fig. 4d,h). This seasonality is very similar to the dynamics discovered by Hager et al. (2022) in LeConte Bay, where seasonal changes in subglacial discharge drive significantly different circulation patterns Alaska, suggesting that this is a typical seasonality pattern for high-silled glacial fjords. We find that seasonal surface warming in Disko Bay has little impact in the fjord when icebergs are included (Figs 3g,i and 7).

Subglacial discharge is also the driver of seasonality in of the freshwater flux from icebergs in our model, which we estimate to be $660\text{--}1050\text{ m}^3\text{ s}^{-1}$ in winter and $1270\text{--}1810\text{ m}^3\text{ s}^{-1}$ in peak summer (Fig. 6). This is in line with previous satellite-derived estimates of iceberg melt of $678\text{--}1346\text{ m}^3\text{ s}^{-1}$ and $1300\text{--}1700\text{ m}^3\text{ s}^{-1}$ for Sermeq Kujalleq in peak summer (Enderlin et al., 2016; Mankoff et al., 2020). Wintertime freshwater flux from iceberg melt is estimated to be several hundreds of $\text{m}^3\text{ s}^{-1}$ (Enderlin et al., 2016). Moon et al. (2018) estimate the peak iceberg freshwater flux for Sermilik-Sermilik Fjord to take place later in the season, September to November, since the volume of icebergs in the fjord accumulates throughout the calving season due to an increase in the ocean temperature. We do not consider variability of the iceberg distribution during the experiments; however, which could contribute to a larger late season freshwater flux from icebergs. However, our sensitivity experiments show that the increase in iceberg melt due to the plume is larger than the variability of the freshwater flux due to the iceberg distribution is $200\text{--}400\text{ m}^3\text{ s}^{-1}$, depending on the plume width, less than the increase due to subglacial discharge (Fig. 6).

Subglacial discharge drives the seasonality of the glacier-plume melt rate, both of the maximum melt rate and of the vertical reach of the plume (Fig. 5). ~~Observational studies of Sermeq Kujalleq indicate that there is a link between iceberg melange and seasonal changes in calving rate, as a reduction of calving during winter and the growth of rigid melange seem to correlate (Joughin et al., 2008; Cassotto et al., 2015; Joughin et al., 2020). Melt rate is~~ Meanwhile, melt rate is found to be connected to calving through either the calving-multiplier effect or a destabilizing influence as a moderator, but the exact processes are elusive and location dependent (O’Leary and Christoffersen, 2013; Rignot et al., 2015; Benn and Åström, 2018; Cowton et al., 2019; Cook et al., 2021; Slater et al., 2021). ~~Our results show that melt is localized at depth, particularly for the early season.~~ Melt localized at depth promotes undercutting, which is considered to be the determining factor for calving style (Benn and Åström, 2018; Slater et al., 2021). ~~Currently, bottom-out rotation is preferential for Sermeq Kujalleq (Amundson et al., 2010; Cassotto et al., 2019), suggesting that undercutting driven by melt is the main driver of calving.~~ Thus, we speculate that as subglacial discharge drives seasonality in melt rate, it will also drive seasonality of undercutting and thus calving of this that will be reflected in calving of the glacier. Our results are in line with previous studies, indicating that subglacial discharge has a strong control of the melt rate, both through discharge volume (Jenkins, 2011; Truffer and Motyka, 2016), but also through the subglacial drainage system properties that ultimately control the plume height (Khazendar et al., 2019; Straneo et al., 2020) the combination of subglacial discharge volume and drainage width — through the subglacial drainage system properties — control plume velocity and the reach of the plume (Jenkins, 2011; Cowton et al., 2015).

Observational studies of Sermeq Kujalleq indicate that there is a link between iceberg ~~mélange~~ and seasonal changes in calving rate, as a reduction of calving during winter and the growth of rigid ~~mélange~~ seem to correlate (Joughin et al., 2008; Cassotto et al., 2020).

The connection between iceberg ~~mélange~~ and the suppression of calving is sometimes attributed to mechanical support by a rigid ~~mélange~~ that prevents calving (Joughin et al., 2008; Burton et al., 2018; Joughin et al., 2020). However, mechanical support disappears once the rigidity is lost, as the non-rigid ~~mélange~~ is a granular material, able to compact and relax under calving events (Peters et al., 2015). The presence of reliably rigid ~~mélange~~ in Ilulissat Icefjord varies inter-annually, and it has been present only sporadically in the previous decade (Cassotto et al., 2015; Joughin et al., 2020), and thus does not give a comprehensive explanation to the link between ~~mélange~~ and calving. ~~Since our results show that icebergs limit melt to the deep part of the glacier front, we hypothesize that icebergs moderate calving by strengthening undercutting. Furthermore, our~~ Our results indicate that the plume ~~transports heat up to the~~ ~~mélange~~ and significantly increases currents within the iceberg ~~mélange~~ (Fig. S3), ~~likely inhibiting rigid mélange formation which likely inhibits rigid mélange formation during high subglacial discharge.~~ Thus, we see the seasonal growth of rigid ~~mélange~~ as a consequence of ~~decreased melting and calving due to a weakening plume.~~ Mernild et al. (2015) ~~find autumn cooling in the~~ ~~weakening plume, with concomitant weakening of the circulation and the supply of heat to the~~ ~~mélange enables formation of rigid mélange.~~ Our study covers only one year with constant iceberg conditions. However, inter-annually, the heat transported by the plume into the ~~mélange~~ will depend on the deep basin ~~of Ilulissat Icefjord, which could be a consequence of increased entrainment and iceberg modification as the plume weakens, similarly as for the weak spring plume in our model~~ water temperature, as discussed in (Joughin et al., 2020), but also on changes in the subglacial discharge volume and the outlet properties (Jenkins, 2011; Slater et al., 2017a). On the other hand, changes in the iceberg coverage will promote variability in the formation of the rigid ~~mélange~~, as the ability of the ~~mélange~~ to remove heat depends on the iceberg coverage.

Our results indicate that icebergs impact the glacier front irrespective of ~~mélange~~ rigidity. Icebergs suppress glacier melt at the top 100–300 m of the glacier front (Fig. 4g). ~~In this case cooling due to increased entrainment and modification would further reduce the melt rate of the glacier, amplifying the winter suppression of melt.~~

5) by depressing the buoyancy of the plume. This suppression strengthens the uneven vertical distribution of glacier melt and thus promotes undercutting. We hypothesize that icebergs moderate calving by increasing undercutting throughout the runoff season. Iceberg control over the neutral buoyancy depth of the plume provides a mechanism for the iceberg ~~mélange~~ to influence melt, even during peak summer discharge when the ~~mélange~~ is not rigid. ~~This suppression is not sensitive to maximum iceberg keel depth but is sensitive to the density of icebergs at the depth of neutral buoyancy (Fig. 10), indicating that iceberg concentration within the intermediate layer is a controlling factor in the suppression of melt of the glacier.~~ In the absence of a ~~physical comprehensive~~ understanding of calving ~~and the role of undercutting~~ it is difficult to make quantitative estimates of how a certain deepening of the neutral buoyancy depth of the plume would impact calving, and we see this as a key point of further study. Further study is also needed on iceberg-plume-GMW dynamics, and on how ~~this~~ ~~these~~ dynamics will evolve in a warming climate. Much will depend on how the subglacial drainage system will react to longer and more intense melt seasons. Increased calving has the potential to strengthen the iceberg-induced modification in the Ilulissat Icefjord: As Sermeq

Kujalleq is predicted to retreat into even deeper geometry (Bondzio et al., 2017). Therefore (Bondzio et al., 2018), maximum iceberg depth, overall discharge volume and thus iceberg modification in the deep basin can be expected to increase.

5.3 Model uncertainties

565 One of the key features of the IceBerg package is that the icebergs do not move or change volume when melting, and we choose not to edit the distribution manually during experiments. While the iceberg cover of the Ilulissat Icefjord is a relatively constant feature, due to the large calving flux of the glacier, observations indicate some annual variability (Cassotto et al., 2015; Enderlin et al., 2016; Joughin et al., 2020). We expect the range of iceberg distributions applied in the sensitivity experiments to span the natural variability, and also along-fjord variations of the iceberg distribution in the Ilulissat Icefjord (Sec. 4.4). We
570 do not expect temporary reductions in the iceberg cover to impact the hydrography of the fjord to a large degree. Observations of summer surface conditions in Ilulissat Icefjord in 2000–2011 show consistently an iceberg-dominated surface at melting point (Cassotto et al., 2015), which makes us assume that any deviations are short in duration.

Our modelled iceberg distribution produces slightly cooler and fresher deep basin than what is observed (Fig. 3f–mg, 8), which seems to be a combination of slightly different properties of the deepest water in the basin, and an overestimated
575 contribution of subglacial discharge within the deep basin (Fig. 7). In the model, GMW in the basin is largely a product of early season mixing-refluxing at the sill, which in reality could be reduced due to shallow or sparse iceberg distribution at the sill early in the season. The good correspondence between the observations and sensitivity experiment *IB200P* (Figs. 3m–i and 9b) indicates that icebergs could have been shallower in 2014 than modelled. (Enderlin et al., 2016) shows that in June 2014 icebergs in Ilulissat Icefjord were indeed relatively shallow, further supporting this interpretation. ~~We~~ For simplicity, we keep
580 the properties of Disko Bay water at sill depth constant~~for simplicity, but variability~~. However, the variability in the water properties at the sill depth could cause dense water inflows ~~that contribute to~~, influencing the deep basin properties. Also, ~~we assume that each phase of the seasonal evolution of the subglacial discharge is equal — 30 days of duration — which could potentially overestimate the early season contribution. The seasonality of the duration of the early season can vary, and a rapid early season would reduce early-season refluxing at the sill. The choice of model parameters can influence the deep basin~~
585 modification, as horizontal diffusivity and the restoration timescale have a slight impact to the deep basin properties (Fig. S4). However, this does not impact the overall results or processes discussed in this study. A thorough observational study focusing on the early-season processes in Ilulissat Icefjord is needed to constrain these uncertainties.

Our model setup is designed for the runoff season, ignoring potentially important wintertime processes, such as large tidal events, sea ice formation and decrease in iceberg draft. Thus, if extended over the following winter, the model does not
590 fully recover from the summer conditions (Fig. S2). However, properties within the iceberg extent recover rapidly (Fig. S3), and dense water renewal in the deep basin continues over winter (Fig. S2c). We keep the iceberg distribution constant also during winter, although in reality iceberg drafts would significantly decrease (Enderlin et al., 2016), reducing modification of the inflowing water during winter and enabling faster recovery of the deep basin properties. Furthermore, it is possible that weak subglacial discharge is ~~likely to vary greatly both intra and interannually (Cook et al., 2021)~~, also present during winter,

595 contributing to the deep basin circulation (Cook et al., 2021). This would mean an underestimated contribution of frontal melt from the glacier during winter.

The IceBerg package implements freshening due to iceberg melt by applying a negative salinity anomaly. In reality we would expect the icebergs to be a source of freshwater input to the fjord (Mernild et al., 2015; Enderlin et al., 2016; Moon et al., 2018). The negative salinity approach forces the modelled inflow and outflow of the iceberg-melt driven circulation to
600 be equal in volume, leading instead of a net outflow as expected. Including icebergs leads to a 50% increase in the volume flux over the sill ~~-, when including icebergs due to the added freshwater export~~ (Fig. 4d,h). ~~This creates a rapid inflow at the base of the icebergs, increasing drift-induced melt. More observations are needed to analyse if the negative salinity approach overestimates the inflow over the sill, which we would expect to be smaller than outflow if icebergs are considered as a source of freshwater.~~ It is uncertain, however, if this increase is realistic for the inflow. We do not include tides or rotational effects,
605 since their ~~impacts~~ impacts are small in our high-silled setup (Carroll et al., 2017). However, both of these effects should be considered in a bathymetrically accurate study of Ilulissat Icefjord, since they could potentially impact the mixing at the sill: rotation could redistribute the GMW outflow laterally, while tides would increase mixing at the sill (Carroll et al., 2017), potentially increasing iceberg melt at the sill.

Icebergs are represented by cell-averaged temperature and salinity forcings in the IceBerg package (Davison et al., 2020) ~~-, and~~
610 ~~thus we do not consider icebergs as~~ and are obstacles for flow only in a grid-cell-average sense. However, Hughes (2022) show that individual icebergs alter the circulation by creating a complicated flow network, which slows down the near-surface current(?). This mechanical effect of the ~~icebergs~~ iceberg network would also cause depression of the GMW outflow, complementing our modelled stratification-driven depression. Furthermore, we do not consider variations to the melt rate parameterizations due to different flow regimes, as described in (Fitzmaurice et al., 2018), which could be a further development point for
615 the IceBerg package. For simplicity, we choose to use standard values of the turbulent transfer coefficients for both heat and salt transfer both for the glacier and for the icebergs (Table. S1). Jackson et al. (2020) suggest new, higher values for the turbulence parameters, and ~~(Davison et al., 2020) tests~~ Davison et al. (2020) test the possibility of varying these parameters for ~~iceberg~~ iceberg melt. Given that their study shows that increased values of turbulent transfer enhance the impacts of icebergs, we consider the standard values to be a conservative estimate. The ~~turbulence parameters at~~ contribution of direct melt from the glacier
620 front ~~have negligible impact to the results as frontal melt is such a small contributor in the fjord is negligible in our simulations, representing only 2% of the total peak freshwater flux~~ (Fig. 6). This is likely a lower-end estimate, since melt outside of the plume is likely underestimated with the standard turbulent transfer parameters (Sutherland et al., 2019; Jackson et al., 2020). This will have most impact during winter when the other freshwater sources are at a minimum.

6 Conclusions

625 We find that ~~icebergs are~~ iceberg melt is the key factor modifying the water mass properties of the Ilulissat Icefjord, and that omitting icebergs can lead to misleading interpretations of both water mass properties and the ~~glacier's response to external forcing.~~ Icebergs impact dynamics of the ice-ocean interface. Iceberg melt impacts Ilulissat Icefjord in three main ways: First,

icebergs cool and freshen the water column ~~within~~over their vertical extent; Second, ~~icebergs limit the vertical reach of the plume, by deepening iceberg melt causes~~ the neutral buoyancy depth ~~and the outflow route of the plume and the export~~
630 of glacially modified ~~water~~waters to be deeper; Third, icebergs modify the deep basin, below their vertical extent, ~~due to both increased mixing of glacially modified outflow into the fjord at the high sill, and iceberg modification of~~by driving
mixing of the glacially modified waters with the deep basin waters and by modifying the incoming ambient ~~water from Disko Bay. Entrainment is particularly important early in the season, when subglacial discharge is weak. The seasonal~~waters. We
635 describe the evolution of the fjord properties and circulation over a runoff season and find the subglacial discharge flux ~~is the primary control of the seasonality of the plume and frontal melt rate, which further supports the idea of the plume as a driver of seasonality in undercutting and thus calving. During peak summer discharge, icebergs depress the plume and thus limit glacier~~
to be the driver of seasonality of both glacier and iceberg melt. Changes in the water column properties due to iceberg submarine melt decrease the total melt of the glacier front and depress the subglacial discharge plume. We postulate that submarine iceberg melt increases undercutting by limiting plume melt to the deep part of the glacier front and, through changes
640 in undercutting, iceberg melt also moderates glacier calving in Ilulissat Icefjord. We postulate that the impact of ~~icebergs on the neutral buoyancy depth of the plume is a key mechanism moderating glacier calving in Ilulissat Icefjord, and submarine iceberg melt on the frontal melt of the glacier~~ provides a comprehensive link between iceberg ~~mélange~~mélange and glacier calving throughout the runoff season.

Code availability. MITgcm is freely available at <https://github.com/MITgcm/MITgcm/archive/master.zip> and <https://doi.org/10.5281/zenodo.4968496> (Marshall et al., 1997). The IcePlume module is available from Tom Cowton on request. The IceBerg module is available at <https://doi.org/10.5281/zenodo.3979647> (Davison et al., 2020).

Data availability. The model domain, boundary conditions and output are available from the corresponding author on request.

Author contributions. KK and KHN conceived the study. KK designed and conducted the simulations and analysis and wrote the original manuscript. FS and KHN supported the interpretation of the model results and contributed to the preparation of the manuscript.

650 *Competing interests.* KHN is a member of the editorial board of The Cryosphere. The authors declare that no other conflicts of interest are present.

Acknowledgements. KK and KHN were supported by the European Research Council under the European Community's Seventh Framework Program (FP7/2007-2013) / ERC grant agreement 610055 as part of the ice2ice project. KHN is also supported by the ClimateNarratives NRC

grant (324520). FS is supported by the NSF and the Heising Simons Foundation. The simulations were performed on resources provided by
655 Sigma2 — the National Infrastructure for High Performance Computing and Data Storage in Norway. Data from the Greenland Ecosystem
Monitoring Programme were provided by the Greenland Institute of Natural Resources, Nuuk, Greenland in collaboration with Department
of Bioscience, Aarhus University, Denmark.

References

- Åkesson, H., Nisancioglu, K. H., and Nick, F. M.: Impact of Fjord Geometry on Grounding Line Stability, *Frontiers in Earth Science*, 6, 71, <https://doi.org/10.3389/feart.2018.00071>, 2018.
- Amundson, J. M., Fahnestock, M., Truffer, M., Brown, J., Lüthi, M. P., and Motyka, R. J.: Ice mélange dynamics and implications for terminus stability, Jakobshavn Isbræ, Greenland, *Journal of Geophysical Research: Earth Surface*, 115, <https://doi.org/10.1029/2009JF001405>, 2010.
- Beaird, N., Straneo, F., and Jenkins, W.: Characteristics of meltwater export from Jakobshavn Isbræ and Ilulissat Icefjord, *Annals of Glaciology*, 58, 107–117, <https://doi.org/10.1017/aog.2017.19>, 2017.
- Benn, D. I. and Åström, J. A.: Calving glaciers and ice shelves, *Advances in Physics: X*, 3, 1048–1076, <https://doi.org/10.1080/23746149.2018.1513819>, 2018.
- Bondzio, J. H., Morlighem, M., Seroussi, H., Kleiner, T., Rückamp, M., Mouginot, J., Moon, T., Larour, E. Y., and Humbert, A.: The mechanisms behind Jakobshavn Isbræ's acceleration and mass loss: A 3-D thermomechanical model study, *Geophysical Research Letters*, 44, 6252–6260, <https://doi.org/10.1002/2017GL073309>, 2017.
- Bondzio, J. H., Morlighem, M., Seroussi, H., Wood, M. H., and Mouginot, J.: Control of Ocean Temperature on Jakobshavn Isbræ's Present and Future Mass Loss, *Geophysical Research Letters*, 45, 12,912–12,921, <https://doi.org/10.1029/2018GL079827>, 2018.
- Böning, C. W., Behrens, E., Biastoch, A., Getzlaff, K., and Bamber, J. L.: Emerging impact of Greenland meltwater on deepwater formation in the North Atlantic Ocean, *Nature Geoscience* 2016 9:7, 9, 523–527, <https://doi.org/10.1038/ngeo2740>, 2016.
- Burton, J. C., Amundson, J. M., Cassotto, R., Kuo, C. C., and Dennin, M.: Quantifying flow and stress in ice mélange, the world's largest granular material, *Proceedings of the National Academy of Sciences of the United States of America*, 115, 5105–5110, <https://doi.org/10.1073/PNAS.1715136115/VIDEO-1>, 2018.
- Carroll, D., Sutherland, D. A., Hudson, B., Moon, T., Catania, G. A., Shroyer, E. L., Nash, J. D., Bartholomäus, T. C., Felikson, D., Stearns, L. A., Noël, B. P. Y., and van den Broeke, M. R.: The impact of glacier geometry on meltwater plume structure and submarine melt in Greenland fjords, *Geophysical Research Letters*, 43, 9739–9748, <https://doi.org/10.1002/2016GL070170>, 2016.
- Carroll, D., Sutherland, D. A., Shroyer, E. L., Nash, J. D., Catania, G. A., and Stearns, L. A.: Subglacial discharge-driven renewal of tidewater glacier fjords, *Journal of Geophysical Research: Oceans*, 122, 6611–6629, <https://doi.org/10.1002/2017JC012962>, 2017.
- Cassotto, R., Fahnestock, M., Amundson, J. M., Truffer, M., and Joughin, I.: Seasonal and interannual variations in ice mélange and its impact on terminus stability, Jakobshavn Isbræ, Greenland, *Journal of Glaciology*, 61, 76–88, <https://doi.org/10.3189/2015JoG13J235>, 2015.
- Cassotto, R., Fahnestock, M., Amundson, J. M., Truffer, M., Boettcher, M. S., De La Peña, S., and Howat, I.: Non-linear glacier response to calving events, Jakobshavn Isbræ, Greenland, *Journal of Glaciology*, 65, 39–54, <https://doi.org/10.1017/JOG.2018.90>, 2019.
- Catania, G. A., Stearns, L. A., Moon, T. A., Enderlin, E. M., and Jackson, R. H.: Future Evolution of Greenland's Marine-Terminating Outlet Glaciers, *Journal of Geophysical Research: Earth Surface*, 125, <https://doi.org/10.1029/2018JF004873>, 2020.
- Cavanagh, J. P., Lampkin, D. J., and Moon, T.: Seasonal Variability in Regional Ice Flow Due to Meltwater Injection Into the Shear Margins of Jakobshavn Isbræ, *Journal of Geophysical Research: Earth Surface*, 122, 2488–2505, <https://doi.org/10.1002/2016JF004187>, 2017.
- Cook, S. J., Christoffersen, P., and Todd, J.: A fully-coupled 3D model of a large Greenlandic outlet glacier with evolving subglacial hydrology, frontal plume melting and calving, *Journal of Glaciology*, pp. 1–17, <https://doi.org/10.1017/jog.2021.109>, 2021.

- Cowton, T., Slater, D., Sole, A., Goldberg, D., and Nienow, P.: Modeling the impact of glacial runoff on fjord circulation and submarine melt rate using a new subgrid-scale parameterization for glacial plumes, *Journal of Geophysical Research: Oceans*, 120, 796–812, <https://doi.org/10.1002/2014JC010324>, 2015.
- Cowton, T. R., Todd, J. A., and Benn, D. I.: Sensitivity of Tidewater Glaciers to Submarine Melting Governed by Plume Locations, *Geophysical Research Letters*, 46, 11 219–11 227, <https://doi.org/10.1029/2019GL084215>, 2019.
- Cuffey, K. M. and Paterson, W. S. B.: *The Physics of Glaciers*, Elsevier Science & Technology Books, 4th editio edn., 2010.
- Davison, B. J., Cowton, T. R., Cottier, F. R., and Sole, A. J.: Iceberg melting substantially modifies oceanic heat flux towards a major Greenlandic tidewater glacier, *Nature Communications*, 11, 1–13, <https://doi.org/10.1038/s41467-020-19805-7>, 2020.
- Davison, B. J., Cowton, T., Sole, A., Cottier, F., and Nienow, P.: Modelling the effect of submarine iceberg melting on glacier-adjacent water properties, *The Cryosphere*, 16, 1181–1196, <https://doi.org/10.5194/TC-16-1181-2022>, 2022.
- De Andrés, E., Slater, D. A., Straneo, F., Otero, J., Das, S., and Navarro, F.: Surface emergence of glacial plumes determined by fjord stratification, *Cryosphere*, 14, 1951–1969, <https://doi.org/10.5194/TC-14-1951-2020>, 2020.
- Enderlin, E. M., Hamilton, G. S., Straneo, F., and Sutherland, D. A.: Iceberg meltwater fluxes dominate the freshwater budget in Greenland’s iceberg-congested glacial fjords, *Geophysical Research Letters*, 43, 11,287–11,294, <https://doi.org/10.1002/2016GL070718>, 2016.
- Fenty, I., Willis, J., Khazendar, A., Dinardo, S., Forsberg, R., Fukumori, I., Holland, D., Jakobsson, M., Moller, D., Morison, J., Meunchow, A., Rignot, E., Schodlock, M., Thompson, A., Tino, K., Rutherford, M., and Trenholm, N.: Oceans Melting Greenland: Early results from NASA’s ocean-ice mission in Greenland, *Oceanography*, 29, 72–83, <https://doi.org/10.5670/ocean.2016.100>, 2016.
- Fitzmaurice, A., Cenedese, C., and Straneo, F.: A Laboratory Study of Iceberg Side Melting in Vertically Sheared Flows, *Journal of Physical Oceanography*, 48, 1367–1373, <https://doi.org/10.1175/JPO-D-17-0185.1>, 2018.
- Frank, T., Åkesson, H., de Fleurian, B., Morlighem, M., and Nisancioglu, K.: Geometric Controls of Tidewater Glacier Dynamics, *The Cryosphere Discussions*, pp. 1–32, <https://doi.org/10.5194/TC-2021-81>, 2021.
- Gladish, C. V., Holland, D. M., Rosing-Asvid, A., Behrens, J. W., and Boje, J.: Oceanic boundary conditions for Jakobshavn Glacier. Part I: Variability and renewal of Ilulissat Icefjord waters, 2001–14, *Journal of Physical Oceanography*, 45, 3–32, <https://doi.org/10.1175/JPO-D-14-0044.1>, 2015.
- Goelzer, H., Nowicki, S., Payne, A., Larour, E., Seroussi, H., Lipscomb, W. H., Gregory, J., Abe-Ouchi, A., Shepherd, A., Simon, E., Agosta, C., Alexander, P., Aschwanden, A., Barthel, A., Calov, R., Chambers, C., Choi, Y., Cuzzone, J., Dumas, C., Edwards, T., Felikson, D., Fettweis, X., Golledge, N. R., Greve, R., Humbert, A., Huybrechts, P., Le Clec’h, S., Lee, V., Leguy, G., Little, C., Lowry, D., Morlighem, M., Nias, I., Quiquet, A., Rückamp, M., Schlegel, N. J., Slater, D. A., Smith, R., Straneo, F., Tarasov, L., Van De Wal, R., and Van Den Broeke, M.: The future sea-level contribution of the Greenland ice sheet: A multi-model ensemble study of ISMIP6, *Cryosphere*, 14, 3071–3096, <https://doi.org/10.5194/TC-14-3071-2020>, 2020.
- Greenland Ecosystem Monitoring: MarineBasis Disko - Water column - CTD measurements (Version 1.0) [Data set], Tech. rep., Greenland Ecosystem Monitoring, <https://doi.org/https://doi.org/10.17897/WH30-HT61>, 2020.
- Hager, A. O., Sutherland, D. A., Amundson, J. M., Jackson, R. H., Kienholz, C., Motyka, R. J., and Nash, J. D.: Subglacial Discharge Reflux and Buoyancy Forcing Drive Seasonality in a Silled Glacial Fjord, *Journal of Geophysical Research: Oceans*, 127, e2021JC018355, <https://doi.org/10.1029/2021JC018355>, 2022.
- Hughes, K. G.: Pathways , form drag , and turbulence in simulations of an ocean flowing through an ice m ´ elange, 2022.

- 730 Jackson, R. H., Shroyer, E. L., Nash, J. D., Sutherland, D. A., Carroll, D., Fried, M. J., Catania, G. A., Bartholomaeus, T. C., and Stearns, L. A.: Near-glacier surveying of a subglacial discharge plume: Implications for plume parameterizations, *Geophysical Research Letters*, 44, 6886–6894, <https://doi.org/10.1002/2017GL073602>, 2017.
- Jackson, R. H., Nash, J. D., Kienholz, C., Sutherland, D. A., Amundson, J. M., Motyka, R. J., Winters, D., Skillingstad, E., and Pettit, E. C.: Meltwater Intrusions Reveal Mechanisms for Rapid Submarine Melt at a Tidewater Glacier, *Geophysical Research Letters*, 47, e2019GL085335, <https://doi.org/10.1029/2019GL085335>, 2020.
- 735 Jenkins, A.: Convection-driven melting near the grounding lines of ice shelves and tidewater glaciers, *Journal of Physical Oceanography*, 41, 2279–2294, <https://doi.org/10.1175/JPO-D-11-03.1>, 2011.
- Joughin, I., Howat, I. M., Fahnestock, M., Smith, B., Krabill, W., Alley, R. B., Stern, H., and Truffer, M.: Continued evolution of Jakobshavn Isbrae following its rapid speedup, *Journal of Geophysical Research*, 113, F04006, <https://doi.org/10.1029/2008JF001023>, 2008.
- 740 Joughin, I., E. Shean, D., E. Smith, B., and Floricioiu, D.: A decade of variability on Jakobshavn Isbræ: Ocean temperatures pace speed through influence on mélange rigidity, *Cryosphere*, 14, 211–227, <https://doi.org/10.5194/TC-14-211-2020>, 2020.
- Khazendar, A., Fenty, I. G., Carroll, D., Gardner, A., Lee, C. M., Fukumori, I., Wang, O., Zhang, H., Seroussi, H., Moller, D., Noël, B. P. Y., van den Broeke, M. R., Dinardo, S., and Willis, J.: Interruption of two decades of Jakobshavn Isbrae acceleration and thinning as regional ocean cools, *Nature Geoscience*, 12, 277–283, <https://doi.org/10.1038/s41561-019-0329-3>, 2019.
- 745 Mankoff, K. D., Solgaard, A., Colgan, W., Ahlstrøm, A. P., Abbas Khan, S., and Fausto, R. S.: Greenland Ice Sheet solid ice discharge from 1986 through March 2020, *Earth System Science Data*, 12, 1367–1383, <https://doi.org/10.5194/ESSD-12-1367-2020>, 2020.
- Marshall, J., Adcroft, A., Hill, C., Perelman, L., and Heisey, C.: A finite-volume, incompressible Navier Stokes model for studies of the ocean on parallel computers, *Journal of Geophysical Research: Oceans*, 102, 5753–5766, <https://doi.org/10.1029/96JC02775>, 1997.
- Meire, L., Mortensen, J., Meire, P., Juul-Pedersen, T., Sej, M. K., Rysgaard, S., Nygaard, R., Huybrechts, P., and Meysman, F. J.: Marine-terminating glaciers sustain high productivity in Greenland fjords, *Global Change Biology*, 23, 5344–5357, <https://doi.org/10.1111/GCB.13801>, 2017.
- 750 Mernild, S. H., Holland, D. M., Holland, D., Rosing-Asvid, A., Yde, J. C., Liston, G. E., and Steffen, K.: Freshwater flux and spatiotemporal simulated runoff variability into Ilulissat Icefjord, West Greenland, linked to salinity and temperature observations near tidewater glacier margins obtained using instrumented ringed seals, *Journal of Physical Oceanography*, 45, 1426–1445, <https://doi.org/10.1175/JPO-D-14-0217.1>, 2015.
- 755 Mojica, J. F., Djoumna, G., Holland, D. M., and Holland, D.: Interannual summer mixing processes in the Ilulissat Icefjord, Greenland, *Journal of Marine Systems*, 214, 103476, <https://doi.org/10.1016/j.jmarsys.2020.103476>, 2021.
- Moon, T., Sutherland, D. A., Carroll, D., Felikson, D., Kehrl, L., and Straneo, F.: Subsurface iceberg melt key to Greenland fjord freshwater budget, *Nature Geoscience*, 11, 49–54, <https://doi.org/10.1038/s41561-017-0018-z>, 2018.
- 760 Morlighem, M., Williams, C. N., Rignot, E., An, L., Arndt, J. E., Bamber, J. L., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L., Mouginot, J., Noël, B. P. Y., O’Cofaigh, C., Palmer, S., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., and Zinglensen, K. B.: BedMachine v3: Complete Bed Topography and Ocean Bathymetry Mapping of Greenland From Multibeam Echo Sounding Combined With Mass Conservation, *Geophysical Research Letters*, 44, 11,051–11,061, <https://doi.org/10.1002/2017GL074954>, 2017.
- 765

- Mortensen, J., Rysgaard, S., Bendtsen, J., Lennert, K., Kanzow, T., Lund, H., and Meire, L.: Subglacial Discharge and its Down-Fjord Transformation in West Greenland Fjords With an Ice Melange, *Journal of Geophysical Research: Oceans*, 125, e2020JC016301, <https://doi.org/10.1029/2020JC016301>, 2020.
- 770 Motyka, R. J., Truffer, M., Fahnestock, M., Mortensen, J., Rysgaard, S., and Howat, I.: Submarine melting of the 1985 Jakobshavn Isbræ floating tongue and the triggering of the current retreat, *Journal of Geophysical Research: Earth Surface*, 116, <https://doi.org/10.1029/2009JF001632>, 2011.
- Mouginot, J., Rignot, E., Björk, A. A., van den Broeke, M., Millan, R., Morlighem, M., Noël, B., Scheuchl, B., and Wood, M.: Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018, *Proceedings of the National Academy of Sciences of the United States of America*, 116, 9239–9244, https://doi.org/10.1073/PNAS.1904242116/SUPPL_FILE/PNAS.1904242116.SD02.XLSX, 2019.
- 775 Muilwijk, M., Straneo, F., Slater, D. A., Smedsrud, L. H., Holte, J., Wood, M., Andresen, C. S., and Harden, B.: Export of ice sheet meltwater from Upernavik Fjord, West Greenland, *Journal of Physical Oceanography*, -1, <https://doi.org/10.1175/JPO-D-21-0084.1>, 2021.
- O’Leary, M. and Christoffersen, P.: Calving on tidewater glaciers amplified by submarine frontal melting, *Cryosphere*, 7, 119–128, <https://doi.org/10.5194/TC-7-119-2013>, 2013.
- Peters, I. R., Amundson, J. M., Cassotto, R., Fahnestock, M., Darnell, K. N., Truffer, M., and Zhang, W. W.: Dynamic jamming of iceberg-choked fjords, *Geophysical Research Letters*, 42, 1122–1129, <https://doi.org/10.1002/2014GL062715>, 2015.
- 780 Rezvanbehbahani, S., Stearns, L. A., Keramati, R., Shankar, S., and van der Veen, C. J.: Significant contribution of small icebergs to the freshwater budget in Greenland fjords, *Communications Earth & Environment*, 1, 1–7, <https://doi.org/10.1038/s43247-020-00032-3>, 2020.
- Rignot, E., Fenty, I., Xu, Y., Cai, C., and Kemp, C.: Undercutting of marine-terminating glaciers in West Greenland, *Geophysical Research Letters*, 42, 5909–5917, <https://doi.org/10.1002/2015GL064236>, 2015.
- 785 Sciascia, R., Straneo, F., Cenedese, C., and Heimbach, P.: Seasonal variability of submarine melt rate and circulation in an East Greenland fjord, *Journal of Geophysical Research: Oceans*, 118, 2492–2506, <https://doi.org/10.1002/jgrc.20142>, 2013.
- Slater, D., Nienow, P., Sole, A., Cowton, T., Mottram, R., Langen, P., and Mair, D.: Spatially distributed runoff at the grounding line of a large Greenlandic tidewater glacier inferred from plume modelling, *Journal of Glaciology*, 63, 309–323, <https://doi.org/10.1017/jog.2016.139>, 2017a.
- 790 Slater, D. A., Goldberg, D. N., Nienow, P. W., and Cowton, T. R.: Scalings for Submarine Melting at Tidewater Glaciers from Buoyant Plume Theory, *Journal of Physical Oceanography*, 46, 1839–1855, <https://doi.org/10.1175/JPO-D-15-0132.1>, 2016.
- Slater, D. A., Nienow, P. W., Goldberg, D. N., Cowton, T. R., and Sole, A. J.: A model for tidewater glacier undercutting by submarine melting, *Geophysical Research Letters*, 44, 2360–2368, <https://doi.org/10.1002/2016GL072374>, 2017b.
- Slater, D. A., Nienow, P. W., Goldberg, D. N., Cowton, T. R., and Sole, A. J.: A model for tidewater glacier undercutting by submarine melting, *Geophysical Research Letters*, 44, 2360–2368, <https://doi.org/10.1002/2016GL072374>, 2017c.
- 795 Slater, D. A., Straneo, F., Das, S. B., Richards, C. G., Wagner, T. J. W., and Nienow, P. W.: Localized Plumes Drive Front-Wide Ocean Melting of A Greenlandic Tidewater Glacier, *Geophysical Research Letters*, 45, 12,350–12,358, <https://doi.org/10.1029/2018GL080763>, 2018.
- Slater, D. A., Straneo, F., Felikson, D., Little, C. M., Goelzer, H., Fettweis, X., and Holte, J.: Estimating Greenland tidewater glacier retreat driven by submarine melting, *Cryosphere*, 13, 2489–2509, <https://doi.org/10.5194/TC-13-2489-2019>, 2019.
- 800 Slater, D. A., Benn, D. I., Cowton, T. R., Bassis, J. N., and Todd, J. A.: Calving Multiplier Effect Controlled by Melt Undercut Geometry, *Journal of Geophysical Research: Earth Surface*, 126, e2021JF006191, <https://doi.org/10.1029/2021JF006191>, 2021.

- Straneo, F. and Cenedese, C.: The Dynamics of Greenland's Glacial Fjords and Their Role in Climate, <http://dx.doi.org/10.1146/annurev-marine-010213-135133>, 7, 89–112, <https://doi.org/10.1146/ANNUREV-MARINE-010213-135133>, 2015.
- 805 Straneo, F., Heimbach, P., Sergienko, O., Hamilton, G., Catania, G., Griffies, S., Hallberg, R., Jenkins, A., Joughin, I., Motyka, R., Pfeffer, W. T., Price, S. F., Rignot, E., Scambos, T., Truffer, M., and Vieli, A.: Challenges to understanding the dynamic response of Greenland's marine terminating glaciers to oceanic and atmospheric forcing, *Bulletin of the American Meteorological Society*, 94, 1131–1144, <https://doi.org/10.1175/BAMS-D-12-00100.1>, 2013.
- Straneo, F., Sutherland, D. A., Stearns, L., Catania, G., Heimbach, P., Moon, T., Cape, M. R., Laidre, K. L., Barber, D., Rysgaard, S.,
 810 Mottram, R., Olsen, S., Hopwood, M. J., and Meire, L.: The case for a sustained Greenland Ice sheet-Ocean Observing System (GrIOOS), *Frontiers in Marine Science*, 6, 138, <https://doi.org/10.3389/FMARS.2019.00138/BIBTEX>, 2019.
- Sutherland, D. A., Jackson, R. H., Kienholz, C., Amundson, J. M., Dryer, W. P., Duncan, D., Eidam, E. F., Motyka, R. J., and Nash, J. D.: Direct observations of submarine melt and subsurface geometry at a tidewater glacier, *Science*, 365, 369–374, https://doi.org/10.1126/SCIENCE.AAX3528/SUPPL_FILE/AAX3528-SUTHERLAND-SM.PDF, 2019.
- 815 Truffer, M. and Motyka, R. J.: Where glaciers meet water: Subaqueous melt and its relevance to glaciers in various settings, *Reviews of Geophysics*, 54, 220–239, <https://doi.org/10.1002/2015RG000494>, 2016.
- Wood, M., Rignot, E., Fenty, I., An, L., Bjørk, A., van den Broeke, M., Cai, C., Kane, E., Menemenlis, D., Millan, R., Morlighem, M., Mougnot, J., Noël, B., Scheuchl, B., Velicogna, I., Willis, J. K., and Zhang, H.: Ocean forcing drives glacier retreat in Greenland, *Science Advances*, 7, https://doi.org/10.1126/SCIADV.ABA7282/SUPPL_FILE/ABA7282_TABLE_S1.XLSX, 2021.