

Allometric scaling of retrogressive thaw slumps

Jurjen van der Sluijs¹, Steven V. Kokelj², Jon F. Tunncliffe³

¹Northwest Territories Centre for Geomatics, Yellowknife, NT, X1A 2L9, Canada

²Northwest Territories Geological Survey, Yellowknife, NT, X1A 2L9, Canada

5 ³School of Environment, University of Auckland, Auckland, NZ

Correspondence to: Jurjen van der Sluijs (Jurjen_vandersluijs@gov.nt.ca)

Abstract. In the warming Arctic, retrogressive thaw slumping (RTS) has emerged as the primary thermokarst modifier of ice-rich permafrost slopes, raising urgency to investigate the distribution and intensification of disturbances and ~~to determine trajectories of landscape evolution and~~ the cascade of effects. Tracking RTS is challenging due to the constraints of remote sensing products and a narrow understanding of complex, thaw-driven landforms, however, high-resolution elevation models provide new insights into geomorphic change. Structural traits, such as RTS depth-of-thaw or volume, can be obtained through allometric scaling. To address fundamental knowledge gaps related to area-volume scaling of RTS, a suitable surface interpolation technique was first needed to model pre-disturbance topography upon which volume estimates could be based. Among 8 methods with 32 parameterizations, Natural Neighbour surface interpolation achieved the best precision in reconstructing pre-disturbed slope topography (90th percentile Root Mean Square Difference \pm 1.0 m). An inverse association between RTS volume and relative volumetric error was observed, with uncertainties <10% for large slumps and <20% for small-to-medium slumps. Second, a Multisource Slump Inventory (MSI) for two study areas in the Beaufort Delta (Canada) was required to characterize the diverse range of disturbance morphologies and activity levels, which provided temporally-consistent ~~information on characterization of~~ thaw slump-~~affected slopes and attributes~~. The MSI delineation of three high-resolution hillshade DEMs (airborne stereo-imagery, LiDAR, ArcticDEM) revealed temporal and spatial trends in these ~~multi-year~~-chronic mass-wasting features. For example, in the Tuktoyaktuk Coastal Plains, a +38% increase in active RTS counts and +69% increase in total active surface area were observed between 2004 and 2016. However, the total disturbance area of RTS did not change considerably (+3.5%) because the vast majority of active thaw slumping processes have occurred in association with past disturbances. Interpretation of thaw-driven change is thus dependent on how active RTS are defined to support disturbance inventories. Our results highlight that active RTS are tightly linked to past disturbances, underscoring the importance of inventorying these features. Third, the pre-disturbance topographies, MSI digitizations, and DEMs were integrated to explore allometric scaling relationships between RTS area and eroded volume. The power-law model indicated non-linearity in the rates of RTS expansion and intensification across scale (adj-R² of 0.85, n=1,522), but also revealed that elongated, shoreline RTS reflects outliers poorly represented by the modelling. These results indicate that variation in the allometric scaling of RTS populations is based on morphometry, terrain position, and complexity of the disturbance area, as well as the method and ontology by which slumps are inventoried. This study highlights the importance of linking field-based knowledge to feature identification and the utility of high-resolution DEMs in quantifying rates of RTS erosion beyond tracking change in the planimetric area. ~~Observations further suggested variation in depth scaling of RTS populations is based on morphometry, terrain position, and complexity of the disturbance area, as well as the method and ontology by which slumps are inventoried.~~

1 Introduction

Warming air temperatures and enhanced precipitation is altering the stability of ice-rich permafrost slopes, creating the need to track geomorphic change over a diverse array of Arctic landscapes (Kokelj and Jorgenson, 2013; Treharne et al., 2022). Monitoring remote Arctic landscapes is challenging due to limited calibration and field verification data, constraints of the available remote sensing products, and a narrow field-based understanding of rapidly changing thermokarst landforms. The frequency and areal extent of terrain affected by retrogressive thaw slumping have increased significantly in the last 25 years in ice-rich permafrost regions across northwestern Canada, Alaska, Siberia, and Tibet (Lantz and Kokelj, 2008; Segal et al., 2016; Rudy et al., 2017; Swanson and Nolan, 2018; Lewkowicz and Way, 2019; Luo et al., 2019; Runge et al., 2022). These chronic landslides grow by ablation of an ice-rich headwall (Kokelj et al., 2015; Treharne et al., 2022). Monitoring remote Arctic landscapes is challenging due to limited calibration and field verification data, constraints of the available remote sensing products, and a narrow field-based understanding of rapidly changing thermokarst landforms. The frequency and areal extent of terrain affected by retrogressive thaw slumping have increased significantly in the last 25 years in ice-rich permafrost regions across northwestern Canada, Alaska, Siberia, and Tibet (Lantz and Kokelj, 2008; Segal et al., 2016; Swanson and Nolan, 2018; Lewkowicz and Way, 2019; Luo et al., 2019; Runge et al., 2022). These chronic landslides grow by ablation of an ice-rich headwall (Lewkowicz, 1987), while the terrain and ground ice conditions combine with a suite of geomorphic and climate feedbacks to influence the nature and intensity of downslope sediment transfer and trajectories of slump growth (Fig. 1a; Kokelj et al., 2015). Cryosphere researchers have leveraged advances in remote sensing capabilities, using new optical, laser, and synthetic aperture radar data at fine spatial and temporal resolutions to better detect thaw-driven landscape change (Fraser et al., 2014; Brooker et al., 2014; Huang et al., 2020; Nitze et al., 2021; Runge et al., 2022; Xia et al., 2022). Although improved sensors and machine learning methods have rapidly advanced the capacity to detect Arctic change over broad areas, the transferability of methods remains a challenge (Nitze et al., 2021; Huang et al., 2022). The majority of studies have tracked disturbance count or area and the newest information is derived through remote sensing investigations. Fewer recent studies contribute to knowledge on the rapidly evolving morphological characteristics of slumps, and the processes and feedback accelerating or arresting thaw-driven disturbance development (Kokelj et al., 2015; Swanson and Nolan, 2018; Zwieback et al., 2018; Jones and Pollard, 2021). The dynamics of thaw slump-affected terrain heightens the need to develop robust monitoring frameworks that better. Cryosphere researchers have leveraged advances in remote sensing capabilities, using new optical, laser, and synthetic aperture radar data at fine spatial and temporal resolutions to better detect thaw-driven landscape change (Brooker et al., 2014; Huang et al., 2020; Nitze et al., 2021; Runge et al., 2022; Xia et al., 2022). Although improved sensors and machine learning methods have rapidly advanced the capacity to detect Arctic change over broad areas, the transferability of methods remains a challenge (Nitze et al., 2021; Huang et al., 2022). The majority of recent studies that track disturbance count or area derive information through remote sensing methods. Fewer recent studies investigate the processes and feedbacks influencing the evolution of thaw-driven disturbances (Kokelj et al., 2015; Swanson and Nolan, 2018; Zwieback et al., 2018; Ward Jones and Pollard, 2021). The dynamics of thaw slump-affected terrain heightens the need for monitoring frameworks that link remote sensing outputs with empirical knowledge.




Figure 1 – Conceptual diagrams of retrogressive thaw slumps showing the influence of area-volume model coefficients on disturbance morphology. The effect of the scaling exponent (δ) is illustrated by the slope schematics (a) and portrayed graphically in (b).

75


Mass-wasting inventories characterize the densityspatial and size-frequency distributions of erosional sites, providing a basis for determining dominant modes of erosion and estimating the volumetric rates of sediment, chemical, or nutrient mobilization from hillslopes (Oguchi, 1997; Campbell and Church, 2003; Tunnicliffe and Church, 2011; Kokelj et al., 2021). In temperate or mountainous regions, most landslides are rapid, discrete events that take place over a short time frame (seconds to days), so inventory studies typically assume a temporal window of relevance where the scar remains visible, before being forested over (Brardinoni et al., 2003). In contrast, retrogressive thaw slumps are chronic sites of thaw-driven erosion that modify slopes over months, years, decades, or even millennia (Burn and Friele, 1989; Lantuit and Pollard, 2008; Lacelle et al., 2010, 2015). These landslides initiate by mechanical or thermal erosion as small disturbances at the base or break in slope, along flow tracks, or as shallow slides that remove surface insulating materials and expose ice-rich permafrost at the ground surface (Lewkowicz, 1987; Lacelle et al., 2015). Ground ice exposed in the slump headwall melts causing the disturbance to enlarge and a thawed slurry to accumulate in the scar zone. Thawed materials are transferred downslope seasonally by gradual creep or episodic surface or deeper-seated flows at rates controlled by material properties and volume, local topography, and climate drivers (Kokelj et al., 2015). Warmer temperatures or greater rainfall have accelerated the evacuation of scar zone materials, driving positive feedbacks that maintain the headwall height and upslope growth potential of the retrogressive disturbance (Kokelj et al., 2015, 2021). Variations in terrain factors, the intensity of thaw-driven processes, and climate give rise to a diverse range of disturbance morphologies (Fig. 2b-e). Regardless of the setting or size of the retrogressive thaw slump, the gradual decline in headwall height with upslope growth and material accumulation results in stabilization (Burn and Lewkowicz, 1990). Diffusive processes eventually produce gentle slope-side concavities colonized by luxuriant vegetation that distinguish old thaw slump scars from the adjacent terrain (Lantz et al., 2009). The association of retrogressive thaw slump activity with favourable ground ice, slope, and initiating conditions suggests that stabilized thaw slump scars are good indicators of climate-sensitive slopes (Fig. 2b-e). Indeed,

85

90

95

100 thaw slump-affected slopes are typically subject to cycles of periodic activity punctuated by stability (Mackay, 1966; Burn and Lewkowicz, 1990; Burn, 2000; Lantuit and Pollard, 2008; Lantz and Kokelj, 2008; Kokelj et al., 2009). The climate-driven increase in polycyclic behaviour has occurred with the intensification of fluvial, coastal, and thermal erosion, and alteration of slope stability thresholds, which has modified the evolution of permafrost slopes (Kokelj et al., 2015, 2021; Jones et al., 2019);(Kokelj et al., 2015; Ward Jones et al., 2019; Kokelj et al., 2021).



105 Figure 2 – Conceptual diagram of the Multisource Slump Inventory (MSI) database and complex lakeside retrogressive thaw slump
 growth patterns in the Anderson Plain – Tuktoyaktuk Coastlands region. (a) The MSI database incorporates a time series of thaw-
 slump disturbance delineations from several high-resolution DEMs covering variable spatial extents yielding an integrated data
 product of landscape change. Inset (b) shows a sequence of DEM-based slump digitization (2004-2011) (68.529° N, -133.743° W)
 following the rapid enlargement of a lakeside thaw slump into previously undisturbed terrain. AttributionThe attribution
 110 of the hillshade DEM-based digitizations is supported by high-resolution optical imagery. (c) Polycyclic behaviour of retrogressive thaw
 slumps showing variously aged disturbances occurring within the footprint of a larger historical disturbance evident in the circa 2016
 ortho imagery (© ESRI World Imagery base map, providers: ESRI and Maxar), (d) hillshaded 2011 LiDAR DEM showing distinct
 and subtle relief of multi-aged disturbance footprints, and e), oblique photograph, 2020 (68.608° N, -133.599° W). Slump boundaries
 represent 2004 (yellow), 2011 (orange), and circa 2016 conditions (red) in insets (b) and (c), to illustrate change between respective
 115 DEM datasets (Supplement S2),(S2; Van der Sluijs and Kokelj, 2023).

The planimetric area of active thaw slump disturbances in northwestern Canada span at least 6 orders of magnitude (Kokelj et al., 2021). Small cusped disturbances with a shallow headwall that retreats upslope have typically affected 0.1 to 1 ha of terrain before stabilizing (Fig. 2b-e). However, climate-driven intensification of thaw has caused disturbances to develop that exceed 10s of ha in area and translocate millions of cubic metres of slope materials to downstream environments (Kokelj et al., 2021). The quantitative metrics change as a disturbance progresses through the stages of geomorphic development or as governing climate conditions change. The dynamic nature of retrogressive thaw slumping suggests that the regional distribution, size, and activity metrics of a permafrost landslide population at any given point in time will reflect the combination of terrain and climate conditions. Digitized spatial datasets yield snapshots of these dynamic disturbance populations (Fig. 2a), but they can vary significantly with project purpose, and factors such as slope disturbance definition, the quality, and resolution of the underlying base data layer(s), mapper experience, and field-based understanding of periglacial geomorphology in the Anthropocene (Aylsworth et al., 2000; Lantz and Kokelj, 2008; Guzzetti et al., 2012; Segal et al., 2016; Huang et al., 2020; Kokelj et al., 2021). Even ~~in cases where if the~~ geomorphic definitions are ~~unambiguousclear~~, several challenges ~~inherent~~ with remote sensing ~~detection~~ remain, such as undetected slumps (i.e., false negatives) and the inclusion of landscape features ~~or noise~~ that are not ~~actually slumps~~ a thaw slump (i.e., false positives) (~~Bernhard et al., 2020; Huang et al., 2020; Nitze et al., 2021; Runge et al., 2022~~)(Bernhard et al., 2020; Huang et al., 2020; Nitze et al., 2021; Runge et al., 2022). There may also be discrepancies among digitization efforts, where differing delineations are an ~~artifact of disparate data products and slump ontology rather than actual slump growth~~. These challenges indicate the need and opportunity for developing standardized methods to identify, delineate, and attribute thaw-driven landslides informed by field knowledge of rapidly evolving thaw-driven landforms (Jones et al., 2019; Kokelj et al., 2021)-~~artifact of disparate data products and slump ontology~~. These challenges indicate the need for standardized methods to identify, delineate, and attribute thaw-driven landslides informed by field-based knowledge.

An improved understanding of permafrost landslides also requires advancing a three-dimensional ~~spatio-temporal~~ framework for quantifying their geomorphic ~~evolution~~ and environmental impacts. The statistical relationships between landslide scar area and eroded volume provide a foundation for quantifying the geomorphic change ~~that occurs as a result of slope mass wasting~~ (Klar et al., 2011). These relationships are typically expressed as power-law models; they have been explored for a range of failure types, material properties, and geological settings across temperate landslide environments (~~Larsen et al., 2010; Klar et al., 2011~~)(Larsen et al., 2010; Klar et al., 2011). The volume of material that is thawed by retrogressive thaw slumping ~~is a critical variable that~~ scales exponentially with scar area (Fig. 1) (Kokelj et al., 2021), but how ~~such scaling~~ these relationships transfer to a diverse array of thaw-driven mass wasting features across a range of permafrost environments is unknown. Furthermore, since climate can alter the lithological controls on permafrost ~~landscape evolution~~ terrain, it is unclear if ~~these~~ scaling parameters ~~that describe thaw-driven landslides~~ will change as slope stability thresholds and permafrost physical properties are altered by a warmer and wetter circumpolar climate.

Initial An investigation of area-volume (A/V) relationships for active thaw slumps in the Beaufort Delta region of northwestern Canada indicated a power-law relationship between ~~disturbance~~ area and the total evacuated sediment volume ($R^2 = 0.9$), providing a first quantitative basis for estimating thaw-driven denudation as a function of disturbance area (Kokelj et al., 2021). There are two coefficients in the power-law relationship (Eq. 1) used to predict eroded volume (V) based on planimetric scar

area (A_s): 1) the scaling factor (a) and 2) the scaling exponent (δ). The scaling factor reflects which represents the intercept of the log-transformed linear relationship between disturbance area and volume (Fig. 1a) (Jaboyedoff et al., 2020). The; and 2) the scaling coefficient exponent (δ) which reflects the elastic distortion (or slope) of the area-volume (A/V) relation with changing area (Chaytor et al., 2009; Tseng et al., 2013). For example, if the scaling exponent is close to $\delta = 1$ then the thaw slump volume grows primarily with increasing scar area, and there is with little to no increase in the absolute scar depth (i.e., linear growth model; Fig. 1a). However, as the exponent approaches $\delta = 1.5$ the relative depth of the concavity grows in proportion to the scar area. Exponents $\delta > 1.5$ reflect thaw slumps that erode more deeply, relative to the planform area (Fig. 1a). Small differences in δ lead to substantial variance in volume predictions, thus the exponent scaling coefficient δ is critical information in quantifying the geomorphic and environmental impacts of thaw-driven mass wasting (Fig. Volume predictions are very sensitive to small differences in scaling exponent (δ) (Fig. 1b).

$$\text{Log}(V) = a + \delta (\log(A_s)) \quad (1)$$


The goal of this study is to couple knowledge of thaw slump processes and form with remote-sensing tools to develop more holistic approaches to quantifying and tracking thaw-driven mass wasting, improve the quality of mass-wasting inventories, and refine population-wide estimates of total sediment yield. To accomplish our project goal we: A) evaluate surface interpolation methods necessary to derive slump volume estimates from high-resolution Digital Elevation Models (DEM); B) improve slump delineation, morphological descriptions, and activity status using available base maps (LiDAR, ArcticDEM, satellite imagery); and C) explore the statistical properties of allometric relationships for a large sample of thaw slumps slump observations (n=2661), using the results from objectives (A) and (B). The 5,948 km² study area extends across a range of terrain types in the Beaufort Delta region of northwestern Canada. The data and analyses enabled us to explore the uncertainty in volumetric yield estimates, characterize the spectrum of thaw slump morphometry and activity levels across different geomorphic settings, and evaluate the utility of area-volume models required to quantify the cumulative landscape effects of climate change.

2 Study area and Methods

2.1 Study area

The general study area spans the Peel Plateau (Kokelj et al., 2017b), the Anderson, lower Mackenzie Plain, and the Tuktoyaktuk Coastlands physiographic regions (Rampton, 1988; Burn and Kokelj, 2009). The ice-rich terrain contains abundant excess ground ice in the form of segregated (Rampton, 1988; Burn, 1997), relict (Murton, 2005; Lacelle et al., 2019), and wedge ice (Mackay, 1990; Kokelj et al., 2014). The spatial distribution of slumps in this region confirms the abundance of ice-rich permafrost and the overall sensitivity of ice-marginal, permafrost-preserved glacial terrain (Kokelj et al., 2017a).

We examined two specific study areas: (1) the lake-rich rolling tundra of Anderson Plain/Tuktoyaktuk Coastlands (APTC) (Rampton, 1988) with an area of 3,278 km², and (2) the fluviially-incised Vittrekwa and Stony Creek watersheds of the Peel Plateau (PP) (Kokelj et al., 2017b) with an area of 2,670 km². These encompass regions host the communities of Tuktoyaktuk, Inuvik, and Fort McPherson, the Dempster and Inuvik to Tuktoyaktuk Highway corridors, and host the greatest density of historical oil and gas infrastructure in Arctic Canada (Fig. 3) (Burn and Kokelj, 2009).






Figure 3: Map of study areas (a) and DEM-based digitization of circa 2016 slumps (yellow) in the Anderson Plain/Tuktoyaktuk Coastlands (b; APTC) and Peel Plateau (c; PP; Vittrekwa and Stony Creek watersheds). Basemap is a 1:250k hillshaded Canadian DEM (CDEM; Natural Resources Canada 2013 – Open Government License). Black lines define the highways (bold) and LiDAR

extents (narrow). ArcticDEM tiles for the PP area included data voids (brown ~~diagonal lines; totaling cross-hatch; totalling~~ 510 km²; Porter et al., 2018).

2.2 Study design

200 This study developed ~~and implemented~~ methods of characterizing thaw-driven landslides to ~~support the estimation of estimate~~ disturbance volumes so that ~~the geomorphic implications of retrogressive thaw slumping impacts~~ can be ~~robustly~~ quantified. The study builds on a long-term permafrost research and landscape change monitoring program by the NWT Geological Survey and its research partners in the Mackenzie Delta region (Kokelj et al., 2005, 2009, 2013, 2015, 2017a; Lacelle et al., 2015; Lantz and Kokelj, 2008; Lantz et al., 2009; Van der Sluijs et al., 2018; Sladen et al., 2021; Kokelj et al., 2021). ~~To improve thaw slump~~ The

205 improvement of mapping methods ~~we~~ leveraged geomorphological knowledge of the study area, existing landslide inventories (Segal et al., 2016), and high-resolution ortho-mosaics and DEM datasets obtained over the past decade and a half (Table 1). The first objective (A) was to determine the uncertainty in surface interpolation methods used to reconstruct pre-erosion hillslope morphology (Table 1 ~~ba~~). The second objective (B) was to improve methods to detect, delineate, and characterize ~~thaw slump~~ RTS-affected terrain using high-resolution DEMs and optical imagery ~~to establish and develop~~ a multi-temporal dataset of

210 disturbances (Table 1 ~~a1b~~) to explore the variation in morphology and activity levels of slump-affected terrain. Utilizing ~~the~~ is DEM-derived disturbance dataset ~~developed in objective (Objective B-)~~, and applying the best interpolation method determined in objective A, we derived area-volume relationships for a large population of thaw slump disturbances and determined morphological and terrain factors that contribute to scatter in the relation (Table 1c). A methods flowchart detailing these inter-related workflows are shown in Fig. S1.

215

Table 1: Summary of research goals, remote sensing datasets and related methods sections, geographic locations, and sources of the datasets used in this study.

Research Objectives	Remote sensing datasets	Study area (Fig. 3)	Slumps (n)	Source/Reference
(A) Interpolation methods to reconstruct pre-erosion topography	2011 Airborne LiDAR, same as B	PP, APTC	34, 37	Van der Sluijs et al., (2018), Kokelj et al., (2021)
	2004 Airborne stereo-photogrammetry (3 m DEM, 0.5 m ortho)	APTC	789	NWT Centre for Geomatics (2008) ¹ <u>S2</u> ¹
	2011 Airborne LiDAR (1 m DEM, 0.2 m ortho)	PP; APTC	127, 503	Van der Sluijs et al., (2018)
(B) Development of multi-temporal slump dataset	Circa 2016 stereo satellite DEM (2 m, ArcticDEM) and circa 2017 base imagery layer (ESRI World Imagery; 0.5 m)	PP; APTC	457, 785	Porter et al., (2018)
	1984-2019 Landsat time-series of tasseled cap indices	PP; APTC	N/a	NWT Centre for Geomatics (2021) ² <u>S2</u> ²
(C) A/V model and outlier detection	Same as (B)	PP; APTC	2,661	

220 ¹ Mackenzie Valley Airphoto Project (MVAP) of Indian and Northern Affairs Canada, now Government of Northwest Territories with NWT Centre for Geomatics as publisher.

² Method based on Fraser et al., (2014).

2.2.1 Pre-disturbance terrain methods

225 The volume of a thaw slump scar is the product of thaw subsidence from thermoerosion and ground ice loss (Lewkowicz, 1987), and downslope sediment transport (Kokelj et al., 2015, 2021; Figs. 2, 3). The volume of materials displaced by thaw slump activity can be estimated through DEM differencing (~~Lantuit and Pollard, 2008; Van der Sluijs et al., 2018; Kokelj et al., 2021;~~

Turner et al., 2021)(Lantuit and Pollard, 2008; Van der Sluijs et al., 2018; Kokelj et al., 2021; Turner et al., 2021). In cases where both DEMs reflect hillslopes in the disturbed state the derived difference reflects the episodic sediment yield (e.g., inter-survey) whereas the total evacuated sediment volume of a thaw slump can be determined if a 'pre-disturbance' DEM is available (i.e., since initiation of the feature). Typically these multi-decadal disturbances predate high-resolution DEMs, requiring reconstruction of the disturbed topography through manual or semi-automated DEM void-filling techniques (ten Brink et al., 2006; Van der Sluijs et al., 2018; Kokelj et al., 2021). To build robust regional slump-volume datasets we investigated the effectiveness of surface interpolation methods for obtaining pre-disturbance DEMs and compared how well the reconstructed ice-rich permafrost terrain surfaces conform to the original topography. To develop a dataset that would enable us to evaluate the performance of interpolation methods we generated a series of slump-shaped data voids that were randomly placed on undisturbed terrain considered conducive to slump development within the 2011 LiDAR DEM extents (Lacelle et al., 2015; areas with 2-12° slopes and within the maximum extent of Laurentide Ice Sheet). We ~~applied~~used the digitizations and size distribution of active-slump populations in fluviially-incised settings to generate random void areas in the PP study area (n=34, Kokelj et al., 2021). The randomization was iterated 30 times so that the 34 ~~slump-shapes~~voids were each positioned in 30 different terrain locations. A similar approach was applied to assess reconstruction fidelity within predominantly lacustrine environments in the APTC area (n=37, Kokelj et al., 2021). Data voids were only retained if the random placement was within 400 m of a lake. These procedures produced a dataset consisting of 1,020 voids for the PP area (n=34 slumps and 30 iterations) and 359 voids in the APTC area (study total n = 1,379).

The magnitude of interpolation error can vary considerably among methods due to underlying model assumptions, parameterization, sensitivities to the quality of the input data, and the morphologic setting of the void (Reuter et al., 2007; Bergonse and Reis, 2015; Boreggio et al., 2018). Eight interpolator suites commonly available and described in GIS software manuals and used for digital terrain modelling (ArcGIS Pro™ v.2.7) were identified to reconstruct the synthetic DEM data voids: 1) Inverse Distance Weighted (IDW), 2) TIN-to-Raster (TR), 3) Regularized Spline with low weights (RSL), 4) Regularized Spline with high weights (RSH), 5) Spline with tension (ST), 6) Empirical Bayesian Kriging with no data transformation (EBK), 7) EBK with empirical data transformation (~~EM~~EBK-EMP), and finally 8) EBK with empirical transformation and detrended semivariograms (EBK-EMPD). The eight different interpolation suites were tested with a range of parameterizations (a total of 32 interpolation methods; Table ~~S2S3~~) to assess performance within and between methods. The topography of each void was reconstructed independently following an automated Python workflow (S4) that buffered each void by 50 m, retained the pixels outside of the void, and converted them to boundary elevations that were used by each of the 32 interpolation methods. Then re-interpolated void surfaces (n=1,379) were compared with the original LiDAR terrain surface to test the accuracy of surface interpolators (Reuter et al., 2007). Elevation differences between true (actual LiDAR DEM) and modelled (interpolated terrain) elevations were summarized for each void by calculating the root mean square difference (RMSD), mean absolute error (MAE), and the summed absolute topographic difference (T_{sum}). In this case, the RMSD is a measure of error (in metres) describing how well the modelled terrain fits the actual terrain. MAE measures the average absolute difference of the error. T_{sum} is an indicator of three-dimensional topographic uncertainty expected for a reconstructed void surface in m^3 , with 0 m^3 reflecting a perfect fit. Non-parametric statistical testing (Kruskal–Wallis and Dunn's post hoc tests with Bonferroni adjustment) was implemented to assess potential differences in RMSD between interpolator suites and between individual methods.

2.2.2 Multisource Slump Inventory (MSI)

In this study we developed a Multisource Slump Inventory (MSI), using several high-resolution observation platforms and data sources to assemble an “evolving” geodatabase of digitized and attributed thaw-driven mass-wasting features and volumes (Fig. 2a). An MSI is Slump disturbances were identified and digitized at very large mapping scales ($>1:2,000$) utilizing compilations of the best available high-resolution hillshade DEMs (≤ 3 m; see Supplement S2 for dataset details) and imagery (≤ 0.5 m). Due to the evolving nature of thaw slumps an approach based on a multi-temporal inventory map (Guzzetti et al., 2012), where disturbances are identified, digitized, and attributed at very large mapping scales ($>1:2,000$) utilizing compilations of the best available hillshade DEMs (≤ 3 m; see Supplement S1 for dataset details) and high-resolution imagery (≤ 0.5 m). The MSI approach is pursued was pursued, namely the Multisource Slump Inventory (MSI). The MSI approach was implemented to 1) acquire temporally consistent, high-resolution digitizations for each slump-affected slope area based on the best available DEM; (Fig. 2a); 2) obtain more descriptive information on the geomorphic activity to better track disturbance evolution by inspecting high-resolution imagery; and 3) reduce the number of small undetected, inactive or shallow slumps as typically the case with landslide inventories (Stark and Hovius, 2001; Guzzetti et al., 2012). We note that many inventories in this study region have explicitly mapped disturbances assessed to be active or recently active (Brooker et al., 2014; Segal et al., 2016; Kokelj et al., 2021) using coarse to high-resolution imagery. The use of high-resolution hillshade DEMs in this study enabled detailed digitization of disturbances and numerous revegetated stable slump scars to be identified and mapped with greater confidence, which has been constrained in the past when using optical or coarser base-data layers (Fig. 2d). The inventory pursued MSI included mapping partially or fully-vegetated old and ancient slump scar areas because they are important indicators of ice-rich thaw sensitive terrain (Fig. 2b-e) (Kokelj et al., 2009).

Digitized features within the MSI are composite entities, whereby planimetric area and activity level attributes can be updated in reference to past observations when newer hillshade DEMs and supporting high-resolution optical imagery become available (Fig. 2a). A case study for the APTC region was formulated to demonstrate how the MSI helps in conveying and linking field-based observations of slump process to a digital inventory intended to track the area, volume, and dynamics of thaw-driven landslides through time. The APTC MSI was compiled using high-resolution ortho-mosaics and DEMs for three observation periods, whereas the PP database incorporated only two observation periods (Tables 1 and S1). Interpretations were informed by fieldwork and previous point- and polygon-based slump detections, most of which focused on active disturbance areas identified using optical imagery (Lantz and Kokelj, 2008; Segal et al., 2016; Kokelj et al., 2021). For each observation period, the MSI disturbance polygons comprised a scar zone, consisting of active and inactive sections, as well as a deposition area when this was indistinguishable from the scar zone. In cases where a debris tongue could be identified as a downslope depositional feature distinct from the active scar zone, only the scar zone was digitized (Kokelj et al., 2021). Shallow slumps where lower slopes revegetate while headwall retreat remains active were included in this inventory for compatibility with those recognized by Lacelle et al., (2010). In addition, shallow landslides are common in fluvially incised environments of the study area, and with the intensification of thaw-driven processes over the past decade, many of these landslides continue to enlarge upslope by retrogressive failure resulting in their classification as thaw slumps. Several steps were undertaken to integrate the datasets (Fig. 2) by applying consistent delineation rules when different data sources were utilized (Supplemental S2). Rather than an independent delineation effort for each observation year (2004, 2011, 2016) the DEMs belonging to the observation periods were used in reference to each other to iteratively delineate slump boundaries by manually identifying stable and active edges (Fig. 2). This ensured that changes in the area reflected headwall retreat and not edge artifacts that commonly arise from independent digitization efforts or data from sources with different resolutions. Over the MSI timescale, slump activity or polycyclic behavior

305 means that slump-disturbed areas will only increase through time, and therefore a slump cannot drop out of the inventory once it is detected, even if it is old. Exceptions may potentially arise in situations where shoreline erosion equals or outpaces upslope slump enlargement, leading to similar or declining planimetric disturbance areas (Obu et al., 2016).

310 The second step of the MSI involved using near-coincident high-resolution imagery to attribute activity levels of the disturbances digitized from the DEM so that landslide dynamics could be tracked through time. Attributes include landscape descriptors (surficial geology, geomorphology). Steps were undertaken to integrate the datasets (Fig. 2) by applying delineation rules formulated for partial or fully-stabilized slumps as well as highly polycyclic slumps. These MSI methods are described in detail in Van der Sluijs and Kokelj (2023) so only a brief summary is provided here. The delineation rules were based on: 1) iterative delineation with reference to past data, 2) slump-disturbed areas can only increase through time, 3) once detected a slump feature cannot drop out of the inventory. Each delineated feature was attributed by landscape descriptors, two-dimensional geometry estimates, and three-dimensional hypsometry estimates, along with a characterization of activity levels (Table 2). The surface area percentage of activity (in 10% increments) was determined by DEM evidence of near-vertical headwalls, and the optical imagery was inspected for headwall shadows with supersaturated materials at the base of the headwall, fresh cohesive or plastic mud flows in the form of unvegetated wet lobes, and the zonal contrast of wet areas (darker) with areas of stabilized bare ground (brighter) or vegetated terrain (Lantuit and Pollard, 2008; Brooker et al., 2014). The imagery datasets reflected peak summer conditions (July/August) for comparable observations of activity over time. It should be noted that the based on surface area percentage of activity (in 10% increments) using visual clues in the optical imagery and hillshaded DEM (Table 2; Fig. 6c; Van der Sluijs and Kokelj, 2023). Percentages were also grouped into three classes to facilitate summary statistics among stable slumps (0%), minimally active slumps (10%) and moderately-to-highly active slumps (20-90%). It should be noted that the total slump-affected area of disturbances in the MSI may not change if the recent activity is taking place entirely within the footprint of an older disturbance (Fig. 2c). However, in such instances the percent activity estimates would indicate an intensification of slumping (e.g., from a 0% inactive slump to a 20% reactivated slump). The “active surface area” of an individual disturbance was based on the total slump-affected area multiplied by the estimated percentage of the active area.

Table 2. Summary of slump geomorphic attributes included in this inventory.

Attribute	Description	Interpretation or notes
UniqueID	Sequential number of each landslide entry in the database, per region	
Region	APTC or PP	
Year	Year of observation	
Datasource	MVAP, LiDAR or ArcticDEM	
Location	Easting and northing of slump centre coordinate	
Geometry ¹		
<i>Area (A)</i>	The projected area enclosed by slump boundary	
<i>Perimeter (P)</i> [*]	Length of the horizontal projection of slump boundary	
<i>Length (L)</i>	Length of the long axis	Equal to the major axis, the longest line that can be drawn through the object unrelated to the slope
<i>Width (W)</i> [*]	$W = A/L$	
<i>Orientation</i> [*]	Orientation (azimuth) of the long axis (L)	
<i>Elongation ratio</i> [*]	$1.128\sqrt{A/L}$	Values range from 0 (highly elongated) to 1 (circular)
<i>Circulatory ratio</i>	$4\pi A/P^2$	Values <0.5 indicate elongation. Values near 1 indicate high circularity
<i>Compactness coefficient</i>	$0.282P/\sqrt{A}$	The ratio of the perimeter of a form to the circumference of the circular area, which equals the area of the form
<i>Form factor</i>	A/L^2	Perfect circle = 0.754; smaller values more elongated
<i>Shape factor</i>	1/Form factor	

Hypsometry		
<i>Elevation</i> ^{1,*}	Minimum, maximum and mean elevation of the slump	
<i>Slope</i> ^{1,*}	Minimum, maximum and mean slope of the slump	<u>In degrees</u>
<i>Aspect</i> ^{1,*}	Mean aspect	
<i>TRI</i> ^{1,*}	Minimum, maximum, range, mean and stdev of Terrain Ruggedness Index (Riley et al., 1999)	Expresses amount of elevation difference between adjacent cells within a slump
<i>Curvature</i> ^{2,*}	Geometric normal curvature along the slope line, calculated for the entire digitized feature. Curvature metrics provided in distribution percentiles (p10, p20, p50, etc.).	Visualized as the shape of a vertical (profile) cross-section through the slump. Positive (negative) values indicate convex (<u>concave</u>) surfaces, negative values indicate concave surfaces.
Surficial geology ^{3,*}		
	Alluvial, colluvial, eolian, fluvial, glaciofluvial, lacustrine, morainal, organic, bedrock, slump, upland, marine	
Geomorphology [*]		
Activity [*]	Percent area attributed to active slump processes, in 10 <u>In 10 percent increments</u>	

¹ Derived using Morphometry Assessment Tools for ArcGIS (Gudowicz and Paluszkiwicz, 2021).

330 ² Derived using Surface Parameters tool in ArcGIS Pro based on 10 m neighborhood distance.

³ Derived from Smith and Duong (2012) and Cote et al., (2013).

* Variable used in random forest modelling after removal of highly correlated variables (Sect. 2.2.3).

2.2.3 Area-volume (A/V) allometry and outlier detection

In this study, we explored ~~two distinct~~ relationships between thaw slump area and two distinct volume parameters. First, we
335 examined relationships between slump area and the difference between real and modelled slump topography (T_{sum}) to determine the uncertainty of reconstructed surfaces and how these scale with slump area (Objective A). The slump area dataset consisted of a randomly placed population of slump voids following procedures described above (Sect. 2.2.1), with uncertainty in the surface reconstruction expressed in both absolute and relative terms. Secondly, we explored the relationships between slump area and
340 the modelled pre-disturbance DEMs determined using the best performing interpolator (Objective A). For this analysis, data on slump area, activity level, and disturbance volume from the MSI (Objective B) were analyzed in a multiple linear regression framework. Slump area and volume data were logarithmically transformed to meet assumptions of normality. The software package R Statistics (v. 4.0.2) was used to produce descriptive statistics and regression analyses. Slumps were digitized from up to 3 different periods, according to the availability of various DEM data sources. Temporal auto-correlation between multiple
345 observations of the area, activity level, and volume for the same slump in the MSI dataset was addressed in the multiple regression by including a categorical ‘Datasource’ term in the models (Table 2).




We explored A/V model performance and residuals to determine whether the nature of the outliers could be described to inform constraints on the model’s application. Areas affected by thaw slumping that yielded problematic volume estimates were
350 identified as: (1) Thaw slumps where the modelled pre-disturbance elevations were lower than post-disturbance elevations, producing ~~differenced DEMs with~~ an *incorrect* positive volumetric change, and (2) thaw slumps with a *correct* negative mean depth of thaw and volume, but with a volume estimate below the T_{sum} uncertainty threshold, determined from testing surface interpolation methods described in Sect. 2.2.1. The morphometry and geomorphic setting of the volumetric outliers were examined in a random forest (RF) classification (Breiman, 2001) to explore the attributes that most commonly distinguished
355 these volumetric outliers relative to the entire thaw slump population. In our analysis, RF was used in a binary schema to separate volumetric inliers from outliers using the *randomForests* package in R, based on a randomly balanced selection of 80% of the MSI observations ($n = 2,130$, retaining 20%, or 531 observations, for independent validation), 1000 trees, and 6 explanatory variables randomly sampled at each split. Highly correlated explanatory variables from the MSI, defined as an absolute Spearman’s correlation of 0.75 or higher, were removed before classification (Table 2). The influence of the three


360 explanatory variables identified by RF variable importance with the largest overall impact on accuracy was assessed using partial
dependence plots showing the probability of being classified as a volumetric inlier or outlier.

3 Results

3.1 Pre-disturbance terrain methods

To determine the most desirable interpolation techniques for reconstructing terrain surfaces affected by thaw slumping we
365 examined the performance of 32 surface interpolation methods by comparing differences between observed (actual) and
modelled (interpolated terrain) surfaces for 1,379 terrain voids representing a population of simulated slump disturbances in the
study regions. Interpolation performance is parameterization-dependent, so methods were grouped into suites to identify
variation in accuracy associated with general algorithmic implementations. Kruskal-Wallis test results indicated significant
differences in RMSD distributions between interpolator suites ($H(7) = 7,637, p < 0.001$). The TIN-to-Raster (TR) suite exhibited
370 the lowest RMSDs (median: 0.3 m, 90th percentile: 1.0 m; Fig. 4a), while Empirical Bayesian Kriging (EBK) with data
transformation and detrended semivariograms (EBK-EMPD) achieved second-lowest RMSDs (median: 0.4 m, 90th percentile:
1.8 m; Fig. 4a). Dunn's post hoc tests indicated that TR's RMSD distribution was significantly different from all other suites ($p <$
 0.001). For all interpolator suites, the RMSD increased non-linearly with the void area, with the TR suite exhibiting the most
gradual increases (Fig. 4b). Even though RMSD increased with void size for all suites, relative uncertainties for all interpolators
375 declined with the void area (Fig. 4c). We further evaluated the Natural Neighbours (NN) and Linear (LIN) parameterizations
within the TR Suite. Statistical testing indicated that RMSD ($H(1) = 2.25, p = 0.133$), Mean Absolute Error distributions (MAE;
 $H(1) = 2.26, p = 0.132$) or summed topographic difference distributions (T_{sum} ; $H(1) = 0.145, p = 0.703$) were not significantly
different from one another, but were significantly different from parameterization results in all other interpolation methods
380 (Table [S2S3](#), Figs. S3, S4). These findings show that the NN had the most robust performance for reconstructing topography
affected by active retrogressive thaw slumping in fluvially-incised glaciated permafrost terrain.






385 **Figure 4: (a) Boxplot of differences between actual and modelled elevation (expressed as Root Mean Square Difference; RMSD) for**
 390 **each interpolator suite and (b) scatterplot between void surface area and RMSD with smoothing line indicating area-dependent**
uncertainty of the respective interpolator suites, and (c) scatterplot between void area and normalized RMSD (RMSD / Area * 1,000)
with loess smoothing line indicating how relative uncertainty of the respective interpolator suites varies with the void area. Together
these graphs indicate that the Topo-to-Raster (TR) suite of methods (linear or natural neighbour interpolation) achieved the lowest
RMSDs among the interpolation suites and exerted the weakest influence of void surface area on RMSD for both study areas. Due to
outliers in the more poorly performing ST suite the axis portraying RMSD was limited to 6 m in (a) and 2 m in (b), respectively.

Using the measure of void area and summed topographic difference (T_{sum}), the precision of interpolated surface estimates for a given disturbance area could be determined, and an uncertainty estimate could then be assigned to the modelled slump volumes. The synthetic disturbances were grouped into three void area classes, and as anticipated, there were significant increases in T_{sum} with increasing void area ($H(2) = 839, p < 0.001$; Fig. 5a). Figure 5b shows a linear model fit through the logarithmically transformed void area (A_s) and summed topographic difference estimates (T_{sum}), which is described by a power-law relationship (Eq. 2).

$$\text{Log}(T_{sum}) = -2.10(\pm 0.04; 95\% \text{ C.I.}) + 1.38(\pm 0.01; 95\% \text{ C.I.}) \cdot (\text{Log}(A_s)) \quad (2)$$

400 Applying Eq. 2 to the real-world slump samples from Kokelj et al., (2021) indicated that the influence of pre-disturbance interpolation on derived slump volume estimates was relatively small, with uncertainties <10% for large slumps and typically

within 10-20% of small to medium disturbances (Fig. 5c), which is expressed as an inverse association between slump volume and relative volumetric error (Fig. 5d).








Figure 5: Results showing performance of the NN interpolator in modelling pre-disturbance terrain. (a) Box-and-whisker plots showing medians, 25% and 75% quantiles of summed topographic difference by void area class. (b) Relationship between void area and summed topographic differences (T_{sum}). The regression is described by Eq. 2 with a 95% prediction interval for future observations (dashed lines). (c) Box-and-whisker plots showing medians, 25% and 75% quantiles of volumetric difference expressed as a percentage based on modelled volume of thaw slumps, and (d) relationship between modelled slump volume and relative volumetric uncertainty based on data from Kokelj et al. (2021).

3.2 Multisource Slump Inventory (MSI)

The digitization and attribution of slump-affected terrain utilizing 2016 DEMs and following MSI methods is summarized for the two study areas ($n=1242$) (Fig. 6). The activity levels of slopes affected by thaw slumping are characterized by similar cumulative distribution functions for the two study regions (Fig. 7a). In PP and APTC regions 40 % to 60 % of the digitized slump affected areas were classified as inactive and an additional 20-30% of disturbed areas were estimated to have only 10% geomorphically active surface (Fig. 7a). ~~Less than 20~~ Between 80 % to 90 % of the digitized features were characterized by bare, geomorphically active surfaces ~~exceeding~~ contributing less than 20 % of the total scar area. In many cases, active headwall retreat was associated with a small, cusped-shaped active scar zone comprised of mud slurry and a larger vegetated accumulation zone on the mid and lower slope where materials have accumulated (Fig. 6c). This common thaw slump morphology is contrasted by the fewer, highly active disturbances with high headwalls and dynamic scar surfaces where large volumes of material are being actively transported from the slope to a downstream environment (Fig. 6c). Singular scars can grow into major ‘complexes’ with multiple lobate scar zones. These geomorphically distinct features are orders of magnitude larger in size, yielding deep cavities and significant downslope debris tongue deposits. They reflect a tipping point in the thaw-driven evolution of ice-rich terrain and have been referred to as “mega slumps” (Kokelj et al., 2015).





430 **Figure 6: Plate of examples of slump form, activity ratings, and morphological variability as documented in the MSI. Morphological and environmental differences between active slumps with compact scar zones and inactive slumps with highly elongated scar zones are shown on 2011 hillshade DEM in insets (a) and (b), respectively. In Inset (c) PP slumps are shown exhibiting various activity levels circa 2016 conditions ((© ESRI World Imagery base map, providers: ESRI and Maxar). In cases of small lateral differences in headwall positions between hillshade ArcticDEM and ESRI World Imagery the final boundaries were digitized based on hillshade ArcticDEM because elevations outside the slump scar zone were used for pre-disturbance surface interpolation (Sec 3.1). The extent of white dots in (c) indicates the surfaces affected by active mass-wasting processes used to estimate percent activity.**

435




Figure 7: The circa 2016 summary statistics showing the a) cumulative distribution of slump count by activity rating for the two study regions. Insets b-d show frequency density plots of the activity levels of slump-affected terrain in the two study regions for scar zone area, elongation ratio, and terrain ruggedness (TRI), respectively.

445 In the APTC study area 785 features defined as thaw slump affected areas were digitized (24.0 slumps/100 km²), but only 40 %
or 313 (9.5/100 km²) areas are sites where the *process* of retrogressive thaw slumping was active. For the PP study area, 457
features were digitized in areas where ArcticDEM was available (21.1 slumps/100 km²; Fig. 3), with ~~a final count of 272 active~~
features (12.6/100 km²) ~~when fully stabilized slumps were excluded~~, or 60% of the total population. When slumps with less than
≤10 % active area were removed, disturbance density decreased further to 4.4/100 km² for APTC and 6.4/100 km² for PP. Taken
together the MSI statistics highlighted that in 2016 the majority of slump-affected slope areas in the APTC and PP regions
consisted of stabilized disturbances or areas of active thaw slumping within the footprint of a much larger, geomorphically
inactive disturbance area.

450

~~To demonstrate the potential application of the MSI method for regional monitoring we examine the time series of disturbances
for the APTC study area. Multi-temporal results for the APTC region demonstrated how the MSI helps in conveying and linking
field-based observations of slump process to a digital inventory intended to track the area, volume, and dynamics of thaw-driven
landslides through time.~~ Figure 8a, and b shows that the regional density of thaw slumps (24.1 slumps/100 km² in 2004 to 24.0
455 slumps/100 km² in 2016) and total disturbed area of thaw slump affected terrain (23.7 ha slump affected terrain/100 km² to 24.7
ha slump affected terrain/100 km²) varied little between 2004 and 2016 (+ 3.5%). However, the regional density of active thaw
slumps (activity ≥ 0%) increased by 38% (~~from~~ 6.9 slumps/100 km² in 2004 to 9.5 slumps/100 km² in 2016)~~;~~, and the estimated
area affected by active slumping increased by 69% (~~from~~ 1.4 ha/100 km² to 2.4 ha/100 km²)~~;~~. The density of moderate-to-highly
active slumps increased by 49% from 3.0 slumps/100 km² to 4.4 slumps/100 km². The results reinforce that slump activity in the
460 APTC region between 2004 and 2016 has increased and occurred within areas of past disturbance.




Figure 8: Temporal change in (a) count and (b) total active area of slump-affected terrain in the APTC study region documented by MSI methods. (c) Examples of APTC slump activity levels and trends in the MSI (e), where past disturbances are reactivated.

We also use the MSI dataset to examine variation in the morphology of RTS-affected terrain across the study regions. Slump area varied between regions and among slump activity ratings (Kruskal-Wallis ($H(5) = 109, p < 0.001$)). As reported in Kokelj et al., (2021), slumps in the PP region were on average larger than those in the APTC region, and exhibited a greater standard deviation (Table 3; Fig. 7b). Dunn's post hoc tests indicated that disturbance areas with active slumping were significantly larger than completely stable scar areas in each region (Table 3). Statistical distributions of elongation ratio, expressing the overall plan form of the disturbance, were also significantly different between regions and among slump activity ratings (Kruskal-Wallis ($H(5) = 322, p < 0.001$)). Slump-affected areas predominantly in the PP region had considerably higher elongation ratios (medians ranging between 0.67-0.73) compared to the lacustrine or coastal slumps located in the APTC region (medians: 0.50-0.54). A higher ratio indicating a more circular or compact morphology increased with slump activity level in the PP (Fig. 7c). Activity level did not have the same influence on the morphometry of shoreline disturbances in the APTC (Table 3). Hypsometric indices derived from the DEM also varied between regions and activity groups. For example, statistical distributions of the Terrain Ruggedness Index (TRI; Riley et al., 1999), describing the amount of local terrain relief in a slump, were significantly different between regions and among slump activity ratings (Kruskal-Wallis ($H(5) = 135, p < 0.001$)). However, significant differences in TRI were only observed at the regional PP versus APTC level, and further only among PP slumps grouped by activity (Table 3). The highest average TRI values were observed for moderate-to-highly active slumps (Fig. 7d). These observations highlight diversity in the morphology of slump-affected terrain, which varies with region and activity

480 level, and when stable and larger polycyclic disturbance footprints are delineated, complex morphologies beyond the simple, “cusped” slope disturbance form emerge.

Table 3: Summary statistics for select two- and three-dimensional metrics of the MSI ¹

	APTC			PP			Kruskal-Wallis tests (df=5)	
	0%	10%	20-90%	0%	10%	20-90%	Chi-square	p-value
Area (m)								
<i>Median</i>	4,729 ^a	6,977 ^b	5,842 ^b	8,156 ^b	12,794 ^c	13,944 ^c	109.36	< 0.001
<i>SD</i>	12,584	13,786	18,691	27,583	58,690	38,147		
<i>Skewness</i>	4.3	2.2	3.2	5.4	7.3	4.1		
<i>Kurtosis</i>	29.9	8.3	15.3	40.0	68.4	26.1		
Elongation ratio								
<i>Median</i>	0.54 ^a	0.50 ^b	0.50 ^b	0.67 ^c	0.70 ^c	0.73 ^d	321.77	< 0.001
<i>SD</i>	0.16	0.15	0.16	0.10	0.11	0.10		
<i>Skewness</i>	0.1	0.0	0.0	-0.1	-0.3	-0.1		
<i>Kurtosis</i>	2.0	2.4	2.1	2.8	2.6	2.3		
TRI (mean)								
<i>Median</i>	0.37 ^a	0.36 ^a	0.40 ^a	0.27 ^b	0.26 ^c	0.30 ^c	134.79	< 0.001
<i>SD</i>	0.15	0.15	0.17	0.08	0.15	0.17		
<i>Skewness</i>	0.7	1.5	1.0	1.3	1.5	1.6		
<i>Kurtosis</i>	3.7	5.9	4.3	5.1	6.2	6.0		

¹ Coding (a,b,c,d) with letters of the groups that are significantly different based on Dunn’s posthoc.

485 3.3 Area-volume (A/V) allometry and outliers

We evaluated the area-volume relationship for a large population of thaw slumps in the two study areas (Fig. 1) utilizing disturbance area and activity attributes from the MSI, and the volumes estimated by differencing the disturbed terrain surfaces against the modelled pre-disturbance DEMs derived using the NN interpolator (Fig. 5). Digitized slump area was the continuous predictor variable, and ‘percent activity’ and ‘Datasource’ were categorical variables used to model scar volumes of slopes affected by retrogressive thaw slumping (Table 2). Volume estimates that yielded a negative value indicating an increase in terrain relief (n=855), which is not possible, were omitted from this analysis, yielding a total sample of 1,806 slump-affected areas with an associated disturbance volume. The resulting model, with an adjusted R² model fit (adj-R²) of 0.66 (*p* < 0.001), was characterized by non-normally distributed residuals, heteroscedasticity, as well as data points that unduly influenced the relationship (Table 3, Model #1). Both ‘Datasource’ and percent activity were found to be significant explanatory variables (*p* < 0.001; Table 3). Exclusion of the datasource and activity terms did not appreciably affect the adj-R² or change the scaling coefficients, or regression diagnostics of the simplified model described by Eq. 3 (adj-R² of 0.65; Table 3, Model #2; Fig. 9a). These model results and diagnostics highlighted the challenges of thaw slump volume predictions for complex disturbances that occur across a diverse range of terrain, slump morphologies and activity levels captured by the MSI.

$$500 \quad \text{Log}(V) = -1.45 (\pm 0.09; 95\% \text{ C.I.}) + 1.30 (\pm 0.02; 95\% \text{ C.I.}) \cdot (\text{Log}(As)) \quad (3)$$

To better understand the application of the A/V models for predicting geomorphic change and volume of permafrost thawed as a result of retrogressive thaw slump disturbances we explore the characteristics of outliers and inliers in the relationship described by Eq. 3 (Fig. 9a). The initial MSI inventory of the study areas yielded 2,661 observations of slump affected terrain, but the NN re-interpolation procedure resulted in 855 occurrences of terrain surface estimates predicting a net positive change in post-

disturbance surface elevation rather than the surface lowering, which is the logical result of retrogressive thaw slumps-[in ice-rich terrain](#). Therefore we determined outliers as thaw slump disturbance areas characterized by incorrect, positive estimates of disturbance volumes (n=855), and those disturbances with a volume estimate that did not exceed the T_{sum} uncertainty threshold defined by the relationship shown in Fig. 5b and Eq. 2 (n=284). By this criteria, inliers were slump-affected areas with modelled volumes exceeding the uncertainty threshold associated with terrain surface reconstruction (Fig. 5b, n=1522).




Figure 9: Relationship between (a) slump area and slump fill volume. Scatterplot in (a) shows the regression models #2 and #4 (Table 3) as well as volumetric inliers (blue) and outliers (green), with observed model improvements when volumetric outliers were filtered out. (b) Variable importance in the random forest classification (predicting the probability of outliers) measured as the mean decrease in model accuracy scaled by the standard error of the change in model accuracy, and (c) Partial dependence plots for the top three variables in order of importance: median profile curvature (50th percentile), mean aspect and elongation ratio. The dashed red lines on the Y-axis in (c) indicate the point at which median curvature (> 0), mean aspect (less than 50 degrees and greater than 270 degrees), and highly elongated slumps (e.g., < 0.5) were more likely to be classified as volumetric outliers.

The RF classification model to explore the morphological or environmental factors that discriminate between volumetric inliers (n=1522) and outliers (n=1139) indicated that median profile curvature, mean aspect, and elongation ratio were the top three predictors distinguishing the two groups (Fig. 9b). This binary classification model had an overall accuracy of 84% (95% CI: 80 to 87%), and classified slumps as outlier and inlier with user's accuracy of 82% and 85%, respectively. Partial dependence plots showed that disturbance areas characterized by a positive median profile curvature that were highly elongated, and for our study area, with a mean aspect ranging from 270 degrees (west) to 50 degrees (northeast) were more likely to be classified as volumetric outliers (Fig 9c). These characteristics were predominantly associated with slump-affected areas extending along shorelines [in the APTC \(Table 3\)](#) where wave-erosion and shoreline retreat has driven long-term polycyclic activity (Kokelj et al., 2009; Ramage et al., 2017) and where active slumping often occurs within larger inactive shoreline scar areas digitized as a single polycyclic disturbance in the MSI (Figs. 2, 6b). The results highlight that the pre-disturbance surface interpolation, and

530 therefore the estimation of volume, did not perform well with disturbance areas that were elongated perpendicular to the local slope common to slope disturbances along eroding shorelines in the study region.

Removal of the volumetric outliers that did not exceed the T_{sum} uncertainty threshold improved model fit from an $adj-R^2 = 0.66$ to an $adj-R^2$ of 0.86 ($n = 1,522$; Table 3, Model #3) for the model including percent activity and datasource, and an $adj-R^2$ of 0.85 for the simplest model between slump area and volume (Model #4; Eq. 4). For the full model, percent activity was a significant secondary explanatory variable ($p < 0.001$), but Datasource ($p = 0.354$) was not. Both models had normally distributed residuals and no evidence of heteroscedasticity or influential samples.

$$\text{Log}(V) = -1.50 (\pm 0.06; 95\% \text{ C.I.}) + 1.36 (\pm 0.01; 95\% \text{ C.I.}) \cdot (\text{Log}(A_s)) \quad (4)$$

540 **Table 4: Statistics of multiple linear regression models.**

Model	Sample (n)	Variables	Unstandardized Coefficients				Model			Tests	
			B	S.E.	t	Pr(>t)	Adj-R ²	S.E.	P value	K-S ²	BP ³
1	1,806	(Constant)	-1.4618	0.0868	-16.83	<0.001	0.66	0.484	<0.001	<0.001	<0.001
		Log(Area)	1.2771	0.0226	56.59	<0.001					
		Datasource ¹									
		LiDAR	0.0838	0.0312	2.689	0.007					
		ArcticDEM	0.0863	0.0288	3.001	0.003					
2	1,806	(Constant)	-1.4456	0.0862	-16.76	<0.001	0.65	0.492	<0.001	<0.001	<0.001
		Log(Area)	1.3013	0.0224	58.10	<0.001					
3	1,522	(Constant)	-1.4809	0.0549	-26.97	<0.001	0.86	0.287	<0.001	0.1697	0.6540
		Log(Area)	1.3407	0.0144	93.28	<0.001					
		Datasource ¹									
		LiDAR	0.0028	0.0203	0.140	0.889					
		ArcticDEM	0.0049	0.0190	0.256	0.798					
4	1,522	(Constant)	-1.4992	0.0552	-27.15	<0.001	0.85	0.296	<0.001	0.1324	0.3291
		Log(Area)	1.3577	0.0144	94.33	<0.001					

¹ DEM Datasource: 2004 MVAP, 2011 LiDAR or circa 2016 ArcticDEM, a categorical variable included to address temporal auto-correlation between multiple observations of the same slump across the DEM datasets and digitization periods.

² Kolmogorov-Smirnov test for normality of residuals (p-value).

³ Breusch-Pagan test for constant variance or heteroscedasticity (p-value).

545 4 Discussion

4.1 Pre-disturbance terrain methods

Thaw slumps are typically chronic, multi-year mass-wasting features, an important distinction from landslide sites most landslides that are typically have associated with a single triggering event. The, and the majority of scar volume from such features represents a one-time translocation of mass on the landscape. For our thaw slump inventory, we are considering both inter-survey changes, as well as volume eroded since initiation, which could be centennial time-scales or greater. Obtaining the latter quantity volume of a chronic erosion site can be challenging if the slump scar pre-dates regional topographic surveys. In this study, the performance of common surface interpolation methods varied markedly for reconstructing slump-affected slopes in the Beaufort Delta region. Utilizing data from fluviially-incised ice-rich terrain (Kokelj et al., 2021), the approximation of pre-disturbance topography using Natural Neighbour (NN) interpolation achieved satisfactory results without complex parameterization (Fig. 4a). In contrast, Empirical Bayesian Kriging approaches elicited twice the error, requiring more complex parameterization and significantly greater computational time than NN. These findings compare well with other studies that identified NN elevation void-filling interpolation as the most appropriate technique, balancing accuracy and shape reliability

560 across a range of natural environments (Bater and Coops, 2009; Boreggio et al., 2018). NN likely achieved the best results due to its simplicity and tendency to adapt to the structure of the elevation data by finding the closest subset of known elevation points to an unknown point and applying weights based on proportionate areas to interpolate a value. These results (Fig. 4) validate the choice of NN methodology implemented in Kokelj et al. (2021) and contribute a level of precision to the previously derived volumes whereby known uncertainties (Fig. 5c) can now be applied in estimations of material mobilization from hillslopes (Tunncliffe and Church, 2011) and in earth system models. Like Tseng et al. (2013), we determined error in absolute volume estimation to be less than 20% for small features and less than 10% for larger slope disturbances. The variability in volume estimates attributable to interpolation error decreases with increasing disturbance area and there is an inverse association between slump area and relative volumetric error (Fig. 5c, d). For example, the total modelled volume of the 71 active thaw-slump disturbances ($7.5 \times 10^{-6} \text{ m}^3$) from Kokelj et al. (2021) is associated with an interpolation-induced total volumetric uncertainty of $6.1 \times 10^{-5} \text{ m}^3$ or about 8% of estimated eroded volume. These volumetric uncertainties may be different depending on the accuracy and resolution of the source DEM, whereby the use of coarser-resolution DEMs (e.g., 10 m, 30 m) will likely be associated with higher relative volumetric uncertainties. In the following sections we evaluate whether NN is a suitable interpolation approach across a broad range of RTS-affected terrains including more complex disturbance morphologies and different activity levels.

4.2 Multisource Slump Inventory (MSI)

575 Retrogressive thaw slumps are dynamic mass-wasting features that develop over periods of years to decades to millennia as thaw-driven processes and feedbacks interact with topography, ground ice and substrate conditions, and climate (Lacelle et al., 2013; Kokelj et al., 2015). ~~Important patterns in the distribution of slump affected slopes and variation in the activity level of these disturbed areas may be obfuscated due to the difficulties with detection and delineation of active and stabilized slopes, limiting our understanding of the processes modifying the landscape and their spatial distribution. The development and implementation of the MSI have shown that retrogressive thaw slumps can be mapped and attributed consistently using a time-composite of high resolution hillshade DEMs supported by optical imagery (Figs. 2, 6). These procedures and a visual assessment of activity and field based knowledge of thaw slump process and form can be implemented at a regional scale to inventory disturbances, detect previously unmapped disturbances, and track change through time (Fig. 6c). Precise manual measurement of surface area and a consistent characterization scheme were important in developing a multi-temporal spatial dataset to explore area volume relationships. The MSI concept does not negate other inventory initiatives that compile the distribution of active slumps over large areas via manual or semi-automated boundary placement (e.g., Segal et al., 2016; Huang et al., 2020; Nitze et al., 2021; Swanson, 2021; Runge et al., 2022), rather it contributes to knowledge of differences in slump ontology and detectability, and provides novel data to consider how we define thaw slump disturbances, demarcates their boundaries, and explore their scar characteristics and allometric properties, all of which has become possible as improved data resolution allows for greater topographic detail to be captured and visualized (Guzzetti et al., 2012; Clare et al., 2019).~~

585 ~~The MSI dataset, summarized for the two study regions indicates that about half of the thaw slump affected slopes were stable (Figs. 7,8), the vast majority of recent thaw slump activity can be associated with areas of past disturbance, and that there is significant morphometric diversity in the active and stable landforms. The unique nature of the MSI is illustrated by general summaries of the dataset revealing that in the two well-studied areas).~~ Patterns in the distribution of slump-affected slopes and variation in the activity levels may be obfuscated due to the difficulties with detection and delineation of active and stabilized slopes, limiting our understanding of the processes modifying the landscape and their spatial distribution. The analysis of MSI

600 results demonstrate that retrogressive thaw slumps can be mapped and attributed consistently using a time-composite of high-resolution hillshade DEMs supported by optical imagery (Figs. 2, 6). These procedures and a visual assessment of activity and field-based knowledge of thaw slump process and form can be implemented at a regional scale to inventory disturbances, detect previously unmapped disturbances, and track change through time (Fig. 6c). Precise manual measurement of surface area and a consistent characterization scheme were important in developing a multi-temporal spatial dataset to explore area-volume relationships (Van der Sluijs and Kokelj, 2023). The MSI concept does not negate other manual or semi-automated inventory initiatives (e.g., Segal et al., 2016; Huang et al., 2020; Nitze et al., 2021; Swanson, 2021; Runge et al., 2022), rather it contributes to knowledge of differences in slump ontology and detectability, and provides a new approach on how to define and demarcate thaw slump disturbances. These methods and data enable us to explore scar characteristics and allometric properties, which has become possible as improved data resolution allows for greater topographic detail to be captured and visualized (Guzzetti et al., 2012; Clare et al., 2019).

610 The MSI dataset, summarized for the two study regions indicates that about half of the thaw slump affected slopes were stable (Figs. 7,8), the vast majority of recent thaw slump activity can be associated with areas of past disturbance, and that there is significant morphometric diversity in the active and stable landforms. The MSI dataset reveals that between 40% and 60% of slump-affected slopes detected by this inventory were inactive disturbances. The median area of stable thaw slump disturbance is lower than for the population with varying activity levels in both study areas (Fig. 7; Table 3). The data also show that total area of active thaw slumping has increased in APTC by 69% between 2004 and 2016, yet the total disturbance area has remained the same because the majority of activity has occurred within old or ancient scar areas that could be confidently identified by the MSI (Figs. 2c, 2d, 8). The increase in polycyclic activity is evacuating greater amounts of materials from within scar concavities, which has implications for the evolution of thawing slopes and area-volume relationships where thaw slump activity levels may account for scattering in area-volume models. There is growing evidence that contemporary slumps in PP and APTC are enlarging to surpass the extents of historical disturbances from which they originated (Kokelj et al., 2015, 2021). These stabilized disturbance areas typically have higher ground temperatures and greater active layer thicknesses relative to undisturbed terrain, and they are situated in areas of ice-rich permafrost that predispose the surface for future thaw-driven instability (Fig. 2, 8) (Kokelj et al., 2009). Our observations highlight how contemporary disturbances are tightly linked to past disturbance, which require inventory methods such as the MSI to detect and monitor these features. Additionally, the MSI observations may inspire additional satellite remote sensing avenues to detect these important landscape features over larger spatial extents.

625 The MSI These results indicate highlight the importance of considering distinctions distinguishing between process and form when mapping thaw-driven mass wasting features. The delineation of slump-affected areas accomplished in the past through examining ortho-mosaics, is less common now with the focus on thaw-driven change, and utilizing planform, coarser-resolution remote sensing data with a focus on covering broad spatial scales (Brooker et al., 2014; Lewkowicz and Way, 2019; Runge et al., 2022)(Brooker et al., 2014; Lewkowicz and Way, 2019; Runge et al., 2022). The delineation of areas affected by a continuum of active, stable and old thaw-slump scars can produce a robust regional account of “slump affected areas”, whereas inventories that map “active slumping” are implicitly documenting process rather than forms. Mapping focused on change differs from approaches typically implemented in landslide mapping, where most features have occurred as the result of episodic events (Guzzetti et al., 2012). The distinction between active process and form is particularly important to consider when inventorying, mapping, and monitoring chronic slope failures such as retrogressive thaw slumping, where progressive growth and polycyclic

are inherent due to the underlying terrain conditions, the nature of the process, and positive feedbacks associated with disturbed permafrost terrain (Kokelj et al., 2009, 2021).

640 The MSI demonstrates that most geomorphically-active disturbance areas are comprised of varying proportions of geomorphically active and stabilized terrain, emphasizing the complex nature of polycyclic retrogressive thaw slump activity-slumping. Utilizing high-resolution DEMs supported by optical imagery and methods implemented here also makes it possible to examine the relative levels of activity and the variation in morphometry of slump-affected terrain. For example, slumps in fluvially-incised terrain of the PP had higher elongation ratios reflecting compact or rounded morphology more
645 common to classic, cusate-shaped thaw slumps (Fig. 7c; Table 3), with ratios increasing with activity level. In the APTC both classic cusate features and elongated thaw slump disturbance occur, with the latter morphology typically characterizing eroding shorelines (Fig. 2b-e, 7b). The underlying DEMs required to generate the MSI data also enable topographic indices such as terrain roughness to be estimated. Descriptive statistics show that topographic roughness of slumps varied between the two study regions with increasing roughness in more highly active disturbances on PP. Together these analyses indicate variability in the
650 morphology of RTS-affected slopes, leading to challenges in thaw slump detection and determination of scaling behaviour.

4.3 Area-volume relationships for thaw slumps

The interpolation methods and the MSI database provided the foundation for area-volume models for slumps of varying activity levels, spanning 6 orders of magnitude in volume, and occurring across a wide range of geomorphic settings to be explored. The scaling coefficient for populations of inactive and active slumps obtained in this study ($\delta = 1.36 \pm 0.01$) is positioned near the lower end of deep-seated bedrock landslides ($\delta = 1.3-1.6$) and the higher end of soil landslides ($\delta = 1.1-1.4$), and exceeds submarine landslides ($\delta = 1.0-1.1$), regardless of differences in failed material, slope processes, sediment pressures or diagenesis (Chaytor et al., 2009; Larsen et al., 2010; Tseng et al., 2013; Jaboyedoff et al., 2020; and references therein). Due to the climate geomorphic feedbacks associated with slump development (Kokelj et al., 2015, 2021) the empirical evidence gathered in this study suggests that a heterogeneous disturbance area population consisting of ancient, recently active, and highly active slumps does not have a proportional erosion depth that is constant, and that scar volume has a non-linear relationship with the area (i.e., $\delta > \pm 1.0$). failure properties (Chaytor et al., 2009; Larsen et al., 2010). The empirical evidence gathered in this study confirms that a population of disturbance areas consisting of ancient, recently active, and highly active
660 slumps have a non-linear relationship with the volume (i.e., $\delta > \pm 1.0$), however due to the dynamics of slump evolution (Kokelj et al., 2015) the proportional erosion depth is heterogeneous. The scaling coefficient of inactive/old and active/modern slumps ($\delta = 1.36$, $n = 1,522$) observed in this study is lower than the $\delta = 1.42$ obtained by Kokelj et al. (2021) which was based on active slumps ($n = 71$) with a “classic” cusate or bowl-shaped form. These differences suggest that slump sub-populations grouped by terrain conditions, disturbance geometry, and activity levels may produce different scaling coefficients within and between geographic regions. Further study is required to investigate, warranting the influence investigation of increasingly active/increasing slump populations activity on disturbance morphometry and excavation volumes.
670

This study also indicates that variation in the scaling factor δ can be expected due to differences in slump inventory approaches and guiding ontologies. Recently, Bernhard et al., (2022) determined a $\delta = 1.17$ and $\delta = 1.27$ for the Tuktoyaktuk and Peel Plateau regions, respectively, based on differencing winter TanDEM-X DEMs and determining relationships between the “area showing an elevation decrease” and “the volume loss between measurement dates”. The differences in δ with our study bring attention to
675 the problem of scar definition and DEM resolution, as small differences in δ lead to substantial variation in volume predictions.

(Fig. 1). For example, ~~applying $\delta = 1.5$ instead of $\delta = 1.4$ produces a volume difference of a factor of two (Larsen et al., 2010; Tseng et al., 2013)~~ applying $\delta = 1.5$ instead of $\delta = 1.4$ produces a volume difference of a factor of two (Larsen et al., 2010; Tseng et al., 2013). Differences in δ may be attributed to data sources with coarser spatial resolution and greater vertical uncertainty, or due to the temporal window of data acquisition where differencing winter DEMs have a potential elevation bias due to incomplete radar penetration of snowdrifts or varying snow depths between years of observation. The δ reported by Bernhard et al. (2022) highlights a critical point of distinction between this study and Kokelj et al. (2021) as well as the larger landslide area-volume literature; (Jaboyedoff et al., 2020), as scaling based on the total area and fill volume of a scar zone (i.e., *fill volume scaling*) is different from their model which is based on annual changes in surface area regressed against annual volumetric change (i.e., *episodic volume scaling*). Both approaches are valid for assessing the impacts of thaw slumping as they provide different means of understanding the environmental implications of slump enlargement. Models based on episodic or annual change may provide an annual yield from the thawing headwalls of active slumps. ~~However during a periodic wave of activity.~~ In contrast, estimating erosion depth from the full scar area is aimed at measuring the time-integrated changes ~~in morphology of complex scar areas,~~ to yield models that elucidate the longer-term trajectory of the slump-affected landscape whereby concavities integrate erosion over decades to millennia. The method and ontological definition by which the process of slumping, and active and inactive disturbance areas are inventoried and represented volumetrically are therefore important considerations that require clear definition when scoping future studies.

Through mapping a broad range of slump-affected slopes, numerous volumetric outliers were detected, where the modelling of pre-disturbance terrain (Fig. 5) did not fit the highly elongated and convex slump topographies (e.g., Fig. 6b). These outliers are related to uncertainty in defining the pre-disturbed slope configuration, especially for coastal slumps or lacustrine slumps which typically exhibit narrow, elongated extents due to long-term shoreline retreat and complex interactions between simultaneous headwall retreat and wave-form erosion (Obu et al., 2016; Clark et al., 2021). (Obu et al., 2016; Clark et al., 2021). Shoreline slumps along lakes typically initiate due to lateral talik expansion, thaw of ice-rich permafrost ~~sub-adjacent~~ subadjacent to the lakeshore, and lake-bottom subsidence (Kokelj et al., 2009) whereas in coastal settings rapid erosion of ice-rich shorelines driven by wave action produces dynamic conditions that extend beyond those associated with thaw slump development alone (Lantuit et al., 2012; Ramage et al., 2017; Leibman et al., 2021; Berry et al., 2021)(Lantuit et al., 2012; Ramage et al., 2017; Leibman et al., 2021; Berry et al., 2021). Lake-bottom subsidence is effectively 'invisible', skewing volumetric estimates of yield. Attempts at inferring and repositioning ancient lake edges or former coast edges (e.g., Ramage et al., 2017) were not made in this study, nor was bathymetric data available to improve digitizations, hence estimates of many lacustrine/coastal slumps volumes were underestimated to the point of not meeting the volumetric uncertainty threshold. The relationship between planimetric and volumetric erosion measurements for highly elongated slumps and coastal erosion sections can be complex (Obu et al., 2016)(Obu et al., 2016), where improved prediction for shoreline slumps likely requires data on sub-surface conditions and knowledge of topographic profiles of undisturbed lake edges to parameterize interpolation algorithms. The challenges encountered with pre-disturbance terrain modelling are in part related to slump ontology, as well as a dichotomy between event-based mass-wasting and chronic denudation of shoreline slumps where ice-rich topography is prone to progressive failure over long time-periods, under seasonal climate forcing.

5 Conclusion

715 Retrogressive thaw slumps are multi-year, chronic disturbances that are increasingly important modifiers of ice-rich permafrost terrain. Quantifying geomorphic characteristics of thaw-driven landslides is required to understand their influence on landscape evolution, downstream sedimentary and geochemical effects, and the release or sequestration of organic carbon. In this research, our goal was to couple knowledge of thaw slump processes and form with remote-sensing tools to advance holistic approaches to monitoring and quantifying the geomorphic effects of retrogressive thaw slumps. We evaluated surface interpolation techniques to derive slump volumes based on DEM differencing (Objective A) and we developed a high-resolution DEM-based slump inventory method to track surface area and activity levels (Objective B). These methods were integrated to explore area-volume relationships for retrogressive thaw slump-affected slopes (Objective C). In summary:

- 720 1. This study improved tools to estimate slump volume and calculate uncertainty arising from pre-disturbance terrain reconstructions. Natural Neighbour interpolation achieved the best precision for modelled pre-disturbance topography. Error estimates in slump volume were <10% for large disturbances and less than 10-20% for small to medium slumps.
 - 725 2. A robust consistent method (MSI) for delineating slump-affected slopes and determining activity levels using high-resolution DEMs and optical imagery was developed. Significant variation in the morphology of slumps ~~described in this dataset~~ occurs in association with terrain type, geomorphic setting, and activity level. The MSI inventory ~~also documents~~ widespread evidence of historical disturbances, most of which are currently inactive. ~~The, and that the~~ majority of active slumping is associated with areas of past disturbance indicating that ~~assessments of~~ stable thaw slumps ~~provide a useful~~ are an important indicator of sensitive terrain. This study corroborated past results indicating a significant (69%) increase in active slumping between 2004-2016.
 - 730 3. The datasets generated in the first two sections of the paper enabled area-volume relationships of RTS-affected slopes to be explored. The resulting power-law model (adj-R² of 0.85, n=1,522) enables a robust assessment of thaw-driven landslide impacts for regions where disturbance area is determined. Analyses of outliers in the A/V relationship were elongated coastal slumps, providing constraints on model application.
- 735 Future directions of research should involve revisiting thaw-driven landslide ontology to support scoping and informed interpretation of remote sensing outputs, exploring factors driving variation in slump morphometry and activity level, and determining whether area-volume relations vary between regions, among other thaw-driven disturbance types, and between volumetric representations of slumps.

Data availability

740 ~~Datasets used in this publication and sources are summarized in Table 1. Framework and geoprocessing steps for developing the Multisource Slump Inventory and pre-disturbance terrain methods are available in the Supplement.~~
Datasets used in this publication and sources are summarized in Table 1. Slump delineations of the MSI are accessible via Van der Sluijs and Kokelj (Van der Sluijs and Kokelj, 2023). Flowchart and geoprocessing steps for developing the input DEMs and the pre-disturbance terrain methods are available in the Supplement. Python code to derive slump volumes is available in S4 (DEM PredisturbanceGeneration AreaVolumeCalc Batch NN Py3.py). Derived slump volumes are available in S5 (MSI 2022 10 Volumes Outlierclass.shp).

745

Supplement

The supplement related to this article is available online at: URL.

Author contributions

750 All authors developed the paper concept. JV and SVK contributed to field data collection and formulation of the Multisource Slump Inventory concept and implementation. JV processed the geospatial datasets, and developed and interpreted statistical analyses with guidance from SVK and JT. JV drafted and revised the paper and figures with input from all authors.

Competing interests

The authors declare that they have no conflict of interest.

755 Disclaimer

No disclaimer.

Acknowledgments

This work is part of a long-term permafrost monitoring and research program within the Government of Northwest Territories (GNWT) - NWT Geological Survey with partners including the NWT Centre for Geomatics. We are grateful to the Indigenous
760 peoples of the Gwich'in Settlement Area and Inuvialuit Settlement Region of the Northwest Territories for the opportunity to work collaboratively and to learn and gather knowledge on their lands. Long-term support from the Tetl'it and Ehdiiat Renewable Resource Councils, the Inuvik, Tuktoyaktuk Hunters and Trappers Committees, the Inuvialuit Joint Secretariat, the Inuvialuit Land Administration, [the Inuvialuit Regional Corporation \(Charles Klengenberg\)](#) and the Aurora Research Institute
765 [\(Joel McAlister\)](#) is gratefully acknowledged. Field support from Christine Firth, Eugene Pascale, Steven Tetlitchi, Alice Wilson, [Ashley Rudy](#), and Billy Wilson are also sincerely acknowledged. William Woodley (NWT Centre for Geomatics) processed MVAP contour lines into the 3-m DEM.

Financial support

The work was supported by the Department of Environment and Natural Resources Climate Change and Northwest Territories Cumulative Impact Monitoring Program of the GNWT (grant nos. 164 and 186, Steven V. Kokelj), the Natural Science and
770 Engineering Research Council of Canada, and the Polar Continental Shelf Program, Natural Resources Canada (projects 313-18, 316-19, 318-20, and 320-20 to Steven V. Kokelj).

References

Aylsworth, J. M., Rodkin-Duk, A., Robertson, T., and Traynor, J. A.: Landslide of the Mackenzie valley and adjacent mountainous and coastal regions, in: The Physical Environment of the Mackenzie Valley: A Baseline for the Assessment of
775 Environmental Change, L.D. Dyke and G.R. Brooks (eds.), 167–176, 2000.

- Bater, C. W. and Coops, N. C.: Evaluating error associated with lidar-derived DEM interpolation, *Computers & Geosciences*, 35, 289–300, <https://doi.org/10.1016/j.cageo.2008.09.001>, 2009.
- Bergonse, R. and Reis, E.: Reconstructing pre-erosion topography using spatial interpolation techniques: A validation-based approach, *J. Geogr. Sci.*, 25, 196–210, <https://doi.org/10.1007/s11442-015-1162-2>, 2015.
- 780 Bernhard, P., Zwieback, S., Leinss, S., and Hajnsek, I.: Mapping Retrogressive Thaw Slumps Using Single-Pass TanDEM-X Observations, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 13, 3263–3280, <https://doi.org/10.1109/JSTARS.2020.3000648>, 2020.
- Bernhard, P., Zwieback, S., Bergner, N., and Hajnsek, I.: Assessing volumetric change distributions and scaling relations of retrogressive thaw slumps across the Arctic, *The Cryosphere*, 16, 1–15, <https://doi.org/10.5194/tc-16-1-2022>, 2022.
- 785 Berry, H. B., Whalen, D., and Lim, M.: Long-term ice-rich permafrost coast sensitivity to air temperatures and storm influence: lessons from Pullen Island, Northwest Territories, Canada, *Arctic Science*, 7, 723–745, <https://doi.org/10.1139/as-2020-0003>, 2021.
- Boreggio, M., Bernard, M., and Gregoretti, C.: Evaluating the Differences of Gridding Techniques for Digital Elevation Models Generation and Their Influence on the Modeling of Stony Debris Flows Routing: A Case Study From Rovina di Cancia Basin (North-Eastern Italian Alps), *Front. Earth Sci.*, 6, <https://doi.org/10.3389/feart.2018.00089>, 2018.
- 790 Brardinoni, F., Slaymaker, O., and Hassan, M. A.: Landslide inventory in a rugged forested watershed: a comparison between air-photo and field survey data, *Geomorphology*, 54, 179–196, [https://doi.org/10.1016/S0169-555X\(02\)00355-0](https://doi.org/10.1016/S0169-555X(02)00355-0), 2003.
- Breiman, L.: Random Forests, *Machine Learning*, 45, 5–32, <https://doi.org/10.1023/a:1010933404324>, 2001.
- ten Brink, U. S., Geist, E. L., and Andrews, B. D.: Size distribution of submarine landslides and its implication to tsunami hazard in Puerto Rico, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2006GL026125>, 2006.
- 795 Brooker, A., Fraser, R. H., Olthof, I., Kokelj, S. V., and Lacelle, D.: Mapping the activity and evolution of retrogressive thaw slumps by tasselled cap trend analysis of a Landsat satellite image stack, *Permafrost and periglacial processes*, 25, 243–256, <https://doi.org/doi:10.1002/ppp.1819>, 2014.
- Burn, C. R.: Cryostratigraphy, paleogeography, and climate change during the early Holocene warm interval, western Arctic coast, Canada, *Can. J. Earth Sci.*, 34, 912–925, <https://doi.org/10.1139/e17-076>, 1997.
- 800 Burn, C. R.: The thermal regime of a retrogressive thaw slump near Mayo, Yukon Territory, *Can. J. Earth Sci.*, 37, 967–981, <https://doi.org/10.1139/e00-017>, 2000.
- Burn, C. R. and Friele, P. A.: Geomorphology, Vegetation Succession, Soil Characteristics and Permafrost in Retrogressive Thaw Slumps near Mayo, Yukon Territory, *Arctic*, 42, 31–40, 1989.
- 805 Burn, C. R. and Kokelj, S. V.: The environment and permafrost of the Mackenzie Delta area, *Permafrost and periglacial processes*, 20, 83–105, <https://doi.org/doi:10.1002/ppp.655>, 2009.
- Burn, C. R. and Lewkowicz, A. G.: Canadian landform examples - 17 retrogressive thaw slumps, *The Canadian Geographer / Le Géographe canadien*, 34, 273–276, <https://doi.org/doi:10.1111/j.1541-0064.1990.tb01092.x>, 1990.
- 810 Campbell, D. and Church, M.: Reconnaissance sediment budgets for Lynn Valley, British Columbia: Holocene and contemporary time scales, *Can. J. Earth Sci.*, 40, 701–713, <https://doi.org/10.1139/e03-012>, 2003.
- Chaytor, J. D., ten Brink, U. S., Solow, A. R., and Andrews, B. D.: Size distribution of submarine landslides along the U.S. Atlantic margin, *Marine Geology*, 264, 16–27, <https://doi.org/10.1016/j.margeo.2008.08.007>, 2009.
- Clare, M., Chaytor, J., Dabson, O., Gamboa, D., Georgiopolou, A., Eady, H., Hunt, J., Jackson, C., Katz, O., Krastel, S., León, R., Micallef, A., Moernaut, J., Moriconi, R., Moscardelli, L., Mueller, C., Normandeau, A., Patacci, M., Steventon, M., Urlaub, M., Völker, D., Wood, L., and Jobe, Z.: A consistent global approach for the morphometric characterization of subaqueous landslides, *Geological Society, London, Special Publications*, 477, 455–477, <https://doi.org/10.1144/SP477.15>, 2019.

- Clark, A., Moorman, B., Whalen, D., and Fraser, P.: Arctic coastal erosion: UAV-SfM data collection strategies for planimetric and volumetric measurements, *Arctic Science*, 7, 605–633, <https://doi.org/10.1139/as-2020-0021>, 2021.
- 820 Côté, M. M., Duchesne, C., Wright, J., and Ednie, M.: Digital compilation of the surficial sediments of the Mackenzie Valley corridor, Yukon Coastal Plain, and the Tuktoyaktuk Peninsula, Natural Resources Canada, Geological Survey of Canada, 2013.
- Fraser, R. H., Olthof, I., Kokelj, S. V., Lantz, T. C., Lacelle, D., Brooker, A., Wolfe, S., and Schwarz, S.: Detecting landscape changes in high latitude environments using Landsat trend analysis: 1. visualization, *Remote Sensing*, 6, 11533–11557, <https://doi.org/10.3390/rs6111533>, 2014.
- 825 Gudowicz, J. and Paluszkiwicz, R.: MAT: GIS-Based Morphometry Assessment Tools for Concave Landforms, *Remote Sensing*, 13, <https://doi.org/10.3390/rs13142810>, 2021.
- Guzzetti, F., Mondini, A. C., Cardinali, M., Fiorucci, F., Santangelo, M., and Chang, K.-T.: Landslide inventory maps: New tools for an old problem, *Earth-Science Reviews*, 112, 42–66, <https://doi.org/10.1016/j.earscirev.2012.02.001>, 2012.
- Huang, L., Luo, J., Lin, Z., Niu, F., and Liu, L.: Using deep learning to map retrogressive thaw slumps in the Beiluhe region (Tibetan Plateau) from CubeSat images, *Remote Sensing of Environment*, 237, 111534, <https://doi.org/10.1016/j.rse.2019.111534>, 2020.
- 830 Huang, L., Lantz, T. C., Fraser, R. H., Tiampo, K. F., Willis, M. J., and Schaefer, K.: Accuracy, Efficiency, and Transferability of a Deep Learning Model for Mapping Retrogressive Thaw Slumps across the Canadian Arctic, *Remote Sensing*, 14, <https://doi.org/10.3390/rs14122747>, 2022.
- Jaboyedoff, M., Carrea, D., Derron, M.-H., Oppikofer, T., Penna, I. M., and Rudaz, B.: A review of methods used to estimate initial landslide failure surface depths and volumes, *Engineering Geology*, 267, 105478, <https://doi.org/10.1016/j.enggeo.2020.105478>, 2020.
- 835 ~~Jones, M. K. W. and Pollard, W. H.: Daily Field Observations of Retrogressive Thaw Slump Dynamics in the Canadian High Arctic, *ARCTIC*, 74, 339–354, <https://doi.org/10.14430/arctic73377>, 2021.~~
- 840 ~~Jones, M. K. W., Pollard, W. H., and Jones, B. M.: Rapid initialization of retrogressive thaw slumps in the Canadian high Arctic and their response to climate and terrain factors, *Environ. Res. Lett.*, 14, 055006, <https://doi.org/10.1088/1748-9326/ab12fd>, 2019.~~
- Klar, A., Aharonov, E., Kalderon-Asael, B., and Katz, O.: Analytical and observational relations between landslide volume and surface area, *Journal of Geophysical Research: Earth Surface*, 116, <https://doi.org/10.1029/2009JF001604>, 2011.
- 845 ~~Kokelj, S. V. and Jorgenson, M. T.: Advances in Thermokarst Research, *Permafrost and periglacial processes*, 24, 108–119, <https://doi.org/doi:10.1002/ppp.1779>, 2013.~~
- Kokelj, S. V., Jenkins, R. E., Milburn, D., Burn, C. R., and Snow, N.: The influence of thermokarst disturbance on the water quality of small upland lakes, Mackenzie Delta region, Northwest Territories, Canada, *Permafrost and periglacial processes*, 16, 343–353, <https://doi.org/doi:10.1002/ppp.536>, 2005.
- 850 Kokelj, S. V., Lantz, T. C., Kanigan, J., Smith, S. L., and Coutts, R.: Origin and polycyclic behaviour of tundra thaw slumps, Mackenzie Delta region, Northwest Territories, Canada, *Permafrost and periglacial processes*, 20, 173–184, <https://doi.org/doi:10.1002/ppp.642>, 2009.
- Kokelj, S. V., Lacelle, D., Lantz, T. C., Tunnicliffe, J., Malone, L., Clark, I. D., and Chin, K. S.: Thawing of massive ground ice in mega slumps drives increases in stream sediment and solute flux across a range of watershed scales, *Journal of Geophysical Research: Earth Surface*, 118, 681–692, <https://doi.org/doi:10.1002/jgrf.20063>, 2013.
- 855 Kokelj, S. V., Lantz, T. C., Wolfe, S. A., Kanigan, J. C., Morse, P. D., Coutts, R., Molina-Giraldo, N., and Burn, C. R.: Distribution and activity of ice wedges across the forest-tundra transition, western Arctic Canada, *Journal of Geophysical Research: Earth Surface*, 119, 2032–2047, <https://doi.org/doi:10.1002/2014JF003085>, 2014.

- 860 Kokelj, S. V., Tunnicliffe, J., Lacelle, D., Lantz, T. C., Chin, K. S., and Fraser, R.: Increased precipitation drives mega slump development and destabilization of ice-rich permafrost terrain, northwestern Canada, *Global and Planetary Change*, 129, 56–68, <https://doi.org/10.1016/j.gloplacha.2015.02.008>, 2015.
- Kokelj, S. V., Lantz, T. C., Tunnicliffe, J., Segal, R., and Lacelle, D.: Climate-driven thaw of permafrost preserved glacial landscapes, northwestern Canada, *Geology*, 45, 371–374, <https://doi.org/10.1130/g38626.1>, 2017a.
- 865 Kokelj, S. V., Tunnicliffe, J. F., and Lacelle, D.: The Peel Plateau of northwestern Canada: an ice-rich hummocky moraine landscape in transition, in: *Landscapes and Landforms of Western Canada*, edited by: Slaymaker, O., Springer International Publishing, Cham, Switzerland, 109–122, 2017b.
- Kokelj, S. V., Kokoszka, J., van der Sluijs, J., Rudy, A. C. A., Tunnicliffe, J., Shakil, S., Tank, S. E., and Zolkos, S.: Thaw-driven mass wasting couples slopes with downstream systems, and effects propagate through Arctic drainage networks, *The Cryosphere*, 15, 3059–3081, <https://doi.org/10.5194/tc-15-3059-2021>, 2021.
- 870 Lacelle, D., Bjornson, J., and Lauriol, B.: Climatic and geomorphic factors affecting contemporary (1950–2004) activity of retrogressive thaw slumps on the Aklavik Plateau, Richardson Mountains, NWT, Canada, *Permafrost and Periglacial Processes*, 21, 1–15, <https://doi.org/10.1002/ppp.666>, 2010.
- Lacelle, D., Brooker, A., Fraser, R. H., and Kokelj, S. V.: Distribution and growth of thaw slumps in the Richardson Mountains–Peel Plateau region, northwestern Canada, *Geomorphology*, 235, 40–51, <https://doi.org/10.1016/j.geomorph.2015.01.024>, 2015.
- 875 Lacelle, D., Fontaine, M., Pellerin, A., Kokelj, S. V., and Clark, I. D.: Legacy of Holocene Landscape Changes on Soil Biogeochemistry: A Perspective From Paleo-Active Layers in Northwestern Canada, *Journal of Geophysical Research: Biogeosciences*, 124, 2662–2679, <https://doi.org/10.1029/2018JG004916>, 2019.
- Lantuit, H. and Pollard, W. H.: Fifty years of coastal erosion and retrogressive thaw slump activity on Herschel Island, southern Beaufort Sea, Yukon Territory, Canada, *Geomorphology*, 95, 84–102, <https://doi.org/10.1016/j.geomorph.2006.07.040>, 2008.
- 880 Lantuit, H., Pollard, W. H., Couture, N., Fritz, M., Schirmer, L., Meyer, H., and Hubberten, H.-W.: Modern and Late Holocene Retrogressive Thaw Slump Activity on the Yukon Coastal Plain and Herschel Island, Yukon Territory, Canada, *Permafrost and Periglacial Processes*, 23, 39–51, <https://doi.org/10.1002/ppp.1731>, 2012.
- Lantz, T. C. and Kokelj, S. V.: Increasing rates of retrogressive thaw slump activity in the Mackenzie Delta region, N.W.T., Canada, *Geophysical Research Letters*, 35, <https://doi.org/10.1029/2007GL032433>, 2008.
- 885 Lantz, T. C., Kokelj, S. V., Gergel, S. E., and Henry, G. H. R.: Relative impacts of disturbance and temperature: persistent changes in microenvironment and vegetation in retrogressive thaw slumps, *Global Change Biology*, 15, 1664–1675, <https://doi.org/10.1111/j.1365-2486.2009.01917.x>, 2009.
- Larsen, I. J., Montgomery, D. R., and Korup, O.: Landslide erosion controlled by hillslope material, *Nature Geosci*, 3, 247–251, <https://doi.org/10.1038/ngeo776>, 2010.
- 890 Leibman, M., Kizyakov, A., Zhdanova, Y., Sonyushkin, A., and Zimin, M.: Coastal Retreat Due to Thermodenudation on the Yugorsky Peninsula, Russia during the Last Decade, Update since 2001–2010, *Remote Sensing*, 13, 4042, <https://doi.org/10.3390/rs13204042>, 2021.
- Lewkowicz, A. G.: Headwall retreat of ground-ice slumps, Banks Island, Northwest Territories, *Canadian Journal of Earth Sciences*, 24, 1077–1085, <https://doi.org/10.1139/e87-105>, 1987.
- 895 Lewkowicz, A. G. and Way, R. G.: Extremes of summer climate trigger thousands of thermokarst landslides in a High Arctic environment, *Nat Commun*, 10, 1329, <https://doi.org/10.1038/s41467-019-09314-7>, 2019.
- Luo, J., Niu, F., Lin, Z., Liu, M., and Yin, G.: Recent acceleration of thaw slumping in permafrost terrain of Qinghai-Tibet Plateau: An example from the Beiluhe Region, *Geomorphology*, 341, 79–85, <https://doi.org/10.1016/j.geomorph.2019.05.020>, 2019.
- 900 Mackay, J. R.: Segregated epigenetic ice and slumps in permafrost Mackenzie Delta area, N.W.T., *Geographical Bulletin*, 8, 59–80, 1966.

- Mackay, J. R.: Some observations on the growth and deformation of epigenetic, syngenetic and anti-syngenetic ice wedges, *Permafrost Periglacial Processes*, 1, 15–29, <https://doi.org/doi:10.1002/ppp.3430010104>, 1990.
- Murton, J. B.: Ground-ice stratigraphy and formation at North Head, Tuktoyaktuk Coastlands, western Arctic Canada: a product of glacier–permafrost interactions, *Permafrost and Periglacial Processes*, 16, 31–50, <https://doi.org/10.1002/ppp.513>, 2005.
- 905 Nitze, I., Heidler, K., Barth, S., and Grosse, G.: Developing and Testing a Deep Learning Approach for Mapping Retrogressive Thaw Slumps, *Remote Sensing*, 13, 4294, <https://doi.org/10.3390/rs13214294>, 2021.
- Obu, J., Lantuit, H., Fritz, M., Pollard, W. H., Sachs, T., and Günther, F.: Relation between planimetric and volumetric measurements of permafrost coast erosion: a case study from Herschel Island, western Canadian Arctic, *Polar Research*, <https://doi.org/10.3402/polar.v35.30313>, 2016.
- 910 Oguchi, T.: Drainage Density and Relative Relief in Humid Steep Mountains with Frequent Slope Failure, *Earth Surface Processes and Landforms*, 22, 107–120, [https://doi.org/10.1002/\(SICI\)1096-9837\(199702\)22:2<107::AID-ESP680>3.0.CO;2-U](https://doi.org/10.1002/(SICI)1096-9837(199702)22:2<107::AID-ESP680>3.0.CO;2-U), 1997.
- Porter, C., Morin, P., Howat, I., Noh, M.-J., Bates, B., Peterman, K., Keeseey, S., Schlenk, M., Gardiner, J., Tomko, K., Willis, M., Kelleher, C., Cloutier, M., Husby, E., Foga, S., Nakamura, H., Platson, M., Wethington, M., Williamson, C., Bauer, G., Enos, J., Arnold, G., Kramer, W., Becker, P., Doshi, A., D’Souza, C., Cummens, P., Laurier, F., and Bojesen, M.: ArcticDEM, 2018.
- 915 Ramage, J. L., Irrgang, A. M., Herzsuh, U., Morgenstern, A., Couture, N., and Lantuit, H.: Terrain controls on the occurrence of coastal retrogressive thaw slumps along the Yukon Coast, Canada, *Journal of Geophysical Research: Earth Surface*, 122, 1619–1634, <https://doi.org/doi:10.1002/2017JF004231>, 2017.
- 920 Rampton, V. N.: Quaternary Geology of the Tuktoyaktuk Coastlands, Northwest Territories., Geological Survey of Canada. Memoir 423, Ottawa, ON, Canada, 1988.
- Reuter, H. I., Nelson, A., and Jarvis, A.: An evaluation of void-filling interpolation methods for SRTM data, *International Journal of Geographical Information Science*, 21, 983–1008, <https://doi.org/10.1080/13658810601169899>, 2007.
- Riley, S., Degloria, S., and Elliot, S. D.: A Terrain Ruggedness Index that Quantifies Topographic Heterogeneity, *International Journal of Science*, 5, 23–27, 1999.
- 925 ~~Rudy, A. C. A., Lamoureux, S. F., Kokelj, S. V., Smith, I. R., and England, J. H.: Accelerating thermokarst transforms ice-cored terrain triggering a downstream cascade to the ocean, *Geophysical Research Letters*, 44, 11080–11087, <https://doi.org/doi:10.1002/2017GL074912>, 2017.~~
- Runge, A., Nitze, I., and Grosse, G.: Remote sensing annual dynamics of rapid permafrost thaw disturbances with LandTrendr, *Remote Sensing of Environment*, 268, 112752, <https://doi.org/10.1016/j.rse.2021.112752>, 2022.
- 930 Segal, R. A., Lantz, T. C., and Kokelj, S. V.: Acceleration of thaw slump activity in glaciated landscapes of the Western Canadian Arctic, *Environmental Research Letters*, 11, 034025, 2016.
- Sladen, W. E., Parker, R. J. H., Kokelj, S. V., and Morse, P. D.: Geomorphologic feature mapping methodology developed for the Dempster Highway and Inuvik to Tuktoyaktuk Highway corridors, 2021.
- 935 Smith, I. R. and Duong, L.: An assessment of surficial geology, massive ice, and ground ice, Tuktoyaktuk Peninsula, Northwest Territories: application to the proposed Inuvik to Tuktoyaktuk all-weather highway, Natural Resources Canada, Geological Survey of Canada, 2012.
- Stark, C. P. and Hovius, N.: The characterization of landslide size distributions, *Geophysical Research Letters*, 28, 1091–1094, <https://doi.org/10.1029/2000GL008527>, 2001.
- 940 Swanson, D. and Nolan, M.: Growth of retrogressive thaw slumps in the Noatak Valley, Alaska, 2010–2016, measured by airborne photogrammetry, *Remote Sensing*, 10, 983, 2018.

- Swanson, D. K.: Permafrost thaw-related slope failures in Alaska's Arctic National Parks, c. 1980–2019, *Permafrost and Periglacial Processes*, 32, 392–406, <https://doi.org/10.1002/ppp.2098>, 2021.
- 945 Treharne, R., Rogers, B. M., Gasser, T., MacDonald, E., and Natali, S.: Identifying Barriers to Estimating Carbon Release From Interacting Feedbacks in a Warming Arctic, *Frontiers in Climate*, 3, 2022.
- Tseng, C.-M., Lin, C.-W., Stark, C. P., Liu, J.-K., Fei, L.-Y., and Hsieh, Y.-C.: Application of a multi-temporal, LiDAR-derived, digital terrain model in a landslide-volume estimation, *Earth Surface Processes and Landforms*, 38, 1587–1601, <https://doi.org/10.1002/esp.3454>, 2013.
- 950 Tunnicliffe, J. F. and Church, M.: Scale variation of post-glacial sediment yield in Chilliwack Valley, British Columbia, *Earth Surface Processes and Landforms*, 36, 229–243, <https://doi.org/10.1002/esp.2093>, 2011.
- Turner, K. W., Pearce, M. D., and Hughes, D. D.: Detailed Characterization and Monitoring of a Retrogressive Thaw Slump from Remotely Piloted Aircraft Systems and Identifying Associated Influence on Carbon and Nitrogen Export, *Remote Sensing*, 13, 171, <https://doi.org/10.3390/rs13020171>, 2021.
- 955 ~~Van der Sluijs, J., Kokelj, S. V., Fraser, R. H., Tunnicliffe, J., and Lacelle, D.: Permafrost Terrain Dynamics and Infrastructure Impacts Revealed by UAV Photogrammetry and Thermal Imaging, *Remote Sensing*, 10, 1734, 2018.~~ and ~~Kokelj, S. V.: High-resolution inventory of retrogressive thaw slump affected slopes using high spatial resolution Digital Elevation models and imagery, *Peel Plateau and Anderson Plain – Tuktoyaktuk Coastlands, Northwest Territories, Northwest Territories Geological Survey*, <https://doi.org/10.46887/2023-013>, 2023.~~
- 960 ~~[Van der Sluijs, J., Kokelj, S. V., Fraser, R. H., Tunnicliffe, J., and Lacelle, D.: Permafrost Terrain Dynamics and Infrastructure Impacts Revealed by UAV Photogrammetry and Thermal Imaging, *Remote Sensing*, 10, 1734, 2018.](https://doi.org/10.46887/2023-013)~~
- ~~[Ward Jones, M. K. and Pollard, W. H.: Daily Field Observations of Retrogressive Thaw Slump Dynamics in the Canadian High Arctic, *ARCTIC*, 74, 339–354, <https://doi.org/10.14430/arctic73377>, 2021.](https://doi.org/10.14430/arctic73377)~~
- 965 ~~[Ward Jones, M. K., Pollard, W. H., and Jones, B. M.: Rapid initialization of retrogressive thaw slumps in the Canadian high Arctic and their response to climate and terrain factors, *Environ. Res. Lett.*, 14, 055006, <https://doi.org/10.1088/1748-9326/ab12fd>, 2019.](https://doi.org/10.1088/1748-9326/ab12fd)~~
- Xia, Z., Huang, L., Fan, C., Jia, S., Lin, Z., Liu, L., Luo, J., Niu, F., and Zhang, T.: Retrogressive thaw slumps along the Qinghai-Tibet Engineering Corridor: a comprehensive inventory and their distribution characteristics, *Earth Syst. Sci. Data Discuss.*, 2022, 1–19, <https://doi.org/10.5194/essd-2021-439>, 2022.
- 970 Zwieback, S., Kokelj, S. V., Günther, F., Boike, J., Grosse, G., and Hajnsek, I.: Sub-seasonal thaw slump mass wasting is not consistently energy limited at the landscape scale, *The Cryosphere*, 12, 549–564, <https://doi.org/10.5194/tc-12-549-2018>, 2018.