

# Estimation of streamwater components and water residence time in a permafrost catchment in the Central Tibetan Plateau using long-term water stable isotopic data

Shaoyong Wang<sup>1,2#</sup>, Xiaobo He<sup>1#</sup>, Shichang Kang<sup>1,2</sup>, Hui Fu<sup>3</sup>, Xiaofeng Hong<sup>4</sup>

5 <sup>1</sup>State Key Laboratory of Cryospheric Science, Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences, Lanzhou 730000, China

<sup>2</sup>College of Resources and Environment, University of Chinese Academy of Sciences, Beijing 100049, China

<sup>3</sup>China Institute of Water Resources and Hydropower Research, Beijing 100049, China

<sup>4</sup>Water Resources Department, Yangtze River Scientific Research Institute, Wuhan 430010, China

10 ~~#These authors contribute equally to this work~~

**Correspondence to:** Xiaobo He (hxb@lzb.ac.cn)

**Abstract.** Global warming has ~~significantly~~greatly impacted the hydrological processes and ecological environment in permafrost regions. Mean residence time (MRT) is a fundamental catchment descriptor that ~~provides~~reveals hydrological information ~~regarding about~~ storage, flow pathways, and water source within a particular catchment. However, water stable  
15 isotopes and MRT have ~~rarely~~scarcely been investigated due to limited data collection in the high-altitude permafrost regions. This study ~~used~~uses the long-term stable isotopic observations to identify runoff components and applied the sine-wave exponential model to estimate water MRT in a high-altitude permafrost catchment (5,300 m a.s.l.) in the central Tibetan Plateau (TP). We found that the isotope composition in precipitation, stream, and supra-permafrost water exhibited obvious seasonal variability. Freeze-thaw ~~process~~cycles of permafrost active layer and direct input of precipitation significantly modified the  
20 stable isotope compositions in supra-permafrost and stream water. The hydrograph separation revealed that precipitation and supra-permafrost water accounted for ~~6235 ± 13.2~~ % and ~~3865 ± 13.2~~ % of the total discharge of stream water, respectively. ~~We estimated that~~ MRT for stream and supra-permafrost water was estimated at 100 and 255 days, respectively. Such shorter MRT of supra-permafrost and stream water (compared to the non-permafrost catchments) might reflect the unique characteristics of hydrological process in permafrost catchments. Moreover, the MRT of supra-permafrost water was ~~found to~~  
25 ~~be~~ more sensitive to environmental change than that of stream water. Climate and vegetation factors affected the MRT of stream and supra-permafrost water mainly by changing the thickness of permafrost active layer. Our results suggest ~~We conclude~~ that ~~global climate~~ warming might retard the rate of water cycle in permafrost regions. Overall, our study ~~expands~~deepened the our understanding of hydrological processes in high-altitude permafrost catchments under global warming, and ~~provided a decision making basis for ecological environmental protection and water resources safety in the~~  
30 ~~source of rivers on the TP.~~

## 1 Introduction

Permafrost, ~~considered is counted~~ among the regions that are most sensitive to global warming (Wang et al., 2009). ~~The permafrost,~~ acts as an aquiclude, which governs the surface runoff and its hydraulic connection with groundwater ~~and alters groundwater flow paths due to its thawing (Woo, 1990; Hinzman et al., 2005; Sjöberg et al., 2021) (Gao et al., 2021)~~. The

35 spatiotemporal variability of permafrost freeze-thaw cycles alongside the development of an active layer influence the catchment hydrology in a time-dependent manner, ~~thereby~~ affecting parameters such as soil water movement direction, storage capacity, and hydraulic conductivity (Tetzlaff et al., 2018; Gao et al., 2021). ~~In-Over~~ the past decades, permafrost has experienced significant, rapid and extensive degradation, which profoundly and extensively affected regional and even continental hydrological regimes, as well as alpine ecology (Cheng et al., 2019; Jin et al., 2021) ~~(Gao et al., 2021)~~.

40 ~~Permafrost~~ This degradation has released valuable meltwater for recharging groundwater (Ma et al., 2019a). ~~This degradation, accompanied by progressive increase of the active layer, could higher water content in the soil column, thereby increasing the groundwater storage capacity (Woo, 1990; Hinzman et al., 2005; Woo et al., 2008). The progressive increase of the permafrost active layer thickness has exacerbated the increased water storage capacity of permafrost and exerted a higher contribution of groundwater to river water (Streletskiy et al., 2015)~~. Thus, compared to non-permafrost regions, the presence of an aquiclude

45 and freeze-thaw cycle of the permafrost active layer, contributes to more complex hydrological processes in permafrost regions. ~~However, it remains unclear how permafrost changes would alter water storage and movement in permafrost catchments.~~

~~The mechanisms of runoff processes in permafrost region (e.g., flow pathways, water source, and residence time) have attracted attention (Tetzlaff et al., 2018; Carey and Quinton, 2005; Li et al., 2020a; Yang et al., 2016). Isotope hydrograph separation is a technique that separates the contributions of new and old waters to stream hydrograph, which has been~~

50 ~~extensively used to track the runoff components and water flow in permafrost catchment (Taylor et al., 2002; Carey and Quinton, 2005; Li et al., 2020a; Yang et al., 2016)~~. Mean residence time (MRT) ~~refers to is defined as~~ the average time required for input water to travel through water channels and to reach a catchment outlet in either a vertical or a horizontal flow path (McDonnell et al., 2010; Shah et al., 2017). MRT is a fundamental hydraulic descriptor that can be used to reveal information

55 ~~regarding about~~ water storage, flow paths as well as water sources within a particular catchment (Shah et al., 2017; Mcguire and McDonnell, 2006; Dunn et al., 2007). Furthermore, catchment water MRT is crucial for examining catchment response to environmental changes (Mcguire et al., 2002). Therefore, by quantifying the MRT, the hydrological sensitivity to climatic changes can be identified (Zhou et al., 2021a). Moreover, this parameter provides new insights into catchment functions and runoff processes (Farrick and Branfireun, 2015). The influencing factors associated with MRT have been ~~analysed analyzed~~

60 by previous studies; however, most of them focused on the influencing factors of the catchment area, topographical factors, groundwater contribution, soil properties, and flow path length (Dunn et al., 2007; Ma et al., 2019b; Soulsby and Tetzlaff, 2008; Rodgers et al., 2005b). Therefore, the influence of permafrost changes, climatic factors, and vegetation variations on catchment MRT in a high-altitude permafrost catchment is ~~rarely seldom~~ evaluated. ~~For instance, Song et al. (2017) and Yang et al. (2021) have investigated water age in permafrost catchment in the hinterland of TP and explored the influence of~~


65 vegetation and climatic factors on water age.— Effects of permafrost freeze–thaw cycles on water MRT in Arctic permafrost  
catchment have also been reported (Tetzlaff et al., 2018). Meanwhile, given the complexities of underlying surface alongside  
the remoteness and logistical difficulties associated with data collection in high-altitude permafrost catchments, few studies  
have examined such permafrost catchments to estimate water MRT using isotopic tracers by long time series sampling.

The Tibetan Plateau (TP) is the source region of many large Asian rivers and is often referred as the “Water Tower of Asia”.  
However, the TP is the most vulnerable among the world's water towers under the changing climate in terms of water-supplying  
70 role, the downstream dependence of ecosystems, and societal impacts (Immerzeel et al., 2020). The permafrost with the highest  
elevation and largest area in the mid-latitudes is mainly distributed in the TP, which has experienced significant and extensive  
degradation (Cheng et al., 2019). The climatic influences of MRT are non-stationary and manifested at an annual scale (Soulsby  
and Tetzlaff, 2008). Therefore, in this study, we estimated MRT ~~by~~ using long-term water stable isotopic data (8-year data for  
stream water and 5-year data for supra-permafrost water) from a high-altitude (5,300 m a.s.l.) permafrost catchment in the  
75 central TP. The main objectives of this study ~~were~~ are to: (1) characterize isotope composition of the catchment water; (2)  
elucidate the potential drivers of isotope variations; (3) identify runoff components ~~by~~ using two-component hydrograph  
separations; (4) estimate an approximation of water MRT of permafrost catchment ~~by~~ using a sine-wave exponential model;  
~~and~~ (5) quantify the climate and permafrost changes effects on permafrost hydrological process ~~by~~ using the estimated  
catchment water MRT. The findings from our study will expand deepen our understanding of the hydrological process in  
80 permafrost regions under global warming. ~~and will be important for water resources supply and safety in the TP.~~

## 2 Materials and methods

### 2.1 Study area

This study was conducted in the Xiaoliuyu catchment located in the source region of the Yangtze River in the central TP. ~~It,~~  
with ~~has~~ an average altitude of 5,300 m a.s.l. and a drainage area of 2.17 ~~km<sup>3</sup>~~ km<sup>2</sup>, which falls under the transition zone between  
85 the monsoon and non-monsoon regions (Fig. 1). This study area is unique given the presence of extensive permafrost. In  
particular, the maximum active layer thickness of permafrost can be > 300 cm, according to the observational data. The area  
is covered by sedge and wormwood, including species such as *Kobresia capilifolia*, *Kobresia pygmaea*, and *Kobresia humilis*  
(Wang et al., 2020a).



90

Figure 1. (a) map of permafrost thermal stability on the Tibetan Plateau; (b) map of the study area and the sampling sites; (c-d) images of the frozen river in this catchment during the cold season and the runoff generated during the warm season. (e) **Photograph of the underlying surface of the catchment and meteorological station.** Thermal stability of permafrost distribution data are from

The climate in the study area features a cold and dry season from October to May and is primarily controlled by westerlies, while warm and wet season from June to September is controlled by the Indian monsoon. The catchment receives > 90 % of the total annual precipitation during the warm and wet seasons (Li et al., 2016). The annual mean air temperature of this area is  $-5.3\text{ }^{\circ}\text{C}$ , while the annual mean precipitation is 491.9 mm. The river in the catchment freezes during the cold season and only generates runoff during the warm season (see Fig. 1b and d). Figure 2 shows the daily variations of precipitation, air temperature, soil moisture and temperature (0–300 cm). The air temperature increases to  $0\text{ }^{\circ}\text{C}$  in early May, indicating the onset of frozen soil thaw, while the peak air temperature is observed in July–August. Soil temperature is also an important factor, that governs the freezing and thawing of soil water in permafrost regions. Notably, the variations of soil moisture agree with the variations of soil temperature in the region. In early May, soil temperature increases to  $>0\text{ }^{\circ}\text{C}$ , thereby, rapidly increasing soil moisture and manifesting the onset of rapid thawing. These seasonal dynamics of the soil water content are altered by freeze-thaw processes in the active layer (Wang et al., 2009).




Figure 2. Temporal variation of soil moisture, soil temperature, air temperature, and precipitation from 2017 to 2020.

## 110 2.2 Permafrost and meteorological data

The soil temperature data (active layer bottom temperature) and active layer thickness of permafrost were obtained from the book of *Blue Book on Climate Change in China 2021*, which published long-term data, related to the permafrost in TP (Cma Climate Change Centre, 2021). The normalized differential vegetation index (NDVI) can reflect the growth status and coverage of vegetation. We collected NDVI data from MOD13A2 products provided by the NASA, USA, with temporal and spatial

115 resolutions of 16 days and 1-~~km~~ × 1 km, respectively (<https://adsweb.modaps.eosdis.nasa.gov/>). In our study, the spatial  
 resolution of NDVI data was unified to 25 × 25 m by resampling. The growing-season NDVI was determined using the  
 maximum synthesis method. Meteorological data, including precipitation amount, ~~relative air humidity~~ as well as air  
 temperature are available from the weather stations within the Xiaoliuyu catchment ([Fig. 1](#)). The soil temperature, active layer  
 thickness of permafrost, NDVI, and the meteorological data were obtained for 2012–2020, as shown in [Table 1](#).

120

**Table 1. Annual permafrost, normalized differential vegetation index (NDVI), and meteorological data from 2012–2020. The symbol “—” indicates no data were available for that year.**

Data type	2012	2013	2014	2015	2016	2017	2018	2019	2020
Active layer thickness (cm)	223	229	228	237	240	240	245	243	237
Soil temperature (°C)	-1.20	-1.19	-1.10	-1.30	-1.00	-0.91	-0.80	-1.31	-1.40
Air temperature (°C)	-5.69	—	-5.97	-4.84	-4.71	-5.00	-5.15	-5.60	-5.50
Precipitation(mm)	527	547	674	307	541	486	319	455	626
<del>Relative humidity (%)</del>	<del>53.78</del>	<del>—</del>	<del>54.8</del>	<del>49.03</del>	<del>50.37</del>	<del>53.55</del>	<del>53.32</del>	<del>57.38</del>	<del>56.29</del>
NDVI	0.2312	0.1336	0.1362	0.1842	0.1880	0.1771	0.1400	0.1807	0.1769

### 2.3 Field sampling and isotope analysis

125 ~~Note that the high sampling frequency can more efficiently identify substantial modification of the catchment hydrology and  
 can provide more information of hydrological processes~~ (Stockinger et al., 2016). In this study, the continuous sampling and  
 high sampling frequency of stream water, supra-permafrost water, and precipitation were conducted from June to October.  
 The sampling sites are shown in Figure 1- and detailed information of sampling sites is summarised in- [Table 2](#). In total, the  
 416 precipitation samples were collected during the observation period in bulk collectors at the Tanggula Cryosphere and  
 Environment Observation Station (TaCOS), Chinese Academy of Sciences at the altitude of 5,050 m a.s.l. [Liquid precipitation](#)  
 130 ~~samples were collected immediately following every precipitation event using bulk collector to minimize the effects of  
 evaporation. Solid precipitation (snow) samples were collected into a plastic bag and taken to a warm place to be thawed,  
 following which water samples were transferred into 50-mL PE bottles. The precipitation samples were immediately collected  
 after every precipitation event to reduce the impact of evaporation.~~ The groundwater in the permafrost region can be classified  
 into three categories: supra-permafrost water, intra-permafrost water as well as sub-permafrost water (Gao et al., 2021; Cheng  
 135 and Jin, 2013). Supra-permafrost water is the most widely distributed groundwater type in the permafrost regions of TP, which  
 is mainly stored in the permafrost active layer (Li et al., 2020a). We collected 755 stream water samples and 296 supra-  
 permafrost water samples at approximately 1-day intervals in the Xiaoliuyu catchment from June to October for each year ([Fig.  
 1](#)). As the stream water and the permafrost layer are frozen due to low temperature in the cold season, the stream and supra-  
 permafrost water is simply generated during the warm season ([Fig. 1](#)). The supra-permafrost water samples were taken from

140 a well, that was drilled at a depth of 1.5 m. Given the logistical constraints in this study area, only a single stream and supra-permafrost water sampling site were collected.

All samples were collected in individual 50-~~ml~~-ml PE bottles that were rinsed three times with the water from the source itself before sampling. The bottles were then sealed and stored in a refrigerator at a temperature of  $-10^{\circ}\text{C}$  to minimize the possibility of contamination from the external environment and isotopic fractionation, induced by liquid water evaporation.

145 Hydrogen ( $\delta\text{D}$ ) and oxygen ( $\delta^{18}\text{O}$ ) stable isotopes composition of all water samples were measured by using Liquid-Water Isotope Analyzer (DLT 100, Los Gatos, USA) at the State Key Laboratory of Cryospheric Sciences, Chinese Academy of Sciences. The water samples were analysed ~~analyzed~~ six times. The first two results were discarded to eliminate “memory effects”, and the average of the last four results was used as the isotope estimate of the water samples. The results were reported in per mil (‰) units relative to the Vienna Standard Mean Ocean Water (V-SMOW). The precision of measurement for  $\delta^{18}\text{O}$

150 and  $\delta\text{D}$  was  $\pm 0.2\text{‰}$  and  $\pm 0.6\text{‰}$ , respectively.

**Table 2. Information of samples and sampling sites.**

Sample type	Years	Altitude (m a.s.l.)	No. of samples
Precipitation	2012, 2014–2020	5129	416
Stream water	2012, 2014–2020	5139	756
Supra-permafrost water	2012, 2017–2020	5146	296

## 2.4 Isotope hydrograph separation method

The isotopic hydrograph separation (IHS) method is based on a mass balance approach. It can be used to estimate the contributions of diverse potential water sources contributing to streamflow ~~sources to terminal water~~ using isotope ( $\delta^{18}\text{O}$  or

155  $\delta\text{D}$ ) as a tracer (Genereux, 1998; Uhlenbrook and Hoeg, 2003). The two-component method can be formalized by using Eqs. (1) and (2), shown below:

$$Q_s = Q_p + Q_e, \quad (1)$$

$$Q_s C_s = Q_p C_p + Q_e C_e, \quad (2)$$

where  $Q_s$ ,  $Q_p$  and  $Q_e$  represent stream water, pre-event (supra-permafrost water) and event water (precipitation) volumes, respectively;  $C_s$ ,  $C_p$  and  $C_e$  are the corresponding values of  $\delta^{18}\text{O}$  or  $\delta\text{D}$ . The contributions of pre-event water and event water to stream water can be calculated by combining Eqs. (1) and (2) to obtain Eqs. (3) and (4):

165 
$$f_p = \frac{C_e - C_t}{C_e - C_p} \times 100\%, \quad (3)$$

$$f_e = \frac{C_t - C_p}{C_e - C_p} \times 100\%, \quad (4)$$

where  $f_p$  and  $f_e$  represent relative contribution ratio of pre-event water and event water to stream water, respectively. In this study, the precipitation-weighted average of precipitation isotopes was used to assess stream water components on a monthly scale to determine the mechanism of runoff process in permafrost catchments.

The uncertainty in hydrograph separations generally included two aspects, one is the analysis error of tracer concentrations, while the other is the spatial and temporal variations of the tracer of components (Uhlenbrook and Hoeg, 2003), calculated ~~The uncertainty of the calculated fractions of event and pre-event water~~ using the Gaussian error propagation technique (Genereux, 1998):

$$w_y = \sqrt{\left(\frac{\partial y}{\partial x_1} w_{x1}\right)^2 + \left(\frac{\partial y}{\partial x_2} w_{x2}\right)^2 + \dots + \left(\frac{\partial y}{\partial x_n} w_{xn}\right)^2} \quad (5)$$

where  $w$  represents the uncertainty in the variable specified in the subscript and  $y$  is the contribution of a specific streamflow component  $x$  to streamwater.

## 2.5 Estimation of mean residence times

Earlier studies have demonstrated that the sine-wave exponential model can be used to estimate MRT using the seasonal variations of isotope composition in precipitation and stream water (Mcguire et al., 2002; Mcguire and Mcdonnell, 2006; Soulsby and Tetzlaff, 2008; Ma et al., 2019b; Zhou et al., 2021a). Mathematically, the movement of conservative tracer through a catchment can be expressed by the convolution integral (Mcguire and Mcdonnell, 2006; Simin et al., 2013), which states that the tracer concentration of output water  $\delta_{out}(t)$  at any time and input tracer  $\delta_{in}(t-\tau)$  that enter uniformly into the catchment in the past  $t-\tau$ , which becomes lagged by its transit time distribution  $g(\tau)$ . The common transit time distribution model  $g(\tau)$  used in hydrologic systems include: exponential, exponential-piston flow, and dispersion models (Simin et al., 2013). The exponential model describes a catchment with flow times that are exponentially distributed, including pathways with very short transit times (Mcguire and Mcdonnell, 2006), which assumed that the system is in steady-state conditions and operates as a perfect mixer (Sánchez-Murillo et al., 2015; Smith, 1984; Chiogna et al., 2014). In our study, the exponential model was used to estimate water MRT:

$$\delta_{out}(t) = \int_0^{\infty} g(\tau) \delta_{in}(t - \tau) d\tau, \quad (65)$$

$$g(\tau) = \tau_m^{-1} \exp\left(-\frac{\tau}{\tau_m}\right), \quad (76)$$

where  $\tau$  is the transit time,  $t$  is the time of tracer exit from the catchment and  $t-\tau$  represents the time of tracer enter into the catchment, and  $\tau_m$  is mean residence time (MRT).

In our study, the seasonal variations of  $\delta^{18}\text{O}$  in precipitation, supra-permafrost water, and stream water were modelled using sine-wave function~~the seasonal variations of  $\delta^{18}\text{O}$  were modelled using sine-wave function to fit the seasonal  $\delta^{18}\text{O}$  variations in precipitation, supra permafrost water, and stream water~~, defined by Eq. (78):

$$\delta(t) = X + A [\cos(ct - \theta)], \quad (87)$$

where  $\delta(t)$  is the modelled  $\delta^{18}\text{O}$  (‰),  $X$  is the mean measured  $\delta^{18}\text{O}$  (‰),  $A$  is the amplitude of the measured  $\delta^{18}\text{O}$ ,  $c$  is the radial frequency constant given as  $c=2\pi/153 \text{ days}=0.04105 \text{ rad d}^{-1}$  in our study (the total number of days from June to October



considered in this study amounted to 153) and  $\theta$  is the phase lag of modelled  $\delta^{18}\text{O}$  in radians. The overall performance of the sine-wave model was evaluated ~~by~~ using the goodness of fit ( $R^2$ ) and root mean square error (RMSE).

Analytical solution of the MRT ( $\tau_m$ ) for the exponential model can be derived by combining Eq. (57) with Eq. (78) (Mcguire and McDonnell, 2006):

$$MRT = c^{-1} \sqrt{\left[ \left( \frac{A_{z1}}{A_{z2}} \right)^2 - 1 \right]}, \quad (98)$$

where  $A_{z1}$  is the amplitude of precipitation  $\delta^{18}\text{O}$ ,  $A_{z2}$  is the amplitude of modelled  $\delta^{18}\text{O}$  in supra-permafrost or stream water in our study, and  $c$  is the radial frequency, defined in Eq. (78). The uncertainty of the MRT estimates were quantified by determining the 95% confidence of the fitted sine-wave's amplitude. Specifically, applying the 95% confidence of the fitted sine-wave's amplitude to Eq. (89) producing MRT error ~~(the upper and lower 95% confidence boundaries of the estimated MRT)~~ (Morales and Oswald, 2020).

### 3 Results ~~and discussion~~

#### 3.1 Stable isotope composition ~~in different waters and $\delta\text{D}$ - $\delta^{18}\text{O}$ relationships of precipitation, stream, and supra-permafrost water~~

~~The hydrological examination can benefit from the following phenomenon. Meteoric water is the main driver of hydrological processes at the catchment scale, while the hydrogen and oxygen isotope composition in different waters are correlated (Chiogna et al., 2014). For a detailed examination of the differences of isotopic composition in the different waters and their mutual transformation, the relationships between  $\delta^{18}\text{O}$  and  $\delta\text{D}$  for precipitation, stream water, and supra-permafrost water were quantified (Fig. 3). During the sampling periods, the isotopic composition in precipitation ( $-30.9$  to  $2.9\text{‰}$  for  $\delta^{18}\text{O}$ ;  $-244.2$  to  $34\text{‰}$  for  $\delta\text{D}$ ) strongly varied, compared with that in stream water ( $-19.6$  to  $-4.0\text{‰}$  for  $\delta^{18}\text{O}$ ;  $-146.3$  to  $-39.8\text{‰}$  for  $\delta\text{D}$ ) and in supra-permafrost water ( $-17.5$  to  $4.6\text{‰}$  for  $\delta^{18}\text{O}$ ;  $-127.6$  to  $50.9\text{‰}$  for  $\delta\text{D}$ ) (Table 3). This finding indicates that precipitation was an important recharge source for supra-permafrost and stream water in the analysed permafrost catchment. In addition, ice meltwater from deeper soil layers is also a source of water replenishment (Wang et al., 2022; Sugimoto et al., 2003; Throckmorton et al., 2016), although a previous study found that melting ground ice in permafrost had little contribution to the observed runoff variations (Landerer et al., 2010).~~


We identified the substantial difference among the water line in different water: the stream water (SWL:  $\delta\text{D}=6.66\delta^{18}\text{O}-7.61\text{‰}$ ), supra-permafrost water (PWL:  $\delta\text{D}=6.10\delta^{18}\text{O}-15.89\text{‰}$ ), and local meteoric water line (LMWL:  $\delta\text{D}=8.01\delta^{18}\text{O}+13.48\text{‰}$ ). The lower slopes of SWL (6.66) and PWL (6.1) strikingly indicate the non-equilibrium fractionation caused by evaporation, reducing the slope of  $\delta\text{D}$ - $\delta^{18}\text{O}$  correlation line (Throckmorton et al., 2016) ~~(Xia et al., 2021)~~. Alternatively, we surmise that there could be precipitation mixed with water from previous events stored in the permafrost active layer. In contrast, supra-permafrost water exhibited the lower slope, which was most likely caused by thawing and

freezing of the active soil layer (Wang et al., 2009). During freezing and thawing, with the thickening of the active layer, part of ice melt water (old water) will be mixed with precipitation. This old water tends to have a large water age and is subjected to evaporation over a long period, thereby lowering the slope of water lines (Throckmorton et al., 2016; Song et al., 2017)–.

This can be explained by the longer residence time of water in the permafrost active layer, resulting in lower slope of water lines. These findings are similar to previous results obtained for the Zuomaokong watershed in the hinterland of the TP (Song et al., 2017). Hence, the lower slope of the water line of supra-permafrost water may indicate relatively longer catchment water MRT.

**Table 3. Mean, range, and standard deviation of  $\delta^{18}\text{O}$  (‰) and  $\delta\text{D}$  (‰) for precipitation, stream water, and supra-permafrost water.**

Sample type	Mean		Minimum		Maximum		Standard deviation	
	$\delta^{18}\text{O}$	$\delta\text{D}$	$\delta^{18}\text{O}$	$\delta\text{D}$	$\delta^{18}\text{O}$	$\delta\text{D}$	$\delta^{18}\text{O}$	$\delta\text{D}$
Precipitation	-13.2	-92.6	-30.9	-244.2	2.9	34.0	7.1	58.0
Supra-permafrost water	-14.0	-101.5	-17.5	-127.6	-4.6	50.9	1.7	10.8
Stream water	-13.5	-97.4	-19.6	-146.3	-4.0	-39.8	2.4	16.3




**Figure 3. Relationships between  $\delta^{18}\text{O}$  and  $\delta\text{D}$  of precipitation, stream water, and supra-permafrost water. (a) all isotope data; (b) the data, constrained by yellow square.**

### 3.2 Temporal variations of stable isotopes in different waters and their potential driversstream and supra-permafrost

The variations of  $\delta^{18}\text{O}$  and  $\delta\text{D}$  compositions in different water samples exhibited similar tendencies, as seen in Figure 3–, therefore, Due to this,  $\delta^{18}\text{O}$  was selected as the representative isotope in the following analysis. The temporal variations of  $\delta^{18}\text{O}$  in stream and supra-permafrost water are shown in Figure 4. The analysis revealed that substantial seasonal variability of stable isotope signature was identified in different waters. The precipitation isotopes deplete heavy isotopes during the period of the Indian summer monsoon (from late May to early September) and enrich heavy isotopes at the beginning and end of the summer monsoon period (late May and mid-October). This pattern is similar to those of the monsoon region of the southern TP, due to

245 the shifting moisture source between the Bay of Bengal and the southern Indian Ocean (Yao et al., 2013). The stream water samples also exhibited a depletion in the isotopic signature, which is virtually consistent with the precipitation results from the overall variation trend (Fig. 4). The isotope compositions in supra-permafrost water for the certain year (such as 2018 and 2019) also reflected the isotope signals of precipitation, but exhibited the least variability for overall supra-permafrost water (lowest standard deviation shown in [Table 3](#)), compared to precipitation and stream water. This finding suggests that mixing  
250 processes within the active layer attenuate variations of  $\delta^{18}\text{O}$  signal from precipitation.




**Figure 4. Comparison of variations of precipitation  $\delta^{18}\text{O}$  (‰), stream and supra-permafrost water  $\delta^{18}\text{O}$  (‰) during the warm seasons from 2017 to 2020.**

The seasonal dynamics of the soil temperature and moisture of the active layer are the most important drivers of hydrological processes in the permafrost regions (Sugimoto et al., 2003; Woo and Xia, 1996; Wang et al., 2015) (~~Wang et al., 2009; Wang et al., 2015~~). Rising air temperature promotes permafrost thawing and thickening of the active layer, thereby increasing soil moisture since more water sources with different isotopic compositions, such as atmospheric precipitation and meltwater from subsurface ice, input or release into the permafrost active layer (Sugimoto et al., 2003; Throckmorton et al., 2016; Song et al., 2017). In our study area, the stable isotope compositions in supra permafrost and stream water may have been significantly altered by the freeze thaw cycle of permafrost active layer. Hence, we conducted the correlation analysis between the isotope compositions in supra-permafrost and stream water, as well as soil temperature and soil moisture (Fig. 5).


The isotopic compositions in supra-permafrost water were ~~found to be~~ significantly associated with soil temperature (Fig. 5c:  $R^2=0.18$ ,  $P<0.000$ ) ( $r=0.201$ ,  $P<0.01$ ). Moreover, we identified a variable relationship between the supra-permafrost water isotopes and soil temperature (first increasing and then decreasing). ~~A similar relationship was observed for soil moisture (but not significant).~~ When soil temperature was  $< 4^\circ\text{C}$ , the active layer was relatively thinner, and less water was stored in the  
265

permafrost active layer, which is more susceptible to evaporation, causing isotope enrichment. With the further increase of soil temperature ( $> 4^{\circ}\text{C}$ ), the active layer becomes thicker and the soil moisture also increases, yielding the supra-permafrost water recharged by greater amounts of precipitation that depleted heavy isotopes (precipitation depleted heavy isotopes in August). The positive correlation between supra-permafrost water with precipitation isotopes (Fig. 5f:  $R^2 = 0.39$ ,  $P < 0.000$ ) ( $r = 0.510$ ,  $P < 0.01$ ), and the negative correlation between precipitation isotope with soil temperature ( $r = -0.447$ ,  $P < 0.01$ ) and with soil moisture ( $r = -0.564$ ,  $P < 0.01$ ) also confirms this finding. These findings reveal the differences in the water movement mechanisms at different stages of permafrost freezing and thawing processes.


The correlation analysis revealed that the isotopic compositions in stream water are negatively correlated with soil temperature (Fig. 5a:  $R^2 = 0.08$ ,  $P < 0.000$ ) (Pearson's correlation coefficient ( $r$ ) =  $-0.316$ ,  $P < 0.01$ ) and soil moisture (Fig. 5b:  $R^2 = 0.31$ ,  $P < 0.000$ ) ( $r = -0.560$ ,  $P < 0.01$ ). They are also strongly positively correlated with the isotopic compositions in precipitation (Fig. 5e:  $R^2 = 0.44$ ,  $P < 0.000$ ) ( $r = 0.644$ ,  $P < 0.01$ ). These findings indicate that the stream water was controlled by precipitation and freeze-thaw cycle of the permafrost active layer. Notably, our correlation results are in line with the previous studies in the Zuomaokong watershed of central TP (Song et al., 2017). At the same time, a very strong correlation was observed between stream water and supra-permafrost water isotopes (Fig. 5d:  $R^2 = 0.23$ ,  $P < 0.000$ ). These significant positive correlations between precipitation and permafrost isotopes and stream water isotopes indicates that precipitation and supra-permafrost water are important recharge sources of stream water. Although this may be related to the fact that thawing of permafrost releases frozen soil water that depleted in isotopes. But previous study found that melting ground ice in permafrost was not a source of the observed runoff variations, but rather the thickening of the active layer (Landerer et al., 2010). In contrast, the isotope compositions in stream water were much more responsive to precipitation than to supra-permafrost water. As the active layer of permafrost can effectively store precipitation and the groundwater storage is a long term accumulation of local precipitation. Thus, the isotope compositions in groundwater generally express integrative characteristics of seasonal precipitation inputs, which are less sensitive to the recharge of precipitation. However, the stream water is mostly affected by seasonal precipitation, thereby, somewhat following the variability of precipitation characteristics. This indicates that precipitation was essential for stream water, whereas the freezing and thawing process of the active layer was pivotal in controlling supra-permafrost water, which reflects the influence of permafrost as a specific surface aquifer on the hydrological process.



**Figure 5. Correlation analysis between stable isotopes and influencing factors. (a) and (b) correlations between stream water  $\delta^{18}\text{O}$  with: soil temperature and moisture, respectively; (c) and (d) correlations between supra-permafrost water  $\delta^{18}\text{O}$  with: soil temperature and stream water  $\delta^{18}\text{O}$ , respectively; (e) and (f) correlations between precipitation  $\delta^{18}\text{O}$  with: stream and supra-permafrost water  $\delta^{18}\text{O}$ , respectively; Red line is the fitted line; Soil moisture and temperature data used in correlation analysis were normalized. “-” presents dimensionless.**



300 **Figure 5. Correlation analysis between stable isotopes and influencing factors. Coloured rectangle shows the significance of the correlation ( $P < 0.01$ ). The larger the value and the darker the colour the more significant of the correlation between variables. Abbreviations: P = Precipitation  $\delta^{18}\text{O}$  (‰); SW = Stream water  $\delta^{18}\text{O}$  (‰); SP = Supra-permafrost water  $\delta^{18}\text{O}$  (‰); ST = Soil temperature ( $^{\circ}\text{C}$ ); SM = Soil moisture (%).**




305 **Figure 6. Correlation between  $\delta^{18}\text{O}$  in supra-permafrost water with: (a) soil temperature (b) soil moisture.**

### 3.3 Two-component hydrograph separations of stream water


The differences of the isotopic composition and water lines between the different waters allow distinguishing the sources of stream water. The EMMA model has been used to identify the mixing processes and quantify the contribution of each

endmember. The monthly mixing diagram using the mean  $\delta^{18}\text{O}$  and  $\delta\text{D}$  showed that the isotope values in stream water are very close to the other endmember (precipitation or supra-permafrost water), indicating that the component of stream water was dominated by different sources at different stages (Fig. 6). However, in some mixing diagrams, stream water was located outside the range composed of the two end-members (atmospheric precipitation and supra-permafrost water), possibly due to the influence of vegetation transpiration and soil evaporation on precipitation and ground ice that were mixed and stored in the active layer (Li et al., 2020b). Nevertheless, the isotopic composition in stream water is very close to one of the endmembers (Fig. 6). Overall, supra-permafrost water and precipitation can be treated as the two end-member in hydrograph separations of stream water. For quantitative evaluation of the results above, the source proportion of stream water was quantitatively determined using the isotopic data and the IHS model (Eqs. 1–4). The results indicate that precipitation and supra-permafrost water contributed  $35 \pm 2\%$  and  $65 \pm 2\%$  of the total discharge of stream water, respectively. Seasonal patterns showed that the precipitation contribution decreased from June to August, then increased in September; for the supra-permafrost water, the contribution to streamflow increased from June to August, then slightly decreased in September (Fig. 7). During the initial thawing stage of permafrost (June), precipitation recharge was the primary source of stream water, approximately accounting for 78% of the total discharge of streamflow. Notably, in June 2018 and 2019, we observed no supra-permafrost water in the sampling well, therefore, the stream water during these periods is almost solely attributed to precipitation. However, the stream water is primarily derived from the active layer water in the thawing and end thawing stages of permafrost (July–September), approximately accounting for 79% of the streamflow. Especially in August, the contribution of supra-permafrost water to stream water can reach 98%. These findings suggested that supra-permafrost water was the dominant source of the stream water during the warm season in the study area.



**Figure 6. Monthly mixing diagram using the mean  $\delta^{18}O$  and  $\delta D$  values for stream water. The orange dotted ellipse indicates that the isotopic value in stream water is very close to the other end member.**







**Figure 7. Monthly variation of components in stream water. “~” indicates approximate estimation of component contributions to stream water.**

### 3.4 Estimation of mean residence ~~and its association with environmental variables~~ ~~time using seasonal variability of water isotopes~~


The modelled amplitudes can reflect the observed patterns of variability of water isotope compositions due to great mixing; meanwhile, seasonal variations in the output waters are far more significant with larger amplitude, thereby, indicating greater responsiveness to recent precipitation inputs (Rodgers et al., 2005a). Moreover, we found that different water isotopes showed obvious seasonal variation, and precipitation was an important input of stream and supra-permafrost water. Thus, we fit seasonal sine-wave curves to the annual  $\delta^{18}\text{O}$  variations in precipitation, stream, and supra-permafrost water. The sine-wave regression parameters for  $\delta^{18}\text{O}$  in precipitation, supra-permafrost water, and stream water are shown in Figures 8–10. We found that the modelled  $\delta^{18}\text{O}$  fit well to the observed isotope values, with RMSE of 4.35–6.58‰ for precipitation, 0.37–1.95‰ for supra-permafrost water, and 0.72–1.99‰ for stream water. Meanwhile, the results of periodic regression analysis of isotope compositions in different waters were all statistically robust ( $P < 0.01$ ). The  $\delta^{18}\text{O}$  seasonal variations of supra-permafrost and stream water exhibited weak amplitudes, compared with the precipitation, which is a consequence of the mixing processes and longer residence time. The mean amplitude of stream water (1.54‰) in our study reasonably agrees with the previous results in the Zuomaokong watershed of hinterland of the TP (1.76‰), which applied the same method to calculate the mean amplitude of stream water  $\delta^{18}\text{O}$  in five permafrost catchments (Song et al., 2017).



**Figure 8. Fitted sine-wave regression models to  $\delta^{18}\text{O}$  in precipitation in Xiaoliuyu catchment during 2012, 2014–2020.  $A$  is the amplitude and RMSE is root mean square error for the modelled isotopic signature.**



**Figure 9. Fitted sine-wave regression models to  $\delta^{18}\text{O}$  in supra-permafrost water in Xiaoliuyu catchment during 2012, 2017–2020.  $A$  is the amplitude and RMSE is root mean square error for the modelled isotopic signature.**



360 **Figure 10. Fitted sine-wave regression models to  $\delta^{18}\text{O}$  in stream water in Xiaoliuyu catchment during 2012, 2014–2020. A is the amplitude and RMSE is root mean square error for the modelled isotopic signature.**

Then, the sine-wave regression parameters were translated into the estimates of water MRT ~~by~~ using Eq. (69). The estimated results of MRT are shown in Figure 12.11. The calculated MRT for the stream water ranged from 42 to 270 days, with a mean value of 100 days and a standard deviation of 68 days. Meanwhile, for supra-permafrost water, the MRT varied from 23 to 596 days, with a mean value of 255 days and a standard deviation of 229 days. The estimated MRT revealed a large annual variability, regardless of stream water or supra-permafrost water. ~~It should be noted that the estimated MRT of supra-permafrost water was distinctly longer than that of stream water. The longer MRT reflects more complex soil water retention and recharge processes (Ma et al., 2019b). In turn, a short MRT of the stream water indicates a relatively rapid response of surface water to precipitation. The estimated MRT of stream and supra-permafrost water in our catchment were shorter, compared to those estimated in most previous studies from non-permafrost or high-latitude permafrost catchments (Table 4).~~

370 ~~This may be related to whether there is groundwater recharge or not. Permafrost acts as an aquiclude, while being usually characterized by rapid hydrograph responses (Tetzlaff et al., 2018). Moreover, it cuts off the interaction and mixing between deep groundwater and surface water and supra-permafrost water. For instance, (Rodgers et al., 2005b; Soulsby et al., 2006b) have reported that catchment water MRT was correlated with the percentage groundwater contributions to stream water in non-permafrost catchment. Furthermore, previous studies have revealed longer MRT for deep groundwater (Table 4). For~~

375 instance, the MRT of the mixture of supra-permafrost and intra-permafrost water ranges from ~1 to 55 years in the permafrost  
 regions of the Arctic region (Hiyama et al., 2013). In particular, deep groundwater MRT in the mountainous Brugga basin  
 ranged from 6 to 9 years. Thus, the mixing or interaction between stream water and groundwater increases the stream water  
 MRT in the non-permafrost catchments. However, this interaction is suspended by the permafrost layer in our catchment,  
 causing the shorter MRT. This implies that the hydrological processes in high-altitude permafrost region are unique, compared  
 380 with non-permafrost or high-latitude permafrost regions.

**Table 4. Statistics of MRT-related research results. The symbol “—” indicates no data were available in the references.**

Site	Altitude (m)	Land type	Water type	MRT	References
<b>Our study site</b>	5100–5435	Permafrost	Stream water	100 days	This study
Siksik Creek	50–100	Permafrost	Stream water	548 days	(Tetzlaff et al., 2018)
Mandava	383	Non-permafrost	Stream water	444 days	(Sanda et al., 2017)
Allt Mhareaidh	699	Non-permafrost	Stream water	263–445 days	(Rodgers et al., 2005a)
Upper Váh	1500	Non-permafrost	Stream water	390–570 days	(M. et al., 2011)
Dee	1000	Non-permafrost	Stream water	600 days	(Soulsby et al., 2010)
Minjiang	300–7100	Non-permafrost	Stream water	293–870 days	(Xia et al., 2021)
<b>Our study site</b>	5100–5435	Permafrost	Groundwater	255 days	This study
Lena River	130–160	Permafrost	Groundwater	1–55 years	(Hiyama et al., 2013)
Himalaya	1600–5200	Non-permafrost	Groundwater	4.5 months	(Shah et al., 2017)
Vermigliana	1221	Non-permafrost	Groundwater	1.3 years	(Chiogna et al., 2014)
Allt a' Mhareaidh	300–1111	Non-permafrost	Groundwater	>5 years	(Soulsby et al., 2000)
Brugga	434–493	Non-permafrost	Groundwater	6–9 years	(Uhlenbrook et al., 2002)
Karst	—	Non-permafrost	Groundwater	161–1407 days	(Wang et al., 2020b)

**The symbol “—” indicates no data were available in the references.** **3.5 Factors determining the MRT of stream and supra-permafrost water**

385 Generally Theoretically, multiple factors affect the water storage in the active layer, including precipitation and soil temperature  
 in permafrost regions (Wright et al., 2008). The climate differences and variability may have substantially affected the MRT  
 estimates (Tetzlaff et al., 2007). In this study, Table 5 shows the quantified correlations between estimated MRT and soil  
parameters (active layer thickness and soil temperature), climate factors (air temperature, and precipitation, and relative  
humidity), and vegetation index as well (NDVI) were quantified (Table 4). The uncertainty of the estimated MRT may  
 390 influence these correlations (Hu et al., 2020), hence we considered the uncertainty of the MRT in regression analysis. The  
results showed that after considering the uncertainty of MRT, the  $R^2$  of the regression analysis for active layer thickness

was improved, with less obvious differences for the other factors. In the following analysis, we use the regression analysis results after considering the uncertainty.

As the buffer layer between the permafrost and atmosphere, the active layer is vulnerable to climate change (Xu and Wu, 2021). The increase of air temperature can alter the temperature of shallow permafrost due to strong land surface-atmosphere interactions. This, in turn, induces an increase of the active layer thickness of permafrost, allowing soil water to be moved and stored in the deeper soil layer. The correlation analysis showed that the MRT of supra-permafrost water exhibits a strong positive correlation with air temperature ( $R^2 = 0.9067, P < 0.05$ ) and soil temperature ( $R^2 = 0.7778, P < 0.05$ ) (Table 4). These strong correlations indicate that soil and air temperature are potentially efficient predictors of supra-permafrost water MRT. In contrast, Meanwhile, the relatively weaker-stronger positive correlations were identified between stream water MRT and air temperature ( $R^2 = 0.16$ ) and soil temperature ( $R^2 = 0.3879, P < 0.001$ ). These strong correlations may relate to the thickening of the active layer due the increase of air and soil temperature. Moreover, Because we also observed that MRT of supra-permafrost and stream water were negatively-weakly correlated with the active layer thickness ( $R^2 = 0.3559, P < 0.01$  and  $R^2 = 0.44-0.23, P < 0.05$ , respectively). The deeper active layer somewhat triggers longer catchment water MRT, because the increase of the thickness of active layer increases the soil and deeper flow paths, thereby slowing the catchment water response to the precipitation. In addition, the deeper active layer allows more water to be stored, precluding the accumulation of soil moisture deficits. These results also support the findings from previous studies in terms of a relationship between permafrost changes and residence time. In particular, (Wright et al., 2009) have stated that MRT of permafrost catchments is highly dependent on the annual development of the active layer. Moreover, (Lyon et al., 2010) have reported that the potential thaw of permafrost layers due to climate change could increase MRT at the catchment scale by 20–45%. A similar study in the non-frozen regions reported that MRT of forestland and shrubland water both increased with soil depth (Ma et al., 2019b).

Precipitation is an important part of the water cycle and is the main input for catchment water sources. However, precipitation in the permafrost region of TP significantly increased in recent decades (Zhao et al., 2019). Does increased precipitation affect runoff processes in permafrost watersheds? In our study, Meanwhile, the longest estimated MRTs (270 days for stream water and 596 days for supra-permafrost water) were observed in 2018 with the relatively lower precipitation amount (319 mm). Thus, we analyzed-analysed the correlation between MRT and precipitation-and relative humidity and found that supra-permafrost and stream water MRT are both negatively correlated with precipitation ( $R^2 = 0.6964, P < 0.01$ -and  $R^2 = 0.32, P < 0.01-0.49$ , respectively) (Table 4). Moreover, supra-permafrost water MRT also has a relatively weak negative correlation with relative humidity ( $R^2 = 0.36$ ) (Table 5). This indicates the increased precipitation and wetter climatic conditions may accelerate the water cycle process in permafrost regions. The larger precipitation corresponds to lower temperature, yielding a thinner active layer, which, in turn making the active layer water to be saturated sooner. This phenomenon subsequently triggers more water to rapidly flow into the river channel in the form of surface runoff, thereby, reducing the MRT of catchment water in the end. However, previous study suggested that much wetter climate than average probably causes higher MRT in the low altitude

temperate regions (Soulsby et al., 2006b). Our study does not resonate with these results, possibly because the aquiclude effect of permafrost reduces surface infiltration and enhances surface runoff generation (Gao et al., 2021).

Soil layer is the main source of nutrients and water required for vegetation growth (Xu et al., 2019). Water shortage in soil layer may occur when water MRT is too short, but longer MRT can hamper the water infiltration consequently to root anoxia, thus, affecting the plant growth (Ma et al., 2019b). Interestingly, we found that the stream and supra-permafrost water MRT are both negatively correlated with NDVI ( $R^2 = 0.2926, P < 0.05$  and  $R^2 = 0.53065, P < 0.01$ , respectively) (Table 2). In other words, Alternatively, an increase in vegetation coverage (high NDVI values) might lead to a longer-shorter MRT in our catchment.




Figure 11. Inter-annual variation of stream water MRT during 2012, 2014–2020, and supra-permafrost water MRT during 2012, 2017–2020.

Previous studies have noted the vegetation cover was one of the most important factors, that governs the hydrological processes and thermal cycles in permafrost catchment (Wang et al., 2012a). The decline of vegetation coverage induced increases of soil temperature and moisture, at which they accelerate the active soil thawing (Wang et al., 2012e). Therefore, the effect of vegetation on MRT is actually driven by the variations of thickness of the permafrost active layer. However, it remains unclear whether a positive feedback mechanism exists between vegetation and permafrost active layer changes or not. Moreover, the optimum residence time for vegetation growth should be elucidated in future studies as well.

In this study, comparisons of the controlling factors of MRT between stream and supra permafrost water indicated that supra permafrost water is more sensitive to environmental change, compared with stream water. Previous studies suggested that supra permafrost water in the seasonally thawed layer is sensitive to climate (Cheng and Jin, 2013), and is, therefore, significantly impacted by precipitation, temperature, and vegetation. From the influencing mechanism perspective, climate and

445 vegetation factors affected the MRT of stream and supra-permafrost water by modifying the active layer thickness of permafrost. It can be deduced that the estimated MRT of supra permafrost water is valuable for evaluating the extent of permafrost degradation. Most importantly, it can be used to infer the effects of long term climate, permafrost changes, and vegetation on the hydrologic regime in permafrost regions.

450 **Table 4. ~~Correlations~~ Relationships between active layer thickness, soil temperature, air temperature, precipitation, ~~relative humidity~~, NDVI, and MRT.  $x$  indicates the factor as the independent variable.**

	Factor	Regression based on mean MRT (days)			Regression based on MRT uncertainty (days)		
		Regression equation	$R^2$	Sig	Regression equation	$R^2$	Sig
<u>Supra-permafrost water</u>	ALT	$y=4.23\exp(0.24x)+48.62$	0.11	$P>0.05$	$y=1.43\exp(0.63x)+150.68$	0.44 ↑	$P<0.05$
	ST	$y=929x+1299$	0.87 ↑	$P<0.05$	$y=956x+1337$	0.78 ↑	$P<0.05$
	AT	$y=752x+4307$	0.69 ↑	$P<0.05$	$y=767x+4283$	0.67 ↑	$P<0.05$
	P	$y=-1.89x+1169$	0.58	$P=0.08$	$y=3479\exp(-0.005x)-125$	0.64 ↓	$P<0.01$
	NDVI	$y=0.103x^{-4.405}$	0.51	$P=0.07$	$y=0.143x^{-4.286}$	0.65 ↓	$P<0.01$
<u>Stream water</u>	ALT	$y=4.67\exp(1.11x)+71.56$	0.81 ↑	$P<0.01$	$y=3.02\exp(0.37x)+59.23$	0.59 ↑	$P<0.01$
	ST	$y=(1.7x+1.35)^{-0.26}$	0.81 ↑	$P<0.001$	$y=3.79E+14\exp(35.47x)+76$	0.79 ↑	$P<0.001$
	AT	$y=51x+372$	0.05	$P>0.05$	$y=42x+325$	0.01	$P>0.05$
	P	$y=4.94E+5(x^{-1.77})$	0.47 ↓	$P<0.01$	$y=1.67E+10(1.7x)^{-3.25}+52.6$	0.32 ↓	$P<0.01$
	NDVI	$y=1.422x^{-2.394}$	0.20 ↓	$P<0.05$	$y=1.50x^{-2.384}$	0.26 ↓	$P<0.05$

Note: ALT = active layer thickness, ST = Soil temperature (°C), AT = air temperature (°C), P = Precipitation (mm), NDVI= normalized differential vegetation index; Sig indicates statistical significance; ↑ and ↓ indicates significant trend of increase and decrease, respectively; Bold font indicates that it passed significance test of 0.05.

Factor	Stream water	Supra-permafrost water
Active layer thickness(cm)	$y=4.63x-995.14 R^2=0.23$	$y=3E-8e^{0.0943x} R^2=0.35$
Soil temperature (°C)	$y=211.96x+339.02 R^2=0.38$	$y=929.35x+1300.1 R^2=0.90$
Air temperature (°C)	$y=1231.1e^{0.5049x} R^2=0.16$	$y=752.19x+4307.9 R^2=0.77$
Precipitation (mm)	$y=378.76e^{-0.003x} R^2=0.49$	$y=-1.90x+1169.7 R^2=0.69$
Relative humidity (%)	$y=-7.62x+508.03 R^2=0.08$	$y=-83.38x+4829.4 R^2=0.36$
NDVI	$y=-1288.6x+327.59 R^2=0.27$	$y=-5723.6x+1292.2 R^2=0.53$

## 4 Discussion

### 4.1 Contribution of supra-permafrost water to stream water

460 Quantifying the components of stream water can provide insights into the hydrological effects of permafrost degradation (Li et al., 2020a). In this study, differences in the seasonal contributions of runoff components to stream water were observed. The contribution of supra-permafrost water during the thawing and end thawing stages of permafrost to stream water was higher

than that of the initial thawing stage. In June, the active layers of permafrost were gradually thawing as temperature increased, yet at this time, the active layer remained relatively thin, while the precipitation increased, resulting in most precipitation directly converging into the river. Under higher temperature and precipitation conditions in July and August, the strong thawing of permafrost occurred. The thickening of the permafrost active layer functions as a water reservoir, thereby allowing for more precipitation recharge into the active layer. Previous studies found that summer rain was the predominant source for water within the active layer in permafrost catchment (Throckmorton et al., 2016; Li et al., 2020b; Zhu et al., 2019), which is attributed to the relatively high permeability of the active layer (Li et al., 2020a). Then the active layer water produces a direct recharge to stream water with a contribution rate of 81% on average at this stage. As temperatures drop, the surface soil layer gradually begins to freeze and the bottom of the active layer approaches freezing in late September, thereby gradually decreasing the volume of supra-permafrost water due to the freezing processes of the aquifer. This phenomenon slightly decreased the contribution of supra-permafrost water to stream in September (~75%). More succinctly, seasonal variations in the freezing and thawing of permafrost directly trigger the runoff process (Li et al., 2020a).

In this study, approximately two-thirds of the stream water was attributed to supra-permafrost water. Previous studies have also reported that precipitation and thawing permafrost water contributes 55.2% and 44.8%, respectively, to the thermokarst lakes in the Beiluhe basin of interior TP (4,600 m a.s.l.) (Yang et al., 2016). A recent study has reported a greater contribution of supra-permafrost water (49%), compared with precipitation (34%), in the whole source region of the Yangtze River (SRYR) (Li et al., 2020a). The contribution rate of the supra-permafrost water to stream water in our catchment was relatively high compared with the whole SRYR. This finding is potentially related to the replenishment from other water sources, since for the whole SRYR, in addition to the precipitation and supra-permafrost water, there is a large amount of replenishment of glacier meltwater, which, to a certain extent, reduces the contribution of supra-permafrost water to stream water. In subarctic permafrost catchment, pre-event water was also a primary contributor of stream water (~90%) (Carey and Quinton, 2005). These studies demonstrated the significant contribution from cryosphere meltwater to water resources in high-altitude permafrost catchments. Considering that permafrost is widely distributed in the central TP and plays an important role in surface/groundwater exchange within the catchment, permafrost degradation will significantly influence the hydrological processes in alpine permafrost region in the context of climate warming.

#### **4.2 Potential driving mechanism for MRT variability** ~~Factors determining the MRT of stream and supra-permafrost water~~

In this study, the estimated MRT of supra-permafrost water was distinctly longer than that of stream water, which reflects the more complex water movement and recharge processes for supra-permafrost water. On the one hand, this is because supra-permafrost water stored in active layers is replenished by more old water compared with surface runoff. On the other hand, it is related to the longer flow path for supra-permafrost water since the active layer increases the length of water flow path (Frampton and Destouni, 2015; Ma et al., 2019b). In turn, a short MRT of the stream water indicates a relatively rapid



response of surface water to precipitation. Overall, the estimated MRT of stream and supra-permafrost water in our catchment were shorter, compared to those estimated in most previous studies from non-permafrost ~~or high latitude permafrost~~ catchments (Table 5). This may be related to whether there is groundwater recharge or not. Permafrost acts as an aquiclude, while being usually characterized by rapid hydrograph responses (Tetzlaff et al., 2018). Moreover, it cuts off the interaction and mixing between deep groundwater and surface water and supra-permafrost water. Rodgers et al (2005b) and Soulsby et al (2006a) have reported that catchment water MRT was correlated with the percentage groundwater contributions to stream water in non-permafrost catchment. Furthermore, previous studies have revealed longer MRT for deep groundwater (Table 5). For instance, deep groundwater MRT in the mountainous Brugga basin reached more than five years (Soulsby et al., 2000). Thus, the mixing or interaction between stream water and groundwater increases the stream water MRT in the non-permafrost catchments. However, this interaction is suspended by the permafrost layer in our catchment, causing the shorter MRT. This implies that the hydrological processes in high-altitude permafrost region are unique, compared with non-permafrost ~~or high-~~ latitude permafrost regions.

**Table 5. Statistics of MRT-related research results.**

Site	Altitude (m)	Water type	Model type	Tracer	Data length (years)	MRT	References
<u>Our study site<sup>a</sup></u>	<u>5100–5435</u>	<u>Stream water</u>	<u>Exponential</u>	<u><math>\delta^{18}\text{O}</math></u>	<u>8</u>	<u>100 days</u>	<u>This study</u>
<u>Huanjiang<sup>t</sup><sup>b</sup></u>	<u>272–627</u>	<u>Stream water</u>	<u>Exponential</u>	<u><math>\delta^{18}\text{O}</math></u>	<u>2</u>	<u>300 days</u>	<u>(Wang et al., 2020b)</u>
<u>Mandava<sup>b</sup></u>	<u>383</u>	<u>Stream water</u>	<u>Exponential</u>	<u><math>\delta^{18}\text{O}</math></u>	<u>2</u>	<u>444 days</u>	<u>(Sanda et al., 2017)</u>
<u>Upper Váh<sup>b</sup></u>	<u>1500</u>	<u>Stream water</u>	<u>Exponential</u>	<u><math>\delta^{18}\text{O}</math></u>	<u>4</u>	<u>390–570 days</u>	<u>(Dosa et al., 2011)</u>
<u>Dee<sup>b</sup></u>	<u>1000</u>	<u>Stream water</u>	<u>Exponential</u>	<u><math>\delta^{18}\text{O}</math></u>	<u>3</u>	<u>601 days</u>	<u>(Soulsby et al., 2010)</u>
<u>Minjiang upper<sup>b</sup></u>	<u>300–7100</u>	<u>Stream water</u>	<u>Exponential</u>	<u><math>\delta^{18}\text{O}</math></u>	<u>1</u>	<u>698 days</u>	<u>(Xia et al., 2021)</u>
<u>Our study site<sup>a</sup></u>	<u>5100–5435</u>	<u>Groundwater</u>	<u>Exponential</u>	<u><math>\delta^{18}\text{O}</math></u>	<u>5</u>	<u>255 days</u>	<u>This study</u>
<u>Himalaya<sup>a</sup></u>	<u>1600–5200</u>	<u>Groundwater</u>	<u>Exponential</u>	<u><math>\delta\text{D}</math></u>	<u>1</u>	<u>4.5 months</u>	<u>(Shah et al., 2017)</u>
<u>Vermigliana<sup>b</sup></u>	<u>1221</u>	<u>Groundwater</u>	<u>Exponential</u>	<u><math>\delta^{18}\text{O}</math></u>	<u>1</u>	<u>1.3 years</u>	<u>(Chiogna et al., 2014)</u>
<u>Allt Mharcaidh<sup>b</sup></u>	<u>300–1111</u>	<u>Groundwater</u>	<u>Exponential</u>	<u><math>\delta^{18}\text{O}</math></u>	<u>4</u>	<u>&gt;5 years</u>	<u>(Soulsby et al., 2000)</u>
<u>Huanjiang<sup>b</sup></u>	<u>272–627</u>	<u>Groundwater</u>	<u>Exponential</u>	<u><math>\delta^{18}\text{O}</math></u>	<u>2</u>	<u>161–1407 days</u>	<u>(Wang et al., 2020b)</u>

Note: “a” indicates the catchment covered by permafrost; “b” indicates the catchment not covered by permafrost. The symbol “—” indicates no data were available in the references.

As the buffer layer between the permafrost and atmosphere, the active layer is vulnerable to climate ~~change–change~~ (Xu and Wu, 2021). The increase of air temperature can alter the temperature of shallow permafrost due to strong land-atmosphere interactions. This, in turn, increases the thickness of permafrost active layer, thereby allowing soil water to move into the deeper soil layer.

In this study, the significant positive correlation between MRT and active layer thickness support previous findings showing that the MRT of permafrost catchments is highly dependent on the depth of the active layer due to the warming effects (Frampton and Destouni, 2015). From the mechanism perspective, the deepening of the active layer can increase the length of

water flow pathway and reduce transport velocities due to a shift in flow direction from horizontal saturated groundwater flow to vertical flow infiltrate into deeper subsurface, thereby increasing water MRT in permafrost catchment (Frampton and Destouni, 2015). Moreover, a previous study have reported that the potential thaw of permafrost layers due to climate change could increase MRT at the catchment scale by 20–45% (Lyon et al., 2010). A similar study in the non-frozen regions reported that MRT of forestland and shrubland water both increased with soil depth (Ma et al., 2019b).

Precipitation is an important part of the water cycle and is the main input for catchment water sources. However, precipitation in the permafrost region of TP significantly increased in recent decades (Zhao et al., 2019). In this study, Significant positive correlation between precipitation and MRT was observed, indicating that the increased precipitation or wetter climatic conditions may accelerate the water cycle process in permafrost regions. In central TP, the increase in precipitation thinned the permafrost active layer by decreasing soil heat flux, thereby cooling the soil and alleviating permafrost degradation (Zhou et al., 2021b; Luo et al., 2020). This phenomenon subsequently triggered more water to rapidly flow into the river channel in the form of surface runoff, thereby, reducing the MRT of catchment water in the end. However, a previous study suggested that much wetter climate than average probably causes higher MRT in the low-altitude temperate regions (Soulsby et al., 2006a). Our study does not resonate with these results, possibly because the aquiclude effect of permafrost reduces surface infiltration and enhances surface runoff generation (Woo, 1990; Hinzman et al., 2005; Gao et al., 2021).

Soil layer is the main source of nutrients and water required for vegetation growth (Xu et al., 2019). Water shortage in soil layer may occur when water MRT is too short, but longer MRT can hamper the water infiltration consequently to root anoxia, thus, affecting the plant growth (Ma et al., 2019b). -In this study, stream and supra-permafrost water MRT both showed significant and negative correlations with NDVI, indicating that vegetation coverage may influence water residence time in permafrost catchment. Previous studies have noted the vegetation cover was one of the most important factors, that governs the hydrological processes and thermal cycles in permafrost catchment (Wang et al., 2012b). The decline of vegetation coverage elevated soil temperature and moisture, which in turn accelerated the permafrost thawing and thickened the active layer (Wang et al., 2012c). Moreover, water MRT variability can be influenced by the development of the active layer; therefore, we believed that the effect of vegetation on MRT is also driven by the variations of thickness of permafrost active layer.

In this study, comparisons of the controlling factors of MRT between stream and supra-permafrost water indicated that supra-permafrost water is more sensitive to environmental change, compared with stream water. Previous findings have also suggested that the supra-permafrost water in the seasonally thawed layer is sensitive to climate (Cheng and Jin, 2013), and is, therefore, significantly impacted by precipitation, temperature, and vegetation. From a mechanistic perspective, climate and vegetation factors affected the MRT of stream and supra-permafrost water by modifying the active layer thickness of permafrost. Considering that MRT is a fundamental descriptor of hydrological function within catchment (Shah et al., 2017; McGuire and McDonnell, 2006). Therefore, changes in hydrological processes in permafrost watersheds can be investigated by assessing water MRT in the context of climate and environmental change.

550 **4.3 Uncertainty and limitations**

In this study, long-term stable isotopic data of stream and supra-permafrost water were used to estimate water MRT and determine the mechanism of MRT variability in a high-altitude permafrost catchment of the TP. Nonetheless, some uncertainty remains in the results of MRT estimation, including model assumptions, spatial variability of isotope input and output and isotopic fractionation.

555 Different transit time distribution (TTD) models are applicable to different watershed conditions (Małozzewski and Zuber, 1998), which may affect the assessment of residence time. The exponential model, a commonly used model for MRT estimation, describes the catchment with flow times that are exponentially distributed (Mcguire and McDonnell, 2006), which assumed that the system is in steady-state conditions and operates as a perfect mixer (Sánchez-Murillo et al., 2015; Smith, 1984; Chiogna et al., 2014). This perfect mixer indicates that the mixing between input and baseflow is rapid and complete,  
560 whereas an ideal mixing cannot exist in an aquifer, which is an important uncertainty source of the applied model (Małozzewski et al., 1983; Fenicia et al., 2010). Nevertheless, exponential model is suitable for MRT estimation in unconfined aquifers with shallow sampling points (Małozzewski and Zuber, 1998; Małozzewski et al., 1983; Stewart and McDonnell, 1991). In effect, the exponential TTD model could also approximate TTD in some non-steady cases (Haitjema, 1995; Rodhe et al., 1996). In this study area, the underlying surface was relatively uniform with less landscape heterogeneity and characterized by rapid  
565 hydrological processes. Moreover, the active layer of permafrost belonged to an unconfined aquifer and functioned as a water reservoir, thereby allowing for more precipitation recharge into the active layer to mix with old water. The amplitudes of output isotopes (stream and supra-permafrost water) were much lower than those of input (precipitation) and the dominant contribution of supra-permafrost water to stream water, both of which indicated that the precipitation was well mixed with other water within the catchment. Thus, the exponential model is suitable for application in permafrost catchment to some  
570 extent.

In general, measurement inputs represent spatial and temporal inputs for the entire catchment (Mcguire and McDonnell, 2006). At the catchment scale, elevation, air temperature, and rainfall intensity may cause considerable variation in isotopic composition of precipitation, particularly in mountainous areas (Ingraham, 1998). Thus, inputs of tracer to the catchment system are highly variable in space and time, which is an important sources of uncertainty in interpretation of catchment  
575 response (Mcguire and McDonnell, 2006; Hrachowitz et al., 2009). A previous study suggested that precipitation at high altitudes is characterized by high isotopic amplitudes (Jasechko et al., 2016), which may result in underestimation of MRT in our study area due to one sampling site for precipitation. In practice, the isotopic composition of precipitation is often sampled at one site (Mcguire and McDonnell, 2006). Considering the catchment area of our study was relatively small (2.7 km<sup>2</sup>) with an altitude drop of 300 m. The size of the selected catchment in this study was much smaller than that of most catchments  
580 previously reported (Mcguire and McDonnell, 2006). Therefore, the effects of elevation on meteorological data and precipitation isotopic variability are minor and one precipitation sampling location could represent the whole catchment to some extent. Additionally, this study only collected supra-permafrost water from one sampling point due to economic and

585 logistical constraints in the alpine regions, which is a limitation in estimating MRT. Given that the supra-permafrost water is primarily derived from precipitation, the spatial variability of isotopes in supra-permafrost water may also be minor in such small catchment. Even so, the spatial variability of isotopes in supra-permafrost water may result in underestimation of MRT in the study area.

590 The fractionation effects attributed to evaporation may potentially increase the uncertainty of water age estimation due to its impact on isotopic compositions and signals (Richardson and Kimberley, 2010; Mcdonnell et al., 2010; Song et al., 2017). Hence, the fractionation effects during the transformation from actual precipitation to effective input must be considered (Mcdonnell et al., 2010; Rusjan et al., 2019). In the study area, the atmospheric precipitation is primarily solid; the solid precipitation will be melted rapidly over a short period following deposition to form liquid water that enters soil and river channels, therefore it is difficult for snowpack to exist within this catchment. Thus, we did not collect snowpack or snow melt water as an input signal for MRT estimation. Nonetheless, solid precipitation may be subjected to evaporative fractionation to some degree when melted to be surface and subsurface runoff, thereby increasing the uncertainty of MRT estimation.  
595 Considering the rapid transformation of snow into infiltrated water and low air temperature, the potential effect of evaporation on the isotopic composition in precipitation, and consequently on MRT estimates, is relatively limited, which was not considered in the MRT estimation in this catchment.

600 To further analyse the uncertainty of MRT derived from the seasonal variability of isotope composition in hydrological component, we used the amplitude coefficient of input and output to estimate the uncertainty of MRT and found it larger for water with long residence time. Regression analysis showed that after considering the uncertainty of MRT, the  $R^2$  of the regression analysis for active layer thickness was improved, with less obvious differences for the other factors. This suggests that uncertainty of estimated MRT may affects the sensitivity of MRT to specific factors (Hu et al., 2020), indicating that the uncertainty of estimated MRT should be considered when discussing MRT influencing factors. Therefore, future research should consider the uncertainty of MRT and improve the accurate assessment of MRT in alpine catchments.

605 Overall, although there remain uncertainty and limitations for MRT estimation in our study, isotope-based MRT estimation is valuable for identifying changes in hydrological processes of the permafrost regions, where there is a lack of observational data. Thus, it is necessary to utilize more measurements in different sub-catchments to augment the data representativeness in future research.

610 Thus, it is necessary to utilize more measurements in different sub catchments to augment the representativeness the future research. Our findings will deepen the understanding of hydrological processes in high altitude permafrost catchments and provide a decision-making basis for ecological environmental protection and water resource safety and exploitation in the source of rivers on the TP

## 4.5 Conclusions

In this study, long-term observational stable isotopic data were used to estimate runoff components and water MRT in a high-  
615 altitude permafrost catchment of the TP. We found that the isotope composition in precipitation, stream and supra-permafrost  
water exhibited obvious seasonal variability. The freeze-thaw cycles of permafrost active layer and direct input of precipitation  
significantly modified the stable isotope compositions in supra-permafrost and stream water. The two-component IHS model  
indicated that the supra-permafrost water precipitation was the dominant main-contribution to the total discharge of stream  
water. We estimated that the MRT ranged from 42 to 270 days (mean 100 days) and 23 to 596 days (mean 255 days) for stream  
620 water and supra-permafrost water, respectively. Such shorter MRT of supra-permafrost and stream water (compared to the  
non-permafrost catchments) might reflect the unique characteristics of hydrological process in permafrost catchments. These  
results can be compared with the previous findings, reported in non-permafrost catchments. Namely, evaluation of seasonal  
variations of stable isotope composition can be efficiently applied to estimate water MRT in high-altitude permafrost catchment.  
Furthermore, the analysis of influencing factors revealed that the MRT of supra-permafrost water was more sensitive to  
625 environmental change than stream water. From the perspective of influencing mechanisms, climate and vegetation factors  
affected the water MRT in permafrost catchment which were mainly driven by changing the thickness of the permafrost active  
layer. It should be emphasized that the estimated MRT of supra-permafrost water is valuable for evaluating the extent of  
permafrost degradation. Under the influence of global warming, the permafrost degenerates and active layer deepens may slow  
down the rate of water cycle in permafrost regions. The findings of this study expand our understanding of the hydrological  
630 processes in high-altitude permafrost catchments under climate warming.

~~Although our study has confirmed that isotope techniques can be utilized to estimate the MRT of water in high-altitude  
permafrost catchments, there were some limitations. As the hydrological investigation at high-altitude region was very  
challenging, we only conducted long-term isotope monitoring on one sub-catchment, which may result in sampling biases.  
Thus, it is necessary to utilize more measurements in different sub-catchments to augment the representativeness of the  
635 research. Future work is required to evaluate the impact of topography and soil property on water MRT in permafrost  
catchments. To conclude, our study provided unique insights and will encourage further research in permafrost regions. Our  
findings will deepen the understanding of hydrological processes in high-altitude permafrost catchments and provide a  
decision-making basis for ecological environmental protection and water resource safety and exploitation in the source of  
rivers on the TP.~~

640 *Data availability.* We have uploaded the long-term isotope data and estimated MRT results in figshare with the URL of <https://figshare.com/articles/dataset/IsoMRTTP/19172765>.

*Author contribution.* XH and SK provided initial ideas of this work. SW and XH performed the water isotope data collection and processing and result analysis. XH provided the meteorological data of the study area. HF and XFH helped in ~~analyzing~~ analysing the results and revised the manuscript. All authors contributed to the writing.

645 *Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* This work was supported by the Joint Research Project of Three-River Headwaters National Park, Chinese Academy of Sciences and the People's Government of Qinghai Province (LHZX-2020-11), the Second Tibetan Plateau Scientific Exploration (Grant 2019QZKK0605), IWHR Research & Development Support Program (HY110145B0012021), ~~and~~ the project of State Key Laboratory of Cryospheric Science (SKLCS-ZZ-2022), and Science and technology Project of  
650 Tibet Autonomous Region (XZ202101ZD0009G):-

## References

- Carey, S. K. and Quinton, W. L.: Evaluating runoff generation during summer using hydrometric, stable isotope and hydrochemical methods in a discontinuous permafrost alpine catchment, Hydrological Processes, 19, 95-114, <https://doi.org/10.1002/hyp.5764>, 2005.
- 655 Cheng, G., Zhao L., Li, R., Wu X., Sheng, Y., Hu, G., Zou, D., Jin, H., Li, X., and Wu Q.: Characteristic, changes and impacts of permafrost on Qinghai-Tibet Plateau, Chinese Science Bulletin, 64, 2783–2795, 2019.
- Cheng, G. D and Jin, H. J.: Groundwater in the permafrost regions on the Qinghai-Tibet Plateau and it changes, Hydrogeology & Engineering Geology, 40, 1–11, <https://doi.org/10.16030/j.cnki.issn.1000-3665.2013.01.017>, 2013.
- Chiogna, G., Santoni, E., Camin, F., Tonon, A., Majone, B., Trenti, A., and Bellin, A. : Stable isotope characterization of the  
660 Vermigliana catchment, Journal of Hydrology, 509, 295-305, <https://doi.org/10.1016/j.jhydrol.2013.11.052>, 2014.
- CMA Climate Change Centre: Blue Book on Climate Change in China Beijing: Science Press, Beijing, pp. 67-68.2021.
- Dosa, M., Holko, L., and Kostka, Z.: Estimation of the mean transit times using isotopes and hydrograph recessions, Die Bodenkultur, 62, 47-52, 2011.
- Dunn, S. M., McDonnell, J. J., and Vaché, K. B.: Factors influencing the residence time of catchment waters: A virtual  
665 experiment approach, Water Resources Research, 43, 1639-1650. <https://doi.org/https://doi.org/10.1029/2006WR005393>, 2007.

- Farrick, K. K and Branfireun, B. A.: Flowpaths, source water contributions and water residence times in a Mexican tropical dry forest catchment, *Journal of Hydrology*, 529, 854–865, <https://doi.org/10.1016/j.jhydrol.2015.08.059>, 2015.
- 670 [Fenicia, F., Wrede, S., Kavetski, D., Pfister, L., Hoffmann, L., Savenije, H. H. G., and McDonnell, J. J.: Assessing the impact of mixing assumptions on the estimation of streamwater mean residence time, \*Hydrological Processes\*, 24, 1730-1741, <https://doi.org/10.1002/hyp.7595>, 2010.](#)
- [Frampton, A. and Destouni, G.: Impact of degrading permafrost on subsurface solute transport pathways and travel times, \*Water Resources Research\*, 51, 7680-7701, <https://doi.org/10.1002/2014WR016689>, 2015.](#)
- 675 Gao, H., Wang J., Yang Y., Pan X., and Duan Z.: Permafrost Hydrology of the Qinghai-Tibet Plateau: A Review of Processes and Modeling, *Frontiers in Earth Science*, 8, 1–13, <https://doi.org/10.3389/feart.2020.576838>, 2021.
- [Genereux, D.: Quantifying uncertainty in tracer-based hydrograph separations, \*Water Resources Research\*, 34, 915-919, <https://doi.org/10.1029/98WR00010>, 1998.](#)
- [Haitjema, H. M.: On the residence time distribution in idealized groundwatersheds, \*Journal of Hydrology\*, 172, 127-146, \[https://doi.org/10.1016/0022-1694\\(95\\)02732-5\]\(https://doi.org/10.1016/0022-1694\(95\)02732-5\), 1995.](#)
- 680 [Hinzman, L. D., Kane, D. L., and Woo, M. K.: Permafrost Hydrology, in: \*Encyclopedia of Hydrological Sciences\* \(eds M.G. Anderson and J.J. McDonnell\), Wiley, <https://doi.org/10.1002/0470848944.hsa178>, 2005.](#)
- Hiyama, T., Hiyama, T., Asai, K., Kolesnikov, A., Gagarin, L., and Shepelev, V.: Estimation of the residence time of permafrost groundwater in the middle of the Lena River basin, eastern Siberia. *Environmental Research Letters*, 8, <https://doi.org/10.1088/1748-9326/8/3/035040>, 2013.
- 685 [Hrachowitz, M., Soulsby, C., Tetzlaff, D., Dawson, J. J. C., and Malcolm, I. A.: Regionalization of transit time estimates in montane catchments by integrating landscape controls, \*Water Resources Research\*, 45, 207-213, <https://doi.org/10.1029/2008WR007496>, 2009.](#)
- [Hu, M., Zhang, Y., Wu, K., Shen, H., Yao, M., Dahlgren, R. A., and Chen, D.: Assessment of streamflow components and hydrologic transit times using stable isotopes of oxygen and hydrogen in waters of a subtropical watershed in eastern China, \*Journal of Hydrology\*, 589, 125363, <https://doi.org/10.1016/j.jhydrol.2020.125363>, 2020.](#)
- 690 Immerzeel, W., Immerzeel, W. W., Lutz, A. F., Andrade, M., Bahl, A., and Baillie, J.: Importance and vulnerability of the world's water towers, *Nature*, 577, 364–369, <https://doi.org/10.1038/s41586-019-1822-y>, 2020.
- [Ingraham, N. L.: Chapter 3 - Isotopic Variations in Precipitation, in: \*Isotope Tracers in Catchment Hydrology\*, edited by: Kendall, C., and McDonnell, J. J., Elsevier, Amsterdam, 87-118, <https://doi.org/10.1016/B978-0-444-81546-0.50010-0>, 1998.](#)
- 695 [Jasechko, S., Kirchner, J., Welker, J., and McDonnell, J.: Substantial proportion of global streamflow less than three months old, \*Nature Geoscience\*, 9, 126–129, <https://doi.org/10.1038/NGEO2636>, 2016.](#)

- Jin, X. Y., Jin, H. J., Iwahana, G., Smarchenko, S., Luo, D. L., Li, X.Y., and Liang, S. H.: Impacts of climate-induced permafrost degradation on vegetation: A review, *Advances in Climate Change Research*, 12, 29–47, <https://doi.org/10.1016/j.accre.2020.07.002>, 2021.
- ~~Wang, L., He, X., Fsteiner, J., Zhang, D. W., Wu, J., Wang, S. Y., and Ding, Y. J.: Models and measurements of seven years of evapotranspiration on a high elevation site on the Central Tibetan Plateau, *Journal of Mountain Science*, 17, 3039–3053, <https://doi.org/10.1007/s11629-020-6051-1>, 2020.~~
- Landerer, F. W., Dickey, J. O., and Güntner, A.: Terrestrial water budget of the Eurasian pan-Arctic from GRACE satellite measurements during 2003–2009, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2010JD014584>, 2010.
- Li, X., He X., Kang, S., Mika, Sillanp., Ding, Y., Han T., Wu, Q., and Yu, Z.: Diurnal dynamics of minor and trace elements in stream water draining Dongkemadi Glacier on the Tibetan Plateau and its environmental implications, *Journal of Hydrology*, 541, 1104–1118, <https://doi.org/10.1016/j.jhydrol.2016.08.021>, 2016.
- Li, Z., Li, Z., Qi, F., Zhang, B., and Gao, W.: Runoff dominated by supra-permafrost water in the source region of the Yangtze river using environmental isotopes, *Journal of Hydrology*, 582, <https://doi.org/10.1016/j.jhydrol.2019.124506>, 2020.
- ~~Li, Z., Li, Z., Fan, X., Wang, Y., Song, L., Gui, J., Xue, J., Zhang, B., and Gao, W.: The sources of supra-permafrost water and its hydrological effect based on stable isotopes in the third pole region, *Science of The Total Environment*, 715, 136911, <https://doi.org/10.1016/j.scitotenv.2020.136911>, 2020b.~~
- ~~Luo, D., Jin, H., Bense, V. F., Jin, X., and Li, X.: Hydrothermal processes of near-surface warm permafrost in response to strong precipitation events in the Headwater Area of the Yellow River, Tibetan Plateau, *Geoderma*, 376, 114531, <https://doi.org/10.1016/j.geoderma.2020.114531>, 2020.~~
- Lyon, S. W., Steve W. Lyon., Hjalmar Laudon., Jan Seibert., Magnus Mrth., Doerthe Tetzlaff., and Kevin H. B.: Controls on snowmelt water mean transit times in northern boreal catchments, *Hydrological Processes*, 24, <https://doi.org/10.1002/hyp.7577>, 2010.
- M., L., D., Holko, Z., and Kostka: Estimation of the mean transit times using isotopes and hydrograph recessions, *Die Bodenkultur*, 62, 47–52, 2011.
- Ma, J., Song, W., Wu, Jinkui., Liu, Z., and Wei, Z: Identifying the mean residence time of soil water for different vegetation types in a water source area of the Yuanyang Terrace, southwestern China, *Isotopes in Environmental and Health Studies*, 55, 1–18, <https://doi.org/10.1080/10256016.2019.1601090>, 2019.
- Ma, Q., Jin, H. J., Bense, V. F., Luo, D. L., and Lan, Y. C.: Impacts of degrading permafrost on streamflow in the source area of Yellow River on the Qinghai-Tibet Plateau, China, *Advances in Climate Change Research*, 10, 225–239, <https://doi.org/10.1016/j.accre.2020.02.001>, 2019.
- ~~Małoszewski, P. and Zuber, A.: A general lumped parameter model for the interpretation of tracer data and transit time calculation in hydrologic systems, *Journal of Hydrology*, 204, 297–300, [https://doi.org/10.1016/S0022-1694\(97\)00122-4](https://doi.org/10.1016/S0022-1694(97)00122-4), 1998.~~



- Małozzewski, P., Rauert, W., Stichler, W., and Herrmann, A.: Application of flow models in an alpine catchment area using tritium and deuterium data, Journal of Hydrology, 66, 319-330, [https://doi.org/10.1016/0022-1694\(83\)90193-2](https://doi.org/10.1016/0022-1694(83)90193-2), 1983.
- 735 McDonnell, J. J.; McGuire, K.; Aggarwal, P.; Beven, K. J.; Biondi, D.; Destouni, G.; Dunn, S.; James, A.; Kirchner, J.; and Kraft, P.: How old is streamwater? Open questions in catchment transit time conceptualization, modelling and analysis, Hydrological Processes, 24, 1745–1754, <https://doi.org/10.1002/hyp.7796>, 2010.
- McGuire, K. J.; DeWalle, D. R.; and Gburek, W. J.: Evaluation of mean residence time in subsurface waters using oxygen-18 fluctuations during drought conditions in the mid-Appalachians, Journal of Hydrology, 261, 132–149, 2002.
- 740 McGuire, K. J and McDonnell, J. J.: A review and evaluation of catchment transit time modeling. Journal of Hydrology, 330, 543–563, <https://doi.org/10.1016/j.jhydrol.2006.04.020>, 2006.
- Morales, K. and Oswald, C.: Water Age in Stormwater Management Ponds and Stormwater Management Pond Treated Catchments, Hydrological Processes, 34, 1854-1867, <https://doi.org/10.1002/hyp.13697>, 2020.
- 745 Martin, Š., Pavlína, S., Tomáš, V., Christina, S., Matthias, K., Jakub, J., Alena, K., and František, P.: Pre-event water contributions and streamwater residence times in different land use settings of the transboundary mesoscale Lužická Nisa catchment, Journal of Hydrology and Hydromechanics, 65, 154-164, <https://doi.org/10.1515/johh-2017-0003>, 2017.
- Ran, Y., X Li., G Cheng., Z Nan., and X Wu.: Mapping the permafrost stability on the Tibetan Plateau for 2005-2015, Science China Earth Sciences, 63, 62–79, <https://doi.org/10.1007/s11430-020-9685-3>, 2020.
- 750 Richardson, B. and Kimberley, M.: Lessons Learned from Monitoring the Effectiveness of the Asian Gypsy Moth Aerial Spraying Eradication Program, Applied Engineering in Agriculture, 26, 355-361, [10.13031/2013.29950](https://doi.org/10.13031/2013.29950), 2010.
- Rodgers, P., Soulsby, C., and Waldron, S.: Stable isotope tracers as diagnostic tools in upscaling flow path understanding and residence time estimates in a mountainous mesoscale catchment. Hydrological Processes, 19, 2291–2307, <https://doi.org/10.1002/hyp.5677>, 2005a.
- Rodgers, P., Soulsby, C., Waldron, S., and Tetzlaff, D.: Using stable isotope tracers to assess hydrological flow paths, residence times and landscape influences in a nested mesoscale catchment. Hydrology and Earth System Sciences, 9, 139–155, <https://doi.org/10.5194/hess-9-139-2005>, 2005b.
- 755 Rodhe, A., Nyberg, L., and Bishop, K.: Transit Times for Water in a Small Till Catchment from a Step Shift in the Oxygen 18 Content of the Water Input, Water Resources Research, 32, 3497-3511, <https://doi.org/10.1029/95WR01806>, 1996.
- Rusjan, S., Sapač, K., Petrič, M., Lojen, S., and Bezak, N.: Identifying the hydrological behavior of a complex karst system using stable isotopes, Journal of Hydrology, 577, 123956, <https://doi.org/10.1016/j.jhydrol.2019.123956>, 2019.
- 760 Sánchez-Murillo, R., Brooks, E., Elliot, W., and Boll, J.: Isotope hydrology and baseflow geochemistry in natural and human-altered watersheds in the Inland Pacific Northwest, USA, Isotopes in environmental and health studies, 51, 1-24, <https://doi.org/10.1080/10256016.2015.1008468>, 2015.
- 765 Sanda, M., Sedlmaierová, P., Vitvar, T., Seidler, C., Kändler, M., Jankovec, J., Kulasova, A., and Paška, F.: Pre-event water contributions and streamwater residence times in different land use settings of the transboundary mesoscale Lužická Nisa catchment, Journal of Hydrology and Hydromechanics, 65, 154-164, <https://doi.org/10.1515/johh-2017-0003>, 2017.

- Shah, R., Jeelani, G., and Noble, J.: Estimating mean residence time of karst groundwater in mountainous catchments of Western Himalaya, India. *Hydrological Sciences Journal*, 62, 1230–1242, <https://doi.org/10.1080/02626667.2017.1313420>, 2017.
- 770 Qu, S., Tao, W., Bao, W., Peng, S., Peng, J., Zhou, M., and Yu Z.: Evaluating Infiltration Mechanisms Using Breakthrough Curve and Mean Residence Time, *Water Resources Management*, 27, 4579–4590, <https://doi.org/10.1007/s11269-013-0427-8>, 2013.
- 775 [Simin, Q., Tao, W., Weimin, B., Peng, S., Peng, J., Minmin, Z., and Zhongbo, Y.: Evaluating Infiltration Mechanisms Using Breakthrough Curve and Mean Residence Time. \*Water Resources Management\*, 27, 4579-4590, 10.1007/s11269-013-0427-8, 2013.](#)
- [Sjöberg, Y., Jan, A., Painter, S. L., Coon, E. T., Carey, M. P., O'Donnell, J. A., and Koch, J. C.: Permafrost Promotes Shallow Groundwater Flow and Warmer Headwater Streams, \*Water Resources Research\*, 57, e2020WR027463, <https://doi.org/10.1029/2020WR027463>, 2021.](#)
- 780 [Smith, J.: Mixing in continuous flow systems \*Chemical Engineering Science\*, 39, 1643-1644, \[https://doi.org/10.1016/0009-2509\\(84\\)80094-9\]\(https://doi.org/10.1016/0009-2509\(84\)80094-9\), 1984.](#)
- Song, C., Wang, G., Liu, G., Mao, T., Sun, X., and Chen, X.: Stable isotope variations of precipitation and streamflow reveal the young water fraction of a permafrost watershed, *Hydrological Processes*, 31, 935–947, <https://doi.org/10.1002/hyp.11077>, 2017.
- 785 Soulsby, C., R. Malcolm., R. Helliwell., R. C. Ferrier., and A. Jenkins.: Isotope hydrology of the Allt a' Mharcaidh catchment, Cairngorms, Scotland: implications for hydrological pathways and residence times, *Hydrological Processes*, 14, 747–762, [https://doi.org/10.1002/\(SICI\)1099-1085\(200003\)14:4<747::AID-HYP970>3.0.CO;2-0](https://doi.org/10.1002/(SICI)1099-1085(200003)14:4<747::AID-HYP970>3.0.CO;2-0), 2000.
- Soulsby, C and Tetzlaff, D.: Towards simple approaches for mean residence time estimation in ungauged basins using tracers and soil distributions, *Journal of Hydrology*, 363, 60–74, <https://doi.org/10.1016/j.jhydrol.2008.10.001>, 2008.
- 790 Soulsby, C., Tetzlaff, D., and Hrachowitz, M.: Spatial distribution of transit times in montane catchments: conceptualization tools for management, *Hydrological Processes*, 24, 3283–3288, <https://doi.org/10.1002/hyp.7864>, 2010.
- Soulsby, C., Tetzlaff, D., Rodgers, P., Dunn, S., and Waldron, S.: Runoff processes, streamwater residence times and controlling landscape characteristics in a mesoscale catchment: an initial assessment, *Journal of Hydrology*, 325, 197–221, 2006.
- 795 [Stewart, M. K. and McDonnell, J. J.: Modeling Base Flow Soil Water Residence Times From Deuterium Concentrations, \*Water Resources Research\*, 27, 2681-2693, <https://doi.org/10.1029/91WR01569>, 1991.](#)
- Stockinger, M. P., Bogen, H. R., Lücke, A., Diekkrüger, B., Cornelissen, T., and Vereecken, H.: Tracer sampling frequency influences estimates of young water fraction and streamwater transit time distribution, *Journal of Hydrology*, 541, 952–964, <https://doi.org/10.1016/j.jhydrol.2016.08.007>, 2016.
- 800 Streletskiy, D., Tananaev, N., Opel, T., Shiklomanov N., Nyland, K., Streletskaya, I., Tokarev, I., and Shiklomanov, A.: Permafrost hydrology in changing climatic conditions: Seasonal variability of stable isotope composition in rivers in

discontinuous permafrost, *Environmental Research Letters*, 10, 095003, <https://doi.org/10.1088/1748-9326/10/9/095003>, 2015.

805 [Sugimoto, A., Naito, D., Yanagisawa, N., Ichyanagi, K., Kurita, N., Kubota, J., Kotake, T., Ohata, T., Maximov, T. C., and Fedorov, A. N.: Characteristics of soil moisture in permafrost observed in East Siberian taiga with stable isotopes of water, \*Hydrological Processes\*, 17, 1073-1092, <https://doi.org/10.1002/hyp.1180>, 2003.](#)

[Taylor, S., Feng, X., Williams, M., and McNamara, J.: How isotopic fractionation of snowmelt affects hydrograph separation, \*Hydrological Processes\*, 16, 3683-3690, \[10.1002/hyp.1232\]\(https://doi.org/10.1002/hyp.1232\), 2002.](#)

Tetzlaff, D., Malcolm, I. A., and Soulsby, C.: Influence of forestry, environmental change and climatic variability on the hydrology, hydrochemistry and residence times of upland catchments, *Journal of Hydrology*, 346, 93-111, 810 <https://doi.org/10.1016/j.jhydrol.2007.08.016>, 2007.

Tetzlaff, D., Piovano, T., Ala-Aho, P., Smith, A., Carey, S. K., Marsh, P., Wookey, P. A., Street, L. E., and Soulsby, C.: Using stable isotopes to estimate travel times in a data-sparse Arctic catchment: Challenges and possible solutions. *Hydrological Processes*, 32, 1936–1952. <https://doi.org/10.1002/hyp.13146>, 2018.

815 [Throckmorton, H. M., Newman, B. D., Heikoop, J. M., Perkins, G. B., Feng, X., Graham, D. E., O'Malley, D., Vesselinov, V. V., Young, J., Wullschlegel, S. D., and Wilson, C. J.: Active layer hydrology in an arctic tundra ecosystem: quantifying water sources and cycling using water stable isotopes, \*Hydrological Processes\*, 30, 4972-4986, <https://doi.org/10.1002/hyp.10883>, 2016.](#)

Uhlenbrook, S., Frey, M., Leibundgut, C., and Maloszewski, P.: Hydrograph separations in a mesoscale mountainous basin at event and seasonal timescales, *Water Resources Research*, 38, <https://doi.org/10.1029/2001WR000938>, 2002.

820 Uhlenbrook, S and Hoeg, S.: Quantifying uncertainties in tracer-based hydrograph separations: a case study for two-, three- and five-component hydrograph separations in a mountainous catchment, *Hydrological Processes*, 17, 431–453, <https://doi.org/10.1002/hyp.1134>, 2003.

[Wang, He, X., Jakob, F. S., Zhang, D., Wu, J., Wang, S., and Ding, Y.: Models and measurements of seven years of evapotranspiration on a high elevation site on the Central Tibetan Plateau, \*Journal of Mountain Science\*, 17, 3039-3053, 825 <https://doi.org/10.1007/s11629-020-6051-1>, 2020a.](#)

Wang F., Chen H., Lian J., Fu Z., and Nie Y.: Seasonal recharge of spring and stream waters in a karst catchment revealed by isotopic and hydrochemical analyses, *Journal of Hydrology*, 591, <https://doi.org/10.1016/j.jhydrol.2020.125595>, 2020.

Wang, G., Hu, H., and Li, T.: The influence of freeze-thaw cycles of active soil layer on surface runoff in a permafrost watershed, *Journal of Hydrology*, 375, 438–449, <https://doi.org/10.1016/j.jhydrol.2009.06.046>, 2009.

830 Wang, G., L, G., and Chunjie.: Effects of changes in alpine grassland vegetation cover on hillslope hydrological processes in a permafrost watershed, *Journal of Hydrology*, 444–445, 22–33. <https://doi.org/10.1016/j.jhydrol.2012.03.033>, 2012a.

Wang G., Liu G., Li C., and Yan Y.: The variability of soil thermal and hydrological dynamics with vegetation cover in a permafrost region, *Agricultural and Forest Meteorology*, 162-163, 44–57, <https://doi.org/10.1016/j.agrformet.2012.04.006>, 2012b.

- 835 Wang, G., Mao, T., Chang, J., and Liu, G.: Soil temperature-threshold based runoff generation processes in a permafrost catchment, *The Cryosphere Discussions*, 9, 5957-5978, <https://doi.org/10.5194/tcd-9-5957-2015>, 2015.
- Woo, M. K.: Consequences of Climatic Change for Hydrology in Permafrost Zones, *Journal of Cold Regions Engineering*, 4,
- Woo, M. K., Kane, D. L., Carey, S. K., and Yang, D.: Progress in permafrost hydrology in the new millennium, *Permafrost and Periglacial Processes*, 19, 237-254, [https://doi.org/10.1029/2008WR006880](https://doi.org/10.1002/ppp.613, 2008.</a></u></p><p>Wright, N., Hayashi, M., and Quinton, W. L.: Spatial and temporal variations in active layer thawing and their implication on runoff generation in peat-covered permafrost terrain, <i>Water Resources Research</i>, 45, <a href=), 2009.
- 845 Wright, N., Quinton, W. L., and Hayashi, M.: Hillslope runoff from an ice-cored peat plateau in a discontinuous permafrost basin, Northwest Territories, Canada, *Hydrological Processes*, 22, 2816–2828, <https://doi.org/10.1002/hyp.7005>, 2008.
- Xia, C., Liu, G., Zhou, J., Meng, Y., and Mei, J.: Revealing the impact of water conservancy projects and urbanization on hydrological cycle based on the distribution of hydrogen and oxygen isotopes in water, *Environmental Science and*
- 850 *Pollution Research*, 28, 1–18, <https://doi.org/10.1007/s11356-020-11647-6>, 2021.
- Xu, H., Qu, Q., Peng, L., Guo, Z., and Wulan, E.: Stocks and stoichiometry of soil organic carbon, total nitrogen, and total phosphorus after vegetation restoration in the Loess Hilly Region, China, *Forests*, 10, 27, <https://doi.org/10.3390/f10010027>, 2019.
- Xu, X and Wu, Q.: Active layer thickness variation on the qinghai-tibetan plateau: historical and projected trends. *Journal of Geophysical Research: Atmospheres*, 126, e2021JD034841, <https://doi.org/10.1029/2021JD034841>, 2021.
- 855 Yang, Y., Wu, Q., Yun, H., Jin, H., and Zhang, Z.: Evaluation of the hydrological contributions of permafrost to the thermokarst lakes on the Qinghai–Tibet Plateau using stable isotopes, *Global and Planetary Change*, 140, 1–8. <https://doi.org/10.1016/j.gloplacha.2016.03.006>, 2016.
- Yao, T., Masson-Delmotte, V., Gao, J., Yu, W., Yang, X., Risi, C., Sturm, C., Werner, M., Zhao, H., He, Y., Ren, W., Tian, L., Shi, C., and Hou, S.: A review of climatic controls on  $\delta^{18}\text{O}$  in precipitation over the Tibetan Plateau: Observations and simulations, *Reviews of Geophysics*, 51, 525–548, <https://doi.org/10.1002/rog.20023>, 2013.
- 860 Yang, Y., Wu, Q., Jin, H., Wang, Q., Huang, Y., Luo, D., Gao, S., and Jin, X.: Delineating the hydrological processes and hydraulic connectivities under permafrost degradation on Northeastern Qinghai-Tibet Plateau, China. *Journal of Hydrology*, 569, <https://doi.org/10.1016/j.jhydrol.2018.11.068>, 2018.
- 865 Zhao, L., Hu, G., Zou, D., Wu, X., and Liu, S.: Permafrost Changes and Its Effects on Hydrological Processes on Qinghai-Tibet Plateau. *Bulletin of the Chinese Academy of Sciences*, 34, 1233–1246, <https://doi.org/10.16418/j.issn.1000-3045.2019.11.006>, 2019.

Zhou J., Liu, G., Meng, Y., Xia, C. C., and Chen, Y.: Using stable isotopes as tracer to investigate hydrological condition and estimate water residence time in a plain region, Chengdu, China. Scientific Reports, 11, <https://doi.org/10.1038/s41598-021-82349-3>, 2021.

870

Zhu, X., Wu, T., Zhao, L., Yang, C., Zhang, H., Xie, C., Li, R., Wang, W., Hu, G., Ni, J., Du, Y., Yang, S., Zhang, Y., Hao, J., Yang, C., Qiao, Y., and Shi, J.: Exploring the contribution of precipitation to water within the active layer during the thawing period in the permafrost regions of central Qinghai-Tibet Plateau by stable isotopic tracing, Science of The Total Environment, 661, 630-644, <https://doi.org/10.1016/j.scitotenv.2019.01.064>, 2019.

875