

# ~~Observed~~ An evaluation of a physics-based and modeled Greenland a semi-empirical firn properties model across the Greenland Ice Sheet (1980–2020)

Megan Thompson-Munson<sup>1,2</sup>, Nander Wever<sup>1,3</sup>, C. Max Stevens<sup>4</sup>, Jan T. M. Lenaerts<sup>1</sup>, and Brooke Medley<sup>4,5</sup>

<sup>1</sup>Department of Atmospheric and Oceanic Sciences, University of Colorado Boulder, Boulder, CO, USA

<sup>2</sup>Cooperative Institute for Research in Environmental Sciences (CIRES), University of Colorado Boulder, Boulder, CO, USA

<sup>3</sup>WSL Institute for Snow and Avalanche Research SLF, Davos, Switzerland

<sup>4</sup>Earth System Science Interdisciplinary Center, University of Maryland College Park, College Park, MD, USA

<sup>5</sup>NASA Goddard Space Flight Center, Greenbelt, MD, USA

**Correspondence:** Megan Thompson-Munson (megan.thompson-munson@colorado.edu)

**Abstract.** The Greenland Ice Sheet's (GrIS) firn layer buffers the ice sheet's contribution to sea level rise by storing meltwater in its pore space. However, available pore space and meltwater retention capability is lost due to ablation of the firn layer and refreezing of meltwater as near-surface ice slabs in the firn. Understanding how firn properties respond to climate is important for constraining the GrIS's future contribution to sea level rise in a warming climate. Observations of firn density provide 5 detailed information about firn properties, but they are spatially and temporally limited. Here we use two firn models, the physics-based SNOWPACK model and the ~~semi-empirical~~ Community Firn Model (CFM) ~~configured with a semi-empirical densification equation (CFM-GSFC)~~, to quantify firn properties across the GrIS from 1980 through 2020. We use an identical forcing (MERRA-2 atmospheric reanalysis) for SNOWPACK and the ~~CFM~~ CFM-GSFC in order to isolate ~~firn~~ model differences. To evaluate the models, we compare simulated firn properties, including firn air content (FAC), to measurements from the 10 SUMup dataset of snow and firn density. Both models perform well ~~(mean absolute percentage errors of 14 % in SNOWPACK and 16 % in the CFM-GSFC)~~, though their performance is hindered by ~~meltwater percolation and~~ the spatial resolution of the atmospheric forcing. In the ~~full ice-sheet ice-sheet-wide~~ simulations, the ~~1980–1995 average~~ spatially-integrated FAC (i.e., air volume in the firn) for the upper 100 m is 34,645 km<sup>3</sup> from SNOWPACK and 28,581 km<sup>3</sup> from the ~~CFM~~ CFM-GSFC. The discrepancy in the magnitude of the modeled FAC stems from differences in densification with depth and variations in the ~~models'~~ 15 ~~treatment of atmospheric inputs~~ sensitivity of the models to atmospheric forcing. In more recent years (2005–2020), both models simulate substantial depletion of pore space. During this period, the spatially-integrated FAC across the entire GrIS decreases by ~~2.8 % and 1.2 % in SNOWPACK and the CFM, respectively. The differing magnitudes of the 2005–2020 spatially-integrated FAC of 3.2 %~~ (-66.6 km<sup>3</sup> yr<sup>-1</sup>) in SNOWPACK and ~~1.5 %~~ (-17.4 km<sup>3</sup> yr<sup>-1</sup> in the CFM) in the CFM-GSFC. These differing magnitudes demonstrate how model differences propagate throughout the FAC record. Over the full modeled record (1980– 20), SNOWPACK simulates a loss of pore space equivalent to 3 mm of sea level rise buffering, while the ~~CFM~~ CFM-GSFC simulates a loss of 1 mm. The greatest depletion in FAC is along the margins, and especially along the western margin where observations and models show the formation of near-surface, low-permeability ice slabs that ~~may~~ inhibit meltwater storage.

## 1 Introduction

Most of the Greenland Ice Sheet (GrIS) is covered ~~in-by~~ a thick, porous layer of partially compacted snow known as firn.

25 The density of firn varies ~~with depth and~~ across the ice sheet and is sensitive to surface mass balance (SMB) processes like accumulation and melt, which cause firn density to also vary ~~in-time and~~ on several different temporal scales (e.g., daily, seasonal, annual). Since firn is porous, it is capable of storing meltwater in its pore space in a solid or liquid form, which can buffer ~~the contribution of increased melt rates to~~ sea-level rise (Harper et al., 2012). ~~The mechanisms for meltwater entering into and remaining stored in firn are complex and varied.~~ Meltwater can remain in a liquid form and remain near the surface in weathered ice crusts (Cooper et al., 2018), pool into subsurface lakes (Dunmire et al., 2021), or percolate into the snowpack and remain in a liquid form in firn aquifers (e.g., Forster et al., 2014). Additionally, meltwater can percolate into the snowpack where it refreezes deeper in the firn layer or is stored in the firn's pore space (Pfeffer et al., 1991; Harper et al., 2012). Refreezing of meltwater within the firn frequently occurs and creates ice lenses (<0.1 m thick) and layers (0.1-1 m thick) that can accumulate into low-permeability ice slabs (>1 m thick) (MacFerrin et al., 2022). These ice slabs make 30 deeper pore space ~~potentially~~ inaccessible to meltwater produced at the surface, which reduces the buffering capacity of the firn (Machguth et al., 2016) and increases the elevation below which meltwater runs off (Tedstone and Machguth, 2022). Rapid depletion of the pore space on the GrIS is expected to accelerate mass loss in the 21st century and increase the ice sheet's contribution to sea level rise (van Angelen et al., 2013). Approximately half of GrIS mass loss is due to an increase in meltwater runoff ~~(van den Broeke et al., 2009; ?)~~~~(van den Broeke et al., 2009; Enderlin et al., 2014)~~, making the firn's buffering capacity 35 increasingly important to the GrIS contribution to sea-level rise.

40 In recent decades, Greenland's firn layer has begun to show evidence of climate change. The 2012 extreme melt season produced expansive ice slabs that persisted for several years and reduced permeability (Culberg et al., 2021). At lower elevations, where significant melt occurs, ~~firn has lost its capacity to store meltwater~~~~the meltwater storage capacity of firn has abruptly decreased~~ (Machguth et al., 2016). In higher-elevation areas where less meltwater is generated, the firn structure has 45 still changed through enhanced densification from warmer temperatures and the presence of liquid water (Machguth et al., 2016), which causes ice-sheet surface height lowering (de la Peña et al., 2015). Changes in the amount of air-filled pore space within the firn, known as the firn air content (FAC), have been investigated in both observations (e.g., Vandeverux et al., 2019) 50 (e.g., Vandecrux et al., 2019; Benson, 1996; Braithwaite et al., 1994; Sørensen et al., 2011; Kuipers Munneke et al., 2015) and models (e.g., Medley et al., 2022). Although regeneration of this pore space is slow (Harper et al., 2012), consecutive years of average or below-average melt have shown to temporarily pause FAC depletion (Rennermalm et al., 2021). These complex interactions between melt and pore space depletion, including ~~the memory effect of changes to the firn from previous years~~~~firn evolving on multi-year timescales~~, motivate the use of detailed firn models to capture these processes and to enhance our 55 understanding of the changes occurring in the firn.

Modeling firn has become important for estimating mass balance (MB) from satellite altimetry, since this method relies on 55 firn models to interpret the causes of surface height changes (e.g., Li and Zwally, 2011) (e.g., Li and Zwally, 2011; Arthern and Wingham, 2011). Changes in surface height cannot be attributed to ice mass or firn density changes without additional information from firn

models (Smith et al., 2022). Accurate estimates of ice sheet firn density over time and space are necessary to constrain the uncertainty in MB assessments relying on surface height changes measured ~~in from~~ satellite altimetry. Additionally, understanding the limits and deficiencies in firn models is essential for quantifying uncertainties in altimetry-based MB estimates  
60 [\(Morris and Wingham, 2014; Verjans et al., 2021\)](#).

Firn models can also be used to fill in gaps in firn density observations and study how firn properties vary on larger spatial and temporal scales, especially since in-situ and remotely sensed observations can only provide snapshots of firn properties in space and time. Semi-empirical firn models have been used to simulate firn properties on both Greenland (e.g., Medley et al., 2022) and Antarctica (e.g., Ligtenberg et al., 2011). These models use empirical relationships between densification, accumulation, 65 and temperature, and they are often tuned to observations [\(e.g., Ligtenberg et al., 2011; Medley et al., 2022; Li and Zwally, 2011\)](#) [\(e.g., Ligtenberg et al., 2011; Medley et al., 2022; Li and Zwally, 2011; Herron and Langway, 1980; Arthern et al., 2010; Simonsen et al., 2011\)](#). Semi-empirical models are beneficial because they ~~do not rely on the physics of firn densification, which are not fully understood~~ [can simulate more accurate depth-density profiles by calibration, which removes the uncertainties introduced by poorly understood densification processes in firn](#). On the other hand, the observations these models rely on for calibration 70 may not be representative of future firn properties in a warming climate (Ligtenberg et al., 2018). Additionally, these models are often tuned using depth-density profiles, which requires a steady-state assumption and thereby lends uncertainty in their ability to simulate firn changes in a transient climate (Lundin et al., 2017). The alternatives to semi-empirical models are physics-based models that use the ~~eonstitutive relationship~~ [material properties of snow and firn to simulate densification based constitutive relations](#) between stress and strain ~~for snow to simulate densification~~ (e.g., Vionnet et al., 2012; Bartelt 75 and Lehning, 2002; Lehning et al., 2002b, a). While physics-based models do not rely on observations ~~from~~ [for](#) tuning, they generally need more detailed meteorological forcing data and are limited by uncertainties in [the representation of](#) physical processes. ~~Physics-based models are typically developed to simulate seasonal snow processes rather than firn processes since snow physics have been more thoroughly studied. Still, both~~ [The wealth of snow physics studies allowed for the development of more complex, physics-based, seasonal snow models like SNOWPACK. Both](#) semi-empirical as well as physics-based 80 firn models have been [successfully](#) used in Greenland [\(e.g., Vandecrux et al., 2020a; Dunmire et al., 2020; Medley et al., 2022\)](#) [\(e.g., Vandecrux et al., 2020a; Dunmire et al., 2020; Medley et al., 2022; Sørensen et al., 2011; Kuipers Munneke et al., 2015\)](#).

The ~~semi-empirical~~ Community Firn Model (CFM) and ~~physics-based~~ SNOWPACK firn model have seen significant development ~~for in~~ polar regions in recent years [\(e.g., Medley et al., 2022; Keenan et al., 2021; Stevens et al., 2020\)](#). [In SNOWPACK, there have been modifications to the settling and microstructure schemes \(Groot Zwaaftink et al., 2013; Steger et al., 2017\), inclusion of drifting snow impacts on near surface density \(Groot Zwaaftink et al., 2013; Keenan et al., 2021; Wever et al., 2022\), and optimizations for computational efficiency by improving the layer merging scheme \(Steger et al., 2017\). The CFM has recently been used over both ice sheets using the GSFC-FDMv1.2.1 densification equation \(Medley et al., 2022\), which we will subsequently refer to as CFM-GSFC.](#) At an ice-sheet scale, few comparisons of semi-empirical and physics-based models exist [\(e.g. Steger et al., 2017\)](#). In many studies investigating firn models, only a single atmospheric forcing is used with a single 85 model (e.g., Medley et al., 2022). Reported results are often not directly comparable between studies since statistics are calculated over different spatial extents or time periods. This approach makes it difficult to attribute differences between modeled 90

and observed firn properties to model forcing (i.e., boundary conditions) or the model itself (i.e., the representation of physical processes by the model).

Here we use the ~~CFM~~<sup>CFM-GSFC</sup> and SNOWPACK to simulate firn properties on the GrIS. Importantly, we use the same atmospheric forcing for both models in order to identify model differences that are ~~completely~~ independent from the forcing data. We compare the simulated firn properties to point observations and then extend the model domain to the entire ice sheet and simulate evolving firn properties across Greenland from 1980 through 2020. In Section 2, we describe the methods used and include descriptions of the firn models, atmospheric forcing, and observational dataset. Section 3 reports the results, which we partition into the model evaluation (Section 3.1), a description of firn properties in a steady state climate (Section 3.2), and a description of firn properties in a changing climate (Section 3.3).

## 2 Methods and data

### 2.1 MERRA-2 atmospheric forcing

The Modern-Era Retrospective Analysis for Research and Applications, version 2 (MERRA-2) is a global atmospheric reanalysis spanning the period from 1980 to present day (Gelaro et al., 2017). We use MERRA-2's hourly 2-m air temperature, relative humidity, 10-m wind speed, incoming shortwave radiation, incoming longwave radiation, and precipitation rate for the 1980–2020 period (Global Modeling and Assimilation Office (GMAO), 2015a, b, c, d) to force both firn models. MERRA-2 is a gridded product with a horizontal resolution of  $0.5^{\circ}$  latitude by  $0.625^{\circ}$  longitude. Due to convergence of meridians toward the poles, the MERRA-2 grid becomes more tightly spaced at higher latitudes. To account for this, we weight model grid ~~points~~<sup>cells</sup> by the cosine of the latitude when calculating ice-sheet-wide or basin-wide averages of firn and atmospheric properties. We choose to use MERRA-2 since it is publicly available, regularly updated and released, and spans a temporal window that captures recent climate change (Fig. A2). ~~Regional climate models are not always widely available or regularly updated, and no single reanalysis clearly outperforms others over the GrIS (Zhang et al., 2021). Additionally, our focus is comparing two firn models with the same forcing in order to isolate model differences. As such, the choice of the A different reanalysis product or regional climate model could also be used here, though the exact choice of forcing dataset is less important for the interpretation of our results relevant for the firn model intercomparison since we aim to compare the output from the two firn models forced with identical input.~~

In order to evaluate and compare the firn models, we first run them using forcing data from only the grid cells nearest to firn density observations. We use 177 MERRA-2 grid cells for this first step. For the ~~full ice-sheet~~<sup>ice-sheet-wide</sup> simulations, we mask out ~~grid~~<sup>grid</sup> cells with an ice coverage of less than 50 % ~~in the MERRA-2 ice cover map~~, which leaves us with a model domain of 1784 MERRA-2 grid cells. The total area of the model domain is 1.81 million km<sup>2</sup>, which is greater than the actual ice-sheet area of 1.71 million km<sup>2</sup> (Rignot and Mouginot, 2012).

## 2.2 SNOWPACK firn model

SNOWPACK is a single-column, physics-based, multi-layer snow and firn model originally designed for avalanche warning in alpine environments (Bartelt and Lehning, 2002; Lehning et al., 2002b, a). In recent years, this model has been ~~successfully~~

125 applied over the GrIS (e.g., Van Tricht et al., 2016; Steger et al., 2017; Izeboud et al., 2020; Dunmire et al., 2021), as well as the Antarctic Ice Sheet (Dunmire et al., 2020; Keenan et al., 2021). It uses a Lagrangian framework and adds layers when snowfall occurs with a new-snow density based on atmospheric conditions (e.g., Keenan et al., 2021), including explicit treatment of the wind compaction under drifting and blowing snow conditions (Wever et al., 2022). Firn densification is calculated using ~~the a~~ constitutive relationship between stress and strain in snow (Bartelt and Lehning, 2002; Lehning et al., 2002b). For 130 calculating firn temperatures, the upper boundary condition is determined from a surface energy balance scheme. The model describes snow microstructure based on four parameters: grain radius, bond radius, sphericity, and dendricity. These evolve over time primarily based on temperature, temperature gradients, and liquid water content ~~of the snow layers~~ (Lehning et al., 2002b). SNOWPACK uses the MeteoIO library (Bavay and Egger, 2014) for preparing the meteorological forcing data for 135 the simulations. ~~The library reads the meteorological forcing from the MERRA-2 grids and provides data to SNOWPACK for each grid cell, at each of the SNOWPACK time steps. SNOWPACK is run at smaller times steps than MERRA-2 data is available, and nearest neighbor interpolation (for wind speed) and linear interpolations (for all other variables) are used to provide meteorological forcing at higher frequency than provided by MERRA-2.~~

We run SNOWPACK with half-hourly time steps. To conserve computational expenses and ~~keep reduce~~ model output sizes ~~manageable~~, detailed model output for all firn layers is stored with 7-day temporal resolution. SNOWPACK's simulated 140 firn layers have variable thicknesses, impacted by the layer-merging scheme ~~use used~~ to reduce computational cost. ~~However, As described in more detail in Steger et al. (2017), depending on depth below the surface and similarity of snow properties in adjacent layers, the layers may be merged to reduce computational costs. If those merged layers come closer to the surface, they can be split again to maintain sufficient spatial resolution to capture the steep gradients near the surface. The higher vertical resolution of a few centimeters per layer is ~~maintained required~~ in the near-surface layers where the firn is more sensitive to 145 short-term atmospheric fluctuations. We set the surface roughness to 0.002 m for ~~calculating turbulent energy fluxes when~~ solving the energy balance ~~with~~. Here, we account for atmospheric stability using the Michlmayr et al. (2008) stability correction when a stable boundary layer is diagnosed. ~~For unstable boundaries, which happen rather infrequently (Schlögl et al., 2017), Eq. 8 in Stearns and Weidner (1993) is used. We apply a bucket scheme to represent vertical water percolation in SNOWPACK.~~~~

150 Since the thickness of the firn layer varies ~~across the GrIS~~ and we aim to simulate processes for the full firn column depth, we spin up the model to build a firn layer until a thickness of at least 150 m is reached, or when the bottom 3 m of the simulated firn consist of solid ice with a total thickness of at least 10 m (whichever condition is reached first). To perform the spin-up, we run the model using forcing data from a reference climate interval (RCI) and repeat the model runs until the desired thickness is reached. Once reached, we perform a final model run using the full-length record (1980–2020). We choose an RCI of 1 January

155 1980 through 31 December 1995 (Fig. A2), which is the same period used in Medley et al. (2022). We make the assumption  
that this period is representative of the longer-term Greenland climate.

### 2.3 The Community Firn Model (CFM)

The Community Firn Model (CFM, Stevens et al., 2020) is an open-source model framework that simulates ~~the evolution of  
firn. It is designed to be modular, allowing the model user to choose which physical processes in firn. Its modularity~~  
160 ~~allows users to choose which~~ processes to simulate and which parameterizations to use (e.g., thermal conductivity, densification  
rate) in a given model run. ~~The core CFM modules simulate firn density and temperature evolution as well as meltwater  
refreezing and runoff. Additionally, the user can easily choose different parameterizations to use. As the CFM provides a  
high degree of flexibility, it is important to specific how the CFM has been configured~~ for a particular ~~model run, including the  
firn densification equation, thermal conductivity, surface density, and water percolation scheme.~~ run. Any pertinent parameterizations  
165 used for the model runs in this paper can be found in Medley et al. (2022). As previously noted, in this paper we refer to our  
particular CFM configuration as “CFM-GSFC” to highlight that we are using the CFM with the semi-empirical GSFC-FDMv1.2.1  
firn densification equation. This equation is based on the firn densification equation proposed by Arthern et al. (2010), but  
optimized using firn-core data from 226 sites across the Antarctic and Greenland Ice Sheets.

Like SNOWPACK, the ~~CFM~~ CFM-GSFC uses a Lagrangian numerical framework; each accumulation event adds a new  
170 layer to the grid and one is removed from the bottom. The ~~CFM~~ CFM-GSFC uses a layer-merging scheme at 5- and 10-m  
depth to reduce computational demands. ~~The CFM-GSFC is coded so that each model time step adds a new layer. As such,  
daily time stepping generates many thin layers. To reduce computational demands, we use the CFM-GSFC’s layer merging  
scheme. For this study’s simulations, we merge 30 of the high-resolution (daily) layers at 5-m depth into mid-resolution  
(approximately monthly) layers. At 10-m depth, 12 layers are merged into coarser (approximately annual) layers.~~ Model  
175 outputs are interpolated onto a 0.25-m regular grid to reduce output file size.

For the present work, we run the ~~CFM~~ CFM-GSFC over the GrIS at daily resolution. The ~~CFM~~ CFM-GSFC’s required  
forcing inputs for these model runs are surface temperature, precipitation (rain and snow), sublimation, and melt. However,  
MERRA-2 does not explicitly provide a melt flux. As such, we force the ~~CFM~~ CFM-GSFC with outputs from SNOWPACK’s  
180 surface energy balance scheme, including skin temperature, melt flux, and sublimation. This method ensures that the surface  
boundary conditions (i.e., mass and energy fluxes) for the ~~CFM~~ CFM-GSFC and SNOWPACK model runs are consistent.

~~The CFM is modular and offers the user several parameterization choices, but in this work we select a single configuration for  
all model runs. We use a constant surface density of  $350 \text{ kg m}^{-3}$  for the ~~CFM~~ CFM-GSFC runs, which is a reasonable estimate  
when compared to observed and SNOWPACK-modeled surface density (Fig. A1). The densification rate is determined with the  
NASA GSFC-FDMv1 firn densification equation (Medley et al., 2022). This equation is based on the firn densification equation  
proposed by Arthern et al. (2010), but optimized using firn-core data from 226 sites across the Antarctic and Greenland Ice  
Sheets. We use a bucket scheme with an enthalpy-based heat flow module to handle meltwater percolation and refreezing.~~

The simulations for each grid cell are initialized with a depth/density profile predicted by the Herron and Langway (1980)  
steady state model. ~~The depth of the model domain varied for each grid cell but was chosen to be near the depth at which the~~

190 ~~firm reaches the ice density. We design the CFM-CFM-GSFC spin-up to repeat the RCI of 1980–1995 until the entire initial firm column is refreshed. For example, if the firm needs 1000 years to spin up, the RCI would repeat 63 times. Once the spin-up is completed, the main model run (1980–2020) commences.~~

## 2.4 SUMup observations

The Surface Mass Balance and Snow on Sea Ice Working Group (SUMup) dataset is a compilation of Arctic and Antarctic observations of SMB components and includes in-situ observations of firn density on both ice sheets (Montgomery et al., 195 2018). The 2022 SUMup release contains data from 845 locations in Greenland (Thompson-Munson et al., 2022) that we use to compare with modeled firn properties. The measurements have been taken over the past several decades and the depths of the cores range from 0.03 m to 334.53 m. Of the 845 measurements, 78 are single point measurements of surface density only. For calculating depth-integrated properties, we only use the 767 observations that contain data points from at least two different depths.

## 200 2.5 Firm air content (FAC) calculation

To compare the observations with the model results, we select the modeled density profile from the closest location and date to when the measurement was taken. The CFM-CFM-GSFC produces daily output, which means the observed and model date are the same. SNOWPACK output for the model evaluation is weekly, which means that the date of the modeled profile can differ from the date of the observed profile by as much as 3 days.

205 From the observed and modeled density profiles, we calculate the FAC(~~also known as “depth-integrated porosity (DIP)”,~~ which has units of meters. The FAC is a quantification of the air-filled pore space in the firn. For any depth interval from  $z_j$  (upper bound) to  $z_k$  (lower bound) where  $z = 0$  m represents the surface ~~and is increasing downwards~~, the FAC is calculated as

$$FAC(z_k - z_j) = \int_{z_j}^{z_k} \frac{\rho_{ice} - \rho(z)}{\rho_{ice}} dz \quad (1)$$

210 where  $\rho_{ice}$  is the density of ice ( $917 \text{ kg m}^{-3}$ ) and  $\rho(z)$  is the layer density at a given depth.

To evaluate the firm models and ensure a direct comparison with the observations, we first calculate the FAC only over the depths represented by the observations (Section 3.1). For these comparisons,  $z_j$  is the uppermost observation depth and  $z_k$  is the lowermost observation depth. When examining ice-sheet-wide firn properties (Sections 3.2 and 3.3), we calculate FAC from the surface to a depth of 100 m ( $z_j = 0$  m,  $z_k = 100$  m), and for 10-m-thick vertical intervals from 0 m down to 50 m ( $z_j = 0, 10, 20, 30, 40$  m;  $z_k = 10, 20, 30, 40, 50$  m).

## 215 2.6 Metrics of model evaluation

We use several metrics to compare the modeled density and FAC to the observations in order to evaluate model performance. The Nash-Sutcliffe efficiency (NSE) coefficient is a measure of a model’s goodness of fit and is typically employed in hydro-

logical modeling (NSE; Nash and Sutcliffe, 1970). A value of 1 indicates perfect model performance, whereas a value of 0

220 indicates that the model's predictive ability is the same as using the observations' means. It is calculated as

$$NSE = 1 - \frac{\sum_{i=1}^n (O_i - M_i)^2}{\sum_{i=1}^n (O_i - \bar{O})^2} \quad (2)$$

where  $O_i$  is the observation value,  $M_i$  is the model value,  $\bar{O}$  is the observation mean, and  $i$  iterates over the  $n$  is the number of values. For quantifying the accuracy of the models, we use the mean absolute percentage error (MAPE), which is calculated as

$$225 \quad MAPE = \frac{1}{n} \sum_{i=1}^n \left| \frac{O_i - M_i}{O_i} \right| \times 100\% \quad (3)$$

Finally, we use the relative bias to understand the fractional degree of under- or over-estimation of the models. At each observation location, we calculate relative bias as

$$Relative\ bias_i = \frac{M_i - O_i}{O_i} \times 100\% \quad (4)$$

and for the full set of locations, we calculate a single, bulk bias value as

$$230 \quad Bulk\ relative\ bias = \frac{\sum_{i=1}^n (M_i - O_i)}{\sum_{i=1}^n (O_i)} \times 100\% \quad (5)$$

### 3 Results

#### 3.1 Model evaluation

Since we use two models in this study, we first discuss the shared and distinct features in simulated density profiles for two

example observations from the SUMup database (Wilhelms, 2000; Machguth et al., 2016). Figure 1a compares a 16.3-m-deep

235 core with several ice layers collected in southwest Greenland on 12 May 2013 (gray, Machguth et al., 2016) to model results

from SNOWPACK (blue) and the CFM-CFM-GSFC (green). In this example, both SNOWPACK and the CFM-CFM-GSFC

simulate high-density layers  $\sim 1$  m below the surface that formed as a result of high melt in 2012. The observed profile

also reaches high densities ( $>700$  kg m $^{-3}$ ) near the surface. However, neither model captures the even higher-density observed

ice layer ( $>800$  kg m $^{-3}$ ) at  $\sim 5$  m depth. The models show higher agreement with one another in near the surface and begin

240 to diverge with depth. SNOWPACK simulates more variability between layers compared to the CFM-CFM-GSFC. This partly

results from the fixed surface density of 350 kg m $^{-3}$  set for the CFM-CFM-GSFC, while the surface density in SNOWPACK

varies based on atmospheric conditions, and partially because the CFM-CFM-GSFC outputs are interpolated onto a grid.

For polar regions in particular, temporal variations in wind and the presence of drifting snow translate into vertical density variations with increasing accumulation. Figure 1b compares a 102.4-m-deep core from the high-elevation interior measured

245 with gamma-ray attenuation (Wilhelms, 2000) with outputs from both models. The modeled profile shapes generally match the

observed profile with an NSE of 0.96 for both SNOWPACK and the CFM-CFM-GSFC (Fig. 1b). SNOWPACK's inter-layer


variability is present in the upper  $\sim$ 25 m and matches the degree of variability seen in the observations. SNOWPACK's density variability stems from the effect of microstructure on the settling, which disappears at depth due to the model's layer merging algorithm.

250 Since most observations are from shallow cores (median depth = 2.0 m; Fig. 2a) the observed FAC values are relatively low (median FAC = 1.3 m; Fig. 2b) and do not represent the FAC of the full firn column. ~~Generally, modeled FAC~~ In these shallow cores where densification has little impact on FAC, the model performance is a reflection of the models' representations of the surface density. In SNOWPACK, the surface density is modeled from the atmospheric input, while in the CFM-GSFC the surface density is fixed at  $350 \text{ kg m}^{-3}$ . We compare observed and modeled FAC for all 767 points (Fig. 3a, b), but we 255 also partition the dataset into bins based on core depth (Fig. 3c-j) to evaluate model performance in terms of both the surface density parameters (shallow cores) and the densification schemes (deeper cores). We use the following core depth thresholds for binning the data: 0 to 1 m ( $n = 253$ ), 1 to 2 m ( $n = 112$ ), 2 to 10 m ( $n = 242$ ), and  $>10$  m ( $n = 160$ ).

260 In the comparison of all 767 points, modeled FAC from both SNOWPACK and the ~~CFM~~ CFM-GSFC agree very well with the calculated FAC from the observed density profiles (Fig. 3a, b). The NSE is 0.90 for SNOWPACK and 0.94 for the ~~CFM~~ CFM-GSFC, and the MAPE is 14 % for SNOWPACK and 16 % for ~~CFM~~ CFM-GSFC (Fig. 3a, b). The bulk relative biases of +7.9 % for SNOWPACK and +0.2 % for the ~~CFM~~ CFM-GSFC show that both models are overestimating FAC on average (Fig. 2c, d), though this overestimation is not statistically significant ( $p = 0.13$  for SNOWPACK,  $p = 0.49$  for the ~~CFM~~ CFM-GSFC). For both models, the relative bias calculated at each observation site is less than 10 % for the majority of points (69 % of SNOWPACK points and 64 % of the ~~CFM~~ CFM-GSFC points). Only 2 % of points in SNOWPACK and 1 % of points in the ~~CFM~~ CFM-GSFC have relative biases greater than 100 %. Many of the large biases occur along the margins of the ice sheet, particularly in the southeast and southwest (Fig. 2c, d).

265 When the evaluation is performed for the four bins of core depths, we find varying performance in each bin (Fig. 3c-j). SNOWPACK performs best in the shallowest cores (NSE = 0.84, MAPE = 9 %) where the FAC is tightly coupled to the surface density scheme (Fig. 3c). As densification becomes more important with depth, the SNOWPACK model performance decreases, as expressed by both the NSE and MAPE (Fig. 3d-f). For the CFM-GSFC, the FAC in the shallower bins (Fig. 3g, h) is impacted by the fixed surface density and vertical interpolation that together prevent the fine resolution necessary for comparisons with observations. At depth, the CFM-GSFC generally performs well and has NSE and MAPE values comparable to SNOWPACK (Fig. 3i, j).

270 For both models, five of the highest absolute biases occur in five cores located in the same MERRA-2 grid cell (Fig. 4). The ~~This~~ grid cell is located in an area with observed firm aquifers (Miller et al., 2018), and it contains large topographic gradients. Here, the models use the same MERRA-2 climatology to simulate firn properties for five observations. Both SNOWPACK and the ~~CFM~~ CFM-GSFC overestimate FAC; absolute biases exceed 5 m and relative biases exceed 100 % in all but one of these cases (Fig. 3, 4). Within this grid cell, the observations substantially differ from the models, especially as the distance between the center of the MERRA-2 grid cell and the observation location increases. The biases are highest furthest from the grid cell center and where the topography is steepest (Fig. 4).




**Figure 1.** Modeled and observed density profiles at two example locations from (a) Machguth et al. (2016) and (b) Wilhelms (2000). The locations of the observations are shown as stars on the map and the profile dates, latitudes, longitudes, and elevations for the observations and models are reported in the legends. Observations are shown in gray, SNOWPACK results are in blue, and the [CFM-CFM-GSFC](#) results are in green.

### 3.2 Firm properties in a steady-state climate

In this section, we compare the modeled results over the entire GrIS during the RCI. The RCI used for model spin-up spans 1980 through 1995, and since we assume that this period represents a relatively steady-state, long-term Greenland climate (Fig. A2), the modeled firm is considered to be in steady state as well. This RCI has been used previously by Medley et al.

285 Since the forcing data from SNOWPACK and the [CFM-CFM-GSFC](#) are identical, differences in the simulated firm properties can be directly attributed to the differences between models. We calculate FAC for the upper 100 m of the firm column in both SNOWPACK and the [CFM-CFM-GSFC](#) to reduce any possible biases arising from inconsistent maximum depths between the two models. Unless otherwise stated, all FAC values reported in the remainder of the manuscript have been calculated for the upper 100 m. [Additionally, when calculating spatial means in FAC, we weight the values to account for the converging meridians that lead to variable sizes of MERRA-2 grid cells.](#)




**Figure 2.** (a) Locations of the 767 SUMup cores used for analysis with shading indicating the depth of the core. (b) Firn air content (FAC) calculated from the observed densities in SUMup. (c) Relative bias between SNOWPACK-modeled FAC and observed FAC. (d) Relative bias between the ~~CFM-modeled~~ CFM-GSFC-modeled FAC and observed FAC. Positive bias indicates an overestimation of FAC by the model, while negative bias indicates an underestimation of FAC by the model.


We use the liquid-to-solid ratio (LTSR) averaged over the RCI to investigate how both models respond to liquid water input from snowmelt and rain, and solid input from accumulation in a relatively steady state climate. The LTSR is calculated as

$$LTSR = \frac{melt + rain}{snow} \quad (6)$$

and is the same for SNOWPACK and the ~~CFM~~ CFM-GSFC since they use the same forcing data. We examine the extent 295 to which LTSR predicts FAC, and find a higher  $r^2$  value for the CFM-GSFC ( $r^2 = 0.89$ ) compared to SNOWPACK ( $r^2 = 0.77$ ). Overall, the range of SNOWPACK FAC is greater than the range of the ~~CFM~~ CFM-GSFC FAC (Fig. 5a). At very low LTSR values (<0.10) where snowfall is the dominant component of surface mass fluxes, FAC in SNOWPACK ranges from 9.7 to 32.1 m, whereas FAC in the ~~CFM~~ CFM-GSFC ranges from 9.9 to only 23.6 m. As the LTSR increases and liquid input dominates, the FAC in both models approaches zero. However, SNOWPACK FAC decreases more rapidly and reaches 300 lower FAC values than the ~~CFM~~ CFM-GSFC at high LTSR values. The ~~CFM~~ CFM-GSFC FAC decreases more gradually with increasing LTSR. In SNOWPACK, 127 points have ~~an~~ a FAC less than 1 m, and in the ~~CFM~~ CFM-GSFC only 2 points are as low (Fig. 5a). We also examine how FAC responds to the summer (June, July, and August) 2-m air temperature during the RCI (Fig. 5b). For summer temperatures below  $\sim -4^\circ\text{C}$ , both models consistently simulate the FAC values exceeding 10 m. For these low temperatures, the range of FAC is greater in SNOWPACK (9.9 to 32.1 m) compared to the ~~CFM~~ CFM-GSFC (10.7



**Figure 3.** Observed versus modeled firn air content (FAC) for all core depths for (a) SNOWPACK and (b) the CFMCFM-GSFC. The smaller panels Panels c-j show the same comparison but for the four bins of observations partitioned by core depth for (c-f) SNOWPACK and (g-j) the CFM-GSFC. The core depth bins are shown in **bold** above each panel. The number of points (n), the Nash-Sutcliffe Efficiency (NSE), and the mean absolute percentage error (MAPE) are reported for each model in the lower right. The gray dashed line is a 1:1 line, and the black solid line is the linear regression. Points with biases greater than 5 m are circled in red and correspond to the density profiles shown in Figure 4.



**Figure 4.** Density profiles and locations of high model biases from Fig. 3. (a) Map showing the five SUMup cores plotted in (c-g), the MERRA-2 domain used for SNOWPACK and the [CFM](#)-[CFM-GSFC](#) simulations shown in (c-g), and 100-m surface elevation contours from the BedMachine dataset (Morlighem et al., 2017). (b) Location map of Greenland with the black plus sign indicating the region shown in (a). Maps in panels (a) and (b) were created with QGreenland v2.0.0 (Moon et al., 2022). (c-g) Observed and modeled density profiles for five locations in southeast Greenland where both SNOWPACK and the [CFM](#)-[CFM-GSFC](#) overestimate firn air content by at least 5 m. The modeled profiles come from the same simulation for the MERRA-2 grid [point cell](#) closest to the firn cores, but differ in date and time to match the observation date as closely as possible. In each panel, we report the FAC absolute bias (modeled minus observed) and the relative bias (in parentheses) for SNOWPACK (“SP”) and the [CFM-GSFC](#) (“CFM”).

305 to 23.5 m). Both models show a rapid decline in FAC as summer temperatures exceed -4°C, and the modeled response FAC is similar at these warmer temperatures.

During the RCI, SNOWPACK and the CFM produce generally similar spatial patterns in FAC across the ice sheet, though the magnitudes differ (Fig. 6), with higher FAC (>10 m) in the ice-sheet interior and lower FAC (<10 m) in the margins (Fig. 6). SNOWPACK simulates an However, on average, FAC is greater in SNOWPACK than in the CFM-GSFC. SNOWPACK simulates a FAC of  $19.1 \pm 8.0$  m (mean  $\pm$  standard deviation) and the CFM simulates an CFM-GSFC simulates a FAC of  $15.8 \pm 6.0$  m, which constitutes a 19 % difference (Fig. 6; Table 1). The root-mean-squared deviation (RMSD), which represents the averaged difference in FAC between the two models, is 4.0 m. The spatially-integrated FAC, which is the total air volume within the firn layer, is  $34,645 \text{ km}^3$  for SNOWPACK and  $28,581 \text{ km}^3$  for the CFM-GSFC. These values represent all modeled grid cells, meaning they include some areas outside of the six basins defined by Rignot and Mouginot (2012). A few areas of missing data exist and are due to one or both of the firn models encountering an issue in the simulation (Fig. 6). On very few occasions, the simulations were unsuccessful because of numerical instabilities and are treated as missing data. For example, if a grid cell is located in the ablation zone and does not receive enough accumulation to build up a firn layer, we treat that grid point as missing data.

We also calculate FAC for vertical segments in steps of 10-m for the ice sheet from a depth of 0 m (surface) down to 50 m (Fig. 7). In both models, the FAC is highest in the shallowest segment (closest to the surface) and lowest in the deepest segment, consistent with densification. In each vertical segment, the SNOWPACK FAC is on average greater than the CFM CFM-GSFC FAC, but the percent difference between the models increases with depth. The models produce similar FAC values between 0 and 10 m, with a difference of 7 %. However, from 40 to 50 m depth, there is a 29 % difference between SNOWPACK and the CFM CFM-GSFC (Fig. 7).

To better understand spatial patterns represented by both models, we compare FAC in each basin of the GrIS as defined by Rignot and Mouginot (2012). The six basins and their abbreviations used in subsequent figures and tables are the northwest (NW), central west (CW), southwest (SW), north (NO), northeast (NE), and southeast (SE) (e.g., Fig. 6). These basins have distinct climatological features captured by MERRA-2 (Table A1). The southeast and southwest are the warmest and wettest basins as they have the highest temperatures, melt, and precipitation. The precipitation in the southeast is substantially higher (>37 %) than in any other basin. The coldest and driest basins are the north and northeast, which are characterized by very low temperatures, melt, and precipitation. Lastly, the central west and northwest basins fall in the middle with have moderate amounts of precipitation and melt but relatively low temperatures (Table A1).

The FAC means for each basin are shown for both models in Table 1. The central west has the highest basin-mean FAC in both SNOWPACK ( $24.8 \pm 4.6$  m) and the CFM CFM-GSFC ( $19.5 \pm 3.7$  m). In both models, the southeast and northwest have similarly high basin-mean FAC values. The lowest basin-mean FAC occurs in the southwest in SNOWPACK ( $17.4 \pm 7.6$  m) and the CFM CFM-GSFC ( $13.9 \pm 5.8$  m). The best model agreement is in the northeast and north basins where the modeled FAC differs by 13 % and the RMSD is 2.7 m, and in the north where the difference is 1315 % and 15 %, respectively the RMSD is 3.0 m (Table 1). The 1980–1995 trends in the spatially-integrated FAC is the total air volume within the firn layer and is are shown in Table 2 averaged for the first (1980) and last (1995) years of the RCI for each basin. We compare these two

340 ~~years to confirm that no~~ The trends are very small (maximum =  $3.1 \pm 0.3 \text{ km}^3 \text{y}^{-1}$ , which confirms that no substantial change in FAC occurs ~~between the start and end of the RCI that is used for the spin-up. In both SNOWPACK and the CFM, all basins experience less than a 1 % change between the two years, which verifies the steady-state assumption of the RCI and the design of the spin-up (Table 2) over the RCI.~~


345 We examine temporal patterns in FAC spatially integrated across the full ice sheet (Fig. 8) and over each basin (Fig. 9).  
350 During the 1980–1995 RCI, there is no strong trend in the full ice sheet’s air content in either model (Fig. 8), which is consistent with ~~our assignment of that time period as the RCI. By the~~ design of the spin-up, ~~there are no substantial changes in the spatially-integrated FAC between the start and end of the RCI, which means the change to Greenland’s sea level rise buffering capacity is also negligible.~~ Both SNOWPACK and the ~~CFM~~ CFM-GSFC simulate a short-term increase in FAC from 1982 until 1987, followed by a decrease in FAC until 1990 (Fig. 8). Patterns of short-term ( $\sim 1$ –5 years) variability are more prevalent in each basin (Fig. 9). The same increasing then decreasing pattern in Fig. 8 is evident in the basin-averaged FAC in the northeast ~~and~~ southeast, and ~~somewhat detectable in other basins~~ southwest (Fig. 9). Only in the north basin are the short-term trends absent.

355 To examine how the models represent the seasonal cycle in spatially-integrated FAC, ~~we remove during the RCI, we subtract the annual means and from each year to isolate the seasonal signal. We~~ then fit a sine wave to the data (Fig. A3). ~~We and~~ quantify a seasonal “breathing” signal from the amplitude following methods from Ligtenberg et al. (2012). The amplitudes of the seasonal signals in the spatially-integrated FAC for each basin are reported in Table 3. In SNOWPACK, the strongest signal during the RCI is in the southeast, and the weakest is in the northeast. In the ~~CFM~~ CFM-GSFC, the strongest signal is also in the southeast, whereas the weakest is in the north where the seasonality was undetectable by the chosen methods (*i.e.* ~~some basins contain too much intra-annual variability for the sine fitting function to detect a seasonal cycle~~). Integrated across 360 the full ice sheet, the seasonal signal is about two times greater in the ~~CFM~~ CFM-GSFC ( $129 \text{ km}^3$ ) compared to SNOWPACK ( $61 \text{ km}^3$ ) (Table 3; Figs. 8, A3).


### 3.3 Firm properties in a changing climate

365 We now turn our attention to a period where the GrIS was undergoing ~~realtively~~ ~~relatively~~ more change. In the 2005–2020 period, the ice-sheet-wide mean FAC is slightly less than that of the 1980–1995 period for both models, but the difference is not statistically significant ( $p = 0.08$  for SNOWPACK,  $p = 0.30$  for the ~~CFM~~ CFM-GSFC) (Table 1). Between the two periods, the basin-averaged FAC values are not ~~statistically significantly~~ different ( $p > 0.05$  in all basins). In both models, the highest basin-averaged FAC is still in the central west, and the lowest is still in the southwest. Additionally, the best model agreement (13 % difference) still occurs in the northeast (Table 1).


370 After the 1980–1995 RCI, trends in spatially-averaged FAC begin to emerge in the full ice sheet signal (Fig. 8). SNOWPACK models a decreasing trend of  $-66.6 \text{ km}^3 \text{ yr}^{-1}$  during the 2005–2020 period, which is significantly greater in magnitude than the trend of  $1.7 \text{ km}^3 \text{ yr}^{-1}$  throughout the RCI ( $p < 0.05$ ). ~~A consistent decreasing trend is modeled from ~2002 and through ~2011. (Table 2)~~. Two extreme FAC depletion events are captured in 2012 and 2019 (Fig. 8a). Between 1980 and 2020, SNOWPACK simulates a loss of  $1043 \text{ km}^3$  of firn pore space, which could store liquid water equivalent to 2.9 mm of sea level



**Figure 5.** Modeled firn air content (FAC) in SNOWPACK (blue) and the [CFM-GSFC](#) (green) as a function of (a) the liquid-to-solid ratio (LTSR, Eq. 6), and (b) the summer 2-m air temperature, all calculated for 1980 through 1995.



**Figure 6.** Modeled reference climate interval (RCI, 1980–1995) mean [Mean](#) firn air content (FAC) calculated over the upper 100 m of the firn column from (a) SNOWPACK and (b) the [CFM-GSFC](#) for the reference climate interval (RCI, 1980–1995). Panel (c) [SNOWPACK minus](#) shows the [CFM-calculated difference between the modeled](#) FAC values (SNOWPACK minus CFM-GSFC). The values in the bottom right of each panel are the mean FAC and spatially-integrated FAC. Panel (c) also includes the percent difference. Areas where one or both of the models [having](#) [have](#) missing data are shown in white. Black outlines show the six basins defined by Rignot and Mouginot (2012).



**Figure 7.** Modeled reference climate interval (RCI, 1980–1995) mean firn air content (FAC) calculated for 10-m-thick vertical segments over the GrIS. Top row: SNOWPACK, middle row: the ~~CFM~~CFM-GSFC, bottom row: SNOWPACK minus the ~~CFM~~CFM-GSFC. The values in the bottom right of each panel are the mean FAC and spatially-integrated FAC. The bottom row also includes the percent difference averaged over the GrIS.

**Table 1.** Mean modeled firn air content (FAC) for the 1980–1995 reference climate interval (RCI) and for the 2005–2020 period, averaged across each of the six basins shown in Fig. 6. FAC is reported as mean±standard deviation, and the [average](#) percent difference [and](#) [root-mean-square deviation \(RMSD\)](#) between SNOWPACK and the [CFM is CFM-GSFC are](#) also shown. The last row shows the statistics for [all basins together](#) [the full GrIS](#).

Basin	1980–1995				2005–2020			
	CFM-GSFC	SNOWPACK	Diff.	RMSD	CFM-GSFC	SNOWPACK	Diff.	RMSD
	FAC (m)	FAC (m)	(%)	(m)	FAC (m)	FAC (m)	(%)	(m)
NW	18.5±4.1	23.3±5.2	23	5.1	18.5±4.4	23.1±5.8	22	4.9
CW	19.5±3.7	24.8±4.6	24	5.4	19.8±4.0	24.8±5.2	22	5.2
SW	13.9±5.8	17.4±7.6	23	4.2	13.7±6.1	16.9±8.0	21	4.0
NO	15.5±4.6	18.1±5.1	15	3.0	15.2±4.8	17.5±5.5	14	2.7
NE	16.6±3.5	19.0±4.4	13	2.7	16.5±3.8	18.7±4.8	13	2.6
SE	18.8±4.8	23.7±7.7	23	5.9	18.8±5.1	23.4±7.9	22	5.6
GrIS	15.8±6.0	19.1±8.0	19	4.0	15.7±6.3	18.8±8.4	18	3.8

**Table 2.** Modeled spatially-integrated firn air content (FAC) [trends and standard errors of the trends](#) for each of the six basins (Fig. 6) [at for](#) the [beginning and end of the](#) 1980–1995 reference climate interval (RCI) and the 2005–2020 period. The [percent change between 1980 and 1995, and 2005 and 2020 is reported for each model](#). The last row shows the [statistics trends](#) for [all basins summed together](#) [the full GrIS](#).

Basin	1980–1995		2005–2020	
	CFM-GSFC	SNOWPACK	CFM-GSFC	SNOWPACK
	trend ( $\text{km}^3 \text{ y}^{-1}$ )			
NW	-0.1±0.1	-1.1±0.1	-2.6±0.1	-12.3±0.2
CW	+0.2±0.0	+1.5±0.1	-2.1±0.0	-10.3±0.1
SW	-1.0±0.1	-1.2±0.2	-6.9±0.1	-16.7±0.2
NO	+0.6±0.0	+0.4±0.1	-5.7±0.0	-11.4±0.1
NE	-1.3±0.1	-0.8±0.3	+1.9±0.1	-2.5±0.2
SE	+3.1±0.1	+3.1±0.3	+2.4±0.2	-4.5±0.4
GrIS	+1.2±1.0	+1.7±0.9	-17.4±1.2	-66.6±1.2

**Table 3.** The seasonal **breathing** signal in each of the six basins (Fig. 6) for the 1980–1995 reference climate interval (RCI) and the 2005–2020 period. The seasonal signal is the amplitude of the best fit sine curve to the **the** spatially-integrated FAC anomalies (Fig. A3). Undetectable signals are reported as “N/A”, and the last row shows the values for all basins together.

Basin	1980–1995		2005–2020	
	CFM-GSFC signal (km <sup>3</sup> )	SNOWPACK signal (km <sup>3</sup> )	CFM-GSFC signal (km <sup>3</sup> )	SNOWPACK signal (km <sup>3</sup> )
NW	4	9	8	N/A
CW	5	6	9	4
SW	17	5	23	13
NO	N/A	5	4	N/A
NE	11	4	15	N/A
SE	49	25	53	33
GrIS	129	61	165	117

rise. The pattern is similar but dampened in the **CFM**–**CFM-GSFC** record (Fig. 8b). The 2005–2020 trend of  $-17.4 \text{ km}^3 \text{ yr}^{-1}$  is less than that of SNOWPACK ( $p < 0.05$ ) but is still statistically different from the RCI trend of ( $p < 0.05$ ; Table 2). The **CFM**–**CFM-GSFC** shows less depletion of FAC, with only  $356 \text{ km}^3$  of pore space lost (1.0 mm sea level rise) by the end of 2020 compared to 1980.

The marginal areas of the GrIS experience the greatest amount of FAC depletion between 2005 and 2020 (Fig. 10). Both models simulate the same spatial patterns in loss, but the trends vary by basin (Table 2). SNOWPACK simulates a negative trend in spatially-integrated FAC in all basins during this time, with the strongest trend of  $-16.7 \pm 0.2 \text{ km}^3 \text{ yr}^{-1}$  in the southwest. The negative trend is weakest in the northeast ( $-2.5 \pm 0.2 \text{ km}^3 \text{ yr}^{-1}$ ) and southeast ( $-4.5 \pm 0.4 \text{ km}^3 \text{ yr}^{-1}$ ), which are also the only two basins where the CFM-GSFC simulates positive trends ( $1.9 \pm 0.1$  and  $2.4 \pm 0.2 \text{ km}^3 \text{ yr}^{-1}$ , respectively). The CFM-GSFC also simulates the strongest negative trend in the southwest where the spatially-integrated FAC change is  $-6.9 \pm 0.1 \text{ km}^3 \text{ yr}^{-1}$  (Table 2).

Similarities and differences in short-term trends are modeled across each basin (Fig. 9). In the north and northeast, both models show that FAC generally decreases with time and is lower in the more recent 2005–2020 period compared to the reference climate interval, consistent with FAC depletion due to increased melt and associated water percolation and refreezing processes. The interannual variability is strongest in the southeast where there is no clear increasing or decreasing trend following the RCI. In the southwest, FAC is fairly constant until it rapidly drops in 2012. Following this, FAC continues to show no clear trend. The northwest and central west show similar patterns in FAC with values increasing from 1995 to until 2005. After 2005, SNOWPACK FAC decreases while the **CFM**–**CFM-GSFC** FAC remains relatively constant (Fig. 9).


With the exceptions of the northwest and central west, SNOWPACK and the **CFM**–**CFM-GSFC** generally simulate the same trends in FAC (Fig. 9). However, in all cases, differences arise in seasonal variability and magnitudes of change. In all basins, the two models begin to diverge between 2005 and 2012. SNOWPACK simulates greater negative changes in FAC

395 Following this divergence, the magnitude of FAC change is greater in SNOWPACK compared to the CFM once the models diverge CFM-GSFC. In most basins, both models capture an extreme drop in FAC in 2012 associated with the extreme melt occurring in that year, followed by an increase in FAC only in the northeast and southeast. The rapid depletion is greatest in the southwest but less pronounced in the central west and north basins. The magnitude of the 2012 depletion is similar in SNOWPACK and the CFM-CFM-GSFC (Fig. 9).

400 The seasonal breathing signal is stronger in the CFM-CFM-GSFC ( $165 \text{ km}^3$ ) compared to SNOWPACK ( $117 \text{ km}^3$ ) in the 2005–2020 period (Table 3; Fig. A3). In both models, the southeast basin has the strongest seasonality (52 and  $33 \text{ km}^3$  in the CFM-GSFC and SNOWPACK, respectively). The weakest signal in the CFM-CFM-GSFC is in the north, and in SNOWPACK the signal is too weak to be detected with the chosen methods in the northwest, northeast, and north basins. These three basins also have the lowest annual precipitation (Table A2). Additionally, the amplitude of the seasonal signal is greater in the 405 2005–2020 period compared to the 1980–1995 RCI for most basins and for the full ice sheet (Table 3; Fig. A3).

In the 2005 to 2020 period, both models simulate a decrease in FAC, especially in many of the marginal areas of the ice sheet (Fig. 10). The spatially-integrated FAC across the full ice sheet decreases by  $\text{km}^3$  in SNOWPACK, and by  $\text{km}^3$  in the CFM (Table 2). Both SNOWPACK and the CFM show substantial FAC loss in the western basins. In SNOWPACK, the southwest basin's spatially-integrated FAC decreases by  $\text{km}^3$ , which is a 8.3 % change. In the CFM, there is a decrease in the southwest basin of  $\text{km}^3$ , which is a 5.4 % change. The models simulate increasing FAC in some parts of the eastern basins, leading to small percent changes in the northeast and southeast. In fact, the only instance of a positive net change is the 0.2 % increase simulated by the CFM in the northeast basin (Table 2). Individual grid cells showing positive change in FAC during this period are mostly in the northeast and southeast basins (Fig. 10).

410 Observations have shown that FAC is not completely indicative of available pore space to store meltwater when thick near-surface ice slabs are present in the firn. To evaluate how both firn models reproduce the formation of near-surface ice layers, we identify which grid cells simulate ice slabs in the top 20 m of the ice column in April 2014. We choose this depth and date in order to directly compare to ice slabs detected by IceBridge accumulation radar (MacFerrin et al., 2019). We define an ice slab as a layer with a density of at least  $830 \text{ kg m}^{-3}$  and a thickness of at least 1 m. However, we do not distinguish between an ice slab that has formed within the firn and any solid ice exposed at the surface of the ice sheet's ablation zone. Our algorithm 420 outputs the depth and thickness of the ice slab solid ice nearest to the surface, which in some cases could be bare ice at the ablation surface since there is no condition that the ice slab must be beneath a layer of snow or firn. The distribution of modeled ice slabs and ablation surfaces is largely constrained to the marginal regions of the ice sheet (Fig. 11). Both SNOWPACK and the CFM-CFM-GSFC simulate high concentrations of ice slabs solid ice in the margins and ablation zone. Most of the ice slabs detected by IceBridge accumulation radar in 2014 (MacFerrin et al., 2019) overlap with the simulated ice slabs solid ice 425 surfaces. SNOWPACK simulates ice slabs and ablation surfaces in 459 grid points, and the mean depth of those slabs is 2.49 depths to those surfaces is 2.5 m. The CFM-CFM-GSFC simulates ice slabs and ablation surfaces in 369 grid points, and the mean depth to those slabs is 3.81 surfaces is 3.8 m.




**Figure 8.** Weekly firn air content (FAC) spatially-integrated across the full ice sheet from (a) SNOWPACK and (b) the [CFM-GSFC](#). The left y-axis shows the FAC, and the right y-axis shows the equivalent change in sea level buffering capacity relative to 1980. The gray shading represents the 1980–1995 reference climate interval (RCI). The solid black lines are the trends in spatially-integrated FAC for 1980–1995, and the dashed lines are the trends for 2005–2020.

## 4 Discussion

### 4.1 Model evaluation

430 Both In the evaluation of the models with observations, both firn models perform ~~equally well, and their well when evaluated~~ across all SUMup core depths and within each core depth bin. Their ~~overall~~ high NSE coefficients ( $\geq 0.90$ ) and low errors (MAPEs  $\leq 16\%$ ) demonstrate their generally good agreement with observed FAC ~~from the SUMup dataset~~. Despite the overall good agreement between both models, differences in model performance can also be identified. For example, the model performance is not uniform across all core depths (Fig. 3). In deeper cores of at least 10-m depth, the performance is worse than in the full set of cores, and the MAPE is higher than in any subset for both SNOWPACK (27 %) and the CFM-GSFC (19 %). More deep-firn observations may be needed to better evaluate the models at depth.

435 The nature of the discrepancies between the modeled and observed properties tend to differ between the two models. The +7.9 % relative bias in SNOWPACK FAC demonstrates that the model tends to overestimate FAC. This bias in SNOWPACK has been shown in both Greenland (Steger et al., 2017) and Antarctica (Keenan et al., 2021). However, Keenan et al. (2021) also 440 showed that SNOWPACK outperformed other semi-empirical models in the uppermost 10 m in locations where the models



**Figure 9.** (a-c, e-g) Time series of the basin-averaged weekly firn air content (FAC) anomaly from the reference climate interval (RCI, 1980–1995; gray shading) mean, which is shown in the bottom left of each panel. Results from SNOWPACK (“SP”) are shown as the darker lines and the results from the CFM-GSFC (“CFM”) are shown as the lighter lines. Note the different y-axis scales. (d) Greenland’s six basins defined by Rignot and Mouginot (2012) used for calculating trends in firn air content (FAC).


were uncalibrated to the observations, which underscores the utility of SNOWPACK in locations where observations of firn properties are sparse or rapidly changing. In contrast, the CFM-GSFC tends to produce smaller-lower FAC values and has a smaller bias of +0.2 %. The signature of model biases differs across the ice sheet as climate, topography, and the impact of firn hydrology vary. Below we explore the highest model biases and consider the conditions that cause SNOWPACK and the CFM-GSFC to differ from each other and from the observations.

445


Some of the highest model biases ( $>100\%$ ) in SNOWPACK and the CFM-GSFC occur in southeast Greenland and are likely a result of two factors. First, some of the observed density profiles are from cores that were drilled directly into a perennial firn aquifer (Miller et al., 2018). These particular cores approach bulk densities of  $1000 \text{ kg m}^{-3}$  due to the liquid water contained within the pore space of the firn. Neither In this study, neither model captures the high densities resulting from the firn aquifer because the use of bucket scheme in the models prevents full saturation in the firn. Even though the conditions for firn aquifer development have been previously investigated in models using the bucket scheme by analyzing the presence of layers where water remains liquid throughout winter (Kuipers Munneke et al., 2014), thus far only more advanced water percolation schemes based on Richards equation are able to substantially increase the degree of saturation in firn, congruent with firn aquifer formation (Verjans et al., 2019). A firn hydrology model intercomparison study that included outputs from the CFM (but not SNOWPACK) two different densification schemes in the CFM found that none of the nine models included

450

455



**Figure 10.** Modeled difference in mean firn air content (FAC) between 2005 and 2020 for (a) SNOWPACK and (b) the ~~CFM~~CFM-GSFC. Black outlines show the six basins defined by Rignot and Mouginot (2012).



**Figure 11.** Modeled and observed ice slabs in 2014. Red polygons show ice slabs detected by IceBridge accumulation radar (MacFerrin et al., 2019). Modeled ice slabs and ablation surfaces are shaded by the depth to the first ice layer that is at least 1 m thick.

460 accurately simulated meltwater infiltration at the four study sites (Vandecrux et al., 2020b). Moreover, Steger et al. (2017) found that the largest model differences between SNOWPACK and IMAU-FDM (a semi-empirical firn model) occur in the southeast margin of the ice sheet where firn aquifers form. ~~Our results agree with In this study, the high model bias in the five cores in southeast Greenland supports~~ these findings that model differences are highest where liquid water is ~~present, indicating abundant. This indicates~~ abundant. This indicates that poor representation of meltwater percolation processes is still a substantial limiting factor in firn model performance (Verjans et al., 2021).

465 An additional reason why the model bias is high in the southeast is likely the coarseness of the forcing grid in relation to the steep ice sheet topography in this area. The ~~observed cores~~ ~~five locations shown~~ in Figure 4 ~~are all closest to lie within~~ the same MERRA-2 grid ~~point and therefore all have cell and thus share~~ the same atmospheric forcing. However, the cores are located along a transect that spans a steep elevation gradient. ~~Within For~~ Within these five cores, the lowest relative bias occurs at the point that is closest to the MERRA-2 grid ~~point cell~~ center (i.e., furthest west along the transect) and where the elevation is highest (Fig. 4c). The highest relative bias occurs at the point furthest from the MERRA-2 grid ~~point cell~~, where the elevation is lowest, and where the ~~observed local~~ vs. MERRA-2 elevation difference is greatest (Fig. 4g). The steep topography may also lead to strong spatial variations in atmospheric processes such as orographic precipitation, which is not well represented on 470 such a coarse grid (van Kampenhout et al., 2019). This demonstrates the limitations of a coarsely-gridded forcing, especially in steeply sloped areas where climate is likely to be highly variable within a single grid cell.

475 Despite the few instances of relatively high disagreement between the models and observations, the overall good performance of both models in ~~simulating reproducing~~ observations gives confidence in the models' abilities to simulate firn properties across ~~the full ice sheet. Model uncertainty in a wide spatial domain. However, most observations are constrained to the accumulation zone, which limits model validation in the hydrologically complex percolation zone. Compared to the dry and flat ice-sheet interior, areas with firn aquifers or steep surface topography is likely to be higher compared to results from flatter, dry firn zones in the ice-sheet interior topography are likely to have higher model uncertainty.~~

## 4.2 Firm air content response to atmospheric forcing

480 Since we use identical atmospheric data to force the models, differences in the modeled firn properties are purely due to differences in the ~~firm~~ models themselves. The models ~~simulate complex FAC responses to the forcing variables (Fig. A4), but they~~ are particularly distinct in their response to the LTSR and summer air temperature (Fig. 5). Both models show an inverse, non-linear response between LTSR and FAC (Fig. 5a). However, there is more spread in SNOWPACK FAC compared to the ~~CFM CFM-GSFC~~ FAC, especially for low values of LTSR where snowfall is a larger contributor to surface input. Because there is less spread in the ~~CFM CFM-GSFC~~ FAC, the LTSR is a stronger predictor of FAC in the ~~CFM CFM-GSFC ( $r^2 = 0.89$ )~~ compared to SNOWPACK ( $r^2 = 0.77$ ). The large range of possible SNOWPACK-simulated FAC values at low LTSR values is likely due to ~~the model~~ a combination of (1) ~~the model~~'s sophisticated new-snow density scheme that uses more than only air temperature and accumulation to determine near-surface density, and (2) ~~the densification scheme. The CFM-GSFC uses a fixed new-snow density and its densification scheme is empirically based, which likely explains the differences between the two models.~~ In the presence of more liquid surface input (i.e., when the LTSR is high), SNOWPACK produces consistently

490 lower FAC values compared to the ~~CFM~~. This indicates ~~CFM-GSFC~~. Our results show that when more liquid water is present compared to snowfall, the ~~CFM~~ ~~CFM-GSFC~~ still simulates available pore space while SNOWPACK's FAC is near-zero in most cases. The response of FAC to the LTSR is important in our consideration of future climate change since we may see the LTSR increase with future warming. In a transition from lower to higher LTSR values, and by proxy, in a transition to a warmer and wetter climate, the ~~CFM~~ ~~CFM-GSFC~~ shows a more gradual decline in FAC and SNOWPACK shows a more pronounced drop 495 toward near-zero FAC values. These responses to the LTSR can be seen in western Greenland where the ~~CFM~~ ~~CFM-GSFC~~ FAC gradually decreases moving from the interior to the ice sheet margin (Fig. 6b), and SNOWPACK FAC sharply drops off in the same area (Fig. 6a). This is an area vulnerable to modern climate change where increased ice slab formation and decreased firn storage capacity ~~has~~ ~~have~~ already been detected (de la Peña et al., 2015).

500 Additionally, we examine the summer air temperature since it can be directly derived from climate model output unlike the LTSR, which, in the case of MERRA-2, requires more detailed output from a dedicated firn model. We find that a temperature threshold appears to control FAC; at  $\sim 4^{\circ}\text{C}$ , FAC in both models rapidly drops (Fig. 5b). The greatest range in FAC and the highest FAC values occur near this temperature for SNOWPACK and the ~~CFM~~ ~~CFM-GSFC~~. Between  $\sim 4$  and  $\sim 0^{\circ}\text{C}$ , the models simulate almost the full range of FAC. Therefore, outside of the  $-4$ – $0^{\circ}\text{C}$  window, temperature is a relative good predictor of FAC, especially in the ~~CFM~~ ~~CFM-GSFC~~ where the spread in FAC (0 to  $\sim 24$  m) is less than in SNOWPACK (0 505 to  $\sim 32$  m). In both summer air temperature and the LTSR, the spread in the SNOWPACK values shows that FAC is more than just a function of a single variable or metric, and it points to the complexity of the model. Generally, the FAC simulated by the ~~CFM is easier to predict using~~ ~~CFM-GSFC~~ shows less variability for a given LTSR or summer temperature value when compared to SNOWPACK.

#### 4.3 Spatial and temporal patterns in firn air content

510 To place our results in the context of other firn studies, we explore how the modeled FAC compares to existing estimations. We note that this is not a direct comparison since choices of atmospheric forcing, model domain boundaries, and temporal periods will generate differences in FAC that are independent from the firn model choice. Still, we use this comparison to validate the order of magnitude of our results. Our modeled FAC integrated across the full ice sheet for the upper 100 m from the ~~CFM~~ ~~CFM-GSFC~~ ( $28,581 \text{ km}^3$ ) is ~~very~~ similar to a 2010–2017 value ( $26,800 \text{ km}^3$ ) calculated from observations (Vandecrux et al., 515 2019). SNOWPACK's FAC of  $34,645 \text{ km}^3$  is on the same order but still larger than the observations and the ~~CFM-modeled~~ ~~CFM-GSFC-modeled~~ value. SNOWPACK's spatially-integrated FAC in the upper 100 m is close to a regional climate model's (HIRHAM5\_MOD) ~~estimate~~ ~~long-term estimate of~~  $\sim 34,000 \text{ km}^3$  for this period (Vandecrux et al., 2019).

520 SNOWPACK and the ~~CFM~~ ~~CFM-GSFC~~ simulate reasonable ice-sheet-wide FAC when compared to other studies, but the two models still differ in magnitude (Fig. 6). In the upper 10 m, the models agree within  $\pm 10\%$ . However, with increasing depth, the model agreement worsens (Fig. 7). This suggests that the difference in spatially-integrated FAC between the two models arises from the differences in densification with depth. SNOWPACK uses ~~the~~ ~~a~~ constitutive relationship between stress and strain in snow to calculate firn densification, while the ~~CFM~~ ~~CFM-GSFC~~'s densification rate is determined using a semi-empirical equation tuned with firn depth-density data. The higher FAC at greater depths predicted by SNOWPACK

indicates that its modeled densification rate is slower in the deeper firn than the rate in the ~~CFM~~CFM-GSFC. This could be  
525 related to the fact that SNOWPACK was developed using data from seasonal alpine snow, which may not be representative of  
the physical processes driving deep firn ~~densification~~densification (Maeno and Ebinuma, 1983; Arnaud et al., 2000). Similarly,  
Stevens et al. (2020) found that the physically-based snow model Crocus predicted slower densification at Summit, Greenland  
~~than compared to~~ other firn densification equations. Determining which model performs better at depth ~~(and by proxy, whether~~  
~~the physics-based or empirical approach is recommended) is~~ hindered by the paucity of deep firn observations that could  
530 provide insight into densification processes.

Further differences in SNOWPACK and the ~~CFM~~CFM-GSFC are seen in the modeled time series of FAC. Both models  
show a substantial depletion of FAC from 2005 through 2020, but the ~~CFM~~CFM-GSFC's response is smaller than that of  
SNOWPACK. During this time, SNOWPACK's simulated trend of  $-66.6 \text{ km}^3 \text{ yr}^{-1}$  corresponds ~~with a -2.8 to a -3.2 %~~ change  
535 in spatially-integrated FAC. The ~~CFM~~CFM-GSFC's simulated trend of  $-17.4 \text{ km}^3 \text{ yr}^{-1}$  is smaller and the change is also  
~~less at -1.2 to -1.5 %~~. While the magnitudes of trends and FAC changes are greater in SNOWPACK, seasonal signals are greater  
in the ~~CFM~~. ~~The stronger seasonality in the CFM is indicative of the model's more simple treatment of forcing data like~~  
~~CFM-GSFC~~. The temporal patterns in FAC are directly related to the atmospheric forcing. Atmospheric input like accumulation  
and temperature have strong seasonal patterns, which likely makes them the strongest drivers of FAC seasonality. Both models'  
FAC seasonal signals are primarily driven by the forcing, but strength of the signal is tied to how each model treats the  
540 forcing. SNOWPACK's FAC seasonality is weaker than that of the CFM-GSFC, which points to a more complex treatment  
of accumulation and temperature, ~~which have strong seasonal patterns~~. SNOWPACK's same sophisticated in SNOWPACK.  
In particular, these forcing variables are used in SNOWPACK's new-snow density scheme but not in the constant new-snow  
density assumed in the CFM-GSFC. As such, the same new-snow density scheme that leads to ~~a~~ the complex relationship  
between LTSR and FAC ~~also results in this smaller~~ may also dampen SNOWPACK's seasonal signal.

545 Partitioning the record of FAC into climatologically distinct basins reveals further differences in the models and spatial  
variability. Notably, the signature of the 2012 extreme melt season can be seen as an abrupt drop in FAC in ~~most~~three basins  
(Fig. 9). In the southeast, northeast, and northwest, both models show that FAC begins to rebound immediately after the  
2012 depletion. This replenishing of the firn pore space has also been observed in shallow firn cores following the extreme  
2012 melt season (Rennermalm et al., 2021). In the southwest, this rebound is only detected in the ~~CFM~~CFM-GSFC and  
550 not SNOWPACK, which again demonstrates some of the discrepancies between the models. ~~Still, both models generally~~The  
~~replenishing of the firn layer is closely tied to high accumulation and low melt. As such, the models' different responses point~~  
~~to the dissimilar treatment of the atmospheric forcing and densification schemes. While they do not agree in every basin, both~~  
~~models are able to~~ capture the ice sheet's rebuilding of some of the porous firn layer lost during an extreme melt event.

555 While spatially-integrated FAC describes the total volume of pore space in the GrIS, permeability and access to pore space is  
important for fully understanding the buffering capacity. Ice slabs, which ~~may~~ render deep pore space inaccessible to meltwater  
(Machguth et al., 2016), are simulated in the ice sheet's marginal areas where the highest FAC depletion occurs between 2005  
and 2020. ~~In fact, the largest percent change~~The strongest negative trend is in the southwest, ~~which~~ has the warmest temper-  
atures and highest melt compared to other basins during this period (Table A2). These findings agree well with observations

that reveal significant FAC depletion in the low-accumulation percolation zone in western Greenland related to increased melt  
560 (Vandecrux et al., 2019). Pore space depletion can also be ~~a sign of caused by~~ firn densification, which ~~has been found to increase cold content in the firn and amplify meltwater freezing and ice slab formation in the near-surface in turn modifies the meltwater refreezing and retention capacities of the firn in a complex manner~~ (Vandecrux et al., 2020b). FAC depletion is found both where ice slabs ~~are simulated, and~~ and ablation surface are simulated and in areas up-glacier from these ~~ice slabs~~ solid ice surfaces (Figs. 10, 11). This FAC depletion and firn densification may prime the firn for future migration of ice slabs toward 565 the ice-sheet interior. The differences in the modeled ~~ice-slab~~ solid ice locations and depths is likely attributed to the overall diverging behavior of the models in wet firn zones, which agrees with findings in the RetMIP firn model intercomparison study (Vandecrux et al., 2020a). ~~SNOWPACK and the CFM-GSFC both rely on the bucket scheme to govern the vertical percolation of meltwater within the pore space, and both have the same applied surface melt from SNOWPACK's surface energy balance scheme. This means that the differences in ice slab and ablation surface locations and depths are related to the modeled firn structure and temperature, and the models' treatment of atmospheric input. SNOWPACK simulates 24 % more grid cells with ice slabs and ablation surfaces, which is in line with the finding that SNOWPACK's FAC decreases more rapidly (and thus the firn column densifies more rapidly) than the CFM-GSFC's in recent years.~~

570

#### 4.4 Study limitations

The use of identical forcing data (MERRA-2) allows for direct comparison between the SNOWPACK and CFM-GSFC models  
575 in this work. We also use the same approaches to the spin-up, which uses an RCI of 1 January 1980 through 31 December 1995. Although we assume that the steady-state conditions of the RCI represent the Greenland climate preceding 1980, we acknowledge that they are not representative of true conditions. For example, in the ~100 years before the RCI, significant trends in climate over the GrIS have been found (Hanna et al., 2011). Our steady-state assumption does not allow for such 580 trends to appear in the spin-up, but the lack of pre-1980 data necessitates such assumptions. Since the focus of this work is to compare firn model results that are independent of the choice of forcing, the RCI assumptions do not impact the intercomparison. When comparing the firn models to the observations, the steady-state assumption may have an impact at depth. Deeper firn in the models is simulated from the repeated 1980–1995 climate, but real firn in the observations is older and may have formed during times when trends in the climate are apparent (e.g., in the pre-1980 20th century; Hanna et al., 2011)

585 ~  
The firn observations themselves are valuable snapshots of firn properties for a specific time and place, but their lack of temporal and spatial continuity limits the extent of this study's evaluation. Density and firn air content from the SUMup observational dataset do not provide sufficient information about how firn evolves through time. Moreover, the timing of when the observations were collected is not uniformly distributed throughout the year, which means there is less information on, for example, winter firn properties versus summer firn properties. However, a key feature of the SNOWPACK and the CFM-GSFC 590 firn models is their ability to simulate the evolution of firn properties on fine temporal scales, even though validation of these model results is very restricted. Similarly, validating the model sensitivities to changes in climate forcing is hampered by the lack of temporal continuity in the available field data.

Additionally, the fact that the field data are spatially limited means that they do not fully capture the variety of potential firn regimes. This is probably most crucial in the percolation zone where few observations exist, yet where meltwater processes are complex and not often well-represented in models (Vandecrux et al., 2020b). For example, the present study relies on the bucket scheme for vertical meltwater transport, which is a simple choice using only the density to control downward water percolation, but it is severely limited in reproducing more complex melt water processes related to firn aquifer and ice slab formation (Verjans et al., 2021). While more sophisticated water percolation schemes exist (particularly those based on Richards equation) and can obtain better results (Verjans et al., 2019), they also show a stronger dependence on firn properties, including for example grain size. The paucity of percolation zone observations limits the opportunities for evaluating model performance, in particular regarding the choice of meltwater percolation scheme.

## 5 Conclusions

An evaluation of the physics-based firn model SNOWPACK, and the ~~semi-empirical CFM~~ CFM-GSFC (the CFM configured with a semi-empirical densification equation) reveals overall high model performance when compared with ~~observations from~~ 605 ~~FAC derived from density observations in the SUMup dataset. The Nash-Sutcliffe efficiency (NSE) coefficient for SNOWPACK is 0.90 and for the CFM is 0.94~~ Model error is higher in comparisons with deeper cores than in shallower ones, and the mean average percentage error (MAPE) is 14 % for SNOWPACK and 16 % for the CFM. The highest model biases occur in the marginal areas of the ~~ice sheet, which is where conditions for predicting GrIS where conditions controlling~~ FAC (e.g., melt, snowfall, topography) are most variable. Comprehensive model evaluation ~~in this work~~ is hindered by the shortage of deep firn 610 observations ~~and the limited~~ spatial coverage of observations. ~~Still, the overall success of the models and understanding of their limitations allow us to perform full ice sheet simulations and place our results in a context of uncertainty, and the lack of time-varying measurements that could give insight into the evolution of firn properties with time. With the available and chosen evaluation data (i.e., density and FAC), SNOWPACK and the CFM-GSFC differ but still perform well.~~

The use of identical atmospheric forcing lets us ~~isolate the differences in the models themselves and examine how they respond~~ examine how the models respond differently to the same forcing ~~due to structural model differences and parameterization choices~~. The summer air temperature ~~acts as a metric~~ is a factor for examining the impact of forcing conditions on ~~FAC, and reveals high model agreement in warmer temperatures where FAC~~ abruptly decreases above a threshold temperature ~~the~~ the FAC. We find that summer air temperature is related to modeled FAC, which exhibits an abrupt decrease at a summer air temperature threshold of  $\sim 4^{\circ}\text{C}$  ~~in both models~~. For average summer air temperatures above  $\sim 0^{\circ}\text{C}$ , both models simulate low FAC values. 620 We also employ the liquid-to-solid ratio (LTSR) ~~LTSR~~ as a metric summarizing the climatological regime of accumulation and melt. While FAC ~~in the CFM~~ more gradually decreases ~~in the CFM-GSFC~~ as the LTSR increases, SNOWPACK FAC decreases more rapidly and reaches near-zero FAC values ~~that the CFM does not capture~~ not simulated by the CFM-GSFC. These different responses in air content to increasing liquid water will become important in future warming scenarios ~~and we expect the models to diverge even more~~. As such, further work is needed to improve our understanding of the physics of firn hydrology, which 625 ~~will consequently improve the ability of models to simulate FAC response to increased melt and liquid water. This will in turn~~

allow us to better predict the firm's response to future warming. Additionally, the CFM-GSFC's stronger relationship between LTSR and FAC, and the model's stronger seasonal signal point to a more direct impact of forcing variables like accumulation and temperature on the CFM-GSFC compared to SNOWPACK. The models also differ in their response to recent climate change during the 2005–2020 period with SNOWPACK simulating more FAC depletion and more ice slab and ablation area development.

The spatially-integrated FAC during the 1980–1995 RCI is 34,645 km<sup>3</sup> from SNOWPACK and 28,581 km<sup>3</sup> from the CFM-GSFC, which are both reasonable estimations when compared to other studies (e.g., 26, though the 800 km<sup>3</sup> from observations and ~34,000 km<sup>3</sup> from a model (Vandecrux et al., 2019)). However inconsistencies in atmospheric forcing data, model domain area, and temporal periods render direct comparisons between studies difficult. Our spin-up is designed such that no significant change occurs in FAC between the start and end of the 16-year period. However, in a more recent period of the same length (2005–2020), substantial loss of pore space is modeled as a 2.8 % change FAC depletes at a rate of 66.6 km<sup>3</sup> y<sup>-1</sup> in SNOWPACK and -1.2 % change in the CFM. The highest loss of pore space occurs in the ice sheet's 17.4 km<sup>3</sup> y<sup>-1</sup> in the CFM-GSFC. The greatest pore space depletion is along the western margins where near-surface ice slabs are detected both by the both models and observations show development of ice slabs and ablation surfaces. This highlights the vulnerability of the firm layer's meltwater storage capacity, especially in these low-elevation and high-melt areas. Notably, the pore space depletion is more extreme is more depleted in SNOWPACK. It simulates a larger magnitude of change and a more negative 2005–2020 trend (-66.6 km<sup>3</sup> yr<sup>-1</sup>) compared to the CFM (-17.4 km<sup>3</sup> yr<sup>-1</sup>). Over the full 41-year period, SNOWPACK simulates a loss of pore space equivalent to storing 3 mm of sea level rise, while the CFM-GSFC firm loses only an equivalent of 1 mm. Although these longer-term changes are greater in SNOWPACK, the strength of the seasonal signal in the CFM is much greater, again pointing to the different treatment of atmospheric input to the models and the differing complexities of the models.

In the present work, neither model clearly outperforms the other within the scope of the evaluation. Even though we identify disparities between results from both models, we are restricted by the limited availability of the required observational data to draw conclusions about the accuracy of one model over the other. Based on our work, we can draw conclusions about the potential benefits and drawback of each model. The physics-based design of SNOWPACK means that it is not tuned to observations and consequently not biased toward available observational data. This may result in more realistic simulations of firm properties under future climate conditions, whose effects are not capture in existing firm observations. However, firm physics are not fully understood and knowledge gaps limit the accuracy of the model. The CFM's modular design allows for the user to easily choose from several densification schemes. Semi-empirical densification schemes such as the one used in the present work are tuned to observations, which means that realistic densification relationships are built into the model and there is less need to rely on poorly understood physics. Nevertheless, firm models in general are limited by knowledge gaps in firm hydrological processes such as vertical meltwater percolation, lateral flow, and conditions for firm aquifer and ice slab formation. Additional research focusing on obtaining detailed observations of these processes would provide opportunities for important developments in firm modeling.


## Appendix A

**Table A1.** Mean atmospheric forcing variables for the 1980–1995 reference climate interval (RCI), averaged across each of the six basins shown in Fig. 6. Values are reported as mean $\pm$ basin standard deviation.


Basin	Temperature (°C)	Precip. (mm w.e. yr $^{-1}$ )	RH (%)	Wind (m s $^{-1}$ )	ISWR (W m $^{-2}$ )	ILWR (W m $^{-2}$ )	Melt (mm w.e. yr $^{-1}$ )
NW	-23.5 $\pm$ 4.8	390.5 $\pm$ 204.4	92.2 $\pm$ 5.5	4.6 $\pm$ 1.2	115.8 $\pm$ 6.7	169.8 $\pm$ 15.6	43.1 $\pm$ 117.2
CW	-23.4 $\pm$ 4.0	427.3 $\pm$ 132.0	92.6 $\pm$ 4.8	5.7 $\pm$ 1.4	127.2 $\pm$ 4.0	171.9 $\pm$ 12.1	31.5 $\pm$ 103.0
SW	-17.0 $\pm$ 3.7	659.9 $\pm$ 368.4	87.1 $\pm$ 5.3	7.9 $\pm$ 1.7	138.4 $\pm$ 7.8	191.1 $\pm$ 12.8	255.2 $\pm$ 345.6
NO	-24.3 $\pm$ 3.2	192.5 $\pm$ 63.0	91.5 $\pm$ 4.8	3.9 $\pm$ 0.9	111.0 $\pm$ 5.3	164.2 $\pm$ 10.2	23.7 $\pm$ 61.6
NE	-25.4 $\pm$ 4.2	188.0 $\pm$ 89.0	90.1 $\pm$ 7.7	4.0 $\pm$ 1.7	122.7 $\pm$ 5.9	158.6 $\pm$ 11.6	15.9 $\pm$ 50.6
SE	-16.8 $\pm$ 5.6	1049.4 $\pm$ 681.3	85.4 $\pm$ 7.2	6.0 $\pm$ 2.5	132.6 $\pm$ 7.4	191.4 $\pm$ 20.2	262.5 $\pm$ 427.4

**Table A2.** Mean atmospheric forcing variables for the 2005–2020 period, averaged across each of the six basins shown in Fig. 6. Values are reported as mean $\pm$ basin standard deviation.


Basin	Temperature (°C)	Precip. (mm w.e. yr $^{-1}$ )	RH (%)	Wind (m s $^{-1}$ )	ISWR (W m $^{-2}$ )	ILWR (W m $^{-2}$ )	Melt (mm w.e. yr $^{-1}$ )
NW	-22.1 $\pm$ 4.7	399.1 $\pm$ 212.3	91.5 $\pm$ 6.2	4.7 $\pm$ 1.3	116.8 $\pm$ 6.8	173.8 $\pm$ 14.9	78.2 $\pm$ 184.2
CW	-22.0 $\pm$ 4.1	427.7 $\pm$ 137.3	92.1 $\pm$ 5.1	5.8 $\pm$ 1.5	128.2 $\pm$ 4.1	176.4 $\pm$ 12.5	58.0 $\pm$ 166.2
SW	-15.5 $\pm$ 3.6	641.5 $\pm$ 363.3	86.9 $\pm$ 5.4	7.7 $\pm$ 1.6	138.7 $\pm$ 7.6	196.3 $\pm$ 13.0	353.3 $\pm$ 426.4
NO	-22.8 $\pm$ 3.2	164.1 $\pm$ 46.2	90.5 $\pm$ 5.9	3.7 $\pm$ 0.9	113.5 $\pm$ 4.8	167.2 $\pm$ 9.5	48.0 $\pm$ 104.6
NE	-24.0 $\pm$ 4.3	190.5 $\pm$ 95.3	89.6 $\pm$ 8.3	3.9 $\pm$ 1.8	124.0 $\pm$ 5.6	162.8 $\pm$ 11.6	29.8 $\pm$ 81.3
SE	-15.7 $\pm$ 5.6	1079.2 $\pm$ 745.9	85.1 $\pm$ 7.8	6.0 $\pm$ 2.6	133.0 $\pm$ 7.2	195.9 $\pm$ 20.5	313.3 $\pm$ 477.6




**Figure A1.** (a) Observed surface density ( $\rho_0$ ) from SUMup versus SNOWPACK. Since some observations begin farther below the surface, in this figure, observed  $\rho_0$  is defined as the uppermost density measurement that is within 0.1 m from the surface. The SNOWPACK  $\rho_0$  is calculated over the same vertical segment as the SUMup observation. The CFM-GSFC uses a prescribed surface density of 350 kg m<sup>-3</sup> (green vertical line), which falls near many of the observed surface densities (green vertical line). (b) Histogram of observed surface density with the mean represented by the black line. Also plotted is the CFM-GSFC surface density of 350 kg m<sup>-3</sup>.



**Figure A2.** Time series of annually-averaged MERRA-2 forcing variables: (a) temperature, (b) precipitation, (c) relative humidity, (d) wind, (e) incoming shortwave radiation (ISWR), and (f) incoming longwave radiation (ILWR) averaged across the [full-ice-sheetGrIS](#). Also shown is (g) the annually-summed melt output from SNOWPACK used as forcing in the [CFMCFM-GSFC](#). Gray shading represents the reference climate interval from 1980 to 1995.



**Figure A3.** The seasonal **breathing** signal in spatially-integrated weekly firn air content (FAC) for the full ice sheet. (a) SNOWPACK seasonal signal during the 1980–1995 period, (b) SNOWPACK seasonal signal during the 2005–2020 period, (c) the **CFM-GSFC** seasonal signal during the 1980–1995 period, (d) the **CFM-GSFC** seasonal signal during the 2005–2020 period. The thinner lines show the anomaly, which is calculated by subtracting each year’s mean spatially-integrated FAC from the record. The thicker lines are the best fit sine curve. The amplitudes of the sine curves represent the seasonal **breathing** signal. Note that the y-axis scales are the same in all four panels.



**Figure A4.** Modeled firn air content (FAC) in SNOWPACK (blue) and the CFM-GSFC (green) as a function of the forcing variables: (a) temperature, (b) precipitation, (c) relative humidity, (d) wind, (e) incoming shortwave radiation (ISWR), and (f) incoming longwave radiation (ILWR) all calculated for 1980 through 1995. Also shown is (g) the melt, which is calculated by SNOWPACK's surface energy balance model and used as a forcing in the CFM-GSFC.

660 *Code and data availability.* The NASA GSFC MERRA-2 data are available at <https://disc.gsfc.nasa.gov/>. The SUMUp snow density sub-dataset can be found at <https://arcticdata.io/catalog/view/doi%3A10.18739%2FA2NP1WK6M>. Model code is available on GitHub; the code to run the SNOWPACK firn model is available at <https://github.com/snowpack-model/snowpack> and the code to run the Community Firn Model (CFM-GSFC) is available at <https://github.com/UWGlaciology/CommunityFirnModel>. Data for the figures can be found on Zenodo at <https://doi.org/10.5281/zenodo.7671892>.

665 *Author contributions.* MTM and NW ran the SNOWPACK model, and CMS ran the CFM-GSFC. MTM processed and analyzed observational data as well as output from both models, and led the manuscript writing. JTML and BM led the study design. All authors contributed to the writing of the manuscript.

*Competing interests.* The contact author has declared that none of the authors has any competing interests.

670 *Acknowledgements.* The authors acknowledge Eric Keenan for his assistance in accessing MERRA-2 data and compiling the SNOWPACK model. This work used the RMACC Summit supercomputer, which is supported by the National Science Foundation (awards ACI-1532235 and ACI-1532236), the University of Colorado Boulder, and Colorado State University. The Summit supercomputer is a joint effort of the University of Colorado Boulder and Colorado State University.

## References

- Arnaud, L., Barnola, J. M., and Duval, P.: Physical modeling of the densification of snow/firn and ice in the upper part of polar ice sheets, in: 675 Physics of Ice Core Records, edited by Hondoh, T., pp. 285–305, Hokkaido University Press, Sapporo, Japan, 2000.
- Arthern, R. J. and Wingham, D. J.: The Natural Fluctuations of Firn Densification and Their Effect on the Geodetic Determination of Ice Sheet Mass Balance, *Climatic Change*, 40, 605–624, <https://doi.org/10.1023/A:1005320713306>, 1998.
- Arthern, R. J., Vaughan, D. G., Rankin, A. M., Mulvaney, R., and Thomas, E. R.: In situ measurements of Antarctic snow compaction compared with predictions of models, *Journal of Geophysical Research*, 115, F03 011, <https://doi.org/10.1029/2009JF001306>, 2010.
- 680 Bartelt, P. and Lehning, M.: A physical SNOWPACK model for the Swiss avalanche warning Part I: numerical model, *Cold Regions Science and Technology*, p. 23, 2002.
- Bavay, M. and Egger, T.: MeteoIO 2.4.2: a preprocessing library for meteorological data, *Geoscientific Model Development*, 7, 3135–3151, <https://doi.org/10.5194/gmd-7-3135-2014>, publisher: Copernicus GmbH, 2014.
- Benson, C. S.: Stratigraphic Studies in the Snow and Firn of the Greenland Ice Sheet:, Tech. rep., Defense Technical Information Center, 685 Fort Belvoir, VA, <https://doi.org/10.21236/ADA337542>, 1996.
- Braithwaite, R. J., Laternser, M., and Pfeffer, W. T.: Variations of near-surface firn density in the lower accumulation area of the Greenland ice sheet, Pâkitsoq, West Greenland, *Journal of Glaciology*, 40, 477–485, <https://doi.org/10.3189/S002214300001234X>, 1994.
- 690 Cooper, M. G., Smith, L. C., Rennermalm, A. K., Miège, C., Pitcher, L. H., Ryan, J. C., Yang, K., and Cooley, S. W.: Meltwater storage in low-density near-surface bare ice in the Greenland ice sheet ablation zone, *The Cryosphere*, 12, 955–970, <https://doi.org/10.5194/tc-12-955-2018>, 2018.
- Culberg, R., Schroeder, D. M., and Chu, W.: Extreme melt season ice layers reduce firn permeability across Greenland, *Nature Communications*, 12, 2336, <https://doi.org/10.1038/s41467-021-22656-5>, 2021.
- 695 de la Peña, S., Howat, I. M., Nienow, P. W., van den Broeke, M. R., Mosley-Thompson, E., Price, S. F., Mair, D., Noël, B., and Sole, A. J.: Changes in the firn structure of the western Greenland Ice Sheet caused by recent warming, *The Cryosphere*, 9, 1203–1211, <https://doi.org/10.5194/tc-9-1203-2015>, 2015.
- Dunmire, D., Lenaerts, J. T. M., Banwell, A. F., Wever, N., Shragge, J., Lhermitte, S., Drews, R., Pattyn, F., Hansen, J. S. S., Willis, I. C., Miller, J., and Keenan, E.: Observations of Buried Lake Drainage on the Antarctic Ice Sheet, *Geophysical Research Letters*, 47, e2020GL087970, <https://doi.org/10.1029/2020GL087970>, 2020.
- Dunmire, D., Banwell, A. F., Wever, N., Lenaerts, J. T. M., and Datta, R. T.: Contrasting regional variability of buried meltwater extent over 700 2 years across the Greenland Ice Sheet, *The Cryosphere*, 15, 2983–3005, <https://doi.org/10.5194/tc-15-2983-2021>, 2021.
- Enderlin, E. M., Howat, I. M., Jeong, S., Noh, M.-J., van Angelen, J. H., and van den Broeke, M. R.: An improved mass budget for the Greenland ice sheet, *Geophysical Research Letters*, 41, 866–872, <https://doi.org/10.1002/2013GL059010>, 2014.
- 705 Forster, R. R., Box, J. E., van den Broeke, M. R., Miège, C., Burgess, E. W., van Angelen, J. H., Lenaerts, J. T. M., Koenig, L. S., Paden, J., Lewis, C., Gogineni, S. P., Leuschen, C., and McConnell, J. R.: Extensive liquid meltwater storage in firn within the Greenland ice sheet, *Nature Geoscience*, 7, 95–98, <https://doi.org/10.1038/ngeo2043>, *bandiera\_abtest: a Cg\_type: Nature Research Journals Number: 2 Primary\_atype: Research Publisher: Nature Publishing Group Subject\_term: Climate-change impacts;Cryospheric science;Hydrology Subject\_term\_id: climate-change-impacts;cryospheric-science;hydrology*, 2014.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., Silva, A. M. d., Gu, W., Kim, G.-K., Koster, R.,

- 710 Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *Journal of Climate*, 30, 5419–5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- 715 Global Modeling and Assimilation Office (GMAO): MERRA-2 tavg1\_2d\_slv\_Nx: 2d,1-Hourly,Time-Averaged,Single-Level,Assimilation,Single-Level Diagnostics V5.12.4, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC), <https://doi.org/10.5067/VJAFPLI1CSV>, type: dataset, 2015a.
- Global Modeling and Assimilation Office (GMAO): MERRA-2 tavg1\_2d\_int\_Nx: 2d,1-Hourly,Time-Averaged,Single-Level,Assimilation,Vertically Integrated Diagnostics V5.12.4, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC), <https://doi.org/10.5067/Q5GVUVUVGO7>, type: dataset, 2015b.
- 720 Global Modeling and Assimilation Office (GMAO): MERRA-2 tavg1\_2d\_rad\_Nx: 2d,1-Hourly,Time-Averaged,Single-Level,Assimilation,Radiation Diagnostics V5.12.4, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC), <https://doi.org/10.5067/Q9QMY5PBNV1T>, type: dataset, 2015c.
- Global Modeling and Assimilation Office (GMAO): MERRA-2 tavg1\_2d\_flx\_Nx: 2d,1-Hourly,Time-Averaged,Single-Level,Assimilation,Surface Flux Diagnostics V5.12.4, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC), <https://doi.org/10.5067/7MCPBJ41Y0K6>, type: dataset, 2015d.
- 725 Groot Zwaftink, C. D., Cagnati, A., Crepaz, A., Fierz, C., Macelloni, G., Valt, M., and Lehning, M.: Event-driven deposition of snow on the Antarctic Plateau: analyzing field measurements with SNOWPACK, *The Cryosphere*, 7, 333–347, <https://doi.org/10.5194/tc-7-333-2013>, publisher: Copernicus GmbH, 2013.
- Hanna, E., Huybrechts, P., Cappelen, J., Steffen, K., Bales, R. C., Burgess, E., McConnell, J. R., Peder Steffensen, J., Van den Broeke, M., Wake, L., Bigg, G., Griffiths, M., and Savas, D.: Greenland Ice Sheet surface mass balance 1870 to 2010 based on Twentieth Century Reanalysis, and links with global climate forcing, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2011JD016387>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2011JD016387>, 2011.
- 730 Harper, J., Humphrey, N., Pfeffer, W. T., Brown, J., and Fettweis, X.: Greenland ice-sheet contribution to sea-level rise buffered by meltwater storage in firn, *Nature*, 491, 240–243, <https://doi.org/10.1038/nature11566>, 2012.
- Herron, M. M. and Langway, C. C.: Firn Densification: An Empirical Model, *Journal of Glaciology*, 25, 373–385, <https://doi.org/10.3189/S0022143000015239>, 1980.
- Izeboud, M., Lhermitte, S., Van Tricht, K., Lenaerts, J. T. M., Van Lipzig, N. P. M., and Wever, N.: The Spatiotemporal Variability of Cloud Radiative Effects on the Greenland Ice Sheet Surface Mass Balance, *Geophysical Research Letters*, 47, e2020GL087315, <https://doi.org/10.1029/2020GL087315>, 2020.
- 735 Keenan, E., Wever, N., Dattler, M., Lenaerts, J. T. M., Medley, B., Kuipers Munneke, P., and Reijmer, C.: Physics-based SNOWPACK model improves representation of near-surface Antarctic snow and firn density, *The Cryosphere*, 15, 1065–1085, <https://doi.org/10.5194/tc-15-1065-2021>, 2021.
- Kuipers Munneke, P., M. Ligtenberg, S. R., van den Broeke, M. R., van Angelen, J. H., and Forster, R. R.: Explaining the presence of perennial liquid water bodies in the firn of the Greenland Ice Sheet, *Geophysical Research Letters*, 41, 476–483, <https://doi.org/10.1002/2013GL058389>, 2014.
- 740 Kuijpers Munneke, P., Ligtenberg, S. R. M., Noël, B. P. Y., Howat, I. M., Box, J. E., Mosley-Thompson, E., McConnell, J. R., Steffen, K., Harper, J. T., Das, S. B., and van den Broeke, M. R.: Elevation change of the Greenland Ice Sheet due to surface mass balance and firn processes, 1960–2014, *The Cryosphere*, 9, 2009–2025, <https://doi.org/10.5194/tc-9-2009-2015>, publisher: Copernicus GmbH, 2015.

- Lehning, M., Bartelt, P., Brown, B., and Fierz, C.: A physical SNOWPACK model for the Swiss avalanche warning Part III: meteorological forcing, thin layer formation and evaluation, *Cold Regions Science and Technology*, p. 16, 2002a.
- 750 Lehning, M., Bartelt, P., Brown, B., Fierz, C., and Satyawali, P.: A physical SNOWPACK model for the Swiss avalanche warning Part II. Snow microstructure, *Cold Regions Science and Technology*, p. 21, 2002b.
- Li, J. and Zwally, H. J.: Modeling of firn compaction for estimating ice-sheet mass change from observed ice-sheet elevation change, *Annals of Glaciology*, 52, 1–7, <https://doi.org/10.3189/172756411799096321>, publisher: Cambridge University Press, 2011.
- Ligtenberg, S. R. M., Helsen, M. M., and van den Broeke, M. R.: An improved semi-empirical model for the densification of Antarctic firn, 755 *The Cryosphere*, 5, 809–819, <https://doi.org/10.5194/tc-5-809-2011>, 2011.
- Ligtenberg, S. R. M., Horwath, M., van den Broeke, M. R., and Legréy, B.: Quantifying the seasonal “breathing” of the Antarctic ice sheet, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL053628>, 2012.
- Ligtenberg, S. R. M., Kuipers Munneke, P., Noël, B. P. Y., and van den Broeke, M. R.: Brief communication: Improved simulation of the present-day Greenland firn layer (1960–2016), *The Cryosphere*, 12, 1643–1649, <https://doi.org/10.5194/tc-12-1643-2018>, 2018.
- 760 Lundin, J. M. D., Stevens, C. M., Arthern, R., Buiertz, C., Orsi, A., Ligtenberg, S. R. M., Simonsen, S. B., Cummings, E., Essery, R., Leahy, W., Harris, P., Helsen, M. M., and Waddington, E. D.: Firn Model Intercomparison Experiment (FirnMICE), *Journal of Glaciology*, 63, 401–422, <https://doi.org/10.1017/jog.2016.114>, 2017.
- MacFerrin, M., Machguth, H., As, D. v., Charalampidis, C., Stevens, C. M., Heilig, A., Vandecrux, B., Langen, P. L., Mottram, R., Fettweis, X., Broeke, M. R. v. d., Pfeffer, W. T., Moussavi, M. S., and Abdalati, W.: Rapid expansion of Greenland’s low-permeability ice slabs, 765 *Nature*, 573, 403–407, <https://doi.org/10.1038/s41586-019-1550-3>, 2019.
- MacFerrin, M. J., Stevens, C. M., Vandecrux, B., Waddington, E. D., and Abdalati, W.: The Greenland Firn Compaction Verification and Reconnaissance (FirnCover) dataset, 2013–2019, *Earth System Science Data*, 14, 955–971, <https://doi.org/10.5194/essd-14-955-2022>, 2022.
- 770 Machguth, H., MacFerrin, M., van As, D., Box, J. E., Charalampidis, C., Colgan, W., Fausto, R. S., Meijer, H. A. J., Mosley-Thompson, E., and van de Wal, R. S. W.: Greenland meltwater storage in firn limited by near-surface ice formation, *Nature Climate Change*, 6, 390–393, <https://doi.org/10.1038/nclimate2899>, 2016.
- Maeno, N. and Ebinuma, T.: Pressure sintering of ice and its implication to the densification of snow at polar glaciers and ice sheets, *The Journal of Physical Chemistry*, 87, 4103–4110, <https://doi.org/10.1021/j100244a023>, publisher: American Chemical Society, 1983.
- Medley, B., Neumann, T. A., Zwally, H. J., Smith, B. E., and Stevens, C. M.: Simulations of firn processes over the Greenland and Antarctic 775 ice sheets: 1980–2021, *The Cryosphere*, 16, 3971–4011, <https://doi.org/10.5194/tc-16-3971-2022>, 2022.
- Michlmayr, G., Lehning, M., Koboltschnig, G., Holzmann, H., Zappa, M., Mott, R., and Schöner, W.: Application of the Alpine 3D model for glacier mass balance and glacier runoff studies at Goldbergkees, Austria, *Hydrological Processes*, 22, 3941–3949, <https://doi.org/10.1002/hyp.7102>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/hyp.7102>, 2008.
- 780 Miller, O., Solomon, D. K., Miège, C., Koenig, L., Forster, R., Schmerr, N., Ligtenberg, S. R. M., and Montgomery, L.: Direct Evidence of Meltwater Flow Within a Firn Aquifer in Southeast Greenland, *Geophysical Research Letters*, 45, 207–215, <https://doi.org/10.1002/2017GL075707>, 2018.
- Montgomery, L., Koenig, L., and Alexander, P.: The SUMup dataset: compiled measurements of surface mass balance components over ice sheets and sea ice with analysis over Greenland, *Earth System Science Data*, 10, 1959–1985, <https://doi.org/10.5194/essd-10-1959-2018>, 2018.
- 785 Moon, T., Fisher, M., Harden, L., Simonoko, H., and Stafford, T.: QGreenland (v2.0.0), National Snow and Ice Data Center, 2022.

- Morlighem, M., Williams, C. N., Rignot, E., An, L., Arndt, J. E., Bamber, J. L., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L., Mouginot, J., Noël, B. P. Y., O'Cofaigh, C., Palmer, S., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., and Zingleren, K. B.: BedMachine v3: Complete Bed Topography and Ocean Bathymetry Mapping of Greenland From Multibeam Echo Sounding Combined With Mass Conservation, *Geophysical Research Letters*, 44, 11,051–11,061, <https://doi.org/10.1002/2017GL074954>, *\_eprint*: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2017GL074954>, 2017.
- 790 Morris, E. M. and Wingham, D. J.: Densification of polar snow: Measurements, modeling, and implications for altimetry, *Journal of Geophysical Research: Earth Surface*, 119, 349–365, <https://doi.org/10.1002/2013JF002898>, *\_eprint*: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2013JF002898>, 2014.
- 795 Morris, E. M. and Wingham, D. J.: Uncertainty in mass-balance trends derived from altimetry: a case study along the EGIG line, central Greenland, *Journal of Glaciology*, 61, 345–356, <https://doi.org/10.3189/2015JoG14J123>, publisher: Cambridge University Press, 2015.
- Nash, J. E. and Sutcliffe, J. V.: River flow forecasting through conceptual models part I — A discussion of principles, *Journal of Hydrology*, 10, 282–290, [https://doi.org/10.1016/0022-1694\(70\)90255-6](https://doi.org/10.1016/0022-1694(70)90255-6), 1970.
- 800 Pfeffer, W. T., Meier, M. F., and Illangasekare, T. H.: Retention of Greenland runoff by refreezing: Implications for projected future sea level change, *Journal of Geophysical Research: Oceans*, 96, 22 117–22 124, <https://doi.org/10.1029/91JC02502>, 1991.
- Rennermalm, A. K., Hock, R., Covi, F., Xiao, J., Corti, G., Kingslake, J., Leidman, S. Z., Miège, C., Macferrin, M., Machguth, H., Osterberg, E., Kameda, T., and McConnell, J. R.: Shallow firn cores 1989–2019 in southwest Greenland’s percolation zone reveal decreasing density and ice layer thickness after 2012, *Journal of Glaciology*, pp. 1–12, <https://doi.org/10.1017/jog.2021.102>, 2021.
- 805 Rignot, E. and Mouginot, J.: Ice flow in Greenland for the International Polar Year 2008–2009, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL051634>, 2012.
- Schlögl, S., Lehning, M., Nishimura, K., Huwald, H., Cullen, N. J., and Mott, R.: How do Stability Corrections Perform in the Stable Boundary Layer Over Snow?, *Boundary-Layer Meteorology*, 165, 161–180, <https://doi.org/10.1007/s10546-017-0262-1>, 2017.
- 810 Simonsen, S. B., Stenseng, L., Adalgeirsdsóttir, G., Fausto, R. S., Hvidberg, C. S., and Lucas-Picher, P.: Assessing a multilayered dynamic firn-compaction model for Greenland with ASIRAS radar measurements, *Journal of Glaciology*, 59, 545–558, <https://doi.org/10.3189/2013JoG12J158>, publisher: Cambridge University Press, 2013.
- Smith, B. E., Medley, B., Fettweis, X., Sutterley, T., Alexander, P., Porter, D., and Tedesco, M.: Evaluating Greenland Surface-Mass-Balance and Firn-Densification Data Using ICESat-2 Altimetry, *preprint*, Ice sheets/Mass Balance Obs, <https://tc.copernicus.org/preprints/tc-2022-44/>, 2022.
- 815 Stearns, C. R. and Weidner, G. A.: Sensible and Latent Heat Flux Estimates in Antarctica, in: *Antarctic Meteorology and Climatology: Studies Based on Automatic Weather Stations*, pp. 109–138, American Geophysical Union (AGU), <https://doi.org/10.1029/AR061p0109>, *\_eprint*: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/AR061p0109>, 1993.
- Steger, C. R., Reijmer, C. H., van den Broeke, M. R., Wever, N., Forster, R. R., Koenig, L. S., Kuipers Munneke, P., Lehning, M., Lhermitte, S., Ligtenberg, S. R. M., Miège, C., and Noël, B. P. Y.: Firn Meltwater Retention on the Greenland Ice Sheet: A Model Comparison, *Frontiers in Earth Science*, 5, 3, <https://doi.org/10.3389/feart.2017.00003>, 2017.
- 820 Stevens, C. M., Verjans, V., Lundin, J. M. D., Kahle, E. C., Horlings, A. N., Horlings, B. I., and Waddington, E. D.: The Community Firn Model (CFM) v1.0, *Geoscientific Model Development*, 13, 4355–4377, <https://doi.org/10.5194/gmd-13-4355-2020>, 2020.

- Sørensen, L. S., Simonsen, S. B., Nielsen, K., Lucas-Picher, P., Spada, G., Adalgeirsdottir, G., Forsberg, R., and Hvidberg, C. S.: Mass balance of the Greenland ice sheet (2003–2008) from ICESat data – the impact of interpolation, sampling and firn density, *The Cryosphere*, 5, 173–186, <https://doi.org/10.5194/tc-5-173-2011>, publisher: Copernicus GmbH, 2011.
- 825 Tedstone, A. J. and Machguth, H.: Increasing surface runoff from Greenland's firn areas, *Nature Climate Change*, 12, 672–676, <https://doi.org/10.1038/s41558-022-01371-z>, number: 7 Publisher: Nature Publishing Group, 2022.
- Thompson-Munson, M., Montgomery, L., Lenaerts, J., and Koenig, L.: Surface Mass Balance and Snow Depth on Sea Ice Working Group (SUMup) snow density subdataset, Greenland and Antarctica, 1952–2019, Arctic Data Center, <https://doi.org/10.18739/A24Q7QR58>, 2022.
- van Angelen, J. H., M. Lenaerts, J. T., van den Broeke, M. R., Fettweis, X., and van Meijgaard, E.: Rapid loss of firn pore space accelerates 830 21st century Greenland mass loss, *Geophysical Research Letters*, 40, 2109–2113, <https://doi.org/10.1002/grl.50490>, 2013.
- van den Broeke, M., Bamber, J., Ettema, J., Rignot, E., Schrama, E., van de Berg, W. J., van Meijgaard, E., Velicogna, I., and Wouters, B.: Partitioning Recent Greenland Mass Loss, *Science*, 326, 984–986, <https://doi.org/10.1126/science.1178176>, 2009.
- van Kampenhout, L., Rhoades, A. M., Herrington, A. R., Zarzycki, C. M., Lenaerts, J. T. M., Sacks, W. J., and van den Broeke, M. R.: Regional grid refinement in an Earth system model: impacts on the simulated Greenland surface mass balance, *The Cryosphere*, 13, 835 1547–1564, <https://doi.org/10.5194/tc-13-1547-2019>, publisher: Copernicus GmbH, 2019.
- Van Tricht, K., Lhermitte, S., Lenaerts, J. T. M., Gorodetskaya, I. V., L'Ecuyer, T. S., Noël, B., van den Broeke, M. R., Turner, D. D., and van Lipzig, N. P. M.: Clouds enhance Greenland ice sheet meltwater runoff, *Nature Communications*, 7, 10266, <https://doi.org/10.1038/ncomms10266>, 2016.
- Vandecrux, B., MacFerrin, M., Machguth, H., Colgan, W. T., van As, D., Heilig, A., Stevens, C. M., Charalampidis, C., Fausto, R. S., Morris, 840 E. M., Mosley-Thompson, E., Koenig, L., Montgomery, L. N., Miège, C., Simonsen, S. B., Ingeman-Nielsen, T., and Box, J. E.: Firn data compilation reveals widespread decrease of firn air content in western Greenland, *The Cryosphere*, 13, 845–859, <https://doi.org/10.5194/tc-13-845-2019>, 2019.
- Vandecrux, B., Fausto, R. S., van As, D., Colgan, W., Langen, P. L., Haubner, K., Ingeman-Nielsen, T., Heilig, A., Stevens, C. M., MacFerrin, 845 M., Niwano, M., Steffen, K., and Box, J.: Firn cold content evolution at nine sites on the Greenland ice sheet between 1998 and 2017, *Journal of Glaciology*, 66, 591–602, <https://doi.org/10.1017/jog.2020.30>, 2020a.
- Vandecrux, B., Mottram, R., Langen, P. L., Fausto, R. S., Olesen, M., Stevens, C. M., Verjans, V., Leeson, A., Ligtenberg, S., Kuipers Munneke, P., Marchenko, S., van Pelt, W., Meyer, C. R., Simonsen, S. B., Heilig, A., Samimi, S., Marshall, S., Machguth, H., MacFerrin, M., Niwano, M., Miller, O., Voss, C. I., and Box, J. E.: The firn meltwater Retention Model Intercomparison Project (RetMIP): evaluation of nine firn models at four weather station sites on the Greenland ice sheet, *The Cryosphere*, 14, 3785–3810, 850 <https://doi.org/10.5194/tc-14-3785-2020>, 2020b.
- Verjans, V., Leeson, A. A., Stevens, C. M., MacFerrin, M., Noël, B., and van den Broeke, M. R.: Development of physically based liquid water schemes for Greenland firn-densification models, *The Cryosphere*, 13, 1819–1842, <https://doi.org/10.5194/tc-13-1819-2019>, 2019.
- Verjans, V., Leeson, A. A., Nemeth, C., Stevens, C. M., Kuipers Munneke, P., Noël, B., and van Wessem, J. M.: Bayesian calibration of firn 855 densification models, *The Cryosphere*, 14, 3017–3032, <https://doi.org/10.5194/tc-14-3017-2020>, 2020.
- Verjans, V., Leeson, A. A., McMillan, M., Stevens, C. M., van Wessem, J. M., van de Berg, W. J., van den Broeke, M. R., Kittel, C., Amory, C., Fettweis, X., Hansen, N., Boberg, F., and Mottram, R.: Uncertainty in East Antarctic Firn Thickness Constrained Using a Model Ensemble Approach, *Geophysical Research Letters*, 48, e2020GL092060, <https://doi.org/10.1029/2020GL092060>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2020GL092060>, 2021.

- Vionnet, V., Brun, E., Morin, S., Boone, A., Faroux, S., Le Moigne, P., Martin, E., and Willemet, J.-M.: The detailed snowpack scheme  
860 Crocus and its implementation in SURFEX v7.2, *Geoscientific Model Development*, 5, 773–791, <https://doi.org/10.5194/gmd-5-773-2012>, publisher: Copernicus GmbH, 2012.
- Wever, N., Keenan, E., Amory, C., Lehning, M., Sigmund, A., Huwald, H., and Lenaerts, J. T. M.: Observations and simulations of new snow  
865 density in the drifting snow-dominated environment of Antarctica, *Journal of Glaciology*, pp. 1–18, <https://doi.org/10.1017/jog.2022.102>, 2022.
- Wilhelms, F.: Density of ice core ngt03C93.2 from the North Greenland Traverse, PANGAEA, <https://doi.org/10.1594/PANGAEA.56560>,  
865 publisher: PANGAEA Type: dataset, 2000.
- Zhang, W., Wang, Y., Smeets, P. C. J. P., Reijmer, C. H., Huai, B., Wang, J., and Sun, W.: Estimating near-surface climatology of multi-  
870 reanalyses over the Greenland Ice Sheet, *Atmospheric Research*, 259, 105 676, <https://doi.org/10.1016/j.atmosres.2021.105676>, 2021.