

# Biogeochemical evolution of ponded meltwater in a High Arctic subglacial tunnel

Ashley J. Dubnick<sup>1,2</sup>, Rachel L. Spietz<sup>3</sup>, Brad D. Danielson<sup>4</sup>, Mark L. Skidmore<sup>1</sup>, Eric S. Boyd<sup>3</sup>, Dave Burgess<sup>4</sup>, Charvanaa Dhoonmoon<sup>2</sup>, Martin Sharp<sup>2</sup>.

- 5 <sup>1</sup>Department of Earth Sciences, Montana State University, Bozeman, Montana, 59717, USA  
<sup>2</sup>Department of Earth and Atmospheric Sciences, University of Alberta, Edmonton, Alberta, T6G 2E3, Canada  
<sup>3</sup>Department of Microbiology and Cell Biology, Montana State University, Bozeman, MT, 59717, USA  
<sup>4</sup>Geological Survey of Canada, 601 Booth Street, Ottawa, Ontario, K1A 0E8, Canada

*Correspondence to:* Ashley J. Dubnick (ashley.dubnick@montana.edu)

- 10 **Abstract.** Subglacial environments comprise ~10% of Earth's land surface, host active microbial ecosystems, and are important components of global biogeochemical cycles. However, the broadly inaccessible nature of subglacial systems has left them vastly understudied, and research to date has been limited to laboratory experiments or field measurements using basal ice or subglacial water accessed through boreholes or from the glacier margin. In this study, we extend our understanding of subglacial biogeochemistry and microbiology to include observations of a slushy pond of water that occupied a remnant  
15 meltwater channel beneath a polythermal glacier in the Canadian High Arctic over winter. The hydraulics and geochemistry of the system suggest that the pond water originated as late-season, ice-marginal runoff with less than ~15% solute contribution from subglacial sources. Over the eight months of persistent sub-zero regional temperatures, the pond gradually froze, cryo-concentrating solutes in the residual water by up to seven times. Despite cryo-concentration and the likely influx of some subglacial solute, the pond was depleted in only the most labile and biogeochemically-relevant compounds, including  
20 ammonium, phosphate, and dissolved organic matter, including a potentially labile tyrosine-like component. DNA amplicon sequencing revealed decreasing microbial diversity with distance into the meltwater channel. The pond at the terminus of the channel hosted a microbial community inherited from late-season meltwater, which was dominated by only six taxa related to known psychrophilic/psychrotolerant heterotrophs that have high metabolic diversity and broad habitat ranges. Collectively, our findings suggest that generalist microbes from the extraglacial or supraglacial environments can become established in  
25 subglacial aquatic systems and deplete reservoirs of nutrients and dissolved organic carbon over a period of months. These findings extend our understanding of the microbial and biogeochemical evolution of subglacial aquatic ecosystems and the extent of their habitability.

## 1 Introduction

30 Subglacial environments currently cover approximately 10% of Earth's land surface and have occupied most of these areas for thousands of years. The broadly inaccessible nature of these systems has left them vastly understudied relative to other terrestrial ecosystems. Yet, over the last two decades efforts to explore subglacial sediment, water, and ice have consistently characterized them as active microbial and chemical systems that are important to global-scale biogeochemical cycles (e.g. Boyd *et al.*, 2010; Hawkings *et al.*, 2014; Wadham *et al.*, 2019; Kellerman *et al.*, 2021).

35 Microorganisms are ubiquitous in subglacial systems. They are found in subglacial water at concentrations ranging from  $1 \times 10^2$  to  $1 \times 10^5$  cells  $\text{ml}^{-1}$  (Christner *et al.*, 2014; Sheridan *et al.*, 2003; Grasby *et al.*, 2003) and are even more abundant in subglacial sediment at concentrations of  $10^6$  to  $10^7$  cells  $\text{g}^{-1}$  of sediment (Sharp *et al.*, 1999; Lanoil *et al.*, 2009). Studies consistently find that among these cells exists a metabolically-active microbial community that is often phylogenetically and functionally diverse (Vick-Majors *et al.*, 2016; Yde *et al.*, 2010; Christner *et al.*, 2014; Hamilton *et al.*, 2013). Some of these  
40 subglacial ecosystems are sustained by reservoirs of organic matter (Hood *et al.*, 2015) acquired as glaciers override soil and vegetation (Christ *et al.*, 2021), marine deposits (Wadham *et al.*, 2012; Michaud *et al.*, 2016), and organic-rich shales (Wadham *et al.*, 2004; Grasby *et al.*, 2003), or from the in-wash of supraglacial or ice-marginal organic matter (Tranter *et al.*, 2005; Andrews *et al.*, 2018). This organic matter can support microorganisms with heterotrophic metabolisms, even under anaerobic conditions involving methanogenesis as a terminal process (Boyd *et al.*, 2010; Stibal *et al.*, 2012a; Wadham *et al.*, 2012).

45 Primary production in many subglacial ecosystems is dominated by lithotrophic microbial metabolisms (Kayani *et al.*, 2018; Boyd *et al.*, 2014; Christner *et al.*, 2014; Dunham *et al.*, 2021), in which organisms utilize solutes or elements from the underlying bedrock to liberate chemical energy. Microbially-mediated redox reactions in subglacial systems occur via the oxidation of reduced inorganic compounds such as ferrous iron ( $\text{Fe}^{2+}$ ), sulfide ( $\text{S}^{2-}$ ), hydrogen ( $\text{H}_2$ ), ammonium ( $\text{NH}_4^+$ ), and methane ( $\text{CH}_4$ ), coupled to the reduction of species such as oxygen ( $\text{O}_2$ ), nitrate ( $\text{NO}_3^-$ ), carbon dioxide ( $\text{CO}_2$ ), or sulfate ( $\text{SO}_4^{2-}$ )  
50 (Miteva *et al.*, 2004; Boyd *et al.*, 2011; Yde *et al.*, 2010; Stibal *et al.*, 2012a; Wadham *et al.*, 2004; Tranter *et al.*, 2002; Boyd *et al.*, 2014; Dunham *et al.*, 2021; Achberger *et al.*, 2016; Christner *et al.*, 2014; Michaud *et al.*, 2017). Glacial comminution serves to liberate, and in some cases even produce (Telling *et al.*, 2015; Macdonald *et al.*, 2018; Gill-Olivas *et al.*, 2021), redox-sensitive species for subglacial ecosystems. Subglacial bedrock and freshly comminuted sediments can also contain high concentrations of labile nutrients required to support microbial activity, including phosphorus (Föllmi *et al.*, 2009; Hawkings  
55 *et al.*, 2016; Hodson *et al.*, 2004), iron (Bhatia *et al.*, 2013; Hawkings *et al.*, 2014; Schroth *et al.*, 2014), and nitrogen (Hodson *et al.*, 2005; Lawson *et al.*, 2014; Wadham *et al.*, 2016; Gill-Olivas *et al.*, 2023).

The research that has informed our understanding of the microbiology and geochemistry of subglacial systems has been conducted using samples of basal ice (e.g. Barker *et al.*, 2010; Montross *et al.*, 2014; Dubnick *et al.*, 2020) and subglacial water accessed via boreholes (e.g. Tranter, 2003; Christner *et al.*, 2014) or from the glacier margin (e.g. Foght *et al.*, 2004;  
60 Skidmore *et al.*, 2005; Boyd *et al.*, 2011; Sheik *et al.*, 2015). Though these environmental samples have advanced our understanding of subglacial biogeochemistry, the inaccessible and dynamic nature of subglacial environments make it difficult

to interpret observations in the context of residence times, water histories, or solute sources, and thus fully resolve the processes that occur in-situ or the rates at which they occur. In this study, we extend our understanding of subglacial biogeochemistry and microbiology to include observations on the evolution of meltwater after it overwinters at the endpoint of a remnant subglacial channel that extends 467 m beneath a glacier. The nature of this system provides us with a unique opportunity to observe the biogeochemical signatures of a relatively isolated subglacial system with strong constraints on the potential solute sources, water histories, and subglacial residence times.


We compared the biogeochemistry and microbiology of water samples from a pond at the terminus of the channel to their water and solute sources: late season runoff (represented by ice samples collected from frozen sections of the channel floor) and basal solute (represented by basal ice exposed along the tunnel walls). We used water isotopes ( $\delta^2\text{H}$ - $\delta^{18}\text{O}$ ) and a conservative geochemical tracer ( $\text{Cl}^-$ ) to quantify the extent to which the late-season water froze and evaporated *in situ*. Using these results, a biogeochemical freeze-fractionation model was developed to 1) identify solutes that appear to behave conservatively and those in which there is an apparent in-situ source or sink, 2) evaluate the extent to which basal solutes may have contributed to the waterbody, and 3) in combination with 16S rRNA gene amplicon sequencing, explore evidence for *in situ* microbial activity and nutrient cycling. Collectively, the biogeochemical and microbial datasets provide evidence of *in situ* microbial activity and show that a distinct microbial community can develop in a subglacial waterbody and deplete reservoirs of the most labile nutrients within months.

## 2 Methods

### 2.1 Field Site

The Sverdrup Glacier is a 25-km long tidewater glacier that overrides metasedimentary rocks and gneiss bedrock (Harrison et al., 2016) while draining a  $\sim 800 \text{ km}^2$  northwest sector of the Devon Ice Cap, Canadian Arctic (Figure 1). Ice velocities along the Sverdrup Glacier are moderate, ranging from  $\sim 30 \text{ m yr}^{-1}$  near the glacier headwall to  $\sim 75 \text{ m yr}^{-1}$  near the terminus (Van Wychen et al., 2017; Cress and Wyness, 1961). While some areas of the glacier are probably frozen to the bed (Van Wychen et al., 2017), flow rates along much of the glacier are influenced by basal sliding or enhanced deformation of basal ice, suggesting that basal ice temperatures, in at least some areas of the bed, approach the pressure-melting point (Van Wychen et al., 2017; Burgess et al., 2005). A two-fold increase in surface velocity of the Sverdrup Glacier first measured during the summer of 1961 (Cress and Wyness, 1961) suggests meltwater reaches the bed and significantly reduces friction.

Consistent with other observations across the Canadian high Arctic, net mass balance of the Sverdrup glacier basin is controlled almost entirely by surface melt (Koerner, 2005). Surface temperatures measured on the Sverdrup Glacier from 2005-2021 indicate a daily average of  $4.5^\circ\text{C}$  over the summer melt season, which extends from early June to late August. Most of the melt generated on the Sverdrup Glacier drains ice-marginally, with the remainder draining via supraglacial streams that reach the bed through moulins near the marine-terminating glacier snout (Keeler, 1964; Koerner, 1961). Winter temperatures average  $-26^\circ\text{C}$ , suggesting that penetration of the winter cold wave likely leaves ice frozen to the bed along the lateral margins, as is the



95 Figure 1: (a) Sverdrup glacier including radar and laser altimeter transects (Transect A-B; Paden et al., 2019) and subglacial tunnel survey. Base map is the Hillshade image from ArcticDEM (Porter et al., 2018), (b) Devon Ice Cap (ESRI satellite base map), showing the location of the Sverdrup Glacier. (c) cross sectional profile showing the bedrock and ice surface along A-B (Paden et al., 2019) with the approximate distance of the subglacial pond from the lateral margin imposed on this transect.

case for many polythermal glaciers (Bingham et al., 2006; Irvine-Fynn et al., 2011). The area explored in this study focuses  
100 on ponded water located at the endpoint of a remnant subglacial tunnel along the eastern margin, ~3 km from the glacier  
terminus (Figure 1).

## 2.2 Field Methods

We visited the Sverdrup subglacial tunnel in May 2019 and mapped the system using a TruPulse®360 (Laser Tech (LTI),  
Colorado USA) rangefinder relative to a known reference point at the glacier margin for which coordinates were collected  
105 using a handheld GPS. The rangefinder device was mounted on a tripod with a known height, and the azimuth, inclination,  
horizontal distance, and vertical distance were collected for a back-sight and fore-sight targets at intervals (of <30 m) along  
the channel. Absolute distances and azimuths were validated using a tape measure and protractor, respectively. Ice temperature  
of the channel wall was measured at ~50 m intervals along the survey track using a BIOS digital contact thermometer.

Samples were collected from the subglacial pond for water isotope, biogeochemical, and microbial analyses at four locations  
110 (S6-S9; Figure 2). The first location (S6) was at the pond edge and was comprised of wet ice, which was collected into a whirl-  
pak® bag using a flame-sterilized ice axe. Though this sample was ice, we consider it to be affiliated with the pond water since  
it was situated in a depression that would receive continual seepage of water from other areas of the pond, and where  
temperatures would be sufficiently cold to freeze it relatively fast and completely. The other three sample sites were collected  
at intervals (~ 25 m apart) into the pond. At each location, pond slush was scooped into a glass beaker that was acid-washed,  
115 rinsed with 18.2 MΩ cm<sup>-1</sup> deionized water (DIW), and furnace (450°C for 4 hrs). Pond ice and water were separated by  
decanting the two fractions into separate whirl-pak® bags. Water samples were immediately frozen, and all samples were kept  
at -20°C until analysis.

Basal ice from the channel walls and ice from frozen sections of the channel floor were collected at five locations at 50-100 m  
intervals (S1-S5; Figure 2). Ice along the channel floor was generally clear and had the appearance of frozen meltwater, in  
120 contrast to the banded opaque glacier and debris-rich basal ice exposed in the tunnel walls. We expect that stagnant late-season  
runoff occupied the subglacial channel floor at the end of the melt season, and while some of this water remained liquid near  
the endpoint (pond), it froze along the outer section of the channel at the end of the melt season, preserving its microbial and  
biogeochemical characteristics. Since freezing would have propagated from the surface to the bottom of the water column  
introducing a biogeochemical gradient in the ice, bulk samples were collected along the vertical profile of the channel ice at  
125 each site. Channel ice samples were collected into whirl-pak® bags using a flame-sterilized ice axe.




Figure 2: Ice temperature (top) and relative elevation, extent, and slope of the Sverdrup subglacial channel including sample locations of basal ice and channel ice (S1-S5), pond edge ice (S6) and pond slush water and ice (S7-S9) (bottom). Note vertical exaggeration of 50:1.

### 130 2.3 Laboratory analyses

Samples were melted in a laminar flow hood, equipped with a HEPA filter, at  $<4^{\circ}\text{C}$ , in acid-washed (10% HCl for 24 hr), DIW-rinsed and furnace (450°C for 4 hrs) glass beakers, covered with furnace aluminum foil. All glassware was acid-washed, DIW-rinsed, and furnace before use. GF/F filters were also furnace, and non-sterile plastic-ware was acid-washed, DIW-rinsed, and autoclaved before use. Glassware, filter papers, and plastic-ware were all sample-rinsed before aliquots were

135 collected for analysis, as follows:

- A 10 ml sample was collected in a 25 ml glass beaker for immediate measurement of pH using a Fisherbrand™ accumet™ liquid-filled pH/ATC electrode (13-620-530A). One ml was also collected for immediate measurement of electrical conductivity (Oakton®).
- A borosilicate glass filter tower (Fisherbrand™) and 0.7 µm 25 mm GF/F filter paper were used to filter aliquots of sample into two 15 ml centrifuge tubes (Cole-Palmer) that were frozen immediately for NH<sub>4</sub><sup>+</sup> and PO<sub>4</sub><sup>3-</sup> analysis, one 20 ml amber glass EPA vial (Fisherbrand™), that was kept at 4°C with no headspace, was used for spectrofluorescence analysis, and one 40 ml amber glass EPA vial that was amended with 5 M HCl to pH = 2, and stored at 4°C until dissolved organic carbon (DOC) analysis.
- A 10 ml sterile BD Luer Lok syringe and 0.2 µm PES syringe filter (Whatman™ Puradisc) was used to filter aliquots of sample into a 15 ml centrifuge tube (Cole-Palmer), which was stored at 4°C until major ion analysis, a 2 ml Eppendorf™ tube, which was stored at 4°C until Si analysis, and a 2 ml Fisherbrand™ cryogenic storage vial, which was frozen until water isotope analysis.
- For debris-rich (basal ice) samples, 10 ml of sample was collected in a 15 ml Falcon® tube, 5 ml of a cell-sediment separation detergent (Morono et al., 2013) was added and the mixture was vortexed for 1 minute, then centrifuged (500 xg) until the sediment separated. The supernatant (for basal ice samples) or 15 ml of water (for all other samples) was transferred to a new 15 ml Falcon® tube, stained with SYBR™ Gold Nucleic Acid Gel Stain (Invitrogen), and then filtered using a borosilicate glass filter tower (Fisherbrand™) and 0.2 µm MF-Millipore™ filter membrane. Filter papers were mounted on a microscope slide and 20 fields of view were counted at 100x magnification using a Nikon Eclipse 80i microscope.
- Remaining sample was filtered through Pall Supor ® 47 mm filter papers using a Thermo Scientific™ Nalgene filtration kit. DNA was extracted from these filter papers using a FastDNA SPIN Kit for Soil (MP Biomedicals, Irvine, CA, USA) by loading the filter paper directly into the lysis tube then following manufacturer's instructions. The DNA extracts were quantified using a Qubit HS DNA kit (Invitrogen, Waltham, MA, USA). The 16S rRNA gene was amplified in triplicate from each sample with 30 cycles of PCR using modified universal primers (515F (Parada et al., 2016) and 806R (Apprill et al., 2015)). The triplicate reactions were pooled, and Illumina sequencing adapters were added with an additional 8 cycles of PCR, in triplicate. The triplicate adapter reactions were pooled for each sample and purified using a Wizard PCR clean-up system (Promega, Madison, WI, USA) then sequenced at the University of Wisconsin Genomics Core Facility using Illumina MiSeq technology.

## 2.4 Sample measurements

- 165 Water isotopes were measured using a Los Gatos Research Liquid Water Isotope Analyzer (LWIA-45-EP), and calibrated with USGS reference standards. Water isotopes are reported by reference to the Vienna standard mean ocean water (VSMOW), in δ notation (Pinti, 2011). Analytical reproducibility for δ<sup>2</sup>H and δ<sup>18</sup>O was 0.6‰ and 0.4‰, respectively.

170 Samples were analyzed for major ions ( $F^-$ ,  $Cl^-$ ,  $Br^-$ ,  $NO_3^-$ ,  $SO_4^{2-}$ ,  $Na^+$ ,  $Mg^{2+}$ , and  $Ca^{2+}$ ) by ion chromatography using a Metrohm Compact IC Flex ion chromatograph equipped with a C4 cation column and an aSupp5 anion column. Determination of reactive phosphorus and ammonium were measured using principles of spectroscopy following methods outlined by (Strickland and Parsons, 1972) and a Thermo Scientific™ Genesys™ 150 UV-Visible Spectrophotometer with a 10 cm path length quartz cuvette. Silica was also determined using this spectrometer, but with 1 cm path length disposable cuvettes, following methods for heteropoly blue (APHA et al., 2017). Precision and accuracy were better than 5% for all analyses. Non-purgeable DOC was measured by high temperature combustion with a Shimadzu TOC-V (CPH) analyzer equipped with a high sensitivity platinum catalyst. We used fluorescence spectroscopy to characterize fluorescent dissolved organic matter (FDOM) using a HORIBA Aqualog with a quartz cuvette (1 cm path length) and a xenon lamp as an excitation source. Excitation was measured at 5 nm intervals from 230 nm to 600 nm and emission was measured every 2.33 nm from 245 nm to 826 nm, using an integration time of 10 seconds to produce excitation emission matrices (EEMs) for samples and blanks.

## 2.5 Data Analysis

### 180 2.5.1 Parallel factor analysis

We used parallel factor analysis (PARAFAC) to decompose the EEMs and identify/quantify FDOM characteristics (Murphy et al., 2013). PARAFAC was completed using the N-way and drEEM toolboxes in Matlab (The Mathworks Inc., 2020), following methods described by Murphy et al (2013). EEMs were corrected for instrument bias, inner filter effects, and regions of scatter were excised after a DIW blank was subtracted from the measured sample. Components derived from PARAFAC modeling were compared to those identified in other studies using the online spectral library of auto-fluorescence by organic compounds in the environment, OpenFluor (Murphy et al., 2014).

### 2.5.2 Bioinformatics

190 The 16S rRNA gene amplicon sequences were processed using *mothur* (Kozich et al., 2013) to filter reads that did not meet minimum quality control thresholds (maxambig=0, maxlength=315, maxhomop=8), join paired reads, cluster sequences into operational taxonomic units (OTUs) using 97% sequence identity, and assign taxonomic identification to OTUs using the SILVA SSU database v138. A total of 2868 singleton OTUs, or those that appeared once across the entire dataset, were removed prior to further analysis. Diversity calculations were completed using the phyloseq package (v 1.28.0) in R (v 3.6.0).

### 2.5.3 Water isotope model

195 Using the principles of isotopic fractionation, a model was developed to estimate the isotopic composition of incremental ice, incremental vapor, and residual water as a hydraulically isolated waterbody progressively freezes and evaporates (Supplementary Methods S1). The degree to which isotopes fractionate during freezing depends on the rate of freezing and can be described by Eq. (1):

$$\varepsilon_{I-W} = \delta_I - \delta_W = 1000(\alpha - 1) \quad (1)$$

Where  $\delta_I$  and  $\delta_W$  are the isotopic composition of the ice and water, respectively,  $\varepsilon_{I-W}$  is the isotopic difference between ice and water, and  $\alpha$  is the respective equilibrium fractionation factor. The degree to which isotopes fractionate during evaporation ( $\varepsilon_{W-V}$ ) depends on the temperature of the water during evaporation and can also be described by Eq. (1) using the isotopic composition of water and vapor instead of ice and water, respectively.

The isotope model equations are described in detail in Supplementary Methods S1 and were processed in Matlab (The Mathworks Inc. 2020) using custom scripts. Fixed variables included: 1) the initial isotopic composition of the waterbody, which was set to the average isotopic composition of channel ice samples, 2) equilibrium values for evaporation, which were set to experimental values obtained for evaporation at 0°C (Majoube, 1971), 3) equilibrium values for freezing, which were set to the average of the isotopic fractionation between ice and water in pond slush, and 4) the step interval, which was set to simulate 0.1% of the waterbody freezing in each step of the model. We adjusted the rate of evaporation vs freezing, to optimize the model's fit with the observed  $\delta^2\text{H}$ - $\delta^{18}\text{O}$  of channel ice and pond ice along the incremental ice line, and the pond water samples along the residual water line (Figure S2). We then fixed the evaporation vs freezing rate and ran the model to estimate the stage of freezing (i.e. the fraction of residual water vs frozen and evaporated water) at each sample site in the pond. To do so, the observed  $\delta^2\text{H}$ - $\delta^{18}\text{O}$  of ice and water samples at each site in the pond were compared to the modeled incremental ice line and the modeled residual water line, respectively. The step with the most similar  $\delta^2\text{H}$ - $\delta^{18}\text{O}$  was identified for each site and the corresponding stage of freezing was determined.

#### 2.5.4 Cl<sup>-</sup> model

Cl<sup>-</sup> does not readily precipitate as salts, interact with rocks, or become assimilated in significant quantities by microorganisms (Davis et al., 1998). Since there were no identifiable sources or sinks for Cl<sup>-</sup> in the system, we used the principles of solute fractionation to develop a model that estimates the [Cl<sup>-</sup>] in incremental ice and residual water as an isolated waterbody progressively freezes and evaporates. Cl<sup>-</sup> is largely rejected from the ice crystal lattice during freezing (Killawee et al., 1998; Clayton et al., 1990) and from water vapor during evaporation, and thus it concentrates in the residual water. Though effectively all non-volatile solutes are excluded from water vapor, the degree to which solutes are excluded from the ice is described by the effective segregation coefficient ( $K_{\text{eff}}$ ) in Eq. (2):

$$K_{\text{eff}(X)} = \frac{[X]_{\text{ice}}}{[X]_{\text{water}}} \quad (2)$$

Where  $K_{\text{eff}(X)}$  is the effective segregation coefficient for biogeochemical species  $X$ , and  $[X]_{\text{ice}}$  and  $[X]_{\text{water}}$  are the corresponding concentrations of that species in coincident ice and water samples.

The Cl<sup>-</sup> model equations are described in detail in Supplementary Methods S2 and were processed in Matlab (The Mathworks Inc. 2020) using custom scripts. Fixed variables included: 1) the initial [Cl<sup>-</sup>] of the waterbody, which was set to the average [Cl<sup>-</sup>] of channel ice (Simulation 1) or with the addition of Cl<sup>-</sup> affiliated when 5% (Simulation 2) or 15% (Simulation 3) of the initial solute is comprised of basal solute sources (represented by the average solute concentrations in basal ice samples), 2)

230  $K_{\text{eff}}$ , which was derived from coincident ice and water samples in the pond slush using Eq. (2), 3) the step interval, which was set to simulate 0.1% of the waterbody freezing in each step of the model, and 4) the rate of evaporation vs freezing, which was set to the ratio derived from isotope modeling.

Results from the  $\text{Cl}^-$  model were used to estimate the stage of freezing (i.e. the fraction of residual water vs frozen and evaporated water) at each sample site in the pond (Figure S2). To do so, the observed  $[\text{Cl}^-]$  in ice and water at each pond sample site were compared to modeled  $[\text{Cl}^-]$  in incremental ice and residual water, respectively. The step with the most similar  $[\text{Cl}^-]$  was identified for each site and the corresponding stage of freezing was determined.

### 2.5.5 Biochemical model

Similar to  $\text{Cl}^-$ , other dissolved impurities are excluded from water vapor during evaporation and the extent to which they are also rejected from the ice crystal lattice during freezing can be quantified using Equation 2 (Killawee et al., 1998; Clayton et al., 1990). We therefore used the same theory and equations as the  $\text{Cl}^-$  model to produce a biogeochemical model (using custom scripts in Matlab (The Mathworks Inc. 2020)) that simulates the concentration of other solutes and impurities (e.g. cells) in residual water and incremental ice (Supplementary Methods S2; Figure S2) at the stage of freezing affiliated with each sample site (determined from the  $\text{Cl}^-$  model). We then compared measured concentrations at each pond sampling site to modeled concentrations. Since the model only simulates the effects of freezing and evaporation, we interpreted  $[X]_{\text{measured}} > [X]_{\text{modeled}}$  as evidence for potential net sources of chemical species 'X' and  $[X]_{\text{measured}} < [X]_{\text{modeled}}$  as evidence for potential net sinks of chemical species 'X'.

## 3 Results

### 3.1 Physical system

The ponded water from which samples were retrieved was located at the endpoint of a remnant subglacial tunnel that extended 467 m horizontally into the glacier at an angle of  $\sim 18^\circ$  relative to the direction of ice flow (NNW; Figure 1). The location of the ponded water was therefore  $\sim 150$  m in perpendicular distance from the margin and  $\sim 79$  m below the glacier surface (Figure 1). Along its 467 m length, the channel floor dropped 5.3 m in elevation (Figure 2) and terminated at an ice-wall that is located near the edge of a subglacial 'cliff', which drops almost 200 m over a horizontal distance of  $\sim 150$  m towards the glacier centerline (Figure 1c). The glacier bed reaches a maximum depth of -190 m a.s.l. (or  $\sim 300$  m below the glacier surface). The ambient air temperature at the surface of the glacier remained  $< 0^\circ\text{C}$  from mid-September 2018 until fieldwork in May 2019, with temperatures falling to  $< -30^\circ\text{C}$  in 6 of these months (Figure S1). At the time of the field survey, ice temperatures along the channel wall increased from  $-21^\circ\text{C}$  at the tunnel entrance to  $-0.4^\circ\text{C}$  at the endpoint, indicating the presence of a subglacial heat source. The channel floor was completely frozen from the entrance to  $\sim 405$  m (Figure 2). Liquid water was observed on the channel floor from  $\sim 405$  m and transitioned into a slushy pond, which extended to the endpoint at 467 m. A higher water:ice ratio was observed in the pond with distance towards the endpoint. The surface of the slushy pond increased

in elevation by 0.4 m towards the endpoint (Figure 2). Hydraulic head at the ice-bed interface in the vicinity of the pond was higher than at any other point across the glacier cross-section (Supplementary Methods S3 and Figure S3).

### 3.2 Water isotopes

265 The isotopic composition of ice samples from the channel floor (S1-S5) lie below the local meteoric water line (LMWL) for the Sverdrup catchment (Copland et al., 2021), which is consistent with the isotopic composition of meltwater originating from local snow/glacier ice. Channel pond and basal ice samples fall above the LMWL and exhibit more negative  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  values relative to the channel ice. The difference between respective ice and water samples at sites 8 and 9 was 1.8-1.9 ‰ for  $\delta^{18}\text{O}$  and 10.9 ‰ for  $\delta^2\text{H}$ , which is consistent with equilibrium values measured in other slow-freezing environments (Table S1) (Jouzel et al., 1999; Ferrick et al., 2002).

270 Our isotope modelling experiments indicate that a freeze:evaporation rate of ~40:1 was required to best fit the model to the observed water and ice data points (Figure 3a). This model also indicates that ~10% of the initial water remained residual at pond sampling sites 8 and 9 (Figure 3c). The  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$  values at Site 7 show that a similar fraction of water remained residual. The  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$  values at Site 6 showed the most extensive freezing, with only 2% remaining as residual water (Figure 3c).

### 275 3.3 Cl<sup>-</sup> concentrations

An average  $K_{\text{eff}(\text{Cl}^-)}$  value (Equation 2) of 0.22 was derived from coincident pond ice (2.0  $\mu\text{M}$  and 2.2  $\mu\text{M}$ ) and water (15.8  $\mu\text{M}$  and 15.7  $\mu\text{M}$ ) (Figure 4) samples at sites S8 and S9, respectively. This  $K_{\text{eff}(\text{Cl}^-)}$  falls within the range observed in other studies (Santibáñez et al., 2019). The average  $[\text{Cl}^-]$  in channel ice samples was 2.1  $\mu\text{M}$ , which is consistent with concentrations in regional glacier ice and snow (Dubnick et al., 2020). Cl<sup>-</sup> fractionation modeling (Supplementary Methods S1) of water with 280 initial  $[\text{Cl}^-]$  of 2.1  $\mu\text{M}$  suggests that ~9% of the water in the pond at S8 and S9 remained as residual to yield the  $[\text{Cl}^-]$  measured at these sites. These results are in close agreement (within 1%) with corresponding results from the isotope fractionation model at S8 and S9 (Figure 3c). Cl<sup>-</sup> comprised a very small portion of the solutes in basal ice, so simulations that included contributions of basal solutes (up to 15% of the solute load) had little effect (<0.1%) on the estimated stage of freezing at each site.






Figure 3: (a)  $\delta^2\text{H} - \delta^{18}\text{O}$  plot showing the modelled residual water and incremental ice as water progressively freezes and evaporates, as well as measured concentrations for channel ice and pond samples. LMWL is from Copland et al. (2021) (b) modelled  $\text{Cl}^-$  concentrations in residual water and incremental ice as water progressively freezes and evaporates, as well as measured concentrations for channel ice and pond samples fit to respective lines, and (c) the average corresponding percent frozen and evaporated for ice and water samples at sampling sites S6, S7, S8 and S9, derived from the isotope model in (a) and the  $[\text{Cl}^-]$  model in (b). Error bars indicate the minimum and maximum modelled results for the ice and water samples at each location.






Figure 4: Biogeochemical content of basal ice, channel ice and pond water/ice. Error bars represent  $\pm$  root sum of squares of precision and accuracy for solute measurements,  $\pm$  root sum of the squares of the errors in the quantities being added or subtracted to derive total solutes and  $\text{HCO}_3^-$ , and  $\pm 1$  standard deviation for cell counts. Error for C1 and C2 is not quantified due to the nature of the data.

### 3.4 Major ion chemistry


Channel ice samples from S1-S5 were generally dilute ( $\bar{x} = 0.91 \text{ mg L}^{-1}$ ) but displayed considerable variability in the concentration of total measured solutes. Pond ice samples (at sites 8-9) had similar total measured solute concentrations (1.1 and 1.2  $\text{mg L}^{-1}$ ) but coincident pond water samples had concentrations ~7-fold higher (7.0  $\text{mg L}^{-1}$  and 7.2  $\text{mg L}^{-1}$ , respectively). Pond edge ice (site 6) had the highest total measured solute load of all channel samples (11.6  $\text{mg L}^{-1}$ ). Basal ice had highly variable total measured solute concentrations (1.5-5.4  $\text{mg L}^{-1}$ ) (Figure 4).  $K_{\text{eff}}$  values derived using coincident ice and water samples at sites S8 and S9 are  $<0.3$  for each major ion, and consistent with those reported by Santibanez et al. (2019) (Figure S4).

Biogeochemical modeling (Supplementary Methods S2) using late season runoff as the sole solute source (Simulation 1) underestimated the concentration of many of the dominant solutes in pond water, including  $\text{HCO}_3^-$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$  and  $\text{SiO}_2$  (Figure 5a). The solute: $\text{Cl}^-$  ratios for all biogeochemical species were greater in basal ice than in channel ice (Figure 5b), so simulations that also incorporated basal solute contributions to the pond (comprising 5-15% of the initial solute load; Simulations 2 and 3) generally yielded more accurate concentrations of these solutes (Figure 5b).

### 3.5 FDOM components

PARAFAC modeling resolved two fluorescent components that comprised 92 % of the variability in the dataset (Figure S5). Component 1 (C1) overlaps with “humic-like” fluorescence (Coble, 1996) and has been associated with fulvic acid DOM fractions (Ohno and Bro, 2006). While this fluorescence feature is not consistently identified in glacier ice, similar fluorescence signatures are ubiquitous in a wide range of terrestrial (Ohno and Bro, 2006; Yamashita et al., 2011) and marine (Jørgensen et al., 2011; Retelletti Brogi et al., 2018; Walker et al., 2009) environments. Further, this fluorescence component has been identified in Antarctic watersheds without higher plants (Cory and McKnight, 2005; Barker et al., 2013) and has been correlated with microbial activity in ocean waters (Jørgensen et al., 2011) indicating that it may also be produced by microbial reworking of organic matter (OM). Further, this fluorophore is persistent in many environments suggesting that, once formed, it may not be readily altered (Yamashita et al., 2010).

Component 2 (C2) overlaps with the fluorescence of amino acids, in particular as either free or bound tyrosine (Peak B, Coble, 1996). Similar FDOM is widely detected in glacial ice, including englacial and basal ice from Greenland, Antarctica, and across the Canadian Arctic from the supraglacial snowpack to ice dating back to the Last Glacial Maximum (D’Andrilli and McConnell, 2021; D’Andrilli et al., 2017; Pautler et al., 2012; Dubnick et al., 2010, 2020; Barker et al., 2006, 2009; Fellman et al., 2010). Tyrosine-like fluorescence is commonly associated with OM of low molecular weight and aromaticity and chemical species that are readily degraded by microorganisms (Coble, 2014). Unlike C1, C2 is rapidly transformed in proglacial and downstream aquatic systems (Wu et al., 2003; Saadi et al., 2006; Fellman et al., 2008; Barker et al., 2013).




360 Figure 5: (a) Biogeochemical modelling results compared to observed concentrations in pond samples. Bars represent simulations that assume channel water as the sole water and solute source (Simulation 1) while error bars represent simulations with an additional 5-15% solutes from basal ice (Simulations 2 and 3). (b) provides a measure of the relative abundance of the different solutes in basal ice compared to channel ice according to the formula  $y = \frac{X_{BI} \cdot Cl_{BI}}{X_{CI} \cdot Cl_{CI}}$  where X is the average concentration of an analyte in either basal ice (BI) or channel ice (CI) and Cl<sup>-</sup> is the average chloride concentration, and y is the relative abundance of that analyte. Error bars represent 0.1 standard deviations.

365

### 3.6 Nutrients

Channel ice samples contained relatively low concentrations of organic and inorganic nutrients, including NO<sub>3</sub><sup>-</sup> ( $\bar{x}$  = 0.68 μM), NH<sub>4</sub><sup>+</sup> ( $\bar{x}$  = 0.31 μM), PO<sub>4</sub><sup>3-</sup> ( $\bar{x}$  = 0.03 μM), and DOC ( $\bar{x}$  = 5.8 μM) (Figure 4). Concentrations of these nutrients were all higher in the pond water than in coincident pond ice samples, yielding K<sub>eff</sub> values < 1 (Figure S4). Compared to modeled concentrations, pond waters were depleted in all nutrients (i.e., NH<sub>4</sub><sup>+</sup>, PO<sub>4</sub><sup>3-</sup>, DOC, and DOM components C1 and C2) except NO<sub>3</sub><sup>-</sup> (Figure 5). Basal ice had higher nutrient:Cl<sup>-</sup> ratios than did the channel ice, especially in the case of PO<sub>4</sub><sup>3-</sup> and NH<sub>4</sub><sup>+</sup> (Figure 5b). Consequently, simulations that involve basal solute contributions to the pond (Simulations 2 and 3; Figure 5b) increase the amount by which these nutrients appear depleted in pond water (Figure 5a).

375 Cell concentrations in channel ice ranged from 1.6 x 10<sup>4</sup> to 11.2 x 10<sup>4</sup> while concentrations in the pond water ranged from 6.3 x 10<sup>4</sup> to 11.9 x 10<sup>4</sup> cells mL<sup>-1</sup> (Figure 4), comparable to the range found in other subglacial waters (Christner et al., 2014; Gaidos et al., 2004; Mikucki et al., 2009). Over 85% of the 16S rRNA gene sequences in pond water samples were affiliated with one of six OTUs that fell in the class Gammaproteobacteria: *Massilia* (28-40% of total reads), two *Polaromonas* OTUs (25-27%), an unclassified Oxalobacteraceae (14-18%), *Rhodoferax* (5-15%), and *Undibacterium* (3%) (Figure 6). While these OTUs (except the unclassified Oxalobacteraceae) were all detected in basal ice, they were more abundant in channel ice samples (including the unclassified Oxalobacteraceae), comprising 23-32% of the assemblage (Figure 6). Chao1, Shannon's (H) and Simpsons (D) diversity indices all indicate taxonomic diversity was lower in pond water than in channel ice or basal ice (Figure 6).



385 Figure 6: (a) Chao1, (b) Shannon's and (c) Simpson's diversity index for microbial assemblages in basal ice, channel ice and pond water samples. (d) The composition of microbial assemblages detected in each sample. Assemblages are grouped by phylum, except for the 6 most dominant OTUs in the pond water, which were all Gammaproteobacteria and are further classified here by genus (i.e. *Massilia*, *Polaromonas* (x2), an unclassified Oxalobacteraceae, *Rhodoferrax*, and *Undibacterium*). Insufficient sample was available for Channel Ice S2, Pond Edge S6, and Pond Water S7.

**4.1 Evolution of the subglacial tunnel and pond**

Subglacial channels develop when inflowing waters have sufficient heat and pressure to melt ice at a rate that exceeds that of creep closure from the overlying ice (Röthlisberger, 1972; Nye, 1976). These channels grow during spring-summer to deliver meltwater to the glacier terminus and are closed by ice creep as runoff rates recede in the fall. Closure rates are typically lower  
395 near the ice margin where overlying ice is thin and overburden pressure is low and are faster at internal locations where overlying ice is thick and overburden pressure is high (Röthlisberger, 1972). At the time of the field survey, the subglacial tunnel terminated at approximately the upper edge of a ‘cliff’, or rapid deepening along the glacier bed profile (Figure 1). Rates of creep closure would be much higher at internal locations beyond this subglacial ‘cliff’ due to the ice overburden pressure (Figure S3). Creep closure near the pond was likely responsible for squeezing the subglacial pond slush into a mound,  
400 resulting in the positive slope of the pond surface in this area at the end of winter (Figure 2).

Most of the melt generated on the Sverdrup Glacier drains ice-marginally and both the water isotopes and geochemistry of channel ice samples indicate snow and glacier ice as predominant water sources. We have no evidence to suggest that late season runoff draining into this subglacial tunnel originated from an upstream subglacial drainage network, but recognize subglacial water could contribute to late season water, for example, from melting basal ice along the glacier margin, or  
405 subglacial drainage from upstream. As ambient air temperatures fell towards the end of the melt season of 2018 (remaining below 0°C from September 2018 – June 2019; Figure S1) drainage into the subglacial channel from the ice margin would have ceased, leaving ponded water along the channel floor in areas with low slope. The cold, dense ambient air drained into the tunnel, progressively freezing the stagnant water on the channel floor. Although the freezing process along the channel floor may have released some latent heat, additional heat source(s) would have been required to maintain liquid water and the near  
410 0°C ice and air temperatures that were observed at the tunnel endpoint in May. The Sverdrup Glacier is marine-terminating ~3 km downstream of the subglacial tunnel and displays a retrograde subglacial slope (Paden et al., 2019), resulting in ice that is grounded below sea level across most of the glacier’s width near the subglacial tunnel (Figure 1). This geometry allows heat from the ocean to more effectively warm areas of the glacier bed. Additional heat may also originate from friction near the glacier bed. Prior studies suggest regions of the Sverdrup Glacier have flow regimes that involve enhanced deformation of  
415 basal ice (Van Wychen et al., 2017), which produce frictional heat. Geothermal heat may also contribute to the subglacial heat flux but the regional geothermal heat flux is relatively low (65 +/- 5 m W m<sup>-2</sup>; (Grasby et al., 2012)).

**4.2 Concentration effects**

The water isotope and Cl<sup>-</sup> models produced consistent estimates of the extent of freezing at the pond sampling sites (Figure 3c). The portion of residual water increased with distance into the pond, from 6% remaining as residual water at the pond edge  
420 (S6) to 9% remaining as residual water at S9. Sites S6 and S7 are at lower elevations than S8 and S9 (Figure 2), and would have experienced colder overlying air in this depression. Freezing processes (and to a lesser degree, evaporation) were,

therefore, the dominant controls on the concentration of solutes in the pond, resulting in concentrations up to 7 times those observed in late-season runoff.

425 The temperature gradient along the downward-sloping subglacial tunnel (Figure 2) facilitates the drainage of cold, dense ambient air into the tunnel and drainage of warm buoyant air out of the tunnel. Regional air masses are cold and have low relative (65%) and specific ( $0.2 \text{ g kg}^{-1}$ ) humidity during the winter (Vincent et al., 2007). The convection and warming of this ambient air through the tunnel would therefore facilitate continual evaporation from the subglacial pond throughout the cold season. The tunnel showed evidence of evaporation; the roof/upper section of the walls were covered in a blanket of intricate ice crystals formed by condensation of water vapor as the buoyant/warm/moist air mass traveled out of the tunnel and cooled.

430 During evaporation, solutes are preferentially excluded from the vapor phase, so evaporation at the pond surface increased solute concentrations in the residual water.

The drainage of cold, dense air along the tunnel floor would also have frozen the late-season runoff pooled on the channel floor, from the surface to the bed, and from the channel entrance towards the endpoint. The pond was comprised of a slushy mixture of ice and water, indicating a slow freezing rate (Michel and Ramseier, 1971) near the tunnel endpoint, and isotope modeling suggests that freezing occurred at a rate  $\sim 40$  times that of evaporation. Suspended and dissolved impurities and light water isotopes are preferentially rejected from ice crystals during the freezing process, resulting in residual water that becomes increasingly concentrated in impurities (Clayton et al., 1990; Killawee et al., 1998) and isotopically light.

435

### 4.3 In-situ geochemical sources

Unlike the waters contained beneath many other polythermal glaciers, our biogeochemical modeling simulations suggest that the pond only received small ( $\sim 5$ -15%) contributions of solutes from basal sources. As discussed above, the channel floor ice and pond water originated as late-season runoff that pooled along the channel floor in areas of low slope. Compared to biogeochemical modeling that simulates late-season runoff as the sole solute and water source to the pond, pond water samples were enriched in total solutes, including  $\text{SO}_4^{2-}$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  (Figure 5a) and Si. In glacial environments, these elements are commonly derived from rock-water reactions (Tranter et al., 2002) and are found at high concentrations in basal ice (Dubnick et al., 2020) and subglacial discharge (Wadham et al., 2010; Li et al., 2022). Importantly, these solutes were enriched in basal ice samples in this study (Figure 5b), suggesting they are liberated from basal processes.

440

445

Basal solute contributions to the pond most likely originated as seepage from basal ice along the pond perimeter. The interstitial water content of basal ice scales with temperature, so the relatively warm ice in this area (Figure 2) could promote drainage of solute-rich water into the pond. Though the pond could also receive basal solutes from a distributed drainage system at the ice-bed interface, hydraulic modeling suggests that if waters occupied a distributed drainage system, they were more likely to drain away from the pond than towards the pond (Figure S3). The pond could also acquire basal solutes by rock-water interactions at its base. However, this channel receives high volumes of seasonal meltwater over consecutive years, so reactive components of the bedrock would be depleted (including those with slow reaction kinetics).

450

455 Simulations that incorporated basal solutes improved the accuracy of modeled solute concentrations, though the accuracy of the models were inconsistent among geochemically-relevant compounds (Figure 5a). The solute composition of basal ice, subglacial water, and species liberated from *in situ* rock weathering, can vary dramatically within a subglacial catchment (Dubnick et al., 2020; Yde et al., 2010; Tranter et al., 2002) due to variation in the composition of the underlying bedrock, redox conditions, ice/water temperatures, and freeze-thaw histories. The five basal ice samples explored in this study also showed high variability in solute chemistry (Figure 4), confirming that potential basal solute sources in the subglacial tunnel  
460 are heterogeneous and that our model may not accurately represent the precise composition of the basal solutes entering the pond.

#### 4.4 Microbiology and biogeochemical nutrient sinks

Sequencing of 16S rRNA gene amplicons indicates that the microbial assemblage in the subglacial pond was dominated by organisms in the order Burkholderiales, which are commonly found in subglacial systems (e.g. Cheng and Foght, 2007; Achberger et al., 2016; Dubnick et al., 2020). More specifically, the most dominant OTUs, *Massilia*, *Polaromonas*, *Rhodoferrax* and *Undibacterium* (Figure 6) are all related to microbes detected at high abundance in other glacial environments (Foght et al., 2004; Lanoil et al., 2009; Darcy et al., 2011; Mitchell et al., 2013; Perini et al., 2019; Dubnick et al., 2020; Dunham et al., 2021) including supraglacial, proglacial, and/or subglacial aquatic systems worldwide. Moreover, these OTUs are related to psychrophilic or psychrotolerant species (e.g. Foght et al., 2004; Darcy et al., 2011; Wang et al., 2018; Perini et al., 2019), and  
470 many of these taxa have been shown to have a range of structural and functional adaptations for survival in cold temperatures (Margesin and Miteva, 2011).

All six dominant OTUs in the pond were closely related (>98%) to globally distributed environmental sequences and cultured representatives, suggesting these organisms are habitat generalists with broad environmental tolerances. For example, *Polaromonas* are dominant in polar and high-elevation environments (Darcy et al., 2011), are thought to rapidly evolve to new environments through extensive horizontal gene transfer (Yagi et al., 2009), and are considered metabolically diverse  
475 “opportunitrophs” (Meyer et al., 2004; Polz et al., 2006). The meltwaters that drained into the subglacial system towards the end of the melt season originated from cryoconite holes, supraglacial streams, englacial ice, precipitation, and extraglacial aquatic or terrestrial sources, and potentially upstream subglacial sources (e.g. basal ice melt from ice marginal cliffs or subglacial water drainage). These environments are exposed to distinct nutrient pools, redox conditions, and, in some cases,  
480 high levels of solar radiation and/or warmer temperatures. The shift in environmental conditions to those in the cold, dark, oligotrophic subglacial pond may have decimated populations not capable of survival in this system, selecting for generalist organisms that are better adapted to those conditions (Xu et al., 2021), and yielding lower cell concentrations in the pond than the models predicted (Figure 5a).

Consistent with the notion of being habitat generalists, the six dominant OTUs in the pond water are closely related to genera  
485 with extreme metabolic versatility. For example, despite sharing nearly identical 16S rRNA genes, *Polaromonas* strains from different glaciers can have very different phenotypes (Gawor et al., 2016), with some strains variably showing evidence of an

ability to oxidize H<sub>2</sub> (Sizova and Panikov, 2007; Dunham et al., 2021), arsenite (Osborne et al., 2010), and/or organic matter (Jeon et al., 2003; Mattes et al., 2008; Osborne et al., 2010). Furthermore, several cold-tolerant species of *Rhodoferrax* (e.g. *R. antarcticus* and *R. fermentans*) have been shown to be facultative anoxygenic phototrophs that can grow photoheterotrophically (using a variety of organic and fatty acids or glucose) and photoautotrophically (using CO<sub>2</sub>) or that can grow via aerobic respiration in dark environments (Madigan et al., 2000). *R. fermentans* has also been shown to be capable of nitrate reduction (Hougardy and Klemme, 1995) and a close relative of *R. ferrireducens* from glacial sediments in Iceland has been shown to fix CO<sub>2</sub> with energy derived from the H<sub>2</sub>-ferric iron redox couple (Dunham et al., 2021). The high abundance of *Rhodoferrax* in the pond may be due to its metabolic flexibility which allows it to compete for nutrients despite their changing availabilities as they are transported from the supraglacial/extraglacial to the subglacial system, and in the subglacial pond as it becomes increasingly cold, concentrated in solutes, and depleted in labile nutrients over the winter.

Our microbial and biogeochemical datasets both suggest the pond functioned as a hotspot for C-cycling over winter. We measured ~43% less DOC than our modeling predicted and a more depleted reservoir of the labile tyrosine-like (C2) fluorophore, which is often rapidly transformed in proglacial and downstream aquatic systems (Wu et al., 2003; Saadi et al., 2006; Barker et al., 2013; Fellman et al., 2010) (Figure 5). Previous studies of *Polaromonas*, *Rhodoferrax*, and *Massilia* strains, which comprised ~70% of the pond microbial assemblage, suggest they are fuel their metabolisms by coupling the oxidation of organic matter with reduction of O<sub>2</sub> (Hougardy and Klemme, 1995; Jeon et al., 2003; Mattes et al., 2008; Osborne et al., 2010), and thus could be responsible for depletion of DOM. Assuming 70% of the cells in the pond were active heterotrophs ( $\bar{x} = 7.5 \times 10^4$  cells mL<sup>-1</sup>), and that the size of this population was relatively stable throughout the ~240 days over winter, the observed DOC deficit (3.9 μM) equates to an average rate of C-remineralization of 0.2 fmol C cell<sup>-1</sup> day<sup>-1</sup>. This rate is comparable to metabolic rates sufficient for microbial growth at ~0°C and is orders of magnitude higher than metabolic rates required for cellular maintenance or survival at 0°C (Price and Sowers, 2004). Though this is only an estimate for C-remineralization in the pond, rates could be higher if less than 70% of the microbial assemblage was engaged in heterotrophic energy metabolisms, or if additional OC pools were also degraded, such as particulate OM, basally-derived OM, or autochthonous OM (our calculation above only accounts for DOC supplied in the late-season runoff).

Microbial activity in the subglacial pond could also explain the depleted reservoir of other inorganic nutrients, including PO<sub>4</sub><sup>3-</sup> and NH<sub>4</sub><sup>+</sup> (Figure 5). Past work has shown that heterotrophic microbial biomass is C-poor yet P- and N-rich (Makino et al., 2003; Godwin and Cotner, 2015) relative to many terrestrial DOM sources, and that microbial heterotrophic activity has been linked to a simultaneous assimilation of mineral nutrients (Fenchel and Blackburn, 1979; Martinussen and Thingstad, 1987). While PO<sub>4</sub><sup>3-</sup> is considered to be a preferred and universal source of phosphorus to microbes (Björkman and Karl, 1994), N can be assimilated as NO<sub>3</sub><sup>-</sup>/NO<sub>2</sub><sup>-</sup> or as the preferred reduced state, NH<sub>4</sub><sup>+</sup>/NH<sub>3</sub> (Paul and Clark, 1996; Nyysönen et al., 2014). Lithoautotrophic activity can contribute to microbial assimilation of inorganic nutrients, including NH<sub>4</sub><sup>+</sup> and PO<sub>4</sub><sup>3-</sup>, as has been demonstrated from laboratory incubations of subglacial sediments and water (Stibal et al., 2012b; Montross et al., 2014; Boyd et al., 2011).

## 520 **5 Conclusions**

This research documents the evolution of ponded meltwater and its resident microbial community at the endpoint of a 467 m long remnant subglacial channel through a Canadian Arctic winter. Solute concentrations in the pond were controlled by: 1) freezing processes, which functioned to cryo-concentrate solutes in the residual water by up to 7 times; 2) seepage of small amounts of basal solutes (comprising <15% of the total solutes) into the pond; and 3) microbial activity, which functioned to  
525 deplete the pond's reservoir of the most labile and biogeochemically-relevant compounds, including  $\text{NH}_4^+$ ,  $\text{PO}_4^{3-}$ , and DOM. Sequencing of the 16S rRNA gene revealed decreasing taxonomic diversity among microbial communities with distance into the channel. Six OTUs dominated the microbial community in the pond. These microorganisms likely originated from the extraglacial or supraglacial (rather than subglacial) environment and were related to taxa that are psychrophilic/psychrotolerant, exhibit extreme metabolic diversity, and have broad habitat ranges. Collectively, our findings  
530 suggest that generalist microorganisms from the extraglacial or supraglacial environments can become established in subglacial aquatic systems and deplete reservoirs of nutrients over a period of months. The inferred generalist lifestyle of these microorganisms may help them survive the extreme selection pressures imposed by the environment, allowing for not only their persistence but activity. These findings extend our understanding of the microbiology and biogeochemistry of subglacial ecosystems.

## 535 **Data Availability**

Sequence data has been deposited in NCBI SRA under BioProject ID PRJNA907039 and biogeochemical data are available via Zenodo DOI 10.5281/zenodo.7384156.

## **Author contribution**

AJD designed and carried out the fieldwork, laboratory work, data analysis, developed the model codes and performed the  
540 simulations. RLS extracted DNA from field samples and conducted the bioinformatics, BDD prepared temperature data record, DB prepared radar and laser altimeter transect data, and CD conducted laboratory DOC analysis. AJD, RLS and BDD prepared the manuscript with contribution from all co-authors. MJS, MLS, ESB funded and supervised the work.

## **Competing Interests**

The authors declare that they have no conflict of interest.

## 545 **Acknowledgements**

We thank the Nunavut Research Institute and the communities of Grise Fjord and Resolute Bay for permission to conduct research on Devon Island, Patrick Williams and Claire Bernard-Grand'Mason for field assistance, the staff at the Polar Continental Shelf Program for field logistics support, Maya Bhatia for laboratory facilities for DOC analysis at the University of Alberta Molecular Biogeochemistry Lab, Montana State University Environmental Analytical Lab for water isotope  
550 measurements, and the University of Wisconsin Genomics Core Facility for 16S rRNA gene sequencing.

This research was supported by the Natural Sciences and Engineering Research Council of Canada (Discovery grant no. 05234-2015), the Polar Continental Shelf Program (grant no. 68519), and the National Aeronautics and Space Administration (NNX16AJ64G and 80NSSC20K1134).

## 555 **References**

- Achberger, A. M., Christner, B. C., Michaud, A. B., Priscu, J. C., Skidmore, M. L., Vick-Majors, T. J., and the WISSARD Science Team: Microbial community structure of subglacial Lake Whillans, West Antarctica, *Front. Microbiol.*, 7, 1–13, <https://doi.org/10.3389/fmicb.2016.01457>, 2016.
- Andrews, M. G., Jacobson, A. D., Osburn, M. R., and Flynn, T. M.: Dissolved Carbon Dynamics in Meltwaters From the  
560 Russell Glacier, Greenland Ice Sheet, *J. Geophys. Res. Biogeosciences*, 123, 2922–2940, <https://doi.org/10.1029/2018JG004458>, 2018.
- APHA, AWWA, and WEF: Standard Methods for the Examination of Water and Wastewater, 22nd ed., edited by: Rice, E. W., Baird, R. B., Eaton, A. D., and Clesceri, L. S., American Public Health Association, American Water Works Association, Water Environment Federation, 1–5 pp., 2017.
- 565 Apprill, A., McNally, S., Parsons, R., and Weber, L.: Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton, *Aquat. Microb. Ecol.*, 75, 129–137, 2015.
- Barker, J. D., Sharp, M. J., Fitzsimons, S. J., and Turner, R. J.: Abundance and dynamics of dissolved organic carbon in glacier systems, *Arctic, Antarct. Alp. Res.*, 38, 163–172, [https://doi.org/10.1657/1523-0430\(2006\)38\[163:AADODO\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2006)38[163:AADODO]2.0.CO;2), 2006.
- Barker, J. D., Sharp, M. J., and Turner, R. J.: Using synchronous fluorescence spectroscopy and principal components analysis  
570 to monitor dissolved organic matter dynamics in a glacier system, *Hydrol. Process.*, 23, 1487–1500, <https://doi.org/10.1002/hyp>, 2009.
- Barker, J. D., Klassen, J. L., Sharp, M. J., Fitzsimons, S. J., and Turner, R. J.: Detecting biogeochemical activity in basal ice using fluorescence spectroscopy, *Ann. Glaciol.*, 51, 47–55, <https://doi.org/10.3189/172756411795931967>, 2010.
- Barker, J. J. D., Dubnick, A., Lyons, W. B., and Chin, Y.: Changes in dissolved organic matter (DOM) fluorescence in  
575 proglacial Antarctic streams, *Arctic, Antarct. Alp. Res.*, 45, 305–317, <https://doi.org/10.1657/1938-4246-45.3.305>, 2013.
- Bhatia, M. P., Kujawinski, E. B., Das, S. B., Breier, C. F., Henderson, P. B., and Charette, M. A.: Greenland meltwater as a

- significant and potentially bioavailable source of iron to the ocean, *Nat. Geosci.*, 6, 1–5, <https://doi.org/10.1038/ngeo1746>, 2013.
- Bingham, R. G., Nienow, P. W., Sharp, M. J., and Copland, L.: Hydrology and dynamics of a polythermal (mostly cold) High Arctic glacier, <https://doi.org/10.1002/esp>, 2006.
- Björkman, K. M. and Karl, D. M.: Bioavailability of inorganic and organic phosphorus compounds to natural assemblages of microorganisms in Hawaiian coastal waters, *Mar. Ecol. Process Ser.*, 111, 265–273, <https://doi.org/10.3354/meps111265>, 1994.
- Boyd, E. S., Skidmore, M. L., Mitchell, A. C., Bakermans, C., and Peters, J. W.: Methanogenesis in subglacial sediments., *Environ. Microbiol. Rep.*, 2, 685–92, <https://doi.org/10.1111/j.1758-2229.2010.00162.x>, 2010.
- Boyd, E. S., Lange, R. K., Mitchell, A. C., Havig, J. R., Hamilton, T. L., Lafrenière, M. J., Shock, E. L., Peters, J. W., and Skidmore, M. L.: Diversity, abundance, and potential activity of nitrifying and nitrate-reducing microbial assemblages in a subglacial ecosystem., *Appl. Environ. Microbiol.*, 77, 4778–87, <https://doi.org/10.1128/AEM.00376-11>, 2011.
- Boyd, E. S., Hamilton, T. L., Havig, J. R., Skidmore, M. L., and Shock, E. L.: Chemolithotrophic primary production in a subglacial ecosystem, *Appl. Environ. Microbiol.*, 80, 6146–6153, <https://doi.org/10.1128/AEM.01956-14>, 2014.
- Burgess, D. O., Sharp, M. J., Mair, D. W. F., Dowdeswell, J. A., and Benham, T. J.: Flow dynamics and iceberg calving rates of Devon Ice Cap, Nunavut, Canada, *J. Glaciol.*, 51, 219–230, <https://doi.org/10.3189/172756505781829430>, 2005.
- Cheng, S. M. and Foght, J. M.: Cultivation-independent and -dependent characterization of Bacteria resident beneath John Evans Glacier, *FEMS Microbiol. Ecol.*, 59, 318–330, <https://doi.org/10.1111/j.1574-6941.2006.00267.x>, 2007.
- Christ, A. J., Bierman, P. R., Schaefer, J. M., Dahl-Jensen, D., Steffensen, J. P., Corbett, L. B., Peteet, D. M., Thomas, E. K., Steig, E. J., Rittenour, T. M., Tison, J. L., Blard, P. H., Perdrial, N., Dethier, D. P., Lini, A., Hidy, A. J., Caffee, M. W., and Southon, J.: A multimillion-year-old record of Greenland vegetation and glacial history preserved in sediment beneath 1.4 km of ice at Camp Century, *Proc. Natl. Acad. Sci. U. S. A.*, 118, <https://doi.org/10.1073/pnas.2021442118>, 2021.
- Christner, B. C., Priscu, J. C., Achberger, A. M., Barbante, C., Carter, S. P., Christianson, K., Michaud, A. B., Mikucki, J. A., Mitchell, A. C., Skidmore, M. L., Vick-Majors, T. J., and the WISSARD Science Team: A microbial ecosystem beneath the West Antarctic ice sheet, *Nature*, 512, 310–313, <https://doi.org/10.1038/nature13667>, 2014.
- Clayton, J. R., Reimnitz, E., Payne, J. R., and Kempema, E. W.: Effects of advancing freeze fronts on distributions of fine-grained sediment particles in seawater- and freshwater-slush ice slurries, *J. Sediment. Petrol.*, 60, 145–151, <https://doi.org/10.1306/212f912e-2b24-11d7-8648000102c1865d>, 1990.
- Coble, P. G.: Characterization of marine and terrestrial DOM in seawater using excitation-emission spectroscopy, *Mar. Chem.*, 51, 325–346, 1996.
- Coble, P. G.: Aquatic organic matter fluorescence, Cambridge University Press, Cambridge, <https://doi.org/10.1017/CBO9781139045452>, 2014.
- Copland, L., Lacelle, D., Fisher, D., Delaney, F., Thomson, L., Main, B., and Burgess, D.: Warmer-wetter climate drives shift in  $\delta\text{D}-\delta^{18}\text{O}$  composition of precipitation across the queen Elizabeth islands, arctic Canada, *Arct. Sci.*, 7, 136–157,

<https://doi.org/10.1139/as-2020-0009>, 2021.

Cory, R. M. and McKnight, D. M.: Fluorescence Spectroscopy Reveals Ubiquitous Presence of Oxidized and Reduced Quinones in Dissolved Organic Matter, *Environ. Sci. Technol.*, 39, 8142–8149, <https://doi.org/10.1021/es0506962>, 2005.

Cress, P. and Wyness, R.: The Devon Island expedition, observations of glacial movements, *Arctic*, 14, 257–259, 1961.

615 D’Andrilli, J. and McConnell, J. R.: Polar ice core organic matter signatures reveal past atmospheric carbon composition and spatial trends across ancient and modern timescales, *J. Glaciol.*, 67, 1028–1042, <https://doi.org/10.1017/jog.2021.51>, 2021.

D’Andrilli, J., Foreman, C. M., Sigl, M., Prisco, J. C., and McConnell, J. R.: A 21000-year record of fluorescent organic matter markers in the WAIS Divide ice core, *Clim. Past*, 13, 533–544, <https://doi.org/10.5194/cp-13-533-2017>, 2017.

620 Darcy, J. L., Lynch, R. C., King, A. J., Robeson, M. S., and Schmidt, S. K.: Global distribution of *Polaromonas* phylotypes - evidence for a highly successful dispersal capacity, *PLoS One*, 6, <https://doi.org/10.1371/journal.pone.0023742>, 2011.

Davis, S. N., Whittemore, D. O., and Fabryka-Martin, J.: Uses of Chloride/Bromide Ratios in Studies of Potable Water, *Groundwater*, 36, 338–350, <https://doi.org/10.1111/J.1745-6584.1998.TB01099.X>, 1998.

625 Dubnick, A., Barker, J. D., Sharp, M. J., Wadham, J. L., Lis, G., Telling, J. P., Fitzsimons, S., and Jackson, M.: Characterization of dissolved organic matter (DOM) from glacial environments using total fluorescence spectroscopy and parallel factor analysis, *Ann. Glaciol.*, 51, 111–122, <https://doi.org/10.3189/172756411795931912>, 2010.

Dubnick, A., Sharp, M. J., Danielson, B., Saidi-Mehrabad, A., and Barker, J. D.: Basal thermal regime affects the biogeochemistry of subglacial systems, *Biogeosciences*, 17, 963–977, <https://doi.org/10.5194/bg-17-963-2020>, 2020.

630 Dunham, E. C., Dore, J. E., Skidmore, M. L., Roden, E. E., and Boyd, E. S.: Lithogenic hydrogen supports microbial primary production in subglacial and proglacial environments, *Proc. Natl. Acad. Sci. U. S. A.*, 118, <https://doi.org/10.1073/PNAS.2007051117>, 2021.

Fellman, J. B., D’Amore, D. V., Hood, E. W., and Boone, R. D.: Fluorescence characteristics and biodegradability of dissolved organic matter in forest and wetland soils from coastal temperate watersheds in southeast Alaska, *Biogeochemistry*, 88, 169–184, <https://doi.org/10.1007/s10533-008-9203-x>, 2008.

635 Fellman, J. B., Spencer, R. G. M., Hernes, P. J., Edwards, R. T., D’Amore, D. V., and Hood, E. W.: The impact of glacier runoff on the biodegradability and biochemical composition of terrigenous dissolved organic matter in near-shore marine ecosystems, *Mar. Chem.*, 121, 112–122, <https://doi.org/10.1016/j.marchem.2010.03.009>, 2010.

Fenchel, T. and Blackburn, T. H.: *Bacteria and mineral cycling*, Academic Press, New York, 1979.

Ferrick, M. G., Calkins, D. J., Perron, N. M., Cragin, J. H., and Kendall, C.: Diffusion model validation and interpretation of stable isotopes in river and lake ice, *Hydrol. Process.*, 16, 851–872, 2002.

640 Foght, J. M., Aislabie, J., Turner, S., Brown, C. E., Ryburn, J., Saul, D. J., and Lawson, W.: Culturable bacteria in subglacial sediments and ice from two Southern Hemisphere glaciers., *Microb. Ecol.*, 47, 329–340, <https://doi.org/10.1007/s00248-003-1036-5>, 2004.

Föllmi, K. B., Hosein, R., Arn, K., and Steinmann, P.: Weathering and the mobility of phosphorus in the catchments and forefields of the Rhône and Oberaar glaciers, central Switzerland: Implications for the global phosphorus cycle on glacial-

- 645 interglacial timescales, *Geochim. Cosmochim. Acta*, 73, 2252–2282, <https://doi.org/10.1016/j.gca.2009.01.017>, 2009.
- Gaidos, E. J., Lanoil, B., Thorsteinsson, T., Graham, A., Skidmore, M. L., Han, S.-K., Rust, T., and Popp, B.: A viable microbial community in a subglacial volcanic crater lake, Iceland., *Astrobiology*, 4, 327–344, <https://doi.org/10.1089/ast.2004.4.327>, 2004.
- Gawor, J., Grzesiak, J., Sasin-Kurowska, J., Borsuk, P., Gromadka, R., Górnica, D., Świątecki, A., Aleksandrak-Piekarczyk, T., and Zdanowski, M. K.: Evidence of adaptation, niche separation and microevolution within the genus *Polaromonas* on Arctic and Antarctic glacial surfaces, *Extremophiles*, 20, 403–413, <https://doi.org/10.1007/s00792-016-0831-0>, 2016.
- Gill-Olivas, B., Telling, J. P., Tranter, M., Skidmore, M. L., Christner, B. C., O’Doherty, S., and Priscu, J.: Subglacial erosion has the potential to sustain microbial processes in Subglacial Lake Whillans, Antarctica, *Commun. Earth Environ.*, 2, <https://doi.org/10.1038/s43247-021-00202-x>, 2021.
- 655 Gill-Olivas, B., Telling, J., Skidmore, M., and Tranter, M.: Abrasion of sedimentary rocks as a source of hydrogen peroxide and nutrients to subglacial ecosystems, *Biogeosciences*, 20, 929–943, <https://doi.org/10.5194/bg-20-929-2023>, 2023.
- Godwin, C. M. and Cotner, J. B.: Aquatic heterotrophic bacteria have highly flexible phosphorus content and biomass stoichiometry, *ISME J.*, 9, 2324–2327, <https://doi.org/10.1038/ismej.2015.34>, 2015.
- Grasby, S. E., Allen, C. C., Longazo, T. G., T, L. J., W, G. D., and B, B.: Supraglacial Sulfur Springs and Associated Biological Activity in the Canadian High Arctic—Signs of Life Beneath the Ice, *Astrobiology*, 3, 583–596, 2003.
- 660 Grasby, S. E., Allen, D. M., Bell, S., Chen, Z., Ferguson, G., Jessop, A., Kelman, M., Ko, M., Majorowicz, J., Moore, M., Raymond, J., and Therrien, R.: Geothermal Energy Resource Potential of Canada, 2012.
- Hamilton, T. L., Peters, J. W., Skidmore, M. L., and Boyd, E. S.: Molecular evidence for an active endogenous microbiome beneath glacial ice., *ISME J.*, 7, 1402–12, <https://doi.org/10.1038/ismej.2013.31>, 2013.
- 665 Harrison, J. C., Lynds, T., Ford, A., and Rainbird, R. H.: Simplified tectonic assemblage map of the Canadian Arctic Islands, *Geol. Surv. Canada, Can. Geosci. Map 80 (preliminary)*, scape12 000 000, 80, 297416, <https://doi.org/10.4095/297416>, 2016.
- Hawkings, J. R., Wadham, J. L., Tranter, M., Raiswell, R., Benning, L. G., Statham, P. J., Tedstone, A., Nienow, P., Lee, K., and Telling, J. P.: Ice sheets as a significant source of highly reactive nanoparticulate iron to the oceans., *Nat. Commun.*, 5, 1–8, <https://doi.org/10.1038/ncomms4929>, 2014.
- 670 Hawkings, J. R., Wadham, J. L., Tranter, M., Telling, J. P., Bagshaw, E. A., Beaton, A., Simmons, S., Chandler, D., Tedstone, A., and Nienow, P.: The Greenland Ice Sheet as a hotspot of phosphorus weathering and export in the Arctic, *Global Biogeochem. Cycles*, 30, 1–22, <https://doi.org/10.1002/2015GB005237>, 2016.
- Hodson, A. J., Mumford, P., and Lister, D.: Suspended sediment and phosphorous in proglacial rivers: Bioavailability and potential impacts upon the P status of ice-marginal receiving waters, *Hydrol. Process.*, 18, 2409–2422, <https://doi.org/10.1002/hyp.1471>, 2004.
- 675 Hodson, A. J., Mumford, P. N., Kohler, J., and Wynn, P. M. P. M.: The High Arctic glacial ecosystem: new insights from nutrient budgets, *Biogeochemistry*, 72, 233–256, <https://doi.org/10.1007/s10533-004-0362-0>, 2005.
- Hood, E., Battin, T. J., Fellman, J., O’neel, S., and Spencer, R. G. M.: Storage and release of organic carbon from glaciers and

- ice sheets, *Nat. Geosci.*, 8, 91–96, <https://doi.org/10.1038/ngeo2331>, 2015.
- 680 Hougardy, A. and Klemme, J. H.: Nitrate reduction in a new strain of *Rhodospirillum rubrum*, *Arch. Microbiol.*, 164, 358–362, <https://doi.org/10.1007/BF02529983>, 1995.
- Irvine-Fynn, T. D. L., Hodson, A. J., Moorman, B. J., Vatne, G., and Hubbard, A. L.: Polythermal glacier hydrology: a review, *Rev. Geophys.*, 49, 1–37, <https://doi.org/10.1029/2010RG000350>.1.INTRODUCTION, 2011.
- Jeon, C. O., Park, W., Padmanabhan, P., DeRito, C., Snape, J. R., and Madsen, E. L.: Discovery of a bacterium, with distinctive dioxigenase, that is responsible for *in situ* biodegradation in contaminated sediment, *Proc. Natl. Acad. Sci.*, 100, 13591–13596, <https://doi.org/10.1073/pnas.1735529100>, 2003.
- 685 Jørgensen, L., Stedmon, C. A., Kragh, T., Markager, S., Middelboe, M., and Søndergaard, M.: Global trends in the fluorescence characteristics and distribution of marine dissolved organic matter, *Mar. Chem.*, 126, 139–148, <https://doi.org/10.1016/j.marchem.2011.05.002>, 2011.
- 690 Jouzel, J., Petit, J. R., Souchez, R., Barkov, N. I., Lipenkov, V. Y., Raynaud, D., Stievenard, M., Vassiliev, N. I., Verbeke, V., and Vimeux, F.: More than 200 meters of lake ice above subglacial Lake Vostok, Antarctica, *Science* (80-. ), 286, 2138–2141, <https://doi.org/10.1126/science.286.5447.2138>, 1999.
- Kayani, M. ur R., Doyle, S. M., Sangwan, N., Wang, G., Gilbert, J. A., Christner, B. C., and Zhu, T. F.: Metagenomic analysis of basal ice from an Alaskan glacier, *Microbiome*, 6, 14–16, <https://doi.org/10.1186/s40168-018-0505-5>, 2018.
- 695 Keeler, C. M.: Relationship between climate, ablation, and runoff on the Sverdrup Glacier 1963, Devon Island, NWT., 1964.
- Kellerman, A. M., Vonk, J., McColaugh, S., Podgorski, D. C., van Winden, E., Hawkings, J. R., Johnston, S. E., Humayun, M., and Spencer, R. G. M.: Molecular Signatures of Glacial Dissolved Organic Matter from Svalbard and Greenland, *Global Biogeochem. Cycles*, 35, <https://doi.org/10.1029/2020GB006709>, 2021.
- Killawee, J. A., Fairchild, I. J., Tison, J. L., Janssens, L., and Lorrain, R.: Segregation of solutes and gases in experimental freezing of dilute solutions: Implications for natural glacial systems, *Geochim. Cosmochim. Acta*, 62, 3637–3655, [https://doi.org/10.1016/S0016-7037\(98\)00268-3](https://doi.org/10.1016/S0016-7037(98)00268-3), 1998.
- 700 Koerner, R. M.: The Devon Island Expedition: Glaciology, *Arctic*, 14, 256–257, <https://doi.org/10.14430/arctic3683>, 1961.
- Koerner, R. M.: Mass balance of glaciers in the Queen Elizabeth Islands, Nunavut, Canada, *Ann. Glaciol.*, 42, 417–423, 2005.
- Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., and Schloss, P. D.: Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform., *Appl. Environ. Microbiol.*, 79, 5112–5120, <https://doi.org/10.1128/AEM.01043-13>, 2013.
- 705 Lanoil, B., Skidmore, M. L., Priscu, J. C., Han, S., Foo, W., Vogel, S. W., Tulaczyk, S., and Engelhardt, H.: Bacteria beneath the West Antarctic ice sheet., *Environ. Microbiol.*, 11, 609–615, <https://doi.org/10.1111/j.1462-2920.2008.01831.x>, 2009.
- Lawson, E. C., Bhatia, M. P., Wadham, J. L., and Kujawinski, E. B.: Continuous summer export of nitrogen-rich organic matter from the Greenland Ice Sheet inferred by ultrahigh resolution mass spectrometry., *Environ. Sci. Technol.*, 48, 14248–14257, <https://doi.org/10.1021/es501732h>, 2014.
- Li, X., Wang, N., Ding, Y., Hawkings, J. R., Yde, J. C., Raiswell, R., Liu, J., Zhang, S., Kang, S., Wang, R., Liu, Q., Liu, S.,

- Bol, R., You, X., and Li, G.: Globally elevated chemical weathering rates beneath glaciers, *Nat. Commun.*, 13, 1–13, <https://doi.org/10.1038/s41467-022-28032-1>, 2022.
- 715 Macdonald, M. L., Wadham, J. L., Telling, J. P., and Skidmore, M. L.: Glacial Erosion Liberates Lithologic Energy Sources for Microbes and Acidity for Chemical Weathering Beneath Glaciers and Ice Sheets, *Front. Earth Sci.*, 6, 1–15, <https://doi.org/10.3389/feart.2018.00212>, 2018.
- Madigan, M. T., Jung, D. O., Woese, C. R., and Achenbach, L. A.: *Rhodoferrax antarcticus* sp. nov., a moderately psychrophilic purple nonsulfur bacterium isolated from an Antarctic microbial mat, *Arch. Microbiol.*, 173, 269–277,   
720 <https://doi.org/10.1007/s002030000140>, 2000.
- Majoube, M.: fractionnement en oxygène-18 et en deuterium entre l'eau et la vapeur, *J. Chim. Phys.*, 68, 1423–1436, 1971.
- Makino, W., Cotner, J. B., Sterner, R. W., and Elser, J. J.: Are bacteria more like plants or animals? Growth rate and resource dependence of bacterial C:N:P stoichiometry, *Funct. Ecol.*, 17, 121–130, <https://doi.org/10.1046/j.1365-2435.2003.00712.x>, 2003.
- 725 Margesin, R. and Miteva, V.: Diversity and ecology of psychrophilic microorganisms, *Res. Microbiol.*, 162, 346–361, <https://doi.org/10.1016/j.resmic.2010.12.004>, 2011.
- Martinussen, I. and Thingstad, T.: Utilization of N, P and organic C by heterotrophic bacteria. II. Comparison of experiments and a mathematical model, *Mar. Ecol. Prog. Ser.*, 37, 285–293, <https://doi.org/10.3354/meps037285>, 1987.
- Mattes, T. E., Alexander, A. K., Richardson, P. M., Munk, A. C., Han, C. S., Stothard, P., and Coleman, N. V.: The genome of   
730 *Polaromonas* sp. strain JS666: insights into the evolution of a hydrocarbon- and xenobiotic-degrading bacterium, and features of relevance to biotechnology., *Appl. Environ. Microbiol.*, 74, 6405–6416, <https://doi.org/10.1128/AEM.00197-08>, 2008.
- Meyer, A. F., Lipson, D. A., Martin, A. P., Schadt, C. W., and Schmidt, S. K.: Molecular and metabolic characterization of cold-tolerant alpine soil *Pseudomonas sensu stricto.*, *Appl. Environ. Microbiol.*, 70, 483–489, <https://doi.org/10.1128/AEM.70.1.483-489.2004>, 2004.
- 735 Michaud, A. B., Skidmore, M. L., Mitchell, A. C., Vick-Majors, T. J., Barbante, C., Turetta, C., Van Gelder, W., and Priscu, J. C.: Solute sources and geochemical processes in Subglacial Lake Whillans, West Antarctica, *Geology*, 44, 347–350, <https://doi.org/10.1130/G37639.1>, 2016.
- Michaud, A. B., Dore, J. E., Achberger, A. M., Christner, B. C., Mitchell, A. C., Skidmore, M. L., Vick-Majors, T. J., and Priscu, J. C.: Microbial oxidation as a methane sink beneath the West Antarctic Ice Sheet, *Nat. Geosci.*, 10, 582–586,   
740 <https://doi.org/10.1038/NGEO2992>, 2017.
- Michel, B. and Ramseier, R. O.: Classification of river and lake ice, *Can. Geotech. J.*, 8, 36–45, <https://doi.org/10.1139/t71-004>, 1971.
- Mikucki, J. A., Pearson, A., Johnston, D. T., Turchyn, A. V., Farquhar, J., Schrag, D. P., Anbar, A. D., Priscu, J. C., and Lee, P. A.: A contemporary microbially maintained subglacial ferrous “ocean”, *Science*, 324, 397–400,   
745 <https://doi.org/10.1126/science.1167350>, 2009.
- Mitchell, A. C., Lafrenière, M. J., Skidmore, M. L., and Boyd, E. S.: Influence of bedrock mineral composition on microbial

- diversity in a subglacial environment, *Geology*, 41, 855–858, <https://doi.org/10.1130/G34194.1>, 2013.
- Miteva, V. I., Sheridan, P. P., and Brenchley, J. E.: Phylogenetic and Physiological Diversity of Microorganisms Isolated from a Deep Greenland Glacier Ice Core, *70*, 202–213, <https://doi.org/10.1128/AEM.70.1.202>, 2004.
- 750 Montross, S., Skidmore, M. L., Christner, B. C., Samyn, D., Tison, J.-L., Lorrain, R., Doyle, S., and Fitzsimons, S.: Debris-Rich Basal Ice as a Microbial Habitat, Taylor Glacier, Antarctica, *Geomicrobiol. J.*, 31, 76–81, <https://doi.org/10.1080/01490451.2013.811316>, 2014.
- Morono, Y., Terada, T., Kallmeyer, J., and Inagaki, F.: An improved cell separation technique for marine subsurface sediments: applications for high-throughput analysis using flow cytometry and cell sorting., *Environ. Microbiol.*, 15, 2841–2849, <https://doi.org/10.1111/1462-2920.12153>, 2013.
- 755 Murphy, K. R., Stedmon, C. A., Graeber, D., and Bro, R.: Fluorescence spectroscopy and multi-way techniques. PARAFAC, *Anal. Methods*, 5, 6557, <https://doi.org/10.1039/c3ay41160e>, 2013.
- Murphy, K. R., Stedmon, C. A., Wenig, P., and Bro, R.: OpenFluor– an online spectral library of auto-fluorescence by organic compounds in the environment, *Anal. Methods*, 6, 658–661, <https://doi.org/10.1039/C3AY41935E>, 2014.
- 760 Nye, J. F.: Water Flow in Glaciers: Jökulhlaups, Tunnels and Veins, *J. Glaciol.*, 17, 181–207, <https://doi.org/DOI:10.3189/S002214300001354X>, 1976.
- Nyysönen, M., Hultman, J., Ahonen, L., Kukkonen, I., Paulin, L., Laine, P., Itävaara, M., and Auvinen, P.: Taxonomically and functionally diverse microbial communities in deep crystalline rocks of the Fennoscandian shield, *ISME J.*, 8, 126–138, <https://doi.org/10.1038/ismej.2013.125>, 2014.
- 765 Ohno, T. and Bro, R.: Dissolved Organic Matter Characterization Using Multiway Spectral Decomposition of Fluorescence Landscapes, *Soil Sci. Soc. Am. J.*, 70, 2028, <https://doi.org/10.2136/sssaj2006.0005>, 2006.
- Osborne, T. H., Jamieson, H. E., Hudson-Edwards, K. A., Nordstrom, D. K., Walker, S. R., Ward, S. A., and Santini, J. M.: Microbial oxidation of arsenite in a subarctic environment: diversity of arsenite oxidase genes and identification of a psychrotolerant arsenite oxidiser, *BMC Microbiol.*, 10, 205, <https://doi.org/10.1186/1471-2180-10-205>, 2010.
- 770 Paden, J., Li, J., Leuschen, C., Rodriguez-Morales, F., and Hale, R.: IceBridge MCoRDS L2 Ice Thickness, Version 1, <https://doi.org/https://doi.org/10.5067/GDQ0CUCVTE2Q> 2019.
- Parada, A. E., Needham, D. M., and Fuhrman, J. A.: Every base matters: Assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples, *Environ. Microbiol.*, 18, 1403–1414, <https://doi.org/10.1111/1462-2920.13023>, 2016.
- 775 Paul, E. A. and Clark, F. E.: *Soil Microbiology and Biochemistry*, 2nd ed., Academic Press, San Diego, 340 pp., 1996.
- Pautler, B. G., Woods, G. C., Dubnick, A., Simpson, A. J., Sharp, M. J., Fitzsimons, S. J., and Simpson, M. J.: Molecular characterization of dissolved organic matter in glacial ice: coupling natural abundance <sup>1</sup>H NMR and fluorescence spectroscopy., *Environ. Sci. Technol.*, 46, 3753–61, <https://doi.org/10.1021/es203942y>, 2012.
- Perini, L., Gostinčar, C., and Gunde-Cimerman, N.: Fungal and bacterial diversity of Svalbard subglacial ice, *Sci. Rep.*, 9, 1–  
780 15, <https://doi.org/10.1038/s41598-019-56290-5>, 2019.

- Pinti, D.: Delta, Isotopic BT - Encyclopedia of Astrobiology, edited by: Gargaud, M., Amils, R., Quintanilla, J. C., Cleaves, H. J. (Jim), Irvine, W. M., Pinti, D. L., and Viso, M., Springer Berlin Heidelberg, Berlin, Heidelberg, 418, [https://doi.org/10.1007/978-3-642-11274-4\\_406](https://doi.org/10.1007/978-3-642-11274-4_406), 2011.
- 785 Polz, M. F., Hunt, D. E., Preheim, S. P., and Weinreich, D. M.: Patterns and mechanisms of genetic and phenotypic differentiation in marine microbes, *Philos. Trans. R. Soc. B Biol. Sci.*, 361, 2009–2021, <https://doi.org/10.1098/rstb.2006.1928>, 2006.
- Porter, C., Morin, P., Howat, I., Noh, Myoung-Jon; Bates, B., Peterman, Kenneth; Keeseey, S., Schlenk, Matthew; Gardiner, Judith; Tomko, K., Willis, M., Kelleher, C., Cloutier, M., Husby, E., Foga, Steven; Nakamura, H., Platson, Melisa; Wethington, Michael, J. ., Williamson, C., Bauer, G., Enos, J., Arnold, G., Kramer, W., Becker, P., Doshi, A., D’Souza, Cristelle; Cummens, 790 P., Laurier, F., and Bojesen, M.: ArcticDEM V1, <https://doi.org/10.7910/DVN/OHHUKH>, 2018.
- Price, P. B. and Sowers, T.: Temperature dependence of metabolic rates for microbial growth, maintenance, and survival, *Proc. Natl. Acad. Sci. U. S. A.*, 101, 4631–4636, <https://doi.org/10.1073/pnas.0400522101>, 2004.
- Retelletti Brogi, S., Ha, S. Y., Kim, K., Derrien, M., Lee, Y. K., and Hur, J.: Optical and molecular characterization of dissolved organic matter (DOM) in the Arctic ice core and the underlying seawater (Cambridge Bay, Canada): Implication for increased 795 autochthonous DOM during ice melting, *Sci. Total Environ.*, 627, 802–811, <https://doi.org/10.1016/j.scitotenv.2018.01.251>, 2018.
- Röthlisberger, H.: Water Pressure in Intra- and Subglacial Channels, *J. Glaciol.*, 11, 177–203, <https://doi.org/DOI:10.3189/S0022143000022188>, 1972.
- Saadi, I., Borisover, M., Armon, R., and Laor, Y.: Monitoring of effluent DOM biodegradation using fluorescence, UV and 800 DOC measurements, *Chemosphere*, 63, 530–539, <https://doi.org/10.1016/j.chemosphere.2005.07.075>, 2006.
- Santibáñez, P. A., Michaud, A. B., Vick-Majors, T. J., D’Andrilli, J., Chiuchiolo, A., Hand, K. P., and Priscu, J. C.: Differential incorporation of bacteria, organic matter, and inorganic ions into lake ice during ice formation, *J. Geophys. Res. Biogeosciences*, 124, 585–600, <https://doi.org/10.1029/2018JG004825>, 2019.
- Schroth, A. W., Crusius, J., Hoyer, I., and Campbell, R.: Estuarine removal of glacial iron and implications for iron fluxes to 805 the oceans, *Geophys. Res. Lett.*, 799–804, <https://doi.org/10.1002/2013GL058954>.Received, 2014.
- Sharp, M. J., Parkes, J., Cragg, B., Fairchild, I. J., Lamb, H., and Tranter, M.: Widespread bacterial populations at glacier beds and their relationship to rock weathering and carbon cycling, *Geology*, 27, 107–110, [https://doi.org/10.1130/0091-7613\(1999\)027<0107:WBPAGB>2.3.CO;2](https://doi.org/10.1130/0091-7613(1999)027<0107:WBPAGB>2.3.CO;2), 1999.
- Sheik, C. S., Stevenson, E. I., Den Uyl, P. A., Arendt, C. A., Aciego, S. M., and Dick, G. J.: Microbial communities of the 810 Lemon Creek Glacier show subtle structural variation yet stable phylogenetic composition over space and time., *Front. Microbiol.*, 6, 495, <https://doi.org/10.3389/fmicb.2015.00495>, 2015.
- Sheridan, P. P., Miteva, V. I., and Brenchley, J. E.: Phylogenetic analysis of anaerobic psychrophilic enrichment cultures obtained from a Greenland glacier ice core, *Appl. Environ. Microbiol.*, 69, 2153–2160, <https://doi.org/10.1128/AEM.69.4.2153>, 2003.

- 815 Sizova, M. and Panikov, N.: *Polaromonas hydrogenivorans* sp. nov., a psychrotolerant hydrogen-oxidizing bacterium from Alaskan soil., *Int. J. Syst. Evol. Microbiol.*, 57, 616–619, <https://doi.org/10.1099/ijs.0.64350-0>, 2007.
- Skidmore, M. L., Anderson, S. P., Sharp, M. J., Foght, J. M., and Lanoil, B.: Comparison of microbial community compositions of two subglacial environments reveals a possible role for microbes in chemical weathering processes, *Appl. Environ. Microbiol.*, 71, 6986–6997, <https://doi.org/10.1128/AEM.71.11.6986>, 2005.
- 820 Stibal, M., Wadham, J. L., Lis, G. P., Telling, J. P., Pancost, R. D., Dubnick, A., Sharp, M. J., Lawson, E. C., Butler, C. E. H., Hasan, F., Tranter, M., and Anesio, A. M.: Methanogenic potential of Arctic and Antarctic subglacial environments with contrasting organic carbon sources, *Glob. Chang. Biol.*, 18, 3332–3345, <https://doi.org/10.1111/j.1365-2486.2012.02763.x>, 2012a.
- Stibal, M., Hasan, F., Wadham, J. L., Sharp, M. J., and Anesio, A. M.: Prokaryotic diversity in sediments beneath two polar  
825 glaciers with contrasting organic carbon substrates., *Extremophiles*, 16, 255–265, <https://doi.org/10.1007/s00792-011-0426-8>, 2012b.
- Strickland, J. D. H. and Parsons, T. R.: *A Practical Handbook of Seawater Analysis*, 2nd ed., Fisheries Research Board of Canada Bulletin, No 167, 310 pp., 1972.
- Telling, J. P., Boyd, E. S., Bone, N., Jones, E. L., Tranter, M., MacFarlane, J. W., Martin, P. G., Wadham, J. L., Lamarche-  
830 Gagnon, G., Skidmore, M. L., Hamilton, T. L., Hill, E., Jackson, M., and Hodgson, D. A.: Rock comminution as a source of hydrogen for subglacial ecosystems, *Nat. Geosci.*, 8, 851–855, <https://doi.org/10.1038/ngeo2533>, 2015.
- The Mathworks Inc. 2020. MATLAB version 9.9.0.1592791 (2020b). Mathworks Inc., Natick, MA.
- Tranter, M.: Geochemical Weathering in Glacial and Proglacial Environments, in: *Treatise on Geochemistry*, vol. 5–9, Elsevier Inc., 189–205, <https://doi.org/10.1016/B0-08-043751-6/05078-7>, 2003.
- 835 Tranter, M., Sharp, M. J., Lamb, H. R., Brown, G. H., Hubbard, B. P., and Willis, I. C.: Geochemical weathering at the bed of Haut Glacier d’Arolla, Switzerland - a new model, *Hydrol. Process.*, 16, 959–993, <https://doi.org/10.1002/hyp.309>, 2002.
- Tranter, M., Skidmore, M. L., and Wadham, J. L.: Hydrological controls on microbial communities in subglacial environments, *Hydrol. Process.*, 19, 995–998, <https://doi.org/10.1002/hyp.5854>, 2005.
- Vick-Majors, T. J., Mitchell, A. C., Achberger, A. M., Christner, B. C., Dore, J. E., Michaud, A. B., Mikucki, J. A., Purcell,  
840 A. M., Skidmore, M. L., Priscu, J. C., and the WISSARD Science Team: Physiological ecology of microorganisms in subglacial lake whillans, *Front. Microbiol.*, 7, <https://doi.org/10.3389/fmicb.2016.01705>, 2016.
- Vincent, L. A., van Wijngaarden, W. A., and Hopkinson, R.: Surface temperature and humidity trends in Canada for 1953–2005, *J. Clim.*, 20, 5100–5113, <https://doi.org/10.1175/JCLI4293.1>, 2007.
- Wadham, J. L., Bottrell, S. H., Tranter, M., and Raiswell, R.: Stable isotope evidence for microbial sulphate reduction at the  
845 bed of a polythermal high Arctic glacier, *Earth Planet. Sci. Lett.*, 219, 341–355, [https://doi.org/10.1016/S0012-821X\(03\)00683-6](https://doi.org/10.1016/S0012-821X(03)00683-6), 2004.
- Wadham, J. L., Tranter, M., Skidmore, M. L., Hodson, A. J., Priscu, J. C., Lyons, W. B., Sharp, M. J., Wynn, P. M., and Jackson, M.: Biogeochemical weathering under ice: Size matters, *Global Biogeochem. Cycles*, 24, 1–11,

- <https://doi.org/10.1029/2009GB003688>, 2010.
- 850 Wadham, J. L., Arndt, S., Tulaczyk, S., Stibal, M., Tranter, M., Telling, J. P., Lis, G. P., Lawson, E. C., Ridgwell, A., Dubnick, A., Sharp, M. J., Anesio, A. M., and Butler, C. E. H.: Potential methane reservoirs beneath Antarctica., *Nature*, 488, 633–637, <https://doi.org/10.1038/nature11374>, 2012.
- Wadham, J. L., Hawkings, J. R., Telling, J. P., Chandler, D., Alcock, J., Lawson, E. C., Kaur, P., Bagshaw, E. a., Tranter, M., Tedstone, a., and Nienow, P.: Sources, cycling and export of nitrogen on the Greenland Ice Sheet, *Biogeosciences Discuss.*,
- 855 13, 1–30, <https://doi.org/10.5194/bg-2015-484>, 2016.
- Wadham, J. L., Hawkings, J. R., Tarasov, L., Gregoire, L. J., Spencer, R. G. M., Gutjahr, M., Ridgwell, A., and Kohfeld, K. E.: Ice sheets matter for the global carbon cycle, *Nat. Commun.*, 10, 1–17, <https://doi.org/10.1038/s41467-019-11394-4>, 2019.
- Walker, S. A., Amon, R. M. W., Stedmon, C., Duan, S., and Louchouart, P.: The use of PARAFAC modeling to trace terrestrial dissolved organic matter and fingerprint water masses in coastal Canadian Arctic surface waters, *J. Geophys. Res.*
- 860 *Biogeosciences*, 114, 1–12, <https://doi.org/10.1029/2009JG000990>, 2009.
- Wang, H., Zhang, X., Wang, S., Zhao, B., Lou, K., and Xing, X. H.: *Massilia violaceinigr* sp. Nov., a novel purple-pigmented bacterium isolated from glacier permafrost, *Int. J. Syst. Evol. Microbiol.*, 68, 2271–2278, <https://doi.org/10.1099/ijsem.0.002826>, 2018.
- Wu, F. C., Tanoue, E., and Liu, C. Q.: Fluorescence and amino acid characteristics of molecular size fractions of DOM in the
- 865 waters of Lake, *Biogeochemistry*, 245–257, 2003.
- Van Wychen, W., Davis, J., Copland, L., Burgess, D. O., Gray, L., Sharp, M. J., Dowdeswell, J. A., and Benham, T. J.: Variability in ice motion and dynamic discharge from Devon Ice Cap, Nunavut, Canada, *J. Glaciol.*, 63, 436–449, <https://doi.org/10.1017/jog.2017.2>, 2017.
- Xu, Q., Luo, G., Guo, J., Xiao, Y., Zhang, F., Guo, S., Ling, N., and Shen, Q.: Microbial generalist or specialist: Intraspecific
- 870 variation and dormancy potential matter, *Mol. Ecol.*, 31, 161–173, <https://doi.org/10.1111/mec.16217>, 2021.
- Yagi, J. M., Sims, D., Brettin, T., Bruce, D., and Madsen, E. L.: The genome of *Polaromonas naphthalenivorans* strain CJ2, isolated from coal tar-contaminated sediment, reveals physiological and metabolic versatility and evolution through extensive horizontal gene transfer, *Environ. Microbiol.*, 11, 2253–2270, <https://doi.org/10.1111/j.1462-2920.2009.01947.x>, 2009.
- Yamashita, Y., Cory, R. M., Nishioka, J., Kuma, K., Tanoue, E., and Jaffé, R.: Fluorescence characteristics of dissolved organic
- 875 matter in the deep waters of the Okhotsk Sea and the northwestern North Pacific Ocean, *Deep. Res. Part II Top. Stud. Oceanogr.*, 57, 1478–1485, <https://doi.org/10.1016/j.dsr2.2010.02.016>, 2010.
- Yamashita, Y., Kloeppel, B. D., Knoepp, J., Zausen, G. L., and Jaffé, R.: Effects of Watershed History on Dissolved Organic Matter Characteristics in Headwater Streams, *Ecosystems*, 14, 1110–1122, <https://doi.org/10.1007/s10021-011-9469-z>, 2011.
- Yde, J. C., Finster, K. W., Raiswell, R., Steffensen, J. P., Heinemeier, J., Olsen, J., Gunnlaugsson, H. P., and Nielsen, O. B.:
- 880 Basal ice microbiology at the margin of the Greenland Ice Sheet, *Ann. Glaciol.*, 51, 71–79, <https://doi.org/10.3189/172756411795931976>, 2010.

