

Annual to seasonal glacier mass balance in High Mountain Asia derived from Pléiades stereo images: examples from the Pamir and the Tibetan Plateau

5

Daniel Falaschi^{1,2}, Atanu Bhattacharya³, Gregoire Guillet⁴, Lei Huang⁵, Owen King^{1,6}, Kriti Mukherjee⁷, Philipp Rastner⁸, Tandong Yao⁹, Tobias Bolch^{1,10}

10 ¹School of Geography and Sustainable Development, University of St Andrews, Irvine Building, North Street St Andrews KY16 9AL, Scotland, UK.

²Instituto Argentino de Nivología, Glaciología y Ciencias Ambientales (IANIGLA), CCT-CONICET Mendoza, C.C. 330, 5500 Mendoza, Argentina

³Department of Earth Sciences and Remote Sensing, JIS University, Kolkata, India-700109

⁴Civil and Environmental Engineering, University of Washington, Seattle, WA, USA

15 ⁵Aerospace Information Research Institute, Chinese Academy of Sciences, Dengzhuang south road 9, Haidian District, Beijing, China-100094

⁶School of Geography, Politics and Sociology, Newcastle University, Newcastle upon Tyne, NE1 7RU.

20 ⁷Cranfield Environment Centre, Cranfield University, College Road, Bedford, MK43 0AL, UK

⁸Department of Geography, University of Zurich, Switzerland

⁹Institute of Tibet Plateau Research, Chinese Academy of Sciences, Beijing, China

¹⁰Institute of Geodesy, Graz University of Technology, Graz, Austria

25 *Correspondence to:* Daniel Falaschi (dfalaschi@mendoza-conicet.gob.ar)

Abstract. Glaciers are crucial sources of freshwater in particular for the arid lowlands surrounding High Mountain Asia. In order to better constrain glacio-hydrological models, annual, or even better, seasonal information about glacier mass changes is highly beneficial. In this study, we test the suitability of very high-resolution Pléiades DEMs to measure glacier-wide mass balance at annual and seasonal scales in two regions of High Mountain Asia (Muztagh Ata in Eastern Pamir and parts of Western Nyainqêntanglha, South-central Tibetan Plateau), where recent estimates have shown contrasting glacier behavior. We find that the average annual mass balance in Muztagh Ata between 2019 and 2022 was -0.07 ± 0.20 m w.e. a^{-1} , suggesting the continuation of a recent phase of slight mass loss following a prolonged period of balanced mass budgets previously observed. The mean annual mass balance in Western Nyainqêntanglha for the same period was highly negative (-0.60 ± 0.15 m w.e. a^{-1}), suggesting increased mass loss rates [compared to the previous six decades](#). The 2022 winter ($+0.13 \pm 0.24$ m w.e.) and summer (-0.35 ± 0.15 m w.e.) mass budgets in Muztagh Ata and Western Nyainqêntanglha (-0.03 ± 0.27 m w.e. in winter; -0.63 ± 0.07 m w.e. in summer) suggest winter and summer accumulation-type regimes, respectively. We support our findings by implementing a Sentinel-1–based Glacier Index to identify the firm and wet snow areas on glaciers and characterize accumulation type. The good match between the geodetic and Glacier Index results demonstrates the potential of very high-resolution Pléiades data to monitor mass balance at short time scales and improves our understanding of glacier accumulation regimes across High Mountain Asia.

1 Introduction

45

Fluctuations of glaciers across High Mountain Asia are at the core of both scientific and public debate due to their major relevance as sources of freshwater for human consumption, their regulatory role of river runoff, and their contribution to sea level rise (Immerzeel et al. 2020; Vishwakarma et al., 2022; Yao et al., 2022). The combination of varying climatic and accumulation regimes (Yao et al., 2012; Huang et al., 2022), debris-cover fraction (Scherler et al., 2018) glacier surges (Farinotti et al., 2020; King et al., 2021; Guillet et al., 2022), presence of supra-glacial and proglacial lakes (Brun et al. 2019; Maurer et al., 2016; King et al., 2019), and inherent dynamic factors interact with ongoing climate change (Armstrong et al. 2021) and result in spatially and temporally variable mass loss rates (Brun et al. 2019; Dehecq et al., 2019). In recent decades, however, consistent increases in glacier wastage have been observed throughout most of High Mountain Asia (Bhattacharya et al., 2021, Hugonnet et al.

50

2021). The most notable exception to this trend is the glaciers within the ‘Pamir-Karakorum’ anomaly, which also extends into Eastern Pamir and the Western Kunlun Shan mountains (Kääb et al., 2015, Liang et al., 2022). In this region, glaciers have been in balance or have had slight mass gains since at least the 1970s (Kääb et al., 2015; Bolch et al., 2017, 2019; Brun et al., 2017; Berthier and Brun, 2019). Recent research, however, hints to an end of this anomaly (Bhattacharya et al., 2021; Hugonnet et al. 2021). Monitoring glacier changes from both remotely-sensed and in situ observations has thus been fundamental for an improved understanding of the relation between climate change and glaciers, and has yielded solid evidence of glacier mass loss worldwide (Zemp et al., 2015, Hugonnet et al., 2021). The measurement of glacier mass budgets involves the summation of accumulation and ablation over a specific period of time (Cogley et al., 2011). Presently, the two most widely used methods for determining glacier mass balance are the glaciological and geodetic methods (Cogley, 2009). On one hand, the glaciological method measures surface mass balance, typically on a seasonal scale, using a combination of distributed snow pits and stakes to measure accumulation and ablation, respectively (Cogley, 2009). On the other hand, the geodetic method consists of the differentiation of multi-temporal and often multi-sourced elevation data (Digital Elevation Models -DEMs), covering usually longer time intervals (>5 years) and larger regions compared to the glaciological method (e.g. Braun et al, 2019; Dussaillant et al., 2019; Shean et al, 2019; Davaze et al., 2020; Hugonnet et al., 2021). In contrast to the glaciological method, the geodetic approach does not only measure the surface mass balance, but also englacial and subglacial processes (Zemp et al., 2013; Andreassen et al., 2016). Moreover, although the glaciological and geodetic methods provide independent observations of glacier mass balance, geodetic surveys of variable temporal resolution can potentially serve to validate and calibrate glaciological records (Zemp et al., 2013; Xu et al., 2018; Wagnon et al., 2021).

When shorter time intervals are considered, the geodetic method has two fundamental limitations: 1) the DEM precision in relation to the magnitude of the elevation change signal during an annual to seasonal time interval, and 2) the uncertainties of the snow, ice and firn density and densification linked to the volume to mass conversion factor, which are high over short timescales (Huss, 2013; Pelto et al., 2019). This information is, as noted, only available for a restricted glacier sample worldwide.

The increased availability of sub-meter, very-high resolution (VHR) satellite imagery (e.g. Pléiades, WorldView) and associated DEMs allows for the quantification of small scale, low magnitude (meter scale) changes at the Earth’s surface whilst retaining a suitable level of precision (Berthier et al., 2014). Testing of these DEMs over several mountain sites across the world has confirmed accuracies ranging between 0.2 to 1 m (Berthier et al., 2014; Shean et al., 2016), indicating high potential for assessing glacier elevation changes over short (<3 yrs) time intervals. Glaciers with relatively high mass balance amplitude and heavily snowbound areas are particularly suited to this purpose (Belart et al., 2017, Deschamps-Berger et al., 2020). Additionally, other very-high resolution DEMs can be derived from aerial photographs captured by terrestrial surveys, unmanned aerial vehicles (UAV) and terrestrial or airborne Lidar scanners and laser altimetry (e.g. ICESat-2) to derive short-term glacier elevation changes (Huss et al., 2013; Fischer et al., 2016; Pelto et al., 2019; Wand et al., 2021; Wu et al., 2022).

The major aims of this paper are therefore to investigate the potential and limitations of geodetic mass balance estimates derived from VHR Pléiades satellite data, using 5 DEMs over the 3-year period 2020-2022. We obtain annual and seasonal mass changes in order to make inferences about glacier accumulation regimes in the selected sites based on the geodetic results and by the SAR-derived glacier index of Huang et al. (2022), supported by climatic data. For these investigations we have selected two specific regions in High Mountain Asia which have displayed dissimilar mass change rates over approximately the last 6 decades: the Muztag Ata (Eastern Pamir), which is predominantly influenced by mid-latitude westerlies and where glaciers have been close to in-balance conditions, and Western Nyainqêntanglha (Central-Eastern Tibet), which is more influenced by the monsoon and has had a strongly negative mass budget (Bhattacharya et al. (2021). Moreover, we aim to monitor the ongoing mass balance trends in the Muztag Ata and Western Nyainqêntanglha regions.


2 Study areas

2.1 Muztag Ata

The Muztag Ata massif is situated in Eastern Pamir, west of the Taklamakan Desert in the Xinjiang Uighur Autonomous province, China (Fig. 1). This mountain range represents an area of anomalously high topography at the northwestern tip of the Tibetan Plateau, rising ~4000 m above the ~3500 m asl. Plateau (Seong et al., 2009). The maximum elevation in the area is Mount Muztag Ata (38°17' N, 75°07' E; 7546 m asl.).

A relevant aspect of the atmospheric circulation over High Mountain Asia is its influence on the glaciers’ thermal regime. Climatically, the Muztag Ata region is characterized by a semi-arid continental-type regime, said to be one of the driest and coldest glacierized regions in low- to mid-latitude China (Zhou et al., 2014). Precipitation is mainly driven by the mid-latitude westerly flow (Yao et al., 2012) and is concentrated between the spring and summer seasons (April-September). Data from the Taxikorgan meteorological station (37°46' N, 75°14' E, ~50 km south of Muztag Ata and placed at ~3100 m asl.), collected between 1957-2010, indicates a mean annual

115 temperature of +3.4 °C (Yan et al., 2013; Yang et al., 2013). Based on an ice core drilled at ~7000 m, Duan et al.
 (2015) estimated mean annual snow accumulation of 605 mm w.e. a⁻¹.
 As per the year 2013, the Muztag Ata massif contains ~273 km² of glacier ice; the largest individual glacier here
 is Kekesayi Glacier at ~86 km² (Holzer et al., 2015). Valley glaciers are cold-based (Zhou et al., 2014), and
 120 several of them grade distally from debris-free ice into debris-covered and subsequently ice-debris complexes
 similar to the landforms in Central Tien Shan (Bolch et al., 2019).
 Previous assessments of glacier mass balance in Muztag Ata (Yao et al., 2012) had detected balanced conditions
 and even slight mass gains in the early 2000's, placing the Muztag Ata area within the Pamir-Karakorum anomaly.
 More recent studies, however, suggest that although glacier area changes are still negligible, the Muztag Ata
 region has recently shifted from approximately balanced conditions [prior to the year 2009](#), to slightly negative
 125 glacier mass budgets [until 2019](#). Overall, Holzer et al. (2015) and Bhattacharya et al. (2021) found slight mass
 losses over the last 6 decades (-0.03 m ±0.33 w.e. a⁻¹ and -0.06 m ±0.07 w.e. a⁻¹, respectively). [The extension of
 this time series of observations will allow for the further monitoring of this transition.](#)



130 **Figure 1. Overview of the study sites with specific glaciers mentioned throughout the study and location in High Mountain Asia. Background images are Pléiades 2019 panchromatic band mosaics overlaid on top of Shuttle Radar Topography Mission (SRTM) DEM hillshades.**

135 **2.2 Western Nyainqentanglha**

The Western Nyainqentanglha (Nyenchen Tanglha in English) mountains are located in the southeastern center
 of the Tibetan Plateau. The SW-NE oriented mountain range [extends over ~230 km](#) in length and reaches a
 maximum elevation of 7162 m asl. [at the top of Mount Nyainqentanglha](#) (Yao et al., 2010). Our study site covers
 a ~750 km² portion situated at the center of the main high mountain range (Fig. 1, right panel).
 140 In general, glaciers to the SE of the Tibetan Plateau are located in a maritime regime under the influence of the
 Indian summer Monsoon with abundant precipitation. However, the location of the Western Nyainqentanglha
 mountains with respect to major atmospheric circulation patterns in High Mountain Asia results in a highly
 complex climate setting. The NW slopes lie on the windward side of the western winds (which prevail during
 the dry season), whereas the SE flank is exposed to the easterly winds of the Indian summer Monsoon that dominate
 145 in the wet season (Kang et al., 2009). Western Nyainqentanglha glaciers therefore lie in a transitional zone
 between the more continental regime to the NW plateau and the maritime regime to the SE, and so polythermal
 glaciers (mixed basal thermal conditions) are common here (Shi and Menenti, 2013). Overall, the climate around

150 these glaciers has been described as continental summer-precipitation-type, with both accumulation and ablation maxima occurring during the summer months (Kang et al., 2009). For their energy balance model, Caidong and Sorteberg (2010) estimated the annual precipitation between 700 mm and >900 mm for Xibu Glacier.

155 Bolch et al. (2010) inventoried >1000 individual glaciers covering about 800 km² in 2001, whereas for our specific study site the glacier area accounted for [more than 150 km²](#). The debris-covered Xibu Glacier (~23 km²) is the largest sampled glacier. In contrast to the Muztag Ata massif, glaciers in the Western Nyainqêntanglha region have been rapidly shrinking (27% between 1970-2014; Wu et al., 2016) and losing mass at an [accelerating](#) pace (from -0.27 ±0.11 m w.e a⁻¹ [between 1968 and 1976](#) to -0.47 m w.e a⁻¹ [between 2012 and 2018](#); Zhang and Zhang, 2017; Luo et al., 2020; Ren et al., 2020; Bhattacharya et al., 2021). [The glaciological mass balance has been surveyed on Zhadang Glacier, a debris-free, ~2 km² mountain glacier located in the northern flank of our study area \(Fig 1\). Both the in situ and modelled mass balance between 2001 and 2017 have shown mass loss rates between -0.43 m w.e. a⁻¹ and -1.97 m w.e. a⁻¹ \(Yao et al., 2012 and unpublished data; Zhang et al., 2013; Huintjes et al., 2015\).](#)

160 3 Data and methods

165 3.1 Geodetic mass balance

To derive annual and seasonal geodetic mass balance estimates, we produced a time series of glacier surface elevation change estimates through DEM differencing. To ensure the robustness of these data, we have considered, and where necessary corrected for, biases related to DEM misalignment, data voids in the elevation change grids, contrasting acquisition dates and seasonal snow conditions. The utilised data and individual processing steps are described in more detail below.

170 3.1.1 DEMs derived from Pléiades tri-stereo imagery

175 The commercial Pléiades 1A and 1B twin satellites were launched in December 2011 and 2012, respectively. Images are acquired at 0.7 m pixel resolution and delivered by Airbus Intelligence at a ground sampling distance (GSD) of 0.5 m for the panchromatic band (Gleyzes et al., 2012). In this study, we used Pléiades tri-stereo panchromatic scenes to cover the pre- and post-monsoon seasons in Muztag Ata and Western Nyainqêntanglha between 2019 and 2022 (Table 1). Because of the limited Pléiades footprint of 20 x 20 km (Berthier et al., 2014), both study areas were covered by two or three acquisitions separated by two weeks in the worst case (Table 1). From these images, we generated 4 m resolution DEMs and 0.5 m orthoimages using the NASA Ames Stereo Pipeline (ASP, Shean and others, 2016), [using](#) the Semi-Global-Matching algorithm (Shean et al., 2020).

180 No field-surveyed ground control points (GCPs) were used in the DEM extraction process. The vertical bias of Pléiades-derived DEMs with no GCPs has been reported typically [to be](#) between 1-2 m (occasionally up to 7 m), [though this can be reduced to a few decimeters after DEM coregistration \(Berthier et al., 2014; Belart et al., 2017; Falaschi et al., 2023\).](#) [Despite the finer \(12-bit\) radiometric resolution of the Pléiades panchromatic bands compared to other sensors used in glaciological applications \(e.g. ASTER, SPOT5 HRS, Worldview\),](#) fresh snow fallen very shortly before image acquisition can lead to a very high reflection over a large area and lack of contrast in the images. In such cases, a relatively large number of data voids can be present in the derived DEMs. [The voids in the September 2021, March 2022 Western Nyainqêntanglha and April 2022 Muztag Ata Pléiades DEMs, account for 20-23% of the glacier area, whilst the remaining DEMs contain less than 9% data voids.](#)

Table 1. Acquisition dates of the Pléiades imagery. *Full area was covered by two scenes acquired on the same day. **Partial coverage only. [W and E are the western and eastern sectors of each study site covered by the Pléiades acquisitions. The values in brackets in the Sector column indicate the time interval in years between consecutive acquisitions on each sector.](#)

Sector	Muztag Ata		Western Nyainqêntanglha	
	Pléiades	Sentinel-2	Pléiades	Sentinel-2
W	11-09-2019	13-09-2019	11-11-2019	11-11-2019
E	05-09-2019	03-09-2019	29-10-2019	27-10-2019
W (~1)	09-09-2020	07-09-2020	08-10-2020*	11-10-2020*
E (1.05)	22-09-2020	22-09-2020		
W (~1)	08-09-2021	07-09-2021	30-09-2021**	06-10-2021
E (~1)	21-09-2021	22-09-2021		
E (1)	22-09-2021			
W (0.59)	10-04-2022	10-04-2022	18-03-2022	20-03-2022*
E (0.56)	17-04-2022	15-04-2022	19-03-2022	
W (0.40)	31-08-2022	02-09-2022	07-11-2022	05-11-2022
E (0.45)	27-09-2022	27-09-2022	01-11-2022	31-10-2022

3.1.2 Classification of snow and ice using Sentinel-2 scenes

The European Copernicus Sentinel-2 satellites (A and B), carrying the MultiSpectral sensor (MSI), were launched in 2015 and 2017. The instruments onboard Sentinel-2A and 2B (swath width = 290 km, orbit repeat rate = 10 days) acquire data in four visible (VIS) and near-infrared (VNIR) bands at 10 m spatial resolution, and six VNIR and short-wave infrared bands (SWIR) at 20 m resolution (Kääb et al, 2016). We used Sentinel-2 imagery to classify surface cover (ice, firn, snow) over Pléiades imagery due to the availability of an automatic, robust method such as the [‘automated snow mapping on glaciers’ tool](#) (ASMAG, Rastner et al., 2019) to identify snow characteristics.

The distribution of snow, firn and ice surfaces is fundamental for a realistic estimation of annual geodetic glacier mass balance (e.g. Pelto et al., 2019). We generated masks that represent the distribution and density of these components on glacier surfaces to convert volume to mass changes (see Sect. 3.1.6). Whilst snow and bare glacier ice are readily identifiable on multispectral satellite imagery using conventional methods (Paul et al., 2015), the distinction between snow, firn and ice on glacier surfaces can be more difficult. Indeed, several methodological approaches to identify snow line altitudes have shown that firn can be classified as either snow or ice, depending for example on the number of impurities, when using normalized snow indexes or thresholding of single bands/band ratios (e.g. Racoviteanu et al., 2019). Here we implemented the ASMAG algorithm (Rastner et al., 2019, Falaschi et al., 2021), using Sentinel-2 images to distinguish between ice and snow/firn areas. In summary, the algorithm converts the raw digital numbers to top of atmosphere reflectance and maps snow cover ratio (SCR). ASMAG then follows the histogram approach of Bippus (2011) applied to a glacier mask to differentiate ice from snow. To this end, it determines automatically a threshold for the near-Infrared (NIR) band values, in turn based on the Otsu (1979) thresholding algorithm (Rastner et al., 2019).

We selected Sentinel-2 scenes acquired as close to the Pléiades imagery as possible. Out of 17 images used for the mapping of snow and ice, the acquisition date of only two Sentinel-2 images differed by more than 2 days (Table 1) from the Pléiades imagery. We inspected ERA5-Land daily temperature and precipitation data to ensure that seasonal snow conditions had not been significantly altered between the Pléiades and corresponding Sentinel-2 scenes, due to snowfall or unusually high temperature events. According to ERA5-Land daily data (Muñoz-Sabater et al., 2021), 3 cm of fresh snow fell between the 30 September and 6 October 2021 Pléiades and Sentinel-2 images in the Western Nyainqêntanglha range. Because the 30 September Pléiades scene is anyhow snowbound, the overall seasonal snow conditions appear consistent among the two scenes. As for temperature, the summers of 2019 to 2021 show monthly temperature anomalies of less than 0.6 °C, which is far below the historical +2.4 °C maximum for the full 1950-2022 series (see Sect S3.3, S4.3).

Despite the topographic correction featured in the ASMAG algorithm, some glacier parts in cast shadows were initially misclassified as ice in the accumulation regions. To remove these misclassifications, we masked out pixels located above the ASMAG-derived mean snowline altitude (SLA) plus 2 standard deviations and reassigned them to the snow class and implemented a low-pass filter.

3.1.3 Glacier outlines

We used the glacier outlines of the year 2019 available from Bhattacharya et al. (2021), which were adjusted based on the Randolph Glacier Inventory 6.0 (Randolph Consortium, 2017), as the basis for our glacier inventory. For geodetic mass balance studies of small regional extent, [the](#) varying quality of glacier polygons may have a relatively large effect on the final results (Sommer et al., 2020; Falaschi et al., 2023). Thus, the RGI glacier polygons were manually adjusted (accounting for glacier length changes, removing non-glacierised stable ground, reinterpreting debris-covered ice) to fit the glacier extent on each of the Pléiades acquisition dates (Table 1) by visual interpretation of the 0.5 m orthophotos. The glacier area uncertainty was conservatively calculated using the approach followed by Wagnon et al. (2021), as the product of the glacier outline initial perimeter of each sampled time interval and two times the 0.5 m ground sampling distance of the Pléiades panchromatic band.

3.1.4 DEM differencing and generation of elevation change maps

We divided the full 2019-2022 study period [into three intervals covering the 2020, 2021 and 2022 hydrological years](#), and two seasonal (2022 winter and summer) time intervals. Before differencing, we coregistered all DEMs to the 2019 DEM following Nuth and Kääb (2011), [using the median value of the elevation changes on stable terrain to correct biases between DEMs](#). This way, any remaining vertical and horizontal shifts between DEMs are minimized before the *dDEM* grids are generated by subtracting the latter from the initial DEMs.

Elevation change grids derived from Pléiades DEMs often show low-frequency biases in off-glacier elevation change residuals, a phenomenon often termed ‘undulations’ (Hugonnet et al., 2022). These undulations have been attributed to satellite along-track attitude oscillations or *jitter* (Girod et al., 2017; Deschamps-Berger et al., 2020). [We applied a spline fit in the along-track direction \(Falaschi et al., 2023\)](#) to remove the low-frequency undulation patterns (Fig. 2a, b).

To eliminate on- and off-glacier anomalous cell values from the *dDEM* grids, we considered all values exceeding ± 150 m as outliers and removed them. A second outlier removal step was carried out where the neighboring cells of data gaps in the *dDEM* grids contained high-magnitude elevation differences, which shared statistical characteristics with real glacier thinning/thickening. To filter this noise, we first implemented a 3-cell buffer around the data gaps and removed the cells within the buffered areas. We then [followed the approach of Gardelle et al. \(2013\), excluding cells where absolute elevation differences differed by more than three standard deviations from the mean elevation change within each altitude band](#). Finally, we filled the resulting data gaps using the glacier-wide hypsometric approach of McNabb et al. (2019), fitting a fifth-degree polynomial function to the mean elevation change on 50 m elevation bins (Fig. 2c-f). [In the specific case of the Western Nyainqêntanglha 2022 winter elevation change grid, data voids concentrated almost exclusively in the uppermost reaches of the sampled glaciers. At similar elevations, the relatively few valid cells available correspond to crevasse movement and show highly negative elevation changes. The glacier-wide void-filling polynomial fitting thus gives too much weight to these negative values and gives the wrong impression that all of the upper part of the glaciers has significantly thinned at high elevation. Therefore, we removed the few highly negative elevation change values in the uppermost reaches of the elevation change plot and adjusted a new function. In doing so, we resampled the elevation bins to 25 m, so that sufficient bins are available for a representative fit.](#)

Lastly, we mosaiced the individual tiles to generate the final elevation change maps. Basic statistical parameters of our *dDEM* grids (Table 2) after coregistration are akin to previously published geodetic mass balance assessments of annual to seasonal scale in other glacierized regions using Pléiades DEMs (e.g. Beraud et al., 2023).

Table 2. Statistics of *dDEM* grids off-glacier after coregistration (SD = standard deviation, SE = standard error, NMAD = normalized median absolute deviation).

	Percentage of data voids on-glacier (%)	off-glacier mean elevation difference (m)	off-glacier Standard Deviation (m)	NMAD (m)
Muztag Ata				
2019-2020	8.6	-0.09	2.99	1.27
2020-2021	10.4	0.06	2.39	1.30
winter 2022	20.8	0.06	2.89	1.30
summer 2022	28.1	-0.11	1.67	1.32
2021-2022	16.4	<0.01	1.94	1.21
2019-2022	14.4	-0.06	2.07	1.30
Western Nyainqêntanglha				
2019-2020	6.8	-0.02	1.36	1.23
2020-2021	23.5	0.04	1.38	1.07
winter 2022	23.7	-0.11	1.41	1.05
summer 2022	14.7	0.03	1.03	0.80
2021-2022	29.7	-0.07	0.99	0.87
2019-2022	9.2	-0.03	1.00	0.56

285

3.1.5 Bulk density

290

Conversion from geodetic volumetric change (in meters) based on the DEM differencing technique to water equivalent (in m w.e.) requires consideration of the densities of the material involved. With no field-surveyed snow density measurements contemporary to our surveyed time periods available, we used the $410 \pm 60 \text{ kg m}^{-3}$ snow density value [retrieved from snow pits](#) in Muztag Ata N15 (Zhu et al., 2018a). We assumed a $\pm 60 \text{ kg m}^{-3}$ uncertainty value, which is a standard value used in geodetic mass balance assessments (Huss., 2013). For glacier ice areas, we used a density of 900 kg m^{-3} (Zhu et al., 2018a) and an uncertainty of $\pm 10 \text{ kg m}^{-3}$ (Clarke et al., 2013).

295




Figure 2. Along-track bias removal in Muztag Ata *dDEM* grids before (a) and after (b) bias correction. The diverse impact of the jitter effect on individual *dDEM* grids can be [observed](#) left and right of the red line in panel (a). (c, d) and (e, f) depict examples of outlier removal and gap-filling before and after the implementation of the buffer approach [in the accumulation area in Muztag Ata and glacier terminus in Western Nyainqêntanglha](#) *dDEM* grids, respectively. Note the crevasse motion captured as alternating blue (elevation gain) and red (elevation loss) in the in the Muztag Ata panels.

305 3.1.6 Mass balance calculation

We present annual mass balance estimates for the Muztag Ata and Western Nyainqêntanglha regions starting in the winter of 2019 up to the summer of 2022 but provide also estimates of the 2022 winter and summer seasons. The annual mass balance B_a can be defined as the algebraic sum of the winter B_w and summer B_s balances

$$310 \quad B_a = B_w + B_s \quad (1)$$

The total volume change Δv (m^3) over a given time interval dt is then derived from the respective elevation difference dh_i of the two grids at pixel i with cell size r of the DEMs, summed over the number of pixels covering the glacier, and is expressed as (cf. Zemp et al., 2013):

$$315 \quad \Delta v = r^2 \times \sum dh_i \quad (2)$$

Applying the snow and ice masks described in Sect. 3.1.2, we then assessed the annual and seasonal geodetic mass balance Δm (m w.e.) considering the volume change Δv , the density of the involved material ρ , and the snow and ice areas A

320

$$\Delta m = \sum \frac{\Delta v \times \rho}{A} \times dt^{-1} \quad (3)$$

3.1.7 Firn densification correction

Lowering of the annual snowpack on glaciers occurs as a result of firn densification, a process that leads to underestimated volume changes (Sold et al., 2013) and is often neglected in geodetic mass balance studies relying on the *dDEM* technique. We implemented a simple firn densification approach following Sold et al. (2013) to estimate the annual elevation change due to firn densification dh_{firn} integrated over the entire firn column [applied on a pixel scale](#)

$$dh_{firn} = \frac{b_n}{\rho_{firn_u}} - \frac{b_n}{\rho_{firn_l}} \quad (4)$$

where b_n is the net mass balance averaged over the previous mass balance year or period (in kg m^{-2}) over the accumulation area. b_n is positive by definition (Belart et al., 2017). ρ_{firn_u} and ρ_{firn_l} are the density values at the upper and lower ends of the firn column, set to $\rho_{firn_u} = 470 \text{ kg m}^{-3}$ and $\rho_{firn_l} = 857 \text{ kg m}^{-3}$, as retrieved from a 41 m-deep ice core drilled at ~7000 m on Muztag Ata (Duan et al., 2015). Because some of our Pléiades imagery was acquired around the end of the accumulation season (Western Nyainqêntanglha) or contains seasonal snow (Muztag Ata), ‘snow’ areas do not exactly match “accumulation” areas and can have an overall negative elevation change.

Using the most recent elevation [change](#) grids available from Bhattacharya et al. (2021; Muztag Ata: 2013-2019; Western Nyainqêntanglha: 2018-2019), in combination with the “snow” areas derived from the 2019 Sentinel-2 scenes to calculate the net mass balance in the “accumulation” area, leads to a negative signal. [Consequently](#), we considered b_n to be the net mass balance of the snow areas as depicted in the September 2021 (Muztag Ata) and October 2020 (Western) Pléiades scenes, which are those showing the least amount of seasonal snow, and thus snow areas should be representative of accumulation areas. Although topoclimatic factors can make firn lines vary spatially and temporally through time (Guo et al., 2014), for the sake of simplicity we assumed that firn densification did not change over different time periods. Our approach is also based on the accumulation rate retrieved from a single year on each study site, whilst some variability is naturally expected (Duan et al., 2015). We then scaled the firn densification correction linearly according to the length of each time span between consecutive imagery acquisition. Because of the simplifications and assumptions of the firn correction (equal net accumulation through the study period, constant densification rate through the hydrological year), we considered a 50% uncertainty in the firn correction when quantifying the total mass balance error.

3.1.8 Seasonal correction

Ideally, mass balance observations should be made repeatedly at the end of the ablation season to ensure that the full, annual budget of ablation and accumulation has been captured, although data availability means this is rarely the case. Correction for the impact of such seasonal bias is therefore commonly required (Belart et al., 2017 and 2019; Abdel Jaber et al., 2019).

Neglecting such seasonality corrections can introduce potential biases, since the effect of seasonal signals are particularly strong when short time intervals are considered. A simple approach to account for the seasonal shift of acquired remotely-sensed data with respect to the hydrological cycle consists of using the daily mass balance rate for the given period to fill in the deficit of missing days or subtract the input of excess days (Abdel Jaber et al., 2019). The extent of the seasonal bias introduced when using this method therefore depends on the fraction of missing or excess days in relation to the onset of the accumulation and ablation periods. We nevertheless chose this approach since our study sites lie in transitional to continental semi-arid environments where, contrary to glaciers in maritime regions, large annual [ice](#) mass turnover is not expected (Duan et al., 2015; Zhu et al., 2015 and 2018a). Moreover, the percentage of missing/excess days relative to the start of the winter (1 October) and summer (1 April) seasons was below 10% for our Pléiades imagery. We considered that any remaining seasonality-related biases [lie](#) within the overall uncertainty range.

3.1.9 Uncertainty

In spite of the coregistration and bias-correction procedures applied, some of the *dDEM* grids showed systematic residuals on stable terrain (off-glacier) (Table 2). We [defined](#) these biases in elevation change off-glacier σ_{sys} as the mean difference between two DEMs (Koblet et al., 2010):

$$380 \quad \sigma_{sys} = \frac{\sum (H_{DEMi} - H_{DEMf})}{n} \quad (5)$$

being H_{DEMi} and H_{DEMf} the elevation of the initial and final DEMs and n the number of cells on stable terrain. For the calculation of the random uncertainty of the volumetric mass balance estimation $\sigma_{\Delta v}$, we considered the volumetric uncertainties on mean elevation change, snow and ice areas and firn densification (see Sect. 4.2), which we summed quadratically to propagate the error:

$$385 \quad \sigma_{\Delta v} = \sum \sqrt{(dh \times \sigma_{Ai})^2 + (A_i \times \sigma_{dh})^2 + (dh_{firn} \times \sigma_{Ai})^2 + (A_i \times \sigma_{dfirn})^2} \quad (6)$$

390 σ_{Ai} being the area uncertainty, σ_{dh} the uncertainty on the rate of elevation change, dh_{firn} the annual elevation change owed to firn densification, and σ_{dfirn} the uncertainty in firn densification rate. In turn, we calculated σ_{dh} using the *patch* method of Berthier et al. (2016), which evaluates the decay of the mean elevation change error on stable terrain (off-glacier) with the averaging area (Wagnon et al., 2021; Falaschi et al., 2023). Since we attributed a different density and related uncertainty to the snow and ice area classes (snow: $\pm 60 \text{ kg m}^{-3}$; ice: $\pm 10 \text{ kg m}^{-3}$), we calculated the uncertainty of each class separately, by adding the density uncertainty $\sigma_{f_{\Delta v}}$ of each surface to the volumetric uncertainty in Eq. (5):

$$395 \quad \sigma_m = \sum \frac{\sqrt{(f_{\Delta v} \times \sigma_{\Delta v})^2 + (\Delta v \times \sigma_{f_{\Delta v}})^2}}{A} \times dt^{-1} \quad (7)$$

[f_{Δv} being the volume to mass conversion factor of ice and snow.](#)

400 Finally, to obtain the overall glacier-wide mass balance uncertainty $\sigma_{m,tot}$, we summed the uncertainties of the snow and ice areas quadratically:

$$\sigma_{m,tot} = \sqrt{\sigma_{m,ice}^2 + \sigma_{m,snow}^2} \quad (8)$$

405 [The void-filling approach used in the Western Nyainqêntanglha 2022 winter elevation change grid \(Sect. 3.1.2\) affects less than 10% of the total glacier area. The mass balance value differed by 0.01 m w.e. only when compared to the mass balance estimate using the “standard” approach used in other grids. We thus did not include any additional uncertainty source in this specific case, and considered that the void-filling related uncertainty is contained within the overall one.](#)

410 [Previous geodetic mass balance research, similar in nature to this study, have considered ice dynamics \(i.e. vertical ice flow\) as an additional uncertainty source. Because the influence of ice dynamics is usually very small \(<5%\) on a year-to-year basis \(e.g. Mukherjee et al. 2022\), we considered the related uncertainty to be beyond the scope of the present study \(Sect. S3.1.10\).](#)

415 3.2 Glacier Index: wet snow and firn area ratios derived from Sentinel-1 and Landsat OLI imagery

We implemented the Glacier Index of Huang et al. (2022) to characterize glacier accumulation regimes and validate our geodetic results. To account for different glacier areas between the study sites, we express firn and wet snow areas on each region as a fraction of the total glacier area (hereafter referred to as firn area ratio and wet snow area ratio). [The snow area](#) ratio can vary to a great extent across different geographic regions through time, whilst interannual variations of the firn area ratio remain relatively small. The Glacier Index I is then defined as the difference between the firn and wet snow area ratios, [calculated in winter and at the end of the summer season, respectively](#):

$$425 \quad I = \frac{A_{firn}}{A_{total}} - \frac{A_{wet.snow}}{A_{total}} \quad (9)$$

where A_{firn} is the firn zone area, $A_{wet.snow}$ the wet snow zone area and A_{total} the total glacier area. The index is expected to be positive for winter accumulation-type glaciers, as summer snowfall is rare and hence $A_{wet.snow} < A_{firn}$. In contrast, the index is more likely to be negative (i.e. $A_{wet.snow} > A_{firn}$) for summer -glaciers due to recurring summer solid precipitation. We classified regions and glaciers as winter accumulation-type where $I \geq 0.05$ and summer-accumulation type where $I < 0.05$ following Huang et al. (2022).

To discriminate wet snow areas in late summer and firn areas in winter over glaciers in our selected study areas, we used Landsat OLI and Sentinel-1 imagery in Google Earth Engine (Gorelick et al., 2017). First, the Landsat scenes are used to identify debris-covered and debris-free areas (ice and snow) on glacier surfaces applying a threshold to the previously computed Normalized Difference Snow Index (Bruns et al., 2014). Then, since the Sentinel-1 C-band is sensitive to snow wetness and roughness (Huang et al., 2013), the backscattering properties between the different glacier facies are sufficiently different to discriminate the frozen firn in winter (here defined as the period between January and March) and wet snow zone in late summer (defined here as the time interval from July 20 to September 10) using backscatter coefficients (Fig. 3).

To account for overestimations of the firn area ratio (due to thin snow over debris-covered ice and crevasses, acting as corner reflectors) we considered that all pixels > -6 dB are misclassified as firn, whereas pixels < -14 dB are erroneously classified as wet snow (Huang et al., 2022). We corrected the firn and wet snow areas accordingly on both winter and summer Sentinel-1 scenes.

4 Results

4.1 Geodetic mass balance in Muztag Ata and Western Nyainqêntanglha

The glacier-wide, annual geodetic mass balances in Muztag Ata varied from moderately negative to slightly positive conditions (Fig. 4; Table 3). B_a estimated from Eq. (3) was negative for the 2020 (when at its most negative, -0.19 ± 0.14 m w.e.) and 2022 glaciological years (-0.17 ± 0.22 m w.e.), but was positive in 2021 ($+0.15 \pm 0.27$ m w.e.). In terms of seasonal mass balance, the 2022 winter mass budget was positive, whereas the summer budget was negative (Table 3).

On an individual glacier basis, both the annual and seasonal mass balances of Kekesayi Glacier matched the overall trends of the broader Muztag Ata massif. Interestingly, the annual mass balance estimates of Muztag Ata No. 15 Glacier yielded positive values for all three surveyed years, whilst the seasonal winter (positive) and summer (negative) budgets closely followed the general pattern in Muztag Ata, respectively (Table 3).




Figure 3. Examples of firn and wet snow cover retrieval in winter and summer 2021 in Muztag Ata (a, b) and Western Nyainqêntanglha (c, d).

465 The hypsometric distribution of elevation changes in Muztag Ata (Fig. 5a) reveals the lack of a clearly defined
 elevation change (i.e. mass balance) gradient. Periods with a positive mass budget (2021, winter 2022) show
 elevation gains at different elevation ranges. They occurred from 4300 m up to the uppermost reaches of the
 glacierized area (7600 m) during 2021, whereas the winter 2022 shows surface elevation increases below 5600
 470 m. Elevation gains (accumulation) during the year 2020 were restricted to elevations above 6100 m, whilst in the
 summer of 2022 elevation gains occurred above 7400 m only.
 In contrast to Muztag Ata, the annual mass balance of glaciers in Western Nyainqentanglha showed a consistent,
 highly negative signal, with the greatest ice loss rate of $(-0.72 \pm 0.22 \text{ m w.e.})$ measured in 2020 (Fig. 6; Table 3).
 While the winter mass budget was roughly balanced $(-0.03 \pm 0.27 \text{ m w.e.})$, the summer mass budget was strongly
 negative (Table 3). It must be noted though, that the interannual and seasonal variability in mass balance are not
 475 fully comparable due to the incomplete coverage of the study site in the 2021 Pléiades acquisition. The largest,
 debris-covered Xibu Glacier had also a highly negative mass loss throughout the study period. In line with
 the overall seasonal budgets in Western Nyainqentanglha, Xibu Glacier lost mass during both the summer and winter
 seasons. Compared to Xibu Glacier, the mass loss of Zhadang Glacier was even greater on the annual scale
 (peaking at $-1.22 \pm 0.21 \text{ m w.e.}$ in 2020), but recovered during the 2022 winter season (Table 3).
 480 In Western Nyainqentanglha, there are clear annual (and 2022 summer) elevation change gradients. They depict
 a steep, positive gradient from lower to higher elevations, with elevation gains starting between 6100-6400 m
 (Fig. 5b). The 2022 annual hypsometric curve, though similar in shape to the two previous years, shows that all
 of the sampled glacier area underwent ice thinning over this interval. The 2022 winter curve shows slight elevation
 gains around 5500 m, coinciding with maximum area per elevation band. This, however, does not counterbalance
 485 elevation losses elsewhere in the glacier area.




Figure 4. Annual (a, b, c), seasonal (d, e) and multi-annual (f) surface elevation change grids over Muztag Ata and associated geodetic mass balance estimates.

490




Figure 5. Hypsometric curve of glacier surface elevation changes over glacierized area in Muztag Ata (a) and Western Nyainqentanglha (b). The light blue bars represent the ice area distribution on 100 m elevation bins for the 2019 glacier area on the right axis.

495




Figure 6. Annual (a, b, c), seasonal (d, e) and multi-annual (f) surface elevation change grids over Western Nyainqentanglha and associated geodetic mass balance estimates.

500

Table 3. Summary of time-interval volume changes (10^6 m^3) and mass balance (m w.e.) for the total glacier area and selected glaciers. Mass balance values in the 2020-2022 column are provided in m w.e. a^{-1} . Values in parenthesis () are calculated using a density of 850 kg m^{-3} and values in brackets [] with a 3-year weighted density (see Sect. 6.4.1 for details).

	2020		2021		winter 2022		summer 2022		2022		2020-2022	
	Δv	Δm	Δv	Δm	Δv	Δm	Δv	Δm	Δv	Δm	Δv	Δm
Muztag Ata												
total glacier area	-86.7 ± 36.9	-0.19 ± 0.14	+105.6 ± 71.1	+0.15 ± 0.27	+13.4 ± 18.4	+0.13 ± 0.24	-152.9 ± 39.0	-0.35 ± 0.15	-95.1 ± 58.0	-0.17 ± 0.22	-52.0 ± 10.7	[-0.15 ± 0.03] (-0.14 ± 0.03)
No. 15	+0.1 ± 0.2	+0.08 ± 0.35	+0.3 ± 0.4	+0.14 ± 0.25	+0.6 ± 0.3	+0.42 ± 0.22	-0.3 ± 0.4	-0.38 ± 0.20	+0.2 ± 0.4	+0.15 ± 0.24	+0.2 ± 0.1	[+0.09 ± 0.04] (+0.08 ± 0.04)
Kekesavi	-37.8 ± 12.2	-0.31 ± 0.17	+25.1 ± 23.5	+0.15 ± 0.33	+3.9 ± 9.8	+0.11 ± 0.29	-56.4 ± 21.0	-0.49 ± 0.30	-37.3 ± 19.4	-0.25 ± 0.27	-22.9 ± 3.6	[-0.29 ± 0.04] (-0.26 ± 0.04)
Western Nvainqêntanglha												
total glacier area	-191.1 ± 16.8	-0.72 ± 0.16	-98.9 ± 8.9	-0.69 ± 0.08	-5.1 ± 63.2	-0.03 ± 0.27	-143.6 ± 6.7	-0.63 ± 0.07	-110.8 ± 15.8	-0.50 ± 0.22	-168.4 ± 4.8	[-1.04 ± 0.11] (-0.93 ± 0.10)
Zhadang	-2.7 ± 0.3	-1.22 ± 0.41	-1.6 ± 0.2	-1.13 ± 0.21	+0.5 ± 1.0	+0.19 ± 0.31	-2.0 ± 0.2	-0.79 ± 0.12	-1.0 ± 0.3	-0.51 ± 0.21	-2.1 ± 0.1	[-1.61 ± 0.16] (-1.44 ± 0.15)
Xibu	-30.8 ± 3.9	-0.71 ± 0.22	-17.1 ± 2.3	-0.61 ± 0.17	-4.8 ± 13.8	-0.11 ± 0.35	-19.9 ± 1.7	-0.55 ± 0.10	-21.5 ± 3.6	-0.44 ± 0.41	-27.4 ± 1.1	[-0.99 ± 0.22] (-0.89 ± 0.20)

505 4.2 Wet summer snow and winter firn area ratios derived from the Glacier Index

Firn area and wet snow-ratios varied between 0.55 - 0.6 and 0.15 - 0.23 in Muztag Ata and 0.25 - 0.28 and 0.33 - 0.66 in Western Nyainqêntanglha, respectively (Fig. 7 and Table S2). These ratios yielded Glacier Index values between 0.32 and 0.45 in Muztag Ata and -0.38 and -0.07 in Western Nyainqêntanglha. To provide a measure of the individual glacier variability, the Glacier Index in Kekesayi and Muztag Ata No. 15 glaciers varied between 0.17 to 0.28 and 0.68 to 0.89, whilst in Xibu and Zhadang glaciers the Index ranged between -0.25 to -0.04 and -0.51 to -0.05, respectively. Based on these assessments, we attribute a winter accumulation regime for Muztag Ata and a summer accumulation regime in Western Nyainqêntanglha. Moreover, the evolution of the summer wet snow area ratio points at 2020 as the most negative mass balance year in Western Nyainqêntanglha and [2022 in Muztag Ata](#).




Figure 7. Evolution of the firn area– (a) and wet snow area–ratio (b) between 2019 and 2022.

520

5. Discussion

5.1 Methodological constraints

525 5.1.1 Internal consistency of the geodetic mass balance estimates

To verify the consistency of our high-resolution geodetic mass balance surveys at annual to seasonal mass balance, we tested the internal robustness of our geodetic surveys. We evaluated the residuals between accumulated vs. the sum of individual survey periods encompassing identical time intervals, in so-called ‘triangulation’ tests (Nuth et al., 2012; Falaschi et al., 2023). This is a way to measure the impact of using different inputs such as *dDEM* grids, snow and ice distribution maps, and the resulting average material density on geodetic mass balance values when evaluating similar periods yet at different time steps.

530 [On a glacier-wide scale, the differences between the sum of the summer and winter mass balances, and accumulated 2022 annual budget yielded differences of \$\pm 0.05\$ m w.e. and \$\pm 0.10\$ m w.e. between both measurements in Muztag Ata and Western Nyainqêntanglha, respectively. In addition, and on an individual glacier scale, we found a discrepancy in geodetic mass balance for Kekesayi Glacier of \$\pm 0.13\$ m w.e. whilst the difference for the Muztag Ata No. 15 was \$\pm 0.11\$ m w.e. Individually, Zhadang and Xibu glaciers in Western Nyainqêntanglha showed differences between \$0.09\$ - \$0.22\$ m w.e. \$\cdot\$ a \$^{-1}\$.](#)

535 Overall, we find that glacier-wide and for the most part individual differences to be within the uncertainty ranges. We attribute the differences to the overall small differences in average density, which in turn derives from the snow and ice distribution. Indeed, for the September 2021 and April 2022 Sentinel-2 snow and ice masks, we used [average](#) densities of 590 kg m^{-3} and 630 kg m^{-3} , respectively. In this sense, Pelto et al. (2019) showed how the spatial distribution of material densities has a larger impact on seasonal mass balances compared to the assumed density values themselves. In contrast, variable density distribution had a greater impact when smaller areas are considered, e.g. in the case of Muztag Ata No. 15.

540 A second consistency test was carried out to test the impact of varying density assumptions in more detail, evaluating the accumulated 2020-2022 mass budget and the sum of all the individual annual periods. We calculated these differences using a single *dDEM* grid and, a) an overall density of 850 ± 60 kg m^{-3} , and b) a 3-year weighted density (Muztag Ata = 774 ± 60 kg m^{-3} ; Western Nyainqêntanglha = 762 ± 60 kg m^{-3}) for all the material involved following Huss (2013) (Table 3). In Muztag Ata, the 2020-2022 glacier-wide estimate (-0.15 ± 0.03 m w.e. a^{-1} ; scenario a) and -0.14 ± 0.03 m w.e. a^{-1} (scenario b) were similar to the averaged mass budget of the

550

individual periods (-0.07 ± 0.21 m w.e. a⁻¹). The accumulated 2020-2022 mass balance of Kekesayi (-0.29 ± 0.04 m w.e. a⁻¹; scenario a) and -0.26 ± 0.04 m w.e. a⁻¹ (scenario b) differed by ± 0.15 and ± 0.12 m w.e. a⁻¹, respectively, from the averaged individual periods (-0.14 ± 0.26 m w.e. a⁻¹). Differences were smaller in the case of Muztag Ata No. 15 Glacier (± 0.04 m w.e. a). In Western Nyainqêntanglha, the differences were much larger (-0.63 ± 0.15 m w.e. a⁻¹ on average vs. -1.04 ± 0.11 m w.e. a⁻¹; scenario a) and -0.93 ± 0.10 m w.e. a⁻¹ (scenario b)). Expectedly, even larger differences were found for Zhadang and Xibu glaciers, since higher thinning rates will have a greater impact on any given density value.

In comparison with end of summer snow densities retrieved from glaciers in other mountain regions (e.g. Pelto et al., 2019; Beraud et al., 2023), our snow density of 410 kg m^{-3} , derived from in-situ surveys (Zhu et al., 2018a) is somewhat lower. Using a snow density of 570 kg m^{-3} (Pelto et al., 2019), the glacier-wide annual mass balance shifts to -0.21 w.e., $+0.24$ w.e. and -0.21 w.e. for the years 2020, 2021 and 2022 in Muztag Ata, and -0.89 w.e., -0.71 w.e. and -0.85 w.e. in Western Nyainqêntanglha. This approach, or the use of a 3-year weighted density as in Huss (2013) tend to reduce the differences between the accumulated vs. added mass budgets. The above confirms that variable density should be used for time spans of 3 years or less, whereas a more thorough density inspection is preferable for shorter timescales (Belart et al., 2017; Klug et al., 2018). Since we covered a period of 3 years only, no conclusion can be drawn on longer time spans.

5.1.2 Potential and limitations of Pléiades DEMs for assessment of short-term geodetic mass balance

Here we discuss the suitability of our Pléiades DEM dataset to quantify both annual and seasonal mass balances over areas with contrasting glacier mass change rates.

We found that 90% of the annual and seasonal glacier elevation changes in Muztag Ata and Western Nyainqêntanglha range between -4.7 m to $+3.3$ m. In consequence, highly accurate and precise DEMs are needed to capture the small magnitude of glacier mass fluctuations. In Sect. 3.1.4 we reported that our DEMs errors are within previously reported errors of Pléiades DEMs used in geodetic mass balance assessments (e.g. Berthier et al., 2014; Denzinger et al., 2021; Falaschi et al., 2023). A fine coregistration between DEMs allows the minimization of vertical and horizontal biases. This is especially important when utilizing DEMs acquired when seasonal snow is present off-glacier, but difficult to circumvent when surveying mass balance during accumulation periods. Under these conditions, vertical shifts of up to 1 m measured over stable terrain have been reported (Beraud et al., 2023). Glaciers close to in-balance conditions such as Muztag Ata are particularly sensitive to an adequate quantification of systematic biases compared to glaciers with a much more negative mass budget, as a given small vertical DEM adjustment may shift from slightly negative to slightly positive elevation changes and vice-versa.

Around our reported DEM errors, the vertical precision (standard deviation on stable terrain) of our Pléiades time series in Muztag Ata and Western Nyainqêntanglha were 2.1 m and 1.2 m, respectively (Table 2). DEM precision can be also described using the normalized median absolute deviation (NMAD), which is less sensitive to outliers compared to standard deviation (Höhle and Höhle, 2009; Dehecq et al., 2016). Independently from seasonal snow conditions, the NMAD over off-glacier terrain was consistently around ± 1.3 m in Muztag Ata, and varied between ± 0.6 m and 1.2 m in Western Nyainqêntanglha (Table 2). These standard deviations and NMAD values are in the same order of magnitude as other very high resolution DEMs used in glaciological applications (Berthier et al., 2014; Belart et al., 2017; Rieg et al., 2018; Pelto et al., 2019; Beraud et al., 2023). Comparatively, the lower precision in the Muztag Ata DEMs result in relatively larger uncertainties in particular cases (Table 3). This becomes more problematic due to the short temporal baseline, and when glaciers are close to balanced conditions. With these caveats in mind, our annual and seasonal geodetic mass balance estimates were robust and consistent with the results retrieved from the Glacier Index (see Sect. 5.2), and hence can provide valuable insight into glacier accumulation regimes in High Mountain Asia.

5.2 Discerning accumulation regimes

The heterogeneity of regional climates displayed in High Mountain Asia, from monsoon-dominated regions with abundant precipitation occurring in the warm season to areas where westerly-induced, winter precipitation prevails, results in diverse glacier behaviour and sensitivity to climate drivers (Bolch et al., 2012; Sakai and Fujita, 2017). The Glacier Index of Huang et al. (2022) showed important discrepancies in accumulation regimes for some regions across High Mountain Asia between 2015 and 2018, as derived from gridded climate datasets, including APHRODITE, ERA5 and HARv2 products. In the specific case of the Muztagh Ata area, there is conflicting evidence about whether glaciers are of winter- or summer-accumulation type. Data from the nearest Taxikorgan station indicate that precipitation occurs mainly during summer time (April-September) (Zhu et al., 2018a), so that a summer accumulation regime may be assumed for the Muztagh Ata glaciers. Indeed, Zhu et al. (2018a, b) suggested that summer precipitation as the main driver of the Muztagh Ata No. 15 Glacier mass balance, and that higher amounts of solid precipitation in summer compared to the cold season were mostly responsible

for the overall positive budget between 1998 and 2012. However, these authors also showed that between 1980 and 1997, and in spite of heavier precipitation in summer compared to winter, other processes affecting glacier mass balance (e.g. snowmelt, sublimation) resulted in a more negative budget in the summer season.

In contrast to the Taxikorgan instrumental records in Muztag Ata, which show higher amounts of precipitation during the summer months, Maussion et al. (2014) used HAR data and found that overall, the Pamir region experiences mostly winter precipitation. Huang et al. (2022) found a conspicuous mismatch between the (transitional) accumulation regime derived from a) gridded climate datasets and climate stations off-glacier and b) the winter accumulation-type according to the SAR analyses. The authors noted the substantial differences in snow accumulation measured at ~7000 asl (605 mm a⁻¹) and those recorded at Taxikorgan station between 1960 and 2002 (60-70 mm; Duan et al., 2015), suggesting that the upper part of Muztag Ata, an anomalously high peak in the region (Seong et al., 2009), may be affected by a different atmospheric circulation system compared to the valley bottoms (i.e. at Taxikorgan station elevation). Sakai et al. (2015) used a summer- to annual-precipitation ratio from APHRODITE data and Temperature-Precipitation plots as indicators of glacier sensitivity to climate changes and categorised the glaciers in Eastern Pamir within the winter accumulation and less sensitive envelope. Although our Pléiades datasets allowed the retrieval of mass changes for two seasonal intervals only, the geodetic estimates of the glacier-wide Muztag Ata 2022 winter (+0.13 ±0.24 m w.e.) and summer (-0.35 ±0.15 m w.e.) mass balance favor a winter accumulation-type scenario for the Muztag Ata glaciers.

In stark contrast to Muztag Ata and Eastern Pamir, Western Nyainqêntanglha is located in a transitional area between the monsoon-dominated glaciers in the SE Tibetan Plateau and Himalaya and the westerlies-dominated regions to the Northwest. Here, glaciers show a low annual temperature range, are highly sensitive to temperature and precipitation changes, and are therefore prone to strong mass losses (Sakai et al., 2015). There is good consensus in characterizing most glaciers in the area as summer accumulation-type (Fujita and Ageta, 2000; Maussion et al., 2014; Sakai et al., 2015; Huang et al., 2022). The mass balance of Zhadang Glacier was found to be particularly sensitive to the onset of the Monsoon period (Kang et al., 2009; Mölg et al., 2012), though mid-latitude westerlies also drive its mass balance (Mölg et al., 2014). Our geodetic results for the Western Nyainqêntanglha glaciers showed high ablation rates prevailing over accumulation in the 2022 summer season (-0.63 ±0.07 m w.e.a⁻¹) and no mass recovery during the winter season (-0.03 ±0.27 m w.e.a⁻¹). However, whilst elevation gains are not visible in the uppermost reaches of the glaciers in the 2022 winter elevation change grid (Fig. 6d), the summer elevation change panel (Fig. 6e) does indeed show elevation changes that we attribute to snow accumulation. The 2009-2011 and 2008-2013 mass balance of Zhadang Glacier modelled by Zhang et al. (2013) and Zhu et al. (2018a) support our findings, indicating that in Western Nyainqêntanglha, accumulation through precipitation is higher in summer than in winter, but contemporary mass losses through runoff and evaporation can exceed mass gains and lead to a strongly negative mass balance.

5.2.1 Insight from the Glacier Index and climate records

The wet snow area- and firn area-ratio showed different patterns and trends in Muztag Ata compared to Western Nyainqêntanglha between 2020 and 2022 (Fig. 7). In Western Nyainqêntanglha, the annual wet snow area- are higher than the firn area-ratios, meaning that there is more accumulation in summer than in winter. The opposite was found in Muztag Ata, where the firn area-ratios are higher than the late summer wet snow area ratio. This is in turn indicative of higher accumulation during winter in Muztag Ata. In addition, the firn area-ratio is much lower in Western Nyainqêntanglha compared to Muztag Ata, and it is therefore not surprising that glacier mass balance has been much more negative for at least the last six decades (Bhattacharya et al., 2021 and references therein, see Table 4). These results agree well with our geodetic mass balance results for the annual and seasonal intervals in the 2020-2022 sampled period (Table 3), which support winter and summer accumulation-type regimes in Muztag Ata and Western Nyainqêntanglha, respectively.

Despite the overall good agreement between the proposed accumulation regime types based on both the geodetic and Glacier Index approaches, we report a minor caveat. On one hand, the wet snow-area ratio in Muztag Ata indicates 2022 as the year with the most negative mass balance year, whilst the geodetic mass balance suggests 2020. Differences in geodetic mass balance and wet snow area-ratios are nevertheless small between both years, and hence we attribute this discrepancy to the geodetic mass balance uncertainty. On the other hand, the wet snow area-ratio in Western Nyainqêntanglha indicates that the year 2022 should be the second most negative mass balance year in the study period (-0.50 ±0.17 m w.e.a⁻¹ as per our geodetic estimate), whereas the geodetic results suggest 2021 (-0.69 ±0.08 m w.e.) as the second most negative year. A possible explanation lies in the differences in the acquisition dates between the Pléiades and Sentinel-1 satellite images used in the geodetic and Glacier Index approaches. The Glacier Index utilizes SAR imagery acquired from July 20 to September 10 each year to guarantee wet snow conditions over accumulation areas (whereas in late September and October, snow may be dry for areas above 5,000 m asl due to low temperatures). On the contrary, the 2021 Pléiades scenes employed in the geodetic method were acquired on 30 September. Our climate data, however, shows that in Western

Nyainqentanglha, precipitation was higher in September [2022](#) compared to August in [2022](#), which will lead to a lower wet snow area–ratio.

Establishing a clear link between [glacier mass balance](#) and the investigated climate datasets and variables is a challenging task. Indeed, Huang et al. (2022) have shown how [both reanalyses](#) and instrumental records, such as the Taxikorgan climate station [in the Muztag Ata region](#), are difficult to link with glacier accumulation regimes in several regions throughout High Mountain Asia. [To understand the influence of precipitation and temperature on the response of the glaciers in the study sites, we used daily Global Precipitation Measurement \(GPM\) IMERG late run precipitation observations and monthly temperature estimates from ERA5 Land \(see Sect. S3.3 and S4.3\). Only the annual temperature anomalies in Western Nyainqentanglha](#) showed a significant correlation with mass balance, which is most probably due to the small number of mass balance observations ($n_{\max} = 7$ in the 1967–2022 correlation tests), [and possibly to the different duration of the sampled time intervals as well](#). In Muztag Ata, our results showed a stronger correlation between glacier mass budget and winter solid precipitation compared to air temperatures, though we stress here that neither correlation was significant. [The fact that winter snowfall appears to be a stronger mass balance driver in relation to air temperature in Muztag Ata is consistent with cold and dry environments, which are influenced by solid precipitation to a greater extent](#) (Zhu et al. 2018a, b).

We observed a different scenario in Western Nyainqentanglha. Correlations were much stronger (and significant in the case of annual temperature) between mass balance and temperature anomalies in comparison to solid precipitation. This is not surprising, since summer temperatures have a greater impact on glacier mass balance in more humid climatic regions such as Western Nyainqentanglha (Zhang et al., 2013, Bhattacharya et al., 2021). Mölg et al., (2014) showed how summer precipitation determines the annual mass balance of summer accumulation glaciers in the area. Summer temperature, however, modulates the solid/liquid precipitation ratio during the summer season, when most of the solid precipitation occurs in the region. [In conclusion, the intricacies and limitations of determining accumulation regimes over our investigated glaciers, based on the available climate records, clearly reveal the need for alternate or complementary approaches](#).

5.3 Mass balance differences between both study sites and in the long-term perspective

Our mass balance estimates between 2020 and 2022 for the Muztag Ata massif suggest a greater variability in mass budget than previously acknowledged in the region (Fig. 8a). During the past six decades, Bhattacharya et al. (2021) found mass balance rates between $+0.03 \pm 0.10$ m w.e. a^{-1} and -0.14 ± 0.10 m w.e. a^{-1} . On annual time steps, our values ranged between $+0.15 \pm 0.27$ m w.e. a^{-1} and -0.19 ± 0.14 m w.e. a^{-1} (Table 3). Averaged between 2020 and 2022, the glacier-wide mass balance of Muztag Ata (-0.07 ± 0.20 m w.e. a^{-1}) is similar to the 2013–2019 mean (-0.12 ± 0.09 m w.e. a^{-1} ; Bhattacharya et al., 2021). This suggests an ongoing [period](#) of slight mass loss in the Muztagh Ata massif for the last three years, meaning a continuation of the very slight mass loss rate observed for most of approximately the last decade (Bhattacharya et al., 2021). A number of studies (Holzer et al., 2015; Lv et al., 2020) report slight, but insignificant positive values between 1999 and 2015/2016, whilst Bhattacharya et al. (2021) found in-balance conditions since 1973.

On an individual glacier basis, glaciological measurements exist for the small (~ 1.1 km²) Muztag Ata No. 15 glacier (Fig. 1) between the years 2002 and 2014 (Yao et al., 2012 and unpublished data; Holzer et al., 2015). The mean 2005–2014 mass balance of this glacier was $+0.11$ m w.e. a^{-1} , which is similar to our $+0.12 \pm 0.28$ m w.e. a^{-1} 2020–2022 average.

In contrast to Muztag Ata, glaciers in the Western Nyainqentanglha range show much higher mass loss rates. The 2020 mass balance of -0.72 ± 0.16 m w.e. is not only much more negative than the -0.07 ± 0.20 w.e. a^{-1} 2020–2022 average found in Muztag Ata, but also represents [the highest recorded rate of ice mass loss](#) for the Western Nyainqentanglha area itself since the late 1960's (Fig. 8b). Indeed, the mass budget has not been more negative than -0.65 ± 0.08 m w.e. a^{-1} [over](#) approximately the last five decades (Hugonnet et al., 2021 and other sources; Table 4). It must be noted, however, that the sampled glaciers vary among these studies, and the mass budget could have differed if data from different temporal scales were considered.

Zhadang Glacier, which had a glaciological mass balance program running between 2006 and 2017 (Yao T, pers. com.), was one of the glaciers with the most negative mass budget in the Western Nyainqentanglha range. [The 2020–2022 mean mass balance was \$-0.95 \pm 0.25\$ m w.e. \$a^{-1}\$, depicting a high mass loss rate consistent with modelled estimates \(\$-1.06\$ m w.e. \$a^{-1}\$ between 2001 and 2011, Huintjes et al., 2015; \$-0.91\$ w.e. \$a^{-1}\$, between 2010 and 2012, Zhu et al., 2015; \$-1.20\$ m w.e. \$a^{-1}\$, between 2009 and 2011, Zhang et al., 2013\)](#). Surprisingly, though, remote sensing–based estimates have yielded significantly less negative mass balances encompassing longer but overlapping periods (-0.50 ± 0.17 m w.e. a^{-1} between 2000 and 2014, Li and Lin., 2017; -0.60 ± 0.19 m w.e. a^{-1} between 2000 and 2017, Ren et al., 2020). The average 2006–2017 in situ mass balance of Zhadang Glacier (-1.35 m w.e. a^{-1} ; Yao et al., 2012 and unpublished data) was found to be closer to our estimate, which may suggest that penetration correction for mass balance estimates based on DEMs originating from SAR data are inadequate in the area.




Figure 9. Comparison of geodetic glacier mass balance for the glacier-wide area and specific glaciers with available glaciological mass balance records in Muztag Ata (a) and Western Nyainqêntanglha (b).

Table 4. Mass balance estimates in Muztag Ata and Western Nyainqêntanglha based on optical and SAR-derived DEM differencing.

	Time period	Mass balance (m w.e.a ⁻¹)	source
Muztag Ata			
	1967-1973	-0.14 ±0.10	Bhattacharya et al. (2021)
	1973-2001	-0.01 ±0.06	
	2001-2009	+0.03 ±0.10	
	2009-2013	-0.08 ±0.12	
	2013-2019	-0.12 ±0.09	
	1967-2019	-0.06 ±0.07	
	1971/1976-1999	-0.15 ±0.18	Zhang et al. (2016)
	1999-2013/2014	-0.13 ±0.23	
	1971/1974-2013-2014	-0.14 ±0.10	
	1973-1999	-0.04 ±0.42	Holzer et al. (2015)
	1999-2009	+0.04 ± 0.45	
	2009-2013	-0.07 ±0.53	
	1973-2009	-0.03 ±0.33	
	2000-2004	+0.19 ±0.08	Hugonnet et al. (2021)
	2005-2009	+0.13 ±0.06	
	2010-2014	+0.09 ±0.07	
	2014-2019	+0.07 ±0.08	
	2000-2019	+0.13 ±0.10	
	2000-2015/2016	+0.16 ±0.03	Lv et al. (2020)
	2000-2018	+0.21 ±0.06	Shean et al. (2020)
W. Nyainqêntanglha			
	1968-1976	-0.27 ±0.11	Bhattacharya et al. (2021)
	1976-2001	-0.24 ±0.13	
	2001-2012	-0.41 ±0.11	
	2012-2018	-0.47 ±0.15	
	2018-2019	-0.39 ±0.18	
	1968-2019	-0.32 ±0.09	
	2000-2004	-0.31 ±0.08	Hugonnet et al. (2021)
	2005-2009	-0.44 ±0.06	
	2010-2014	-0.58 ±0.07	
	2014-2019	-0.65 ±0.08	
	2000-2019	-0.49 ±0.17	
	2000-2018	-0.51 ±0.11	Shean et al. (2020)
(total W. Nyainqêntanglha)	2000-2020	-0.37 ±0.12	Wang et al. (2023)
(total W. Nyainqêntanglha)	2000-2017	-0.30 ±0.19	Ren et al. (2020)
(total W. Nyainqêntanglha)	1975-2000	-0.25 ±0.15	Zhou et al. (2018)
(total W. Nyainqêntanglha)	2000-2013/2014	-0.23 ±0.13	Li and Lin (2017)
(total W. Nyainqêntanglha)	2000-2013/2014	-0.30 ±0.07	Zhang and Zhang (2017)

Conclusions

740

In this study we have assessed the capability of very high-resolution Pléiades DEMs to quantify glacier mass balance over short (annual to seasonal) time steps over regions in High Mountain Asia that have shown contrasting mass balance trends in the last few decades. We find intrinsic DEM quality to be the main source of uncertainty in geodetic mass balance estimates, rather than varying seasonal snow conditions across the two study sites. Indeed, the mean vertical precision (NMAD) of Muztag Ata and Western Nyainqêntanglha DEMs were ±1.3 m and ±0.6 m to ±1.2 m, respectively. These levels of precision are in line with previously reported values of other very-high resolution DEMs used in similar geodetic mass balance estimations worldwide.

745

Two main conclusions can be drawn from the internal consistency tests (utilized to evaluate the differences between accumulated vs. the sum of individual periods). On one side, the tests confirm that the usage of a semi-automated approach such as ASMAG for the identification of different glacier surfaces (ice, snow/firn), and thus the distribution of assumed material densities, is a robust approach when quantifying glacier mass balance at a

750

multi-glacier scale. Greater differences on individual glaciers, however, suggest that manual mapping by visual interpretation might be preferable, especially on small glaciers. Concurrently, the tests reaffirm that using time-weighted densities reduced the residuals between accumulated vs. added mass budgets, stressing the necessity of

755

Mean annual (-0.07 ± 0.20 m w.e. a⁻¹) mass balance estimates in Muztagh Ata between 2020 and 2022 point at the continuation of the slight mass loss trend after a period of apparently balanced conditions. On the contrary, the glacier mass balance in Western Nyainqêntanglha reached a negative peak of -0.72 ± 0.16 m w.e. a⁻¹ in 2022, which represents a new maximum over the last six decades. The analysis of ERA5-land 1950-2021 temperature and GPM 2001-2021 solid precipitation anomalies confirm that the years 2020 and 2021 as, a) average years in terms of air temperature and snowfall in Muztagh Ata, and b) particularly warm and dry years in Western Nyainqêntanglha.

760

The 2022 winter and summer mass balance estimates ($+0.13 \pm 0.24$ m w.e. and -0.35 ± 0.15 m w.e., respectively) in Muztagh Ata suggest a winter accumulation type, whilst mass losses of -0.03 ± 0.27 m w.e. and -0.63 ± 0.07 m w.e. in the winter and summer seasons, respectively, confirm a summer accumulation-type regime in Western Nyainqêntanglha, with ablation prevailing over accumulation in the summer (ablation) season. With the SAR-based Glacier Index, we indirectly validated our geodetic mass balance estimates and the derived inferences that were made in relation to glacier accumulation types in Muztagh Ata and Western Nyainqêntanglha. Whilst the Index does not provide a specific mass balance estimate per se that can be directly compared against geodetic (or glaciological) results, it can provide further insight into accumulation regimes in poorly known regions. Moreover, it fully bypasses the need for instrumental or reanalysis records, which are often unavailable or unrepresentative of climate conditions at glacier locations. Further geodetic and glaciological mass balance measurements in combination with such an Index will open new possibilities in glaciological research.

765

770

Based on the above, we conclude that our DEM time series and mass budget estimates proved to be consistent for making reliable, short-term estimations of glacier mass balance using a remote sensing-based approach. The ever-increasing number and availability of very-high resolution optical satellites with stereo capability and relatively short revisit time, such as [Pléiades](#), [WorldView 1-2](#), [SPOT6-7](#) or the recently launched [ALOS-3](#), will allow for increasing the number of glaciers in isolated regions that can be readily monitored.

775

780

Code availability

We are grateful to Etienne Berthier for kindly providing the along-track bias correction and the elevation change uncertainty “patch” tools.

785

The DEM generation code from Shean et al. (2016) is available from the corresponding github repository at <https://github.com/NeoGeographyToolkit/StereoPipeline>. The DEM coregistration code (Nuth and Kääb, 2011) is available at <https://github.com/GeoUtils/geoutils>.

Data availability

790

The Sentinel-2 scenes were obtained from the USGS EarthExplorer data poll (<https://earthexplorer.usgs.gov/>) and the ESA Copernicus Open Access Hub (<https://scihub.copernicus.eu/>). The Sentinel-1 and Landsat OLI scenes used in the Glacier Index are available online on Google Earth Engine through the Google Cloud Storage and the Google Cloud public data program. The elevation change grids from Bhattacharya et al. (2021) are available from PANGAEA at <https://www.pangaea.de> and <https://www.mountrys.org/datasets/>. The elevation change maps from Hugonnet et al., 2021 are publicly available at <https://doi.org/10.6096/13>. The ERA5-Land data was downloaded from the Copernicus Climate Data Store at <https://cds.climate.copernicus.eu/>.

795

Author contribution

800

DF and TB designed the study. OK contributed to the study design. DF led the study, processed and analyzed data and wrote the manuscript. AB, KM and LH, PR processed and analyzed data. AB, KM, LH, OK and TB contributed to the writing of the manuscript. TB and TY secured the funding. The order of authors from third to seventh is in alphabetical order.

805

Competing interests

Some authors are members of the editorial board of The Cryosphere. The peer-review process was guided by an independent editor, and the authors have also no other competing interests to declare.

810

Acknowledgements

We are grateful to CNES/Airbus DS for the provision of the Pléiades satellite data within the ISIS program for reduced costs. Pléiades © CNES 2020/2021/2022 and AIRBUS DS. We would like to thank Christine Baron, Sylvie Boureausseau, and the whole Airbus Intelligence Team for their assistance during the acquisition of the Pléiades imagery. AB acknowledges funding by the Science & Engineering Research Board (SERB), Department of Science & Technology (DST), India (grant no. CRG/2021/002450). [Special thanks to editor Regula Frauenfelder for handling the challenging task of finding reviewers and to Cesar Deschamps-Berger for his thorough review of the original manuscripts.](#)

Funding

This study was supported by the Strategic Priority Research Program of Chinese Academy of Sciences (XDA20100300) and the Swiss National Science Foundation (200021E_177652/1) within the framework of the DFG Research Unit GlobalCDA (FOR2630) and benefited from the research cooperation within the Dragon 5 program supported by ESA and NRSCC (4000136930/22/I-NB).

References

Abdel Jaber, W., Rott, H., Floricioiu, D., Wuite, J., and Miranda, N.: Heterogeneous spatial and temporal pattern of surface elevation change and mass balance of the Patagonian ice fields between 2000 and 2016, *Cryosphere*, 13(9), 2511–2535, <https://doi.org/10.5194/tc-13-2511-2019>, 2019.

Andreassen, L. M., Elvehøy, H., Kjøllmoen, B., and Engeset, R. V.: Reanalysis of long-term series of glaciological and geodetic mass balance for 10 Norwegian glaciers, *Cryosphere*, 10(2), 535–552, <https://doi.org/10.5194/tc-10-535-2016>, 2016.

Armstrong, W. H., Polashenski, D., Truffer, M., Horne, G., Hanson, J. B., Hawley, R. L., Hengst, A. M., Vowels, L., Menounos, B., and Wychen, W. V.: Declining basal motion dominates the long-term slowing of Athabasca Glacier, Canada, *J. Geophys. Res. Earth. Surf.*, 127(10), <https://doi.org/10.1029/2021JF006439>, 2022.

Belart, J. M. C., Berthier, E., Magnússon, E., Anderson, L. S., Pálsson, F., Thorsteinsson, T., Howat, I. M., Aðalgeirsdóttir, G., Jóhannesson, T., and Jarosch, A. H.: Winter mass balance of Drangajökull ice cap (NW Iceland) derived from satellite sub-meter stereo images, *Cryosphere*, 11(3), 1501–1517, <https://doi.org/10.5194/tc-11-1501-2017>, 2017.

Belart, J. M. C., Magnússon, E., Berthier, E., Pálsson, F., Aðalgeirsdóttir, Gu., and Jóhannesson, T.: The geodetic mass balance of Eyjafjallajökull ice cap for 1945–2014: Processing guidelines and relation to climate, *J. Glaciol.* 65(251), 395–409, <https://doi.org/10.1017/jog.2019.16>, 2019.

Beraud, L., Cusicanqui, D., Rabatel, A., Brun, F., Vincent, C., and Six, D.: Glacier-wide seasonal and annual geodetic mass balances from Pléiades stereo images: Application to the Glacier d’Argentière, French Alps, *J. Glaciol.*, 69(275), 525–537, <https://doi.org/10.1017/jog.2022.79>, 2023.

Berthier, E., and Brun, F.: Karakoram geodetic glacier mass balances between 2008 and 2016: Persistence of the anomaly and influence of a large rock avalanche on Siachen Glacier, *J. Glaciol.*, 65(251), 494–507, <https://doi.org/10.1017/jog.2019.32>, 2019.

Berthier, E., Vincent, C., Magnússon, E., Gunnlaugsson, Á. Þ., Pitte, P., Le Meur, E., Masiokas, M., Ruiz, L., Pálsson, F., Belart, J. M. C., and Wagnon, P.: Glacier topography and elevation changes derived from Pléiades sub-meter stereo images, *Cryosphere*, 8(6), 2275–2291, <https://doi.org/10.5194/tc-8-2275-2014>, 2014.

Berthier, E., Cabot, V., Vincent, C., and Six, D.: Decadal region-wide and glacier-wide mass balances derived from multi-temporal ASTER satellite Digital Elevation Models. Validation over the Mont-Blanc area, *Front. Earth Sci.*, 4, <https://doi.org/10.3389/feart.2016.00063>, 2016.

Bhattacharya, A., Bolch, T., Mukherjee, K., King, O., Menounos, B., Kapitsa, V., Neckel, N., Yang, W., and Yao, T.: High Mountain Asian glacier response to climate revealed by multi-temporal satellite observations since the 1960s, *Nat. Commun.*, 12(1), 4133, <https://doi.org/10.1038/s41467-021-24180-y>, 2021.

Bippus, G.: Characteristics of summer snow areas on glaciers observed by means of Landsat data, Ph. D. thesis, University of Innsbruck, Austria, 2011.

Bolch, T., Yao, T., Kang, S., Buchroithner, M. F., Scherer, D., Maussion, F., Huintjes, E., and Schneider, C.: A glacier inventory for the western Nyainqentanglha Range and the Nam Co Basin, Tibet, and glacier changes 1976–2009, *Cryosphere*, 4(3), 419–433, <https://doi.org/10.5194/tc-4-419-2010>, 2010.

Bolch, T., Kulkarni, A., Käab, A., Huggel, C., Paul, F., Cogley, J. G., Frey, H., Kargel, J. S., Fujita, K., Scheel, M., Bajracharya, S., and Stoffel, M.: The state and fate of Himalayan glaciers, *Science*, 336(6079), 310–314, <https://doi.org/10.1126/science.1215828>, 2012.

Bolch, T., Rohrbach, N., Kutuzov, S., Robson, B. A., and Osmonov, A.: Occurrence, evolution and ice content of ice-debris complexes in the Ak-Shiirak, Central Tien Shan revealed by geophysical and remotely-sensed

- investigations: Ice-debris complexes in Ak-Shiirak, *Earth Surf. Process Landf.*, 44(1), 129–143, <https://doi.org/10.1002/esp.4487>, 2019.
- 875 Braun, M. H., Malz, P., Sommer, C., Fariás-Barahona, D., Sauter, T., Casassa, G., Soruco, A., Skvarca, P., and Seehaus, T. C.: Constraining glacier elevation and mass changes in South America, *Nat. Clim. Chang.*, 9(2), 130–136, <https://doi.org/10.1038/s41558-018-0375-7>, 2019.
- Brun, F., Berthier, E., Wagnon, P., Kääb, A., and Treichler, D.: A spatially resolved estimate of High Mountain Asia glacier mass balances from 2000 to 2016, *Nat. Geosci.*, 10(9), 668–673. <https://doi.org/10.1038/ngeo2999>, 2017.
- 880 Brun, F., Wagnon, P., Berthier, E., Jomelli, V., Maharjan, S. B., Shrestha, F., and Kraaijenbrink, P. D. A.: Heterogeneous influence of glacier morphology on the mass balance variability in High Mountain Asia, *J. Geophys. Res. Earth Surf.*, 2018JF004838, <https://doi.org/10.1029/2018JF004838>, 2019.
- Burns, P., and Nolin, A.: Using atmospherically-corrected Landsat imagery to measure glacier area change in the Cordillera Blanca, Peru from 1987 to 2010, *Remote Sens. Environ.*, 140, 165–178. <https://doi.org/10.1016/j.rse.2013.08.026>, 2014.
- 885 Caidong, C., and Sorteberg, A.: Modelled mass balance of Xibu glacier, Tibetan Plateau: sensitivity to climate change, *J. Glaciol.*, 56(196), 235–248. <https://doi.org/10.3189/002214310791968467>, 2010.
- Clarke, G. K. C., Anslow, F. S., Jarosch, A. H., Radić, V., Menounos, B., Bolch, T., and Berthier, E.: Ice volume and subglacial topography for Western Canadian glaciers from mass balance fields, Thinning Rates, and a Bed Stress Model, *J. Clim.*, 26(12), 4282–4303, <https://doi.org/10.1175/JCLI-D-12-00513.1>, 2013.
- 890 Cogley, G. J.: Geodetic and direct mass-balance measurements: Comparison and joint analysis, *Ann. Glaciol.*, 50(50), 96–100. <https://doi.org/10.3189/172756409787769744>, 2009.
- Cogley, J. G., Hock, R., Rasmussen, L. A., Arendt, A. A., Bauder, A., Braithwaite, R. J., Jansson, P., Kaser, G., Möller, M., Nicholson, L., and Zemp, M.: Glossary of glacier mass balance and related terms, IHP-VII Technical Documents in Hydrology No. 86, IACS Contribution No. 2, UNESCO-IHP, Paris.
- 895 Davaze, L., Rabatel, A., Dufour, A., Hugonnet, R., and Arnaud, Y.: Region-wide annual glacier surface mass balance for the European Alps from 2000 to 2016, *Front. Earth Sci.*, 8, 149, <https://doi.org/10.3389/feart.2020.00149>, 2020.
- Dehecq, A., Millan, R., Berthier, E., Gourmelen, N., Trouve, E., and Vionnet, V.: Elevation changes inferred from TanDEM-X data over the Mont-Blanc area: Impact of the X-Band Interferometric Bias, *IEEE J. Sel. Top. Appl. Earth Obs.*, 9(8), 3870–3882, <https://doi.org/10.1109/JSTARS.2016.2581482>, 2016.
- 900 Dehecq, A., Gourmelen, N., Gardner, A. S., Brun, F., Goldberg, D., Nienow, P. W., Berthier, E., Vincent, C., Wagnon, P., and Trouvé, E.: Twenty-first century glacier slowdown driven by mass loss in High Mountain Asia, *Nat. Geosci.*, 12(1), 22–27, <https://doi.org/10.1038/s41561-018-0271-9>, 2019.
- 905 Denzinger, F., Machguth, H., Barandun, M., Berthier, E., Girod, L., Kronenberg, M., Usabaliev, R., and Hoelzle, M.: Geodetic mass balance of Abramov Glacier from 1975 to 2015, *J. Glaciol.*, 67(262), 331–342, <https://doi.org/10.1017/jog.2020.108>, 2021.
- Deschamps-Berger, C., Gascoïn, S., Berthier, E., Deems, J., Gutmann, E., Dehecq, A., Shean, D., and Dumont, M.: Snow depth mapping from stereo satellite imagery in mountainous terrain: evaluation using airborne laser-scanning data, *Cryosphere*, 14(9), 2925–2940, <https://doi.org/10.5194/tc-14-2925-2020>, 2020.
- 910 Duan, K., Xu, B., and Wu, G.: Snow accumulation variability at altitude of 7010 m a.s.l. in Muztag Ata Mountain in Pamir Plateau during 1958–2002, *J. Hydrol.*, 531, 912–918, <https://doi.org/10.1016/j.jhydrol.2015.10.013>, 2015.
- Dussaillant, I., Berthier, E., Brun, F., Masiokas, M., Hugonnet, R., Favier, V., Rabatel, A., Pitte, P., and Ruiz, L.: Two decades of glacier mass loss along the Andes, *Nat. Geosci.*, 12(10), 802–808, <https://doi.org/10.1038/s41561-019-0432-5>, 2019.
- Falaschi, D., Rivera, A., Lo Vecchio Repetto, A., Moragues, S., Villalba, R., Rastner, P., Zeller, J., and Salcedo, A. P.: Evolution of surface characteristics of three debris-covered glaciers in the Patagonian Andes from 1958 to 2020, *Front. Earth Sci.*, 9, 671854, <https://doi.org/10.3389/feart.2021.671854>, 2021.
- 920 Falaschi, D., Berthier, E., Belart, J. M. C., Bravo, C., Castro, M., Durand, M., and Villalba, R.: Increased mass loss of glaciers in Volcán Domuyo (Argentinian Andes) between 1962 and 2020, revealed by aerial photos and satellite stereo imagery, *J. Glaciol.*, 69(273), 40–56, <https://doi.org/10.1017/jog.2022.43>, 2023.
- Farinotti, D., Immerzeel, W. W., de Kok, R. J., Quincey, D. J., and Dehecq, A.: Manifestations and mechanisms of the Karakoram glacier anomaly, *Nat. Geosci.*, 13(1), 8–16, <https://doi.org/10.1038/s41561-019-0513-5>, 2020.
- 925 Fischer, M., Huss, M., Kummert, M., and Hoelzle, M.: Application and validation of long-range terrestrial laser scanning to monitor the mass balance of very small glaciers in the Swiss Alps, *Cryosphere*, 10(3), 1279–1295, <https://doi.org/10.5194/tc-10-1279-2016>, 2016.
- Fujita, K., and Ageta, Y.: Effect of summer accumulation on glacier mass balance on the Tibetan Plateau revealed by mass-balance model, *J. Glaciol.*, 46(153), 244–252, <https://doi.org/10.3189/172756500781832945>, 2000.
- 930

- Gardelle, J., Berthier, E., Arnaud, Y., and Kääb, A.: Region-wide glacier mass balances over the Pamir-Karakoram-Himalaya during 1999-2011, *Cryosphere*, 7(4), 1263–1286, <https://doi.org/10.5194/tc-7-1263-2013>, 2013.
- 935 Girod, L., Nuth, C., Kääb, A., McNabb, R., and Galland, O.: MMASTER: Improved ASTER DEMs for elevation change monitoring, *Remote Sens.*, 9(7), 704, <https://doi.org/10.3390/rs9070704>, 2017.
- Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., and Moore, R.: Google Earth Engine: Planetary-scale geospatial analysis for everyone, *Remote Sens. Environ.*, 202, 18–27, <https://doi.org/10.1016/j.rse.2017.06.031>, 2017.
- 940 Gleyzes, M. A., Perret, L., Kubik, P.: Pléiades system architecture and main performances, in: International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences, Volume XXXIX-B1, XXII ISPRS Congress, Melbourne, Australia, 25 August – 01 September 2012, 2012.
- Guo, Z., Wang, N., Kehrwald, N. M., Mao, R., Wu, H., Wu, Y., and Jiang, X.: Temporal and spatial changes in Western Himalayan firn line altitudes from 1998 to 2009, *Glob. Planet. Change*, 118, 97–105, <https://doi.org/10.1016/j.gloplacha.2014.03.012>, 2014.
- 945 Höhle, J., and Höhle, M.: Accuracy assessment of digital elevation models by means of robust statistical methods. *ISPRS J. Photogramm. Remote Sens.*, 64(4), 398–406. <https://doi.org/10.1016/j.isprsjprs.2009.02.003>, 2009.
- Holzer, N., Vijay, S., Yao, T., Xu, B., Buchroithner, M., and Bolch, T.: Four decades of glacier variations at Muztag Ata (eastern Pamir): a multi-sensor study including Hexagon KH-9 and Pléiades data, *Cryosphere*, 9(6), 2071–2088, <https://doi.org/10.5194/tc-9-2071-2015>, 2015.
- 950 Huang, L., Li, Z., Tian, B., Chen, Q., and Zhou, J.: Monitoring glacier zones and snow/firn line changes in the Qinghai–Tibetan Plateau using C-band SAR imagery, *Remote Sens. Environ.*, 137, 17–30, <https://doi.org/10.1016/j.rse.2013.05.016>, 2013.
- 955 Huang, L., Hock, R., Li, X., Bolch, T., Yang, K., Wang, N., Yao, T., Zhou, J., Dou, C., and Li, Z.: Winter accumulation drives the spatial variations in glacier mass balance in High Mountain Asia, *Sci. Bull.*, S2095927322003644, <https://doi.org/10.1016/j.scib.2022.08.019>, 2022.
- Huintjes, E., Sauter, T., Schröter, B., Maussion, F., Yang, W., Kropáček, J., Buchroithner, M., Scherer, D., Kang, S., and Schneider, C.: Evaluation of a coupled snow and energy balance model for Zhadang Glacier, Tibetan Plateau, using glaciological measurements and time-lapse photography, *Arct. Antarct. Alp. Res.*, 47(3), 573–590, <https://doi.org/10.1657/AAAR0014-073>, 2015.
- 960 Huss, M.: Density assumptions for converting geodetic glacier volume change to mass change, *Cryosphere*, 7(3), 877–887, <https://doi.org/10.5194/tc-7-877-2013>, 2013.
- Huss, M., Sold, L., Hoelzle, M., Stokvis, M., Salzmann, N., Farinotti, D., and Zemp, M.: Towards remote monitoring of sub-seasonal glacier mass balance, *Ann. Glaciol.*, 54(63), 75–83, <https://doi.org/10.3189/2013AoG63A427>, 2013.
- 965 Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., Farinotti, D., Huss, M., Dussailant, I., Brun, F., and Kääb, A.: Accelerated global glacier mass loss in the early twenty-first century, *Nature*, 592(7856), 726–731, <https://doi.org/10.1038/s41586-021-03436-z>, 2021.
- Hugonnet, R., Brun, F., Berthier, E., Dehecq, A., Mannerfelt, E. S., Eckert, N., and Farinotti, D.: Uncertainty analysis of digital elevation models by spatial inference from stable terrain, *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.*, 1–17, <https://doi.org/10.1109/JSTARS.2022.3188922>, 2022.
- 970 Immerzeel, W.W., Lutz, A.F., Andrade, M. et al.: Importance and vulnerability of the world’s water towers, *Nature*, 577, 364–369, <https://doi.org/10.1038/s41586-019-1822-y>, 2020.
- 975 Kääb, A., Treichler, D., Nuth, C., and Berthier, E.: Brief Communication: Contending estimates of 2003–2008 glacier mass balance over the Pamir–Karakoram–Himalaya, *Cryosphere*, 9(2), 557–564, <https://doi.org/10.5194/tc-9-557-2015>, 2015.
- Kääb, A., Winsvold, S., Altena, B., Nuth, C., Nagler, T., and Wuite, J.: Glacier remote sensing using Sentinel-2. Part I: radiometric and geometric performance, and application to ice velocity, *Remote Sens.*, 8(7), 598, <https://doi.org/10.3390/rs8070598>, 2016.
- 980 Kang, S., Chen, F., Gao, T., Zhang, Y., Yang, W., Yu, W., and Yao, T.: Early onset of rainy season suppresses glacier melt: A case study on Zhadang glacier, Tibetan Plateau, *J. Glaciol.*, 55(192), 755–758, <https://doi.org/10.3189/002214309789470978>, 2009.
- King, O., Bhattacharya, A., Bhambri, R., and Bolch, T.: Glacial lakes exacerbate Himalayan glacier mass loss, *Sci. Rep.*, 9(1), 18145, <https://doi.org/10.1038/s41598-019-53733-x>, 2019.
- 985 King, O., Bhattacharya, A., and Bolch, T.: The presence and influence of glacier surging around the Geladandong ice caps, North East Tibetan Plateau, *Adv. Clim. Chang. Res.*, 12(3), 299–312, <https://doi.org/10.1016/j.accre.2021.05.001>, 2021.
- Koblet, T., Gärtner-Roer, I., Zemp, M., Jansson, P., Thee, P., Haeberli, W., and Holmlund, P.: Reanalysis of multi-temporal aerial images of Storglaciären, Sweden (1959-99) - Part 1: determination of length, area, and volume changes, *Cryosphere*, 4(3), 333–343, <https://doi.org/10.5194/tc-4-333-2010>, 2010.
- 990

- Li, G., and Lin, H.: Recent decadal glacier mass balances over the Western Nyainqêntanglha Mountains and the increase in their melting contribution to Nam Co Lake measured by differential bistatic SAR interferometry, *Glob. Planet. Change*, 149, 177–190, <https://doi.org/10.1016/j.gloplacha.2016.12.018>, 2017.
- 995 Liang, Q., Wang, N., Yang, X., Chen, A., Hua, T., Li, Z., and Yang, D.: The eastern limit of ‘Kunlun-Pamir-Karakoram Anomaly’ reflected by changes in glacier area and surface elevation, *J. Glaciol.*, 68(272), 1167–1176, <https://doi.org/10.1017/jog.2022.30>, 2022.
- Luo, W., Zhang, G., Chen, W., and Xu, F.: Response of glacial lakes to glacier and climate changes in the Western Nyainqêntanglha range, *Sci. Total Environ.*, 735, 139607, <https://doi.org/10.1016/j.scitotenv.2020.139607>, 2020.
- 1000 Lv, M., Quincey, D. J., Guo, H., King, O., Liu, G., Yan, S., Lu, X., and Ruan, Z.: Examining geodetic glacier mass balance in the eastern Pamir transition zone, *J. Glaciol.*, 66(260), 927–937, <https://doi.org/10.1017/jog.2020.54>, 2020.
- Maussion, F., Scherer, D., Mölg, T., Collier, E., Curio, J., and Finkelnburg, R.: Precipitation seasonality and variability over the Tibetan Plateau as resolved by the High Asia Reanalysis, *J. Clim.*, 27(5), 1910–1927, <https://doi.org/10.1175/JCLI-D-13-00282.1>, 2014.
- 1005 McNabb, R., Nuth, C., Kääh, A., and Girod, L.: Sensitivity of glacier volume change estimation to DEM void interpolation, *Cryosphere*, 13(3), 895–910, <https://doi.org/10.5194/tc-13-895-2019>, 2019.
- Mölg, T., Maussion, F., Yang, W., and Scherer, D.: The footprint of Asian monsoon dynamics in the mass and energy balance of a Tibetan glacier, *Cryosphere*, 6(6), 1445–1461, <https://doi.org/10.5194/tc-6-1445-2012>, 2012.
- 1010 Mölg, T., Maussion, F., and Scherer, D.: Mid-latitude westerlies as a driver of glacier variability in monsoonal High Asia, *Nat. Clim. Chang.*, 4(1), 68–73, <https://doi.org/10.1038/nclimate2055>, 2014.
- Mukherjee, K., Menounos, B., Shea, J., Morteza pour, M., Ednie, M., and Demuth, M. N.: Evaluation of surface mass-balance records using geodetic data and physically-based modelling, *Place and Peyto glaciers, western Canada*, *J. Glaciol.*, 1–18, <https://doi.org/10.1017/jog.2022.83>, 2022.
- 1015 Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta, S., Choulga, M., Harrigan, S., Hersbach, H., Martens, B., Miralles, D. G., Piles, M., Rodríguez-Fernández, N. J., Zsoter, E., Buontempo, C., and Thépaut, J.-N.: ERA5-Land: A state-of-the-art global reanalysis dataset for land applications, *Earth System Science Data*, 13(9), 4349–4383, <https://doi.org/10.5194/essd-13-4349-2021>, 2021.
- Nuth, C., and Kääh, A.: Co-registration and bias corrections of satellite elevation data sets for quantifying glacier thickness change, *Cryosphere*, 5(1), 271–290, <https://doi.org/10.5194/tc-5-271-2011>, 2011.
- 1020 Nuth, C., Schuler, T. V., Kohler, J., Altena, B., and Hagen, J. O.: Estimating the long-term calving flux of Kronebreen, Svalbard, from geodetic elevation changes and mass-balance modeling. *J. Glaciol.*, 58(207), 119–133. <https://doi.org/10.3189/2012JoG11J036>, 2012.
- Otsu, N.: A Threshold Selection Method from Gray-Level Histograms, *IEEE Trans. Syst. Man Cybern. Syst.*, 9(1), 62–66, <https://doi.org/10.1109/TSMC.1979.4310076>, 1979.
- 1025 Paul, F., Bolch, T., Kääh, A., et al.: The glaciers climate change initiative: methods for creating glacier area, elevation change and velocity products, *Remote Sens. Environ.*, 162, 408–426, <https://doi.org/10.1016/j.rse.2013.07.043>, 2015.
- Pelto, B. M., Menounos, B., and Marshall, S. J.: Multi-year evaluation of airborne geodetic surveys to estimate seasonal mass balance, Columbia and Rocky Mountains, Canada, *Cryosphere*, 13(6), 1709–1727, <https://doi.org/10.5194/tc-13-1709-2019>, 2019.
- 1030 Racoviteanu, A. E., Rittger, K., and Armstrong, R.: An automated approach for estimating snowline altitudes in the Karakoram and Eastern Himalaya from Remote Sensing, *Front. Earth Sci.*, 7, 220, <https://doi.org/10.3389/feart.2019.00220>, 2019.
- 1035 Rastner, P., Prinz, R., Notarnicola, C., Nicholson, L., Sailer, R., Schwaizer, G., and Paul, F.: On the automated mapping of snow cover on glaciers and calculation of snow line altitudes from multi-temporal Landsat data, *Remote Sens.*, 11(12), 1410, <https://doi.org/10.3390/rs11121410>, 2019.
- Ren, S., Menenti, M., Jia, L., Zhang, J., Zhang, J., and Li, X.: Glacier mass balance in the Nyainqêntanglha Mountains between 2000 and 2017 retrieved from ZiYuan-3 stereo images and the SRTM DEM, *Remote Sens.*, 12(5), 864, <https://doi.org/10.3390/rs12050864>, 2020.
- 1040 RGI Consortium: Randolph Glacier Inventory—A dataset of global glacier outlines, Version 6 [data set], <https://doi.org/10.7265/4M1F-GD79>, 2017.
- Rieg, L., Klug, C., Nicholson, L., & Sailer, R.: Pléiades Tri-Stereo Data for Glacier Investigations—Examples from the European Alps and the Khumbu Himal, *Remote Sens.*, 10(10), 1563, <https://doi.org/10.3390/rs10101563>, 2018
- 1045 Sakai, A., Nuimura, T., Fujita, K., Takenaka, S., Nagai, H., and Lamsal, D.: Climate regime of Asian glaciers revealed by GAMDAM glacier inventory, *Cryosphere*, 9(3), 865–880, <https://doi.org/10.5194/tc-9-865-2015>, 2015.
- Sakai, A., and Fujita, K.: Contrasting glacier responses to recent climate change in high-mountain Asia, *Sci. Rep.*, 7(1), 13717, <https://doi.org/10.1038/s41598-017-14256-5>, 2017.
- 1050

- Scherler, D., Wulf, H., and Gorelick, N.: Global assessment of supraglacial debris-cover extents, *Geophys. Res. Lett.*, 45(21), 11,798–11,805, <https://doi.org/10.1029/2018GL080158>, 2018.
- Seong, Y. B., Owen, L. A., Yi, C., Finkel, R. C., and Schoenbohm, L.: Geomorphology of anomalously high glaciated mountains at the northwestern end of Tibet: Muztag Ata and Kongur Shan, *Geomorphology*, 103(2), 227–250, <https://doi.org/10.1016/j.geomorph.2008.04.025>, 2009.
- 1055 Shean, D. E., Alexandrov, O., Moratto, Z. M., Smith, B. E., Joughin, I. R., Porter, C., and Morin, P.: An automated, open-source pipeline for mass production of digital elevation models (DEMs) from very-high-resolution commercial stereo satellite imagery, *ISPRS J. Photogramm. Remote Sens.*, 116, 101–117, <https://doi.org/10.1016/j.isprsjprs.2016.03.012>, 2016.
- 1060 Shean, D. E., Bhushan, S., Montesano, P., Rounce, D. R., Arendt, A., and Osmanoglu, B.: A systematic, regional assessment of High Mountain Asia glacier mass balance, *Front. Earth Sci.*, 7, 363, <https://doi.org/10.3389/feart.2019.00363>, 2020.
- Shi, J., and Menenti, M.: Monitoring recent variations of the movements on the polythermal glaciers -a case study in the Nyainqêntanglha Mountains, in: 2013 IEEE International Geoscience and Remote Sensing Symposium - IGARSS, 2013, 3622–3625, <https://doi.org/10.1109/IGARSS.2013.6723614>, 2013.
- 1065 Sommer, C., Malz, P., Seehaus, T. C., Lippl, S., Zemp, M., and Braun, M. H.: Rapid glacier retreat and downwasting throughout the European Alps in the early 21st century, *Nat. Commun.*, 11(1), 3209, <https://doi.org/10.1038/s41467-020-16818-0>, 2020.
- 1070 Vishwakarma, B. D., Ramsankaran, R., Azam, Mohd. F. et al.: Challenges in understanding the variability of the cryosphere in the Himalaya and its impact on regional water Resources, *Front. Water*, 4, 909246, <https://doi.org/10.3389/frwa.2022.909246>, 2022.
- Wagnon, P., Brun, F., Khadka, A., Berthier, E., Shrestha, D., Vincent, C., Arnaud, Y., Six, D., Dehecq, A., Ménégou, M., and Jomelli, V.: Reanalysing the 2007–19 glaciological mass-balance series of Mera Glacier, Nepal, Central Himalaya, using geodetic mass balance, *J. Glaciol.*, 67(261), 117–125, <https://doi.org/10.1017/jog.2020.88>, 2021.
- 1075 Wang, S., Liu, J., Pritchard, H. D., Ke, L., Qiao, X., Zhang, J., Xiao, W., and Zhou, Y.: Characterizing 4 decades of accelerated glacial mass loss in the West Nyainqêntanglha Range of the Tibetan Plateau, *Hydrol. Earth Syst. Sci.*, 27(4), 933–952, <https://doi.org/10.5194/hess-27-933-2023>, 2023.
- 1080 Wu, K., Liu, S., Guo, W., Wei, J., Xu, J., Bao, W., and Yao, X.: Glacier change in the western Nyainqêntanglha Range, Tibetan Plateau using historical maps and Landsat imagery: 1970–2014, *J. Mt. Sci.*, 13(8), 1358–1374, <https://doi.org/10.1007/s11629-016-3997-0>, 2016.
- Wu, K., Liu, S., Zhu, Y., Xie, F., Gao, Y., Qi, M., Miao, W., Duan, S., Han, F., and Grünwald, R.: Monitoring the surface elevation changes of a monsoon temperate glacier with repeated UAV surveys, Mainri Mountains, China, *Remote Sens.*, 14(9), 2229, <https://doi.org/10.3390/rs14092229>, 2022.
- 1085 Xu, C., Li, Z., Wang, P., Anjum, M. N., Li, H., and Wang, F.: Detailed comparison of glaciological and geodetic mass balances for Urumqi Glacier No.1, eastern Tien Shan, China, from 1981 to 2015, *Cold Reg Sci Technol.*, 155, 137–148, <https://doi.org/10.1016/j.coldregions.2018.08.006>, 2018.
- Yan, S., Guo, H., Liu, G., and Ruan, Z.: Mountain glacier displacement estimation using a DEM-assisted offset tracking method with ALOS/PALSAR data, *Remote Sens. Lett.*, 4(5), 494–503, <https://doi.org/10.1080/2150704X.2012.754561>, 2013.
- 1090 Yang, H., Yan, S., Liu, G., and Ruan, Z.: Fluctuations and movements of the Kuksai Glacier, western China, derived from Landsat image sequences, *J. Appl. Remote. Sens.*, 8(1), 084599, <https://doi.org/10.1117/1.JRS.8.084599>, 2013.
- 1095 Yao, T., Li, Z., Yang, W., Guo, X., Zhu, L., Kang, S., Wu, Y., and Yu, W.: Glacial distribution and mass balance in the Yarlung Zangbo River and its influence on lakes, *Chinese Science Bulletin*, 55(20), 2072–2078, <https://doi.org/10.1007/s11434-010-3213-5>, 2010.
- Yao, T., Thompson, L., Yang, W., Yu, W., Gao, Y., Guo, X., Yang, X., Duan, K., Zhao, H., Xu, B., Pu, J., Lu, A., Xiang, Y., Kattel, D. B., and Joswiak, D.: Different glacier status with atmospheric circulations in Tibetan Plateau and surroundings, *Nat. Clim. Chang.*, 2(9), 663–667, <https://doi.org/10.1038/nclimate1580>, 2012.
- 1100 Yao, T., Bolch, T., Chen, D., Gao, J., Immerzeel, W., Piao, S., Su, F., Thompson, L., Wada, Y., Wang, L., Wang, T., Wu, G., Xu, B., Yang, W., Zhang, G., and Zhao, P.: The imbalance of the Asian water tower, *Nat. Rev. Phys.*, <https://doi.org/10.1038/s43017-022-00299-4>, 2022.
- 1105 Zemp, M., Thibert, E., Huss, M., Stumm, D., Rolstad Denby, C., Nuth, C., Nussbaumer, S. U., Moholdt, G., Mercer, A., Mayer, C., Joerg, P. C., Jansson, P., Hynek, B., Fischer, A., Escher-Vetter, H., Elvehøy, H., and Andreassen, L. M.: Reanalysing glacier mass balance measurement series, *Cryosphere*, 7(4), 1227–1245, <https://doi.org/10.5194/tc-7-1227-2013>, 2013.
- Zemp, M., Frey, H., Gärtner-Roer, I.: Historically unprecedented global glacier decline in the early 21st century, *J. Glaciol.*, 61(228), 745–762, <https://doi.org/10.3189/2015JoG15J017>, 2015.

- 1110 Zhang, G., Kang, S., Fujita, K., Huintjes, E., Xu, J., Yamazaki, T., Haginoya, S., Wei, Y., Scherer, D., Schneider, C., and Yao, T.: Energy and mass balance of Zhadang glacier surface, central Tibetan Plateau, *J. Glaciol.*, 59(213), 137–148, <https://doi.org/10.3189/2013JoG12J152>, 2013.
- Zhang, Q., and Zhang, G.: Glacier elevation changes in the western Nyainqêntanglha Range of the Tibetan Plateau as observed by TerraSAR-X/TanDEM-X images, *Remote Sens. Lett.*, 8(12), 1142–1151, <https://doi.org/10.1080/2150704X.2017.1362123>, 2017.
- 1115 Zhang, Z., Liu, S., Wei, J., Xu, J., Guo, W., Bao, W., and Jiang, Z.: Mass change of glaciers in Muztag Ata–Kongur Tagh, eastern Pamir, China from 1971/76 to 2013/14 as derived from remote sensing data, *PLoS One*, 11(1), e0147327, <https://doi.org/10.1371/journal.pone.0147327>, 2016.
- Zhou, J., Li, Z., and Guo, W.: Estimation and analysis of the surface velocity field of mountain glaciers in Muztag Ata using satellite SAR data, *Environ. Earth Sci.*, 71(8), 3581–3592, <https://doi.org/10.1007/s12665-013-2749-5>, 2014.
- 1120 Zhou, Y., Li, Z., Li, J., Zhao, R., and Ding, X.: Glacier mass balance in the Qinghai–Tibet Plateau and its surroundings from the mid-1970s to 2000 based on Hexagon KH-9 and SRTM DEMs, *Remote Sens. Environ.*, 210, 96–112, <https://doi.org/10.1016/j.rse.2018.03.020>, 2018.
- Zhu, M., Yao, T., Yang, W., Maussion, F., Huintjes, E., and Li, S.: Energy- and mass-balance comparison between Zhadang and Parlung No. 4 glaciers on the Tibetan Plateau, *J. Glaciol.*, 61(227), 595–607, <https://doi.org/10.3189/2015JoG14J206>, 2015.
- 1125 Zhu, M., Yao, T., Yang, W., Xu, B., Wu, G., Wang, X., and Xie, Y.: Reconstruction of the mass balance of Muztag Ata No. 15 glacier, eastern Pamir, and its climatic drivers, *J. Glaciol.*, 64(244), 259–274, <https://doi.org/10.1017/jog.2018.16>, 2018a.
- 1130 Zhu, M., Yao, T., Yang, W., Xu, B., Wu, G., and Wang, X.: Differences in mass balance behavior for three glaciers from different climatic regions on the Tibetan Plateau, *Clim. Dyn.*, 50(9–10), 3457–3484, <https://doi.org/10.1007/s00382-017-3817-4>, 2018b.