

1 Significant underestimation of peatland permafrost along the 2 Labrador Sea coastline in northern Canada

3 Yifeng Wang¹, Robert G. Way¹, Jordan Beer¹, Anika Forget¹, Rosamond Tutton^{1,2}, Meredith C. Purcell³

4 ¹Northern Environmental Geoscience Laboratory, Department of Geography and Planning, Kingston, K7L 3N6, Canada

5 ²Global Water Futures, Wilfrid Laurier University, Yellowknife, X1A 2P8, Canada

6 ³Tornat Wildlife, Plants, and Fisheries Secretariat, Happy Valley-Goose Bay, A0P 1E0, Canada

7 *Correspondence to:* Yifeng Wang (yifeng.wang@queensu.ca)

8 **Abstract.** Northern peatlands cover approximately four million km², and about half of these peatlands are estimated to contain
9 permafrost and periglacial landforms, like palsas and peat plateaus. In northeastern Canada, peatland permafrost is predicted
10 to be concentrated in the western interior of Labrador but is assumed to be largely absent along the Labrador Sea coastline.
11 However, the paucity of observations of peatland permafrost in the interior, coupled with traditional and ongoing use of
12 perennially frozen peatlands along the coast by Labrador Inuit and Innu, suggests a need for re-evaluation of the reliability of
13 existing peatland permafrost distribution estimates for the region. In this study, we develop a multi-stage consensus-based
14 point inventory of peatland permafrost complexes in coastal Labrador and adjacent parts of Quebec using high-resolution
15 satellite imagery, and we validate it with extensive field visits and low-altitude aerial photography and videography. A subset
16 of 2092 wetland complexes that potentially contained peatland permafrost were inventoried, of which 1119 were classified as
17 likely containing peatland permafrost. Likely peatland permafrost complexes were mostly found in lowlands within 22 km of
18 the coastline where mean annual air temperatures often exceed +1 °C. A clear gradient in peatland permafrost distribution
19 exists from the outer coasts, where peatland permafrost is more abundant, to inland peatlands, where permafrost is generally
20 absent. This coastal gradient may be attributed to a combination of climatic and geomorphological influences which lead to
21 lower insolation, thinner snowpacks, and poorly drained, frost-susceptible materials along the coast. The results of this study
22 suggest that existing estimates of permafrost distribution for southeastern Labrador require adjustments to better reflect the
23 abundance of peatland permafrost complexes to the south of the regional sporadic discontinuous permafrost limit. This study
24 constitutes the first dedicated peatland permafrost inventory for Labrador and provides an important baseline for future
25 mapping, modelling, and climate change adaptation strategy development in the region.

26 1 Introduction

27 Near the southern boundary of latitudinal permafrost zonation, lowland perennially frozen ground is primarily
28 restricted to wetlands in the form of palsas (peat mounds with a frozen core of mineral and organic material) and peat plateaus
29 (fields of frozen peat elevated above the general surface of the surrounding peatland) (Payette, 2004; International Permafrost
30 Association Terminology Working Group, 2005; Zoltai, 1972; Zoltai and Tarnocai, 1975). Persistence of these cryotic

landforms at the extreme limits of their viability is facilitated by a large temperature offset between the ground surface and the top of permafrost, caused by the thermal properties of thick layers of overlying peat and the buffering effect of ground ice (Burn and Smith, 1988; Williams and Smith, 1989). In recent years, many studies have shown that peatland permafrost can be very sensitive to climate warming and ecosystem modifications (Beilman et al., 2001; Borge et al., 2017; Thibault and Payette, 2009). Understanding the distribution of these ice-rich, thaw-sensitive periglacial environments is important for assessing thermokarst potential (Gibson et al., 2021; Olefeldt et al., 2016), local hydrological and vegetation change (Zuidhoff and Kolstrup, 2005), regional infrastructure or land-use planning, and global carbon stores and carbon cycling activities (Hugelius et al., 2014).

Palsas and peat plateaus are primarily thought to occur in continental locations (Fewster et al., 2020; Hustich, 1939) where colder winters allow deeper frost penetration and drier summers promote less thaw. As such, palsas and peat plateaus have been described in many continental locations in Canada, including Yukon Territory, the Northwest Territories, and the Prairie provinces (e.g., Beilman et al., 2001; Coultish and Lewkowicz, 2003; Mamet et al., 2017; Thie, 1974; Zoltai, 1972). However, these landforms have also been documented in coastal locations including the Hudson Bay Lowlands in northern Manitoba, Ontario, and Quebec (e.g., McLaughlin and Webster, 2014; Ou et al., 2016; Pironkova, 2017). In the Labrador region of northeastern Canada, continental- to hemispheric-scale studies have depicted peatland permafrost as present in the region's continental interior but as far less abundant or completely absent along most of the Labrador Sea coastline (Fewster et al., 2020; Hugelius et al., 2020; Olefeldt et al., 2021). However, historic and contemporary use of coastal peatland permafrost environments by Labrador Inuit and Innu is well documented (Anderson et al., 2018), and published field-based observations (e.g., Anderson et al., 2018; Andrews, 1961; Brown, 1975, 1979; Davis et al., 2020; Dionne, 1984; Elias, 1982; Hustich, 1939; Seguin and Dionne, 1992; Smith, 2003; Way et al., 2018; Wenner, 1947) suggest that peatland permafrost is abundant along some sections of the coast. This recurring misestimation of peatland permafrost has an impact on predictions of ground ice content (O'Neill et al., 2019), thermokarst potential (Olefeldt et al., 2016), and carbon content (Hugelius et al., 2014) in the region.

Locally, preservation of peatland permafrost complexes is relevant to Labrador Inuit and Innu because these areas are frequented for traditional activities such as bakeapple (cloudberry; Inuttitut: appik; Innu-aimun: shikuteu; *Rubus chamaemorus*) berry-picking (Anderson et al., 2018; Karst and Turner, 2011; Norton et al., 2021), goose hunting, and fox trapping (Way et al., 2018). Improvements to our understanding of regional peatland permafrost distribution will provide an important baseline for local and regional climate change adaptation strategy development, while better representation of the distribution of thaw-sensitive terrain will inform future development of linear and built infrastructure in coastal Labrador (Way et al., 2021b; Bell et al., 2011).

Previous peatland permafrost mapping in Labrador has been limited to scattered observations of palsa bogs from the National Topographic Database (Natural Resources Canada, 2005) and the Ecological Land Classification (Environment Canada, 1999), with no comprehensive peatland permafrost inventorying efforts completed to date (Way et al., 2018). In this study, we develop a multi-stage, consensus-based point inventory of contemporary peatland permafrost complexes within 100

65 km of the Labrador Sea coastline (Figure 1), comprising the region of Nunatsiavut and surrounding areas, including the land
66 claims agreement-in-principle of the Labrador Innu Nation (Nitassinan) and coastal areas claimed by the NunatuKavut
67 Community Council (NunatuKavut). The goal of this inventory is to map and contextualize the contemporary distribution of
68 peatland permafrost complexes throughout coastal Labrador, using extensive validation efforts from a combination of field
69 visits and low-altitude image and video acquisitions. We hypothesize that this point-based inventory will reinforce the local
70 understanding of a high abundance of peatland permafrost landforms in coastal locations, which will be relevant for carbon
71 modelling, land use planning, infrastructure development, and climate change adaptation strategy development at local to
72 regional scales in northeastern Canada. This contribution will also provide insights into the reliability of relevant peatland
73 permafrost and permafrost distribution products, which currently claim an absence or low abundance of both peatland
74 permafrost and permafrost along the Labrador Sea coastline. Based on our results, we also propose amendments to the current
75 limits of the sporadic discontinuous and isolated patches of permafrost distribution zones in southeastern Labrador. This point-
76 based inventory is a first step towards understanding the distribution of peatland permafrost in Labrador and will contribute to
77 refined regional and global estimates of ground ice content, thermokarst potential, and carbon storage in northern Canada.




Figure 1. (a) Permafrost zonation in Labrador (Heginbottom et al., 1995) with the boundary for the inventory study area (black line) corresponding to areas within 100 km of the Labrador Sea coastline. Map is annotated with locations of the Torngat Mountains National Park, Akami-Uapishkuk-KakKasuak-Mealy Mountains National Park Reserve, and selected communities; (b) Inset map showing Labrador's position in Canada.

83 2 Study area

84 2.1 Bioclimatic setting

85 Labrador's climate is strongly influenced by atmosphere-ocean interactions from the adjacent Labrador Sea (Barrette
86 et al., 2020; Way and Viau, 2015). In coastal Labrador, long, cold winters and short, cool summers are largely dictated by the
87 Labrador Current that carries cold Arctic waters down the eastern coast of mainland Canada (Banfield and Jacobs, 1998;
88 Foster, 1983; Roberts et al., 2006; Way et al., 2017). Mean annual air temperatures (1980-2010) decrease with continentality
89 and latitude, ranging from -12°C in parts of the Torngat Mountains National Park to +1.5°C near the community of Blanc-
90 Sablon (Karger et al., 2017, 2021). Labrador is also characterized by some of the highest precipitation amounts in the North
91 American boreal zone (Banfield and Jacobs, 1998; Hare, 1950) due to its varying relief, high moisture availability from the
92 adjacent Atlantic Ocean, and high frequency of passing winter storm systems (Brown and Lemay, 2012). Precipitation totals
93 as high as ~2700 mm per year are estimated for some locations at high elevations along the coast (Karger et al., 2017, 2021),
94 with solid precipitation fractions increasing with both latitude and elevation (~0.35 at Blanc-Sablon; ~0.5 at Nain)
95 (Environment and Climate Change Canada, 2022).

96 Ecologically, Labrador is characterized by taiga forests in the interior, tundra in the north, and wind-swept coastal
97 barrens along the coastline of the Labrador Sea (Roberts et al., 2006). Tree cover is sparse in the coastal barrens because of
98 climatic and physiographic limitations, but dense patches of black spruce (*Picea mariana*), white spruce (*Picea glauca*),
99 tamarack (*Larix laricina*), and balsam fir (*Abies balsamea*), interspersed with deciduous trees, like paper birch (*Betula*
100 *papyrifera*) and trembling aspen (*Populus tremuloides*), exist in sheltered locations and on some slopes (Roberts et al., 2006).
101 Wetlands are found throughout Labrador, but total wetland abundance is difficult to assess given widespread disagreement
102 between existing estimates of wetland and peatland extents for this region (Supplement Sect. S1). Generally, wetlands in
103 Labrador tend to decrease in abundance but increase in size as latitude increases. Most wetlands along the southern Labrador
104 coast are classified as raised bogs, while inland, most wetlands are string and blanket bogs (Foster and Glaser, 1986).

105 2.2 Physical environment

106 Labrador is mostly underlain by igneous and metamorphic bedrock (Roberts et al., 2006). Extensive blankets of
107 glacial till were deposited during the retreat of the Laurentide Ice Sheet (12-6 k years BP) (Bell et al., 2011; Dyke, 2004),
108 along with thin layers of medium- to fine-grained marine and glaciomarine sediments in coastal lowland areas below the
109 marine limit (Fulton, 1995). The post-glacial marine limit decreases with latitude, from ~150 m a.s.l. in southeastern Labrador
110 and along the Quebec Lower North Shore to 0 m a.s.l. at the northernmost tip of Labrador in the Torngat Mountains (Dyke et
111 al., 2005; Occhietti et al., 2011; Vacchi et al., 2018). The broad distribution of near-surface bedrock and hardpans (Smith,
112 2003) results in poor drainage that has facilitated peatland development across large areas of southern Labrador, particularly
113 in depressions and over flat deposits.

114 **2.3 Permafrost distribution**

115 While permafrost conditions in Labrador, including the presence of peatland permafrost landforms, have been noted
116 during ecological, palynological, glaciological, and archeological surveys and studies (Anderson et al., 2018; Andrews, 1961;
117 Hustich, 1939; Smith, 2003; Wenner, 1947), permafrost-specific field investigations are limited to R.J.E. Brown’s (1975)
118 helicopter survey in the late 1960s and the Labrador Permafrost Project that began in 2013 (Way, 2017). Our understanding
119 of permafrost distribution in Labrador has relied on extensive extrapolation of limited field observations and broad assumptions
120 of the interactions between air temperature, vegetation cover, snow cover, and permafrost presence (Ives, 1979). According to
121 the Permafrost Map of Canada (Heginbottom et al., 1995), the area underlain by permafrost in Labrador is less extensive than
122 comparable regions in northern Canada like Yukon Territory or the Northwest Territories. Approximately two-thirds of
123 Labrador is classified in the isolated patches of permafrost zone (<10 % permafrost by area), but the distribution of permafrost
124 does become more widespread farther north (Figure 1). Along the Labrador coastline, the sporadic discontinuous permafrost
125 zone (10-50 % permafrost by area) extends slightly further south along the outer edge of the
126 Akami–Uapishk–KakKasuak–Mealy Mountains National Park Reserve than in the interior, though the justification for this
127 departure is not clarified in published literature. Continuous permafrost (>90 % permafrost by area) is expected to persist only
128 at high elevations and latitudes, mostly in the Torngat Mountains (Heginbottom et al., 1995).

129 **2.4 Inventory extent**

130 This study is focused on the coastal areas of Labrador and Quebec, within 100 km of the Labrador Sea coastline
131 (Figure 1). This area of interest was informed by knowledge gained from prior works in the region (Anderson et al., 2018;
132 Andrews, 1961; Brown, 1975, 1979; Davis et al., 2020; Dionne, 1984; Elias, 1982; Hustich, 1939; Seguin and Dionne, 1992;
133 Smith, 2003; Way, 2017; Way et al., 2018; Wenner, 1947) that indicated a greater abundance of peatland permafrost landforms
134 along the coast as compared to the interior of Labrador. Exhaustive descriptions of records of peatland permafrost and other
135 periglacial landforms in Labrador have been presented by Brown (1979) and Way (2017), both of whom found limited evidence
136 of peatland permafrost in Labrador’s interior.

137 **3 Methods**

138 Palsas and peat plateaus are typically found in bogs and may measure up to 4 m higher than their surrounding
139 wetlands, so large peatland permafrost landforms can be identified and mapped from high-resolution satellite imagery (Borge
140 et al., 2017; Gibson et al., 2020, 2021). Our point inventory, which includes only the largest and most visually apparent peatland
141 permafrost complexes within 100 km of the Labrador Sea coastline, was generated through a multi-stage mapping and
142 consensus-based review process, supported by extensive validation efforts mostly completed between 2017 and 2022. Mapping
143 and identification activities were informed by existing wetland and peatland distribution products (Supplement Sect. S1), but

significant disagreement between these products limited their direct application and utility during the inventorying process. An initial inventory of wetlands of interest (WOIs) was developed as a subset of the wetlands in coastal Labrador deemed potentially suitable (e.g., bogs and fens) for the development and persistence of peatland permafrost landforms. The presence of peatland permafrost landforms within the WOIs was then evaluated through a consensus-based review of high-resolution satellite imagery by three mappers with permafrost-specific field experience in the region. Final interpretation of peatland permafrost presence or absence within the WOIs was based on reviewer agreement and was informed by field- and imagery-based validation of peatland permafrost landform presence or absence.

3.1 Data sources

WOIs were identified and evaluated using Maxar (Vivid) optical satellite imagery, available as the World Imagery basemap via ArcGIS Online (0.5 m ground sampling distance; 5 m absolute spatial accuracy) (Esri, 2022). These satellite imagery mosaics consisted of summer imagery with minimal cloud and snow cover, with acquisition dates for Labrador ranging from 2010 to 2020.


Topographic data from Natural Resources Canada covering the WOIs were extracted from the Canadian Digital Elevation Data (CDED; 50 m spatial resolution), with a small gap near the provincial border between Labrador and the Quebec Lower North Shore that was filled in using the Canadian Digital Surface Model (CDSM). Gridded mean annual air temperature (MAAT) and mean annual thawing degree days (TDD) for the 1981 to 2010 climate normal were extracted from CHELSA V2.1 (~1 km spatial resolution) (Karger et al., 2017, 2021) at the WOI locations. Mean annual freezing degree days (FDD) for the WOI locations for 1981 to 2010 were calculated from MAAT and TDD over the same climate normal, following prior work in the region (Way et al., 2017; Way and Lewkowicz, 2018).

3.2 Inventorying peatland permafrost complexes

3.2.1 Identifying wetlands of interest (WOIs)

A team of three mappers used ArcGIS Online to identify and place point features within WOIs throughout coastal Labrador (Figure 2). The point-based nature of the inventorying process allowed for evaluation of the entire study area by incorporating field- and imagery-based validation for many WOIs over a large study area, as opposed to detailed validation of peatland permafrost areal coverage within a given WOI. These WOIs were restricted to include only those that contained prospective peatland permafrost landforms exceeding 2 m in length or width (~4 m²), which was determined to be the smallest detectable feature based on the 0.5 m spatial resolution of the satellite imagery. Mappers were instructed to identify WOIs based on local geomorphology, local hydrology and drainage patterns, the presence of a white or grey lichen surface cover corresponding to *Cladonia* and/or *Ochrolechia* spp. lichens, shadows indicative of elevated landform edges and surface uplift, and thermokarst ponding or exposed peat indicative of thaw processes. The inventory sought to only include contemporary peatland permafrost landforms, so WOIs with extensive thermokarst ponding but no evident peatland permafrost landforms

175 were not included in the database. Individual WOIs ranged in size from ~0.2 km² to larger than ~3.5 km². However, the total
176 area underlain by peatland permafrost within each WOI was not able to be reliably evaluated using satellite imagery. WOIs
177 near one another were sometimes difficult to discern due to potential connectivity between adjacent systems, but contiguous
178 WOIs could generally be identified by differences in drainage, vegetation, and morphology, or because of separation by linear
179 infrastructure like roads, airstrips, and trails (Figure 2). Mappers also assigned each WOI a self-assessed score to reflect their
180 confidence in their interpretation of permafrost presence within the wetland complex (1 = low confidence, 2 = medium
181 confidence, 3 = high confidence).



182
183 **Figure 2. Examples of wetland complexes of interest (WOIs) in Labrador that were identified by the mapping team using high-**
184 **resolution satellite imagery available via Esri ArcGIS Online. Examples of WOI boundaries are shown in red and were determined**
185 **based on differences in drainage or vegetation from adjacent WOIs or based on separation following linear infrastructure, such as**
186 **roads, airstrips, or trails. Identification was restricted to WOIs that contained prospective peatland permafrost landforms.**

187 3.2.2 Quality control of WOI database

188 The WOI inventory was subjected to a quality control check, during which complexes were reviewed and duplicates
189 or points clearly not corresponding to wetlands were removed. In some cases, non-wetland locations may have been retained
190 because of difficulties discerning peat plateaus from surface peat over bedrock or coastal tundra.

191 3.2.3 Consensus-based review of WOI database

192 The quality-controlled WOI inventory was sent back to the mappers for a consensus-based review, similar to Way et
193 al.'s (2021a) approach for rock glacier inventorying in northern Labrador. Each WOI was independently reviewed by two team
194 members, both of whom had access to the mapper's initial confidence rating, and one of whom had access to a field-validated
195 dataset of WOIs (see Sect. 3.3 Validation of subset of WOI database). Both team members were asked to indicate whether
196 each WOI contained peatland permafrost landforms. WOIs evaluated by both reviewers as containing peatland permafrost
197 were considered likely to contain palsas or peat plateaus, while WOIs evaluated by both reviewers as not containing peatland
198 permafrost were considered unlikely to contain palsas or peat plateaus. WOIs with conflicting classifications were considered
199 to possibly contain palsas or peat plateaus. This consensus-based review process resulted in a full inventory of WOIs that were
200 classified as likely, possibly, or unlikely to contain peatland permafrost.

201 3.3 Validation of subset of WOI database

202 The full, consensus-based inventory results were compared with a field- and imagery-validated dataset of 557 WOIs,
203 with and without contemporary peatland permafrost landforms. From July to September 2021 and 2022, field evaluations of
204 WOIs were undertaken via in-person field visits, remotely piloted aircraft (RPA) image acquisitions (DJI Mini 2 microdrone,
205 weighing less than 250 g), video clip acquisition from a helicopter survey, and image acquisitions from commercial Twin Otter
206 aircraft flights. Interpretation of the presence or absence of permafrost landforms within each WOI that was visited or aerially
207 surveyed was also determined through consensus between two mappers. Any WOIs with disagreements in interpretation were
208 re-evaluated and discussed until consensus could be reached between the two mappers.

209 Field visits to WOIs were undertaken at road-accessible locations within 500 m of the Trans-Labrador Highway and
210 other accessible side roads via truck or ATV and at coastal locations via speedboat from the nearby communities of Black
211 Tickle, Cartwright, Rigolet, and Nain. The number of WOIs that could be visited for field validation was restricted by weather
212 conditions, tides, the availability of local guides and boat drivers with location-specific expertise, and other logistical and
213 operational constraints. During field visits, team members probed the soil to the depth of refusal (maximum of 125 cm). The
214 nature of refusal, interpreted as frozen ground, compact sediment, clasts, rock, or not applicable (N/A; >125 cm), was noted
215 and used to assess permafrost presence or absence. Where the cause of probe refusal was unclear, instantaneous ground
216 temperature measurements were collected using vertically arranged thermistors connected to an Onset Hobo UX120-006M 4-
217 Channel Analog Data Logger (accuracy ± 0.15 °C) (Davis et al., 2020; Holloway and Lewkowicz, 2020; Way et al., 2021b;

218 Way and Lewkowicz, 2015). Ground temperatures were recorded within the probed hole for a minimum of 10 minutes to allow
219 for thermal equilibration. Frost probing and instantaneous ground temperature measurements were targeted towards locations
220 considered most likely to contain frozen ground and thus mostly occurred on elevated peat-covered microtopography within
221 each WOI.

222 Low-altitude RPA imagery of prospective peatland permafrost complexes were collected using a DJI Mini 2
223 microdrone when weather conditions were suitable (i.e., no rain, no fog, low wind). Low-altitude georeferenced video footage
224 was collected using a GoPro Hero9 camera mounted onto a helicopter during a fuel cache mission in northern Labrador in July
225 and August 2021, led by the Torngat Wildlife, Plants, and Fisheries Secretariat. The camera was set to record real-time video
226 (1080 p, 60 fps, wide) at an oblique angle (~45°). The flight altitude was between 90 m and 120 m a.g.l., similar to coastal
227 Nunavik transects performed by Boisson and Allard (2018), and the flight plan between the Goose Bay Airport and the Torngat
228 Mountains National Park was designed to fly over WOIs in coastal locations north of the community of Makkovik (55.0° N)
229 (Supplement Sect. S2). Low-altitude georeferenced aerial images were also collected using handheld digital cameras (Nikon
230 Coolpix W300 or Olympus Tough TG-6) during commercial Air Borealis Twin Otter flight segments between Cartwright and
231 Black Tickle and between Goose Bay, Rigolet, Makkovik, Postville, Hopedale, Natuashish, and Nain. The Twin Otter flights
232 only crossed over WOIs along existing commercial flight routes.

233 **3.4 Compilation of final WOI database**

234 The final WOI database of likely, possible, or unlikely peatland permafrost complexes was developed following the
235 incorporation of the field-validated dataset. WOIs that were classified as likely or possibly to contain peatland permafrost were
236 subject to a final round of review in which the peatland permafrost landforms were identified as palsas, peat plateaus, or both
237 palsas and peat plateaus (mixed).

238 **3.5 Statistical analyses of final WOI database**

239 ANOVA (analysis of variance) and *post hoc* Tukey's HSD (honest significant difference) tests were performed to
240 determine whether the mean latitude, distance from coastline, elevation, MAAT, TDD, and FDD were statistically significantly
241 different between the final classes of likely, possibly, and unlikely peatland permafrost complexes. Statistical analyses were
242 performed in R 4.0.3 (R Core Team, 2020).

243 **4 Results**

244 **4.1 Peatland permafrost complex identification and review**

245 A total of 2092 unique WOIs, limited to the largest and most visually apparent prospective peatland permafrost
246 complexes within the study area, were included in the full inventory. Reviewer agreement was very high (89 %) during the
247 consensus-building review process, with 1116 complexes classified by both reviewers as likely containing peatland permafrost

248 and 750 complexes classified by both reviewers as unlikely to contain peatland permafrost, and only 226 complexes with
249 conflicting classifications of permafrost presence or absence (11 %) (Supplement Sect. S2).

250

251 **4.2 Validation of peatland permafrost complexes**

252 In Summer 2021 and 2022, in-person field visits (n=63 WOIs), RPA visits (n=141 WOIs), helicopter video clips
253 (n=69 WOIs), and Twin Otter images (n=314 WOIs) were combined to evaluate peatland permafrost presence at 531 WOIs,
254 49 of which were cross-validated using multiple methods (Figure 3; Supplement Sect. S2). Previous work from 2017 to 2020,
255 including field visits (n=23 WOIs) and RPA image collection (n=19 WOIs), were also used to validate palsa or peat plateau
256 presence at an additional 19 complexes and peatland permafrost absence at an additional seven complexes (Anderson et al.,
257 2018; Way, 2017). Out of the 557 WOIs evaluated via field and/or imagery validation methods, 311 were interpreted to contain
258 peatland permafrost landforms. Comparison between the validation dataset and the consensus-based inventory resulted in re-
259 classification of 39 of the 226 possible peatland permafrost complexes (17 %) to either likely (n=3) or unlikely (n=36) peatland
260 permafrost complexes.

261




Figure 3. (a) Locations of validated peatland permafrost complexes in coastal Labrador from field-based activities and imagery acquisition using RPA, helicopter, and Twin Otter from 2017 to 2022; Example of peatland complexes containing palsas and/or peat plateaus near (b) Rigolet, (c) Cartwright, (d) Black Tickle-Domino, and (e) Red Bay.

4.3 Peatland permafrost complex inventory

A total of 1119 out of 2092 WOIs were classified as likely containing peatland permafrost landforms, with an additional 187 wetland complexes classified as possibly containing peatland permafrost landforms (Figure 4). The largest clusters of likely and possible peatland permafrost complexes were located between Makkovik (55.0° N) and Black Tickle (53.5° N) (Figure 4; Figure 5A). The likely peatland permafrost complexes were at low elevation (mean elevation of 29 m a.s.l.) (Figure 5C) within 22 km of the coastline (mean distance from coastline of 2.6 km) (Figure 4; Figure 5B). Likely peatland

272 permafrost complexes were distributed from 51.4° N near Blanc-Sablon to 58.6° N in the Torngat Mountains National Park
273 (Figure 4; Figure 5A), with most complexes located in southeastern Labrador (mean latitude of 54.1° N) (Supplement Sect.
274 S3). Comparison against gridded climate products showed that the MAAT at peatland permafrost complexes ranged from -7.5
275 °C to +1.2 °C, with corresponding ranges for FDDs of 1126 degree days to 3471 degree days and TDDs of 733 degree days to
276 1704 degree days (Figure 5D-F). Despite the wide range in MAAT, the majority of the likely peatland permafrost complexes
277 (90 %) were found in locations with MAATs between -2 °C and +1 °C (Figure 5D).




Figure 4. (a) Spatial distribution of inventoried peatland complexes (n=2092) classified as likely containing peatland permafrost landforms (n=1119), possibly containing peatland permafrost landforms (n=187), and unlikely to contain peatland permafrost landforms (n=786); (b) Inset map showing density of peatland permafrost complexes within 20 by 20 km (400 km²) grid cells.




Figure 5. Distribution of wetland complexes likely or possibly containing peatland permafrost landforms by (a) latitude; (b) distance from the coastline; (c) elevation; and (d) mean annual air temperature; (e) mean annual freezing degree days; and (f) mean annual thawing degree days for the 1981 to 2010 climate normal.

286

287 ANOVA and *post hoc* Tukey's HSD tests revealed that the mean distance from coastline, elevation, MAAT, FDD,
288 and TDD were statistically different between the likely, possibly, and unlikely peatland permafrost complexes at the 95 %
289 confidence level. When compared with the complexes that likely contained peatland permafrost, the 187 complexes that
290 possibly contained peatland permafrost were similarly distributed all along the coastline but were skewed further north (mean
291 latitude of 54.5° N) and extended as far as 60.2° N (Supplement Sect. S3). These less certain features were at greater distances
292 from the coastline (mean distance from coast of 7.8 km) and at higher elevations (mean elevation of 66 m a.s.l.). The 786
293 complexes that were unlikely to contain peatland permafrost were well distributed between 51.4° N and 60.2° N (Figure 4) but
294 were located further from the coastline (mean distance from coastline of 10.7 km), at higher elevations (mean elevation of 78
295 m a.s.l.), and at higher MAATs (mean MAAT of -0.5 °C) than the complexes that likely or possibly contained peatland
296 permafrost (Supplement Sect. S3).

297 Likely and possible peatland permafrost complexes were also classified according to the type of peatland permafrost
298 landforms found within the wetland complex. Complexes that were exclusively comprised of palsas accounted for half of the
299 likely and possible peatland permafrost complexes (50 %) and were distributed along the entire study area. Complexes with
300 exclusively peat plateaus were less common (29 %) and were spatially concentrated between ~53° N and ~55° N. The
301 remaining 21 % of the likely and possible peatland permafrost complexes were interpreted to contain a combination of palsas
302 and peat plateaus, but it is possible that many of these complexes contain dissected and heavily degraded peat plateaus that
303 now resemble palsas. Further field-based investigations would be required to differentiate these degradational landforms.




Figure 6. (a) Spatial distribution of likely and possible peatland permafrost complexes classified by peatland permafrost landform type as palsas, peat plateaus, or a mix of both palsas and peat plateaus for coastal Labrador. (b) Inset map showing existing permafrost distribution zones (Heginbottom et al., 1995) for a subsection of coastal Labrador and the location of a new proposed location for the southern limit of the sporadic discontinuous permafrost zone.

310 **5 Discussion**

311 **5.1 Distribution of peatland permafrost in Labrador**

312 Peatland permafrost complexes in Labrador and adjacent portions of Quebec are abundant in lowlands within 22 km
313 of the Labrador Sea coastline (Figure 5B). A geographic gradient is especially apparent between Rigolet (54.2° N) and Black
314 Tickle (53.5° N), where peat plateaus are abundant along the coast but absent from wetlands farther inland (Figure 4). The
315 higher density of peatland permafrost complexes along the coast could be linked to climatic factors like persistent fog and
316 cloud cover leading to less incoming solar radiation (Way et al., 2018) or thinner and denser snowpacks (Seppälä, 1994; Vallée
317 and Payette, 2007) in the wind-exposed barrens along the coast (Way et al., 2018). Further work should focus on exploring the
318 role of local climate conditions in the formation and persistence of peatland permafrost in coastal Labrador and similar northern
319 coastal locations. Peatland permafrost was found across a large range of MAATs, spanning from -7.5 °C to +1.2 °C. Permafrost
320 persistence at MAATs above +1 °C in southeastern Labrador was previously noted in a field study at five palsa complexes
321 (Way et al., 2018). Peatland permafrost complexes in Labrador were located at higher MAATs than is predicted for other
322 northern coastal regions like northern Finland, Norway, and Sweden (approximately +0.4 °C) (Parviainen and Luoto, 2007).
323 Our results also suggest that the MAAT threshold of +0.2 °C for peatland permafrost areas previously applied to North America
324 (Fewster et al., 2020) is too low for Labrador and adjacent parts of Quebec where peatland permafrost landforms continue to
325 persist due to their relict and resilient nature (Dionne, 1984; Way et al., 2018). Large thermal offsets (up to and often exceeding
326 2.0 °C in southeastern Labrador) (Way and Lewkowicz, 2018) are typical of organic-rich landscapes like peatlands and may
327 promote continued landform persistence despite a warming climate (Jorgenson et al., 2010). This may further exacerbate
328 discrepancies between peatland permafrost observations and regional estimates, calling into question the utility of simplified
329 threshold-based approaches when modelling with future climate scenarios. Information on the timing of peatland initiation
330 following deglaciation (Gorham et al., 2007), rates of peat deposition (Tarnocai, 2009; Gorham, 1991), and corresponding peat
331 thicknesses should also be considered in studies of peatland permafrost distribution, as thicker peat deposits may influence
332 permafrost development and protect permafrost persistence through a larger thermal offset (Smith and Riseborough, 2002).

333 The regional distribution of fine-grained sediments and local depositional history are expected to play an important
334 role in landscape suitability for peatland permafrost landforms (O'Neill et al., 2019; Seppälä, 1986; Zoltai, 1972). For example,
335 differences in the distribution of palsa versus peat plateau landforms have previously been attributed to varying thicknesses
336 and extents of the underlying sediment, with thicker sediment deposits leading to the development of palsas and thinner
337 sediment deposits linked to the development of peat plateaus (Allard and Rousseau, 1999). Differences in sediment grain size
338 may also influence the thickness of the ice lenses and the depth at which they form, with thicker ice lenses developing deeper
339 in finer sediments and thinner ice lenses forming at shallower depths in coarser sediments (Allard and Rousseau, 1999). Further
340 examination of how these variables could influence peatland permafrost formation and persistence in coastal Labrador is
341 challenged by the paucity of information on surficial materials and marine limits along most of the Labrador Sea coastline
342 (Hagedorn, 2022; Occhietti et al., 2011). To date, local marine limits have been identified at some individual locations and

study sites in coastal Labrador (e.g., Bell et al., 2011; Dyke et al., 2005; Occhietti et al., 2011; Vacchi et al., 2018), but widespread mapping of marine sediments has only been completed for a small section of northern coastal Labrador from Goose Bay to Hopedale (Hagedorn, 2022). Based on the information that is currently available, we can qualitatively link the distribution of the largest clusters of peatland permafrost complexes, particularly peat plateau complexes, to locations where post-glacial marine invasions had occurred, such as along the lowland-dominated coastline between Makkovik (55.0° N) and Black Tickle (53.5° N), where frost-susceptible, glaciomarine surficial materials are generally widespread (Fulton, 1989, 1995; Hagedorn, 2022; Occhietti et al., 2011). Meanwhile, fewer peatland permafrost complexes were mapped between Makkovik (55.0° N) and Hopedale (55.5° N), where the elevated topography resulted in limited marine invasions and post-glacial marine deposition along the coast. Significant and coordinated advances in surficial mapping will be required before similar links between peatland permafrost distribution and surficial material type, sediment grain size, and elevation relative to the marine limit can be made for other parts of the coastline.

5.2 Implications for peatland permafrost and permafrost distribution in northeastern Canada

Comparisons between our inventory results and several recent national to global wetland, peatland (Supplement Sect. S1; Supplement Sect. S4), and peatland permafrost distribution products (e.g., Fewster et al., 2020; Hugelius et al., 2020; Olefeldt et al., 2021) (Figure 7) provide compelling evidence that peatland permafrost along the Labrador coast is poorly represented by existing datasets. While differences in scale may explain some of this discrepancy, the general pattern presented in most previous datasets, showing relatively greater peatland permafrost in the continental interior and less along the coast, is directly contradicted by the results of this study. This reversed pattern could reflect inaccurate assumptions on the climate limits of peatland permafrost and/or may reflect the absence of field data from many northern coastal peatland permafrost environments (Borge et al., 2017). Inclusion of physiographic variables, like soil conditions, frost-susceptibility of sediments, and more detailed surficial deposit maps are likely necessary for an improved representation of peatland permafrost in northern coastal regions. Recent work by O'Neill et al. (2019), for example, has demonstrated that segregated ice can be reliably modelled along sections of the Labrador Sea coastline (Figure 7D) by incorporating paleogeographic variables like vegetation cover, surficial geology, and glacial lake and marine limits.




Figure 7. Comparison of inventoried peatland permafrost complexes with peatland permafrost presence and percent coverage as modelled by (a) Fewster et al. (2020); (b) Hugelius et al. (2020); and (c) Olefeldt et al. (2021) and with segregated ice content as modelled by (d) O'Neill et al. (2019).

372

373 The results of our inventory also suggest that some amendments to existing representations of permafrost distribution
374 may be required for coastal Labrador. For example, the highest density of peatland permafrost complexes (Figure 4B) was
375 found near the community of Black Tickle (53.5° N) on the Island of Ponds (2 palsa complexes, 19 mixed palsa and peat
376 plateau complexes, and 59 peat plateau complexes within 94 km²) (Figure 6), which is currently classified in the isolated
377 patches of permafrost zone on the Permafrost Map of Canada (Heginbottom et al., 1995) and the no permafrost zone on the
378 2000-2016 Northern Hemisphere Permafrost Map (Obu et al., 2019) (Supplement Sect. S5). The identification of large swaths
379 of likely peatland permafrost complexes, including more than 150 peat plateaus, between Cartwright (53.7° N) and Black
380 Tickle (53.5° N) suggest that the physiography-based Permafrost Map of Canada's limit for the sporadic discontinuous zone
381 along the Labrador coast (Heginbottom et al., 1997) (Supplement Sect. S5), could reasonably be extended south by ~110 km
382 from its current position (~53.7° N) (Figure 6B). This southerly extension of the sporadic discontinuous permafrost limit has
383 previously been suggested by Allard and Seguin (1987) and Payette (2001) who indicated that regional vegetation and
384 geomorphology favoured permafrost along much of this coastline (Payette, 1983). Unexpectedly, large clusters of likely
385 peatland permafrost complexes were also identified near the communities of Red Bay (Supplement Sect. S6) and Blanc-Sablon,
386 both of which are considered to be underlain by little to no permafrost (Heginbottom et al., 1995; Obu et al., 2019) (Supplement
387 Sect. S5). A 15 km extension of the southern limit of the Permafrost Map of Canada's isolated patches permafrost zone to
388 include the Blanc-Sablon region would better reflect contemporary permafrost conditions in this area, especially given that
389 permafrost has been previously detected in mineral soils in the community and in surrounding peatlands below the marine
390 limit (Dionne, 1984).

391 **5.3 Challenges and limitations of a point-based inventory of peatland permafrost complexes in coastal Labrador**

392 The most challenging aspects of the inventorying process involved interpreting peatland permafrost presence in
393 isolated WOIs containing small landforms, while in the case of more obvious peatland permafrost features, there were at times
394 difficulties in determining distinct wetland boundaries (Figure 2). However, we believe that these issues were mitigated
395 through the inclusion of multiple mappers, which facilitated the development of a large initial database and reduced the
396 potential omission of prospective WOIs. The consensus-based review process that followed was designed to minimize the
397 inclusion of false positives in the final dataset of 1119 likely peatland permafrost complexes, but we recognize that this
398 conservative approach may have resulted in the exclusion of some complexes. At the northern end of the study area, where
399 other types of periglacial landforms become more common, misclassification of palsas for other elevated periglacial landforms
400 may have contributed to the designation of a higher number of possible peatland permafrost complexes. It is certainly possible
401 that some segregated ice mounds with less than 40 cm of overlying peat (i.e., lithalsas) may have been included in the inventory,
402 particularly near the northern end of the study area where wetlands are less abundant and peat deposits may be thinner
403 (Supplement Sect. S1). This suggests that the definition of peatlands, as wetlands containing at least 40 cm of surface peat
404 (Tarnocai et al., 2011), and its application to palsas and lithalsas, can introduce some ambiguity during inventorying.

405 While other inventorying approaches, including grid-based methods (Ramsdale et al., 2017; Gibson et al., 2020, 2021;
406 Borge et al., 2017), were considered, a point-based inventory was ultimately developed for this study. The implementation of
407 a grid-based approach with delineation of individual landforms for each WOI could have been useful for estimating ground
408 ice content, thermokarst potential, carbon content, and overall permafrost coverage, but the purpose of this study was to
409 generate an initial inventory to guide future research that will facilitate quantitative assessments of peatland permafrost
410 distribution and coverage in these regions. Our field experience in the region suggests that areal delineations of peatland
411 permafrost complexes in coastal Labrador will require extensive validation, and it is unlikely that even experienced permafrost
412 mappers could accurately map the extents of permafrost throughout some complexes without extensive field investigations.
413 Despite the above limitations, our inventory allowed for the incorporation of dedicated, co-located field- and imagery-based
414 validation information. Post-validation adjustments to the inventory, including reclassification of 39 WOIs highlights the
415 importance of ground-truthing in remote sensing- or modelling-based periglacial landform inventories.

416 Owing to a lack of prior field-based assessments of permafrost conditions in Labrador, it was also difficult to
417 independently validate our peatland permafrost inventory results. However, a detailed aerial photograph- and field-based
418 Ecological Land Classification (ELC) survey undertaken in the late 1970s did cover a subset of our study area in southeastern
419 Labrador (Environment Canada, 1999). The ELC identified a total wetland area of 666 km² which was at least partly covered
420 by inventoried peatland permafrost landforms (Figure 8). Comparison with the present study showed that mappers identified
421 peatland permafrost complexes in 23 of the 24 contiguous ELC wetland areas indicated as containing palsas. Examination of
422 the one remaining ELC peatland permafrost-containing wetland area revealed the presence of irregular ponding patterns
423 indicative of thermokarst and elevated landforms that could be peatland permafrost but, due to their small size, would require
424 in situ field visits for validation. Some of the inventoried likely peatland permafrost complexes that were not captured as part
425 of the peatland permafrost areas from the ELC were instead classified in other wetlands, like string bogs, and in raised marine
426 terrain units. Overall, the results of our inventory are in good agreement with the limited previous overlapping field
427 investigations and inventorying efforts from the ELC.

428




Figure 8. (a) Comparison of inventoried peatland permafrost complexes with palsa bog regions identified in the Ecological Land Classification (ELC) survey (Environment Canada, 1999); (b) Inset map showing the extent of the peatland permafrost area that was mapped in the ELC.

6 Conclusions

This study provides the first detailed point inventory of peatland permafrost landforms along the Labrador Sea coastline. Using high-resolution satellite imagery and extensive field- and imagery-based validation efforts, we applied a multi-stage, consensus-based inventorying approach to identify 1119 likely peatland permafrost complexes. Peatland permafrost complexes were primarily found in lowlands on outer coasts, spanning from 51.4° N to 58.6° N, with the largest clusters of complexes occurring ~110 km south of the previously mapped limit of sporadic discontinuous permafrost in northeastern Canada (Heginbottom et al., 1995).

Comparisons between our point inventory results and existing wetland, peatland, and peatland permafrost distribution products reveal major discrepancies between this study and prior estimates of peatland permafrost in Labrador with implications for ground ice content (O'Neill et al., 2019), thermokarst potential (Olefelt et al., 2016), and carbon content (Hugelius et al., 2014). Significant advances in the development of relevant datasets on surficial materials, marine limits, peatland distribution, and peat ages and thicknesses, along with field-based advances in climate monitoring for cloud cover, fog, and snow, are critically needed to better characterize northern coastal regions like Labrador. Our results highlight the importance of field-based validation for periglacial landform mapping and modelling and of considering physiography and geomorphology for accurate representations of peatland permafrost in larger scale spatial products. The significant underestimation of peatland permafrost along the Labrador Sea coastline shown in this study should inform future permafrost, peatland permafrost, and carbon content mapping efforts, infrastructure and climate change adaptation strategy development, and wildlife management considerations for Labrador and other northern coastal regions.

Data availability. Likely and possible peatland permafrost locations from the coastal Labrador peatland permafrost complex inventory are freely available for download from Nordicana D (Wang et al., 2022).

Author contribution. YW and RW designed the study and drafted the manuscript. YW led the raw data collection and the data analysis. RW contributed to raw data collection and data analysis and was the PI for the NSERC Discovery Grant supporting peatland permafrost research activities in Labrador. JB contributed to raw data collection and data analysis. AF and RT contributed to raw data collection. MP coordinated the collection of helicopter video footage. JB, AF, RT, and MP reviewed and contributed edits to the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The authors would like to acknowledge the Nunatsiavut Government (Rodd Laing), the Nunatsiavut Research Centre (Carla Pamak, Michelle Saunders), the NunatuKavut Community Council (Bryn Wood, George Russell Jr., Charlene Kippenhuck), and the Innu Nation (Jonathan Feldgajer, Jack Penashue) for their guidance on research conducted on traditional Inuit and Innu lands. We thank Caitlin Lapalme, George Way, and Amy Norman of Goose Bay; Freeman Butt and Tanya Barney of Red Bay; Jeffrey and Wendy Keefe of Black Tickle; Barbara Mesher of Cartwright; Jane and Jack Shiwak, Tyler and Harvey Palliser, and Sandi and Karl Michelin of Rigolet; and Caroline Nochasak, Liz Pijogge, Carla Pamak, Kayla Wyatt, Frédéric Dwyer-Samuel, and Patricia Johnson-Castle of Nain for their logistical and in-kind support. We acknowledge Caitlin Lapalme, Victoria Colyn, Kayla Wyatt, Frédéric Dwyer-Samuel, Adrian Earle, Patrick Lauriault, Michaela Smitas-Kraas, and Tara Ryan for their valuable field assistance, and we are grateful to Derrick Pottle, Tyler Palliser, Martin Shiwak, Reginald Maggo, Jeremy Ivany, Martin Andersen, Eldred Andersen, Jeffrey Keefe, Gary Bird, Pat Davis, and Anthony Elson for their boat operation and guiding services. Funding for this research was provided by the Natural Sciences and Engineering

474 Research Council of Canada, the Northern Scientific Training Program, and Queen's University. We thank Dr. Antoni
475 Lewkowicz and Dr. Luise Hermanutz for many insightful conversations about permafrost and peatlands in Labrador. The
476 manuscript has also benefitted from helpful comments and suggestions from Michelle Saunders, Dr. H. Brendan O'Neill, and
477 Dr. Hanna Lee, and we thank two anonymous referees and Dr. Steve Kokelj for their constructive reviews.

478 **References**

- 479 Allard, M. and Rousseau, L.: The international structure of a palsa and a peat plateau in the Rivière Boniface region, Québec:
480 Inferences on the formation of ice segregation mounds, *Géographie Phys. Quat.*, 53, 373–387,
481 <https://doi.org/10.7202/004760ar>, 1999.
- 482 Allard, M. and Seguin, M. K.: Le pergélisol au Québec nordique : bilan et perspectives, *Géographie Phys. Quat.*, 41, 141–152,
483 <https://doi.org/10.7202/032671ar>, 1987.
- 484 Anderson, D., Ford, J. D., and Way, R. G.: The impacts of climate and social changes on cloudberry (bakeapple) picking: A
485 case study from southeastern Labrador, *Hum. Ecol.*, 46, 849–863, <https://doi.org/10.1007/s10745-018-0038-3>, 2018.
- 486 Andrews, J. T.: The glacial geomorphology of the northern Nain-Okak section of Labrador, M.Sc. Thesis, McGill University,
487 Montreal, Canada, 301 pp., 1961.
- 488 Banfield, C. E. and Jacobs, J. D.: Regional patterns of temperature and precipitation for Newfoundland and Labrador during
489 the past century, *Can. Geogr. Géographe Can.*, 42, 354–64, 1998.
- 490 Barrette, C., Brown, R., Way, R. G., Mailhot, A., Diaconescu, E. P., Grenier, P., Chaumont, D., Dumont, D., Sévigny, C.,
491 Howell, S., and Senneville, S.: Nunavik and Nunatsiavut regional climate information update, in: Nunavik and Nunatsiavut:
492 From science to policy, an integrated regional impact study (IRIS) of climate change and modernization, second iteration,
493 edited by: Ropars, P., Allard, M., and Lemay, M., ArcticNet Inc., Quebec City, Canada, 62, 2020.
- 494 Beilman, D. W., Vitt, D. H., and Halsey, L. A.: Localized permafrost peatlands in western Canada: Definition, distributions,
495 and degradation, *Arct. Antarct. Alp. Res.*, 33, 70–77, <https://doi.org/10.1080/15230430.2001.12003406>, 2001.
- 496 Bell, T., Putt, M., and Sheldon, T.: Landscape hazard assessment in Nain, Phase I: Inventory of surficial sediment types and
497 infrastructure damage, Final Report to Nunatsiavut Government and Nain Inuit Community Government, 2011.
- 498 Boisson, A. and Allard, M.: Coastal classification of Nunavik and dynamics of the Arctic/Subarctic coastal environments,
499 Vulnerabilities of the Quebec's Arctic Territory in the Context of Climate Change, Kuujuaq, Canada, 2018.
- 500 Borge, A. F., Westermann, S., Solheim, I., and Etzelmüller, B.: Strong degradation of palsas and peat plateaus in northern
501 Norway during the last 60 years, *The Cryosphere*, 11, 1–16, <https://doi.org/10.5194/tc-11-1-2017>, 2017.
- 502 Brown, R. and Lemay, M.: Chapter 2: Climate variability and change in the Canadian Eastern Subarctic IRIS region (Nunavik
503 and Nunatsiavut), in: Nunavik and Nunatsiavut: From science to policy. An Integrated Regional Impact Study (IRIS) of climate
504 change and modernization, ArcticNet Inc., <https://doi.org/10.13140/2.1.1041.7284>, 2012.
- 505 Brown, R. J. E.: Permafrost investigations in Quebec and Newfoundland (Labrador), National Research Council of Canada,
506 Division of Building Research, Technical Paper 449, <https://doi.org/10.4224/20374659>, 1975.

507 Brown, R. J. E.: Permafrost distribution in the southern part of the discontinuous zone in Québec and Labrador, *Géographie*
508 *Phys. Quat.*, 33, 279–289, <https://doi.org/10.7202/1000364ar>, 1979.

509 Burn, C. R. and Smith, C. A. S.: Observations of the “thermal offset” in near-surface mean annual ground temperatures at
510 several sites near Mayo, Yukon Territory, Canada, *ARCTIC*, 41, 99–104, <https://doi.org/10.14430/arctic1700>, 1988.

511 Coultish, T. L. and Lewkowicz, A. G.: Palsa dynamics in a subarctic mountainous environment, Wolf Creek, Yukon Territory,
512 Canada, *Proc. 8th Int. Conf. Permafr.*, 6, 2003.

513 Davis, E., Trant, A., Hermanutz, L., Way, R. G., Lewkowicz, A. G., Siegwart Collier, L., Cuerrier, A., and Whitaker, D.:
514 Plant–environment interactions in the low Arctic Torngat Mountains of Labrador, *Ecosystems*, 24, 1038–1058,
515 <https://doi.org/10.1007/s10021-020-00577-6>, 2020.

516 Dionne, J.-C.: Paleses et limite méridionale du pergélisol dans l’hémisphère nord : Le cas de Blanc-Sablon, Québec, *Géographie*
517 *Phys. Quat.*, 38, 165–184, <https://doi.org/10.7202/032550ar>, 1984.

518 Dyke, A. S.: An outline of North American deglaciation with emphasis on central and northern Canada, in: *Developments in*
519 *Quaternary Sciences*, vol. 2, Elsevier, 373–424, [https://doi.org/10.1016/S1571-0866\(04\)80209-4](https://doi.org/10.1016/S1571-0866(04)80209-4), 2004.

520 Dyke, A. S., Dredge, L. A., and Hodgson, D. A.: North American deglacial marine- and lake-limit surfaces, *Géographie Phys.*
521 *Quat.*, 59, 155–185, <https://doi.org/10.7202/014753ar>, 2005.

522 Elias, S. A.: Paleoenvironmental interpretation of Holocene insect fossils from northeastern Labrador, Canada, *Arct. Alp. Res.*,
523 14, 311, <https://doi.org/10.2307/1550794>, 1982.

524 Environment and Climate Change Canada: Canadian Climate Normals, 2022.

525 Environment Canada: Audio Tape Transcript of the East-Central Labrador Ecological Land Survey,
526 https://ftp.maps.canada.ca/pub/nrcan_rncan/archive/vector/labrador/, 1999.

527 Esri: World Imagery, 2022.

528 Fewster, R. E., Morris, P. J., Swindles, G. T., Gregoire, L. J., Ivanovic, R. F., Valdes, P. J., and Mullan, D.: Drivers of Holocene
529 palsa distribution in North America, *Quat. Sci. Rev.*, 240, 106337, <https://doi.org/10.1016/j.quascirev.2020.106337>, 2020.

530 Foster, D. R.: The history and pattern of fire in the boreal forest of southeastern Labrador, *Can. J. Bot.*, 61, 2459–2471,
531 <https://doi.org/10.1139/b83-269>, 1983.

532 Foster, D. R. and Glaser, P. H.: The raised bogs of south-eastern Labrador, Canada: Classification, distribution, vegetation and
533 recent dynamics, *J. Ecol.*, 74, 47, <https://doi.org/10.2307/2260348>, 1986.

534 Fulton, R. J. (Ed.): Quaternary Geology of Canada and Greenland, Minister of Supply and Services Canada, Ottawa, Canada,
535 846 pp., 1989.

536 Fulton, R. J.: Surficial Materials of Canada, “A” Series Map 1880A, Natural Resources Canada, Geological Survey of Canada,
537 <https://doi.org/10.4095/205040>, 1995.

538 Gibson, C., Morse, P. D., Kelly, J. M., Turetsky, M. R., Baltzer, J. L., Gingras-Hill, T., and Kokelj, S. V.: Thermokarst mapping
539 collective: Protocol for organic permafrost terrain and preliminary inventory from the Taiga Plains test area, Northwest
540 Territories, N. W. T. Open Rep., 2020, 29, 2020.

541 Gibson, C., Cottenie, K., Gingras-Hill, T., Kokelj, S. V., Baltzer, J. L., Chasmer, L., and Turetsky, M. R.: Mapping and
542 understanding the vulnerability of northern peatlands to permafrost thaw at scales relevant to community adaptation planning,
543 *Environ. Res. Lett.*, 16, 055022, <https://doi.org/10.1088/1748-9326/abe74b>, 2021.

544 Gorham, E.: Northern peatlands: Role in the carbon cycle and probable responses to climatic warming, *Ecol. Appl.*, 1, 182–
545 195, <https://doi.org/10.2307/1941811>, 1991.

546 Gorham, E., Lehman, C., Dyke, A., Janssens, J., and Dyke, L.: Temporal and spatial aspects of peatland initiation following
547 deglaciation in North America, *Quat. Sci. Rev.*, 26, 300–311, <https://doi.org/10.1016/j.quascirev.2006.08.008>, 2007.

548 Hagedorn, G. W.: Preliminary delineation of marine sediments in east-central Labrador: Parts of NTS map areas 13F, -G, -I, -
549 J, -K, -N AND -O, Gov. Nfld. Labrador Dep. Ind. Energy Technol. Curr. Res., 22, 189–201, 2022.

550 Hare, F. K.: Climate and zonal divisions of the boreal forest formation in eastern Canada, *Geogr. Rev.*, 40, 615,
551 <https://doi.org/10.2307/211106>, 1950.

552 Heginbottom, J. A., Dubreuil, M. A., and Harker, P. T.: Canada, Permafrost, National Atlas of Canada, Natural Resources
553 Canada, Geomatics Canada, <https://doi.org/10.4095/294672>, 1995.

554 Heginbottom, J. A., Brown, J., Melnikov, E. S., and Ferrians Jr., O. J.: Circumarctic map of permafrost and ground ice
555 conditions, in: Permafrost: Sixth International Conference, Sixth International Conference on Permafrost, Guangzhou, China,
556 1132–1136, 1997.

557 Holloway, J. E. and Lewkowicz, A. G.: Half a century of discontinuous permafrost persistence and degradation in western
558 Canada, *Permafr. Periglac. Process.*, 31, 85–96, <https://doi.org/10.1002/ppp.2017>, 2020.

559 Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. A. G., Ping, C.-L., Schirrmeister, L., Grosse, G., Michaelson,
560 G. J., Koven, C. D., O'Donnell, J. A., Elberling, B., Mishra, U., Camill, P., Yu, Z., Palmtag, J., and Kuhry, P.: Estimated stocks
561 of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps, *Biogeosciences*, 11, 6573–6593,
562 <https://doi.org/10.5194/bg-11-6573-2014>, 2014.

563 Hugelius, G., Loisel, J., Chadburn, S., Jackson, R. B., Jones, M., MacDonald, G., Marushchak, M., Olefeldt, D., Packalen, M.,
564 Siewert, M. B., Treat, C., Turetsky, M., Voigt, C., and Yu, Z.: Large stocks of peatland carbon and nitrogen are vulnerable to
565 permafrost thaw, *Proc. Natl. Acad. Sci.*, 117, 20438–20446, <https://doi.org/10.1073/pnas.1916387117>, 2020.

566 Hustich, I.: Notes on the coniferous forest and tree limit on the east coast of Newfoundland-Labrador, *Acta Geogr.*, 7, 81,
567 1939.

568 International Permafrost Association Terminology Working Group: Multi-language glossary of permafrost and related ground-
569 ice terms, edited by: van Everdingen, R. O., 2005.

570 Ives, J. D.: A proposed history of permafrost development in Labrador-Ungava, *Géographie Phys. Quat.*, 33, 233–244,
571 <https://doi.org/10.7202/1000360ar>, 1979.

572 Jorgenson, M. T., Romanovsky, V., Harden, J., Shur, Y., O'Donnell, J., Schuur, E. A. G., Kanevskiy, M., and Marchenko, S.:
573 Resilience and vulnerability of permafrost to climate change, *Can. J. For. Res.*, 40, 1219–1236, [https://doi.org/10.1139/X10-](https://doi.org/10.1139/X10-060)
574 060, 2010.

575 Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E., Linder, H. P., and
576 Kessler, M.: Climatologies at high resolution for the earth's land surface areas, *Sci. Data*, 4, 170122,
577 <https://doi.org/10.1038/sdata.2017.122>, 2017.

578 Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E., Linder, H. P., and
579 Kessler, M.: Climatologies at high resolution for the earth's land surface areas V2.1, 2021.

580 Karst, A. L. and Turner, N. J.: Local ecological knowledge and importance of bakeapple (*Rubus chamaemorus* L.) in a
581 southeast Labrador Métis community, *Ethnobiol. Lett.*, 2, 6–18, <https://doi.org/10.14237/eb1.2.2011.28>, 2011.

582 Mamet, S. D., Chun, K. P., Kershaw, G. G. L., Loranty, M. M., and Kershaw, G. P.: Recent increases in permafrost thaw rates
583 and areal loss of palsas in the western Northwest Territories, Canada: Non-linear palsa degradation, *Permafr. Periglac. Process.*,
584 28, 619–633, <https://doi.org/10.1002/ppp.1951>, 2017.

585 McLaughlin, J. and Webster, K.: Effects of climate change on peatlands in the far north of Ontario, Canada: A synthesis, *Arct.*
586 *Antarct. Alp. Res.*, 46, 84–102, <https://doi.org/10.1657/1938-4246-46.1.84>, 2014.

587 Natural Resources Canada: Standards and Specifications of the National Topographic Data Base, Edition 3.1, 2005.

588 Norton, C. H., Cuerrier, A., and Hermanutz, L.: People and plants in Nunatsiavut (Labrador, Canada): Examining plants as a
589 foundational aspect of culture in the Subarctic, *Econ. Bot.*, 75, 287–301, <https://doi.org/10.1007/s12231-021-09530-7>, 2021.

590 Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H. H., Dashtseren, A., Delaloye, R., Elberling, B.,
591 Etzelmüller, B., Kholodov, A., Khomutov, A., Kääb, A., Leibman, M. O., Lewkowicz, A. G., Panda, S. K., Romanovsky, V.,
592 Way, R. G., Westergaard-Nielsen, A., Wu, T., Yamkhin, J., and Zou, D.: Northern Hemisphere permafrost map based on
593 TTOP modelling for 2000–2016 at 1 km² scale, *Earth-Sci. Rev.*, 193, 299–316,
594 <https://doi.org/10.1016/j.earscirev.2019.04.023>, 2019.

595 Occhietti, S., Parent, M., Lajeunesse, P., Robert, F., and Govare, É.: Late Pleistocene–early Holocene decay of the Laurentide
596 Ice Sheet in Québec–Labrador, in: *Developments in Quaternary Sciences*, vol. 15, Elsevier, 601–630,
597 <https://doi.org/10.1016/B978-0-444-53447-7.00047-7>, 2011.

598 Olefeldt, D., Goswami, S., Grosse, G., Hayes, D., Hugelius, G., Kuhry, P., McGuire, A. D., Romanovsky, V. E., Sannel, A.
599 B. K., Schuur, E. A. G., and Turetsky, M. R.: Circumpolar distribution and carbon storage of thermokarst landscapes, *Nat.*
600 *Commun.*, 7, 13043, <https://doi.org/10.1038/ncomms13043>, 2016.

601 Olefeldt, D., Hovemyr, M., Kuhn, M. A., Bastviken, D., Bohn, T. J., Connolly, J., Crill, P., Euskirchen, E. S., Finkelstein, S.
602 A., Genet, H., Grosse, G., Harris, L. I., Heffernan, L., Helbig, M., Hugelius, G., Hutchins, R., Juutinen, S., Lara, M. J.,
603 Malhotra, A., Manies, K., McGuire, A. D., Natali, S. M., O'Donnell, J. A., Parmentier, F.-J. W., Räsänen, A., Schädel, C.,
604 Sonnentag, O., Strack, M., Tank, S. E., Treat, C., Varner, R. K., Virtanen, T., Warren, R. K., and Watts, J. D.: The Boreal–
605 Arctic Wetland and Lake Dataset (BAWLD), *Earth Syst. Sci. Data*, 13, 5127–5149, [https://doi.org/10.5194/essd-13-5127-](https://doi.org/10.5194/essd-13-5127-2021)
606 2021, 2021.

607 O'Neill, H. B., Wolfe, S. A., and Duchesne, C.: New ground ice maps for Canada using a paleogeographic modelling approach,
608 *The Cryosphere*, 13, 753–773, <https://doi.org/10.5194/tc-13-753-2019>, 2019.

609 Ou, C., LaRocque, A., Leblon, B., Zhang, Y., Webster, K., and McLaughlin, J.: Modelling and mapping permafrost at high
610 spatial resolution using Landsat and Radarsat-2 images in Northern Ontario, Canada: Part 2 – regional mapping, *Int. J. Remote*
611 *Sens.*, 37, 2751–2779, <https://doi.org/10.1080/01431161.2016.1151574>, 2016.

- 612 Parviainen, M. and Luoto, M.: Climate envelopes of mire complex types in fennoscandia, *Geogr. Ann. Ser. Phys. Geogr.*, 89,
613 137–151, <https://doi.org/10.1111/j.1468-0459.2007.00314.x>, 2007.
- 614 Payette, S.: The forest tundra and present tree-lines of the northern Québec-Labrador peninsula, *Proc. North. Qué. Tree-Line*
615 *Conf.*, 47, 3–23, 1983.
- 616 Payette, S.: Chapitre 9: Les processus et les formes périglaciaires, in: *Écologie des tourbières du Québec-Labrador*, Presses de
617 l'Université Laval, 42, 2001.
- 618 Payette, S.: Accelerated thawing of subarctic peatland permafrost over the last 50 years, *Geophys. Res. Lett.*, 31, L18208,
619 <https://doi.org/10.1029/2004GL020358>, 2004.
- 620 Pironkova, Z.: Mapping palsa and peat plateau changes in the Hudson Bay Lowlands, Canada, using historical aerial
621 photography and high-resolution satellite imagery, *Can. J. Remote Sens.*, 43, 455–467,
622 <https://doi.org/10.1080/07038992.2017.1370366>, 2017.
- 623 R Core Team: R, 2020.
- 624 Ramsdale, J. D., Balme, M. R., Conway, S. J., Gallagher, C., van Gasselt, S. A., Hauber, E., Orgel, C., Séjourné, A., Skinner,
625 J. A., Costard, F., Johnsson, A., Losiak, A., Reiss, D., Swirad, Z. M., Kereszturi, A., Smith, I. B., and Platz, T.: Grid-based
626 mapping: A method for rapidly determining the spatial distributions of small features over very large areas, *Planet. Space Sci.*,
627 140, 49–61, <https://doi.org/10.1016/j.pss.2017.04.002>, 2017.
- 628 Roberts, B. A., Simon, N. P. P., and Deering, K. W.: The forests and woodlands of Labrador, Canada: Ecology, distribution
629 and future management, *Ecol. Res.*, 21, 868–880, <https://doi.org/10.1007/s11284-006-0051-7>, 2006.
- 630 Seguin, M. K. and Dionne, J. C.: Modélisation géophysique et caractérisation thermique du pergélisol dans les palses de Blanc-
631 Sablon, Quebec, *Geol. Surv. Can. Curr. Res.*, E, 207–216, 1992.
- 632 Seppälä, M.: The origin of palsas, *Geogr. Ann.*, 68, 141–147, 1986.
- 633 Seppälä, M.: Snow depth controls palsa growth, *Permafr. Periglac. Process.*, 5, 283–288,
634 <https://doi.org/10.1002/ppp.3430050407>, 1994.
- 635 Smith, J. S.: Shifting sites and shifting sands: A record of prehistoric human/landscape interactions from Porcupine Strand,
636 Labrador, M.Sc. Thesis, Memorial University of Newfoundland, St. John's, Canada, 288 pp., 2003.
- 637 Smith, M. W. and Riseborough, D. W.: Climate and the limits of permafrost: A zonal analysis, *Permafr. Periglac. Process.*, 13,
638 1–15, <https://doi.org/10.1002/ppp.410>, 2002.
- 639 Tarnocai, C.: The impact of climate change on Canadian peatlands, *Can. Water Resour. J.*, 34, 453–466,
640 <https://doi.org/10.4296/cwrj3404453>, 2009.
- 641 Tarnocai, C., Kettles, I. M., and Lacelle, B.: Peatlands of Canada, Natural Resources Canada, Geological Survey of Canada,
642 Open File 6561, <https://doi.org/10.4095/288786>, 2011.
- 643 Thibault, S. and Payette, S.: Recent permafrost degradation in bogs of the James Bay area, northern Quebec, Canada, *Permafr.*
644 *Periglac. Process.*, 20, 383–389, <https://doi.org/10.1002/ppp.660>, 2009.

645 Thie, J.: Distribution and thawing of permafrost in the southern part of the discontinuous permafrost zone in Manitoba,
646 ARCTIC, 27, 189–200, <https://doi.org/10.14430/arctic2873>, 1974.

647 Vacchi, M., Engelhart, S. E., Nikitina, D., Ashe, E. L., Peltier, W. R., Roy, K., Kopp, R. E., and Horton, B. P.: Postglacial
648 relative sea-level histories along the eastern Canadian coastline, *Quat. Sci. Rev.*, 201, 124–146,
649 <https://doi.org/10.1016/j.quascirev.2018.09.043>, 2018.

650 Vallée, S. and Payette, S.: Collapse of permafrost mounds along a subarctic river over the last 100 years (northern Québec),
651 *Geomorphology*, 90, 162–170, <https://doi.org/10.1016/j.geomorph.2007.01.019>, 2007.

652 Wang, Y., Way, R. G., and Beer, J.: Coastal Labrador peatland permafrost inventory, v. 1.0, Nord. D98,
653 <https://doi.org/10.5885/45762XD-1DB498A49B864CFB>, 2022.

654 Way, R. G.: Field and modelling investigations of permafrost conditions in Labrador, northeast Canada, Ph.D. Thesis,
655 University of Ottawa, Ottawa, Canada, 293 pp., 2017.

656 Way, R. G. and Lewkowicz, A. G.: Investigations of discontinuous permafrost in coastal Labrador with DC electrical resistivity
657 tomography, *Proc. 68th Can. Geotech. Conf. 7th Can. Permafr. Conf.*, 8, <https://doi.org/10.13140/RG.2.1.1647.8803>, 2015.

658 Way, R. G. and Lewkowicz, A. G.: Environmental controls on ground temperature and permafrost in Labrador, northeast
659 Canada, *Permafr. Periglac. Process.*, 29, 73–85, <https://doi.org/10.1002/ppp.1972>, 2018.

660 Way, R. G. and Viau, A. E.: Natural and forced air temperature variability in the Labrador region of Canada during the past
661 century, *Theor. Appl. Climatol.*, 121, 413–424, <https://doi.org/10.1007/s00704-014-1248-2>, 2015.

662 Way, R. G., Lewkowicz, A. G., and Bonnaventure, P. P.: Development of moderate-resolution gridded monthly air temperature
663 and degree-day maps for the Labrador-Ungava region of northern Canada: High-resolution air temperature and degree-day
664 maps for Labrador, *Int. J. Climatol.*, 37, 493–508, <https://doi.org/10.1002/joc.4721>, 2017.

665 Way, R. G., Lewkowicz, A. G., and Zhang, Y.: Characteristics and fate of isolated permafrost patches in coastal Labrador,
666 Canada, *The Cryosphere*, 12, 2667–2688, <https://doi.org/10.5194/tc-12-2667-2018>, 2018.

667 Way, R. G., Wang, Y., Bevington, A. R., Bonnaventure, P. P., Burton, J. R., Davis, E., Garibaldi, M. C., Lapalme, C. M.,
668 Tutton, R., and Wehbe, M. A.: Consensus-Based Rock Glacier Inventorying in the Torngat Mountains, Northern Labrador,
669 *Proc. 2021 Reg. Conf. Permafr. 19th Int. Conf. Cold Reg. Eng.*, 130–141, 2021a.

670 Way, R. G., Lewkowicz, A. G., Wang, Y., and McCarney, P.: Permafrost investigations below the marine limit at Nain,
671 Nunatsiavut, Canada, *Proc. 2021 Reg. Conf. Permafr. 19th Int. Conf. Cold Reg. Eng.*, 38–48, 2021b.

672 Wenner, C.-G.: Pollen diagrams from Labrador, *Geogr. Ann.*, 29, 137–374, 1947.

673 Williams, P. J. and Smith, M. W.: *The Frozen Earth: Fundamentals of Geocryology*, Cambridge University Press, 1989.

674 Zoltai, S. C.: Palsas and peat plateaus in central Manitoba and Saskatchewan, *Can. J. For. Res.*, 2, 291–302,
675 <https://doi.org/10.1139/x72-046>, 1972.

676 Zoltai, S. C. and Tarnocai, C.: Perennially frozen peatlands in the western Arctic and Subarctic of Canada, *Can. J. Earth Sci.*,
677 12, 28–43, <https://doi.org/10.1139/e75-004>, 1975.

678 Zuidhoff, F. S. and Kolstrup, E.: Palsa development and associated vegetation in northern Sweden, *Arct. Antarct. Alp. Res.*,
679 37, 49–60, [https://doi.org/10.1657/1523-0430\(2005\)037\[0049:PDAAVI\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2005)037[0049:PDAAVI]2.0.CO;2), 2005.

680