

1 Significant underestimation of peatland permafrost along the 2 Labrador Sea coastline in northern Canada

3 Yifeng Wang¹, Robert G. Way¹, Jordan Beer¹, Anika Forget¹, Rosamond Tutton^{1,2}, Meredith C. Purcell³

4 ¹Northern Environmental Geoscience Laboratory, Department of Geography and Planning, Kingston, K7L 3N6, Canada

5 ²Global Water Futures, Wilfrid Laurier University, Yellowknife, X1A 2P8, Canada

6 ³Tornat Wildlife, Plants, and Fisheries Secretariat, Happy Valley-Goose Bay, A0P 1E0, Canada

7 *Correspondence to:* Yifeng Wang (yifeng.wang@queensu.ca)

8 **Abstract.** Northern peatlands cover approximately four million km², and about half of these peatlands are estimated to contain
9 permafrost and periglacial landforms, like palsas and peat plateaus. In northeastern Canada, peatland permafrost is predicted
10 to be concentrated in the western interior of Labrador but is assumed to be largely absent along the Labrador Sea coastline.
11 However, the paucity of observations of peatland permafrost in the interior, coupled with traditional and ongoing use of
12 perennially frozen peatlands along the coast by Labrador Inuit and Innu, suggests a need for re-evaluation of the reliability of
13 existing peatland permafrost distribution estimates for the region. In this study, we develop a multi-stage consensus-based
14 point inventory of peatland permafrost complexes in coastal Labrador and adjacent parts of Quebec using high-resolution
15 satellite imagery, and we validate it with extensive field visits and low-altitude aerial photography and videography. A subset
16 of 2092 wetland complexes that potentially contained peatland permafrost were inventoried, of which 1119 were classified as
17 likely containing peatland permafrost. Likely peatland permafrost complexes were mostly found in lowlands within 22 km of
18 the coastline where mean annual air temperatures often exceed +1 °C. A clear gradient in peatland permafrost distribution
19 exists from the outer coasts, where peatland permafrost is more abundant, to inland peatlands, where permafrost is generally
20 absent. This coastal gradient may be attributed to a combination of climatic and geomorphological influences which lead to
21 lower insolation, thinner snowpacks, and poorly drained, frost-susceptible materials along the coast. The results of this study
22 suggest that existing estimates of permafrost distribution for southeastern Labrador require adjustments to better reflect the
23 abundance of peatland permafrost complexes to the south of the regional sporadic discontinuous permafrost limit. This study
24 constitutes the first dedicated peatland permafrost inventory for Labrador and provides an important baseline for future
25 mapping, modelling, and climate change adaptation strategy development in the region.

26 1 Introduction

27 Near the southern boundary of latitudinal permafrost zonation, lowland perennially frozen ground is primarily
28 restricted to wetlands in the form of palsas (peat mounds with a frozen core of mineral and organic material) and peat plateaus
29 (fields of frozen peat elevated above the general surface of the surrounding peatland) (Payette, 2004; International Permafrost
30 Association Terminology Working Group, 2005; Zoltai, 1972; Zoltai and Tarnocai, 1975). Persistence of these cryotic

31 landforms at the extreme limits of their viability is facilitated by a large temperature offset between the ground surface and the
32 top of permafrost, caused by the thermal properties of thick layers of overlying peat and the buffering effect of ground ice
33 (Burn and Smith, 1988; Williams and Smith, 1989). In recent years, many studies have shown that peatland permafrost can be
34 very sensitive to climate warming and ecosystem modifications (Beilman et al., 2001; Borge et al., 2017; Thibault and Payette,
35 2009). Understanding the distribution of these ice-rich, thaw-sensitive periglacial environments is important for assessing
36 thermokarst potential (Gibson et al., 2021; Olefeldt et al., 2016), local hydrological and vegetation change (Zuidhoff and
37 Kolstrup, 2005), regional infrastructure or land-use planning, and global carbon stores and carbon cycling (Hugelius et al.,
38 2014).

39 Palsas and peat plateaus are primarily thought to occur in continental locations (Fewster et al., 2020; Hustich, 1939)
40 where colder winters allow deeper frost penetration and drier summers promote less thaw. As such, palsas and peat plateaus
41 have been described in many continental locations in Canada, including Yukon Territory, the Northwest Territories, and the
42 Prairie provinces (e.g., Beilman et al., 2001; Coultish and Lewkowicz, 2003; Mamet et al., 2017; Thie, 1974; Zoltai, 1972).
43 However, these landforms have also been documented in coastal locations including the Hudson Bay Lowlands in northern
44 Manitoba, Ontario, and Quebec (e.g., McLaughlin and Webster, 2014; Ou et al., 2016; Pironkova, 2017). In the Labrador
45 region of northeastern Canada, continental- to hemispheric-scale studies have depicted peatland permafrost as present in the
46 region's continental interior but as far less abundant or completely absent along most of the Labrador Sea coastline (Fewster
47 et al., 2020; Hugelius et al., 2020; Olefeldt et al., 2021). However, historic and contemporary use of coastal peatland permafrost
48 environments by Labrador Inuit and Innu is well documented (Anderson et al., 2018), and published field-based observations
49 (e.g., Anderson et al., 2018; Andrews, 1961; Brown, 1975, 1979; Davis et al., 2020; Dionne, 1984; Elias, 1982; Hustich, 1939;
50 Seguin and Dionne, 1992; Smith, 2003; Way et al., 2018; Wenner, 1947) suggest that peatland permafrost is abundant along
51 some sections of the coast. This recurring misestimation of peatland permafrost occurrence has an impact on predictions of
52 ground ice content (O'Neill et al., 2019), thermokarst potential (Olefeldt et al., 2016), and carbon content (Hugelius et al.,
53 2014) in the region.

54 Locally, preservation of peatland permafrost complexes is relevant to Labrador Inuit and Innu because these areas are
55 frequented for traditional activities such as bakeapple (cloudberry; Inuttitut: appik; Innu-aimun: shikuteu; *Rubus*
56 *chamaemorus*) berry-picking (Anderson et al., 2018; Karst and Turner, 2011; Norton et al., 2021), goose hunting, and fox
57 trapping (Way et al., 2018). Improvements to our understanding of regional peatland permafrost distribution will provide an
58 important baseline for local and regional climate change adaptation strategy development, while better representation of the
59 distribution of thaw-sensitive terrain will inform future development of linear and built infrastructure in coastal Labrador (Way
60 et al., 2021b; Bell et al., 2011).

61 Previous peatland permafrost mapping in Labrador has been limited to scattered observations of palsa bogs from the
62 National Topographic Database (Natural Resources Canada, 2005) and the Ecological Land Classification (Environment
63 Canada, 1999), with no comprehensive peatland permafrost inventorying efforts completed to date (Way et al., 2018). The
64 objectives of this study are to: 1) use a multi-stage, consensus-based review process, coupled with extensive validation efforts

65 from a combination of field visits and low-altitude image and video acquisitions, to develop a point inventory of contemporary
66 peatland permafrost complexes in coastal Labrador; 2) characterize the distribution of peatland permafrost in coastal Labrador
67 using selected climatic and physiographic variables; and 3) provide insights into the reliability of relevant peatland permafrost
68 and permafrost distribution products, which currently claim an absence or low abundance of both peatland permafrost and
69 permafrost along the Labrador Sea coastline. This study is limited to peatland permafrost complexes located within 100 km of
70 the Labrador Sea coastline (Figure 1), comprising the region of Nunatsiavut and surrounding areas, including the land claims
71 agreement-in-principle of the Labrador Innu Nation (Nitassinan) and coastal areas claimed by the NunatuKavut Community
72 Council (NunatuKavut). We expect that this point-based inventory will reinforce the local understanding of a high abundance
73 of peatland permafrost landforms in coastal locations, which will be relevant for carbon modelling, land use planning,
74 infrastructure development, and climate change adaptation strategy development at local to regional scales in northeastern
75 Canada. Based on our results, we also propose amendments to the current limits of the sporadic discontinuous and isolated
76 patches of permafrost distribution zones in southeastern Labrador. This point-based inventory is a first step towards
77 understanding the distribution of peatland permafrost in Labrador and will contribute to refined regional and global estimates
78 of ground ice content, thermokarst potential, and carbon storage in northern Canada.




Figure 1. (a) Permafrost zonation in Labrador (Heginbottom et al., 1995) with the boundary for the inventory study area (black line) corresponding to areas within 100 km of the Labrador Sea coastline. Map is annotated with locations of the Torngat Mountains National Park, Akami-Uapishku-KakKasuak-Mealy Mountains National Park Reserve, and selected communities; (b) Inset map showing Labrador's position in Canada.

84 2 Study area

85 2.1 Bioclimatic setting

86 Labrador's climate is strongly influenced by atmosphere-ocean interactions from the adjacent Labrador Sea (Barrette
87 et al., 2020; Way and Viau, 2015). In coastal Labrador, long, cold winters and short, cool summers are largely dictated by the
88 Labrador Current that carries cold Arctic waters down the eastern coast of mainland Canada (Banfield and Jacobs, 1998;
89 Foster, 1983; Roberts et al., 2006; Way et al., 2017). Mean annual air temperatures (1980-2010) decrease with continentality
90 and latitude, ranging from -12°C in parts of the Torngat Mountains National Park to +1.5°C near the community of Blanc-
91 Sablon (Karger et al., 2017, 2021). Labrador is also characterized by some of the highest precipitation amounts in the North
92 American boreal zone (Banfield and Jacobs, 1998; Hare, 1950) due to its varying relief, high moisture availability from the
93 adjacent Atlantic Ocean, and high frequency of passing winter storm systems (Brown and Lemay, 2012). Precipitation totals
94 as high as ~2700 mm per year are estimated for some locations at high elevations along the coast (Karger et al., 2017, 2021),
95 with snow fractions increasing with both latitude and elevation (~0.35 at Blanc-Sablon; ~0.5 at Nain) (Environment and
96 Climate Change Canada, 2022).

97 Ecologically, Labrador is characterized by taiga forests in the interior, tundra in the north, and wind-swept coastal
98 barrens along the coastline of the Labrador Sea (Roberts et al., 2006). Tree cover is sparse in the coastal barrens because of
99 climatic and physiographic limitations, but dense patches of black spruce (*Picea mariana*), white spruce (*Picea glauca*),
100 tamarack (*Larix laricina*), and balsam fir (*Abies balsamea*), interspersed with deciduous trees, like paper birch (*Betula*
101 *papyrifera*) and trembling aspen (*Populus tremuloides*), exist in sheltered locations and on some slopes (Roberts et al., 2006).
102 Wetlands are found throughout Labrador, but total wetland abundance is difficult to assess given widespread disagreement
103 between existing estimates of wetland and peatland extents for this region (Supplement Sect. S1). Generally, wetlands in
104 Labrador tend to decrease in abundance but increase in size as latitude increases. Most wetlands along the southern Labrador
105 coast are classified as raised bogs, while inland, most wetlands are string and blanket bogs (Foster and Glaser, 1986).

106 2.2 Physical environment

107 Labrador is mostly underlain by igneous and metamorphic bedrock (Roberts et al., 2006). Extensive blankets of
108 glacial till were deposited during the retreat of the Laurentide Ice Sheet (12-6 kyr BP) (Bell et al., 2011; Dyke, 2004), along
109 with thin layers of medium- to fine-grained marine and glaciomarine sediments in coastal lowland areas below the marine
110 limit (Fulton, 1995). The post-glacial marine limit decreases with latitude, from ~150 m a.s.l. in southeastern Labrador and
111 along the Quebec Lower North Shore to 0 m a.s.l. at the northernmost tip of Labrador in the Torngat Mountains (Dyke et al.,
112 2005; Occhietti et al., 2011; Vacchi et al., 2018). The broad distribution of near-surface bedrock and hardpans (Smith, 2003)
113 results in poor drainage that has facilitated peatland development across large areas of southern Labrador, particularly in
114 depressions and over flat areas.

115 2.3 Permafrost distribution

116 While permafrost conditions in Labrador, including the presence of peatland permafrost landforms, have been noted
117 during ecological, palynological, glaciological, and archeological surveys and studies (Anderson et al., 2018; Andrews, 1961;
118 Hustich, 1939; Smith, 2003; Wenner, 1947), permafrost-specific field investigations are limited to R.J.E. Brown's (1975)
119 helicopter survey in the late 1960s and the Labrador Permafrost Project that began in 2013 (Way, 2017). Our understanding
120 of permafrost distribution in Labrador has relied on extensive extrapolation of limited field observations and broad assumptions
121 of the interactions between air temperature, vegetation cover, snow cover, and permafrost presence (Ives, 1979). According to
122 the Permafrost Map of Canada (Heginbottom et al., 1995), the area underlain by permafrost in Labrador is less extensive than
123 comparable regions in northern Canada like Yukon Territory or the Northwest Territories. Approximately two-thirds of
124 Labrador is classified in the isolated patches of permafrost zone (<10 % permafrost by area), but the distribution of permafrost
125 is more widespread farther north (Figure 1). Along the Labrador coastline, the sporadic discontinuous permafrost zone (10-
126 50 % permafrost by area) extends slightly further south along the outer edge of the Akami-Uapishk-KakKasuak-Mealy
127 Mountains National Park Reserve than in the interior, though the justification for this departure is not clarified in published
128 literature. Continuous permafrost (>90 % permafrost by area) is expected to persist only at high elevations and latitudes, mostly
129 in the Torngat Mountains (Heginbottom et al., 1995).

130 2.4 Inventory extent

131 This study is focused on the coastal areas of Labrador and Quebec, within 100 km of the Labrador Sea coastline
132 (Figure 1). This area of interest was informed by knowledge gained from prior works in the region (Anderson et al., 2018;
133 Andrews, 1961; Brown, 1975, 1979; Davis et al., 2020; Dionne, 1984; Elias, 1982; Hustich, 1939; Seguin and Dionne, 1992;
134 Smith, 2003; Way, 2017; Way et al., 2018; Wenner, 1947) that indicated a greater abundance of peatland permafrost landforms
135 along the coast as compared to the interior of Labrador. Exhaustive descriptions of records of peatland permafrost and other
136 periglacial landforms in Labrador have been presented by Brown (1979) and Way (2017), both of whom found limited evidence
137 of peatland permafrost in Labrador's interior.

138 3 Methods

139 Palsas and peat plateaus are typically found in bogs and may measure up to 4 m higher than their surrounding
140 wetlands, so large peatland permafrost landforms can be identified and mapped from high-resolution satellite imagery (Borge
141 et al., 2017; Gibson et al., 2020, 2021). Our point inventory, which includes only the largest and most visually apparent peatland
142 permafrost complexes within 100 km of the Labrador Sea coastline, was generated through a multi-stage mapping and
143 consensus-based review process, supported by extensive validation efforts mostly completed between 2017 and 2022. Mapping
144 and identification activities were informed by existing wetland and peatland distribution products (Supplement Sect. S1), but

significant disagreement between these products limited their direct application and utility during the inventorying process. An initial inventory of wetlands of interest (WOIs) was developed as a subset of the wetlands in coastal Labrador deemed potentially suitable (e.g., bogs and fens) for the development and persistence of peatland permafrost landforms. The presence of peatland permafrost landforms within the WOIs was then evaluated through a consensus-based review of high-resolution satellite imagery by three mappers with permafrost-specific field experience in the region. Final interpretation of peatland permafrost presence or absence within the WOIs was based on reviewer agreement and was informed by field- and imagery-based validation of peatland permafrost landform presence or absence.

3.1 Data sources

WOIs were identified and evaluated using Maxar (Vivid) optical satellite imagery, available as the World Imagery basemap via ArcGIS Online (0.5 m ground sampling distance; 5 m absolute spatial accuracy) (Esri, 2022). These satellite imagery mosaics consisted of summer imagery with minimal cloud and snow cover, with acquisition dates for Labrador ranging from 2010 to 2020.

Topographic data from Natural Resources Canada covering the WOIs were extracted from the Canadian Digital Elevation Data (CDED; 50 m spatial resolution), with a small gap near the provincial border between Labrador and the Quebec Lower North Shore that was filled in using the Canadian Digital Surface Model (CDSM). Gridded mean annual air temperature (MAAT) and mean annual thawing degree days (TDD) for the 1981 to 2010 climate normal were extracted from CHELSA V2.1 (~1 km spatial resolution) (Karger et al., 2017, 2021) at the WOI locations. Mean annual freezing degree days (FDD) for the WOI locations for 1981 to 2010 were calculated from MAAT and TDD over the same climate normal, following prior work in the region (Way et al., 2017; Way and Lewkowicz, 2018).


3.2 Inventorying peatland permafrost complexes

3.2.1 Identifying wetlands of interest (WOIs)

A team of three mappers used ArcGIS Online to identify and place point features within WOIs throughout coastal Labrador (Figure 2). The point-based nature of the inventorying process allowed for evaluation of the entire study area by incorporating field- and imagery-based validation for many WOIs over a large study area, as opposed to detailed validation of peatland permafrost areal coverage within a given WOI. These WOIs were restricted to include only those that contained prospective peatland permafrost landforms exceeding 2 m in length or width (~4 m²), which was determined to be the smallest detectable feature based on the 0.5 m spatial resolution of the satellite imagery. Mappers were instructed to identify WOIs based on local geomorphology, local hydrology and drainage patterns, the presence of a white or grey lichen surface cover corresponding to *Cladonia* and/or *Ochrolechia* spp. lichens, shadows indicative of elevated landform edges and surface uplift, and thermokarst ponding or exposed peat indicative of thaw processes. The inventory sought to only include contemporary peatland permafrost landforms, so WOIs with extensive thermokarst ponding but no evident peatland permafrost landforms

176 were not included in the database. Individual WOIs ranged in size from ~0.2 km² to larger than ~3.5 km². However, the total
177 area underlain by peatland permafrost within each WOI was not able to be reliably evaluated using satellite imagery. WOIs
178 near one another were sometimes difficult to discern due to potential connectivity between adjacent systems, but contiguous
179 WOIs could generally be identified by differences in drainage, vegetation, and morphology, or because of separation by linear
180 infrastructure like roads, airstrips, and trails (Figure 2). Mappers also assigned each WOI a self-assessed score to reflect their
181 confidence in their interpretation of permafrost presence within the wetland complex (1 = low confidence, 2 = medium
182 confidence, 3 = high confidence).

183



184

185 **Figure 2. (a-b) Examples of wetland complexes of interest (WOIs) in Labrador that were identified by the mapping team using high-**
186 **resolution satellite imagery available via Esri ArcGIS Online. Examples of WOI boundaries are shown in red and were determined**
187 **based on differences in drainage or vegetation from adjacent WOIs or based on separation following linear infrastructure, such as**
188 **roads, airstrips, or trails. Identification was restricted to WOIs that contained prospective peatland permafrost landforms. Example**
189 **WOIs are presented from (a-b) Esri ArcGIS Online and (c-d) Esri ArcScene Online to provide nadir and oblique perspectives.**

190 3.2.2 Quality control of WOI database

191 The WOI inventory was subjected to a quality-control check, during which complexes were reviewed and duplicates
192 or points clearly not corresponding to wetlands were removed. In some cases, non-wetland locations may have been retained
193 because of difficulties discerning peat plateaus from surface peat over bedrock or coastal tundra.

194 3.2.3 Consensus-based review of WOI database

195 The quality-controlled WOI inventory was sent back to the mappers for a consensus-based review, similar to Way et
196 al.'s (2021a) approach for rock glacier inventorying in northern Labrador. Each WOI was independently reviewed by two team
197 members, both of whom had access to the mapper's initial confidence rating, and one of whom had access to a field-validated
198 dataset of WOIs (see Sect. 3.3 Validation of subset of WOI database). Both team members were asked to indicate whether
199 each WOI contained peatland permafrost landforms. WOIs evaluated by both reviewers as containing peatland permafrost
200 were considered likely to contain palsas or peat plateaus, while WOIs evaluated by both reviewers as not containing peatland
201 permafrost were considered unlikely to contain palsas or peat plateaus. WOIs with conflicting classifications were considered
202 to possibly contain palsas or peat plateaus. This consensus-based review process resulted in a full inventory of WOIs that were
203 classified as likely, possibly, or unlikely to contain peatland permafrost.

204 3.3 Validation of subset of WOI database

205 The full, consensus-based inventory results were compared with a field- and imagery-validated dataset of 557 WOIs,
206 with and without contemporary peatland permafrost landforms. From July to September 2021 and 2022, field evaluations of
207 WOIs were undertaken via in-person field visits, remotely piloted aircraft (RPA) image acquisitions (DJI Mini 2 microdrone,
208 weighing less than 250 g), video clip acquisition from a helicopter survey, and image acquisitions from commercial Twin Otter
209 aircraft flights. Interpretation of the presence or absence of permafrost landforms within each WOI that was visited or aerially
210 surveyed was also determined through consensus between two mappers. Any WOIs with disagreements in interpretation were
211 re-evaluated and discussed until consensus could be reached between the two mappers.

212 Field visits to WOIs were undertaken at road-accessible locations within 500 m of the Trans-Labrador Highway and
213 other accessible side roads via truck or ATV and at coastal locations via speedboat from the nearby communities of Black
214 Tickle, Cartwright, Rigolet, and Nain. The number of WOIs that could be visited for field validation was restricted by weather
215 conditions, tides, the availability of local guides and boat drivers with location-specific expertise, and other logistical and
216 operational constraints. During field visits, team members probed the soil to the depth of refusal (maximum of 125 cm). The
217 nature of refusal, interpreted as frozen ground, compact sediment, clasts, rock, or not applicable (N/A; >125 cm), was noted
218 and used to assess permafrost presence or absence. Where the cause of probe refusal was unclear, instantaneous ground
219 temperature measurements were collected using vertically arranged thermistors connected to an Onset Hobo UX120-006M 4-
220 Channel Analog Data Logger (accuracy ± 0.15 °C) (Davis et al., 2020; Holloway and Lewkowicz, 2020; Way et al., 2021b;

Way and Lewkowicz, 2015). Ground temperatures were recorded within the probed hole for a minimum of 10 minutes to allow for thermal equilibration. Frost probing and instantaneous ground temperature measurements were targeted towards locations considered most likely to contain frozen ground and thus mostly occurred on elevated peat-covered microtopography within each WOI.

Low-altitude RPA imagery of prospective peatland permafrost complexes were collected using a DJI Mini 2 microdrone when weather conditions were suitable (i.e., no rain, no fog, low wind). Low-altitude georeferenced video footage was collected using a GoPro Hero9 camera mounted onto a helicopter during a fuel cache mission in northern Labrador in July and August 2021, led by the Torngat Wildlife, Plants, and Fisheries Secretariat. The camera was set to record real-time video (1080 p, 60 fps, wide) at an oblique angle (~45°). The flight altitude was between 90 m and 120 m a.g.l., similar to coastal Nunavik transects performed by Boisson and Allard (2018), and the flight plan between the Goose Bay Airport and the Torngat Mountains National Park was designed to fly over WOIs in coastal locations north of the community of Makkovik (55.0° N) (Supplement Sect. S2). Low-altitude georeferenced aerial images were also collected using handheld digital cameras (Nikon Coolpix W300 or Olympus Tough TG-6) during commercial Air Borealis Twin Otter flight segments between Cartwright and Black Tickle and between Goose Bay, Rigolet, Makkovik, Postville, Hopedale, Natuashish, and Nain. The Twin Otter flights only crossed over WOIs along existing commercial flight routes.

3.4 Compilation of final WOI database

The final WOI database of likely, possible, or unlikely peatland permafrost complexes was developed following the incorporation of the field-validated dataset. WOIs that were classified as likely or possibly to contain peatland permafrost were subject to a final round of review in which the peatland permafrost landforms were identified as palsas, peat plateaus, or both palsas and peat plateaus (mixed).

3.5 Statistical analyses of final WOI database

ANOVA (analysis of variance) and *post hoc* Tukey's HSD (honest significant difference) tests were performed to determine whether the mean latitude, distance from coastline, elevation, MAAT, TDD, and FDD were statistically significantly different between the final classes of likely, possible, and unlikely peatland permafrost complexes. Statistical analyses were performed in R 4.0.3 (R Core Team, 2020).

4 Results

4.1 Peatland permafrost complex identification and review

A total of 2092 unique WOIs, limited to those that prospectively contained the largest (>4 m²) and most visually apparent peatland permafrost landforms, were included in the full inventory. Reviewer agreement was very high (89 %) during the consensus-building review process, with 1116 complexes classified by both reviewers as likely containing peatland

251 permafrost and 750 complexes classified by both reviewers as unlikely to contain peatland permafrost, and only 226 complexes
252 with conflicting classifications of permafrost presence or absence (11 %) (Supplement Sect. S2).

253

254 **4.2 Validation of peatland permafrost complexes**

255 In Summer 2021 and 2022, in-person field visits (n=63 WOIs), RPA visits (n=141 WOIs), helicopter video clips
256 (n=69 WOIs), and Twin Otter images (n=314 WOIs) were combined to evaluate peatland permafrost presence at 531 WOIs,
257 49 of which were cross-validated using multiple methods (Figure 3; Supplement Sect. S2). Previous work from 2017 to 2020,
258 including field visits (n=23 WOIs) and RPA image collection (n=19 WOIs), were also used to validate palsa or peat plateau
259 presence at an additional 19 complexes and peatland permafrost absence at an additional seven complexes (Anderson et al.,
260 2018; Way, 2017). Out of the 557 WOIs evaluated via field and/or imagery validation methods, 311 were interpreted to contain
261 peatland permafrost landforms. Comparison between the validation dataset and the consensus-based inventory resulted in re-
262 classification of 39 of the 226 possible peatland permafrost complexes (17 %) to either likely (n=3) or unlikely (n=36) peatland
263 permafrost complexes.

264




Figure 3. (a) Locations of validated peatland permafrost complexes in coastal Labrador from field-based activities and imagery acquisition using RPA, helicopter, and Twin Otter from 2017 to 2022; Example of peatland complexes containing palsas and/or peat plateaus near (b) Rigolet, (c) Cartwright, (d) Black Tickle-Domino, and (e) Red Bay.

4.3 Peatland permafrost complex inventory

A total of 1119 out of 2092 WOIs were classified as likely containing peatland permafrost landforms, with an additional 187 wetland complexes classified as possibly containing peatland permafrost landforms (Figure 4). The largest clusters of likely and possible peatland permafrost complexes were located between Makkovik (55.0° N) and Black Tickle (53.5° N) (Figure 4; Figure 5A). The likely peatland permafrost complexes were at low elevation (mean elevation of 29 m a.s.l.) (Figure 5C) within 22 km of the coastline (mean distance from coastline of 2.6 km) (Figure 4; Figure 5B). Likely peatland

275 permafrost complexes were distributed from 51.4° N near Blanc-Sablon to 58.6° N in the Torngat Mountains National Park
276 (Figure 4; Figure 5A), with most complexes located in southeastern Labrador (mean latitude of 54.1° N) (Supplement Sect.
277 S3). Comparison against gridded climate products showed that the MAAT at peatland permafrost complexes ranged from -7.5
278 °C to +1.2 °C, with corresponding ranges for FDDs of 1126 degree days to 3471 degree days and TDDs of 733 degree days to
279 1704 degree days (Figure 5D-F). Despite the wide range in MAAT, the majority of the likely peatland permafrost complexes
280 (90 %) were found in locations with MAATs between -2 °C and +1 °C (Figure 5D).




Figure 4. (a) Spatial distribution of inventoried peatland complexes (n=2092) classified as likely containing peatland permafrost landforms (n=1119), possibly containing peatland permafrost landforms (n=187), and unlikely to contain peatland permafrost landforms (n=786); (b) Inset map showing density of peatland permafrost complexes within 20 by 20 km (400 km²) grid cells.




Figure 5. Distribution of wetland complexes likely or possibly containing peatland permafrost landforms by (a) latitude; (b) distance from the coastline; (c) elevation; and (d) mean annual air temperature; (e) mean annual freezing degree days; and (f) mean annual thawing degree days for the 1981 to 2010 climate normal.

290 ANOVA and *post hoc* Tukey's HSD tests revealed that the mean distance from coastline, elevation, MAAT, FDD,
291 and TDD were statistically different between the likely, possible, and unlikely peatland permafrost complexes at the 95 %
292 confidence level. When compared with the complexes that likely contained peatland permafrost, the 187 complexes that
293 possibly contained peatland permafrost were similarly distributed all along the coastline but were skewed further north (mean
294 latitude of 54.5° N) and extended as far as 60.2° N (Supplement Sect. S3). These less certain features were at greater distances
295 from the coastline (mean distance from coast of 7.8 km) and at higher elevations (mean elevation of 66 m a.s.l.). The 786
296 complexes that were unlikely to contain peatland permafrost were well distributed between 51.4° N and 60.2° N (Figure 4) but
297 were located further from the coastline (mean distance from coastline of 10.7 km), at higher elevations (mean elevation of 78
298 m a.s.l.), and at higher MAATs (mean MAAT of -0.5 °C) than the complexes that likely or possibly contained peatland
299 permafrost (Supplement Sect. S3).

300 Likely and possible peatland permafrost complexes were also classified according to the type of peatland permafrost
301 landforms found within the wetland complex (Figure 6). Complexes that were exclusively comprised of palsas accounted for
302 half of the likely and possible peatland permafrost complexes (50 %) and were distributed along the entire study area.
303 Complexes with exclusively peat plateaus were less common (29 %) and were spatially concentrated between ~53° N and ~55°
304 N. The remaining 21 % of the likely and possible peatland permafrost complexes were interpreted to contain a combination of
305 palsas and peat plateaus, but it is possible that many of these complexes contain dissected and heavily degraded peat plateaus
306 that now resemble palsas. Further field-based investigations would be required to differentiate these degradational landforms.




Figure 6. (a) Spatial distribution of likely and possible peatland permafrost complexes classified by peatland permafrost landform type as palsas, peat plateaus, or a mix of both palsas and peat plateaus for coastal Labrador. (b) Inset map showing existing permafrost distribution zones (Heginbottom et al., 1995) for a subsection of coastal Labrador and the location of a new proposed location for the southern limit of the sporadic discontinuous permafrost zone.

314 **5.1 Distribution of peatland permafrost in Labrador**

315 Peatland permafrost complexes in Labrador and adjacent portions of Quebec are abundant in lowlands within 22 km
316 of the Labrador Sea coastline (Figure 5B). A geographic gradient is especially apparent between Rigolet (54.2° N) and Black
317 Tickle (53.5° N), where peat plateaus are abundant along the coast but absent from wetlands farther inland (Figure 4). The
318 higher density of peatland permafrost complexes along the coast could be linked to climatic factors like persistent fog and
319 cloud cover leading to less incoming solar radiation (Way et al., 2018) or thinner and denser snowpacks (Seppälä, 1994; Vallée
320 and Payette, 2007) in the wind-exposed barrens along the coast (Way et al., 2018). Further work should focus on exploring the
321 role of local climate conditions in the formation and persistence of peatland permafrost in coastal Labrador and similar northern
322 coastal locations. Peatland permafrost was found across a large range of MAATs, spanning from -7.5 °C to +1.2 °C. Permafrost
323 persistence at MAATs above +1 °C in southeastern Labrador was previously noted in a field study at five palsa complexes
324 (Way et al., 2018). Peatland permafrost complexes in Labrador were located at higher MAATs than is predicted for other
325 northern coastal regions like northern Finland, Norway, and Sweden (approximately +0.4 °C) (Parviainen and Luoto, 2007).
326 Our results also suggest that the MAAT threshold of +0.2 °C for peatland permafrost areas previously applied to North America
327 (Fewster et al., 2020) is too low for Labrador and adjacent parts of Quebec where peatland permafrost landforms continue to
328 persist due to their relict and resilient nature (Dionne, 1984; Way et al., 2018). Large thermal offsets (up to and often exceeding
329 2.0 °C in southeastern Labrador) (Way and Lewkowicz, 2018) are typical of organic-rich landscapes like peatlands and may
330 promote continued permafrost persistence despite a warming climate (Jorgenson et al., 2010). This may further exacerbate
331 discrepancies between peatland permafrost observations and regional estimates, calling into question the utility of simplified
332 threshold-based approaches when modelling with future climate scenarios. Information on the timing of peatland initiation
333 following deglaciation (Gorham et al., 2007), rates of peat deposition (Tarnocai, 2009; Gorham, 1991), and corresponding peat
334 thicknesses should also be considered in studies of peatland permafrost distribution, as thicker peat deposits may influence
335 permafrost development and support permafrost persistence through a larger thermal offset (Smith and Riseborough, 2002).

336 The regional distribution of fine-grained sediments and local depositional history are expected to play an important
337 role in landscape suitability for peatland permafrost landforms (O'Neill et al., 2019; Seppälä, 1986; Zoltai, 1972). For example,
338 differences in the distribution of palsa versus peat plateau landforms have previously been attributed to varying thicknesses
339 and extents of the underlying sediment, with thicker deposits of frost-susceptible sediments leading to the development of
340 palsas and thinner deposits linked to the development of peat plateaus (Allard and Rousseau, 1999). Differences in sediment
341 grain size may also influence the thickness of the ice lenses, with thicker ice lenses developing in finer sediments, where strong
342 capillarity and cryosuction can be more easily maintained, and thinner ice lenses forming in coarser sediments (Allard and
343 Rousseau, 1999). Further examination of how these variables could influence peatland permafrost formation and persistence
344 in coastal Labrador is challenged by the paucity of information on surficial materials and marine limits along most of the
345 Labrador Sea coastline (Hagedorn, 2022; Occhietti et al., 2011). To date, local marine limits have been identified at some

individual locations and study sites in coastal Labrador (e.g., Bell et al., 2011; Dyke et al., 2005; Occhietti et al., 2011; Vacchi et al., 2018), but widespread mapping of marine sediments has only been completed for a small section of northern coastal Labrador from Goose Bay to Hopedale (Hagedorn, 2022). Based on the information that is currently available, we can qualitatively link the distribution of the largest clusters of peatland permafrost complexes, particularly peat plateau complexes, to locations where post-glacial marine invasions had occurred, such as along the lowland-dominated coastline between Makkovik (55.0° N) and Black Tickle (53.5° N), where frost-susceptible, glaciomarine surficial materials are generally widespread (Fulton, 1989, 1995; Hagedorn, 2022; Occhietti et al., 2011). Meanwhile, fewer peatland permafrost complexes were mapped between Makkovik (55.0° N) and Hopedale (55.5° N), where the elevated topography resulted in limited marine invasions and post-glacial marine deposition along the coast. Significant and coordinated advances in surficial mapping will be required before similar links between peatland permafrost distribution and surficial material type, sediment grain size, and elevation relative to the marine limit can be made for other parts of the coastline.

5.2 Implications for peatland permafrost and permafrost distribution in northeastern Canada

Comparisons between our inventory results and several recent national to global wetland, peatland (Supplement Sect. S1; Supplement Sect. S4), and peatland permafrost distribution products (e.g., Fewster et al., 2020; Hugelius et al., 2020; Olefeldt et al., 2021) (Figure 7) provide compelling evidence that peatland permafrost along the Labrador coast is poorly represented by existing datasets. While differences in scale may explain some of this discrepancy, the general pattern presented in most previous datasets, showing relatively greater peatland permafrost in the continental interior and less along the coast, is directly contradicted by the results of this study. This reversed pattern could reflect inaccurate assumptions on the climate limits of peatland permafrost and/or may reflect the absence of field data from many northern coastal peatland permafrost environments (Borge et al., 2017). Inclusion of physiographic variables, like soil conditions, frost-susceptibility of sediments, and more detailed surficial deposit maps are likely necessary for an improved representation of peatland permafrost in northern coastal regions. Recent work by O'Neill et al. (2019), for example, has demonstrated that segregated ice can be reliably modelled along sections of the Labrador Sea coastline (Figure 7D) by incorporating paleogeographic variables like vegetation cover, surficial geology, and glacial lake and marine limits.




Figure 7. Comparison of inventoried peatland permafrost complexes with peatland permafrost presence and percent coverage as modelled by (a) Fewster et al. (2020); (b) Hugelius et al. (2020); and (c) Olefeldt et al. (2021) and with segregated ice content as modelled by (d) O'Neill et al. (2019).

377 The results of our inventory also suggest that some amendments to existing representations of permafrost distribution
378 may be required for coastal Labrador. For example, the highest density of peatland permafrost complexes (Figure 4B) was
379 found near the community of Black Tickle (53.5° N) on the Island of Ponds (2 palsa complexes, 19 mixed palsa and peat
380 plateau complexes, and 59 peat plateau complexes within 94 km²) (Figure 6), which is currently classified in the isolated
381 patches of permafrost zone on the Permafrost Map of Canada (Heginbottom et al., 1995) and the no permafrost zone on the
382 2000-2016 Northern Hemisphere Permafrost Map (Obu et al., 2019) (Supplement Sect. S5). The identification of large swaths
383 of likely peatland permafrost complexes, including more than 150 peat plateaus, between Cartwright (53.7° N) and Black
384 Tickle (53.5° N) suggest that the physiography-based Permafrost Map of Canada's limit for the sporadic discontinuous zone
385 along the Labrador coast (Heginbottom et al., 1995) (Supplement Sect. S5) could reasonably be extended south by ~110 km
386 from its current position (~53.7° N) (Figure 6B). This southerly extension of the sporadic discontinuous permafrost limit has
387 previously been suggested by Allard and Seguin (1987) and Payette (2001) who indicated that regional vegetation and
388 geomorphology favoured permafrost along much of this coastline (Payette, 1983). Unexpectedly, large clusters of likely
389 peatland permafrost complexes were also identified near the communities of Red Bay (Supplement Sect. S6) and Blanc-Sablon,
390 both of which are considered to be underlain by little to no permafrost (Heginbottom et al., 1995; Obu et al., 2019) (Supplement
391 Sect. S5). A 15 km extension of the southern limit of the Permafrost Map of Canada's isolated patches permafrost zone to
392 include the Blanc-Sablon region would better reflect contemporary permafrost conditions in this area, especially given that
393 permafrost has been previously detected in mineral soils in the community and in surrounding peatlands below the marine
394 limit (Dionne, 1984).

395 **5.3 Challenges and limitations of a point-based inventory of peatland permafrost complexes in coastal Labrador**

396 The most challenging aspects of the inventorying process involved interpreting peatland permafrost presence in
397 isolated WOIs containing small landforms, while in the case of more obvious peatland permafrost features, there were at times
398 difficulties in determining distinct wetland boundaries (Figure 2). However, we believe that these issues were mitigated
399 through the inclusion of multiple mappers, which facilitated the development of a large initial database and reduced the
400 potential omission of prospective WOIs. The consensus-based review process that followed was designed to minimize the
401 inclusion of false positives in the final dataset of 1119 likely peatland permafrost complexes, but we recognize that this
402 conservative approach may have resulted in the exclusion of some complexes. At the northern end of the study area, where
403 other types of periglacial landforms become more common, misclassification of palsas for other elevated periglacial landforms
404 may have contributed to the designation of a higher number of possible peatland permafrost complexes. It is certainly possible
405 that some segregated ice mounds with less than 40 cm of overlying peat (i.e., lithalsas) may have been included in the inventory,
406 particularly near the northern end of the study area where wetlands are less abundant and peat deposits may be thinner
407 (Supplement Sect. S1). This suggests that the definition of peatlands, as wetlands containing at least 40 cm of surface peat
408 (Tarnocai et al., 2011), and its application to palsas and lithalsas, can introduce some ambiguity during inventorying.

409 While other inventorying approaches, including grid-based methods (Ramsdale et al., 2017; Gibson et al., 2020, 2021;
410 Borge et al., 2017), were considered, a point-based inventory was ultimately developed for this study. The implementation of
411 a grid-based approach with delineation of individual landforms for each WOI could have been useful for estimating ground
412 ice content, thermokarst potential, carbon content, and overall permafrost coverage, but the purpose of this study was to
413 generate an initial inventory to guide future research that will facilitate quantitative assessments of peatland permafrost
414 distribution and coverage in these regions. Our field experience in the region suggests that areal delineations of peatland
415 permafrost landforms in coastal Labrador will require extensive validation, and it is unlikely that even experienced permafrost
416 mappers could accurately map the extents of permafrost throughout some complexes without extensive field investigations.
417 Despite the above limitations, our inventory allowed for the incorporation of dedicated, co-located field- and imagery-based
418 validation information. Post-validation adjustments to the inventory, including reclassification of 39 WOIs highlights the
419 importance of ground-truthing in remote sensing- or modelling-based periglacial landform inventories.

420 Owing to a lack of prior field-based assessments of permafrost conditions in Labrador, it was also difficult to
421 independently validate our peatland permafrost inventory results. However, a detailed aerial photograph- and field-based
422 Ecological Land Classification (ELC) survey undertaken in the late 1970s did cover a subset of our study area in southeastern
423 Labrador (Environment Canada, 1999). The ELC identified a total wetland area of 666 km² which was at least partly covered
424 by inventoried peatland permafrost landforms (Figure 8). Comparison with the present study showed that mappers identified
425 peatland permafrost complexes in 23 of the 24 contiguous ELC wetland areas indicated as containing palsas. Examination of
426 the one remaining ELC peatland permafrost-containing wetland area revealed the presence of irregular ponding patterns
427 indicative of thermokarst and elevated landforms that could be peatland permafrost but, due to their small size, would require
428 in situ field visits for validation. Some of the inventoried likely peatland permafrost complexes that were not captured as part
429 of the peatland permafrost areas from the ELC were instead classified in other wetlands, like string bogs, and in raised marine
430 terrain units. Overall, the results of our inventory are in good agreement with the limited previous overlapping field
431 investigations and inventorying efforts from the ELC.

432




Figure 8. (a) Comparison of inventoried peatland permafrost complexes with palsa bog regions identified in the Ecological Land Classification (ELC) survey (Environment Canada, 1999); (b) Inset map showing the extent of the peatland permafrost area that was mapped in the ELC.

6 Conclusions

This study provides the first detailed point inventory of peatland permafrost landforms along the Labrador Sea coastline. Using high-resolution satellite imagery and extensive field- and imagery-based validation efforts, we applied a multi-stage, consensus-based inventorying approach to identify 1119 likely peatland permafrost complexes. Peatland permafrost complexes were primarily found in lowlands on outer coasts, spanning from 51.4° N to 58.6° N, with the largest clusters of complexes occurring ~110 km south of the previously mapped limit of sporadic discontinuous permafrost in northeastern Canada (Heginbottom et al., 1995).

Comparisons between our point inventory results and existing wetland, peatland, and peatland permafrost distribution products reveal major discrepancies between this study and prior estimates of peatland permafrost in Labrador with implications for ground ice content (O'Neill et al., 2019), thermokarst potential (Olefelt et al., 2016), and carbon content (Hugelius et al., 2014). Significant advances in the development of relevant datasets on surficial materials, marine limits, peatland distribution, and peat ages and thicknesses, along with field-based advances in climate monitoring for cloud cover, fog, and snow, are critically needed to better characterize northern coastal regions like Labrador. Our results highlight the importance of field-based validation for periglacial landform mapping and modelling and of considering physiography and geomorphology for accurate representations of peatland permafrost in larger scale spatial products. The significant underestimation of peatland permafrost along the Labrador Sea coastline shown in this study should inform future permafrost, peatland permafrost, and carbon content mapping efforts, infrastructure and climate change adaptation strategy development, and wildlife management considerations for Labrador and other northern coastal regions.

Data availability. Likely and possible peatland permafrost locations from the coastal Labrador peatland permafrost complex inventory are freely available for download from Nordicana D (Wang et al., 2022).

Author contribution. YW and RW designed the study and drafted the manuscript. YW led the raw data collection and the data analysis. RW contributed to raw data collection and data analysis and was the PI for the NSERC Discovery Grant supporting peatland permafrost research activities in Labrador. JB contributed to raw data collection and data analysis. AF and RT contributed to raw data collection. MP coordinated the collection of helicopter video footage. JB, AF, RT, and MP reviewed and contributed edits to the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The authors would like to acknowledge the Nunatsiavut Government (Rodd Laing), the Nunatsiavut Research Centre (Carla Pamak, Michelle Saunders), the NunatuKavut Community Council (Bryn Wood, George Russell Jr., Charlene Kippenhuck), and the Innu Nation (Jonathan Feldgajer, Jack Penashue) for their guidance on research conducted on traditional Inuit and Innu lands. We thank Caitlin Lapalme, George Way, and Amy Norman of Goose Bay; Freeman Butt and Tanya Barney of Red Bay; Jeffrey and Wendy Keefe of Black Tickle; Barbara Mesher of Cartwright; Jane and Jack Shiwak, Tyler and Harvey Palliser, and Sandi and Karl Michelin of Rigolet; and Caroline Nochasak, Liz Pijogge, Carla Pamak, Kayla Wyatt, Frédéric Dwyer-Samuel, and Patricia Johnson-Castle of Nain for their logistical and in-kind support. We acknowledge Caitlin Lapalme, Victoria Colyn, Kayla Wyatt, Frédéric Dwyer-Samuel, Adrian Earle, Patrick Lauriault, Michaela Smitas-Kraas, and Tara Ryan for their valuable field assistance, and we are grateful to Derrick Pottle, Tyler Palliser, Martin Shiwak, Reginald Maggo, Jeremy Ivany, Martin Andersen, Eldred Andersen, Jeffrey Keefe, Gary Bird, Pat Davis, and Anthony Elson for their boat operation and guiding services. Funding for this research was provided by the Natural Sciences and Engineering

478 Research Council of Canada, the Northern Scientific Training Program, and Queen's University. We thank Dr. Antoni
479 Lewkowicz and Dr. Luise Hermanutz for many insightful conversations about permafrost and peatlands in Labrador. The
480 manuscript has also benefitted from helpful comments and suggestions from Michelle Saunders, Dr. H. Brendan O'Neill, Dr.
481 Hanna Lee, and Dr. Christian Hauck, and we thank two anonymous referees and Dr. Steve Kokelj for their constructive
482 reviews.

483 **References**

- 484 Allard, M. and Rousseau, L.: The international structure of a palsa and a peat plateau in the Rivière Boniface region, Québec:
485 Inferences on the formation of ice segregation mounds, *Géographie Phys. Quat.*, 53, 373–387,
486 <https://doi.org/10.7202/004760ar>, 1999.
- 487 Allard, M. and Seguin, M. K.: Le pergélisol au Québec nordique : bilan et perspectives, *Géographie Phys. Quat.*, 41, 141–152,
488 <https://doi.org/10.7202/032671ar>, 1987.
- 489 Anderson, D., Ford, J. D., and Way, R. G.: The impacts of climate and social changes on cloudberry (bakeapple) picking: A
490 case study from southeastern Labrador, *Hum. Ecol.*, 46, 849–863, <https://doi.org/10.1007/s10745-018-0038-3>, 2018.
- 491 Andrews, J. T.: The glacial geomorphology of the northern Nain-Okak section of Labrador, M.Sc. Thesis, McGill University,
492 Montreal, Canada, 301 pp., 1961.
- 493 Banfield, C. E. and Jacobs, J. D.: Regional patterns of temperature and precipitation for Newfoundland and Labrador during
494 the past century, *Can. Geogr. Géographe Can.*, 42, 354–64, 1998.
- 495 Barrette, C., Brown, R., Way, R. G., Mailhot, A., Diaconescu, E. P., Grenier, P., Chaumont, D., Dumont, D., Sévigny, C.,
496 Howell, S., and Senneville, S.: Nunavik and Nunatsiavut regional climate information update, in: Nunavik and Nunatsiavut:
497 From science to policy, an integrated regional impact study (IRIS) of climate change and modernization, second iteration,
498 edited by: Ropars, P., Allard, M., and Lemay, M., ArcticNet Inc., Quebec City, Canada, 62, 2020.
- 499 Beilman, D. W., Vitt, D. H., and Halsey, L. A.: Localized permafrost peatlands in western Canada: Definition, distributions,
500 and degradation, *Arct. Antarct. Alp. Res.*, 33, 70–77, <https://doi.org/10.1080/15230430.2001.12003406>, 2001.
- 501 Bell, T., Putt, M., and Sheldon, T.: Landscape hazard assessment in Nain, Phase I: Inventory of surficial sediment types and
502 infrastructure damage, Final Report to Nunatsiavut Government and Nain Inuit Community Government, 2011.
- 503 Boisson, A. and Allard, M.: Coastal classification of Nunavik and dynamics of the Arctic/Subarctic coastal environments,
504 Vulnerabilities of the Quebec's Arctic Territory in the Context of Climate Change, Kuujuaq, Canada, 2018.
- 505 Borge, A. F., Westermann, S., Solheim, I., and Etzelmüller, B.: Strong degradation of palsas and peat plateaus in northern
506 Norway during the last 60 years, *The Cryosphere*, 11, 1–16, <https://doi.org/10.5194/tc-11-1-2017>, 2017.
- 507 Brown, R. and Lemay, M.: Chapter 2: Climate variability and change in the Canadian Eastern Subarctic IRIS region (Nunavik
508 and Nunatsiavut), in: Nunavik and Nunatsiavut: From science to policy. An Integrated Regional Impact Study (IRIS) of climate
509 change and modernization, ArcticNet Inc., <https://doi.org/10.13140/2.1.1041.7284>, 2012.

510 Brown, R. J. E.: Permafrost investigations in Quebec and Newfoundland (Labrador), National Research Council of Canada,
511 Division of Building Research, Technical Paper 449, <https://doi.org/10.4224/20374659>, 1975.

512 Brown, R. J. E.: Permafrost distribution in the southern part of the discontinuous zone in Québec and Labrador, *Géographie*
513 *Phys. Quat.*, 33, 279–289, <https://doi.org/10.7202/1000364ar>, 1979.

514 Burn, C. R. and Smith, C. A. S.: Observations of the “thermal offset” in near-surface mean annual ground temperatures at
515 several sites near Mayo, Yukon Territory, Canada, *ARCTIC*, 41, 99–104, <https://doi.org/10.14430/arctic1700>, 1988.

516 Coultish, T. L. and Lewkowicz, A. G.: Palsa dynamics in a subarctic mountainous environment, Wolf Creek, Yukon Territory,
517 Canada, *Proc. 8th Int. Conf. Permafr.*, 163–168, 2003.

518 Davis, E., Trant, A., Hermanutz, L., Way, R. G., Lewkowicz, A. G., Siegwart Collier, L., Cuerrier, A., and Whitaker, D.:
519 Plant–environment interactions in the low Arctic Torngat Mountains of Labrador, *Ecosystems*, 24, 1038–1058,
520 <https://doi.org/10.1007/s10021-020-00577-6>, 2020.

521 Dionne, J.-C.: Pales et limite méridionale du pergélisol dans l’hémisphère nord : Le cas de Blanc-Sablon, Québec, *Géographie*
522 *Phys. Quat.*, 38, 165–184, <https://doi.org/10.7202/032550ar>, 1984.

523 Dyke, A. S.: An outline of North American deglaciation with emphasis on central and northern Canada, in: *Developments in*
524 *Quaternary Sciences*, vol. 2, Elsevier, 373–424, [https://doi.org/10.1016/S1571-0866\(04\)80209-4](https://doi.org/10.1016/S1571-0866(04)80209-4), 2004.

525 Dyke, A. S., Dredge, L. A., and Hodgson, D. A.: North American deglacial marine- and lake-limit surfaces, *Géographie Phys.*
526 *Quat.*, 59, 155–185, <https://doi.org/10.7202/014753ar>, 2005.

527 Elias, S. A.: Paleoenvironmental interpretation of Holocene insect fossils from northeastern Labrador, Canada, *Arct. Alp. Res.*,
528 14, 311, <https://doi.org/10.2307/1550794>, 1982.

529 Environment and Climate Change Canada: Canadian Climate Normals,
530 https://climate.weather.gc.ca/climate_normals/index_e.html, 2022.

531 Environment Canada: Audio Tape Transcript of the East-Central Labrador Ecological Land Survey,
532 https://ftp.maps.canada.ca/pub/nrcan_rncan/archive/vector/labrador/, 1999.

533 Esri: World Imagery, 2022.

534 Fewster, R. E., Morris, P. J., Swindles, G. T., Gregoire, L. J., Ivanovic, R. F., Valdes, P. J., and Mullan, D.: Drivers of Holocene
535 palsa distribution in North America, *Quat. Sci. Rev.*, 240, 106337, <https://doi.org/10.1016/j.quascirev.2020.106337>, 2020.

536 Foster, D. R.: The history and pattern of fire in the boreal forest of southeastern Labrador, *Can. J. Bot.*, 61, 2459–2471,
537 <https://doi.org/10.1139/b83-269>, 1983.

538 Foster, D. R. and Glaser, P. H.: The raised bogs of south-eastern Labrador, Canada: Classification, distribution, vegetation and
539 recent dynamics, *J. Ecol.*, 74, 47, <https://doi.org/10.2307/2260348>, 1986.

540 Fulton, R. J. (Ed.): *Quaternary Geology of Canada and Greenland*, Minister of Supply and Services Canada, Ottawa, Canada,
541 846 pp., 1989.

542 Fulton, R. J.: *Surficial Materials of Canada*, “A” Series Map 1880A, Natural Resources Canada, Geological Survey of Canada,
543 <https://doi.org/10.4095/205040>, 1995.

544 Gibson, C., Morse, P. D., Kelly, J. M., Turetsky, M. R., Baltzer, J. L., Gingras-Hill, T., and Kokelj, S. V.: Thermokarst mapping
545 collective: Protocol for organic permafrost terrain and preliminary inventory from the Taiga Plains test area, Northwest
546 Territories, N. W. T. Open Rep., 2020, 29, 2020.

547 Gibson, C., Cottenie, K., Gingras-Hill, T., Kokelj, S. V., Baltzer, J. L., Chasmer, L., and Turetsky, M. R.: Mapping and
548 understanding the vulnerability of northern peatlands to permafrost thaw at scales relevant to community adaptation planning,
549 Environ. Res. Lett., 16, 055022, <https://doi.org/10.1088/1748-9326/abe74b>, 2021.

550 Gorham, E.: Northern peatlands: Role in the carbon cycle and probable responses to climatic warming, Ecol. Appl., 1, 182–
551 195, <https://doi.org/10.2307/1941811>, 1991.

552 Gorham, E., Lehman, C., Dyke, A., Janssens, J., and Dyke, L.: Temporal and spatial aspects of peatland initiation following
553 deglaciation in North America, Quat. Sci. Rev., 26, 300–311, <https://doi.org/10.1016/j.quascirev.2006.08.008>, 2007.

554 Hagedorn, G. W.: Preliminary delineation of marine sediments in east-central Labrador: Parts of NTS map areas 13F, -G, -I, -
555 J, -K, -N AND -O, Gov. Nfld. Labrador Dep. Ind. Energy Technol. Curr. Res., 22, 189–201, 2022.

556 Hare, F. K.: Climate and zonal divisions of the boreal forest formation in eastern Canada, Geogr. Rev., 40, 615,
557 <https://doi.org/10.2307/211106>, 1950.

558 Heginbottom, J. A., Dubreuil, M. A., and Harker, P. T.: Canada, Permafrost, National Atlas of Canada, Natural Resources
559 Canada, Geomatics Canada, <https://doi.org/10.4095/294672>, 1995.

560 Holloway, J. E. and Lewkowicz, A. G.: Half a century of discontinuous permafrost persistence and degradation in western
561 Canada, Permafr. Periglac. Process., 31, 85–96, <https://doi.org/10.1002/ppp.2017>, 2020.

562 Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. A. G., Ping, C.-L., Schirrmeister, L., Grosse, G., Michaelson,
563 G. J., Koven, C. D., O'Donnell, J. A., Elberling, B., Mishra, U., Camill, P., Yu, Z., Palmtag, J., and Kuhry, P.: Estimated stocks
564 of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps, Biogeosciences, 11, 6573–6593,
565 <https://doi.org/10.5194/bg-11-6573-2014>, 2014.

566 Hugelius, G., Loisel, J., Chadburn, S., Jackson, R. B., Jones, M., MacDonald, G., Marushchak, M., Olefeldt, D., Packalen, M.,
567 Siewert, M. B., Treat, C., Turetsky, M., Voigt, C., and Yu, Z.: Large stocks of peatland carbon and nitrogen are vulnerable to
568 permafrost thaw, Proc. Natl. Acad. Sci., 117, 20438–20446, <https://doi.org/10.1073/pnas.1916387117>, 2020.

569 Hustich, I.: Notes on the coniferous forest and tree limit on the east coast of Newfoundland-Labrador, Acta Geogr., 7, 81,
570 1939.

571 International Permafrost Association Terminology Working Group: Multi-language glossary of permafrost and related ground-
572 ice terms, edited by: van Everdingen, R. O., 2005.

573 Ives, J. D.: A proposed history of permafrost development in Labrador-Ungava, Géographie Phys. Quat., 33, 233–244,
574 <https://doi.org/10.7202/1000360ar>, 1979.

575 Jorgenson, M. T., Romanovsky, V., Harden, J., Shur, Y., O'Donnell, J., Schuur, E. A. G., Kanevskiy, M., and Marchenko, S.:
576 Resilience and vulnerability of permafrost to climate change, Can. J. For. Res., 40, 1219–1236, [https://doi.org/10.1139/X10-](https://doi.org/10.1139/X10-060)
577 060, 2010.

578 Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E., Linder, H. P., and
579 Kessler, M.: Climatologies at high resolution for the earth's land surface areas, *Sci. Data*, 4, 170122,
580 <https://doi.org/10.1038/sdata.2017.122>, 2017.

581 Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E., Linder, H. P., and
582 Kessler, M.: Climatologies at high resolution for the earth's land surface areas V2.1, 2021.

583 Karst, A. L. and Turner, N. J.: Local ecological knowledge and importance of bakeapple (*Rubus chamaemorus* L.) in a
584 southeast Labrador Métis community, *Ethnobiol. Lett.*, 2, 6–18, <https://doi.org/10.14237/eb1.2.2011.28>, 2011.

585 Mamet, S. D., Chun, K. P., Kershaw, G. G. L., Loranty, M. M., and Kershaw, G. P.: Recent increases in permafrost thaw rates
586 and areal loss of palsas in the western Northwest Territories, Canada: Non-linear palsa degradation, *Permafr. Periglac. Process.*,
587 28, 619–633, <https://doi.org/10.1002/ppp.1951>, 2017.

588 McLaughlin, J. and Webster, K.: Effects of climate change on peatlands in the far north of Ontario, Canada: A synthesis, *Arct.*
589 *Antarct. Alp. Res.*, 46, 84–102, <https://doi.org/10.1657/1938-4246-46.1.84>, 2014.

590 Natural Resources Canada: Standards and Specifications of the National Topographic Data Base, Edition 3.1, 2005.

591 Norton, C. H., Cuerrier, A., and Hermanutz, L.: People and plants in Nunatsiavut (Labrador, Canada): Examining plants as a
592 foundational aspect of culture in the Subarctic, *Econ. Bot.*, 75, 287–301, <https://doi.org/10.1007/s12231-021-09530-7>, 2021.

593 Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H. H., Dashtseren, A., Delaloye, R., Elberling, B.,
594 Etzelmüller, B., Kholodov, A., Khomutov, A., Kääb, A., Leibman, M. O., Lewkowicz, A. G., Panda, S. K., Romanovsky, V.,
595 Way, R. G., Westergaard-Nielsen, A., Wu, T., Yamkhin, J., and Zou, D.: Northern Hemisphere permafrost map based on
596 TTOP modelling for 2000–2016 at 1 km² scale, *Earth-Sci. Rev.*, 193, 299–316,
597 <https://doi.org/10.1016/j.earscirev.2019.04.023>, 2019.

598 Occhietti, S., Parent, M., Lajeunesse, P., Robert, F., and Govare, É.: Late Pleistocene–early Holocene decay of the Laurentide
599 Ice Sheet in Québec–Labrador, in: *Developments in Quaternary Sciences*, vol. 15, Elsevier, 601–630,
600 <https://doi.org/10.1016/B978-0-444-53447-7.00047-7>, 2011.

601 Olefeldt, D., Goswami, S., Grosse, G., Hayes, D., Hugelius, G., Kuhry, P., McGuire, A. D., Romanovsky, V. E., Sannel, A.
602 B. K., Schuur, E. A. G., and Turetsky, M. R.: Circumpolar distribution and carbon storage of thermokarst landscapes, *Nat.*
603 *Commun.*, 7, 13043, <https://doi.org/10.1038/ncomms13043>, 2016.

604 Olefeldt, D., Hovemyr, M., Kuhn, M. A., Bastviken, D., Bohn, T. J., Connolly, J., Crill, P., Euskirchen, E. S., Finkelstein, S.
605 A., Genet, H., Grosse, G., Harris, L. I., Heffernan, L., Helbig, M., Hugelius, G., Hutchins, R., Juutinen, S., Lara, M. J.,
606 Malhotra, A., Manies, K., McGuire, A. D., Natali, S. M., O'Donnell, J. A., Parmentier, F.-J. W., Räsänen, A., Schädel, C.,
607 Sonnentag, O., Strack, M., Tank, S. E., Treat, C., Varner, R. K., Virtanen, T., Warren, R. K., and Watts, J. D.: The Boreal–
608 Arctic Wetland and Lake Dataset (BAWLD), *Earth Syst. Sci. Data*, 13, 5127–5149, [https://doi.org/10.5194/essd-13-5127-](https://doi.org/10.5194/essd-13-5127-2021)
609 2021, 2021.

610 O'Neill, H. B., Wolfe, S. A., and Duchesne, C.: New ground ice maps for Canada using a paleogeographic modelling approach,
611 *The Cryosphere*, 13, 753–773, <https://doi.org/10.5194/tc-13-753-2019>, 2019.

612 Ou, C., LaRocque, A., Leblon, B., Zhang, Y., Webster, K., and McLaughlin, J.: Modelling and mapping permafrost at high
613 spatial resolution using Landsat and Radarsat-2 images in Northern Ontario, Canada: Part 2 – regional mapping, *Int. J. Remote*
614 *Sens.*, 37, 2751–2779, <https://doi.org/10.1080/01431161.2016.1151574>, 2016.

- 615 Parviainen, M. and Luoto, M.: Climate envelopes of mire complex types in fennoscandia, *Geogr. Ann. Ser. Phys. Geogr.*, 89,
616 137–151, <https://doi.org/10.1111/j.1468-0459.2007.00314.x>, 2007.
- 617 Payette, S.: The forest tundra and present tree-lines of the northern Québec-Labrador peninsula, *Proc. North. Qué. Tree-Line*
618 *Conf.*, 47, 3–23, 1983.
- 619 Payette, S.: Chapitre 9: Les processus et les formes périglaciaires, in: *Écologie des tourbières du Québec-Labrador*, Presses de
620 l'Université Laval, 42, 2001.
- 621 Payette, S.: Accelerated thawing of subarctic peatland permafrost over the last 50 years, *Geophys. Res. Lett.*, 31, L18208,
622 <https://doi.org/10.1029/2004GL020358>, 2004.
- 623 Pironkova, Z.: Mapping palsa and peat plateau changes in the Hudson Bay Lowlands, Canada, using historical aerial
624 photography and high-resolution satellite imagery, *Can. J. Remote Sens.*, 43, 455–467,
625 <https://doi.org/10.1080/07038992.2017.1370366>, 2017.
- 626 R Core Team: R, 2020.
- 627 Ramsdale, J. D., Balme, M. R., Conway, S. J., Gallagher, C., van Gasselt, S. A., Hauber, E., Orgel, C., Séjourné, A., Skinner,
628 J. A., Costard, F., Johnsson, A., Losiak, A., Reiss, D., Swirad, Z. M., Kereszturi, A., Smith, I. B., and Platz, T.: Grid-based
629 mapping: A method for rapidly determining the spatial distributions of small features over very large areas, *Planet. Space Sci.*,
630 140, 49–61, <https://doi.org/10.1016/j.pss.2017.04.002>, 2017.
- 631 Roberts, B. A., Simon, N. P. P., and Deering, K. W.: The forests and woodlands of Labrador, Canada: Ecology, distribution
632 and future management, *Ecol. Res.*, 21, 868–880, <https://doi.org/10.1007/s11284-006-0051-7>, 2006.
- 633 Seguin, M. K. and Dionne, J. C.: Modélisation géophysique et caractérisation thermique du pergélisol dans les palses de Blanc-
634 Sablon, Quebec, *Geol. Surv. Can. Curr. Res.*, E, 207–216, 1992.
- 635 Seppälä, M.: The origin of palsas, *Geogr. Ann.*, 68, 141–147, 1986.
- 636 Seppälä, M.: Snow depth controls palsa growth, *Permafr. Periglac. Process.*, 5, 283–288,
637 <https://doi.org/10.1002/ppp.3430050407>, 1994.
- 638 Smith, J. S.: Shifting sites and shifting sands: A record of prehistoric human/landscape interactions from Porcupine Strand,
639 Labrador, M.Sc. Thesis, Memorial University of Newfoundland, St. John's, Canada, 288 pp., 2003.
- 640 Smith, M. W. and Riseborough, D. W.: Climate and the limits of permafrost: A zonal analysis, *Permafr. Periglac. Process.*, 13,
641 1–15, <https://doi.org/10.1002/ppp.410>, 2002.
- 642 Tarnocai, C.: The impact of climate change on Canadian peatlands, *Can. Water Resour. J.*, 34, 453–466,
643 <https://doi.org/10.4296/cwrj3404453>, 2009.
- 644 Tarnocai, C., Kettles, I. M., and Lacelle, B.: Peatlands of Canada, Natural Resources Canada, Geological Survey of Canada,
645 Open File 6561, <https://doi.org/10.4095/288786>, 2011.
- 646 Thibault, S. and Payette, S.: Recent permafrost degradation in bogs of the James Bay area, northern Quebec, Canada, *Permafr.*
647 *Periglac. Process.*, 20, 383–389, <https://doi.org/10.1002/ppp.660>, 2009.

648 Thie, J.: Distribution and thawing of permafrost in the southern part of the discontinuous permafrost zone in Manitoba,
649 ARCTIC, 27, 189–200, <https://doi.org/10.14430/arctic2873>, 1974.

650 Vacchi, M., Engelhart, S. E., Nikitina, D., Ashe, E. L., Peltier, W. R., Roy, K., Kopp, R. E., and Horton, B. P.: Postglacial
651 relative sea-level histories along the eastern Canadian coastline, *Quat. Sci. Rev.*, 201, 124–146,
652 <https://doi.org/10.1016/j.quascirev.2018.09.043>, 2018.

653 Vallée, S. and Payette, S.: Collapse of permafrost mounds along a subarctic river over the last 100 years (northern Québec),
654 *Geomorphology*, 90, 162–170, <https://doi.org/10.1016/j.geomorph.2007.01.019>, 2007.

655 Wang, Y., Way, R. G., and Beer, J.: Coastal Labrador peatland permafrost inventory, v. 1.0, Nord. D98,
656 <https://doi.org/10.5885/45762XD-1DB498A49B864CFB>, 2022.

657 Way, R. G.: Field and modelling investigations of permafrost conditions in Labrador, northeast Canada, Ph.D. Thesis,
658 University of Ottawa, Ottawa, Canada, 293 pp., 2017.

659 Way, R. G. and Lewkowicz, A. G.: Investigations of discontinuous permafrost in coastal Labrador with DC electrical resistivity
660 tomography, *Proc. 68th Can. Geotech. Conf. 7th Can. Permafr. Conf.*, 8, <https://doi.org/10.13140/RG.2.1.1647.8803>, 2015.

661 Way, R. G. and Lewkowicz, A. G.: Environmental controls on ground temperature and permafrost in Labrador, northeast
662 Canada, *Permafr. Periglac. Process.*, 29, 73–85, <https://doi.org/10.1002/ppp.1972>, 2018.

663 Way, R. G. and Viau, A. E.: Natural and forced air temperature variability in the Labrador region of Canada during the past
664 century, *Theor. Appl. Climatol.*, 121, 413–424, <https://doi.org/10.1007/s00704-014-1248-2>, 2015.

665 Way, R. G., Lewkowicz, A. G., and Bonnaventure, P. P.: Development of moderate-resolution gridded monthly air temperature
666 and degree-day maps for the Labrador-Ungava region of northern Canada: High-resolution air temperature and degree-day
667 maps for Labrador, *Int. J. Climatol.*, 37, 493–508, <https://doi.org/10.1002/joc.4721>, 2017.

668 Way, R. G., Lewkowicz, A. G., and Zhang, Y.: Characteristics and fate of isolated permafrost patches in coastal Labrador,
669 Canada, *The Cryosphere*, 12, 2667–2688, <https://doi.org/10.5194/tc-12-2667-2018>, 2018.

670 Way, R. G., Wang, Y., Bevington, A. R., Bonnaventure, P. P., Burton, J. R., Davis, E., Garibaldi, M. C., Lapalme, C. M.,
671 Tutton, R., and Wehbe, M. A.: Consensus-Based Rock Glacier Inventorying in the Torngat Mountains, Northern Labrador,
672 *Proc. 2021 Reg. Conf. Permafr. 19th Int. Conf. Cold Reg. Eng.*, 130–141, 2021a.

673 Way, R. G., Lewkowicz, A. G., Wang, Y., and McCarney, P.: Permafrost investigations below the marine limit at Nain,
674 Nunatsiavut, Canada, *Proc. 2021 Reg. Conf. Permafr. 19th Int. Conf. Cold Reg. Eng.*, 38–48, 2021b.

675 Wenner, C.-G.: Pollen diagrams from Labrador, *Geogr. Ann.*, 29, 137–374, 1947.

676 Williams, P. J. and Smith, M. W.: *The Frozen Earth: Fundamentals of Geocryology*, Cambridge University Press, 1989.

677 Zoltai, S. C.: Palsas and peat plateaus in central Manitoba and Saskatchewan, *Can. J. For. Res.*, 2, 291–302,
678 <https://doi.org/10.1139/x72-046>, 1972.

679 Zoltai, S. C. and Tarnocai, C.: Perennially frozen peatlands in the western Arctic and Subarctic of Canada, *Can. J. Earth Sci.*,
680 12, 28–43, <https://doi.org/10.1139/e75-004>, 1975.

681 Zuidhoff, F. S. and Kolstrup, E.: Palsa development and associated vegetation in northern Sweden, *Arct. Antarct. Alp. Res.*,
682 37, 49–60, [https://doi.org/10.1657/1523-0430\(2005\)037\[0049:PDAAVI\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2005)037[0049:PDAAVI]2.0.CO;2), 2005.

683