

Variation in bacterial composition, diversity, and activity across different subglacial basal ice types

Shawn M. Doyle¹, Brent C. Christner²

¹Department of Oceanography, Texas A&M University, College Station, TX, 77843, USA

5 ²Department of Microbiology and Cell Science, University of Florida, Gainesville, FL, 32611, USA

Correspondence to: Shawn M. Doyle (shawndoyle@tamu.edu)

Abstract. Glaciers and ice sheets possess ~~layers of~~ basal ice layers characterized by high amounts of entrained debris that can serve as sources of nutrients and organic matter, providing a habitat for microorganisms adapted to the frozen conditions. Basal ice forms through various mechanisms and is classified based on ~~its~~ ice and debris content; however, little is known about variation in microbial composition, diversity, and activity across different basal ice types. We investigated these parameters in 10 four different types of basal ice from a cold-based and temperate glacier and used a meta-analysis to compare our findings with microbiome studies from other frozen environments. We found basal ice environments harbor a diverse range of microbiomes whose composition and activity can vary significantly between basal ice types, even within adjacent facies from the same glacier. In some debris-rich basal ices, elevated ATP concentrations, isotopic gas signatures, and high ~~ratios of~~ 15 ~~amplified sequences for 16S rRNA relative to that for 16S rRNA genes~~ 16S rRNA/rDNA amplicon ratios implicated certain bacterial taxa (e.g., *Paenisporosarcina*, *Desulfocapsa*, *Syntrophus*, *Desulfosporosinus*) as being potentially active, with ice temperature appearing to be an important predictor for the diversity of ~~taxa~~ inferred to be active taxa. Compared to those of other sympagic environments, ~~these the basal ice~~ microbiomes ~~often resembled~~ more closely resemble those found in permafrost or perennial cave ice ~~rather than other glacial ice environments.~~ In contrast, debris-poor basal ices harbored microbiomes 20 more like those found in ~~oligotrophic~~ englacial ice. Collectively, these results suggest that different basal ice types contain distinct microbiomes that are actively structured by ~~the diagenesis~~ physicochemical properties of their habitat.

1. Introduction

At the ice-bed interface of a glacier or ice sheet, basal ice forms when subglacial debris and/or water is incorporated into the base of the ice mass through mechanisms such as freeze-on, regelation, or incorporation of pre-existing ice (Knight, 1997; 25 Souchez et al., 2006~~+~~). Basal ice has a physical structure, chemistry, and microbial composition that is directly affected by its proximity to and interaction with material beneath the glacier (Knight, 1997; Doyle et al., 2013; Montross et al., 2014). When frictional or geothermal heating ~~create temperate conditions at~~ warms the bed, ~~the melting of~~ basal ice can melt and provide inputs of organic matter, nutrients, oxidants, and microorganisms to the subglacial environment (Siegert et al., 2001). ~~Hence, Cycles of~~ basal ice ~~likely represents an important vector for dispersing~~ formation and melting across the bed can thus 30 enable the dispersion of microbes into subglacial ~~aquatic~~ environments ~~that exist within~~ along the glacier's flow path (Achberger et al., 2017).

Basal ice layers contain debris entrained from the underlying substrate that ranges in size from fine-grained clays and silts to coarse sands, gravels, and boulders. Basal ice formation can produce layering and stratification patterns that become deformed by ~~the~~ folding and ~~shearingshear~~ forces associated with glacier flow (Samyn et al., 2008). These processes of entraining and restructuring of subglacial debris combine to produce distinct types of basal ice. This ranges from relatively debris-poor types containing <1% (w/v) sediment to debris-rich types composed almost entirely of sediment with only interstitial ice (Knight, 1997). When compared to the overlying englacial ice (i.e., derived from snowfall), the sediment-rich basal ice contains higher concentrations of potential microbial substrates that include formate, acetate, ferrous iron, and ammonia (Skidmore et al., 2000; Wadham et al., 2004; Tung et al., 2006; Yde et al., 2010). These compounds are hypothesized to be an important source of electron donors for heterotrophic and lithotrophic microorganisms inhabiting subglacial environments, including frozen matrices within the basal ice itself (Montross et al., 2014, 2013).

Although ecosystems associated with sympagic environments were first described nearly 60 years ago (Bunt, 1963), viable microorganisms entrapped within glacial ice have been assumed to exist in a persistent state of metabolic dormancy (Abyzov et al., 1998). This view has been challenged based on evidence from several pioneering studies that revealed anomalies in entrapped gas concentrations within englacial (Sowers, 2001; Campen et al., 2003) and basal ice (Souchez et al., 1995, 1998; Price and Sowers, 2004). The metabolic activity of microorganisms in basal ice is now thought to play an important role in the biogeochemistry of downstream terrestrial, freshwater, and marine environments that receive seasonal discharges of basal ice and subglacial meltwater (Wadham et al., 2010; Barker et al., 2018; Hopwood et al., 2019; Vick-Majors et al., 2020). In fact, transport of basal material is one of the largest sources of labile dissolved organic carbon and iron to marine systems receiving outflow from glaciated watersheds (Hood et al., 2009; Bhatia et al., 2013; Hawkings et al., 2020; [Laufer-Meiser et al., 2021](#)).

~~The~~ [Englacial ice of the](#) world's ice sheets are estimated to contain ~10.2 Pg of organic carbon (Prisco et al., 2008), which represents only ~0.5% of the global soil organic carbon pool (Batjes, 2014). However, the sediments underlying the Greenland and Antarctic ice sheets ~~may contain~~ [are thought to be relatively rich in organic matter \(estimated](#) reservoirs of ~500 and ~21,000 Pg of organic carbon, respectively ~~(~~ Wadham et al., 2008, 2012 ~~)~~ [and the source of material that is incorporated into basal ice when it forms](#). Due to accelerated rates of melt and glacial retreat in the polar regions, the products of organic matter processing by subglacial [and basal ice](#) microorganisms may play an increasingly important role in global carbon cycling. For example, the subglacial production of biogenic methane and its subsequent release to the atmosphere could represent a positive feedback that accelerates further climate warming (Wadham et al., 2008). Indeed, [both subglacial and](#) basal ice sediments have been shown to contain viable methanogens (Skidmore et al., 2000; Boyd et al., 2010) as well as methanotrophs (Dieser et al., 2014; Michaud et al., 2017), indicating the presence of ~~a~~ subglacial methane cycling. Microbial activities in the basal zones of the world's glaciers and ice sheets may thus be an overlooked component of Earth's climate oscillations over geologic timeframes. Despite their potential importance, the logistic challenges associated with accessing these environments have

contributed to the dearth of information on the microbes and biogeochemical processes they perform in the basal zone of large ice masses.

In this study, we used 16S rRNA gene amplicon sequencing to profile the composition and structure of microbial assemblages within basal ice types and from different locations. To explore how these microbial assemblages compare to those in other perennially frozen environments, we performed a meta-analysis that combined our data with that from other studies. Many of these studies sequenced different variable regions of the 16S rRNA gene (e.g., V4, V2, V3V4) that do not fully overlap, hindering the use of traditional OTU-based approaches. To address this, we used the SATé-enabled phylogenetic placement (SEPP) technique to insert exact amplicon sequence variants (ASVs) from these heterogeneous variable regions into the same reference phylogeny (Janssen et al., 2018). This enables a phylogenetically stable comparison of the microbial assemblages inhabiting these different frozen environments. In contrast to the aeolian-sourced microbes entrapped in englacial ice, the microorganisms in basal ice appear to represent “communities” in an ecological sense. ~~We also used~~ These observations, together with data from RNA- and ATP-based approaches to identify potentially active taxa and regions in basal ice types ~~and discuss the, are discussed in relation to their~~ implications ~~of~~ for microbial processes that occur in an important transient environment of the cryosphere.

2. Materials and Methods

2.1 Ice sampling

2.1.1 Taylor Glacier

Taylor Glacier is a cold-based glacier (i.e., ~~has~~ basal ice that is constantly below the pressure-melting point) ~~that is an~~ outlet glacier of the East Antarctic Ice Sheet, ~~terminating that terminates~~ at the western end of Taylor Valley (McMurdo Dry Valleys, Victoria Land, Antarctica). During the austral summers of 2007 and 2009, two tunnels were excavated into the basal zone of the northern lateral margin of the glacier (Fig. S1). Tunnel construction was initiated on fresh ice aprons and extended ~8 m into the glacier interior. At the end of each tunnel, a vertical shaft (2007 expedition) or large chamber (2009 expedition) was excavated to expose a stratigraphic sequence of multiple basal ice types (Montross et al., 2014). Electric chainsaws equipped with unlubricated carbide tipped chains were used to exhume samples of the ice sequence from each basal ice type. Ice samples were sealed in polypropylene flat wrap, packed into insulated containers, and shipped frozen from the field to Louisiana State University where they were subsequently stored in ~~an environmental room with a walk-in freezer temperature set to point of~~ -20 °C. Monitoring ~~of logged data~~ indicated ~~an~~ the actual temperature ~~of -18.9 ± 1.03 °C (n=6800) was~~ maintained during storage: ~~was -18.9 ± 1.03 °C (n=6800).~~

95 2.1.2 Matanuska Glacier

Matanuska Glacier is a 43 km temperate (i.e., englacial and basal ice constantly at the pressure melting point) valley glacier located in south-central Alaska, approximately 140 km northeast of Anchorage. In July 2013, a horizon of debris-rich basal ice exposed near the glacier terminus was sampled using an electric chainsaw (Fig. S2). After collection, the ice samples were packaged in polypropylene flat wrap and shipped frozen to Louisiana State University for storage at -20 °C.

100 2.2 Ice decontamination

Between August and September of 2013, ~~within a temperature controlled environmental room (-5°C),~~ ~2 kg of ice was cut from larger basal ice samples using either a band saw (for the sediment-poor basal ice) or a diamond-bladed masonry saw (sediment-rich basal ice). ~~The surface~~ that was housed within an environmental room (-5°C). The surfaces of freshly cut samples were then cleaned following previous methods (Christner et al., 2005). Briefly, the outermost surface was physically removed by scraping with an autoclaved microtome blade and the newly exposed ice surface was then washed with chilled (-5°C), ~~0.22 µm~~ filtered (0.22 µm pore size) 95% v/v ethanol. Microtome scraping was omitted when processing sediment-rich samples due to the presence of coarse granules and stones embedded in the sediment-layers. Following the ethanol wash, samples were washed extensively with ice-cold, filtered (0.22 µm, pore size), twice-autoclaved deionized water. The samples were washed until a minimum of ~5 mm of the outer surface had been removed based on the ice weight. Large, sterilized surgical forceps were used to handle ~~at~~the ice samples during decontamination and were exchanged frequently during the procedure to prevent carryover contamination. Sample weights before and after decontamination were used to estimate the amount of material removed from each sample during processing, which ~~was typically ranged~~ between 15% ~~and to~~ 25% of the initial mass.

115 The effectiveness of the decontamination procedure was monitored by coating ice samples beforehand with a tracer solution that consisted of three components: [1] ~10⁸ cells mL⁻¹ of *E. coli* JM109 cells transformed with a pETBlue-2 plasmid (Novagen) containing the gene for alcohol dehydrogenase (ADH) from *Drosophila melanogaster*, [2] fluorescein (1000 ppm), and [3] 33% (v/v) glycerol to prevent freezing on the ice at -5°C. The removal of the tracer was monitored after each step of the decontamination procedure (i.e., scraping, ethanol wash, H₂O wash). The fluorescein fluorophore component was monitored by quantifying blue-green fluorescence in 200 µL of rinse-water or meltwater using a BioTurner 20/20n Luminometer (P/N 2030-002) equipped with the blue fluorescence module (BioTurner 2030-041). The pETBlue-2 plasmid component was monitored using a PCR with the following conditions: 1 U of Taq DNA polymerase (5PRIME), 1× MasterTaq buffer, 1× TaqMaster PCR enhancer, 1.5 mM Mg²⁺, 0.2 µM of each primer (TF7: 5'-TAATACGACTCACTATAGGG -3'; pETBlue-DOWN: 5'-GTAAATTGCTAACGCAGTCA-3'), 200 µM dNTPs, and ~100 pg of template DNA. Thirty cycles of PCR were ~~done~~performed with a 30 s denaturation step at 94 °C, 60 s annealing step at 55 °C, and extension at 72 °C for 60 s, followed by a final extension at 72 °C for 10 min. The presence of viable *E. coli* was monitored by spread plating rinse-

water and meltwater on agar-solidified LB media containing ampicillin (100 µg mL⁻¹) followed by overnight incubation at 37 °C. Samples in which any component of the tracer solution was detected in the final meltwater were of a sample, the meltwater was considered contaminated, and the sample was discarded and reprocessed from a new subsample further analysis.

130 2.3 ATP analysis

Immediately after melting at 4 °C, samples were filtered through 0.22 µm pore size polyethersulfone (Supor; Pall) filters using a bench-top vacuum manifold (≤ 20 kPa). Clean ice samples were filtered through a 25 mm diameter filter while banded, solid, and dispersed ice samples were processed through either a 47 mm or 90 mm diameter filter, depending on total sediment content. Identical blank filters that were prepared in parallel served as procedural controls. The filters were cut into small pieces using a sterile scalpel and extracted by vortexing for 10 s in 5 mL of boiling Tris-acetate buffer (TAB; 50mM Tris base adjusted to pH 7.75 with acetic acid). The mixture was then incubated in a boiling water bath for 5 minutes and immediately placed on ice for 30 minutes. Following centrifugation for 5 minutes at 4500×g to remove debris, the supernatant was collected.

The concentration of extracted ATP was measured using a modification of the firefly luciferase-luciferin assay described by (Amato and Christner, 2009). Briefly, 10 µL of extract was added to an equal volume of triethanolamine buffer (TEOA buffer; 200 mM triethanolamine pH 7.6, 2 mM MgCl₂, 240 mM KCL) and incubated at 37°C for 10 minutes. To correct for adenylate adsorption to and luciferase inhibition by the sediment in the sample extracts, two samples were prepared in parallel for each measurement. Following incubation at 37°C, 10 uL of TAB supplemented with 10 µM ATP was added to one of the samples, providing an internal standard for each measurement. All samples were kept on ice during preparation and were equilibrated to room temperature (22 °C) prior to the addition of luciferase and measurement of luminescence.

Weighted linear regression was used to generate calibration curves using concentrations of ATP (in TAB) from 1 pM to 10 µM. The luciferase-luciferin cocktail was prepared fresh for each series of measurements and consisted of 100 U ml⁻¹ luciferase and 0.140 mM D-luciferin. Luminescence was quantified with a 20/20n luminometer (Promega) using auto-injection of 100 µL of the luciferase-luciferin cocktail. Relative luminescence units (RLU) were integrated for three seconds immediately after injection. Sample RLU values were corrected for adsorption and inhibition using the following formula:

$$\frac{1}{\left(\frac{RLU_{\text{sample+std}} - RLU_{\text{sample}}}{RLU_{\text{blank+std}}}\right)} \times RLU_{\text{sample}} = RLU_{\text{corrected}}$$

The corrected RLU values were used together with the standard calibration curve to determine the ATP concentration in the extractant. Final ATP concentrations were corrected for dilution and divided by the mass of the ice sampled. All values reported are the means of replicate samples (n = 3 to 5).

2.4 Nucleic acid extraction

Decontaminated basal ice samples were placed in sterilized containers and melted at 4°C. All samples except one melted in less than 24 hours and were immediately processed. The remaining sample, a large 8.5kg clean ice sample for RNA extraction (see below), took 72 ~~hours~~ to melt. For samples from sediment-rich basal ice (i.e., banded, solid, and dispersed ice), the resulting meltwater was shaken to resuspend ~~any~~the sediment and DNA was extracted from the resulting meltwater-sediment slurry using MoBio PowerMax DNA Isolation kits. To increase DNA extraction yields, two 10g extractions were combined onto a single silica spin column for each sample. For samples from sediment-poor basal ice (i.e., clean ice), meltwater was filtered onto 47 mm, 0.22 µm pore size Supor PES filter membranes (Pall) with ~~≤20-cm-Hg~~ 27 kPa of vacuum. DNA was extracted from the filters using MoBio PowerWater DNA Isolation kits as per the manufacturer's instructions with one exception: the mechanical cell lysis step was performed on a BioSpec MiniBeadbeater-8 for 2 minutes at maximum speed. All DNA samples were stored at -80°C until PCR amplification.

Due to ~~the~~ low cell abundances in the basal ice ($\leq 10^4$ cells g⁻¹; (Doyle et al., 2013)), ~~very~~ large samples (~5 kg of ice; Table S1A) were processed to recover sufficient RNA for reverse transcription PCR (RT-PCR). To facilitate filtration of the large sample sizes, coarse sediment was removed from sediment-rich basal ice samples after melting ~~using~~by low-speed centrifugation (700×g, 10 min, 4°C). Since cells attached to these materials would have been removed during this process, microbes that were specifically associated with coarse sediments and large stones may be underrepresented in the processed samples we analysed. The resulting supernatant containing fine clay and silt-like particles was then concentrated onto 90 mm, 0.22 µm pore size Supor PES filter membranes. Meltwater from clean ice samples did not require ~~this~~ centrifugation step and were directly filtered onto 90mm, 0.22 µm pore size Supor PES filters.

After filtration, the filters were immediately processed for RNA extraction using a modified phenol-chloroform extraction protocol (Dieser et al., 2014). Briefly, filters were sliced into small pieces using a sterile scalpel, immersed in 3 mL of TE buffer (1 mM EDTA, 10 mM Tris; pH 6.3) containing lysozyme (15 mg mL⁻¹), and vortexed for 30 minutes at room temperature. Samples were then homogenized in a BioSpec Mini-Beadbeater for 2 minutes at maximum speed using ~1 g of sterilized 0.1 mm diameter zirconia/silica beads. After bead beating, crude extracts were amended with two volumes of chilled (4°C) denaturing buffer (4M guanidine thiocyanate, 50 mM Tris, 10 mM EDTA, 1% w/v N-lauroylsarcosine, 1% β-mercaptoethanol), the resulting insoluble material was pelleted via centrifugation (4500×g for 5 min at 4°C), and the supernatant was collected. The pellet was washed with 3 mL of chilled denaturing buffer, centrifuged again, and the resulting supernatant pooled with the first. ~~This~~The pooled lysate was then extracted with an equal volume of phenol:chloroform:isoamyl alcohol (25:24:1, pH 6.6), followed by a second extraction with chloroform:isoamyl alcohol (24:1). Nucleic acids were purified from these extracts via an overnight ethanol precipitation with 0.3 M sodium acetate (pH 6.0) ~~Linear~~ that included linear acrylamide (20 µg mL⁻¹; Ambion) ~~was used~~ as a ~~co-precipitant~~ co-precipitant to ~~aid~~ increase

recovery. Genomic DNA was eliminated from RNA extracts by digesting for one hour at 37°C with 4 U of TURBO DNAase
190 (Ambion) followed by purification with MEGAclean Transcription Clean-Up kits (Ambion). Extracted DNA and RNA
concentrations were measured using Quant-it Picogreen and Ribogreen kits (Life Technologies), respectively, per the
manufacturer's instructions.

2.5 16S rRNA amplicon sequencing

From the extracted RNA, complementary DNA (cDNA) was reverse transcribed from ~1 ng of total RNA using SuperScript
195 II reverse transcriptase with the 806R primer (5'-GGACTACVSGGGTATCTAAT-3') following the manufacturer's protocol.
Controls lacking RNA template or reverse transcriptase were analyzed to monitor for contaminating RNA and the persistence
of genomic DNA, respectively, in the samples.

The hyper-variable V4 region of the 16S rRNA gene was PCR amplified from the DNA extracts and cDNA libraries using a
200 barcoded 515F-806R primer pair (Caporaso et al., 2012). Each 50 µL PCR contained ~100 pg of DNA or cDNA, 2.5 U of
AmpliTaq Gold DNA polymerase LD (Invitrogen), 1× Gold Buffer, 2.5 mM MgCl₂, 0.2 µM of each primer, and 0.2 µM
dNTPs. Amplification conditions included [an initial denaturation](#) at 95°C for 9 min, followed by 30 to 40 cycles of denaturation
at 94°C for 1 min, annealing at 50°C for 30 s, and extension at 72°C for 30 s. A final extension was performed at 72°C for 10
min.

205

Following amplification, amplicons were separated by electrophoresis on a 2% agarose gel to assess relative band intensity
and size. The amplicons were quantified with Quant-it fluorometry and pooled at equimolar concentrations. Two extraction
blanks were included in the pooled libraries to serve as procedural controls (Salter et al., 2014; Eisenhofer et al., 2019). Pooled
libraries were purified with a MoBio UltraClean PCR Cleanup Kit and sequenced on the Illumina MiSeq platform (v2
210 chemistry, 2× 250 bp; Georgia Genomics Facility).

2.6 16S rRNA amplicon analysis

Sequence read curation and processing was performed using DADA2 (Callahan et al., 2016) with the following filtering
parameters: ~~maxN=0, trunQ=2, rm.phix=TRUE, maxEE=2, R1 truncLen=240, R2 truncLen=200~~. Error rates for the filtered
and trimmed R1 and R2 reads were calculated using the *learnErrors* function and subsequently used to denoise reads using
215 the DADA2 sample inference algorithm. The denoised reads were merged into amplicon sequence variants (ASV) using a
global ends-free alignment. Paired reads containing mismatches in the overlapping region were removed from the dataset.
Chimeric ASVs were identified and removed ~~by~~ using the consensus method within the *removeBimeraDenovo* function. A
consensus taxonomy for each ASV was then assigned using a naïve Bayesian classifier (Wang et al., 2007) trained on release
[138.1](#) of the SILVA reference database (Quast et al., 2013). The dataset was then subsampled to an even read depth across
220 all samples (59,562 sequences). Any ASV [that](#) classified as a chloroplast, mitochondria, or from which ≥0.1% of its reads were

from a procedural blank were removed (Table S1B). ASVs observed exclusively in the RNA libraries were flagged as a potential ~~contaminant~~[contaminants](#) and removed from the analysis (Table S1C).

2.7 Meta-analyses of microbiomes of perennially frozen ecosystems

Publicly available datasets on the NCBI Sequence Read Archive (SRA) from perennially ~~frozen~~[frozen](#) freshwater ~~frozen~~ environments were identified using a search on August 8th, 2021 with the following syntax: ice metagenome[Organism], glacier metagenome[Organism], and permafrost metagenome[Organism]. Run tables from unambiguously ~~labeled~~[labelled](#) 16S rRNA amplicon datasets were aggregated and studies based on Sanger sequencing were excluded; only datasets based on Illumina, 454, or IonTorrent sequencing were considered in the analysis. Because many studies contained samples from laboratory experiments or field manipulations (e.g., nutrient amendment, thawing, etc.), each dataset was manually curated and cross-referenced with a publication or other data source so that only high quality 16S rRNA surveys from genuine, unmodified permanently frozen environments were included. As part of this effort, datasets were filtered to remove samples from environments that were only seasonally frozen. For example, samples of an active soil layer or transition zone from permafrost datasets were excluded. Additionally, because of their marine origin, we ~~also~~ choose to exclude the wealth of sea ice sample data available and focus the analysis on terrestrial icy environments. Unfortunately, several datasets also had to be excluded because their sequence reads on the SRA were not demultiplexed and barcode information was not provided in the linked metadata or publication. A list of datasets that met these criteria and were included in the meta-analysis are detailed in Table S1D. All runs were downloaded using the SRA Toolbox *fastq-dump* function.

Each dataset was individually processed using DADA2 to infer ASV sequences and produce an ASV table. For each study, the filtering parameters were adjusted to account for variation in fragment length and the 16S rRNA variable region sequenced. Due to higher rates of homopolymer and indel errors with 454 and IonTorrent sequencing, datasets using these platforms used a band size parameter of 32 and homopolymer region (≥ 3 repeated bases) gap cost of -1 during the sample inference algorithm. For Illumina datasets, the band size parameter was set to 16 and gaps in homopolymer regions were treated as normal gaps (gap cost of -8). Nine datasets had to be removed from the meta-analysis after these steps due to the loss of too many sequences during quality control steps within the DADA2 pipeline. Complete parameter details and throughput statistics for each dataset in the final meta-analysis are available in Table S1E and S1F. In datasets containing clearly ~~labeled~~[labelled](#) replicates, read counts from replicate samples were combined and averaged before downstream analyses.

ASV sequences from each study were merged and inserted into the 99% Greengenes reference tree (McDonald et al., 2012) using the SATé-enabled phylogenetic placement (SEPP) technique (Mirarab et al., 2012) within the QIIME2 *fragment-insertion* plugin (Janssen et al., 2018). The merged ASV tables from each study were filtered to remove ASVs rejected from tree insertion by SEPP. [The remaining ASVs which were then classified as described above using the SILVA v138.1 reference](#)

database, and those that could not be taxonomically classified beyond the domain rank (n=2434614) or were classified as mitochondrial sequences (n=419392) were flagged as potential artifacts and removed.

255

The statistical significance of hierarchical clustering patterns between samples in the meta-analysis was performed using Monte Carlo simulations of pairwise unifracs as implemented in the *sigclust2* R package (Kimes et al., 2017). Because parallelized, matrix-based unifracs are incompatible with *sigclust2*, we used the serial, pairwise unifracs calculator from the *scikit-bio* python package. This calculator was embedded into R for the *sigclust2* Monte Carlo simulations using the *reticulate* R package.

260

3. RESULTS

3.1 Description of Basal Ice Samples

Based on the nomenclature of (Hubbard et al., 2009), we identified three distinct types of basal ice in the basal ice profiles sampled from Taylor Glacier: clean ice, banded ice, and solid ice (Table 1; Fig. S4S3; (Montross et al., 2014)). All three types were collected during the 2007 expedition, while only clean ice and banded ice horizons were targeted during the 2009 expedition. The basal ice recovered from Matanuska Glacier was identified as dispersed ice and was the only type observed at the location of sampling.

265

270 **Table 1. Description of basal ice types collected in this study**

Type	Glacier	Temp (°C)	Description ^c
Clean ice	Taylor	-15 ^a	debris-free ice
Banded ice	Taylor	-15 ^a	finely stratified layers of debris and ice, layers range in thickness from a few mm to several cm
Solid ice	Taylor	-15 ^a	composed primarily of frozen debris with only interstitial ice, no visible layering
Dispersed ice	Matanuska	~0 ^b	contains scattered debris aggregated into small clusters, no visible layering

^aMontross *et al.*, 2014. ^bLawson *et al.*, 1998. ^cHubbard *et al.*, 2009

3.2 Biomass in sediment-rich and sediment-poor basal ice

The quantity of DNA and RNA extracted from the basal ices was divided by the sample mass to assess the relative amounts of microbial biomass present within each basal ice type. Approximately 10,000-fold more DNA per gram of basal ice was extractable from the sediment-rich basal ice types (i.e., banded, solid, and dispersed) than in the sediment-poor clean ice (Table 2). Assuming a bacterial genome weight of 2.5 fg, the concentration of DNA extracted from the clean Taylor Glacier basal ice

275

corresponds to ~~560~~[between 600 to 1900](#) genomes per gram of ice, which agrees well with direct cell counts of samples from this ice

facies (Table 2; (Doyle et al., 2013; Montross et al., 2014)). In contrast, genome abundances inferred from DNA concentrations in the sediment-rich Taylor Glacier basal ice samples were two to three orders of magnitude higher than direct cells counts.

280 Among the sediment-rich basal ice types, the temperate dispersed basal ice from Matanuska Glacier contained approximately 6-fold more DNA per gram of ice than the banded and solid basal ices from the cold-based Taylor Glacier (-15°C). The concentration of extractable RNA was very low in all basal ice types (between 0.7 to 3.2 pg RNA g⁻¹; Table 2). The highest RNA concentration—though only marginally—was observed in Matanuska Glacier’s dispersed basal ice. These RNA concentrations were consistent with [trends in the](#) ATP concentration data, a proxy for viable microbial biomass, and in general,
 285 ~~high~~[higher](#) concentrations of ATP g⁻¹ ice were observed in sediment-rich basal ice types versus those with ~~low~~[lower](#) sediment content.

290 **Table 2. Cell counts, biomass estimates, and yield of DNA and RNA extracted from basal ice samples. Sediment-rich basal ice facies are highlighted in gray.**

Sample	Cell Density ^a	Yield (pg g ⁻¹ ice)		Biomass Estimates	
	cells g ⁻¹ ice	DNA	RNA	genomes ^b g ⁻¹ ice	ATP (pg g ⁻¹ ice)
Clean07	0.3 – 0.5 ×10 ³	4.7	0.7	1.9 ×10 ³	<0.01 – 0.04
Clean09	n.d.	1.4	n.d.	0.6 ×10 ³	n.d.
Banded07	1.8 – 18.2 ×10 ³	1.7 ×10 ⁴	2.3	6.8 ×10 ⁶	n.d.
Banded09	n.d.	1.6 ×10 ⁴	0.8	6.5 ×10 ⁶	0.24 – 2.8
Solid	6.7 ×10 ³	1.4 ×10 ⁴	1.2	5.7 ×10 ⁶	<0.01
Dispersed	n.d.	9.9 ×10 ⁴	3.2	4.0 ×10 ⁷	0.06 – 0.14

^aDoyle et al. 2013. ^bNumber of microbial genomes estimated using 2.5 fg of DNA per genome. N.d.= no data

3.3 Composition of microbial assemblages in different ~~types of~~ basal ice [types](#)

A total of 4,227,678 paired-end reads with an average read length of 253 bp were obtained from MiSeq sequencing. After filtration and denoising, 2,883,221 non-chimeric sequences representing 3,310 ASVs remained. Subsampling and culling of ASVs classified as plastids or flagged as potential contaminants produced a final, curated dataset composed of ~~2,005~~[1,987](#)

295 ASVs.





Figure 1. [A] Relative abundances of microbial lineages observed in basal ice based on the use of DNA and RNA templates for detecting 16S rRNA sequences. The graph displays the highest resolution classification for the most abundant taxa and was constructed as follows: First, ASVs were clustered by genus and any having a relative abundance $\geq 20\%$ in at least one of the samples were plotted. This procedure was subsequently repeated with the remaining unplotted ASVs at the rank of family, order, class, and finally phylum. Any remaining rare taxa left after this procedure were not plotted. The bottom plot was scaled to reduce inflated relative abundance biases introduced during curation. [B] Scatter plots of potentially active ASVs within four types of basal ice. Each point represents an individual ASV. The dotted-line indicates a 1:1 ratio between the RNA- and DNA-based 16S rRNA libraries. Gray points denote ASVs observed only in the DNA-based libraries. In the banded basal ice plot, circles denote ASVs from the Banded07 sample while squares denote ASVs from the Banded09 sample. Detected ASVs in each plot denotes the number of ASVs observed in both the RNA- and DNA-based 16S rRNA libraries.

Based on average-neighbor clustering of Bray-Curtis distances, the basal ice samples clustered into three significantly different (AMOVA; $F_{[2,3]}=5.23$, $p=0.017$) groups: (1) Taylor Glacier sediment-poor basal ice, (2) Taylor Glacier sediment-rich basal ice, and (3) Matanuska Glacier dispersed ice. The sediment-poor clean ice samples contained microbial assemblages overwhelmingly composed of *Firmicutes* (68%) and *Proteobacteria* (25%), with smaller proportions of *Bacteroidetes**Bacteroidota* (5%) and *Actinobacteria**Actinobacteriota* (1%). The genus *Tumebacillus* was especially abundant in the clean ice samples (Fig. 1A and S3S4), representing more than half of all taxa in these samples. Assemblages within the banded and solid ice samples were comparatively more diverse than those found in clean ice (Fig. 2), with a wider range of abundant phyla represented: *Actinobacteria* (26%), *Firmicutes* (23%), *Bacteroidetes* (30%), *Actinobacteriota* (27%), *Bacteroidota* (19%), *Caldatribacteriota* (8%), *Proteobacteria* (11%), *Atribacteria* (9%), *Tenericutes* (7%), *Desulfobacterota* (4%), and *Planctomycetes* (*Planctomycetota*) (2%). Notably, ASVs related to the class *Clostridia* composed about 20.21% of the total in these samples. The microbial assemblage within Matanuska's dispersed basal ice was distinct from those observed from Taylor Glacier. Here we observed significant amounts of high abundances of *Nitrospirae**Nitrospirota* (39%) and *Betaproteobacteria**Burkholderiales* (30%), of which several of the most abundant ASVs were related to sulfur and/or iron cycling members of the genera *Thermodesulfovibrio*, *Rhodoferrax*, and *Thiobacillus*. Comparing assemblages from all basal ice samples, only 2419 taxa (~3%) were shared between all four basal ice types (Table S1G). These shared taxa were generally rare members with median relative abundances ranging between 0.03% and 0.31%.






Figure 2. Microbial alpha diversity (Shannon) within various perennally frozen environments.

3.4 Assessing metabolic status of bacterial taxa within different types of basal ice


Of the [2,005,198](#) ASVs observed in this study, we detected 141 (7% of total) within the RNA-derived 16S rRNA libraries (Table S1H), suggesting they were recovered from potentially intact cells rather than eDNA or dead cells. From this group, 80 of the ASVs [had with](#) RNA/DNA abundance log-ratios >1 [and likely could](#) represent metabolically active taxa within the basal ice. [The vast majority \(63 ASVs\)](#) Most of these high-ratio ASVs ([63 ASVs](#)) were observed in the dispersed ice from Matanuska Glacier (Fig. 1B), the warmest and youngest basal ice sampled in this study. Many of the Matanuska Glacier ASVs were members of the *Chloroflexi* and [Deltaproteobacteria](#) *Desulfobacterota*, with some of the largest RNA/DNA abundance log-ratios belonging to *Desulfocapsa*, *Syntrophus*, and members of the family *Anaerolineaceae*. Within the banded basal ice from Taylor Glacier, over half of [the](#) ASVs detected in the RNA-derived libraries were members of the *Firmicutes*. [Among Within](#) this phylum, members of the *Paenisporsarcina*, [Paenibacillaceae](#) *Clostridium*, *Virgibacillus*, *Desulfosporosinus*, *Caldicoprobacter*, and [Virgibacillus](#) *Paenibacillus* had the highest activity potential based on RNA/DNA abundance log-ratios (Table S1H) and represented >90% of the RNA reads in these samples. In comparison, the clean and solid basal ice samples contained the least number of potentially active members—10 ASVs from [only](#) 472 RNA total reads, only 2 of which had RNA/DNA abundance log-ratios >1.

3.5 Meta-Analysis: Microbial populations of various perennially frozen environments

For the meta-analysis, we analyzed 54.6 million 16S rRNA sequences across 256 individual samples from 24 publicly available datasets that represent five different types of freshwater ice environments: (1) supraglacial ice, (2) englacial ice, (3) basal ice, (4) perennial cave ice, and (5) permafrost. After sequence filtering and curation, 7 samples were excluded from meta-analysis due to low read depth. The final meta-analysis dataset contained 25.7 million non-chimeric sequences representing 54,170,577 unique ASVs. Subsampling to equal read depth reduced this to 35,357 ASVs.

We calculated Shannon diversity indices to compare microbial alpha diversity between these frozen environments. Within the basal ice assemblages characterized in this study, alpha diversity was significantly higher in the sediment-rich basal ice types than the low sediment clean ices ($t(4)=3.34$, $p=0.03$) (Fig. 2). Compared to the other frozen environments in the meta-analysis, alpha diversity within Antarctic sediment-rich basal ice samples was most like that observed in basal ice from various coastal glaciers in Svalbard (Tukey HSD mean difference = 0.07, $p=1$) (Perini et al., 2019). The lower alpha diversity values within Taylor Glacier's clean ice were most like those found in englacial ice samples from the GISP2D, NEEM, Guliya Ice Cap, and WAIS Divide ice cores (Tukey HSD mean difference = 0.39, $p=1$) (Miteva et al., 2015, 2016; Price et al., 2015; Zhong et al., 2021).

To parameterize microbial β -diversity between samples, we used hierarchical cluster analyses of both unweighted (UF) and weighted UniFrac (wUF) distances (Fig. 3 and Fig. S5). In the UF cluster analysis (Fig. 3), microbial communities largely clustered by environment type, indicating presence/absence data alone can reasonably discriminate between different parts of






Figure 3. Hierarchical cluster analysis of UF distances between samples using Ward's minimum variance method. The significance of each node was calculated with a Monte Carlo simulation using the 'sigclust2' R package. Family-wise error rate (FWER) was controlled across the dendrogram using a sequential testing procedure starting at the central node. *WISSARD samples were collected via hot-water drilling and thus represent an integrated sample of firn and englacial ice.


the cryosphere. This suggests microbial community composition in permanently frozen environments is strongly influenced by founder effects, possibly due to a population bottleneck effect occurring ~~upon~~ after freezing. In the cluster analysis of wUF, the overall cluster pattern was comparable but many of the branching patterns between samples were more statistically significant, indicating the inclusion of abundance data can help discriminate samples from the same type of environment.

Within the cluster analyses, we also found some samples displayed larger variation and overlap in community structures than others. For example, microbial assemblages within englacial ice were the most similar (UF: 0.47 ± 0.15 ; mean \pm s.d.; $n = 42$). In contrast, perennial cave ice samples (Itcus et al., 2018; Paun et al., 2019) had the highest heterogeneity among samples (UF: 0.85 ± 0.59 ; $n=14$). β -diversity of permafrost microbiomes were intermediate of these endmembers, with relatively tight clustering observed in permafrost from the Storflaket peat bog (northern Sweden) but wider variation in permafrost samples from University Valley (Antarctica) (Goordial et al., 2016; Monteux et al., 2018). The microbial communities within Taylor Glacier's banded and solid basal ice facies clustered closely with perennial cave ices located in both Romania (Itcus et al.,

2018; Paun et al., 2019) and California’s Lava Bed National Monument (O’Connor et al., 2021). ~~Those~~In contrast, those within
 380 the clean basal ice were ~~instead~~ most similar to assemblages observed in englacial ice from the NEEM and GISP2D ice cores
 from Greenland (Miteva et al., 2015, 2016)(Fig. 3).

We used an RDA model to determine how microbial lineages were distributed across the different types of frozen environments
 (Fig. 4, left). The final RDA model explained 50.79% of the variability between samples and effectively separated the samples
 385 into three clusters, echoing patterns observed in the unifracs cluster analysis (Fig. 3). Of the ~~1627~~1745 microbial lineages
 included in the analysis, we identified 3641 with loading vectors greater than the equilibrium contribution (i.e., the proportion
 of variance that would be explained by a random constrained axis), indicating they were associated with one or more of the
 five types of frozen environments. Scalar projections of these lineages onto the centroid factor of each type of frozen
 environment allowed us to estimate how strongly these lineages were associated with each environment (Fig. 4, right).
 390 Unsurprisingly, many supraglacial ice environments harbored comparatively higher proportions of *Cyanobacteria* and plastid-
 harboring taxa than the other frozen environments. Englacial ice environments were enriched with members of the genera
Pseudomonas, *Methylobacterium*, *Acinetobacter*, ~~and~~ *Polaromonas*, *Massilia*, and *Janthinobacterium*. In contrast, basal ice,
 permafrost, and perennial cave ice environments all contained higher membership of *Firmicutes* and
*Actinobacteria*Actinobacteriota related taxa such as *Clostridium*, *Desulfosporosinus*, *Oryzihumus*, and *Cryobacterium*.

395








Figure 4. [Left] RDA triplot of the distribution of [16271745](#) microbial lineages across five different perennally frozen environments. Colored circles represent individual samples. Centroids of each environment type are indicated with an “x”. Plus signs (+) represent microbial lineages. The purple circle denotes the equilibrium contribution of the overall model and delineates lineages which substantially constrained the overall ordination. Those lineages are highlighted in red. [Right] Magnitude of the scalar projections of these [3641](#) lineages onto the centroids for each ice [type](#) approximates that lineage’s association with that frozen environment. Scalar projections smaller than the equilibrium contribution were considered inconclusive and marked gray. Positive and negative associations are black and red, respectively.

405 3.6 Meta-Analysis of early studies based on 16S rRNA clone sequencing

Although 16S rRNA clone libraries lack the read depth to include in our primary meta-analysis, we performed a secondary analysis to compare how many of the microbial taxa observed in several early cryosphere microbiome studies (Yde et al., 2010; Skidmore et al., 2005; Cheng and Foght, 2007; Katayama et al., 2007; Steven et al., 2008) compared to the ‘indicator’ lineages we identified in Fig. 4. To achieve this, we inserted full-length clone sequences into the meta-analysis reference tree using SEPP and taxonomically classified them using Wang's Bayesian inference method with the Silva v. [132138.1](#) database as a reference. Clone sequences were then agglomerated together with meta-analysis ASVs belonging to the same genera.





415 **Figure 5. Distance-based redundancy analysis of Aitchison distances between different perennally frozen environments. Colored circles represent centroids of each study's samples. Black triangles/diamonds represent individual early studies of basal ice and permafrost based on Sanger sequencing of 16S rRNA clone libraries.**

Of the [236287](#) 16S rRNA [unique](#) clone sequences included in this secondary analysis, all but [three](#) five belonged to a taxon that had been observed in the meta-analysis. Despite their extremely low read depth (between 11 and 136 clones), these five studies clustered with similar environments in a distance-based RDA ordination (Fig. 5). Remarkably, of the [3641](#) indicator taxa we identified in the primary meta-analysis, 12 were represented by at least one clone sequence among these five studies. Many of these were among the most abundant sequences observed in these clone libraries (Fig. S6).

420

4. DISCUSSION

4.1 Microbial communities vary in size and composition across basal ice types

Of the four different types of basal ice analyzed in this study, we found that those containing large quantities of sediment or debris contained higher quantities of extractable nucleic acids and microbial abundances than those containing low sediment content. This is consistent with previous microbiological investigations of the GISP2 ice core that found higher cell concentrations in the deepest sections of the core, where clay particles were embedded in the basal ice (Tung et al., 2006). In addition to higher numbers of total cells and biomass, microbial diversity is also notably higher in the sediment-rich basal ice facies relative to clean basal ice or englacial ice, implicating subglacial debris as the primary source of microorganisms in the basal ice.

Our estimates of bacterial cell abundance based on the yield of extracted DNA ($\sim 10^6$ cells g^{-1} ice) were approximately 100-fold higher than those based on direct microscopic counts ($\sim 10^4$ cells g^{-1} ice) in the sediment-rich banded and solid basal ice. However, in the clean basal ice samples, these estimates were highly comparable (Table 2). This difference likely reflects the much greater efficiency of DNA extraction versus that of cell extraction. It may also be related to the increased prevalence of eukaryotic genomes or environmental DNA that is associated with sediment-rich basal ices (Perini et al., 2019; Sonjak et al., 2006; Fraser et al., 2018). Alternatively, Taylor Glacier's debris-rich basal ice may contain significant amounts of necromass. This would be consistent with previous studies that have found some deep, debris-rich horizons of basal ice, such as those of the Greenland ice sheet, contain ancient DNA from buried organisms (Willerslev et al., 2007).

We found microbial assemblages within basal ices of the cold-based (-15°C) Taylor Glacier to contain high abundances of *Firmicutes* and *Actinobacteria* relative to the other frozen habitats examined. These Gram-positive phyla are commonly observed as the dominant taxa in a wide range of frozen environments including ground ice, permafrost, polar saline springs, cryopegs, and sea ice (Steven et al., 2008; Kochkina et al., 2001; Steven et al., 2007; Perreault et al., 2008; Lacelle et al., 2011; Boetius et al., 2015; Spirina et al., 2017). One possible explanation for these high abundances is that the physiochemical stresses faced by cells under frozen conditions (Doyle et al., 2012) favor the prevalence of microorganisms that can form highly resistant endospores (*Firmicutes*) or spores (*Actinobacteria*) (Nicholson et al., 2000; Ensign, 1978). Indeed, the ability to enter a metabolically dormant state and form an environmentally resistant spore likely represents a robust survival strategy for microorganisms that become entrapped inside ice for extended timeframes (Filippidou et al., 2016). However, multiple lines of evidence indicate this hypothesis does not adequately explain the abundance of these spore-forming taxa in our basal ice samples. First, ASVs related to known endospore-forming genera such as *Paenispodosarcina*, *Virgibacillus*, *Clostridium*, and *Desulfosporosinus* were abundant in all our RNA libraries. Because the rRNA content of endospores decreases relatively quickly after sporulation—on the order of days to weeks (Segev et al., 2012; Korza et al., 2016)—their detection in ~~$>10,000$ -year-old~~ ancient basal ice implies these taxa existed as vegetative cells and not

455 as endospores within the ice. Second, populations of *Paenisporosarcina* sp. isolated from the Taylor Glacier banded ice ~~did~~
not sporulate ~~after being when~~ refrozen in basal meltwater at in situ temperatures (-15°C), ~~but~~ and instead ~~continuously~~
~~incorporate~~ ~~incorporated~~ radiolabeled DNA and protein precursors into macromolecules over 70 days of frozen incubation
(Doyle et al., 2013). Third, we found pasteurization of banded basal ice meltwater dramatically reduced isolate cultivability,
indicating these isolates were not ~~being~~ recovered from heat-resistant endospores. (Doyle et al., 2013). Finally, 16S rRNA
460 surveys of permafrost that have selectively depleted DNA from dead or vegetative cells have found many endospore-forming
taxa, especially members of the *Clostridia*, were more likely to persist as vegetative cells rather than endospores in the
permafrost (Burkert et al., 2019). Collectively, these results suggest the high abundances of *Firmicutes* and
Actinobacteria Actinobacteriota in Taylor Glacier's basal ice may not only be related to their environmental durability while
dormant, but also due to properties associated with their vegetative cycle that promote physiological activities within frozen
465 matrices (Doyle et al., 2013).

4.2 Young and warm versus old and cold basal ice

Due to the chemical and biological lability of RNA, paired RNA/DNA 16S rRNA amplicon libraries provide a useful proxy
for discerning viable and metabolically active microorganisms ~~in a community~~ from those that are dead or dormant: in a
community. However, there are limitations to this ~~analysis to be mindful of~~ approach. Variations in sequencing depth, PCR
470 primer bias, differences in DNA/RNA extraction, and inconsistent relationships between rRNA content and activity for
different taxa are all sources of bias that can ~~introduce biases in 16S~~ affect observed ratios. As such, direct comparisons of 16S
rRNA/rDNA ratios as reliable indicators of the relative metabolic activity between different ASVs should be ~~performed~~ viewed
with caution (Blazewicz et al., 2013). This ~~may especially issue could~~ be ~~the case~~ exacerbated in frozen environments, where
RNA may be more stable in dormant cells. As such, we used a conservative approach that focuses primarily on the number of
475 detected taxa in each basal ice type, especially those with unambiguously high RNA/DNA abundance log-ratios, ~~which that~~
are difficult to explain in ancient ice samples without invoking metabolic activity.

On average, only about 2% of the total ASVs in each sample were also detected in the paired RNA library. This low proportion
of overlap suggests that only a small fraction of the microorganisms preserved in the basal ice are potentially metabolically
480 active within the ice matrix, ~~although~~ However, it is possible some taxa may not have been ~~undetected~~ detected due to primer
bias during reverse transcription. We took care to extract nucleic acids from samples as soon as melting was complete (typically
between 12 and 24 h) ~~to limit~~, limiting any impacts the melting process had on ~~which ASVs we detected the microbes released~~
during the melting. Although one sample took ~~much longer~~ 72 h to melt (i.e. Clean07 for RNA), we ~~ultimately~~ did not find
~~much good~~ evidence for active ASVs in this sample, suggesting the effects of ~~the extended~~ melting time ~~(72 h)~~ for this sample
485 ~~were minimal~~ minimally affected our results. The dispersed basal ice from Matanuska Glacier contained a much larger number
of high-ratio ASVs (i.e. rRNA/rDNA abundance log-ratios >1) than any of the basal ice facies recovered from Taylor Glacier

(Fig. 1 and S3S4). One plausible explanation for this observation is the substantial difference in underlying geology, age, and temperature of basal ice between the two glaciers.

490 Assuming similar starting inputs of microbes, older ice would be expected to harbor smaller numbers of potentially active populations than younger ice due to microbial attrition over time (Doyle et al., 2012). Analysis of the δD and $\delta^{18}O$ composition of meltwater from Taylor Glacier surface ice samples estimate ages of 11,500 to 65,000 years BP (Aciego et al., 2007), while recent radiometric ^{81}Kr dating efforts have yielded age estimates near the glacial terminus of 123,500 years BP (Buizert et al., 2014). In comparison, ~~measurements~~ measurement of oxygen isotopes in Matanuska Glacier's dispersed basal ice layer
495 indicated the basal ice originated in the accumulation area (Lawson and Kulla, 1978). Based on the velocity of Matanuska Glacier ($\sim 110 \text{ m yr}^{-1}$), this would represent an approximate time of 250 years between deposition and terminal ablation. Trace amounts of anthropogenic tritium produced by atmospheric thermonuclear weapon tests in the 1950s and 1960s have also been detected in Matanuska Glacier's basal ice zone (Strasser et al., 1996), indicating portions of this ice were formed even more recently by freeze-on of surface-derived meltwater (Lawson, 1979). This indicates implies that the dispersed ice we sampled
500 from the Matanuska Glacier is substantially younger than any of the basal ice types collected from Taylor Glacier, ~~with the latter probably being between 100 to 500 times older than the former.~~

In addition to age of the basal ice, the large difference in ice temperature between a temperate and cold-based glacier would also affect the potential for metabolic activity in the ice habitat (Price and Sowers, 2004; Price, 2000). Taylor Glacier is a cold-
505 based glacier with a basal ice temperature between $-15^{\circ}C$ and $-17^{\circ}C$ based on independent measurements (Montross et al., 2014; Samyn et al., 2008). In contrast, the Matanuska Glacier is temperate, and the basal ice zone is at the pressure melting point of $0^{\circ}C$ (Lawson et al., 1998). As such, cells present in the liquid habitat at grain boundaries (Price, 2000) in Taylor Glacier's basal ice are under considerably higher physiochemical stress than ice near the melting point. For example, the predicted ionic strength of the briny liquid at ice crystal boundaries—where the microorganisms are located—is nearly 4.6M
510 at $-15^{\circ}C$. ComparativelyIn comparison, this is 11.5-fold higher than estimates for unfrozen water in glacial ice at $-1^{\circ}C$ (Doyle et al., 2012). Hence, the wider diversity and increased number of ASVs with a high potential for metabolic activity in the dispersed basal ice may ~~reflect~~ not only the be related to sample age, but the more favorable conditions for metabolism in temperate versus cold-based glacial ice matrices.

4.3 Debris-rich basal ice types are hot spots for microbial activity

515 Multiple studies have reported observations of very low concentrations of oxygen (Souchez et al., 1995) and large excesses of CH_4 in debris-rich basal ice from the Greenland and Antarctica ice sheets (Tung et al., 2006; Wadham et al., 2012; Rhodes et al., 2013; Lamarche-Gagnon et al., 2019; Lee et al., 2020; Stibal et al., 2012). Mirroring these findings, an analysis of gasses entrapped within Taylor Glacier banded basal ice found O_2 was depleted to as low as 4% in horizons where CO_2 concentrations were concurrently enriched as high as 20,000-fold relative to atmospheric concentrations (Montross et al., 2014). Isotopic

520 analysis of $\delta^{13}\text{C}$ -CO₂ revealed this CO₂ was isotopically depleted (-24‰), and therefore, was likely produced by microbial
respiration of organic matter (Montross, 2012). In this study, among all the RNA-based 16S rRNA libraries we sequenced
from Taylor Glacier basal ice, we found the largest ~~readsread~~ counts and largest number of potentially active ASVs in these
same banded basal ice horizons. Together with elevated ATP concentrations, these observations collectively suggest Taylor
Glacier's banded basal ice horizons harbor metabolically active microbiomes that alter entrapped gas compositions by actively
525 metabolizing organic matter to CO₂ at -15°C within the ice matrix.

Alternatively, ~~thesethe~~ trends in gas data could be the result of microbial activity that had occurred before the basal ice formed.
One way to explore this possibility is by considering the total gas content of the ice. If the gases were dissolved in ~~the~~ water
that froze onto the base of the glacier, then total gas volumes as low as 0.02 to 0.08 cc/g would be expected (Cuffey et al.,
530 2000; Sleewaegen et al., 2003). In solid facies of the Taylor Glacier basal ice profile, the lower total gas volumes observed
were indeed low (0.01 to 0.07 cc/g; (Montross et al., 2014)). Hence, although ~~we did not see muchthere is little~~ evidence for
microbial activity in the solid basal ice, we cannot exclude ~~largescalethat large-scale~~ melting/refreezing processes ~~have~~
occurred during its formation. In contrast, ~~in~~ the banded basal ice—where isotopic, ATP, and RNA data provided strong
evidence for microbial activity—~~thehas~~ total gas volumes ~~were consistent withsimilar to~~ those observed in the clean ice (0.09
535 to 0.11 cc/g; (Montross et al., 2014)). ~~These volumes are more similar to those typically seen in)) and reported for~~ englacial
ice of meteoric origin (0.1 cc/g; (Paterson, 1994)). In summary, large-scale melting/refreezing cannot explain the gas data
observed across all facies, and in the banded ice, in situ microbial respiration is the ~~onlybest~~ explanation consistent with its
total gas content.

540 ~~This~~The differences in metabolic activity ~~inferred in different basal ice types~~ may simply be a consequence of substrate
availability for microbial consumption. Previous geochemical analysis of Taylor Glacier's basal ice found dissolved organic
carbon (DOC) concentrations are much higher in the debris-rich basal ice types (~63 mg L⁻¹) than in the debris-poor types
(~0.8 mg L⁻¹; (Montross, 2012; Montross et al., 2014))))). However, our comparison of multiple basal ice types suggests there
are likely additional, unidentified factors besides sediment content controlling whether basal ice harbors an active microbiome.
545 For example, although the banded and solid basal ice samples harbored very similar microbial communities in our DNA-based
libraries, we found very little evidence for active ASVs within the solid basal ice. This suggests—at least in polar glaciers
containing basal ice horizons well below 0°C—elevated debris and/or DOC ~~alone~~ are not ~~alone~~ sufficient for supporting
metabolic activity in the ice.

550 In many of the debris-rich basal ice types ~~we~~ investigated in this study, we observed an enrichment of putatively anaerobic
taxa (e.g., *Clostridia*) within the same samples where O₂ concentrations were depleted relative to atmospheric values. One
possible explanation is that the subglacial debris within these horizons were enriched with these taxa before being entrained
into the basal ice. Alternatively, these patterns represent ecological shifts in microbial community composition that occur

inside the ice matrix as oxygen becomes depleted. However, there is an important nuance to how these shifts probably occur. 555 Although there is ample evidence that some microbial taxa can remain metabolically active while frozen (Doyle et al., 2013; Panikov et al., 2006; Bakermans and Skidmore, 2011; Tuorto et al., 2014; Segura et al., 2017), there is comparatively little concrete evidence that microorganisms can physically grow and divide while frozen inside ice. This suggests that reproduction ~~does not occur, or if it does, is~~ may be imperceptibly slow under the conditions within an ice matrix. ~~Hence~~ Instead of a genuine enrichment or outgrowth of putative anaerobes, the increased relative abundance of many putative anaerobic taxa that we 560 observed in ~~many of the~~ sediment rich basal ice samples ~~likely represents a decline~~ could represent a die-off of strictly aerobic taxa after oxygen has become depleted ~~rather than a genuine enrichment or outgrowth of putative anaerobes~~. This hypothesis may explain why enrichment cultures inoculated with 100,000-year-old Greenland basal ice became turbid substantially faster (2 to 3 weeks versus several months) when incubated anaerobically versus aerobically (Sheridan et al., 2003).

4.4. Basal ice diagenesis likely controls microbiome composition and structure

565 ~~The~~ Phylogenetically, the microbiomes of cold-based, debris-rich basal ice horizons such as the banded and solid basal ice from Taylor Glacier were ~~phylogenetically~~ most like those in perennial cave ice or permafrost. These similarities are likely a reflection of the comparable physiochemical conditions found within these environments. All three are characterized by relatively high concentrations of debris, constant subzero temperatures, and perhaps—as suggested by the co-enrichment of several putative anaerobic taxa (e.g., *Clostridium*, *Desulfosporosinus*) in some samples—depleted or decreased oxygen 570 concentrations. In contrast, clean basal ice appears to harbor microbiomes most like those found in englacial ice. This is consistent with the notion that clean basal ice and englacial ice are diagenetically very similar. Both contain very little to no debris and are typically formed from the accumulation and compression of snow into glacial ice. In fact, except in unique cases where new clean basal ice is formed by the accretion of subglacial lake water onto the base of a glacier, the only real difference between clean basal and englacial ice may be location. In other words, in many cases clean basal ice is simply deep englacial 575 ice. Collectively, these findings suggest the microbiomes of basal ice environments are not simply preserved assemblages, but communities ~~but rather that~~ are actively structured by the diagenesis and physiochemical nature of their habitat.

4.5 Meta-analysis limitations

Comparing 16S rRNA amplicon datasets from different studies is hindered by inconsistencies in the sequencing platform used and the variable region of the 16S rRNA gene that was amplified for sequencing. Using SEPP, heterogeneous ASVs from 580 different studies are inserted into the same reference tree. This creates stable phylogenetic placements that enable integration and meta-analysis of amplicon data from multiple variable regions of the 16S rRNA gene (Janssen et al., 2018). Nevertheless, it is important to note the limitations of this approach. First, integrating previously published 16S rRNA amplicon datasets based on DGGE profiling remains difficult as they lack sufficient read depth or taxonomic resolution to make meaningful comparisons with more modern datasets. As a result, our meta-analysis excluded many of the pioneering works that 585 investigated the microbial ecology of glacial ice. Second, there are potential biases in our meta-analysis due to how we

classified some icy environments. For example, while it is clear supraglacial ice exists on the surface, there is no universal definition of a depth wherein glacial ice should suddenly be considered ‘englacial’. Indeed, supraglacial and englacial ice would probably be more accurately portrayed as opposite ends of a vertical gradient through the body of a glacier rather than discrete categories. This may explain some of the overlap in microbiome composition and structure we observed between
590 supraglacial and englacial environments (Figs. 4 and 5).

Another potential source of bias in the meta-analysis ~~to be cognizant of~~ is how some ice samples ~~are~~were collected, specifically basal ice. Due to the logistical difficulty of accessing the basal ice zone of a glacier, most samples of basal ice included in the meta-analysis were not collected from inside an excavated tunnel like our samples at Taylor Glacier. Instead, most of these
595 samples (including our sample of Matanuska’s dispersed basal ice~~);~~) were collected from easily accessible horizons of basal ice that were exposed at the glacial margin. These samples have been exposed to sunlight and seasonal temperature fluctuations that are not found in the glacial interior. It is difficult to accurately predict how these conditions would affect the basal ice microbiomes. However, it might explain why, for example, phototrophic taxa (e.g., *Cyanobacteria*) are observed in several of the basal ice samples from coastal Svalbard glaciers.

600 **4.6 Conclusions and Implications**

Our findings show that certain microbes do not persist in a state of dormancy while frozen, and coupled with other biogeochemical data (Doyle et al., 2013; Montross et al., 2014), indicate that the sediment-rich basal ice horizons of glaciers harbor microbial communities that actively conduct biogeochemical cycling at subzero temperatures. Although we have limited observations, our data further implies that a larger diversity of bacteria remain metabolically active in temperate basal
605 ice found beneath nonpolar glaciers and ice sheet interiors (Lawson et al., 1998; Bell et al., 2011). Given that the basal temperature of the Antarctic and Greenland ice sheets are near or at the pressure-melting point (Gow et al., 1968), this raises the intriguing possibility that subglacial regions of the cryosphere may be more biogeochemically active than previously thought.

610 Microorganisms in the basal ice of cold-based glaciers may be important biogeochemical actors. Although rates of microbial activity in these environments are extremely low ~~and much lower than~~compared to those of warmer environments (Doyle et al., 2013; Amato and Christner, 2009; Panikov et al., 2006; Christner, 2002; Amato et al., 2010), this may be offset by the long residence time of basal ice. Microbial processing in basal ice could have important implications to material that is transported to and melts at the margin, representing an important source of dissolved organic carbon, nutrients, and trace metals to marine
615 ecosystems (Vick-Majors et al., 2020; Hawkins et al., 2020; Rignot and Jacobs, 2002; Jung et al., 2019).

Finally, our meta-analysis demonstrates that basal ice environments harbor a diverse range of microbial communities ~~which~~
~~ean~~that resemble those found in a wide range of other icy environments. This is likely a reflection of the many different types

of basal ice, which vary considerable in both physical and chemical characteristics (46)Hubbard et al., 2009). Debris-rich
620 basal ice types, with comparatively higher concentrations of potential nutrients and redox substrates for microbial metabolism
resemble other debris-rich icy environments such as permafrost and/or perennial cave ice. Likewise, clean basal ice microbial
communities are more like those found in the oligotrophic, debris-free englacial ice that makes up the bulk of a glacier's mass.
Broadly, these findings indicate basal ice diagenesis plays a major role in microbiome composition and structure.

Data availability

625 Sequence read data generated in this study is available on NCBI SRA under BioProject PRJNA282540.

Author contribution

SD and BCC designed the research, SD performed the research and analyses, SD and BCC wrote the paper.

Competing interests

The authors declare that they have no conflict of interest.

630 **Acknowledgements**

This study was funded by an Office of Polar Programs grant (ANT-0636828) from the National Science Foundation. We thank
Amanda Achberger, Pierre Amato, Tim Brox, Lindsay Knippenberg, Scott Montross, and Mark Skidmore for multiple
discussions and as vital partners in tunnel excavation and sampling in Antarctica. We also thank the staff of McMurdo station
and Petroleum Helicopters International for logistical support in ~~the field~~[Antarctica](#).

635 **References**

- Abyzov, S., Mitskevich, I., and Poglazova, M.: The Microflora of Deep Horizons of the Central Antarctic Ice Sheet, 67,
547–555, 1998.
- Achberger, A. M., Michaud, A. B., Vick-Majors, T. J., Christner, B. C., Skidmore, M. L., Priscu, J. C., and Tranter, M.:
Microbiology of subglacial environments, in: Psychrophiles: from biodiversity to biotechnology, Springer, 83–110, 2017.
- 640 Aciego, S. M., Cuffey, K. M., Kavanaugh, J. L., Morse, D. L., and Severinghaus, J. P.: Pleistocene ice and paleo-strain rates
at Taylor Glacier, Antarctica, 68, 303–313, <https://doi.org/10.1016/j.yqres.2007.07.013>, 2007.
- Amato, P. and Christner, B. C.: Energy Metabolism Response to Low-Temperature and Frozen Conditions in Psychrobacter
cryohalolentis, 75, 711–718, <https://doi.org/10.1128/AEM.02193-08>, 2009.

- Amato, P., Doyle, S. M., Battista, J. R., and Christner, B. C.: Implications of Subzero Metabolic Activity on Long-Term Microbial Survival in Terrestrial and Extraterrestrial Permafrost, 10, 789–798, <https://doi.org/10.1089/ast.2010.0477>, 2010.
- Bakermans, C. and Skidmore, M. L.: Microbial Metabolism in Ice and Brine at -5°C : Microbial metabolism in ice and brine at -5°C , 13, 2269–2278, <https://doi.org/10.1111/j.1462-2920.2011.02485.x>, 2011.
- Barker, J. D., Grottooli, A. G., and Lyons, W. B.: Stable isotope evidence for the biogeochemical transformation of ancient organic matter beneath Suess Glacier, Antarctica, 50, e1448643, <https://doi.org/10.1080/15230430.2018.1448643>, 2018.
- 650 Batjes, N. H.: Total carbon and nitrogen in the soils of the world, 65, 10–21, https://doi.org/10.1111/ejss.12114_2, 2014.
- Bell, R. E., Ferraccioli, F., Creyts, T. T., Braaten, D., Corr, H., Das, I., Damaske, D., Frearson, N., Jordan, T., Rose, K., Studinger, M., and Wolovick, M.: Widespread Persistent Thickening of the East Antarctic Ice Sheet by Freezing from the Base, 331, 1592–1595, <https://doi.org/10.1126/science.1200109>, 2011.
- 655 Bhatia, M. P., Das, S. B., Xu, L., Charette, M. A., Wadham, J. L., and Kujawinski, E. B.: Organic carbon export from the Greenland ice sheet, *Geochimica et Cosmochimica Acta*, 109, 329–344, <https://doi.org/10.1016/j.gca.2013.02.006>, 2013.
- Blazewicz, S. J., Barnard, R. L., Daly, R. A., and Firestone, M. K.: Evaluating rRNA as an indicator of microbial activity in environmental communities: limitations and uses, 7, 2061–2068, <https://doi.org/10.1038/ismej.2013.102>, 2013.
- Boetius, A., Anesio, A. M., Deming, J. W., Mikucki, J. A., and Rapp, J. Z.: Microbial ecology of the cryosphere: sea ice and glacial habitats, 13, 677–690, <https://doi.org/10.1038/nrmicro3522>, 2015.
- 660 Boyd, E. S., Skidmore, M., Mitchell, A. C., Bakermans, C., and Peters, J. W.: Methanogenesis in subglacial sediments: Subglacial methanogenesis, 2, 685–692, <https://doi.org/10.1111/j.1758-2229.2010.00162.x>, 2010.
- Buizert, C., Bagginstos, D., Jiang, W., Purtschert, R., Petrenko, V. V., Lu, Z.-T., Müller, P., Kuhl, T., Lee, J., Severinghaus, J. P., and others: Radiometric ^{81}Kr dating identifies 120,000-year-old ice at Taylor Glacier, Antarctica, 111, 6876–6881, 2014.
- 665 Bunt, J.: Microbiology of Antarctic sea-ice: diatoms of Antarctic sea-ice as agents of primary production, 199, 1255–1257, 1963.
- Burkert, A., Douglas, T. A., Waldrop, M. P., and Mackelprang, R.: Changes in the Active, Dead, and Dormant Microbial Community Structure across a Pleistocene Permafrost Chronosequence, *Appl. Environ. Microbiol.*, 85, e02646-18, <https://doi.org/10.1128/AEM.02646-18>, 2019.
- 670 Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., and Holmes, S. P.: DADA2: High-resolution sample inference from Illumina amplicon data, 13, 581–583, <https://doi.org/10.1038/nmeth.3869>, 2016.
- Campen, R. K., Sowers, T., and Alley, R. B.: Evidence of microbial consortia metabolizing within a low-latitude mountain glacier, 31, 231, [https://doi.org/10.1130/0091-7613\(2003\)031<0231:EOMCMW>2.0.CO;2](https://doi.org/10.1130/0091-7613(2003)031<0231:EOMCMW>2.0.CO;2), 2003.
- 675 Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens, S. M., Betley, J., Fraser, L., Bauer, M., and others: Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms, 6, 1621–1624, 2012.

- Cheng, S. M. and Foght, J. M.: Cultivation-independent and -dependent characterization of Bacteria resident beneath John Evans Glacier: Characterization of subglacial Bacterial communities, 59, 318–330, <https://doi.org/10.1111/j.1574-6941.2006.00267.x>, 2007.
- 680 Christner, B. C.: Incorporation of DNA and Protein Precursors into Macromolecules by Bacteria at -15°C, 68, 6435–6438, <https://doi.org/10.1128/AEM.68.12.6435-6438.2002>, 2002.
- Christner, B. C., Mikucki, J. A., Foreman, C. M., Denson, J., and Priscu, J. C.: Glacial ice cores: A model system for developing extraterrestrial decontamination protocols, 174, 572–584, <https://doi.org/10.1016/j.icarus.2004.10.027>, 2005.
- 685 Cuffey, K. M., Conway, H., Gades, A. M., Hallet, B., Lorrain, R., Severinghaus, J. P., Steig, E. J., Vaughn, B., and White, J. W. C.: Entrainment at cold glacier beds, 28, 351–354, 2000.
- Dieser, M., Broemsen, E. L. J. E., Cameron, K. A., King, G. M., Achberger, A., Choquette, K., Hagedorn, B., Sletten, R., Junge, K., and Christner, B. C.: Molecular and biogeochemical evidence for methane cycling beneath the western margin of the Greenland Ice Sheet, 8, 2305–2316, <https://doi.org/10.1038/ismej.2014.59>, 2014.
- 690 Doyle, S., Dieser, M., Broemsen, E., and Christner, B.: General characteristics of cold-adapted microorganisms, in: Polar Microbiology: Life in a Deep Freeze, American Society of Microbiology, 103–125, 2012.
- Doyle, S., Montross, S., Skidmore, M., and Christner, B.: Characterizing Microbial Diversity and the Potential for Metabolic Function at -15 °C in the Basal Ice of Taylor Glacier, Antarctica, 2, 1034–1053, <https://doi.org/10.3390/biology2031034>, 2013.
- 695 Eisenhofer, R., Minich, J. J., Marotz, C., Cooper, A., Knight, R., and Weyrich, L. S.: Contamination in Low Microbial Biomass Microbiome Studies: Issues and Recommendations, Trends in Microbiology, 27, 105–117, <https://doi.org/10.1016/j.tim.2018.11.003>, 2019.
- [Ensign, J.: Formation, properties, and germination of actinomycete spores., Annual Review of Microbiology, 32, 185–219, https://doi.org/10.1146/annurev.mi.32.100178.001153, 1978.](https://doi.org/10.1146/annurev.mi.32.100178.001153)
- 700 Filippidou, S., Wunderlin, T., Junier, T., Jeanneret, N., Dorador, C., Molina, V., Johnson, D. R., and Junier, P.: A Combination of Extreme Environmental Conditions Favor the Prevalence of Endospore-Forming Firmicutes, Front. Microbiol., 7, <https://doi.org/10.3389/fmicb.2016.01707>, 2016.
- Fraser, C. I., Connell, L., Lee, C. K., and Cary, S. C.: Evidence of plant and animal communities at exposed and subglacial (cave) geothermal sites in Antarctica, Polar Biol, 41, 417–421, <https://doi.org/10.1007/s00300-017-2198-9>, 2018.
- 705 Goordial, J., Davila, A., Lacelle, D., Pollard, W., Marinova, M. M., Greer, C. W., DiRuggiero, J., McKay, C. P., and Whyte, L. G.: Nearing the cold-arid limits of microbial life in permafrost of an upper dry valley, Antarctica, 10, 1613–1624, <https://doi.org/10.1038/ismej.2015.239>, 2016.
- Gow, A. J., Ueda, H. T., and Garfield, D. E.: Antarctic Ice Sheet: Preliminary Results of First Core Hole to Bedrock, 161, 1011–1013, <https://doi.org/10.1126/science.161.3845.1011>, 1968.
- 710 Hawkings, J. R., Skidmore, M. L., Wadham, J. L., Priscu, J. C., Morton, P. L., Hatton, J. E., Gardner, C. B., Kohler, T. J., Stibal, M., Bagshaw, E. A., Steigmeyer, A., Barker, J., Dore, J. E., Lyons, W. B., Tranter, M., Spencer, R. G. M., and Team, the S. S.: Enhanced trace element mobilization by Earth’s ice sheets, PNAS, 117, 31648–31659, <https://doi.org/10.1073/pnas.2014378117>, 2020.

- 715 Hood, E., Fellman, J., Spencer, R. G. M., Hernes, P. J., Edwards, R., D'Amore, D., and Scott, D.: Glaciers as a source of ancient and labile organic matter to the marine environment, 462, 1044–1047, <https://doi.org/10.1038/nature08580>, 2009.
- Hopwood, M. J., Carroll, D., Dunse, T., Hodson, A., Holding, J. M., Iriarte, J. L., Ribeiro, S., Achterberg, E. P., Cantoni, C., Carlson, D. F., Chierici, M., Clarke, J. S., Cozzi, S., Fransson, A., Juul-Pedersen, T., Winding, M. S., and Meire, L.: Review Article: How does glacier discharge affect marine biogeochemistry and primary production in the Arctic?, 1–51, <https://doi.org/10.5194/tc-2019-136>, 2019.
- 720 Hubbard, B., Cook, S., and Coulson, H.: Basal ice facies: a review and unifying approach, 28, 1956–1969, <https://doi.org/10.1016/j.quascirev.2009.03.005>, 2009.
- Itcus, C., Pascu, M. D., Lavin, P., Perşoiu, A., Iancu, L., and Purcarea, C.: Bacterial and archaeal community structures in perennial cave ice, *Sci Rep*, 8, <https://doi.org/10.1038/s41598-018-34106-2>, 2018.
- 725 Janssen, S., McDonald, D., Gonzalez, A., Navas-Molina, J. A., Jiang, L., Xu, Z. Z., Winker, K., Kado, D. M., Orwoll, E., Manary, M., Mirarab, S., and Knight, R.: Phylogenetic Placement of Exact Amplicon Sequences Improves Associations with Clinical Information, 3, e00021-18, <https://doi.org/10.1128/mSystems.00021-18>, 2018.
- Jung, J., Yoo, K.-C., Rosenheim, B. E., Conway, T. M., Lee, J. I., Yoon, H. I., Hwang, C. Y., Yang, K., Subt, C., and Kim, J.: Microbial Fe(III) reduction as a potential iron source from Holocene sediments beneath Larsen Ice Shelf, *Nat Commun*, 10, 1–10, <https://doi.org/10.1038/s41467-019-13741-x>, 2019.
- 730 Katayama, T., Tanaka, M., Moriizumi, J., Nakamura, T., Brouchkov, A., Douglas, T. A., Fukuda, M., Tomita, F., and Asano, K.: Phylogenetic Analysis of Bacteria Preserved in a Permafrost Ice Wedge for 25,000 Years, 73, 2360–2363, <https://doi.org/10.1128/AEM.01715-06>, 2007.
- Kimes, P. K., Liu, Y., Neil Hayes, D., and Marron, J. S.: Statistical significance for hierarchical clustering, 73, 811–821, <https://doi.org/10.1111/biom.12647>, 2017.
- 735 Knight, P. G.: The basal ice layer of glaciers and ice sheets, *Quaternary Science Reviews*, 16, 975–993, [https://doi.org/10.1016/S0277-3791\(97\)00033-4](https://doi.org/10.1016/S0277-3791(97)00033-4), 1997.
- Kochkina, G., Ivanushkina, N., Karasev, S., Gavrish, E. Y., Gurina, L., Evtushenko, L., Spirina, E., Vorob'eva, E., Gilichinskii, D., and Ozerskaya, S.: Survival of micromycetes and actinobacteria under conditions of long-term natural cryopreservation, 70, 356–364, 2001.
- 740 Korza, G., Setlow, B., Rao, L., Li, Q., and Setlow, P.: Changes in *Bacillus* Spore Small Molecules, rRNA, Germination, and Outgrowth after Extended Sublethal Exposure to Various Temperatures: Evidence that Protein Synthesis Is Not Essential for Spore Germination, *J Bacteriol*, 198, 3254–3264, <https://doi.org/10.1128/JB.00583-16>, 2016.
- Lacelle, D., Radtke, K., Clark, I. D., Fisher, D., Lauriol, B., Utting, N., and Whyte, L. G.: Geomicrobiology and occluded O₂–CO₂–Ar gas analyses provide evidence of microbial respiration in ancient terrestrial ground ice, 306, 46–54, <https://doi.org/10.1016/j.epsl.2011.03.023>, 2011.
- 745 Lamarche-Gagnon, G., Wadham, J. L., Sherwood Lollar, B., Arndt, S., Fietzek, P., Beaton, A. D., Tedstone, A. J., Telling, J., Bagshaw, E. A., Hawkings, J. R., Kohler, T. J., Zarsky, J. D., Mowlem, M. C., Anesio, A. M., and Stibal, M.: Greenland melt drives continuous export of methane from the ice-sheet bed, 565, 73–77, <https://doi.org/10.1038/s41586-018-0800-0>, 2019.

- 750 [Laufer-Meiser, K., Michaud, A. B., Maisch, M., Byrne, J. M., Kappler, A., Patterson, M. O., Røy, H., and Jørgensen, B. B.: Potentially bioavailable iron produced through benthic cycling in glaciated Arctic fjords of Svalbard, *Nature communications*, 12, 1–13, <https://doi.org/10.1038/s41467-021-21558-w>, 2021.](https://doi.org/10.1038/s41467-021-21558-w)
- Lawson, D.: Sedimentological Analysis of the Western Terminus Region of the Matanuska Glacier, Alaska, 132, 1979.
- 755 Lawson, D. E. and Kulla, J. B.: An Oxygen Isotope Investigation of the Origin of the Basal Zone of the Matanuska Glacier, Alaska, *The Journal of Geology*, 86, 673–685, <https://doi.org/10.1086/649736>, 1978.
- Lawson, D. E., Strasser, J. C., Evenson, E. B., Alley, R. B., Larson, G. J., and Arcone, S. A.: Glaciohydraulic supercooling: a freeze-on mechanism to create stratified, debris-rich basal ice: I. Field evidence, 44, 547–562, <https://doi.org/10.3189/S0022143000002069>, 1998.
- 760 Lee, J. E., Edwards, J. S., Schmitt, J., Fischer, H., Bock, M., and Brook, E. J.: Excess methane in Greenland ice cores associated with high dust concentrations, *Geochimica et Cosmochimica Acta*, 270, 409–430, <https://doi.org/10.1016/j.gca.2019.11.020>, 2020.
- McDonald, D., Price, M. N., Goodrich, J., Nawrocki, E. P., DeSantis, T. Z., Probst, A., Andersen, G. L., Knight, R., and Hugenholtz, P.: An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea, 6, 610, 2012.
- 765 Michaud, A. B., Dore, J. E., Achberger, A. M., Christner, B. C., Mitchell, A. C., Skidmore, M. L., Vick-Majors, T. J., and Priscu, J. C.: Microbial oxidation as a methane sink beneath the West Antarctic Ice Sheet, 10, 582–586, <https://doi.org/10.1038/ngeo2992>, 2017.
- Mirarab, S., Nguyen, N., and Warnow, T.: SEPP: SATé-enabled phylogenetic placement, in: *Biocomputing 2012*, World Scientific, 247–258, 2012.
- 770 Miteva, V., Rinehold, K., Sowers, T., Sebastian, A., and Brenchley, J.: Abundance, viability and diversity of the indigenous microbial populations at different depths of the NEEM Greenland ice core, 34, 25057, <https://doi.org/10.3402/polar.v34.25057>, 2015.
- Miteva, V., Sowers, T., Schüpbach, S., Fischer, H., and Brenchley, J.: Geochemical and Microbiological Studies of Nitrous Oxide Variations within the New NEEM Greenland Ice Core during the Last Glacial Period, 33, 647–660, <https://doi.org/10.1080/01490451.2015.1074321>, 2016.
- 775 Monteux, S., Weedon, J. T., Blume-Werry, G., Gavazov, K., Jassey, V. E. J., Johansson, M., Keuper, F., Olid, C., and Dorrepaal, E.: Long-term in situ permafrost thaw effects on bacterial communities and potential aerobic respiration, 12, 2129–2141, <https://doi.org/10.1038/s41396-018-0176-z>, 2018.
- 780 Montross, S., Skidmore, M., Christner, B., Samyn, D., Tison, J.-L., Lorrain, R., Doyle, S., and Fitzsimons, S.: Debris-Rich Basal Ice as a Microbial Habitat, Taylor Glacier, Antarctica, 31, 76–81, <https://doi.org/10.1080/01490451.2013.811316>, 2014.
- Montross, S. N.: Biogeochemistry of basal ice from Taylor Glacier, Antarctica, Montana State University, 2012.
- 785 Montross, S. N., Skidmore, M., Tranter, M., Kivimäki, A.-L., and Parkes, R. J.: A microbial driver of chemical weathering in glaciated systems, 41, 215–218, <https://doi.org/10.1130/G33572.1>, 2013.

[Nicholson, W. L., Munakata, N., Horneck, G., Melosh, H. J., and Setlow, P.: Resistance of *Bacillus* endospores to extreme terrestrial and extraterrestrial environments, *Microbiology and molecular biology reviews*, 64, 548–572, <https://doi.org/10.1128/MMBR.64.3.548-572.2000>, 2000.](https://doi.org/10.1128/MMBR.64.3.548-572.2000)

- 790 O'Connor, B. R. W., Fernández-Martínez, M. Á., Léveillé, R. J., and Whyte, L. G.: Taxonomic Characterization and Microbial Activity Determination of Cold-Adapted Microbial Communities in Lava Tube Ice Caves from Lava Beds National Monument, a High-Fidelity Mars Analogue Environment, 21, 613–627, <https://doi.org/10.1089/ast.2020.2327>, 2021.
- Panikov, N. S., Flanagan, P. W., Oechel, W. C., Mastepanov, M. A., and Christensen, T. R.: Microbial activity in soils frozen to below -39°C , 38, 785–794, <https://doi.org/10.1016/j.soilbio.2005.07.004>, 2006.
- Paterson, W. S. B.: *Physics of glaciers*, Butterworth-Heinemann, 1994.
- Paun, V. I., Icaza, G., Lavin, P., Marin, C., Tudorache, A., Perşoiu, A., Dorador, C., and Purcarea, C.: Total and Potentially Active Bacterial Communities Entrapped in a Late Glacial Through Holocene Ice Core From Scarisoara Ice Cave, Romania, 10, 1193, <https://doi.org/10.3389/fmicb.2019.01193>, 2019.
- 800 Perini, L., Gostinčar, C., and Gunde-Cimerman, N.: Fungal and bacterial diversity of Svalbard subglacial ice, 9, 20230, <https://doi.org/10.1038/s41598-019-56290-5>, 2019.
- Perreault, N. N., Greer, C. W., Andersen, D. T., Tille, S., Lacrampe-Couloume, G., Lollar, B. S., and Whyte, L. G.: Heterotrophic and Autotrophic Microbial Populations in Cold Perennial Springs of the High Arctic, *Appl. Environ. Microbiol.*, 74, 6898–6907, <https://doi.org/10.1128/AEM.00359-08>, 2008.
- 805 Price, P. B.: A habitat for psychrophiles in deep Antarctic ice, *PNAS*, 97, 1247–1251, <https://doi.org/10.1073/pnas.97.3.1247>, 2000.
- Price, P. B. and Sowers, T.: Temperature dependence of metabolic rates for microbial growth, maintenance, and survival, *PNAS*, 101, 4631–4636, <https://doi.org/10.1073/pnas.0400522101>, 2004.
- Price, P. B., Morris, J., Bay, R. C., Adhikari, A., Giovannoni, S. J., and Vergin, K. L.: Molecular Fossils from
810 Microorganisms Preserved in Glacial Ice, <https://doi.org/10.1101/019240>, 2015.
- Priscu, J. C., Tulaczyk, S., Studinger, M., Kennicutt, M., Christner, B. C., and Foreman, C. M.: Antarctic subglacial water: origin, evolution and ecology, 119–135, <https://doi.org/10.1093/acprof:oso/9780199213887.003.0007>, 2008.
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glockner, F. O.: The SILVA ribosomal RNA gene database project: improved data processing and web-based tools, 41, D590–D596, <https://doi.org/10.1093/nar/gks1219>, 2013.
- 815 Rhodes, R. H., Faïn, X., Stowasser, C., Blunier, T., Chappellaz, J., McConnell, J. R., Romanini, D., Mitchell, L. E., and Brook, E. J.: Continuous methane measurements from a late Holocene Greenland ice core: Atmospheric and in-situ signals, *Earth and Planetary Science Letters*, 368, 9–19, <https://doi.org/10.1016/j.epsl.2013.02.034>, 2013.
- Rignot, E. and Jacobs, S. S.: Rapid Bottom Melting Widespread near Antarctic Ice Sheet Grounding Lines, 296, 2020–2023, <https://doi.org/10.1126/science.1070942>, 2002.
- 820

- Salter, S. J., Cox, M. J., Turek, E. M., Calus, S. T., Cookson, W. O., Moffatt, M. F., Turner, P., Parkhill, J., Loman, N. J., and Walker, A. W.: Reagent and laboratory contamination can critically impact sequence-based microbiome analyses, 12, 87, 2014.
- 825 Samyn, D., Svensson, A., and Fitzsimons, S. J.: Dynamic implications of discontinuous recrystallization in cold basal ice: Taylor Glacier, Antarctica, 113, <https://doi.org/10.1029/2006JF000600>, 2008.
- Segev, E., Smith, Y., and Ben-Yehuda, S.: RNA Dynamics in Aging Bacterial Spores, *Cell*, 148, 139–149, <https://doi.org/10.1016/j.cell.2011.11.059>, 2012.
- 830 Segura, J. H., Nilsson, M. B., Haei, M., Sparrman, T., Mikkola, J.-P., Gräsvik, J., Schleucher, J., and Öquist, M. G.: Microbial mineralization of cellulose in frozen soils, *Nat Commun*, 8, 1154, <https://doi.org/10.1038/s41467-017-01230-y>, 2017.
- Sheridan, P. P., Miteva, V. I., and Brenchley, J. E.: Phylogenetic Analysis of Anaerobic Psychrophilic Enrichment Cultures Obtained from a Greenland Glacier Ice Core, 69, 2153–2160, <https://doi.org/10.1128/AEM.69.4.2153-2160.2003>, 2003.
- 835 Siegert, M. J., Ellis-Evans, J. C., Tranter, M., Mayer, C., Petit, J.-R., Salamatin, A., and Prisco, J. C.: Physical, chemical and biological processes in Lake Vostok and other Antarctic subglacial lakes, 414, 603–609, <https://doi.org/10.1038/414603a>, 2001.
- Skidmore, M., Anderson, S. P., Sharp, M., Foght, J., and Lanoil, B. D.: Comparison of Microbial Community Compositions of Two Subglacial Environments Reveals a Possible Role for Microbes in Chemical Weathering Processes, 71, 6986–6997, <https://doi.org/10.1128/AEM.71.11.6986-6997.2005>, 2005.
- 840 Skidmore, M. L., Foght, J. M., and Sharp, M. J.: Microbial Life beneath a High Arctic Glacier, *Appl. Environ. Microbiol.*, 66, 3214–3220, <https://doi.org/10.1128/AEM.66.8.3214-3220.2000>, 2000.
- Sleewaegen, S., Samyn, D., Fitzsimons, S. J., and Lorrain, R. D.: Equifinality of basal ice facies from an Antarctic cold-based glacier, 37, 257–262, <https://doi.org/10.3189/172756403781815708>, 2003.
- Sonjak, S., Frisvad, J. C., and Gunde-Cimerman, N.: *Penicillium* Mycobiota in Arctic Subglacial Ice, 52, 207–216, <https://doi.org/10.1007/s00248-006-9086-0>, 2006.
- 845 Souchez, R., Janssens, L., Lemmens, M., and Stauffer, B.: Very low oxygen concentration in basal ice from Summit, central Greenland, 22, 2001–2004, <https://doi.org/10.1029/95GL01995>, 1995.
- Souchez, R., Bouzette, A., Clausen, H. B., Johnsen, S. J., and Jouzel, J.: A stacked mixing sequence at the base of the Dye 3 Core, Greenland, 25, 1943–1946, <https://doi.org/10.1029/98gl01411>, 1998.
- 850 Souchez, R., Jouzel, J., Landais, A., Chappellaz, J., Lorrain, R., and Tison, J.-L.: Gas isotopes in ice reveal a vegetated central Greenland during ice sheet invasion, 33, <https://doi.org/10.1029/2006GL028424>, 2006.
- Sowers, T.: N₂O record spanning the penultimate deglaciation from the Vostok ice core, 106, 31903–31914, <https://doi.org/10.1029/2000jd900707>, 2001.
- Spirina, E., Durdenko, E., Demidov, N., Abramov, A., Romanovsky, V., and Rivkina, E.: Halophilic-psychrotrophic bacteria of an Alaskan cryopeg—a model for astrobiology, 51, 1440–1452, 2017.

- 855 Steven, B., Briggs, G., McKay, C. P., Pollard, W. H., Greer, C. W., and Whyte, L. G.: Characterization of the microbial diversity in a permafrost sample from the Canadian high Arctic using culture-dependent and culture-independent methods: Microbial diversity in Canadian high Arctic permafrost, *59*, 513–523, <https://doi.org/10.1111/j.1574-6941.2006.00247.x>, 2007.
- 860 Steven, B., Pollard, W. H., Greer, C. W., and Whyte, L. G.: Microbial diversity and activity through a permafrost/ground ice core profile from the Canadian high Arctic, *10*, 3388–3403, <https://doi.org/10.1111/j.1462-2920.2008.01746.x>, 2008.
- Stibal, M., Wadham, J. L., Lis, G. P., Telling, J., Pancost, R. D., Dubnick, A., Sharp, M. J., Lawson, E. C., Butler, C. E. H., Hasan, F., Tranter, M., and Anesio, A. M.: Methanogenic potential of Arctic and Antarctic subglacial environments with contrasting organic carbon sources, *18*, 3332–3345, <https://doi.org/10.1111/j.1365-2486.2012.02763.x>, 2012.
- 865 Strasser, J. C., Lawson, D. E., Larson, G. J., Evenson, E. B., and Alley, R. B.: Preliminary results of tritium analyses in basal ice, Matanuska Glacier, Alaska, U.S.A.: evidence for subglacial ice accretion, *22*, 126–133, <https://doi.org/10.3189/1996AoG22-1-126-133>, 1996.
- Tung, H. c., Price, P. b., Bramall, N. e., and Vrdoljak, G.: Microorganisms Metabolizing on Clay Grains in 3-Km-Deep Greenland Basal Ice, *Astrobiology*, *6*, 69–86, <https://doi.org/10.1089/ast.2006.6.69>, 2006.
- 870 Tuorto, S. J., Darias, P., McGuinness, L. R., Panikov, N., Zhang, T., Häggblom, M. M., and Kerkhof, L. J.: Bacterial genome replication at subzero temperatures in permafrost, *ISME J*, *8*, 139–149, <https://doi.org/10.1038/ismej.2013.140>, 2014.
- Vick-Majors, T. J., Michaud, A. B., Skidmore, M. L., Turetta, C., Barbante, C., Christner, B. C., Dore, J. E., Christianson, K., Mitchell, A. C., Achberger, A. M., Mikucki, J. A., and Priscu, J. C.: Biogeochemical Connectivity Between Freshwater Ecosystems beneath the West Antarctic Ice Sheet and the Sub-Ice Marine Environment, *34*, e2019GB006446, <https://doi.org/10.1029/2019GB006446>, 2020.
- 875 Wadham, J. L., Bottrell, S., Tranter, M., and Raiswell, R.: Stable isotope evidence for microbial sulphate reduction at the bed of a polythermal high Arctic glacier, *219*, 341–355, [https://doi.org/10.1016/S0012-821X\(03\)00683-6](https://doi.org/10.1016/S0012-821X(03)00683-6), 2004.
- Wadham, J. L., Tranter, M., Tulaczyk, S., and Sharp, M.: Subglacial methanogenesis: A potential climatic amplifier?, *22*, <https://doi.org/10.1029/2007GB002951>, 2008.
- 880 Wadham, J. L., Tranter, M., Skidmore, M., Hodson, A. J., Priscu, J., Lyons, W. B., Sharp, M., Wynn, P., and Jackson, M.: Biogeochemical weathering under ice: Size matters, *24*, <https://doi.org/10.1029/2009GB003688>, 2010.
- Wadham, J. L., Arndt, S., Tulaczyk, S., Stibal, M., Tranter, M., Telling, J., Lis, G. P., Lawson, E., Ridgwell, A., Dubnick, A., Sharp, M. J., Anesio, A. M., and Butler, C. E. H.: Potential methane reservoirs beneath Antarctica, *488*, 633–637, <https://doi.org/10.1038/nature11374>, 2012.
- 885 Wang, Q., Garrity, G. M., Tiedje, J. M., and Cole, J. R.: Naïve Bayesian Classifier for Rapid Assignment of rRNA Sequences into the New Bacterial Taxonomy, *Appl. Environ. Microbiol.*, *73*, 5261–5267, <https://doi.org/10.1128/AEM.00062-07>, 2007.
- 890 Willerslev, E., Cappellini, E., Boomsma, W., Nielsen, R., Hebsgaard, M. B., Brand, T. B., Hofreiter, M., Bunce, M., Poinar, H. N., Dahl-Jensen, D., Johnsen, S., Steffensen, J. P., Bennike, O., Schwenninger, J.-L., Nathan, R., Armitage, S., de Hoog, C.-J., Alfimov, V., Christl, M., Beer, J., Muscheler, R., Barker, J., Sharp, M., Penkman, K. E. H., Haile, J., Taberlet, P., Gilbert, M. T. P., Casoli, A., Campani, E., and Collins, M. J.: Ancient Biomolecules from Deep Ice Cores Reveal a Forested Southern Greenland, *317*, 111–114, <https://doi.org/10.1126/science.1141758>, 2007.

Yde, J. C., Finster, K. W., Raiswell, R., Steffensen, J. P., Heinemeier, J., Olsen, J., Gunnlaugsson, H. P., and Nielsen, O. B.: Basal ice microbiology at the margin of the Greenland ice sheet, 51, 71–79, <https://doi.org/10.3189/172756411795931976>, 2010.

- 895 Zhong, Z.-P., Tian, F., Roux, S., Gazitúa, M. C., Solonenko, N. E., Li, Y.-F., Davis, M. E., Van Etten, J. L., Mosley-Thompson, E., Rich, V. I., Sullivan, M. B., and Thompson, L. G.: Glacier ice archives nearly 15,000-year-old microbes and phages, *Microbiome*, 9, 160, <https://doi.org/10.1186/s40168-021-01106-w>, 2021.