



Englacial Architecture of Lambert Glacier, East Antarctica

Rebecca J. Sanderson¹, Kate Winter², S. Louise Callard¹, Felipe Napoleoni^{3,4}, Neil Ross¹, Tom A. Jordan⁵,
Robert G. Bingham⁶

5

¹School of Geography, Politics and Sociology, Newcastle University, Newcastle, UK

²Department of Geography and Environmental Sciences, Faculty of Engineering and Environment Northumbria University, Newcastle, UK

10 ³Department of Geography, Durham University, Durham, UK

⁴Centro de Estudios Científicos, Valdivia, Chile

⁵British Antarctic Survey, Cambridge, UK

⁶School of GeoSciences, University of Edinburgh, Edinburgh, UK

Correspondence to: Corresponding author: Rebecca Sanderson (r.sanderson5@newcastle.ac.uk)

15

20

25



Abstract. The analysis of englacial layers using radio-echo sounding data enables the characterisation and reconstruction of current and past ice-sheet flow. Despite the Lambert Glacier catchment being one of the largest in Antarctica, discharging ~16% of East Antarctica’s ice, its englacial architecture has been little analysed. Here, we present a comprehensive analysis of Lambert Glacier’s englacial architecture using radio-echo sounding data collected by the Antarctica's Gamburtsev Province Project (AGAP) North survey. We used an “internal-layering continuity index” (ILCI) to characterise the internal architecture of the ice and identify four macro-scale ILCI zones with distinct glaciological contexts. Whilst the catchment is dominated by continuous englacial layering, disrupted or discontinuous layering is highlighted by the ILCI at both the onset of enhanced ice flow (defined here as $>15 \text{ ma}^{-1}$) and along the shear margin, revealing the transition from internal-deformation-controlled to basal-sliding-dominated ice flow. These zones are characterised by buckled and folded englacial layers which align with the current ice-flow regime, and which we interpret as evidence that the flow direction of the Lambert Glacier trunk has changed little, if at all, during the Holocene. However, disturbed englacial layers along a deep subglacial channel that does not correspond to modern ice-flow routing suggest that ice-flow change has occurred in a former tributary which fed Lambert Glacier from grid north. As large outlet systems such as Lambert Glacier are likely to play a vital role in the future drainage of the East Antarctic Ice Sheet, constraining their englacial architecture to reconstruct their past ice flow and assess basal conditions is important.

1 Introduction

The Antarctic Ice Sheet, particularly in West Antarctica, is currently undergoing rapid thinning (Fricker et al., 2012; Shepherd et al., 2018; Rignot et al., 2019). Whilst changes in East Antarctica have been less widespread and lower magnitude than those observed in West Antarctica (Gardner et al., 2018), East Antarctica accounts for ~75% of the total Antarctic Ice Sheet area and contains 52 m of potential sea-level rise (Fricker et al., 2021; Stokes et al., 2022). As temperatures increase, major outlet glaciers in East Antarctica could see increases in ice discharge and changes to ice-flow (Li et al., 2016; Miles et al., 2021), with subsequent impacts for future sea-level rise (Miles et al., 2013; Stokes et al., 2022). To forecast future sea-level rise, empirical constraints are required to inform ice-sheet models. These constraints include regional flow history, ice-sheet structure and rheology, and basal boundary conditions, which can be derived from radio-echo sounding (RES) characterisation of the englacial architecture and bed of the ice sheet (Holschuh et al., 2017).

Lambert Glacier is part of the wider Lambert-Amery system (Fig. 1a) (Fricker et al., 2000), the largest glacier-ice-shelf system in East Antarctica, making it an important component of overall ice-sheet mass-balance. Mass-balance observations (Wen et al., 2008; Yu et al., 2010) and numerical-modelling studies (Bassis et al., 2005; Glasser et al., 2015) suggest that the Lambert-Amery system currently has an overall positive mass balance of $20.9 \pm 1.9 \text{ Gt a}^{-1}$ (Gong et al., 2014; Pittard et al., 2017; Cui



et al., 2020), and significant future retreat beyond the present-day grounding line is not projected over the next 500 years, even under a range of future climate scenarios (Pittard et al., 2017). However, low-angle retrograde slopes 250 km upstream of Lambert Glacier's current grounding line mean that marine-ice-sheet instability may occur over longer timescales (Gong et al., 2014). A deep trough situated below Lambert Glacier extends ~600 km inland of the grounding line. Subglacial troughs such as these have the ability to drain the ice sheet rapidly (Morlighem et al., 2020). Because of this potential, and also because areas of Antarctica which have been "stable" for a long time can hold vital information about ice-sheet history (Pattyn et al., 2020), this study characterises the flow of Lambert Glacier from analysis of the catchment's englacial stratigraphy and architecture in RES profiles.

Englacial features (e.g. englacial layers and basal ice units) identifiable in RES data (e.g. Siegert, 1999; Bingham et al., 2007a; Bell et al., 2011; Schroeder et al., 2020) record past and present ice flow and internal deformation. The shape, form, presence, and even absence, of radar-imaged englacial features have been used to reconstruct past disruptions in flow regime (e.g. Siegert et al., 2003b; Bingham et al., 2007b) and accumulation rates (e.g. Winter et al., 2019; Ashmore et al., 2020; Bodart et al., 2021); whilst also allowing us to incorporate age tracers into ice-sheet models (e.g. Parrenin et al., 2017; Sutter et al., 2021). In this study, we undertake analysis of the englacial architecture of Lambert Glacier using the Internal Layering Continuity Index (ILCI; after Karlsson et al., (2012); Bingham et al., (2015)) to assess the current flow regime, and better understand past changes in the region, which could elucidate future change.

2 Study Area

Lambert Glacier (71–75° S, 68° E) contains ~1,480,000 km² of ice, comprising 16% of the grounded ice of East Antarctica (Fricker et al., 2000), which is equivalent to ~8 m of global sea level (Rignot et al., 2019). With a width of over 40 km and a length of 400 km from the grounding line to the southernmost point in which ice flow is visible on Landsat, Lambert Glacier is the largest ice stream in the world (Allison, 1979). Draining into Amery Ice Shelf, ice velocity in the Lambert Glacier catchment increases away from the ice divide at Dome A, where ice flow is 0-5 ma⁻¹ (Fig. 1c). The onset of enhanced ice flow is defined as the location of the transition between inland ice flow and streaming ice flow (Bindschadler et al., 2001); in Lambert Glacier we define this transition where ice velocity exceeds 15 ma⁻¹. The BedMachine (v2) compilation of basal topography in Antarctica (Morlighem et al., 2020) (Fig. 1b) shows that this enhanced ice flow is channelised through a deep topographic graben-like rift valley depression, known as the Lambert Rift (Leitchenkov et al., 2018). The Lambert Rift has a topographic relief of ~3500 m (Fig. 1b), with the bed resting > 2000 m below sea-level (Fig. 1b) (Allison, 1979; Morlighem et al., 2020). Where the rift valley is deepest, the ice is >4 km thick (Morlighem et al., 2020)(Fig. 1d). Ice-flow velocity increases markedly (i.e. from 15 to 250 ma⁻¹) as ice flows through the rift, towards the Amery Ice Shelf (Fig. 1c, 1d).




Fig. 1. Location of Lambert Glacier, within the wider Lambert-Amery system of East Antarctica. The Lambert-Amery catchment is marked with a dotted black line (Zwally *et al.*, 2012b). (a) Reference Elevation Model of Antarctica (REMA)-derived ice-surface elevation showing the location of relevant AGAP-N RES flight lines (red). Contours (grey solid line) are at 500 m intervals (Howat *et al.*, 2019). Dome A and B are highlighted by yellow circles. Subglacial-lake locations (Livingstone *et al.*, 2022) are marked with blue circles. (b) Bed elevation of the study area from BedMachine v2 (Morlighem *et al.*, 2020), which includes the AGAP-N RES bed-elevation data. The grounding line is shown with a black line (Bindschadler, *et al.*, 2011). (c) Ice-flow speed averaged over 1992-2017, overlying a raster of the maximum horizontal gradient (65) of ice velocity with a grey-scale colour pallet (Mouginot *et al.*, 2019). (d) Ice thickness from BedMachine v2 (Morlighem *et al.*, 2020). (e) High resolution RAMP RADARSAT-1 radar imagery of Lambert
90
95 Glacier (Jezek, 1999)



3 Methods

3.1 Data

The RES data used for this study were acquired in 2007-2009 over the northern region covered by the internationally collaborative aerogeophysical survey of Antarctica's Gamburtsev Province (AGAP) Project (Bell et al., 2011; Ferraccioli et al., 2011; Rose et al., 2013). The total “AGAP North” survey grid (highlighted in Fig. 1a) comprises 120,000 line km of data, with a line spacing of 5 km and tie lines ~33 km apart.

All data were acquired with the 150 MHz British Antarctic Survey PASIN (Polarmetric Airborne Survey INstrument) radar system, operating over a bandwidth of 15-20 MHz, whilst deploying a 4 μ s, 10MHz linear chirp designed to sound as deeply as possible into the ice. 2-D Synthetic Aperture Radar (SAR) processing was applied to enhance resolution and reduce echo signal noise to generate the freely-available dataset deposited in the BAS Polar Airborne Geophysics Data Portal (Frémand et al., 2022). Previous applications of these data have: (i) used bed echoes to map subglacial topography (Fretwell et al., 2013); (ii) explored the geological and geomorphological evolution of the Gamburtsev Subglacial Mountains (Ferraccioli et al., 2011; Rose et al., 2013); and (iii) analysed the region’s distinctive basal ice units (Bell et al., 2011). Across part of the same region, Luo et al. (2020) used a single traverse of ground-based ice-penetrating radar data from Zhongshan station to Dome A to investigate ice-sheet internal structure and basal roughness, applying similar analysis to that deployed for this paper, but over a much more geographically-limited area. Our study is the first to interrogate comprehensively the englacial architecture of Lambert Glacier catchment using multiple radar survey lines. To place our analysis and findings in a wider context, we combine our analysis of the radar data with freely available MEaSURES ice-surface velocity data (Mouginot et al., 2019), BedMachine bed-topography maps (Morlighem et al., 2020) and RADARSAT Antarctic Mapping Project RAMP satellite imagery (Jezek, 1999).


3.2 Internal layer continuity analysis

Following Karlsson et al. (2012) and Bingham et al. (2015), we applied the “Internal Layer Continuity Index” (ILCI) to classify and map englacial stratigraphy across the Lambert Glacier catchment. The ILCI was initially developed as a tool for objectively and quantitatively capturing the degree to which continuous and flat isochrones had previously been qualitatively observed to be buckled or disrupted by variations in ice flow (e.g. Rippin et al., 2003; Siegert et al., 2003b; Bingham et al., 2007b; Karlsson et al., 2009). The ILCI operates by quantifying the average power of peaks observed through the ice column in each trace, and then further averaging the trace-by-trace results over windows of a predefined length along radar tracks (Fig. 2). Although the ILCI does not directly analyse the continuity of high-power peaks between adjacent traces, the most commonly accepted explanation for consistently high ILCI measured across multiple kms of ice is that englacial layering is continuous and undisturbed over that region, as noted in several previous studies (e.g. Karlsson et al., 2012; Karlsson et al., 2014; Bingham et al., 2015; Winter et al., 2015; Winter et al., 2016). The most common explanation for low ILCI across a region is that the reflection power (as measured trace by trace) has been reduced by buckling, folding, warping, or even the complete absence



130 of englacial layering. This disruption is understood to be the result of ice currently, or previously, experiencing significant deformation as it flows through an ice-stream onset or shear zone, or around/over significant subglacial protuberances (Bingham et al., 2015; Winter et al., 2015; Winter et al., 2016). ILCI therefore provides a useful mechanism for reconnoitring englacial conditions across large regions of an ice sheet, to characterise broad ice-flow conditions and guide more detailed investigations in regions of interest. It is in this spirit that we apply the ILCI technique in this study.

135 Following previous ILCI applications (Karlsson et al., 2012; Bingham et al., 2015; Winter et al., 2016), the upper and lower 20% of the ice column, where englacial layers are often absent in RES returns, was removed. We averaged ILCI over windows of 100 and 1000 traces (2 -20 km along track) to identify both small-scale ‘localised’ anomalies and more regional trends. From the broad range of ILCI values retrieved by ILCI across Lambert Glacier, we define a “high-continuity” ILCI value to be >0.45 . This value is higher than other studies which have applied ILCI to PASIN radar data which have used 0.18 (Karlsson et al., 2012), 0.10 (Bingham et al., 2015), and 0.06 (Winter et al., 2015; Winter et al., 2016) as an indicative threshold after
140 which continuity is defined as “high”. However, our use of a 0.45 threshold is consistent with the recommendation from a recent pan-Antarctic application of ILCI that the threshold will vary depending on the purpose of the study and the region of interest (Frémand et al., 2022). The “high” ILCI value defined over the Lambert Glacier catchment reflects sensitivity of the analysis when applied to a larger region, like ours, which varies significantly in bed elevation and ice velocity.



145 **Fig. 2.** Example radargrams from the AGAP North survey showing: (a) continuous englacial layers along RES survey line A44b, (b)
 discontinuous layering within RES survey line A29h and (c) absent layers in RES survey line A39d. In each panel, the continuity
 index (ILCI) is shown in red (unsmoothed), with a running mean of 10 traces highlighted by the black line. (d) Profile locations
 marked onto site-wide ILCI results (using a running mean of 100 traces). Background is BedMachine v2 bed topography (Morlighem
et al., 2020). (f) A-scope plot demonstrating relative received power (dB) (left) and the absolute value of the gradient, for a trace in
 panel (a) (demonstrating continuous layering). (g) similar to (f), an A-scope of relative received power (left) and the absolute value
 150 of the gradient (right) for a trace in panel (b) (characterising disrupted layering). For both (f) and (g), the black line depicts the
 whole trace whilst the blue is the section used for calculation of the ILCI.



4 Results

155 ILCI results derived from our 100-trace (2 km) and 1000-trace (20 km) windows reveal four macro-scale zones across the Lambert Glacier catchment (Fig. 3):

Zone 1: ILCI values are generally high (>0.45) in the upper catchment, before the onset of enhanced ice flow (i.e. $>15 \text{ ma}^{-1}$). This zone nominally represents an area of continuous englacial layering (e.g. Fig. 2a).

160 Zone 2: Very low ILCI values (<0.3) dominate the glacier onset zone, where ice currently transitions from low flow speeds to faster streaming flow. These low ILCI values likely reflect buckled and disrupted ice structures (see Fig. 2b).

Zone 3: Variable ILCI values (0.25-0.5) are returned in Zone 3, within the lower catchment. Here, areas of high ILCI (>0.4) are transected by two narrow zones of low ILCI (<0.3) which coincide with channelised, fast glacier flow through deep subglacial valleys.


165 Zone 4: ILCI values are relatively low (<0.4) around Lake 90°E and Sovetskaya Lake, where there is a region of slow-flowing thick ice ($>3000 \text{ m}$).

4.1 Englacial continuity, ice-flow and bed topography

A spatial correspondence between ILCI returns and the current ice-flow regime at the glacial surface is apparent across the Lambert Glacier catchment (Fig. 3a). In Zone 1, up-ice of the onset zone (Fig. 3a), no ice flow exceeds 10 ma^{-1} across an area of $38,000 \text{ km}^2$. This slow-flowing inland ice is characterised by high ILCI values (>0.45), where similar slow-flow conditions
170 throughout the ice column help to maintain englacial stratigraphic continuity. Downstream of Zone 1 and into the Lambert Glacier onset zone (Zone 2), the ILCI signature evolves as surface ice-flow speeds increase (Fig. 3a). Low ILCI, associated with disrupted or absent layering (<0.3), begins to occur where ice-flow speeds pick up, to between $\sim 10\text{-}15 \text{ ma}^{-1}$, including through a topographically controlled tributary-like zone (marked with a T in Fig. 3) where ice-flow remains below $<15 \text{ ma}^{-1}$.
175 Through much of the rest of Zone 2, ice is channelled through substantial ice-flow trunk that connects the onset zone of Lambert Glacier to the Amery Ice Shelf (Fig. 4). Fast-flowing ice in this trunk is associated with low ILCI (Fig. 3 (Zone 2)) and disrupted englacial layering (Fig. 4). However, down ice flow, through Zone 3, the ILCI is bimodal. Ice in the main trunk of Lambert Glacier displays the expected association of low ILCI values (<0.3) where ice-flow speeds are high ($> 100 \text{ m a}^{-1}$). Elsewhere, however, and throughout Zone 3, ILCI is often >0.45 . These ILCI values are typically associated with continuous englacial layering, which is surprising, given that surface ice velocities are generally quite fast flowing, and always in excess
180 of 20 ma^{-1} . Zone 4 (Fig. 3) is beyond the main AGAP-N radar survey grid, and therefore characterised instead by six reconnaissance AGAP-N RES survey lines grid-south of Lambert Glacier. Although ice velocity is $<10 \text{ ma}^{-1}$ across the entirety of this zone, ILCI values are <0.4 contrasting the assumption that ice flowing slower will have a high ILCI return. Applying a lower threshold for “high” ILCI might highlight higher resolution changes in continuity in Zone 4, however, as we are primarily looking at the regional picture, this type of detailed analysis is beyond the scope of the present study.



185 The upper catchment bed topography in Zone 1 is dominated by the high relief of the subglacial Gamburtsev Subglacial
Mountains, beneath Dome A (Morlighem et al., 2020) (Figs 1, 3a). When averaged over 1000 traces (Fig. 3b) the ILCI analysis
does not appear to capture the impact of these subglacial mountains on englacial continuity. However, when ILCI is averaged
over 100 traces (Fig. 3a), variations in ILCI across zone 1 suggest an association between the steep subglacial bed slopes and
disrupted or absent englacial layering above. Grid east of the upper catchment there is a region of lower bed relief with higher,
190 more continuous, ILCI values (>0.5). The high ILCI in this region correspond with parts of the catchment which display an
ice thickness of <2000 m (Fig. 1c). As ice enters the deep graben-like rift valley through which ice flow is currently focused,
there is a spatial correspondence between increased ice velocity (i.e. >15 ma^{-1}), the deepening of the valley (~ 400 m asl to \sim
200 m asl) and low ILCI values (<0.3). Grid west of the onset, a tributary-like valley (marked with a T in Fig. 3) demonstrates
low bed elevation (1,000 m b.s.l. at the deepest point), and low ILCI values (<0.3). However, ice velocity here is slower (<10
195 ma^{-1}), which implies that the roughness of the bed topography is the cause of low layer continuity. Downstream of the onset
zone, in Zone 3 (Fig. 3b), the ILCI analysis shows a correspondence between the englacial architecture and bed topography.
Adjacent to the central glacier trunk and rift valley, broad regions (~ 250 km wide) of flatter bed relief and large, relatively flat,
subglacial platforms are associated with higher ILCI values (>0.45), reflecting continuous englacial layering throughout the
ice column, and the layering is disrupted only where fast flow follows the deep geological rift and at the shear margin (Fig. 4).
200 Zone 4 contains a similar area of flatter bed relief (Fig. 3b) but, unlike Zone 3, this zone is characterised by lower ILCI values
(<0.4), suggestive of more disrupted ice flow.



205 Fig. 3. ILCI analysis demonstrating the four zones of ILCI returns in the Lambert Glacier. White line is the Lambert-Amery
catchment boundary (Zwally et al., 2012a) (a) ILCI smoothed over 100 traces, overlying MEaSUREs ice-surface velocity
measurements (Mouginot et al., 2019). (b) ILCI analysis smoothed over 1000 traces, overlying BedMachine v2 bed topography
(Morlighem et al., 2020). Blue circles depict the four dominant ILCI zones within [1] the upper catchment; [2] the onset zone; [3]
downstream; and [4] grid south of the onset. A tributary feeding Lambert Glacier from grid north is denoted with a T and white
dashed lines highlight the tributary. White circles on Fig. 3a mark the location of RES transects displayed in Figs 4 and 7.

210





Fig. 4. Ice velocity (Mouginot et al., 2019) and Internal Layering Continuity Index (ILCI) returns from RES line A29C. White boxes highlight disrupted englacial layering where ice flows through the central rift and at the margin of Lambert Glacier. The location of this transect is noted in Fig. 3a.

215 4.2 Englacial folding

The spatial correspondence between ice flow exceeding 15 ma^{-1} and the marked shift in ILCI values in Zone 2 (Fig. 3a) suggests that the onset of ice flow is important for the englacial structure of Lambert Glacier. Although ILCI values are relatively low in this zone (i.e. <0.4), radargrams show lateral continuity of englacial structures across multiple flight lines, where recognisable units or structures can be mapped from radargram to radargram down-ice (Fig. 5). Englacial layers at the onset of enhanced flow ($>15 \text{ ma}^{-1}$) undulate in distinct trough and crest sequences, similar to folds observed elsewhere in Antarctica and Greenland (Bell et al., 1998; Conway et al., 2002; Ng et al., 2004; Holschuh et al., 2014; Franke et al., 2022a). The axes of these folds are subparallel to the current flow direction, with fold width decreasing down flow from the onset zone, as ice flow converges. The amplitude of folding increases with ice thickness. Eighteen distinct flow bands (defined by prominent englacial layer trough folds) have been identified within the region of ice-flow onset (Fig. 5). All flow bands propagate down-ice as the ice flows through the subglacial rift valley beneath Lambert Glacier. Although the fold axes are typically vertical in orientation, the lowermost parts of some diverge from the vertical close to the bed (e.g. at depths of ~ 2.5 km). An example of this can be seen approximately 25-30 km along RES transect A46B-A46B' and A54B-A54B' (Fig. 5) where the fold axes deflect towards the centre of the valley. In each case, the deflection of the fold axes appears just above a zone of basal ice that is characterised by few or no internal reflections (i.e. an 'echo-free zone').

230



235 Fig. 5. Radar profiles a) (A46B-A46B'), b) (A58D-A58D') and c) (A54B-A54B') showing englacial layering and the subglacial bed. Flow direction is from panels A to C. Eighteen fold bands, defined by a down-ice-warping in englacial layers, are mappable down-ice from panel A to panel C (dashed red lines). d) Current ice velocity and flow vectors (white arrows) (Mouginot et al., 2019) of the onset of Lambert Glacier with flow direction from bottom left to top right; red lines denote profiles in (a), (b) and (c).



In addition to the high-frequency folds observed at the ice-stream onset in Zone 2, large-scale englacial structures are visible in radargrams collected along the western margin of Lambert Glacier as ice flow converges towards its central trunk. A single large-scale fold (>360 km long, 5-10 km wide), characterised by a diagnostic, visually-traceable englacial structure, is apparent in the slower-flowing ice outside the Lambert Glacier tributary shear margin (Fig. 6). This englacial feature is visible in a series of 36 radargrams, through which it can be traced ~360 km down-ice of the onset zone (Fig. 6a). The limits of the survey grid restrict further tracing of this feature, but we are confident that the fold extends further upstream and downstream than the 360 km we map here. As ice velocity increases down flow, from 15 ma^{-1} to $>100 \text{ ma}^{-1}$, the width of the fold reduces, and its amplitude increases. The fold runs parallel to ice flow in an area of accelerating and converging ice-flow velocities ($\sim 15 \text{ ma}^{-1}$ to $\sim 50 \text{ ma}^{-1}$) (Fig. 6b), and is located immediately adjacent to the central rift valley of the enhanced flow zone of Lambert Glacier (Fig. 6c). Assuming that the fold formed and then advected down-ice with flow, we estimate that this englacial fold could have persisted for at least 10.5 ka (based on calculations of the average current ice velocity (Mouginot et al., 2019) and the distance between each fold on individual flightlines). In most radargrams, the fold is unrelated to the bed topography (i.e. the layers do not drape over a localised topographic high) and the fold axis is parallel to macro-scale ice flow. Similar scale folds, associated with convergent ice flow and the shear margins of outlet-glacier tributaries, have been observed in Greenland and Antarctica (Jacobel et al., 2000; Bons et al., 2016; Ross et al., 2020). The amplitude of the folding ($\sim 200 \text{ m}$ vertical relief) is greatest in the lower Lambert catchment (around Zone 3) and corresponds with pronounced convergent ice-surface flow stripes imaged on RADARSAT (Ely et al., 2016) that seem to initiate in response to accelerating ice flow at the onset zone (Fig. 6). One of these flow stripes corresponds with the fold axis of the traceable englacial fold, and extends the full $\sim 360 \text{ km}$ down-ice (Fig. 6). The surface flowline from the peak of the fold is undetectable in RADARSAT imagery in the upper catchment where ice velocity drops below $\sim 10 \text{ ma}^{-1}$. However, the englacial fold extends significantly further up-ice of this point, suggesting that it extends further into the upper catchment than the overlying surface flow stripe with which it is associated.




Fig. 6. (a) Englacial-layer folds within a selection of radargrams from upstream transect A46B-A46B' to down-ice transect A38B-A38B'. Locations of lines A46B-A38B shown on b and c ; (b) Fold axis locations mapped with a yellow circle, along transects A46B to A38B, underlain by ice-flow speed overlying a raster of the maximum horizontal gradient (∇) of ice velocity with a grey-scale colour pallet (Mouginot *et al.*, 2019); (c) Fold axis locations mapped between lines A46B and A38B, underlain by bed elevation (Morlighem *et al.*, 2020). (d) Location of surface flow stripes (Ely and Clark, 2016) (black lines) and peaks of the englacial fold axes (yellow dots) across the wider AGAP-N radar dataset, overlain on RAMP RADARSAT radar imagery of Lambert Glacier (Jezek 1999). In panels b-d, the white line denotes the boundary of the Lambert-Amery catchment.

260



265 5 Discussion

5.1 Catchment-scale englacial architecture of Lambert Glacier

The overall pattern of ILCI across the Lambert catchment has a spatial correspondence to the bed topography. Ice flowing over the subglacial highlands (northern Gamburtsev Province) upstream of the Lambert Glacier onset zone has low ILCI (<0.3) when averaged over 100 traces. This low ILCI in the upper ice-stream catchment is not associated with current enhanced or fast ice flow (as the ice currently flows at <math><5 \text{ ma}^{-1}</math>), and there is no evidence of past enhanced/fast ice-flow features such as buckled layers at depth) in the radargrams (e.g. Winter et al., 2015). There are three likely causes for the low ILCI values, the first of which we favour. First, the low ILCI could be a consequence of thick warmer ice, causing attenuation of the radar signal, resulting in poor imaging of deeper englacial layers (Matsuoka et al., 2012). Weak reflections caused by low amplitude power returned from deep layers would impact the ILCI as the method involves assessing the variability of the internal layer reflection on individual traces (Karlsson et al., 2012). Second, it is possible that disruption of the layers giving rise to low ILCI is a result of power loss from dipping englacial reflectors, as seen elsewhere in Antarctica (Holschuh et al., 2014; Winter et al., 2015). However, although low ILCI (<math><0.3</math>) is apparent in areas of steeply inclined subglacial topography in the upper Lambert catchment, the relationship between ILCI and basal topography is not as evident as that between low ILCI and ice thickness. Third, it is possible that thick ice, converging through topographically controlled bed topography and/or refreezing at the bed under pressure, is causing the formation of folds and basal ice packages (Bell et al., 2011; Wrona et al., 2018). These basal ice units would disrupt the ILCI result as they would not contain traceable reflections between traces. However, they are unlikely to have a significant impact on the ILCI, as the lower 20% of the ice thickness (where these basal ice packages are found) was removed prior to ILCI analysis.

In Lambert Glacier's onset zone, where enhanced flow (>15 ma^{-1}) initiates, the ice is channelised into the deep rift valley that constrains Lambert Glacier (Fig. 3b). Buckling of englacial layers is the most likely explanation for the low ILCI values returned in zones 2 and 3 (Fig. 3), as diagnosed for other areas of Antarctica where ice flow speeds up (e.g. Siegert et al., 2003; Bingham et al., 2007). Discontinuous and buckled ice layers dominate the full ice column here, as a result of relatively high current ice-flow speeds (~50 - 150 ma^{-1}). Higher velocities throughout the ice column, caused in part by the presence and properties of basal temperate ice and associated basal sliding (Pittard et al., 2016), allow older ice in this part of the Lambert system to be transported more rapidly towards the grounding line. Pittard et al. (2016) modelled the basal melt rate for the glacial system and found the average to be just 1.3 mm a^{-1} ; however, they also note a large potential variability in the modelled melt, with a maximum modelled basal melt rate of >500 mm a^{-1} . We propose that variable basal sliding, and therefore variable ice-flow speeds across zones 2 and 3, are the primary reason for high ILCI values in the region (which qualitatively define visible internal layer buckles) (Fig. 5). Although this relation may also hold for ice in the main glacier trunk flowing through the Lambert Rift, ILCI returns here (>0.4) suggest more continuous englacial layering across the rest of lower Lambert Glacier, despite presently enhanced ice-flow at the surface. This suggests that in the lower Lambert Glacier (i.e. zone 3), away from



the main glacier trunk, englacial layers are more continuous than they were in ILCI zone 2 (around the onset zone) (Fig. 3a). The higher ILCI in the wider, lower catchment (i.e. outside the main trunk) may be attributed to (i) a lack of significant subglacial topographic variation disrupting the ice or (ii) colder, slow-flowing ice adjacent to the main trunk (Dawson et al., 2022). Although Payne and Baldwin (1999) found that colder, slow-flowing ice can control the location of major ice streams, Young et al. (2019) have stressed more recently that colder temperatures increase stiffness and can therefore reduce deformation of internal layers and hence induce less buckling and disruption.

Zone 4, the region grid south of the main Lambert Glacier channel (Fig. 3), demonstrates a breakdown of the assumption that low ILCI values and discontinuous layering occur as a result of changes in bed topography and/or enhanced ice velocity. In this region, the bed is flatter than the upper catchment and the ice is slower flowing (i.e. $<10 \text{ ma}^{-1}$) (Fig. 3), although the ice is considerably thicker (typically $>3000 \text{ m}$ thick) than across the majority of the catchment (typically $<2000 \text{ m}$ thick) (e.g. Fig. 7). Similar to the low ILCI region Zone 1 (Fig. 3), we suggest that the thick ice here could be increasing the attenuation of the RES waves, generating the weaker reflections read by our automated ILCI analysis. However, another possibility is that the thicker, warmer ice in this area (Van Liefferinge and Pattyn, 2013; Dawson et al., 2022) could also reduce the amplitude of englacial layer undulations and mask continuous englacial layers.




Fig. 7. Example of RES transect (A33) exhibiting visible continuous layers, but a relatively “low” ILCI return. The mean ILCI return for the transect is 0.38. The location of this transect is detailed in Fig. 3a.

315

5.2 The onset zone of Lambert Glacier



The disrupted englacial-layer geometry at the onset of Lambert Glacier (Zone 2, Fig. 3) is comparable to that observed elsewhere in Antarctica (e.g. the tributary onset zones of Pine Island Glacier (Karlsson et al., 2012)), where englacial layers are similarly folded and buckled. As we assume that these buckled layers are the product of lateral shear stresses at the transition from slow and fast flow (Siebert et al., 2003a), ILCI evidence of disrupted englacial layers up-ice of the onset zone suggests that there may not be an abrupt no-slip to sliding transition beneath Lambert Glacier. Instead, a more gradual shift to basal sliding, over a distance of approx. 200 km, appears to take place. However, an alternative view is that the onset zone could have moved over time. We explore these two ideas using pre-existing research from other Antarctic field sites. At Pine Island Glacier, ILCI analysis of englacial layering suggested that while some areas demonstrated a similar gradual transition to basal sliding, in other areas the transition between internal deformation and basal sliding is very sharp (Karlsson et al., 2012). Karlsson et al. (2012) attributed this abrupt transition to a distinct qualitative difference between disturbed and absent layering at a tributary confluence. This setting is not particularly analogous to Lambert Glacier, where there is a broad onset zone, which is not influenced by multiple tributaries like Pine Island Glacier. In this respect, the onset zone of Lambert Glacier appears to be more analogous to that of Bindschadler Ice Stream, West Antarctica, where an analysis of longitudinal stress gradients has shown that a shift to basal sliding occurs as an integrated response of bed and ice interactions upstream of the onset zone, rather than changes in basal characteristics at a set location (Price et al., 2002). The initiation of flow bands at the onset of Lambert Glacier (Fig. 5) (Ely and Clark, 2016) is not coincident with highs in bed topography and therefore the flow bands are likely to have formed as a result of differential basal conditions causing high basal sliding (and resultant ice flow speed up) (Ely and Clark, 2016). Correspondence of disturbed sloping englacial layers and elevated basal shear stress in the Lambert onset zone (Morlighem et al., 2013; Sergienko et al., 2014) supports the theory of an extended gradual transition from a frozen to temperate bed (i.e. from internal deformation to basal sliding), as suggested for Institute Ice Stream in West Antarctica (Bryant et al., 2019; Mantelli et al., 2019). In contrast, tributary 1 (marked with a T on Fig. 3) returns low ILCI values when ice flows through the subglacial valley. Although these ILCI values suggest increased ice velocity, this is not observed at the present-day ice surface (Mouginot et al., 2019). The disruption to the englacial layers suggested by the ILCI is possible evidence that tributary 1 previously experienced enhanced ice flow. This would have been the result of either previous enhanced ice flow in the tributary, or a migration of the onset region of Lambert Glacier, with ice flow from a more northerly direction. Similar tributary flow switching is thought to have occurred at Institute Ice Stream in West Antarctica (as evidenced by RES transect and ILCI analysis) (Winter et al., 2015). In summary, we posit that the onset zone of Lambert Glacier has a gradual transition from internal deformation to basal sliding, but there is evidence of potential flow changes having occurred there in the past.

The trunk of Lambert Glacier is characterised by low-continuity ILCI values and buckled englacial layers which are first recorded at, or just upstream of, the ice-flow onset zone (Fig. 3). We suggest that buckled layer geometry in the glacier trunk, as it flows through the subglacial rift valley, forms as a result of changes to the internal-strain field upstream, as fast ice becomes channelised and/or as large subglacial protuberances disrupt the strain field when ice flows across them. This is



further evidenced by the presence of flow bands, consisting of troughs and crests in multiple radargrams (Fig. 5). Persistent flow bands and buckled folds across tens of km suggest a steady long-term direction of ice flow, as changes in flow direction would have altered the persistence and orientation of these channel buckles.

355 Basal sliding can cause spatial and temporal differences in ice-flow speeds, such as at the onset of Lambert Glacier, and this is often attributed to the presence of meltwater at the glacial bed. Subglacial hydrology has therefore been inferred to play an important role at the onset of faster-flowing ice or ice streams (Bell et al., 2007; Franke et al., 2020; Siegert and Bamber, 2000). There is limited evidence of subglacial lakes in the onset zone of Lambert Glacier (Zone 2, Fig. 3) (Livingstone et al., 2022). However, there are numerous subglacial lakes in the upper catchment of Lambert Glacier, beneath Domes A and B
360 (Fig. 1) (Livingstone et al., 2022) which could supply basal water to the onset zone, and two very large lakes (e.g. Lake 90°E (2000 km²) and Lake Sovetskaya (1600 km²)) are located grid south in the catchment (Bell et al., 2006). These large lakes are tectonically controlled and considered stable (Bell et al., 2006) so large-scale production of meltwater up-ice from these is unlikely to play an important role in the transition from internal deformation to sliding beneath Lambert Glacier.

5.3 Englacial structure of Lambert Glacier shear margin

365 Large-scale englacial folds are located just outside the Lambert shear margin and persist throughout the ice column, and over a distance of 360 km (Fig. 6). The consistent orientation of the fold axis of these features within multiple radar lines transecting the glacier suggests that no significant past changes in flow direction from the current ice-flow regime are recorded in this part of the ice sheet. This supports Glasser et al. (2015), who noted that ice parcels at the onset of Lambert Glacier and other glaciers in Antarctica such as Recovery, Pine Island and Byrd glaciers may have resided for ~ 2500 to 18,500 years. Likewise, the
370 topographic constraint of the rift valley within which the glacier flows imposes a fixed width to Lambert Glacier, restricting the shear margin from migrating easily in response to perturbations in ice flow from external forces (Grinsted et al., 2022). The large-scale englacial folding we observe at the shear margin is therefore likely to be a result of lateral convergence, driven by enhanced ice flow through the subglacial valley (Franke et al., 2022a). However, we note that the englacial architecture of Lambert Glacier appears to be sensitive to variations in basal thermal conditions; where it is likely that ice compression causing
375 the fold is exaggerated by the transition from cold/frozen conditions outside the shear margin to the warm/ thawed conditions in the central trunk. Furthermore, Hills et al. (2022) note that the temperature gradient across the shear margin is likely to be stronger across the shear margin than along it. This helps to account for the fold advection we record down flow, where we find a similar englacial architecture for hundreds of km, with minimal change or disruption to the fold regime.

380 We have found the large-scale englacial fold along Lambert Glacier's shear margin to be directly manifested on the ice-sheet surface as a flow stripe (Fig. 6d). This study provides supporting evidence demonstrating that non-topographic englacial features and local variations in flow regime can lead to the development of surface features such as flow stripes (Cooper et al., 2019; Glasser et al., 2015; Ross et al., 2020). The fold along the shear margin extends significantly further upstream than the



385 corresponding surface flowline. This is consistent with long-term persistence of ice flow in the Lambert catchment and may
provide evidence that the onset of enhanced flow once occurred further up ice flow than at present. The surface expression of
the englacial folding (i.e the flow stripe), tied to the subsurface evidence from the radar, allows us to characterise and infer
large-scale flow patterns over time (Ely and Clark, 2016). The persistent occurrence and orientation of the englacial fold
suggests that the ice-flow pattern of Lambert Glacier has not changed for a substantial period. Based on calculations of the
average current ice velocity (Mouginot et al., 2019), and the distance between each fold on the individual flightline, the
390 englacial fold at the Lambert Glacier shear margin is likely to have persisted for over 10.5 ka (i.e. throughout the Holocene),
consistent with modelled evidence (Gong et al., 2014) for long-term ice-flow stability of Lambert Glacier. Increased
understanding of ice margins is vital for understanding the mechanics of glaciers and ice streams, and how they can change in
the future. The englacial layers across the margin warp but remain continuous along Lambert Glacier, as is the case across
Whillans Ice Stream (Van Der Veen et al., 2007) indicating the sensitivity of englacial layers to high shear strain.

395 **5.4 Wider implications for large outlet glacier systems**

This study has provided an extensive evaluation of the englacial architecture of a large outlet glacier system in East Antarctica.
Assessing large outlet glacier/ice-stream systems, particularly in East Antarctica, is important for ice-sheet mass-balance and
their contributions to sea-level rise (Stokes et al., 2022). Although extensive research has been undertaken on ice streams
(Winsborrow et al., 2010), the recent acceleration of outlet glaciers as a response to climatic or internal perturbations remains
400 poorly constrained (Jordan et al., 2018; Miles et al., 2021). By investigating the englacial architecture of these systems, and
their relationships to the bed and surface, we can obtain an increased understanding of how perturbations at and near the ice-
stream margin can propagate up-ice (e.g. up to the onset zone), and impact the flow of the interior ice. Likewise, our knowledge
of shear-margin formation and migration has been developed, as has been the case for the North East Greenland Ice Stream
(Franke et al., 2022b; Grinsted et al., 2022; Holschuh et al., 2019). Here we have provided a baseline characterisation of the
405 Lambert Glacier system and have analysed RES data to examine current and past ice flow. If similar studies were applied to
other large outlet systems across Antarctica, we will be able to provide more and better, localised constraints for ice-flow
models, which will improve our current understanding of ice flow and discharge in the past and for the present day, with clear
benefits from each, for predictive forecasting.

6 Conclusions

410 This paper has interrogated and characterised englacial stratigraphy across the Lambert Glacier catchment, the largest glacier
ice-shelf system in East Antarctica. Our results highlight four distinct zones of englacial architecture. Zone 1 returns high ILCI
values (>0.45) in the upper catchment, where englacial stratigraphic continuity is maintained by slow-flowing ice conditions.
Zone 2 comprises of very low ILCI values (<0.3) in the onset zone where ice accelerates from slow flow by internal
deformation to more enhanced flow (defined here as $>15 \text{ ma}^{-1}$), which generates buckles and discontinuity in the englacial
415 stratigraphy. Zone 3 defines an area with variable ILCI values (0.25-0.5) within the lower catchment, where areas of traceable,



continuous englacial layers are recorded, as well as regions of discontinuity or englacial layer absence, particularly when ice-flow speeds are high ($> 100 \text{ m a}^{-1}$). Zone 4 contains relatively low ILCI values (< 0.4) outside the main flow of the Lambert Glacier, around Lake 90°E and Sovetskaya Lake, where ice is $\sim 3000 \text{ m}$ thick, but slow flowing.

420 Whilst our results within Zone 1 are to be expected, reflecting findings from similar upper-catchment ice flows in Antarctica, we note interesting findings in all other zones. The englacial stratigraphy in Zone 2 demonstrates a gradual (rather than abrupt) transition from internal deformation to basal sliding at the onset zone. Combined with the minimal deformation and amplitude change of traced flow band fold axes in the region, this finding helps to reveal long term ice flow stability within the main trunk of the Lambert Glacier. This long-term stability aligns with the persistence of a large-scale englacial fold along the shear margin of the glacier (which we traced for 360 km), where analysis of the fold axis and orientation provide little evidence for migration of the shear margin. These discoveries represent a novel understanding of one of the largest glacial ice-shelf systems in Antarctica, where findings are unlike those found in many other areas of the continent, where englacial stratigraphic analysis has revealed more dynamic ice flow changes within onset zones, and along glacier shear margins.

430 We do however note some evidence of change within the catchment, in ILCI Zone 2, where slow-flowing ice feeds into the main trunk of Lambert Glacier from the north (point T on Fig. 3). Here, we find disrupted and discontinuous englacial layers which do not correspond to changes in the glacial bed, or indeed current ice flow conditions at the glacier surface. We therefore suggest that the disturbed englacial stratigraphic layers here evidence previously enhanced ice flow, within a former tributary of the Lambert Glacier. Whilst this finding will have some implication for former ice flux calculations, it is possibly more notable for future forecasting simulations, as the tributary could provide a conduit for flow again in the future, if ice conditions were to change as a result of internal or external forcings (i.e. climate warming).

440 In Zone 3 more continuous englacial layering is recorded again, despite ice having moved through a region of fast ice flow, within Zone 2, to get to Zone 3. Whilst changes in the bed, and a resultant relaxation in ice flow could account for this transition in englacial layering, we hypothesise that these changes in ILCI may also reflect changing thermal regimes, as ice flows through the catchment and transitions from fast, warm flow, to colder, slower flow. Whilst similar thermal changes may be at play in Zone 4, where in this case, warmer ice could be reducing the amplitude of englacial layer undulations and masking continuous englacial layering (resulting in the unexpectedly low ILCI values we record), we suggest that our ILCI readings could also be impacted by RES attenuation in thick ice flows, which generate weaker reflections, despite any real change in the englacial architecture.

Our findings highlight the applicability of ILCI tracing over large outlet glacier systems, and the importance of characterising englacial architecture across a catchment, in order to more fully understand the past and present flow conditions of the Antarctic Ice Sheet. Our results constrain key ice dynamics and provide flow information which will improve estimates of future mass-



450 balance and sea level rise estimates. This is critical for global forecasting, especially as East Antarctica contains a significant potential contribution to sea level rise.

Data availability

All datasets used in this paper are freely available online. The UK Polar Airborne Geophysics Data Portal hosts SEGYY files of
455 RES data for the AGAP-N survey (<https://www.bas.ac.uk/project/nagdp/>) whilst the National Snow and Ice Data Centre contains BedMachine (Morlighem et al., 2020) data (containing bed elevation, bed topography and ice thickness information): <http://nsidc.org/data/nsidc-0756> as well as ice-surface velocity data from MEaSURES (Mouginot et al., 2019): <https://nsidc.org/data/nsidc-0754/versions/1>. We use the freely available Quantarctica data set (<https://www.npolar.no/quantarctica/>) to view and interrogate RADARSAT mosaic imagery in QGIS
460 (<https://www.qgis.org/en/site/>).

Author contribution.

R.J.S carried out the analysis, interpretation, and drafting of the manuscript, with contributions from all co-authors to interpretation and editing. F.N. advised on the ILCI code.
465

Competing interests.

The authors declare that they have no conflict of interest.

Acknowledgments.

470 We would like to thank the AGAP team for collecting the RES data across the Lambert Glacier system, and the British Antarctic Survey (BAS) for making the data freely available and accessible online. We are also grateful for the collection and continual development of freely available datasets and data interrogation platforms like BedMachine, MEaSURES, Quantarctica and QGIS. This paper was inspired by the AntArchitecture Action Group of the Scientific Committee for Antarctic Research and the RINGS Action Group of the Scientific Committee for Antarctic Research. R.J.S. was supported by
475 the National Environmental Research Council (NERC)-funded ONE Planet Doctoral Training Partnership (NE/S007512/1), hosted jointly by Newcastle and Northumbria Universities. F.N. acknowledges support from the Agencia Nacional de Investigación y Desarrollo (ANID) Programa Becas de Doctorado en el Extranjero, Beca Chile, for a doctoral scholarship.

References

Allison, I.: The mass budget of the Lambert Glacier drainage basin, Antarctica, *Journal of Glaciology*, 22, 223-235,
480 <https://doi.org/10.3189/S0022143000014222>, 1979.



- Ashmore, D. W., Bingham, R. G., Ross, N., Siegert, M., Jordan, T. A., and Mair, D. W. F.: Englacial Architecture and Age-Depth Constraints Across the West Antarctic Ice Sheet, *Geophysical Research Letters*, 47, <https://doi.org/10.1029/2019GL086663>, 2020.
- 485 Bassis, J. N., Coleman, R., Fricker, H. A., and Minster, J. B.: Episodic propagation of a rift on the Amery Ice Shelf, East Antarctica, *Geophysical Research Letters*, 32, <https://doi.org/10.1029/2004GL022048>, 2005.
- Bell, R., Blankenship, D., Finn, C. A., Morse, D., Scambos, T., Brozena, J., and Hodge, S.: Influence of subglacial geology on the onset of a West Antarctic ice stream from aerogeophysical observations, *Nature*, 394, 58-62, <https://doi.org/10.1038/27883>, 1998.
- 490 Bell, R. E., Ferraccioli, F., Creyts, T. T., Braaten, D., Corr, H., Das, I., Damaske, D., Frearson, N., Jordan, T., Rose, K., Studinger, M., and Wolovick, M.: Widespread Persistent Thickening of the East Antarctic Ice Sheet by Freezing from the Base, *Science*, 331, 1592-1595, <https://www.science.org/doi/10.1126/science.1200109>, 2011.
- Bell, R. E., Studinger, M., Fahnestock, M. A., and Shuman, C. A.: Tectonically controlled subglacial lakes on the flanks of the Gamburtsev Subglacial Mountains, East Antarctica, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2005GL025207>, 2006.
- 495 Bell, R. E., Studinger, M., Shuman, C. A., Fahnestock, M. A., and Joughin, I.: Large subglacial lakes in East Antarctica at the onset of fast-flowing ice streams, *nature*, 445, 904-907, <https://doi.org/10.1038/nature05554>, 2007.
- Bindschadler, R., Bamber, J., and Anandakrishnan, S.: Onset of streaming flow in the Siple Coast region, West Antarctica, *The West Antarctic Ice Sheet: Behavior and Environment*, 77, 123-136, <https://doi.org/10.1029/AR077p0123>, 2001.
- Bindschadler, R., Choi, H., and Collaborators, A.: High-resolution Image-derived Grounding and Hydrostatic Lines for the Antarctic Ice Sheet., Boulder, Colorado, USA: National Snow and Ice Data Center., 2011.
- 500 Bingham, R. G., Rippin, D. M., Karlsson, N. B., Corr, H. F., Ferraccioli, F., Jordan, T. A., Le Brocq, A. M., Rose, K. C., Ross, N., and Siegert, M. J.: Ice-flow structure and ice dynamic changes in the Weddell Sea sector of West Antarctica from radar-imaged internal layering, *Journal of Geophysical Research: Earth Surface*, 120, 655-670, <https://doi.org/10.1002/2014JF003291>, 2015.
- 505 Bingham, R. G. and Siegert, M. J.: Radio-echo sounding over polar ice masses, *Journal of Environmental and Engineering Geophysics*, 12, 47-62, <https://doi.org/10.2113/JEEG12.1.47>, 2007a.
- Bingham, R. G., Siegert, M. J., Young, D. A., and Blankenship, D. D.: Organized flow from the South Pole to the Filchner-Ronne ice shelf: An assessment of balance velocities in interior East Antarctica using radio-echo sounding data, *Journal of Geophysical Research-Earth Surface*, 112, <https://doi.org/10.1029/2006JF000556>, 2007b.
- 510 Bodart, J. A., Bingham, R. G., Ashmore, D. W., Karlsson, N. B., Hein, A., and Vaughan, D. G.: Age-depth stratigraphy of Pine Island Glacier inferred from airborne radar and ice-core chronology, *Journal of Geophysical Research: Earth Surface*, 126, e2020JF005927, <https://doi.org/10.1029/2020JF005927>, 2021.



- 515 Bons, P. D., Jansen, D., Mundel, F., Bauer, C. C., Binder, T., Eisen, O., Jessell, M. W., Llorens, M.-G., Steinbach, F., and Steinhage, D.: Converging flow and anisotropy cause large-scale folding in Greenland's ice sheet, *Nature communications*, 7, 1-6, <https://doi.org/10.1038/ncomms11427>, 2016.
- Bryant, M., Mantelli, E., Suckale, J., Castelletti, D., Seroussi, H., Siegert, M., and Schroeder, D.: Observational constraints from englacial layers on fast flow initiation of a West-Antarctic ice stream, 2019.
- Conway, H., Catania, G., Raymond, C., Gades, A., Scambos, T., and Engelhardt, H.: Switch of flow direction in an Antarctic ice stream, *Nature*, 419, 465-467, <https://doi.org/10.1038/nature01081>, 2002.
- 520 Cooper, M. A., Jordan, T. M., Siegert, M. J., and Bamber, J. L.: Surface Expression of Basal and Englacial Features, Properties, and Processes of the Greenland Ice Sheet, *Geophysical Research Letters*, 46, 783-793, <https://doi.org/10.1029/2018GL080620>, 2019.
- Cui, X. B., Du, W. J., Xie, H., and Sun, B.: The ice flux to the Lambert Glacier and Amery Ice Shelf along the Chinese inland traverse and implications for mass balance of the drainage basins, East Antarctica, *Polar Research*, 39, <https://doi.org/10.33265/polar.v39.3582>, 2020.
- 525 Dawson, E. J., Schroeder, D. M., Chu, W., Mantelli, E., and Seroussi, H.: Ice mass loss sensitivity to the Antarctic ice sheet basal thermal state, *Nature communications*, 13, 1-9, <https://doi.org/10.1038/s41467-022-32632-2>, 2022.
- Ely, J. C. and Clark, C. D.: Flow-stripes and foliations of the Antarctic ice sheet, *Journal of Maps*, 12, 249-259, <https://doi.org/10.1080/17445647.2015.1010617>, 2016.
- 530 Ferraccioli, F., Finn, C. A., Jordan, T. A., Bell, R. E., Anderson, L. M., and Damaske, D.: East Antarctic rifting triggers uplift of the Gamburtsev Mountains, *Nature*, 479, 388-U139, <https://doi.org/10.1038/nature10566>, 2011.
- Franke, S., Bons, P. D., Westhoff, J., Weikusat, I., Binder, T., Streng, K., Steinhage, D., Helm, V., Eisen, O., and Paden, J. D.: Holocene ice-stream shutdown and drainage basin reconfiguration in northeast Greenland, *Nature Geoscience*, <https://doi.org/10.1038/s41561-022-01082-2>, 2022a.
- 535 Franke, S., Jansen, D., Binder, T., Dörr, N., Helm, V., Paden, J., Steinhage, D., and Eisen, O.: Bed topography and subglacial landforms in the onset region of the Northeast Greenland Ice Stream, *Annals of Glaciology*, 61, 143-153, <https://doi.org/10.1017/aog.2020.12>, 2020.
- Franke, S., Jansen, D., Binder, T., Paden, J. D., Dörr, N., Gerber, T. A., Miller, H., Dahl-Jensen, D., Helm, V., and Steinhage, D.: Airborne ultra-wideband radar sounding over the shear margins and along flow lines at the onset region of the Northeast Greenland Ice Stream, *Earth System Science Data*, 14, 763-779, <https://doi.org/10.5194/essd-14-763-2022>, 2022b.
- 540 Frémand, A. C., Bodart, J. A., Jordan, T. A., Ferraccioli, F., Robinson, C., Corr, H. F., Peat, H. J., Bingham, R. G., and Vaughan, D. G.: British Antarctic Survey's Aerogeophysical Data: Releasing 25 Years of Airborne Gravity, Magnetic, and Radar Datasets over Antarctica, *Earth System Science Data Discussions*, <https://doi.org/10.5194/essd-14-3379-2022>, 2022.
- 545



- 550 Fretwell, P., Pritchard, H. D., Vaughan, D. G., Bamber, J. L., Barrand, N. E., Bell, R., Bianchi, C., Bingham, R. G., Blankenship, D. D., Casassa, G., Catania, G., Callens, D., Conway, H., Cook, A. J., Corr, H. F. J., Damaske, D., Damm, V., Ferraccioli, F., Forsberg, R., Fujita, S., Gim, Y., Gogineni, P., Griggs, J. A., Hindmarsh, R. C. A., Holmlund, P., Holt, J. W., Jacobel, R. W., Jenkins, A., Jokat, W., Jordan, T., King, E. C., Kohler, J., Krabill, W., Riger-Kusk, M., Langley, K. A., Leitchenkov, G., Leuschen, C., Luyendyk, B. P., Matsuoka, K., Mouginot, J., Nitsche, F. O., Nogi, Y., Nost, O. A., Popov, S. V., Rignot, E., Rippin, D. M., Rivera, A., Roberts, J., Ross, N., Siegert, M. J., Smith, A. M., Steinhage, D., Studinger, M., Sun, B., Tinto, B. K., Welch, B. C., Wilson, D., Young, D. A., Xiangbin, C., and Zirizzotti, A.: Bedmap2: improved ice bed, surface and thickness datasets for Antarctica, *Cryosphere*, 7, 375-393, <https://doi.org/10.5194/tc-7-375-2013>, 2013.
- 555 Fricker, H. A., Arndt, P., Brunt, K. M., Datta, R. T., Fair, Z., Jasinski, M. F., Kingslake, J., Magruder, L. A., Moussavi, M., Pope, A., Spergel, J. J., Stoll, J. D., and Wouters, B.: ICESat-2 Meltwater Depth Estimates: Application to Surface Melt on Amery Ice Shelf, East Antarctica, *Geophysical Research Letters*, 48, <https://doi.org/10.1029/2020GL090550>, 2021.
- 560 Fricker, H. A. and Padman, L.: Thirty years of elevation change on Antarctic Peninsula ice shelves from multitemission satellite radar altimetry, *Journal of Geophysical Research-Oceans*, 117, <https://doi.org/10.1029/2011JC007126>, 2012.
- Fricker, H. A., Warner, R. C., and Allison, I.: Mass balance of the Lambert Glacier-Amery Ice Shelf system, East Antarctica: a comparison of computed balance fluxes and measured fluxes, *Journal of Glaciology*, 46, 561-570, <https://doi.org/10.3189/172756500781832765>, 2000.
- 565 Gardner, A. S., Moholdt, G., Scambos, T., Fahnestock, M., Ligtenberg, S., van den Broeke, M., and Nilsson, J.: Increased West Antarctic and unchanged East Antarctic ice discharge over the last 7 years, *Cryosphere*, 12, 521-547, <https://doi.org/10.5194/tc-12-521-2018>, 2018.
- 570 Glasser, N. F., Jennings, S. J. A., Hambrey, M. J., and Hubbard, B.: Origin and dynamic significance of longitudinal structures ("flow stripes") in the Antarctic Ice Sheet, *Earth Surface Dynamics*, 3, 239-249, <https://doi.org/10.5194/esurf-3-239-2015>, 2015.
- Gong, Y., Cornford, S. L., and Payne, A. J.: Modelling the response of the Lambert Glacier-Amery Ice Shelf system, East Antarctica, to uncertain climate forcing over the 21st and 22nd centuries, *Cryosphere*, 8, 1057-1068, <https://doi.org/10.5194/tc-8-1057-2014>, 2014.
- 575 Grinsted, A., Hvidberg, C. S., Lilien, D. A., Rathmann, N. M., Karlsson, N. B., Gerber, T., Kjær, H. A., Vallelonga, P., and Dahl-Jensen, D.: Accelerating ice flow at the onset of the Northeast Greenland Ice Stream, *Nature communications*, 13, 1-4, <https://doi.org/10.1038/s41467-022-32999-2>, 2022.
- Holschuh, N., Christianson, K., and Anandakrishnan, S.: Power loss in dipping internal reflectors, imaged using ice-penetrating radar, *Annals of Glaciology*, 55, 49-56, <https://doi.org/10.3189/2014AoG67A005>, 2014.



- 580 Holschuh, N., Lilien, D., and Christianson, K.: Thermal weakening, convergent flow, and vertical heat transport in the
Northeast Greenland Ice Stream shear margins, *Geophysical Research Letters*, 46, 8184-8193,
<https://doi.org/10.1029/2019GL083436>, 2019.
- Holschuh, N., Lilien, D., and Christianson, K. A.: Estimating the Heat Production and Distribution across Ice-Stream
Shear Margins Using Surface Velocities, 2017, C41C-1251.
- Howat, I. M., Porter, C., Smith, B. E., Noh, M.-J., and Morin, P.: The reference elevation model of Antarctica, *The
Cryosphere*, 13, 665-674, <https://doi.org/10.5194/tc-13-665-2019>, 2019.
- 585 Jacobel, R., Scambos, T., Nereson, N., and Raymond, C.: Changes in the margin of Ice Stream C, Antarctica, *Journal of
Glaciology*, 46, 102-110, <https://doi.org/10.3189/172756500781833485>, 2000.
- Jezek, K. C.: Glaciological properties of the Antarctic ice sheet from RADARSAT-1 synthetic aperture radar imagery,
Annals of Glaciology, Vol 29, 1999, 29, 286-290, <https://doi.org/10.3189/172756499781820969>, 1999.
- Jordan, J. R., Gudmundsson, G. H., Stokes, C., Jamieson, S., Jenkins, A., and Miles, B.: What's Cooking in Antarctica?
A modeling study of Cook Ice Shelf, East Antarctica, *AGUFM*, 2018, C31C-1527, 2018.
- 590 Karlsson, N. B., Bingham, R. G., Rippin, D. M., Hindmarsh, R. C. A., Corr, H. F. J., and Vaughan, D. G.: Constraining
past accumulation in the central Pine Island Glacier basin, West Antarctica, using radio-echo sounding, *Journal of
Glaciology*, 60, 553-562, <https://doi.org/10.3189/2014JoG13J180>, 2014.
- Karlsson, N. B., Rippin, D. M., Bingham, R. G., and Vaughan, D. G.: A 'continuity-index' for assessing ice-sheet dynamics
595 from radar-sounded internal layers, *Earth and Planetary Science Letters*, 335, 88-94,
<https://doi.org/10.1016/j.epsl.2012.04.034>, 2012.
- Karlsson, N. B., Rippin, D. M., Vaughan, D. G., and Corr, H. F.: The internal layering of Pine Island Glacier, West
Antarctica, from airborne radar-sounding data, *Annals of Glaciology*, 50, 141-146,
<https://doi.org/10.3189/S0260305500250660>, 2009.
- 600 Leitchenkov, G., Belyatsky, B., and Kaminsky, V.: The age of rift-related basalts in East Antarctica,
<https://doi.org/10.1134/S1028334X18010051>, 2018, 11-14.
- Li, X., Rignot, E., Mouginot, J., and Scheuchl, B.: Ice flow dynamics and mass loss of Totten Glacier, East Antarctica,
from 1989 to 2015, *Geophysical Research Letters*, 43, 6366-6373, <https://doi.org/10.1002/2016GL069173>, 2016.
- Livingstone, S. J., Li, Y., Rutishauser, A., Sanderson, R. J., Winter, K., Mikucki, J. A., Bjornsson, H., Bowling, J. S., Chu,
605 W. N., Dow, C. F., Fricker, H. A., McMillan, M., Ng, F. S. L., Ross, N., Siegert, M. J., Siegfried, M., and Sole, A. J.:
Subglacial lakes and their changing role in a warming climate, *Nature Reviews Earth & Environment*, 3, 106-124,
<https://doi.org/10.1038/s43017-021-00246-9>, 2022.
- Luo, K., Liu, S., Guo, J., Wang, T., Li, L., Cui, X., Sun, B., and Tang, X.: Radar-Derived Internal Structure and Basal
Roughness Characterization along a Traverse from Zhongshan Station to Dome A, East Antarctica, *Remote Sensing*, 12,
610 1079, <https://doi.org/10.3390/rs12071079>, 2020.



- Mantelli, E., Bryant, M., Schroeder, D. M., Suckale, J., Castelletti, D., Räss, L., Seroussi, H. L., and Siegert, M. J.: Spatial distribution of englacial layer slope as a constraint on ice sheet basal conditions, 2019, C53B-1349.
- Matsuoka, K., MacGregor, J. A., and Pattyn, F.: Predicting radar attenuation within the Antarctic ice sheet, *Earth and Planetary Science Letters*, 359, 173-183, <https://doi.org/10.1016/j.epsl.2012.10.018>, 2012.
- 615 Miles, B., Stokes, C. R., Vieli, A., and Cox, N. J.: Rapid, climate-driven changes in outlet glaciers on the Pacific coast of East Antarctica, *Nature*, 500, 563-566, <https://doi.org/10.1038/nature12382>, 2013.
- Miles, B. W., Jordan, J. R., Stokes, C. R., Jamieson, S. S., Gudmundsson, G. H., and Jenkins, A.: Recent acceleration of Denman Glacier (1972–2017), East Antarctica, driven by grounding line retreat and changes in ice tongue configuration, *The Cryosphere*, 15, 663-676, <https://doi.org/10.5194/tc-15-663-2021>, 2021.
- 620 Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., Eisen, O., Ferraccioli, F., Forsberg, R., Fretwell, P., Goel, V., Greenbaum, J. S., Gudmundsson, H., Guo, J. X., Helm, V., Hofstede, C., Howat, I., Humbert, A., Jokat, W., Karlsson, N. B., Lee, W. S., Matsuoka, K., Millan, R., Mouginot, J., Paden, J., Pattyn, F., Roberts, J., Rosier, S., Ruppel, A., Seroussi, H., Smith, E. C., Steinhage, D., Sun, B., van den Broeke, M. R., van Ommen, T. D., van Wessem, M., and Young, D. A.: Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet, *Nature Geoscience*, 13, 132+, <https://doi.org/10.1038/s41561-019-0510-8>, 2020.
- 625 Morlighem, M., Seroussi, H., Larour, E., and Rignot, E.: Inversion of basal friction in Antarctica using exact and incomplete adjoints of a higher-order model, *Journal of Geophysical Research: Earth Surface*, 118, 1746-1753, <https://doi.org/10.1002/jgrf.20125>, 2013.
- Mouginot, J., Rignot, E., and Scheuchl, B.: Continent-Wide, Interferometric SAR Phase, Mapping of Antarctic Ice Velocity, *Geophysical Research Letters*, 46, 9710-9718, <https://doi.org/10.1029/2019GL083826>, 2019.
- 630 Ng, F. and Conway, H.: Fast-flow signature in the stagnated Kamb ice stream, West Antarctica, *Geology*, 32, 481-484, 2004.
- Parrenin, F., Cavitte, M. G., Blankenship, D. D., Chappellaz, J., Fischer, H., Gagliardini, O., Masson-Delmotte, V., Passalacqua, O., Ritz, C., and Roberts, J.: Is there 1.5-million-year-old ice near Dome C, Antarctica?, *The Cryosphere*, 11, 2427-2437, <https://doi.org/10.5194/tc-11-2427-2017>, 2017.
- 635 Pattyn, F. and Morlighem, M.: The uncertain future of the Antarctic Ice Sheet, *Science*, 367, 1331-1335, <https://www.science.org/doi/full/10.1126/science.aaz5487>, 2020.
- Pittard, M. L., Galton-Fenzi, B. K., Watson, C. S., and Roberts, J. L.: Future sea level change from Antarctica's Lambert-Amery glacial system, *Geophysical Research Letters*, 44, 7347-7355, <https://doi.org/10.1002/2017GL073486>, 2017.
- 640 Price, S. F., Bindschadler, R. A., Hulbe, C. L., and Blankenship, D. D.: Force balance along an inland tributary and onset to Ice Stream D, West Antarctica, *Journal of Glaciology*, 48, 20-30, <https://doi.org/10.3189/172756502781831539>, 2002.
- Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessem, M. J., and Morlighem, M.: Four decades of Antarctic Ice Sheet mass balance from 1979-2017, *Proceedings of the National Academy of Sciences of the United States of America*, 116, 1095-1103, <https://doi.org/10.1073/pnas.1812883116>, 2019.



- 645 Rippin, D. M., Siegert, M. J., and Bamber, J. L.: The englacial stratigraphy of Wilkes Land, East Antarctica, as revealed by internal radio-echo sounding layering, and its relationship with balance velocities, *Annals of Glaciology*, Vol 36, 36, 189-196, <https://doi.org/10.3189/172756403781816356>, 2003.
- Rose, K. C., Ferraccioli, F., Jamieson, S. S. R., Bell, R. E., Corr, H., Creyts, T. T., Braaten, D., Jordan, T. A., Fretwell, P. T., and Damaske, D.: Early East Antarctic Ice Sheet growth recorded in the landscape of the Gamburtsev Subglacial Mountains, *Earth Planet. Sci. Lett.*, 375, 1-12, <https://doi.org/10.1016/j.epsl.2013.03.053>, 2013.
- 650 Ross, N., Corr, H., and Siegert, M.: Large-scale englacial folding and deep-ice stratigraphy within the West Antarctic Ice Sheet, *Cryosphere*, 14, 2103-2114, <https://doi.org/10.5194/tc-14-2103-2020>, 2020.
- Schroeder, D. M., Bingham, R. G., Blankenship, D. D., Christianson, K., Eisen, O., Flowers, G. E., Karlsson, N. B., Koutnik, M. R., Paden, J. D., and Siegert, M. J.: Five decades of radioglaciology, *Annals of Glaciology*, 61, 1-13, 2020.
- 655 Sergienko, O. V., Creyts, T. T., and Hindmarsh, R. C. A.: Similarity of organized patterns in driving and basal stresses of Antarctic and Greenland ice sheets beneath extensive areas of basal sliding, *Geophysical Research Letters*, 41, 3925-3932, <https://doi.org/10.1002/2014GL059976>, 2014.
- Shepherd, A., Ivins, E., Rignot, E., Smith, B., van den Broeke, M., Velicogna, I., Whitehouse, P., Briggs, K., Joughin, I., Krinner, G., Nowicki, S., Payne, T., Scambos, T., Schlegel, N., Geruo, A., Agosta, C., Ahlstrom, A., Babonis, G., Barletta, V., Blazquez, A., Bonin, J., Csatho, B., Cullather, R., Felikson, D., Fettweis, X., Forsberg, R., Gallee, H., Gardner, A., Gilbert, L., Groh, A., Gunter, B., Hanna, E., Harig, C., Helm, V., Horvath, A., Horwath, M., Khan, S., Kjeldsen, K. K., Konrad, H., Langen, P., Lecavalier, B., Loomis, B., Luthcke, S., McMillan, M., Melini, D., Mernild, S., Mohajerani, Y., Moore, P., Mougintot, J., Moyano, G., Muir, A., Nagler, T., Nield, G., Nilsson, J., Noel, B., Ootosaka, I., Pattle, M. E., Peltier, W. R., Pie, N., Rietbroek, R., Rott, H., Sandberg-Sorensen, L., Sasgen, I., Save, H., Scheuchl, B., Schrama, E., Schroder, L., Seo, K. W., Simonsen, S., Slater, T., Spada, G., Sutterley, T., Talpe, M., Tarasov, L., van de Berg, W. J., van der Wal, W., van Wessem, M., Vishwakarma, B. D., Wiese, D., Wouters, B., and Team, I.: Mass balance of the Antarctic Ice Sheet from 1992 to 2017, *Nature*, 558, 219-+, <https://doi.org/10.1038/s41586-018-0179-y>, 2018.
- 665 Siegert, M. J.: On the origin, nature and uses of Antarctic ice-sheet radio-echo layering, *Progress in physical geography*, 23, 159-179, <https://doi.org/10.1177/030913339902300201>, 1999.
- 670 Siegert, M. J. and Bamber, J. L.: Subglacial water at the heads of Antarctic ice-stream tributaries, *Journal of Glaciology*, 46, 702-703, <https://doi.org/10.3189/172756500781832783>, 2000.
- Siegert, M. J., Payne, A. J., and Joughin, I.: Spatial stability of Ice Stream D and its tributaries, West Antarctica, revealed by radio-echo sounding and interferometry, *Annals of Glaciology*, 37, 377-382, <https://doi.org/10.3189/172756403781816022>, 2003.
- 675 Stokes, C. R., Abram, N. J., Bentley, M. J., Edwards, T. L., England, M. H., Foppert, A., Jamieson, S. S., Jones, R. S., King, M. A., and Lenaerts, J.: Response of the East Antarctic Ice Sheet to past and future climate change, *Nature*, 608, 275-286, <https://doi.org/10.1038/s41586-022-04946-0>, 2022.



- 680 Sutter, J., Fischer, H., and Eisen, O.: Investigating the internal structure of the Antarctic ice sheet: the utility of isochrones for spatiotemporal ice-sheet model calibration, *The Cryosphere*, 15, 3839-3860, <https://doi.org/10.5194/tc-15-3839-2021>, 2021.
- Van Der Veen, C. J., Jezek, K. C., and Stearns, L.: Shear measurements across the northern margin of Whillans Ice Stream, *Journal of Glaciology*, 53, 17-29, <https://doi.org/10.3189/172756507781833929>, 2007.
- 685 Wen, J. H., Wang, Y. F., Liu, J. Y., Jezek, K. C., Huybrechts, P., Csatho, B. M., Farness, K. L., and Bo, S.: Mass budget of the grounded ice in the Lambert Glacier-Amery Ice Shelf system, *Annals of Glaciology*, Vol 48, 48, 193-+, <https://doi.org/10.3189/172756408784700644>, 2008.
- Winsborrow, M. C., Clark, C. D., and Stokes, C. R.: What controls the location of ice streams?, *Earth-Science Reviews*, 103, 45-59, <https://doi.org/10.1016/j.earscirev.2010.07.003>, 2010.
- 690 Winter, A., Steinhage, D., Creyts, T. T., Kleiner, T., and Eisen, O.: Age stratigraphy in the East Antarctic Ice Sheet inferred from radio-echo sounding horizons, *Earth System Science Data*, 11, 1069-1081, <https://doi.org/10.5194/essd-11-1069-2019>, 2019.
- Winter, K., Woodward, J., Dunning, S. A., Turney, C. S. M., Fogwill, C. J., Hein, A. S., Golledge, N. R., Bingham, R. G., Marrero, S. M., Sugden, D. E., and Ross, N.: Assessing the continuity of the blue ice climate record at Patriot Hills, Horseshoe Valley, West Antarctica, *Geophysical Research Letters*, 43, 2019-2026, <https://doi.org/10.1002/2015GL066476>, 2016.
- 695 Winter, K., Woodward, J., Ross, N., Dunning, S. A., Bingham, R. G., Corr, H. F. J., and Siegert, M. J.: Airborne radar evidence for tributary flow switching in Institute Ice Stream, West Antarctica: Implications for ice sheet configuration and dynamics, *Journal of Geophysical Research-Earth Surface*, 120, 1611-1625, <https://doi.org/10.1002/2015JF003518>, 2015.
- Wrona, T., Wolovick, M. J., Ferraccioli, F., Corr, H., Jordan, T., and Siegert, M. J.: Position and variability of complex structures in the central East Antarctic Ice Sheet, *Geological Society, London, Special Publications*, 461, 113-129, <https://doi.org/10.1144/SP461.12>, 2018.
- 700 Yu, J. Y., Liu, H. X., Jezek, K. C., Warner, R. C., and Wen, J. H.: Analysis of velocity field, mass balance, and basal melt of the Lambert Glacier-Amery Ice Shelf system by incorporating Radarsat SAR interferometry and ICESat laser altimetry measurements, *Journal of Geophysical Research-Solid Earth*, 115, <https://doi.org/10.1029/2010JB007456>, 2010.
- 705 Zwally, H. J., Giovinetto, M. B., Beckley, M. A., and Saba, J. L.: Antarctic and Greenland drainage systems, GSFC cryospheric sciences laboratory. 2012a.