

1

2 **Assessment of the impact of dam reservoirs on river ice cover - an**

3 **example from the Carpathians (central Europe)**

4 Maksymilian Fuks¹

5 ¹Department of Geoenvironmental Research, Institute of Geography and Spatial Organization, Polish Academy of Sciences,
6 Poland

7 *Correspondence to:* Maksymilian Fuks (fuksmaksymilian@twarda.pan.pl)

8

9 **Abstract.** This paper presents a method for determining the impact of dam reservoirs on the ice cover of rivers downstream
10 of below their locations based on a long measurement period (1950–2020) and synthetic-aperture radar (SAR) data. Two
11 rivers and two sets of dam reservoirs located in the Carpathian Mountains (central Europe) were selected for this study. In
12 order to estimate the influence of ~~their~~ reservoirs, a logistic regression model was built to describe the relationship between
13 the course of air temperature and the occurrence of ice cover (i.e., total ice cover and border ice) at water gauge cross
14 sections upstream and downstream of above and below the reservoirs. The influence of reservoirs was then defined as the
15 differences between the number of days with ice cover values predicted from air temperature and those observed at the water
16 gauge cross sections. Additionally, the extent of the impact of the reservoirs was estimated based on SAR data (Sentinel-1)
17 by identifying river sections downstream of below the reservoirs on which total ice cover did not form despite the persistence
18 of very low air temperatures. This study demonstrates that dam reservoirs are an important factor in transforming the ice
19 regime of rivers. The occurrence of ice cover as a result of reservoir operations could decrease by over 80% in the sections
20 immediately downstream of reservoirs. The impact of the reservoir on river ice cover diminishes as the distance from the
21 reservoir increases. We found that decreases in the incidence of ice cover as a result of their operation could exceed 80% in
22 the sections immediately below their location, and that this reduction decreased as the distance from the reservoir increases.
23 Using According to SAR data, it was estimated that total ice cover did not form in sections 26–60 kilometers downstream
24 of below the reservoirs, despite the presence of favorable thermal conditions. Based on the results of the study presented here,
25 it can be assumed that in areas where many dam reservoirs are located, the ice regime of rivers is significantly transformed,
26 which should be taken into account when studying river ice cover. Based on the research results presented here, it is
27 reasonable to assume that the rapid increase in the number of dam reservoirs worldwide in the second half of the 20th century
28 is an important factor transforming the ice regime of rivers. This study also demonstrates that the logistic regression model
29 and SAR data are useful tools for assessing the impact of dam reservoirs on river ice cover.

30

31 **Keywords**

— sformatowano: Czcionka: Nie Pogrubienie, Angielski (Stany Zjednoczone)

32 Dam reservoirs, river ice cover, logistic regression, synthetic aperture radar (SAR), Carpathians

34 1. Introduction

35 Over the course of the 20th and 21st centuries, human impact on the natural environment has increased significantly. The
36 transformation of the environment and its effects, previously occurring on a local scale, have now begun to be observed on a
37 regional and global scale. The rapid increase in the impact of human activity since the 1950s has affected both biotic and
38 abiotic aspects of the environment and has been dubbed the Great Acceleration (Lewis and Maslin, 2015). The section of the
39 environment undergoing the most significant change is the terrestrial part of the Earth's cryosphere. This is mainly because
40 of climate change and a significant increase in air temperature, particularly in cold areas (Fox-Kemper et al., 2021). Since the
41 second half of the 20th century, there has been a decline in the extent and mass of ice sheets, mountain glaciers, and snow
42 cover, as well as notable melting of permafrost (Fox -Kemper et al., 2021). Significant changes have also been observed in
43 the river ice phenomena, which due to their periodic nature and the relatively small volume of river ice, are particularly
44 sensitive to climatic variability and human influence (Newton and Mullan, 2021).

45 Dam reservoirs are an example of anthropogenic elements of the geographic environment that can significantly affect the
46 ice regime of rivers. Such structures change the conditions of ice processes in rivers, mainly through alterations in the flow
47 volume and change in the thermodynamics of rivers downstream reservoir location in winter (e.g., Starosolszky, 1990;
48 Takács et al., 2013; Maheu et al., 2014; Takács and Kern, 2015; Pawłowski, 2015; Apsite et al., 2016; Chang et al., 2016). In
49 cold and temperate climate zones, due to the occurrence of thermal stratification in the reservoir and the release of bottom
50 hypolimnion waters, the operations of the reservoir cool the river downstream in summer and warm it in winter, which
51 translates into a reduction in the annual amplitude of water temperature. This effect is most pronounced immediately
52 downstream of the reservoir and decreases with increasing distance (Cai et al. 2018). Higher river water temperatures
53 downstream of the reservoirs during winter have been recorded for both large and relatively small reservoirs (<20m, Maheu
54 et al. 2014). The increase in water temperature downstream of the reservoir may impede the phase transformation and ice
55 cover formation. Reservoirs also act as barriers to moving ice forms. Because ice and ice floes are intercepted, they
56 contribute to the formation of the total ice cover downstream of the dam to a lesser extent (Starosolszky, 1990). Moreover,
57 reservoirs capture ice and sediment which can also result in their reduced amount downstream of the dam. This, in turn,
58 leads to fewer nucleating particles which potentially affects the freezing process (Michel, 1961; Osterkamp and Gilfilian,
59 1975; Carlson, 1981; Chen et al. 2023). Reservoirs also modify the discharge hydrograph: by increasing or decreasing the
60 volume of flow, they can affect the timing of river ice cover formation and breakup (Houkuna et al., 2022).

61 As a result of dam reservoir operation, an ice regime can be transformed over sections of several to even several hundred
62 As a result of dam reservoir operation, an ice regime can be transformed over sections of several to even several hundred
63 As a result of dam reservoir operation, an ice regime can be transformed over sections of several to even several hundred
64 As a result of dam reservoir operation, an ice regime can be transformed over sections of several to even several hundred
65 of dam reservoir operation, an ice regime can be transformed over sections of several to even several hundred kilometers,

Sformatowano: Wcięcie: Pierwszy wiersz: 0.5 cm

66 sometimes causing the complete disappearance of ice phenomena (Maheu et al., 2014; Pawłowski, 2015; Chang et al., 2016).
67 Pawłowski (2015) showed that the construction of the Wloclawek dam reservoir on the Vistula River (Poland, Central
68 Europe) resulted in a 26% reduction in the duration of ice cover and a 47% reduction in all ice phenomena downstream
69 of/below its location. Chang et al. (2016) showed that the construction of the Longyangxia and Liujiaxia reservoirs on the
70 Yellow River (northern China) resulted in a reduction in the ice cover duration of the river by 8–33 days and a reduction in
71 the thickness of the ice cover by 16–25 cm. This effect has also been noted in small rivers with small and medium-sized
72 reservoirs. For example, Maheu et al. (2014) showed that the operation of small dam reservoirs in eastern Canada resulted in
73 change in the thermal regime of rivers and the disappearance of ice phenomena in rivers over a distance of 0.3–2.5 km.

74 Despite the significant role of dam reservoirs in transforming the ice regimes of rivers, the problem is relatively under-
75 researched. Most studies have focused on comparing ice parameters in the riverbed before and after reservoir construction on
76 large lowland rivers for which long measurement series are available. Despite the significant role of dam reservoirs in
77 transforming the ice regimes of rivers, the problem is relatively poorly recognized in many ways. Most studies focus on
78 assessing the impact of individual reservoirs, neglecting the regional and global aspects of this issue. It has been estimated
79 that there are more than 8,000 dam reservoirs in ice regime areas, most of which are located in central and northern Europe,
80 eastern Asia and the central section of North America (Fukś, 2023). The main difficulty in assessing the impact of dam
81 reservoirs on river ice cover comes in distinguishing whether changes are due to operational or climatic factors. So far, in
82 order to study reservoirtheir impact, ice phenomena have been compared during periods with similar thermal conditions
83 (usually determined by the average air temperature of winter) before and after the construction of reservoirs (e.g., Takács
84 et al., 2013; Pawłowski, 2015; Chang et al., 2016). This approach makes it possible to assess the impact of reservoirs only on
85 the basis of selected single years in which the relevant thermal and hydrological conditions occurred. However, in order to
86 accurately characterize the role of reservoirs in transforming the ice regime of rivers, it is necessary to conduct accurate,
87 quantitative assessment of thetheir impact of reservoirs on river ice cover for long periods. Another issue is the small number
88 of studies based on remote sensing data (including radar) for relatively small mountain rivers, where the course of ice
89 processes is poorly understood (Thellman et al. 2021). This results in poor understanding of the extent of the influence of
90 dam reservoirs on river ice cover (especially small mountain rivers), making it difficult to estimate their role on a regional or
91 global scale.

92 The main objective of this study is to determine the impact of Carpathian dam reservoirs on the ice cover of rivers
93 downstream of belowtheir locations based on long observation series and radar (SAR) imaging. Specific objectives include:
94 (1) develop and present a method to quantitatively assess the impact of dam reservoirs on the duration of ice cover based on
95 measurement data from water gauge cross sections; (2) estimate the extent of the impact of dams-of their impact based on
96 satellite radar imagery (SAR) and assess the feasibility of using Sentinel-1 satellite radar data to determine the extent of
97 reservoirs' impact on the ice cover of this type of river. The essential hypothesis tested here is that dam reservoirs, at local
98 and regional scales, have a greater impact on transformations in the occurrence of river ice cover than climate change.

2 Study area, data, and methods

2.1 Study area

The study was based on two sets of dam reservoirs located in the Outer Western Carpathians (Solina-Myczkowce) and the Central Western Carpathians (Dunajec- Sromowce Wyżne) in central Europe (Fig. 1). These reservoirs are located on two second-order mountain rivers whose sources are in the higher reaches of the Carpathians: the Dunajec and the San rivers. The average annual flow of the Dunajec at its mouth averages ~~is more than over~~ $84 \text{ m}^3 \text{ s}^{-1}$ with a catchment area of 6735 km^2 , while that of the San averages $134 \text{ m}^3 \text{ s}^{-1}$ with a catchment area of $16,824 \text{ km}^2$ (Punzet, 1991). The width of the rivers in the sections ~~downstream of below~~ the dams varies from 30 m to more than 100 m. The basic characteristics of both sets of reservoirs are shown in Table 1.

Table 1: Characteristics of the studied reservoirs.

Reservoir	River	Year of completion	Total capacity [million m ³]	Type of dam	Damming height [m]	Average inflow during the winter [m ³ s ⁻¹]	Average outflow during the winter [m ³ s ⁻¹]
Czorsztyn	Dunajec	1997	231,9	ground dam	52	20,6	20,4
Sromowce Wyżne	Dunajec	1994	7,42	ground dam	10		
Solina	San	1968	473,0	concrete dam	58	24,1	24,6
Myczkowce	San	1961	10,0	ground dam	15		


Source of data: Hennig et al. 1991; Bajorek et al. 2003

The two reservoir complexes studied ~~in~~ here consist of a main reservoir and an equalization reservoir. The Czorsztyn and Solina reservoirs are intended mainly for electricity production, and play a role in flood control, while the Sromowce and Myczkowce reservoirs serve as equalizing reservoirs for daily flow fluctuations caused by the operation of hydroelectric power plants. Both reservoirs significantly affect the thermal regime of the rivers ~~downstream of below~~ their locations. It has previously been shown that the operation of the hydroelectric power plant at the Solina reservoir significantly transforms the thermal regime of the Myczkowce reservoir, and consequently of the river ~~downstream below it~~, warming it in winter by about 2 °C (Lewinska and Lewinski, 1972). In the case of the Czorsztyn-Sromowce reservoir complex, studies have shown that ~~reservoir they~~ warm the temperature of the Dunajec waters ~~downstream of below~~ the reservoir by 1-2°C (Wiejaczka et al. 2015, Kędra and Wiejaczka, 2017). In addition, the synchronization between air and water temperatures in the river downstream of the reservoirs was also disrupted by the dam operation (Kędra and Wiejaczka, 2016). Both reservoirs transform the discharge of the river downstream of their locations, relative to natural conditions, due to the operation of hydroelectric power plants and equalization reservoirs. During the winter this is evidenced by the rise in river water levels downstream of the reservoir.

Sformatowano: Wcięcie: Pierwszy wiersz: 0.5 cm

125 The San River catchment area (Solina-Myczkowce reservoir complex) is characterized by low population density (<20
126 people/km²). No large cities are located along the river, and the entire upper part of the catchment area is located within the
127 Bieszczady National Park and is protected. Therefore, it can be assumed that the ice and thermal regime of this river is not
128 significantly impacted by human activity (except for the operation of the reservoir). In the case of the Dunajec River
129 (Czorsztyn-Sromowce reservoir complex), the population density is higher and reaches up to 200 people/km². The upper
130 part of the Dunajec River basin is located within the Tatra National Park and is protected. Several cities and dozens of
131 villages with tourist infrastructure (winter tourism, skiing, thermal pools) are located along the studied river and its
132 tributaries. Therefore, it can be assumed that the temperature of surface waters and ice regime may be affected to some
133 extent (for example, through the emission of thermal pollutants into rivers), but the scale of impact is difficult to estimate
134 due to the lack of data.

135 The main reasons for selecting these sites for the study were the good availability of hydrological and meteorological
136 data, as well as the location and characteristics of these reservoirs. In terms of their size, these reservoirs represent typical
137 facilities for ice cover areas. According to the GRanD database and calculations presented by Fuk^s (2023), in areas where
138 river ice cover occurs, reservoirs with dam heights ranging between 41-60 meters account for about 20% of all reservoirs
139 (Lehner et al. 2011). This allows the results obtained to be applied to other similar facilities and rivers. In addition, there are



not enough studies in the literature on ice phenomena in relatively small mountain rivers due to the lack of observational data, the difficulty of conducting observations with remote sensing data, and greater practical significance of large rivers (Thellman et al. 2021). In addition, a disturbance between the air and water temperatures of the river below the reservoirs caused by their operation has also been found (Kędra and Wiejaczka, 2016). Both sets of reservoirs also experience a transformation of river flow volume due to the operation of hydropower plants and equalization reservoirs, with respect to natural conditions.

Source: Map tiles by Stamen Design, under CC BY 4.0. Data by OpenStreetMap, under ODbL

2.2 Data and methods

In order to estimate the impact of dam reservoirs on river ice cover, data on the daily occurrence of ice cover over the period 1950–2020 were obtained at eight water gauge cross sections (four for each reservoir complex studied) (Figure 1). The occurrence of ice cover at a water gauge cross-section was defined as any occurrence of total ice cover (water surface completely covered by ice) or border ice (partial coverage of the water surface by ice). In both cases, one cross-section was located upstream of the reservoir (C1: 14.6 km, S1: 28.5 km), while the others were located downstream of the reservoir location, at distances ranging from several to tens of kilometers from the reservoir (C2: 1.8 km, C3: 22 km, C4: 52 km, S2: 11.7 km, S3: 33 km, S4: 80.5 km, tab. 2). Data on the occurrence of ice phenomena each day of the winter periods (November to March) in the 1950–1980 period were obtained from hydrological yearbooks published by the Polish State Hydrological and Meteorological Institute. These yearbooks were issued only in printed form, so for this study they were digitized by manually transcribing data on daily ice cover occurrence. Data on the occurrence of ice cover in the 1981–2020 period were obtained from the online public database of the Institute of Meteorology and Water Management - National Research Institute (IMGW-PIB, - Data availability statement <https://danepubliczne.imgw.pl/>). Data on the course of average daily air temperature at climatological stations were also obtained from the IMGW-PIB public database. For the Solina-Myczkowce reservoir complex, climatological data were obtained from five stations (Ts1, Ts2, Ts3, Ts4, Ts5), while for the Czorsztyn-Sromowce reservoir complex they were obtained from two (Tc1, Tc2, Tab. 2). The acquired data on the occurrence of ice cover were subjected to statistical analysis involving a comparison of the average duration of ice cover before and after the formation of the reservoirs at stations upstream of and downstream of reservoirs locations, as well as a comparison of the two studied rivers. Data from periods for which there were no observations, or in which observations were incomplete, were excluded from the statistical analysis.

Table 2: Characteristics of hydrological and climatological stations.

Station name	Geographical coordinates	Height above sea level [meters]	Description
C1	49° 29' 12.169" N 20° 3' 13.869" E	576	Hydrological station. Air temperature data from station Tc1 were used to model ice phenomena.

Sformatowano: Odstęp Przed: 0 pkt, Po: 0 pkt

Sformatowano: Odstęp Po: 6 pkt

— sformatowano: Czcionka: Pogrubienie

— sformatowano: Czcionka: 9 pkt

Sformatowano: Wyrównany do środka

Tabela sformatowana

— sformatowano: Czcionka: 9 pkt

— sformatowano: Czcionka: 9 pkt, Angielski (Stany Zjednoczone)

— sformatowano: Czcionka: 9 pkt, Kolor czcionki: Automatyczny

Sformatowano: Wyrównany do środka, Interlinia: Wielokrotnie 1.15 wrs

— sformatowano: Czcionka: 9 pkt, Kolor czcionki: Automatyczny

— sformatowano: Czcionka: 9 pkt

Sformatowano: Wyrównany do środka

— sformatowano: Czcionka: 9 pkt

— sformatowano: Czcionka: 9 pkt, Kolor czcionki: Automatyczny

Sformatowano: Wyrównany do środka

— sformatowano: Czcionka: 9 pkt

— sformatowano: Czcionka: 9 pkt

177 The acquired data on average daily air temperatures from climatological stations (Tc1, Tc2, Ts2, Ts4, Ts5) were analyzed
178 for trends. The non-parametric Mann-Kendall test was used to detect them, while the Theil-Sen estimator was used to
179 determine the magnitude of changes (Mann 1945, Theil 1950, Sen 1968, Kendall 1975). The trend incidence analysis was
180 carried out for entire periods where data were available, as well as for a uniform period for all stations (1982-2015). Results
181 at the p<0.05 level were accepted as statistically significant. The analysis was carried out for the average values of each
182 month and for the entire winter period (November to December).

183 In order to separate the effects of climate change and reservoir operations on the ice cover of the studied rivers, a logistic
184 regression method was used to model the relationship between air temperature and ice cover occurrence. Logistic regression
185 is a method that allows the classification of a dichotomous explanatory variable based on one or more explanatory variables.
186 The method was first proposed by Cox (1957) and is widely used in classification and prediction in natural science research.
187 Previously, it has been used in studies of river ice phenomena (Yang et al. 2020, Wu et al. 2021), the delineation of flood-
188 prone areas (Lee and Kim, 2021) or of areas susceptible to landslides (Ayalew and Yamagishi, 2005), among other
189 applications, but has not been used in natural studies of the effects of dam reservoirs. Logistic regression analysis is based on
190 the concept of odds that represent the ratio of the probability (p) that an event will occur and the probability that
191 the event will not occur, and which is expressed by equation (1).

$$192 \text{oddsOR} = \frac{p}{1-p} \quad (1)$$

194 In order to carry out a binary classification of the dependent variable on the basis of the continuous independent variable, a
195 logit transformation was applied by logarithmizing the odds ratio, as expressed by equation (2):

$$196 \text{Logit}(\text{oddsOR}) = \log\left(\frac{p}{1-p}\right) = \beta_0 + \beta_1 x_1 \quad (2)$$

197 where x_1 is the explanatory variable and β_0 and β_1 are regression coefficients that are estimated using the maximum
198 likelihood method. The probability of ice cover occurrence on a given day (p) is calculated using equation (3) and
199 classification is done by applying a threshold value of probability that separates the occurrence of ice cover from its absence:

$$200 p = \frac{1}{1 + e^{-(\beta_0 + \beta_1 x_1)}} \quad (3)$$

202 where x_1 is the average air temperature of the 14 days prior to each day. The average air temperature from the 14 days
203 preceding each modelled day was selected based on preliminary tests (not shown). These consisted of testing the predictive
204 ability (accuracy and area under the receiver operating characteristic curve – ROC AUC) of the logistic regression model for
205 the occurrence of ice cover based on the average daily air temperatures from 2 to 20 days prior to each modelled day. The
206 analyses showed that in most cases, taking the average from 14 days resulted in the best predictive ability of the models. The

207
208 The first stage of analysis required coding of the data from the water gauge stations, with a value of 1 assigned to each
209 day in which ice cover occurred and a value of 0 assigned if it did not for the period from November to March over the entire
210 study period (sometimes the period was shorter than 70 years due to data gaps). Each record was assigned an average air
211 temperature from 14 days before the modeled day determined from measurements from the nearest climatological station. In
212 some cases, data from several nearby stations were used simultaneously due to gaps in measurement data. Based on the
213 acquired database, a logistic regression model of the relationship between the course of the average air temperature from 14
214 days before the modeled day and the occurrence of ice cover in the period before the dam reservoirs were built for each
215 water gauge station. For the water gauges on the Dunajec River (C1, C2, C3, C4), data from the period 01.11.1957-
216 31.03.1990 were used as the set for learning the model, while for the San River, data from the period 01.11.1956-31.03.1967
217 (S1) or 14.01.1951-31.03.1967 (S2, S3, S4) were used, depending on the station. The data for building the model was
218 divided into a training set and a test set in the ratio of 70% to 30%. Model learning and hyperparameter optimizations (C -
219 Inverse of regularization strength) were carried out based on the training set. Stratified cross-validation was used to avoid
220 over-fitting the model on the training set, and the quality of the resulting models was determined by the prediction accuracy
221 and the value of the area under the ROC curve (receiver operating characteristic curve) on the test set (ROC-AUC value).
222 Based on the obtained models, for each adopted winter period, the probability of ice cover occurrence at the tested cross
223 sections was calculated based on the average air temperature, from which it was determined whether ice cover could occur
224 on a given day. Then, the ice cover occurrence data calculated from the model based on the course of the average air
225 temperature was compared with actual observations of ice cover at all stations. In this study, it was assumed that the
226 difference between the number of days with ice cover predicted by the model and the number of days with ice observed at
227 the stations was due to the operation of dam reservoirs. In order to validate these results, the results of modeling and
228 observations from stations upstream and downstream of the reservoir were compared. All calculations were
229 carried out using Python and the Scikit-learn library.

230 The period of January to February 2017 was chosen to determine the spatial extent of the impact of dam reservoirs on
231 river ice cover based on remote sensing data. The analysis of radar (SAR) data was aimed at determining the sections of the
232 studied rivers downstream of the reservoirs on which ice cover did not form despite the maintenance of favorable air
233 temperatures. Remote sensing data was used because it was not possible to estimate this parameter based on data from water
234 gauge stations, due to the small number of stations and large distances between them. The year 2017 was chosen due to the
235 persistence of very low air temperatures in the study areas during this period. ~~This choice was due to the persistence of very~~
236 ~~low air temperatures at measuring stations in the study areas during this period.~~ Many stations recorded the lowest average
237 January temperature since 2006 (-7.3°C) at this point, and occasionally, the air temperature reached as low as -20°C . It was
238 found that such air temperature resulted in the persistence of ice cover on the studied rivers before the construction of dam
239 reservoirs. Moreover, during this period, ice cover was observed at the water gauge cross sections upstream of ~~above~~ the two

240 reservoirs from the beginning of December to the end of February.

241 The occurrence of ice cover was determined on the basis of radar (SAR) images acquired by Sentinel-1 satellites. In the
242 The occurrence of ice cover was determined on the basis of radar (SAR) images acquired by Sentinel-1 satellites. In the
243 determined on the basis of radar (SAR) images acquired by Sentinel-1 satellites. In the first stage, the area of rivers (water
244 surface) downstream of the studied reservoirs was determined by manually creating polygons on the basis of aerial
245 photos and cloudless Sentinel-2 satellite images. River areas in the vicinity of hydraulic structures (bridges), narrow river
246 sections (< 30 m in width) and areas close to the banks (about 10 meters from the shore) were excluded from the analysis,
247 which made it possible to exclude pixels partially covering areas other than the water surface. Then, for the designated
248 polygons, Sentinel-1 SAR IW GRD imagery was acquired from five different days, for both studied rivers (Tab. 3)

249 .
250 .
251 **Table 3: Dates of acquisition of radar images for analysis.**

Area of analysis	Dates	Polarization
<u>Dunajec river - Czorsztyn-Sromowce</u> <u>reservoir complex</u>	<u>02.01.2017, 09.01.2017, 14.01.2017, 26.01.2017, 14.02.2017</u>	<u>VV, VH</u>
<u>San River - Solina-Myczkowce</u> <u>reservoir complex</u>	<u>09.01.2017, 16.01.2017, 21.01.2017, 28.01.2017, 14.02.2017</u>	<u>VV, VH</u>

252 Data from descending orbits in VV and VH polarization were used. This study used imagery provided by Google Earth
253 Engine, in which orbit metadata were updated, border noise was removed, thermal noise was removed, radiometric
254 calibration values were applied, and terrain correction was performed. After preprocessing, the data had a spatial resolution
255 of 10 meters. In order to classify the acquired images (water/ice), thresholds of the backscattering coefficient that separated
256 the occurrence of ice cover from water were determined for both studied rivers. This determination of the presence of ice
257 cover was made possible by the marked contrast between the two classes (ice/water) due to the significant effect of ice cover
258 presence on the backscattering coefficient of the microwave radiation beam (Stonevicius et al., 2022; Palomaki and Sproles,
259 2022). Consolidated ice tends to have significantly higher backscattering values than water, mainly due to the roughness of
260 ice's surface. For this purpose, the value of the backscattering coefficient was used for designated sections of rivers
261 completely covered by ice (January 14 for the Dunajec River and January 9 for the San River). These sections were
262 determined on the basis of cloudless images from the Sentinel-2 satellite. The thresholds of the values separating the two
263 classes were calculated separately for the two studied rivers according to equation (4):
264

$$265 \tau_{\sigma} = \sigma - s \quad (4)$$

Sformatowano: Odstęp Po: 6 pkt

– sformatowano: Czcionka: Pogrubienie

Sformatowano: Wcięcie: Pierwszy wiersz: 0 cm

– sformatowano: Czcionka: 9 pkt

Sformatowano: Wyrównany do środka

Tabela sformatowana

– sformatowano: Czcionka: 9 pkt

Sformatowano: Wyrównany do środka

– sformatowano: Czcionka: 9 pkt

Sformatowano: Wyrównany do środka


268 where σ is the average backscatter from selected sections of ice-covered rivers, and s is the average standard deviation of σ
269 (Sobiech and Dierking 2013). For the Dunajec River, the thresholds were set at -19.24 dB for VH polarization and -10.16
270 dB for VV polarization, while for the San River they were set at -21.16 for VH polarization and -11.55 for VV polarization.
271 The results were validated by comparing the classification results of Sentinel-1 radar imagery with optical imagery acquired
272 by Sentinel-2 satellite for the Dunajec area acquired on February 14, 2017.

273 In order to identify the causes of the transformation of the ice regime of the rivers resulting from the operations of the
274 dam reservoirs, data on daily water level and temperature, as well as flow volume at station C3 for the period 1984-2020
275 were obtained (IMGW-PIB, Data availability statement). The choice of the Dunajec River and station C3 for the in-depth
276 analysis was due to the availability of data for the periods before and after reservoir construction. Based on the collected
277 data, the variability of river parameters was analyzed in the period after the construction of the reservoir complex (1996-
278 2020) as compared to the period before the construction (1984-1995). In addition, in order to identify in detail the impact of
279 the dam reservoir on the variability of water temperature and the occurrence of ice cover in the longitudinal profile of the
280 river, field measurements were carried out on 05.12.2023. This day, in the study area, was preceded by a period of
281 approximately 6 days with negative air temperatures reaching -10°C . The assessment of the occurrence of ice phenomena
282 was carried out visually (the type of ice phenomena and the percentage of the channel occupied by ice were determined).
283 Water temperature measurements were conducted at 10 points (3 upstream the reservoir and 7 downstream of the reservoir)
284 using an Elmetron CC-315 conductivity meter.

285 3. Results

286 The average air temperature in winter (November to March) during the studied periods at stations in the Dunajec
287 River basin (Czorsztyn-Sromowce reservoir complex) ranged from -0.54°C (station Tc1) to 0.36°C (station Tc2). At both
288 stations January was the coldest month in the winter period (-3.51°C at station Tc1 and -2.47°C at station Tc2), and the
289 warmest month was November (2.92°C at station Tc1 and 3.43°C at station Tc2). At climatological stations in the San River
290 basin (Solina-Myczkowce reservoir complex), temperatures ranged from -0.04°C (at station Ts5) to 0.2°C (at station Ts2,
291 Fig. 4). The warmest month at these stations was November (3.63°C at station Ts2, 3.71°C at station Ts4 and 3.46°C at


Sformatowano: Numerowanie + Poziom: 1 + Styl numeracji: 1, 2, 3, ... + Rozpocznij od: 3 + Wyrównanie: Na lewo + Wyrównanie: 0.63 cm + Wcięcie: 1.27 cm



320 **Figure 52:** Observed number of days with ice cover on the Dunajec River in each month at water gauge stations (a) and
 322 annual sum of the number of days with ice cover with a 5-year moving average (b). Observed number of days with ice cover
 323 in each month at water gauge stations.

324 At cross sections below the reservoirs, the average annual number of days with ice cover ranged from 45–53 days for the
 325 Czorsztyn-Sromowce reservoir complex (C2, C3, C4) and from 32–46 days for the Solina-Myezkowiec reservoir complex
 326 (S2, S3, S4). In all studied cross sections, a decrease in the frequency of ice cover was observed after the reservoirs were put
 327 into operation (Fig. 2).

Sformatowano: Odstęp Po: 6 pkt



328
 329 **Figure 6:** Observed number of days with ice cover on the San River in each month at water gauge stations (a) and annual
 330 sum of the number of days with ice cover with a 5-year moving average (b).


331 For the San River, at the cross-section located upstream of the reservoir (S1), a 10% decrease in the number of days
 332 with ice cover was observed in the period after its construction (1969–2020) compared to the earlier period (1950–1968). At
 333 cross sections downstream of the reservoir (S2, S3, S4), the decrease in frequency was 77% , 39.8% and 31.2%, respectively.
 334 In the period before the construction of the reservoirs on the two rivers under study, the annual ice pattern followed a similar

Sformatowano: Interlinia: 1.5 wiersza

— sformatowano: Angielski (Stany Zjednoczone)

335 trend, despite the fact that they are approximately 140 kilometers apart. For example, the Pearson correlation coefficient of
336 the annual number of days in the 1950–1968 period between stations C2 and S2 was 0.76 and was 0.86 between stations C3
337 and S3. In the period after the construction of the Solina-Myczkowce reservoir complex (1969–2009), the correlation
338 dropped to 0.04 and 0.47, respectively.

339 The predictive ability of the logistic regression models was relatively high. The accuracy of correctly classifying days in
340 the test set into two groups (days with ice cover/days without ice cover) based on the average air temperature of the 14 days
341 before the modeled day ranged from 80–87%. The developed models had very good predictive ability, as evidenced by the
342 high values of the area under the ROC (receiver operating characteristic) curve ranging from 0.89–0.94 (Fig. 73).




343
344 **Figure 73:** ROC (receiver operating characteristic) curve, AUC (area under ROC curve) values and prediction accuracy. ←

Sformatowano: Odstep Po: 6 pkt

345 In the case of the Dunajec River, there is a noticeable difference in the values observed and predicted by the model
346 downstream and upstream of the Czorsztyn-Sromowce Wyżne reservoir complex (Fig. 6). At the C1 water
347 gauge cross-section, the number of days with ice cover observed and predicted from air temperature before and after the
348 reservoir was very similar (Fig. 8). At this location in the 1957–1992 period, the annual average observed number of days
349 with ice cover was 73.1 days, while the number predicted by the model was 72.3 days.

350



351


Figure 84: Comparison of modeling results based on air temperature with observations of ice cover for the Dunajec River and the Czorsztyn-Sromowce reservoir complex (a) and the density of the distribution of days with ice cover at temperatures before and after the construction of the dam reservoir (b). The year of construction of the reservoir was taken as 1992 (the year before the construction of the Sromowce Wyżne reservoir). The red line indicates the year the dam was built.

Sformatowano: Odstęp Po: 6 pkt

— sformatowano: Czcionka: (Domyślny) Times New Roman

In the post-reservoir period (1993–2020), these values were 59.9 and 59.2 days, respectively. At point C2, located directly downstream of the reservoir, there was a significant discrepancy between the observed and predicted values based on air temperature in the post-reservoir period. In the 1957–1992 period, the average annual totals observed and

359 predicted by the model were very similar: 74.4 and 74.6, respectively. In the period after the construction of the reservoir
 360 (1993–2020), the observed average annual number of days with ice cover was 9.3, while the number of days predicted by the
 361 model was 61.



362


— sformatowano: Kolor czcionki: Automatyczny, Angielski (Zjednoczone Królestwo), Deseń: Przezroczysty

367 At point C3, after the construction of the reservoir, the difference between the observed and predicted average number of
368 days with ice cover was less than at point C2, but still notable (half as much). At point C4 in the period after the construction
369 of the reservoir, values were more similar to each other; the observed number of days was 50.1 while the predicted number
370 of days was 47.5. In the cross sections located [downstream of/below](#) the reservoir (C2, C3, C4), there was a significant shift
371 in the distribution of the number of days with ice cover in the average air temperature in the period after the construction of
372 the reservoir compared to the earlier period. This effect was not observed at point C1.

373 Similar results were obtained regarding the Solina-Myczkowce reservoir system (Fig. 95). In the cross-section located
374 [upstream of/above](#) the reservoir (S1), the temperature-based prediction and the observed average number of days with ice
375 cover were similar both before (1950–1968) and after (1969–2020) the reservoir was built. For cross-section S2, located
376 directly [downstream of/below](#) the reservoir, a significant discrepancy was found in the observed and model-predicted average
377 number of days with ice cover after the reservoir's construction. The average observed number of days with ice cover before
378 the reservoir's construction was 72.1, while the number predicted by the model was 72.8. After the reservoir's construction,
379 the observed average number of days dropped to 15.7 while the predicted number of days was 62.2. A similar trend was
380 observed for cross-section C3. The annual average observed number of days with ice cover in the period after the reservoir's
381 formation was 33.6, while that predicted by the model based on temperature was 52.5. At water gauge cross-section S4, the
382 model-predicted and observed number of days with ice cover were very similar both before (predicted = 52.1; observed =
383 60.6) and after the reservoir was built (predicted = 44.8; observed = 40.5). In the case of the San River, a slight shift in the
384 distribution of the number of days with ice cover in the mean air temperature was observed only at cross-section S2 located
385 directly [downstream of/below](#) the Solina reservoir.


386 It is worth noting that the accuracy of the prediction of ice cover occurrence by the developed models. Although the
387 prediction accuracy determined from the test set varied in the 80–87% range, the multi-year averages of observed and
388 predicted values at stations [upstream of/above](#) the reservoirs were very close to each other (59.2/59.9 in cross-section C1 and
389 76.7/73.5 in cross-section S1 in the period after the construction of the reservoirs). The high agreement of these data suggests
390 a higher accuracy than was determined from the test sets. This is most likely due to the dichotomous nature of the errors
391 made by the model. The overall error includes predictions of the occurrence of ice cover when in reality there was none, and
392 predictions of the absence of ice cover when in fact there was. Most likely, the existence of both types of errors in similar
393 proportions resulted in high agreement over the long term (> 40 years).

394 The results of analysis of Sentinel-1 (SAR) data showed that, during the study period (January-February 2017), total ice
395 cover did not form in a section of about 60 kilometers [downstream of/below](#) the Czorsztyn-Sromowce reservoir complex and
396 26 kilometers [downstream of/below](#) the Solina-Myczkowce reservoir complex (Fig. 106).
397



398
399 **Figure 106:** The extent of ice cover on the Dunajec (a) and San rivers (b).
400

401 Visual analysis of the classification results of SAR imagery, backscatter coefficient distribution maps, and optical
 402 imagery showed that the determination of the area where ice cover was not present downstream of the reservoir was
 403 relatively accurate. River sections without ice cover were characterized by a predominance of pixels classified as water,
 404 while sections with ice cover were characterized by a predominance of pixels classified as ice (Fig. 11). The largest
 405 classification errors were recorded in narrow and shallow sections of the surveyed rivers without ice cover, where there was
 406 an increase in the backscatter coefficient unrelated to the presence of ice cover. This resulted in misclassification of pixels
 407 from this area as ice. Misclassification of pixels was also recorded in transition sections between open water and total
 408 ice cover where border ice was present, especially in narrow river sections.



409
 410 **Figure 11:** Comparison of Sentinel-1 backscatter coefficient (b), classification results (c) and Sentinel-2 data (a) acquired
 411 on February 14, 2017 for the Dunajec River.
 412 Source: own elaboration based on data obtained from Copernicus Sentinel Data (<https://scihub.copernicus.eu/>). Copernicus
 413 Sentinel data (2017), processed by ESA.

414 Analysis of changes in water temperature at station C3 showed that after the construction of the Czorsztyn-Sromowce
 415 reservoir complex (1996-2020), there was a decrease in water temperature in the summer period and an increase in the
 416 winter period compared to the period prior to the dam development (1984-1995, Fig. 12b). In the period after the
 417 construction of the reservoirs, the largest increase in average water temperature was recorded in November (an increase of
 418 about 3°C). During the remaining winter months (December-March), an increase in monthly average water temperature was

also observed, ranging from 0.2°C to 1.6°C. This effect was confirmed by field measurements (Fig. 12a).




Figure 12: Water temperature and the occurrence of ice phenomena in the longitudinal profile of the Dunajec River on December 5, 2023 (a), and the variation in the average water temperature at station C3 on individual days of the year before (1984-1995) and after (1996-2020) the reservoir was built (b).

On December 5, 2023, the average water temperature upstream of the reservoir (measuring points f1, f2, f3) amounted to 0.4°C. In this section, border ice covered up to 20% of the water surface. Downstream of the reservoir there was a sharp increase in water temperature up to 4°C (Fig. 12a). As the distance from the reservoir increased, water temperature decreased up to 0.3°C at a distance of about 75 kilometers downstream of the reservoir (measuring point f10). In the section of the river up to about 45 kilometers downstream of the reservoir, no ice phenomena were observed. At distances further than 50 kilometers from the reservoir, initial ice forms, ice floe and local border ice were observed.

Analysis of changes in flow volume at station C3 showed that in winter periods after the construction of the Czorsztyn-Sromowce reservoir complex (1996-2020), there was an increase in water flow volume (an average increase of 3.7 m³/s⁻¹) compared to the period before the dam development (1984-1995, Fig. 10). During the winter period, an increase in flow volume was recorded in November (an increase of 7.3 m³/s⁻¹), February (an increase of 7.5 m³/s⁻¹) and slightly in December (0.5 m³/s⁻¹). The increase in flow volume resulted in a 2 cm increase in the average water level in winter (Fig. 13).

— sformatowano: Czcionka: 10 pkt, Pogrubienie, Angielski (Stany Zjednoczone)

— sformatowano: Czcionka: Pogrubienie, Angielski (Stany Zjednoczone)

— sformatowano: Czcionka: 10 pkt, Pogrubienie, Angielski (Stany Zjednoczone)

— sformatowano: Angielski (Stany Zjednoczone)

— sformatowano: Angielski (Stany Zjednoczone)

— sformatowano: Indeks górny


— sformatowano: Indeks górny

— sformatowano: Indeks górny

— sformatowano: Indeks górny

— sformatowano: Indeks górny

— sformatowano: Indeks górny



436
437 **Figure 13:** Average water level and discharge at station C3 in the periods before (1984-1995) and after (1996-2020) the
438 construction of the reservoir.
439

440 **4. Discussion**

441 This study demonstrates that the analyzed dam development was an important element in transforming the ice regime of
442 the downstream rivers. This is evidenced by a significant decrease in the incidence of ice cover downstream of the reservoirs
443 in the period after their construction, with minor change at cross sections upstream of the reservoirs. The significance of the
444 reservoirs is also indicated by the lack of sharp increases in air temperature in the post-reservoir period in the study areas. In
445 the case of the Dunajec River, a lower average winter air temperature was observed in the 10-year period after the reservoir
446 construction as compared to the 10-year period prior to the investment. At the same time a decrease in the frequency of ice
447 cover was observed at cross sections C2 and C3 (downstream of the reservoir). This suggests that the recorded disappearance
448 of ice cover on the studied rivers is not due to climatic conditions. The important impact of reservoir operations on the river
449 ice regime is also evidenced by the analysis of water temperature, ice cover occurrence and changes in flow volume. Field
450 studies have shown that during periods of low air temperatures (reaching -10°C), high water temperatures (reaching 4°C) are
451 possible downstream of the reservoir, due to the release of bottom warm water from the reservoir. On the day of the survey,
452 the increase in temperature resulted in the absence of ice phenomena along the 45 kilometers downstream of the reservoir. In
453 the post-reservoir period (1996-2020) at station C3, this effect (and the potential impact of climatic variability) translated
454 into a 1.18°C increase in average winter water temperature compared to the earlier period (1984-1996). The important role of
455 dam reservoirs in the transformation of water temperature is also evidenced by other studies; Kędra and Wiejaczka (2016,
456 2017) and Wiejaczka et al. (2015) have previously shown that the Czorsztyn-Sromowce reservoir system had a significant
457 impact on the water temperature of the Dunajec River, as well as the synchronization of air and water temperature in the

— sformatowano: Nie Wyróżnienie

— sformatowano: Nie Wyróżnienie

— sformatowano: Nie Wyróżnienie

— sformatowano: Nie Wyróżnienie

— sformatowano: Nie Wyróżnienie

458 river. The analysis of flow volume showed that there was an increase in flow volume in the post-reservoir period (1996-
459 2020) at station C3 compared to the period prior to the dam development (1984-1996). Increased discharge in the river
460 results in delayed formation of stable ice cover due to increased water velocity in the riverbed (Houkuna et al. 2022).

461 In the study area, the increase in air temperature most likely manifested itself in a slightly later formation of ice cover and
462 its earlier disappearance. After 2010, at cross-sections located upstream of the reservoirs, ice cover occurred sporadically in
463 November and March, which is partially confirmed by trends in air temperature at the climatological stations studied (an
464 increase in average November temperature in the range of 0.6-0.8°C/decade in the period 1982-2015). Accordingly, rising
465 air temperatures may exacerbate the effects on river ice cover caused by the operation of dam reservoirs. However, the
466 relatively small variation in the annual number of days with ice cover in cross sections upstream of the reservoirs and the
467 lack of significant trends in average air temperature in all months except November suggest that in the study area the
468 increase in air temperature has not significantly changed the frequency of ice cover. Further research based on more detailed
469 data is needed in order to explore the potential impact of climate change on the ice cover.

470 On the basis of observational data and modeling results, it was found that the greatest transformations occurred at cross
471 On the basis of observational data and modeling results, it was found that
472 the greatest transformations occurred at cross sections located closest to the facilities (C2, S2), and reservoir influence
473 decreased with increasing distance from the reservoir. This effect may be interpreted as a gradual decrease in the influence of
474 the reservoir and restoration of the natural course of thermal and ice processes in rivers. The use of a classification method
475 based on logistic regression allowed us to estimate that, in the case of the Czorsztyn-Sromowce reservoir complex, at cross-
476 section C2 (1.8 km downstream of the dam) in the period 1996–2020, the operation of the reservoir reduced the
477 duration of ice cover by 84% on average, while at point C3 (22 km downstream of the dam), reservoir operation
478 reduced ice cover by 46%. Similarly, in the case of the Solina-Myczkowce reservoir complex, the operation of the reservoir
479 reduced the duration of ice cover by about 75% at point S2 (11.7 kilometers downstream of the dam), and by 36% at
480 point S3 (33 kilometers downstream of the dam). These results suggest that in the stretch of rivers about 20-40
481 kilometers downstream of the reservoirs, the influence of the reservoirs was the main factor (it transformed the ice
482 regime more than climatic variability) determining the observed disappearance of ice cover and the course of ice processes.
483 This is supported by the much smaller magnitude of the decrease in the frequency of ice cover at cross sections upstream
484 of the reservoirs (10% and 13.3% after the construction of the reservoirs compared to the earlier period), where the
485 decrease was mainly due to climatic conditions.

486
487 extreme caution should be exercised when analyzing the impact of dam reservoirs using the presented method because of
488 limitations that arise from both the nature of the data and the river ice processes themselves. First of all, the presented
489 method does not take into account other possible factors affecting river ice phenomena; these mainly may include regulation
490 of rivers affecting the conditions of ice formation, all kinds of thermal pollutants emitted into rivers, discharges of municipal

— sformatowano: Nie Wyróżnienie

491
492 A visual comparison of the classification results of radar imagery (Sentinel-1) with optical data (Sentinel-2) showed that
493 it was possible to determine, with relative accuracy, the extent to which there was no ice cover [downstream of below](#) dam
494 reservoirs on mountain rivers with similar characteristics to those analyzed in this study. Based on the threshold of the
495 backscattering coefficient to two classes (water/ice) on rivers similar to those studied here (width of 20–100 meters), it was
496 possible to determine the approximate extent of the river section [downstream of below](#) the reservoir on which the total ice
497 cover did not form. The range of influence of the studied reservoirs on the occurrence of river ice cover on the basis of SAR
498 data was determined to be 26 kilometers for the Solina-Myczkowce reservoir complex, and 60 for the Czorsztyn-Sromowce
499 reservoir complex. An analysis of the river network in the catchments of the studied rivers showed that the smaller extent of
500 the influence of the Solina-Myczkowce reservoir complex was most likely due to the mixing of the waters of the San with
501 two relatively large tributaries in close proximity to the reservoir, the Hoczewka and Osława (average winter flow at the
502 mouth of 2.8 m³/s⁻¹ and 8.6 m³/s⁻¹, respectively). The mixing of these waters with those of the San River (average winter
503 outflow from the reservoir 24.6 m³/s⁻¹) may lead to a drop in water temperature and the appearance of ice phenomena. This is
504 also evidenced by the lack of a clear shift in the number of days with ice cover in the average air temperature at cross
505 sections [downstream of below](#) the reservoir. By comparison, for the Dunajec River, total ice cover appeared during the
506 analyzed period about 60 kilometers [downstream of below](#) the reservoir in the vicinity of a tributary of the Poprad River, one
507 of the larger tributaries of the river.

508 Similar results have been previously obtained for other dam reservoirs located in mountainous areas, including in the
509 Carpathian Mountains. Cyberska (1972, 1975) analyzed the influence of a complex of dam reservoirs (dam height of 32.5
510 meters) on the thermals and occurrence of ice phenomena on the Dunajec River (Poland). Cyberska estimated that in the
511 period after the reservoir's construction, the 12–65 km downstream area of the reservoir saw an average 65% reduction in the
512 duration of ice cover. These values are slightly higher than those obtained in this study, especially for cross sections located
513 far from the reservoir, which may be due to the fact that they were estimated based on the comparison of periods before and
514 after the reservoir's construction without accounting for the possible influence of changing climatic conditions. An estimate
515 based on a similar methodology made by Wiejaczka (2009) showed that, on the Ropa River, at a cross-section located 16 km
516 downstream of the dam, after the construction of the dam reservoir (dam height of 34 meters), there was a 35% decrease in
517 the frequency of ice cover (total and border ice). Chang et al. (2016) compared periods with similar thermal conditions
518 (before and after reservoir construction) and analyzed the impact of large (dams 178 and 147 meters high) dam reservoirs on
519 ice phenomena on the Yellow River. They found that [reservoir](#) operation reduced the duration of ice phenomena at
520 downstream stations by 33, 22, and 8 days, which is less than the value estimated in this study. However, these results are
521 particularly significant given that the gauging stations are located more than 800 kilometers [downstream of](#)
522 the reservoirs. An important role in the transformation of the ice regime was also demonstrated in the case of the Williston
523 Reservoir in Canada on the Peace River (dam height 186 meters). As a result of that operation, total ice cover did not form
524 for 100–300 kilometers downstream of the dam (Jasek and Pryse-Phillips 2015). This is far higher than the value estimated


525 in this study, which may be due to the different sizes of these rivers and reservoirs. Similar results were obtained for the
526 Krasnoyarsk reservoir (Belolipetsky and Genova 1998); downstream of the dam (124 meters), the ice cover also did not form
527 for 100–300 kilometers, depending on hydrometeorological conditions. Transformations of the river ice regime have been
528 observed for both large reservoirs (dams higher than 15 meters) and small ones. For example, Maheu (2016) analyzed the
529 impact of small dam reservoirs (dams 7–13 meters high) on thermals and water ice in eastern Canada. Using two examples,
530 he showed that the operation of these facilities significantly raised water temperatures and reduced ice formation in sections
531 up to 2.5 kilometers downstream.

532 Similar effects on river ice cover have also been reported for lowland reservoirs, which have different characteristics
533 (usually less depth) due to terrain. Takács et al. (2013) analyzed the occurrence of ice cover upstreamabove and downstream
534 ofbelow small dams (< 10 meters) in the Raba River basin (Westen Transdanubia, Hungary). They based their study on
535 selected periods with similar thermal conditions before and after the construction of the reservoirs, showing that, after
536 ~~theirtheir~~ construction, the relative frequency of ice cover downstream ofbelow reservoirtheir location decreased by up to
537 10%, and that anthropogenic factors were crucial in transforming the ice regime of rivers. The significantly smaller impact of
538 these reservoirs than the values estimated in this study can be explained by the smaller size of the damby their smaller size.
539 Pawłowski (2015) showed that the construction of the Włocławek reservoir on the Vistula River (Poland) resulted in a 47%
540 reduction in the duration of ice cover downstream of its location and a 26% reduction in the duration of all ice events,
541 leading to a significant transformation of the river's ice regime. Here, to demonstrate the impact of reservoirs, periods with
542 similar average air temperatures before and after the reservoirs were selected. These values were smaller than those obtained
543 in this work, which is likely due to the fact that the Włocławek reservoir has a damming level that is five-fold lower (11
544 meters). Apsite et al. (2016) analyzed the impact of the operation of three dam reservoirs (dam heights of 18–40 meters) on
545 the phenology of ice phenomena on the Daugava River (Latvia), showing that, at a station 6 kilometers downstream ofbelow
546 the reservoir after its construction, there was a reduction in the duration of ice cover by 91 days. This is a greater decrease in
547 the frequency of ice cover than estimated in this study, but it was not determined how much of this effect was due to the
548 construction of the reservoir in relation to climate change.

549 Despite the relatively high predictive ability of the presented logistic regression models and the high agreement between
550 modeling results and observations at stations upstream of the reservoirs, caution should be exercised when
551 analyzing the impact of dam reservoirs using the presented method because of limitations that arise from both the nature of
552 the data and the river ice processes themselves. First of all, the presented method does not take into account other possible
553 factors affecting river ice phenomena; these mainly may include regulation of rivers affecting the conditions of ice
554 formation, all kinds of thermal pollutants emitted into rivers, discharges of municipal and industrial wastewater that can
555 increase the content of dissolved substances and thus lower the freezing point, and the occurrence of natural changes in the
556 hydrological and morphological characteristics of rivers and their channels. An important problem is also the significant
557 sensitivity of the model to input data on the occurrence of river ice cover; due to its characteristics (large variation of
558 parameters in the longitudinal profile of rivers, non-linear nature of development and disappearance, significant sensitivity to

559 hydrological and meteorological conditions), this is difficult to describe and classify into a rigid framework, which can
560 translate into modeling results.

561 An important issue in analyzing the impact of dam-reservoir operations on the ice regime of rivers lies in distinguishing
562 their impact from that of air-temperature variability, which is fundamental to the course of river-ice processes and is the main
563 determinant of the currently observed shortening of the duration of ice events on rivers (Magnuson et al., 2000; Yang et al.,
564 2020; Newton and Mullan 2020). In the studies cited above, the impact of reservoirs was analyzed by comparing thermally
565 similar periods before and after their formation. Typically, periods have been selected based on average winter temperatures.
566 However, approach appears to be an oversimplification due to the averaging of extreme values over entire periods.
567 Furthermore, this method limits analysis to selected periods only. The method presented in this work made it possible to
568 demonstrate the impact of reservoirs on river ice cover over the entire period after their creation, regardless of climatic
569 variability.



571 **Figure 149:** Location of dam reservoirs (red dots) in areas of river ice cover in Europe (highlighted in blue). The area of this
572 research is marked with an arrow. Compiled from Fukš, 2023.

5. Conclusions

Using two reservoirs located in the Carpathian region as an example, this study presents a method for estimating the impact of dam reservoirs on river ice cover based on measurement data from water gauge cross sections and a logistic regression model. An estimation of the extent of the impact of dam reservoirs based on SAR data acquired by Sentinel-1 was also made here, and this method's use for determining the extent of dam impact was evaluated. The conclusions of the study can be summarized as follows:

1. At the local scale (single river), dam reservoirs have a greater impact on the observed decrease in the occurrence of ice cover of the rivers studied than does climate change. The results presented here suggest that, in areas with a large number of reservoirs, these reservoirs may play an important role at the regional scale. This is evidenced by the modeling results and their comparison to the variability of ice cover occurrence in cross-sections not influenced by a dam reservoir (C1, S1). The decrease in the incidence of ice cover due to the operation of dam reservoirs could exceed 80% in the sections of rivers immediately downstream of dam locations, with this effect decreasing with increasing distance from the reservoir. Based on this study, it can be assumed that the increase in the number of dam reservoirs is an important factor in the currently observed shortening of the duration of river ice cover.

2. The range of river sections downstream of/below the studied reservoirs on which total ice cover does not form was estimated at 60 and 26 kilometers from the reservoir dam location. Based on the results presented in this study and a review of the literature, it can be concluded that the extent of dams' impact varies greatly. This is most likely due to a number of environmental conditions in which the river and reservoir are located, as well as the technical features of the dam and reservoir.

2-3. Due to the significant impact of the studied reservoirs on the occurrence of river ice cover, it is necessary to take into account the influence of such structures when conducting studies addressing the role of climate change in the temporal and spatial variability of river ice cover. Failure to take into account the impact of reservoirs may result in erroneous attribution of the disappearance of river ice cover to an increase in air temperature and misinterpretation of the results. This is important given the significant increase in the number of reservoirs in areas with river ice cover occurrence in the context of accelerating trends towards warmer air temperatures (1980s and 1990s).

3-4. Logistic regression models are a useful tool for studying the impact of dam reservoirs on river ice cover. This is evidenced by the high predictive ability of the created models, the relatively high accuracy determined on the basis of test sets, and the very high agreement of the modeling results with observations at cross sections above-upstream of the reservoirs. After appropriate adaptation, the logistic regression model and the presented procedure can be used to study the impact of dam reservoirs on other elements of the natural environment.

4-5. In relatively narrow (20–100 meters) mountain rivers, SAR data is a useful tool for determining the sections downstream of/below dam reservoirs in which ice cover does not form. Despite the many errors inherent in the classification of SAR imagery, it is possible to estimate how far downstream of/below the reservoirs there is ice cover, which permits study

— sformatowano: Nie Wyróżnienie

608 of the extent of their influence. The usefulness of this type of data is evidenced by the validation of results based on optical
609 imaging of the Sentinel-2 satellite.

610 **Funding**

611 This research was funded in whole by the National Science Center, Poland (grant number: 2020/39/O/ST10/00652). For the
612 purpose of Open Access, the author has applied a CC-BY public copyright license to any Author Accepted Manuscript
613 (AAM) version arising from this submission.

614 **Declaration of Competing Interest**

615 The author declares that he has no known competing financial interests or personal relationships that could influence the
616 work presented in this article.

617 **Data availability**

618 Data on the daily occurrence of ice cover on the studied rivers and daily air temperature were obtained from the repository of
619 the Polish Institute of Meteorology and Water Management (IMGW-PIB, https://danepubliczne.imgw.pl/data/dane_pomiarowo_obserwacyjne/) and hydrological yearbooks of surface waters of the
620 Polish Institute of Meteorology and Water Management from 1949-1980. Sentinel-1 data was obtained from the Earth
621 Engine Data Catalog (https://developers.google.com/earth-engine/datasets/catalog/COPERNICUS_S1_GRD).

623 **References**

- 624 [Apsite, E., Elferts, D., and Latkovska, I.: Long-term changes of Daugava River ice phenology under the impact of the](#)
625 [cascade of hydro power plants., in: Proc. Latv. Acad. Sci., 70, 71–77, <https://doi.org/10.1515/prolas-2016-0012>, 2016.](#)
- 626 [Ayalew, L., and Yamagishi, H.: The application of GIS-based logistic regression for landslide susceptibility mapping in the](#)
627 [Kakuda-Yahiko Mountains, Central Japan, *Geomorphology*, 65\(1-2\), 15-31,](#)
628 [https://doi.org/10.1016/j.geomorph.2004.06.010, 2005.](#)
- 629 [Belolipetsky, V. M., Genova, S. N.: Investigation of hydrothermal and ice regimes in hydropower station bays. *Int. J.*](#)
- 630 [Comput. Fluid Dyn.](#), 10(2), 151-158, <https://doi.org/10.1080/10618569808961681>, 1998.
- 631 [Cai, H., Piccolroaz, S., Huang, J., Liu, Z., Liu, F., Toffolon, M.: Quantifying the impact of the Three Gorges Dam on the](#)
632 [thermal dynamics of the Yangtze River. *Environ. Res. Lett.*, 13\(5\), 054016, \[10.1088/1748-9326/aab9e0\]\(https://doi.org/10.1088/1748-9326/aab9e0\), 2018.](#)
- 633 [Carlson, R.F.: Ice formation on rivers and lakes. *North. Eng.*, 13 \(4\), 4–9, 1981.](#)
- 634 [Chang, J., Wang, X., Li, Y., and Wang, Y.: Ice regime variation impacted by reservoir operation in the Ning-Meng reach of](#)
635 [the Yellow River. *Nat. Hazards*, 80, 1015–1030, <https://doi.org/10.1007/s11069-015-2010-5>, 2016.](#)
- 636 [Chen, Y., Lian, J., Zhao, X., Guo, Q., Yang, D.: Advances in Frazil Ice Evolution Mechanisms and Numerical Modelling in](#)
637 [Rivers and Channels in Cold Regions. *Water*, 15\(14\), 2582, 2023.](#)
- 638 [Cox, D. R.: The regression analysis of binary sequences. *Journal of the Royal Statistical Society: Series B*](#)
- 639 [\(Methodological\), 20\(2\), 215-232, 1958.](#)
- 640 [Cyberska B., Wpływ zbiornika retencyjnego na transformację naturalnego reżimu termicznego rzeki. *Prace IMGW*, 4, 45-](#)
641 [108, 1975. \[in Polish\]](#)

— sformatowano: Czcionka: (Domyślny) Times New Roman, Kolor czcionki: Automatyczny

— sformatowano: Czcionka: (Domyślny) Times New Roman, Kolor czcionki: Automatyczny

— sformatowano: Czcionka: (Domyślny) Times New Roman, Kolor czcionki: Automatyczny

— sformatowano: Czcionka: (Domyślny) Times New Roman, Kolor czcionki: Automatyczny

— sformatowano: Czcionka: (Domyślny) Times New Roman

— sformatowano: Czcionka: (Domyślny) Times New Roman, Nie Kursywa

— sformatowano: Czcionka: (Domyślny) Times New Roman

642 [Cyberska B., Zmiany w temperaturze i zlodzeniu rzek poniżej zbiorników retencyjnych. Gospodarka Wodna, 7, 244-250,](#)
643 [1972. \[in Polish\]](#)

644 [Fox-Kemper, B., Hewitt, H. T., Xiao, C., et al. \(Eds.\) Climate Change 2021: The Physical Science Basis. Contribution of](#)
645 [Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge](#)
646 [University Press, Cambridge, United Kingdom and New York, NY, USA, 1211–1362, \[https://\]\(https://doi.org/10.1017/9781009157896.011\)](#)
647 [doi.org/10.1017/9781009157896.011, 2021.](#)

648 [Fukś M.: Changes in river ice cover in the context of climate change and dam impacts – a review. Aquat. Sci., 85\(4\),](#)
649 [https://doi.org/10.1007/s00027-023-01011-4, 2023.](#)

650 [Hennig, J., Zbiorniki retencyjne, in: Dynowska I., Maciejewski M., \(Eds.\), 1991, Dorzecze górnej Wisły Część 1, PWN,](#)
651 [1991. \[in Polish\]](#)

652 [Huokuna, M., Morris, M., Beltaos, S., Burrell, B. C.; Ice in reservoirs and regulated rivers. Int. J. of River Basin](#)
653 [Manag., 20\(1\), 1-16, <https://doi.org/10.1080/15715124.2020.1719120>, 2022.](#)

654 [Jasek, M., Pryse-Phillips, A.: Influence of the proposed Site C hydroelectric project on the ice regime of the Peace](#)
655 [River. Can. J. Civ. Eng., 42\(9\), 645-655, <https://doi.org/10.1139/cjce-2014-0425>, 2015.](#)

656 [Kędra, M., and Wiejaczka, Ł.: Climatic and dam-induced impacts on river water temperature: Assessment and management](#)
657 [implications. Sci. Total Environ., 626, 1474-1483, <https://doi.org/10.1016/j.scitotenv.2017.10.044>, 2017.](#)

658 [Kendall, M.G.; Rank Correlation Methods, 4th ed.; Charles Griffin: London, UK, 1975.](#)

659 [Kloss, A.: Zespół zbiorników wodnych Czorsztyn-Niedzica i Sromowce Wyżne im. Gabriela Narutowicza. Monografia.](#)
660 [Regionalny Zarząd Gospodarki Wodnej w Krakowie, Hydroprojekt Warszawa, Instytut Meteorologii i Gospodarki](#)
661 [Wodnej, Kraków, 2003. \[in Polish\]](#)

662 [Lee, J. Y., and Kim, J. S.: Detecting areas vulnerable to flooding using hydrological-topographic factors and logistic](#)
663 [regression. Appl. Sci., 11\(12\), 5652, <https://doi.org/10.3390/app11125652>, 2021.](#)

664 [Lehner B, Reidy Liermann C, Revenga C, Vörösmarty C, Fekete B, Crouzet P, Döll P, Endejan M, Frenken K, Magome](#)
665 [J, Nilsson C, Robertson JC, Rodel R, Sindorf N, Wisser D \(2011\) Highresolution mapping of the world’s reservoirs and](#)
666 [dams for sustainable river-fow management. Front Ecol Environ 9:494–502. <https://doi.org/10.1890/100125>.](#)

667 [Lewińska, J., Lewiński, A.: Wpływ energetycznego wykorzystania zbiorników wodnych na zmianę reżimu termicznego](#)
668 [wody. Gosp. Wodna, 8, 295 – 296, 1972. \[in Polish\]](#)

669 [Lewis, S. L., and Maslin, M. A.: Defining the Anthropocene. Nature, 519, 171-180, <https://doi.org/10.1038/nature14258>,](#)
670 [2015.](#)

671 [Magnuson, J. J., Robertson, D. M., Benson, B. J., Wynne, R. H., Livingstone, D. M., Arai, T., Assel, R.A., Barry, R. G.,](#)
672 [Card, V., Kuusisto, E., Granin, N. G., Prowse, T. D., Stewart, K. M., and Vuglinski, V.S.: Historical trends in lake and](#)
673 [river ice cover in the Northern Hemisphere, Science, 289: 1743–1746, <https://doi.org/10.1126/science.289.5485.1743>,](#)
674 [2000.](#)

— sformatowano: Polski

Kod pola został zmieniony

— sformatowano: Czcionka: (Domyślny) Times New Roman, Kolor czcionki: Automatyczny, Angielski (Zjednoczone Królestwo)

— sformatowano: Czcionka: (Domyślny) Times New Roman, Kolor czcionki: Automatyczny, Angielski (Zjednoczone Królestwo)

— sformatowano: Czcionka: 10 pkt, Nie Kursywa, Angielski (Zjednoczone Królestwo)

— sformatowano: Czcionka: (Domyślny) Times New Roman, Nie Kursywa, Kolor czcionki: Automatyczny, Angielski (Zjednoczone Królestwo)

— sformatowano: Czcionka: 10 pkt, Nie Kursywa, Angielski (Zjednoczone Królestwo)

— sformatowano: Czcionka: (Domyślny) Times New Roman, Nie Kursywa, Kolor czcionki: Automatyczny, Angielski (Zjednoczone Królestwo)

— sformatowano: Czcionka: 10 pkt, Nie Kursywa, Angielski (Zjednoczone Królestwo)

— sformatowano: Czcionka: (Domyślny) Times New Roman, Kolor czcionki: Automatyczny, Angielski (Zjednoczone Królestwo)

— sformatowano: Domyślna czcionka akapitu, Czcionka: (Domyślny) Times New Roman, Kolor czcionki: Automatyczny, Angielski (Zjednoczone Królestwo)

Kod pola został zmieniony

— sformatowano: Czcionka: (Domyślny) Times New Roman

— sformatowano: Czcionka: (Domyślny) Times New Roman

— sformatowano: Czcionka: (Domyślny) Times New Roman, Nie Kursywa

— sformatowano: Czcionka: (Domyślny) Times New Roman

— sformatowano: Czcionka: (Domyślny) Times New Roman

Sformatowano: Normalny, Wcięcie: Z lewej: 0 cm, Wysunięcie: 0.5 cm, Bez punktów lub numeracji

— sformatowano: Polski

— sformatowano: Czcionka: (Domyślny) Times New Roman, 10.5 pkt, Kolor czcionki: Automatyczny

- 707 [Wiejaczka, Ł., Kijowska-Strugała, M., Pierwoła, P., and Nowak, M.: Influence of the Czorsztyn-Sromowce Wyżne](#)
708 [Reservoir complex on the Dunajec River thermal-regime, Geogr. Pol., 88\(3\), 467-482,](#)
709 <http://dx.doi.org/10.7163/GPol.0029>, 2015.
- 710 [Wiejaczka, Ł.: Wpływ zbiornika wodnego „Klimkówka” na zlodzenie Ropy, in: Funkcjonowanie środowiska](#)
711 [przyrodniczego w okresie przemian gospodarczych w Polsce, edited by: Bochenek W., and Kijowska M., Szymbark,](#)
712 [172-187, 2009. \[in Polish\]](#)
- 713 [Wu, Y., Duguay, C. R., and Xu, L.: Assessment of machine learning classifiers for global lake ice cover mapping from](#)
714 [MODIS TOA reflectance data, Remote Sensing of Environment, 253, 112206, https://doi.org/10.1016/j.rse.2020.112206,](#)
715 [2021.](#)
- 716 [Yang, X., Pavelsky, T. M., and Allen, G. H.: The past and future of global river ice. Nature, 577\(7788\), 69-73,](#)
717 <https://doi.org/10.1038/s41586-019-1848-1>, 2020.
- 718 [Apsite, E., Elferts, D., and Latkovska, I.: Long-term changes of Daugava River ice phenology under the impact of the](#)
719 [cascade of hydro power plants., in: Proc. Latv. Acad. Sci., 70, 71–77, https://doi.org/10.1515/prolas-2016-0012, 2016.](#)

Sformatowano: Wcięcie: Z lewej: 0 cm, Pierwszy wiersz: 0 cm