

~~Extremes~~ Extreme events of surface snow grains change grain size increase in East Antarctica and their relationship with meteorological conditions

Claudio Stefanini^{1,2}, Giovanni Macelloni², Marion Leduc-Leballeur², Vincent Favier³, Benjamin Pohl⁴, and Ghislain Picard³

¹Dipartimento di Scienze Ambientali, Informatica e Statistica, DAIS, Ca' Foscari University of Venice, 30170 Mestre-Venezia, Italy

²Institute of Applied Physics “Nello Carrara” – National Council of Research, 50019 Sesto Fiorentino, Italy

³Université Grenoble Alpes, CNRS, IRD, Grenoble INP, IGE, 38000 Grenoble, France

⁴Biogéosciences, UMR6282 CNRS / Université de Bourgogne, Dijon, France

Correspondence: Stefanini (claudio.stefanini@unive.it)

Abstract. This study explores the ~~grain-size-seasonal-variations~~ seasonal variations of snow grain size on the East Antarctic Plateau, where dry metamorphism occurs, by using microwave radiometer observations from 2000 to 2022. Local meteorological conditions and large scale atmospheric phenomena have been considered in order to explain some peculiar changes of the snow grains. We find that the highest ice divide is the region with the largest grain size in the summer, mainly because the wind speed is low. Moreover, some extreme grain size values with respect to the average (over $+3\sigma$) were identified. In these cases, the ERA5 reanalysis revealed a high pressure blocking ~~/ridge situation in the proximity of the~~ close to the onsets of the summer increase of the grain size, ~~conveying the relatively warm and moist air coming from the mid-latitudes, often associated with atmospheric rivers~~. It channels moisture intrusions from the mid-latitudes, through atmospheric rivers that cause major snowfall events over the plateau. If weak wind and low temperature conditions occur during the following weeks, dry snow metamorphism is facilitated, leading to grain growth. This determines anomalous high maximums of the snow grain size at the end of summer. These phenomena confirm the importance of moisture intrusion events in the East Antarctica and their impact on the physical properties of the ice sheet surface, with a co-occurrence of atmospheric rivers and seasonal changes of the grain size significant over 95%.

1 Introduction

Understanding and modelling the physical changes in snow properties, and in particular the grain size, is important to predict the changes in snowpack albedo (Grenfell et al., 1994; Domine et al., 2006), and hence of surface energy balance at high latitudes. Moreover, it provides information on the firm properties, useful to retrieve the ice sheet mass balance by space-borne altimetry (Keenan et al., 2021; Kingslake et al., 2022). The seasonal variations of the grain size observed by spaceborne microwave radiometer in Antarctica shows a large summer increase, which is mainly driven by temperature (Picard et al., 2012). The increase in temperature during austral summer leads to coarsening of snow grains due to grain-to-grain transport of water vapour (Col-

beck, 1982, 1993; Sturm and Benson, 1997; Flanner and Zender, 2006; Town et al., 2008), while winter precipitations and wind-transported snow accumulate small grains over the surface (Domine et al., 2007). On the East Antarctic Plateau the mean annual precipitation is low, less than 100 kg m^{-2} in the highest part of the ice sheet, due to its ~~insulation~~ isolation from the rest of the hemisphere (Palerm et al., 2014). However, some relatively warm and moist oceanic air masses occasionally travel far inland, providing higher than usual snow accumulations (Turner et al., 2019). These intrusions could be linked to atmospheric rivers known to transport warm and moist air from ~~the tropics~~ subtropics or mid-latitudes to the polar latitudes (e.g. Wille et al., 2021).

~~Understanding and modelling the physical changes in snow properties, and in particular the grain size, is important to predict the changes in snowpack albedo (Grenfell et al., 1994; Domine et al., 2006), and hence of surface energy balance at high latitudes. Moreover, it provides information on the firm properties, useful to retrieve the ice sheet mass balance by space-borne altimetry (Keenan et al., 2021; Kingslake et al., 2022).~~ On the Antarctic Plateau, in-situ ~~snow structure observations~~ observations of snow structure are rare and the meteorological measurements, needed to interpret the causes of changes in the snowpack, are sparse. This is the reason for the constant need of supplementary data coming, for example, from satellite observations, able to provide continuous information on snow surface characteristics at continental scale. Previous studies investigated the snow grain size retrieved from remote sensing observations. In Jin et al. (2008) near infrared and visible satellite observations were exploited to retrieve the grain size in the top 1 cm during clear sky conditions and observed widespread summer increase of grain size. Picard et al. (2012) combined high frequency passive microwave observations to implement an indicator of grain size in the uppermost ~ 10 cm of the snowpack, and compared it to observations at Dome C and simulation results obtained by modelling water vapour transport near the surface. They observed inter-annual variability in the summer grain growth and linked them to precipitation variability. ~~Also Casado et al. (2021)~~ Casado et al. (2021) also focused on Dome C, deepening the link between water isotopic signature and surface snow metamorphism observing changes in this grain size indicator for two contrasted summers (2014 and 2015). Hence, past studies were limited to clear sky conditions or to a particular location (Dome C). Our work, by means of high frequency microwave observations explores the interaction between surface snow grain size and meteorological conditions, on a continental scale and on a seasonal time frame.

This work focuses on extremes of ~~high grain size~~ large grain sizes. The investigated events were recorded during summer between 2000 and 2022, close to the highest elevation points of the East Antarctica Plateau, along the ice divide. Using both passive microwave observations and atmospheric reanalysis, we investigate the physical processes involved in these events and their link with the local and regional atmospheric conditions.

The paper is organised as follows. Section 2 and Section 3 present the satellite and meteorological datasets and statistical tools. Section 4 shows the results by first describing the common features of the grain size summer increases and their relationship with local meteorological conditions (Section 4.1); second, focusing the analysis to the four most extreme grain size increases (Section 4.2) and lastly, connecting the changes of the grain size with large scale meteorological conditions (Section 4.3). In Section 5 we discuss the results and Section 6 draws conclusions.

2 Materials

55 2.1 Passive microwave observations

We use the Advanced Microwave Sounding Unit B (AMSU-B), which is a 5-channel microwave radiometer operating since 1998 on-board of NOAA-15 to NOAA-19 and METOP satellites. Swath data from the sensors NOAA-15 to NOAA-19 have been downloaded from the National Center for Environmental Information (<https://www.avl.class.noaa.gov/>) and after selecting incidence angles in the range 0–25°, processed to obtain daily mean from January 2000 to May 2022 in Southern polar stereographic projection (EPSG: 3976) at a resolution of 25 km. Brightness temperatures observed at 89 and 150 GHz are then used to compute the snow *Grain Size Index (GSI)* defined as $1 - T_B(150 \text{ GHz})/T_B(89 \text{ GHz})$ by Picard et al. (2012). They focused their study on Dome C, and provided an approximate correspondence between the index and millimeters: 0.00 of GSI is ~ 0.025 mm and 0.20 is ~ 0.150 mm (for the grains in the first 7 cm of the snowpack). Because of the particular and high sensitivity of the microwaves to liquid water, GSI is only effective in regions where snow melt never occurs, such as the East Antarctic Plateau (Picard et al., 2007).

2.2 Meteorological data

2.2.1 ERA5 reanalysis

For this study, we use skin temperature, 2 m temperature, 10 m wind speed, snowfall, surface pressure, cloud cover and surface downward long-wave and short-wave radiation flux over Antarctica from the ~~ERA~~ ERA5 reanalysis produced by the European Centre for Medium-Range Weather Forecasts (ECMWF, Hersbach et al., 2018a, b). Data are provided over a regular latitude-longitude grid of 0.25°, and we reprojected them ~~over using~~ the Southern polar stereographic projection. We used daily ~~mean means~~ from 2000 to ~~2022~~ 2022 but also hourly values as inputs for the thermal model.

2.2.2 Atmospheric river catalogs

The atmospheric river (AR from now on) presence is obtained from the two catalogs built by Wille et al. (2021). The detection is based on the integrated water vapour (IWV) and the meridional component of the integrated vapour transport (vIVT), both from 3 hourly data of ERA5 and of the Modern-Era Retrospective analysis for Research and Applications, Version 2 (MERRA-2) reanalysis (Gelaro et al., 2017). The datasets provide a boolean indicator of the ~~atmospheric river~~ AR presence for each grid cell worldwide with a latitude–longitude resolution of $0.50^\circ \times 0.625^\circ$. Previous work revealed weak sensitivity to the input meteorological dataset used to study the climatology of ARs, between MERRA-2 (used for the tier-2 exercise of the ARTMIP project: Collow et al. (2022)) and ERA5, especially over East Antarctica, even though important differences could arise when considering individual events (Pohl et al., 2021; Wille et al., 2021).

2.2.3 Antarctic Oscillation index

The Antarctic Oscillation index (AAO, also known as the Southern Annular Mode, SAM) (Thompson and Wallace, 2000) is provided by the National Oceanic and Atmospheric Administration (NOAA) and based on the daily anomalies of 700 hPa
85 geopotential height south of 20° S. In this study, we used the daily index over the 2000–2022 period (https://www.cpc.ncep.noaa.gov/products/precip/CWlink/daily_ao_index/ao/ao.shtml, last visited: February 2023).

2.3 Snowpack temperature

The vertical profile of temperature in the upper snowpack is estimated with the surface energy budget and thermal diffusion model called Minimal Firn Model (~~<https://github.com/ghislainp/mfm>, last visited: February 2023~~ Picard et al., 2009, 2012; Domine et al., 2019) inspired by
90 (~~<https://github.com/ghislainp/mfm>~~, ~~<https://github.com/ghislainp/mfm>~~, last visited: February 2023, Picard et al., 2009, 2012; Domine et al., 2019) inspired by the Minimal Snow Model (Essery, 2004). The model takes as input the 2 m air temperature, surface pressure, 10 m wind speed, 2 m specific humidity, surface downwelling long-wave radiation flux, surface downwelling short-wave radiation flux, extracted from the [hourly](#) ERA5 reanalysis. Besides, the required snowpack properties are taken from the literature: the density at the surface and at 10 m depth (e.g. Leduc-Leballeur et al. (2015); Tian et al. (2018)), the thermal conductivity (Boone, 2002),
95 the specific heat capacity of ice (Picard et al., 2009) and the temperature at 10 m depth (equivalent to the mean annual air temperature in the Antarctic dry zones, Wang and Hou (2010)). From the MFM outputs, we used the temperature at 10 cm depth and the temperature gradient in subsurface, defined as the difference between temperatures estimated at 0 and 1 cm in depth (positive gradient when the surface is colder than the subsurface).

3 Methods

100 To investigate the atmospheric processes which generate the highest snow grain size occurrences over the East Antarctic Plateau, we used the GSI defined by Picard et al. (2012) as a proxy for the snow grain size. We selected four case studies. In particular, we studied the GSI summer changes in order to compare the mean variations with the variation during which occurs the highest GSI and we analysed the large-scale atmospheric circulation during each case study.

3.1 Detection of the highest GSI over East Antarctica

105 We define an extreme ~~event of grain growth~~ [grain size](#) as GSI being higher than 0.23. This value is the 99.5th percentile of all the maximum GSI over the dry area in Antarctica in the period 2000–2022 and is significantly higher (p -value = 0.012) than the mean annual maximum values of 0.11 ± 0.01 typical of the East Antarctica ice divide. Fig. 1 shows the maximum GSI over 2000–2022 derived from AMSU-B observations. Four extreme events have been ~~selected~~ [identified](#) (symbols in Fig. 1): the first in 2001 at 81.06° S, 63.12° E, near the Pole of Inaccessibility (Rees et al. (2021), referred to *A2001* as hereafter);
110 the second in 2007 at 79.64° S, 82.97° E, near Dome A (*B2007*); the third in 2016 at 76.60° S, 98.42° E, near Lake Vostok (*C2016*) and the fourth in 2020 at 77.38° S, 39.08° E, very close (~ 15 km) to Dome Fuji (*D2020*).

3.2 Identification of the seasonal increases in GSI

The annual onset and ending dates of the increase in GSI have been determined by means of the algorithm *Bayesian Estimator of Abrupt change, Seasonal change, and Trend* proposed by Zhao et al. (2019). In this algorithm, the **GSI** time-series are decomposed by numerous alternative ~~models, their~~ statistical models which separate the seasonal, trend and residual contributions. Their relative usefulness is quantified and they are combined into a better model via Bayesian model averaging. Thus, an estimation of the probabilities of change-point occurrences are provided by identification of abrupt changes in the slope of the GSI time-series. The seasonal grain size increase (*SGSI* from now on) is defined as the period which covers the interval from the onset to the ending dates.

We applied this algorithm on the 2020–2022 time-series at the four locations. Because of the GSI intrinsic variability and the presence of missing data, the onset and ending dates are not always identifiable. Thus, over a total of 88 summers (i.e. 22 years at four locations), both the onset and ending dates were identified in 55 cases (62.5%) and are used for our analysis.

3.3 Principal Component Analysis

Principal Component Analysis (PCA) is used to identify the main behaviour of variables involved in the SGSI onset and ending. It is performed by using the 55 SGSI identified in Section 3.2 with two configurations: in the first one, day 0 is the onset date and the analysis is performed between 10 days before and 20 days after day 0. In the second, day 0 is the ending date and analysis is performed from 20 days before to 10 days after day 0. PCA is applied to the daily time-series of GSI, skin temperature, 10 m wind speed, snowfall, surface pressure, cloud cover, ice temperature at 10 cm depth and the surface ice temperature gradient between 0 and 1 cm in depth. In order to apply the PCA algorithm, we used the *FactoMineR* library in R (<https://cran.r-project.org/web/packages/FactoMineR/index.html>).

3.4 Statistics of co-occurrences

To assess whether the SGSI onset date is related to an ~~atmospheric river occurrence or not~~ AR occurrence, we selected 1000 random dates in the period between November and December (when the onset usually happens) and we calculated the frequency of the ~~atmospheric rivers~~ ARs (defined by either the IWV or the vIVT classification) in a 5-day interval centred on those dates. Hence, an ~~atmospheric river is associated to~~ AR is associated with an onset date if it occurs in the interval [-2;+2] days around that date. We tested if the ~~atmospheric rivers~~ ARs interval follows a binomial distribution with both a probability p equal to its frequency and a parameter n equal to the number of onsets. From this, a p -value is obtained indicating if we can reject the null hypothesis that ~~atmospheric rivers~~ ARs and onsets occur independently.




Figure 1. Map of GSI maximum for the 2000–2022 period (colours) with the elevation contours for 3000 m and 3500 m (dotted lines); the locations of the four extreme grain size events detected are marked with asterisks; the melting area is masked.

4 Results

140 4.1 SGSI characterization

Fig. 1 shows that, in East Antarctica, the highest GSI occurred on the ice divide where elevation is higher than 3000 m and the slope is less than 0.1° (at 1 km resolution) (Slater et al., 2018).

The GSI time series of the four extreme events are reported in Fig. 2. A clear seasonal oscillation is observed at the four locations with the maximum occurring during summer, but with a large inter-annual variability.




Figure 2. GSI time series for the four selected locations from January 2000 to May 2022: a) A2001; b) B2007; c) C2016; d) D2020. The 0.23 extreme growth threshold (0.23) is shown in red lines.

145 The SGSI is characterized by an increase in GSI of 0.14 ± 0.01 on average with respect to the annual minimum, and it generally takes place between the end of November and the beginning of February. The following decrease in GSI tends to be slower and lasts until the next spring.

From 55 cases selected over 2000–2022 (cf. Section 3.2), we found that the SGSI lasts 54.9 ± 2.4 days on average, with a minimum of 28 days and a maximum of 114 days. The mean Julian day for the onset date is 336.8 ± 1.7 (i.e. 3 December) and for the ending date is 25.6 ± 2.0 (i.e. 26 January). The growing rate, defined as the total increase in GSI during the SGSI divided by its duration (in days), on average is $0.0026 \pm 0.0001 \text{ day}^{-1}$.

We Seasonal increase of the GSI is mainly driven by temperature and the subsequent increase of water vapor available to be transported within the snowpack upper layers (0–20 cm), triggering dry snow gradient metamorphism (Colbeck, 1982). Summer precipitation contributes to inter-annual variability through the deposition of a thin layer of fine grains at the surface, which increases albedo and reduces the penetration of solar radiation and the heating of the topmost layer, thereby reducing the grain growth (Picard et al., 2012). In the same way, the snow transport by wind results in a sorting of the grains by size in the topmost layers, as smaller snow grains are expected to fall out last (Grenfell et al., 1994; Domine et al., 2007). By contrast, the seasonal increase of the grains is favored by high subsurface temperature vertical gradients (Colbeck, 1993), mainly because of the solid-state greenhouse effect (Dombrovsky et al., 2019): the snow cools at the surface by emitting infrared radiation and warms a few centimeters below by absorbing solar radiation. This effect is particularly enhanced when the sky is clear.

For these reasons, we analysed variations of skin temperature, surface temperature gradient, temperature at 10 cm depth, snowfall, cloud cover and wind speed at 10 m during the SGSIs in the locations of the four extreme events. Surface pressure was included to explore the possibility of a common large-scale atmospheric regime. We computed the anomaly of selected meteorological variables with respect to the 2000–2022 period and we performed linear correlations with the seasonal maximum values of GSI. The correlation coefficients with the associated p-values are reported in Table 1. For the 55 selected SGSIs, a negative significant correlation is found with skin temperature (-0.41) and 10 m wind speed (-0.49), as well as a positive significant correlation with surface temperature gradient (0.39). Thus, wind and temperature appear to be the main drivers in determining the maximum GSI. Besides, despite the fact the inter-annual cycle of the grain size is driven by temperature, it emerges that during the SGSI a lower than usual skin temperature leads to larger grain size value.

Some weaker correlation but significant at the 90% level, are obtained with temperature at 10 cm depth, snowfall and cloud cover (-0.28, -0.26 and -0.25, respectively), suggesting that these factors can also be involved in the maximum GSI. Conversely, no significant correlation is found for with surface pressure.

Note that Picard et al. (2012) computed correlation between the increase in GSI during 1 December–15 January and accumulated precipitation during the same period at Dome C (75.10° S , 123.33° E), excluding the years with accumulation larger than 4 kg m^{-2} . Using the same approach, we obtained a correlation of -0.30 at our four locations over 2000–2022. They found a high correlation of -0.83 at Dome C over 1999–2010, but if we extend the calculation to the period 2000–2022, we obtain a much weaker correlation of -0.44 at Dome C. We also Another difference is that we are using the up-to-date ERA5 reanalysis, instead of ERA-Interim (Simmons et al., 2007).

Lastly, we studied the meteorological conditions during the SGSI onset and end by using the Principal Component Analysis applied to the selected 55 cases, as described in Section 3.3. Fig. 3a shows the first components for the beginning of the SGSI. GSI starts to rise around day 0, i.e. the onset date. An increase in surface pressure and in skin temperature is observed with a peak 1-2 days after the onset. The temperature gradient over the 10 cm under the surface features a decrease a few days after

Table 1. Linear correlation coefficients between some meteorological parameters anomalies and the maximum GSI of the identified 55 SGSI. Significance over 95% is grey-marked with two asterisks and over 90% is pale grey with one asterisk.

	Correlation	p-value
Wind speed 10 m	-0.49	0.00016**
Skin temperature	-0.41	0.00210**
Snowfall	-0.26	0.05905*
Cloud cover	-0.25	0.06484*
Surface temperature gradient	0.39	0.00366**
Temperature at 10 cm depth	-0.28	0.04129*
Surface pressure	-0.05	0.68960




Figure 3. First components of the Principal Component Analysis performed over GSI, skin temperature, 10 m wind speed, snowfall, surface pressure, cloud cover, temperature at 10 cm depth and the surface temperature gradient between 0 and 1 cm in depth during a) the onset and b) the end of the SGSI. The proportion of explained variance for each parameter is reported in the legend.

the onset, with a minimum at day 3, and then it increases, while temperature at 10 cm depth keeps increasing for the whole range of 30 days considered, with the maximum rate between day 0 and day 5. Snowfall has a sharp peak during day 0 and day 1, and at the same time, 10 m wind speed and cloud cover rise. Then these three parameters decrease and exhibit lower stable values after day 5. The proportions of variance explained by the first component (i.e. for each parameter the amount of its variance the first component explains) of the GSI and temperatures at skin level and at 10 cm depth are high (74%, 66% and 95%, respectively), a-bit-but they are lower for surface pressure and surface temperature gradient (34% and 37%) and even-much lower for wind speed, snowfall and cloud cover (13%, 26% and 13% respectively), probably-because-these. This is probably because the latter parameters have a much higher-larger natural day-to-day variability.

Fig. 3b reports the first components of PCA for the ending of the SGSI. GSI ~~first component~~ reaches its maximum at day 0, then sharply decreases at day 2. Note that during ~~2-3-2-3~~ days from day 2, all variables ~~have~~ change in behaviour: skin temperature and temperature at 10 cm depth ~~showed~~ ~~show~~ a long term decrease, but then the first rises (during days ~~1-3-1-3~~) and the second stabilizes; the opposite variations are observed for surface temperature gradient, which ~~exhibited~~ ~~exhibit~~ a decreases during days ~~1-5-1-5~~. Snowfall, wind speed, surface pressure and cloud cover, which were low and relatively stable, have a sharp peak during days ~~2-3-2-3~~. This particular combination seems to determine the seasonal halt of the snow grain growth. The proportions of variance explained are quite similar to the onset case, with the exception of the surface temperature gradient which is higher while the temperature at 10 cm depth and the GSI which are lower, indicating a greater variability of these variables during the SGSI end. These peaks in the snowfall occurring both at the onset and at the end of the SGSI suggested to explore more details their characteristics, as a relatively huge precipitation event could be linked to the occurrence of an AR. We investigate this in Section 4.3.

4.2 The four most extreme maximum grain size events

During the selected four extreme events (see Section 3.1), GSI increased by 0.24–0.27 between the onset and end dates, and reached values above 0.23 (Table 2). Moreover, the GSI remained significantly above the climatological average (2000–2022) for the whole summers and subsequent autumns, with extreme standard deviations of over $+3\sigma$ during these maximums. The duration of these events ranges from 39 to 90 days.

In order to evaluate the spatial extent of the regions featuring high GSI events, we ~~defined~~ ~~define~~ the extent of each of the 22 summer seasons as the area with the seasonal maximums GSI larger than 0.20, i.e. the 95th percentile of all the maximum GSI over the dry area in Antarctica in 2000–2022. On average, annually this area is $40\,000 \pm 10\,000$ km². The extent of the most extreme events ranges from 26 000 to 178 000 km². The magnitude, extent and growth rate of the A2001 and D2020 events are similar, while the B2007 and C2016 events feature a slightly larger maximum but smaller growth rate and spatial extent.

Table 2. Coordinates and elevation of the four extreme grain size events, and characterization of their SGSI.

Event	Coordinates	Elevation m	Extent km ²	Max GSI	SGSI	Duration days	Growing rate day ⁻¹
A2001	81.06° S, 63.12° E	3891	178 000	0.234	01/12/2000–25/01/2001	55	0.0046
B2007	79.64° S, 82.97° E	3914	26 000	0.253	05/12/2006–05/03/2007	90	0.0030
C2016	76.60° S, 98.42° E	3669	46 000	0.244	14/12/2015–14/02/2016	62	0.0039
D2020	77.38° S, 39.08° E	3794	140 000	0.237	09/12/2019–17/01/2020	39	0.0060

Fig. Figures 4 to 7 show GSI and some snow and atmospheric variables during the summer seasons of the most extreme events and their 2000–2022 climatology for comparison.

The A2001 event (Fig. 4) started on 1 December 2000 and the SGSI lasted until 25 January 2001. The GSI rate, defined
215 as the daily variation averaged over a 10-day window, was maximum one week after the onset, when the sky was mainly
clear. After the growth ending date, snowfalls became more frequent and the cloud cover and wind speed were often above the
climatological mean. In February, the GSI rate was about zero and began to become negative later in March.

The B2007 SGSI (Fig. 5) lasted ~~longer than the other extreme events, between from~~ 5 December 2006 ~~and to~~ 5 March 2007.
Despite the sudden decrease in GSI on 13 January coinciding with a large snowfall, GSI continues to increase after and the
220 SGSI lasted 90 days, which is much longer than the mean (54.9 days, cf 4.1) and the other extreme events (~~short shorter~~ than 62
days). Note that the GSI rate remained positive but decreased in January when more frequent snowfalls and higher wind speed
happened. ~~This events differs~~ ~~These events differ~~ from the other ~~extreme events~~ ~~extremes~~ by a lower mean snowfall anomaly
~~which of nearly zero~~ during the SGSI ~~was nearly zero~~ (-3% with respect to the 2000–2022 climatology), ~~unlike, against -50%~~
~~or below for~~ the other extreme events ~~which had very negative snowfall anomalies (equal or less than -50%)~~. Besides, excluding
225 the days immediately after the onset, the sky was generally cloudy during the whole SGSI (a non-significant +2% in total with
respect to climatology during the SGSI, while the anomalies for the other ~~extreme events~~ ~~extremes~~ were negative, from -7% to
-23%).

The C2016 event (Fig. 6) started on 14 December 2015 and stopped on 14 February 2016. The GSI rate was steadily positive
until mid-January, ~~and~~ then began to decrease. The wind was generally below the climatological mean, and snowfalls were
230 scarce. Starting from the end date, when a snowfall occurred, the wind speed suddenly rose above average, and remained
steadily high.

Finally, the onset date of the D2020 event was on 9 December 2019 (Fig. 7), and GSI increased until 17 January 2020.
The GSI rate was high (nearly +0.1/10 days) from the onset to the end of December, and the growing rate averaged over the
SGSI was the highest ~~with respect to the other of all~~ extreme events (see Table 2). The low wind speed condition around the
235 onset was favoured by a high pressure situation, above the climatological mean by nearly 20 hPa in the days right before the
onset (the highest value among the extreme events). Overall, calm wind condition was maintained for the rest of the SGSI,
recording the largest negative anomaly (-0.8 m s^{-1}) among the extreme events. ~~Actually, the~~ ~~The~~ onset occurred just after
a large precipitation event observed over the 8–~~10~~–~~10~~ December period (1.1 kg m^{-2} of snowfall), while the other extreme
events had smaller accumulations around the onset. Note that snowfall of 8 and 9 December 2019 were both above 0.4 kg m^{-2} ,
240 corresponding to a precipitation event ranking above the 95th percentile for daily snow accumulation as estimated by Dittmann
et al. (2016). Very ~~few little~~ snowfall was recorded during the rest of the SGSI. The wind speed increased at the SGSI end, and
in the following days wind speed above the climatology often occurred accompanied by snowfalls.

Table 3 summarises the anomalies in wind speed at 10 m, temperatures, snowfall, cloud cover, surface temperature gradient
and surface pressure during the SGSIs of the four most extreme grain size events. Even though these four cases had different
245 durations, some common features appear: the presence of snowfall precludes the onset accompanied by negative surface tem-
perature gradients, a low wind speed situation during a few days following the onset, with also low temperatures, especially at
skin level, and very few snowfalls and low cloud cover, except for the B2007 event (Fig. 5) which had frequent snowfall events.
No clear conclusion can be drawn for the surface pressure anomaly, since two events occurred with positive mean anomalies




Figure 4. The A2001 event timeseries daily from 1 November 2000 to 31 March 2001 of a) GSI, and b) the GSI rate over a 10 day window; c) temperature at 2 m (dark blue), skin temperature (blue), ice temperature at 10 cm depth (pale blue), and d) ice temperature gradient between 0 and -1 cm (grey line); e) wind speed at 10 m; f) surface pressure; g) cloud cover; h) snowfall. The 2000–2022 climatology of each variable is in dotted line. The SGSI period is in grey area.




Figure 5. As in Fig. 4 but for the B2007 event, from 1 November 2006 to 31 March 2007.

and two events with negative ones. However, in the 10-15-10-15 days around the onset, the pressure anomaly was positive in all cases.




Figure 6. As in Fig. 4 but for the C2016 event, from 1 November 2015 to 31 March 2016.

4.3 On the role of atmospheric rivers

The SGSI onset and end is-are generally associated with stormy conditions inducing large precipitation amounts, as shown in Fig. 3. Recently, Wille et al. (2021) demonstrated that atmospheric rivers (ARs)ARs, which facilitate the intrusion of relatively warm and moist air masses from the lower latitudes into Antarctica, generate extreme snowfall events. They also play a leading




Figure 7. As in Fig. 4 but for the D2020 event, from 1 November 2019 to 31 March 2020.

255 role in determining the Antarctic precipitation variability on inter-annual timescales, especially on the East Antarctica Plateau where the snowfall accumulation is overall low. They estimated that, in spite of their rareness (about 1% of the time), ARs are responsible for around 10–20% of the annual accumulation across East Antarctica ~~is caused by atmospheric rivers~~. Along the highest East Antarctic ice divide, large amount of snow can be generated under blocking anticyclonic circulations or ridging

Table 3. Mean anomalies with respect to the 2000–2022 climatology during the SGSI of the most extreme events. Significance over 95% is ~~grey~~ marked with two asterisks.

Event	Wind speed	Skin T	T 2 m	T -10 cm	Snowfall	Cloud cover	Surface T gradient	Surface pressure
A2001	-0.7 m s ⁻¹	-1.1 K	-1.0 K	-0.6 K	-50%	-23%	+0.3 K/m	+2.2 hPa
B2007	-0.7 m s ⁻¹	-1.8 K	-1.4 K	-1.3 K	-3%	+2%	+0.7 K/m	-0.9 hPa
C2016	-0.6 m s ⁻¹	-2.1 K	-1.7 K	-1.5 K	-60%	-7%	+0.8 K/m	-3.5 hPa
D2020	-0.8 m s ⁻¹	-1.2 K	-0.5 K	-0.4 K	-51%	-23%	+0.5 K/m	+1.9 hPa
Average	-0.7 m s ⁻¹ **	-1.6 K**	-1.2 K**	-1.0 K**	-41%	-13%	+0.6 K/m**	-0.1 hPa

situations which ~~accompany channel~~ the moist air ~~going to the pole from the mid-latitudes southwards, towards the interior~~ (Hirasawa et al., 2013; Pohl et al., 2021).

Here, we ~~studied study~~ the possible link between the ARs presence and the SGSI. Assessing the co-occurrence of ARs and the onset date requires to compare ~~the frequency of the atmospheric river their frequency~~ around the onset date, and perform a statistical test to determine if this frequency is significantly different from that obtained with random dates (cf. Section 3.4). ~~The mean frequencies of atmospheric river~~

~~In Table 4, for each ARs catalog and location, we reported the number of onsets available for each location (n), the frequencies of AR occurrence in 1000 random 5-day intervals ~~centred centered~~ between November and December in 2000–2022 ~~are ($p=9.3\%$, 6.8% , 7.2% , 8.9% for the A2001, B2007, C2016, D2020 locations, respectively. $m=5, 3, 5, 3$ onsets were associate) and the number of onsets associated~~ with at least one ~~atmospheric river AR~~ in the 5-day window centered on each onset ~~. Considering the number of onsets available for each location are $n=13, 17, 13, 12$, respectively and comparing~~ (m). ~~m are compared~~ to a binomial distribution with ~~these n and p , we obtained that the m have p -value $=0.0006, 0.0249, 0.0002$ and 0.0173 for location A2001, B2007, C2016 and D2020, respectively. It means that between the p -values are in Table 4. Between 2000 and 2022, the onsets and the atmospheric rivers ARs are very unlikely independent, except at the B2007 location where the relationship using the ERA5 catalogs is not statistically robust.~~ Thus, we can reject the null hypothesis that ~~atmospheric rivers ARs~~ and the onset of the grain SGSI are independent. However, ~~it is important to note that not all the onsets are associated to an atmospheric river. On average, for not associated with an AR. For the four locations, atmospheric rivers the probability that ARs are associated with the onset in 38%, 18%, 38% and 25% of the cases an onset ($P=m/n$) is $\sim 19\%$ and $\sim 30\%$ on average, respectively for ERA5 and MERRA-2 (Table 4).~~ Additionally, no ~~atmospheric river AR~~ approached the four locations during the onset of the most extreme SGSI investigated in Section 4.2. ~~Therefore, atmospheric rivers~~ Note that some discrepancies exist between reanalysis in Antarctica due to the scarce observations available to constrain ~~the models by assimilation. But the fact that the two reanalyses provide convergent results suggests that the signals are robust and reproducible. Therefore, ARs are sometimes involved in the onset, but they are not a required factor necessary condition.~~~~

and they do not play a role in determining the maximum seasonal grain size. ~~A similar analysis was conducted also~~ Similar analyses were conducted to assess the co-occurrence of ARs and the end date of the SGSIs ~~and results were quite similar. We obtained p -value = 0.0360, 0.0013 and 0.0001 for A2001, B2007 and D2020, respectively, and a non-significant 0.1444 for C2016. Hence, at least in~~ (Table 4) considering the months between January and March in 2000–2022. At least in 2 or 3 of the 4 locations, there is a significant connection between ARs and the ending of the SGSI, with the probability that ARs are associated with the ending dates being $\sim 20\%$ and $\sim 22\%$ on average, respectively for ERA5 and MERRA-2, but with larger variability with respect to the onset case.

<u>Location</u>	<u>Reanalysis</u>	<u>Onset</u>				<u>Ending</u>			
		<u>p</u>	<u>m</u>	<u>p-value</u>	<u>P</u>	<u>p</u>	<u>m</u>	<u>p-value</u>	<u>P</u>
A2001	<u>MERRA-2</u>	<u>9.3%</u>	<u>5</u>	<u>0.0006*</u>	<u>38%</u>	<u>5.0%</u>	<u>2</u>	<u>0.0360*</u>	<u>15%</u> -
	<u>ERA5</u>	<u>15.5%</u>	<u>4</u>	<u>0.0390*</u>	<u>31%</u>	<u>14.3%</u>	<u>3</u>	<u>0.1041</u>	<u>23%</u>
B2007	<u>MERRA-2</u>	<u>6.8%</u>	<u>3</u>	<u>0.0249*</u>	<u>18%</u>	<u>5.7%</u>	<u>4</u>	<u>0.0013*</u>	<u>24%</u> -
	<u>ERA5</u>	<u>5.3%</u>	<u>1</u>	<u>0.2268</u>	<u>6%</u>	<u>2.8%</u>	<u>4</u>	<u>0.0010*</u>	<u>24%</u>
C2016	<u>MERRA-2</u>	<u>7.2%</u>	<u>5</u>	<u>0.0002*</u>	<u>38%</u>	<u>5.5%</u>	<u>1</u>	<u>0.1444</u>	<u>8%</u> and-
	<u>ERA5</u>	<u>3.7%</u>	<u>2</u>	<u>0.0110*</u>	<u>15%</u>	<u>2.5%</u>	<u>0</u>	<u>0.2753</u>	<u>0%</u>
D2020	<u>MERRA-2</u>	<u>8.9%</u>	<u>3</u>	<u>0.0173*</u>	<u>25%</u>	<u>7.4%</u>	<u>5</u>	<u>0.0001*</u>	<u>42%</u> of the ending dates associated to an AR, respectively.
	<u>ERA5</u>	<u>9.4%</u>	<u>3</u>	<u>0.0208*</u>	<u>25%</u>	<u>8.5%</u>	<u>4</u>	<u>0.0021*</u>	<u>33%</u>

Table 4. Statistical parameters used in a binomial distribution to assess the significance of the co-occurrence of the ARs and the onset and ending dates. The meaning of each parameter is explained in the text. The p -values over 95% are marked with an asterisk.

5 Discussion

290 5.1 Grain growth

Our analysis shows that the amplitude of the summer increase of GSI is highly variable and large anomalies emerge occasionally during the summer, giving rise to very high values of grain size, localised in specific areas. We found some interactions between grain size evolution and local meteorological conditions at different time scales, from daily to seasonal, along the main East Antarctic ice divide. Wind and skin temperature have been shown to play a leading role in affecting snow metamorphism during the austral summer with ~~correlation of -0.49 and -0.41, respectively~~ significant correlation with the GSI. We observed that the beginning of the growth usually takes place just after a snowfall: at first, the grain size decreases during the precipitation, as new smaller grains are brought onto the surface (Domine et al., 2007). However, the snowfall is usually associated ~~to~~ with a large near-surface temperature gradient which is a driver of the dry snow metamorphism causing a rapid increase of the grain size (e.g. Colbeck (1982), Grenfell et al. (1994)).

300 The East Antarctic Plateau cannot be regarded as thermally homogeneous (González et al., 2021). For ~~example, cold air is transported~~ instance, katabatic winds transport cold air downhill from the domes and the ice divides ~~triggering katabatic winds,~~

and shallow depressions exist on the flank of the ice divides where pooling of drained cold air allows greater radiative heat loss and hence enhanced cooling of the surface (Scambos et al., 2018). Besides, because of ~~this top position~~their top positions, the ice divides are characterised by a general wind divergence (Parish and Bromwich, 2007) and weak wind conditions (Yamanouchi et al., 2003). In addition to the low accumulations, these features are favourable to the presence of large snow grains since high-speed winds tend to transport and deposit small ice particles on the surface, burying larger grains present. Moreover, lower than normal skin temperature observed during the SGSIs of the most extreme events, suggests that a large temperature gradient is possible in the first centimeters of the snowpack and hence an enhanced flux of water vapour could appear from the subsurface to the surface, leading to the formation of recrystallization crystals (Flanner and Zender, 2006; Champollion et al., 2013; Leduc-Leballeur et al., 2017). Cold surface conditions could result from the generally clearer than normal sky conditions observed during these phases, favorable to local intense radiative cooling of the surface and an enhanced solid-state greenhouse effect (Dombrovsky et al., 2019), causing a strong thermal gradient in the uppermost part of the snowpack, as observed in Gallet et al. (2014). The solid-state greenhouse effect could not be appreciated in this study as we used the python version of the MFM, in which its parametrization is absent. The low wind speed also hinders the dispersion into the atmosphere of the sublimated snow favouring local supersaturation conditions and, hence, the growing of grains.

5.2 Link with large scale mean conditions

As demonstrated by Wille et al. (2021), the zonal distribution of the 500 hPa geopotential height anomalies associated with ARs landfalls resembles the negative phase of the SAM. We thus analysed if a possible connection between GSI increase and large scale atmospheric circulation also exists. The 500 hPa geopotential height can help identify large scale atmospheric structures, such as the tropospheric polar vortex (Kwon et al., 2020; Gordon et al., 2022). Fig. 8 shows the ERA5 500 hPa geopotential height mean standardized anomalies (with respect to 2000–2022 climatology) for the 5-day intervals centered on the available onset dates for each location. There are mostly positive anomalies all over the continent, with significant values in the nearby of the considered locations, with waviness in the Southern Ocean. The same analysis was performed using MERRA-2 reanalysis dataset and provided similar results (not shown). For the four study cases, we also observe a positive anomaly in the Amundsen Sea, possibly related to a low activity of the Amundsen Sea Low (see for example Turner et al. (2012)).

Some tests were performed in order to assess a more frequent occurrence of particular zonal wave numbers during the 55 onsets but the results were not conclusive. Transient waviness (wavenumber 3 or 4) can be seen in Fig. 8 even though not at significant level, hence future work will study in more detail the possible involving of mid-latitude atmospheric dynamics and zonal wave activity.

As a matter of fact, studying Fig. 8 we can observe, in all the four cases, that there are some pressure differences between the continent and the ocean that are clearly reminiscent of the SAM. Thus, these seasonal events of grain size increases tend to correspond to negative SAM-like patterns. Further analysis of the Antarctic Oscillation index we found shows that the period October–~~November~~–December 2019 (the months before the D2020 event) recorded the lowest mean value of the AAO index since the beginning of the this time series in 1979 (Lim et al., 2021). This was the result of a minor sudden stratospheric warm-

ing (SSW) that occurred in September 2019 and propagated downwards, thereby causing anomalous tropospheric circulation from the end of October to mid-December 2019 (Shen et al., 2020). Such ~~stratospheric warming SSW~~ events are very rare in the Southern Hemisphere, as their occurrence has been estimated to 4% per year (Wang et al., 2020). However, this did not happen before the other three extreme snow grain size events. ~~Hence, the average AAO around the onset cases considered in this study is not statistically different from any other date during November-December.~~ During the four most extreme events most of the time (three cases out of four) a negative AAO index is present, i.e. a weak polar vortex situation. In particular during the 5-day intervals centred on the onsets of the four most extreme events we ~~have find~~ an average of the AAO index of -1.48 (A2001), 0.18 (B2007), -1.71 (C2016) and -2.09 (D2020). A negative AAO situation seems to ~~appear more preferential to trigger favor~~ SGSI. However, a negative AAO is neither necessary nor ~~is~~ sufficient. Not necessary as other onsets had very negative AAO values (not shown) and they did not ended in high GSI later in the season. Not sufficient ~~as for example the AAO since the AAO index~~ for the B2007 event is slightly positive. In fact, over the 55 selected cases, 31 had negative AAO mean values in the 5-day intervals centred on the onsets, with an average of -0.39 ± 0.19 over all the cases, with a *p-value* of 0.078 on a two-tailed t-test. Thus, a negative AAO index is more present than the positive phase during the onsets, even though this does not reach the 95% significance level.

In the Dome Fuji area during the anticyclonic circulation, a peculiar situation of warm-core eddy emerges near the top of the dome, while cooler air accumulates on the saddle on the eastern side, as described in Section 5.1. Indeed, on 5 December 2019, ERA5 reanalysis recorded the maximum gradient of about 4 K between the warm-core eddy and the saddle. Afterwards, temperature dropped until the beginning of January 2020, with an average anomaly of -4.3 K from 21 December 2019 to 4 January 2020, also observed in Fig. 7. The anticyclonic ridge developed when an intense wet oceanic air flow arrived over the ice sheet plateau through the Dronning Maud Land on 7–9 December and reached Dome Fuji, which became the center of the anticyclonic counterclockwise rotation of the flow. This event is not an AR according to the criteria defined in Wille et al. (2021), and neither the A2001 and B2007 events, when moisture intrusions arrived from Enderby Land and Amery Ice Shelf, respectively. Only the C2016 event is associated to an AR occurrence, which arrived from Victoria Land, on the basis of the ERA5 catalog, but not of the MERRA-2 catalog. Nonetheless, intense moisture advections were present during all these extreme events.

6 Conclusions

We studied the seasonal variations of the grain size in East Antarctica using an index computed using the 150 and 89 GHz observations from the satellite instrument AMSU-B, and previously introduced and analysed at Dome C (Picard et al., 2012). The present study covers the interval 2000–2022, the entire period of availability of the remote-sensed snow grain size index from AMSU-B sensors, over which we identified 55 seasonal grain size increases (SGSI) in four locations with identifiable onset and end dates. Using the ERA5 reanalysis, we established that grain size signal is linked to particular atmospheric conditions. In order to study these favourable circumstances, we identified four events of extreme increase of the grain size, in




Figure 8. Mean anomalies of the 500 hPa geopotential height fields from ERA5 over the period between 2 days before and 2 days after each synchronised onset date available for each location with respect to the 2000–2022 climatology. The number of available onset dates are 13, 17, 13 and 12, respectively. Brighter colours highlight the value with significance over 90% on a two-tailed t-test. Asterisks mark the locations: a) A2001 b) B2007 c) C2016 d) D2020.

different locations along the highest ice divide, and compared them to the usual SGSIs between 2000 and 2022. We searched for connections with local meteorological conditions and then with the synoptic atmospheric background.

370 The growth onset of the snow grains is usually linked to a snowfall event, in one third or one fifth (according respectively to MERRA-2 and ERA5) of the considered 55 cases associated with the presence of an atmospheric river providing warm and moist air into the inner plateau. The final maximum value of the grain size, that is usually reached in ~~January-February~~January-February, is determined by what happens in the weeks following the onset: the common features during the SGSI which emerge from this study are low wind speed conditions and skin temperature below the climatological mean.

375 We observed that ARs and, to a lesser extent, weak polar vortex conditions were related to the SGSI onset and end. However these situations are not necessary conditions. Another potential connection may exist with the Amundsen Sea Low, but our study did not allow to accurately characterize if a co-occurrence of different critical conditions could provoke a large GSI event. Further analysis could be done using the ERA5 reanalysis in order to extend the time interval and evaluate the significance of these anomalous situations and possibly to discover some other similar configurations which had led to intense growth of
380 the snow grain. The microwave data available in the past (e.g. Special Sensor Microwave Imager, SSM/I) miss the 150 GHz channel, but the 85–91 GHz channel could be used to find abrupt decreases in 1987–1999. However, the extension is not trivial as the grain size index is indirect and results from remote sensed observations. In situ observations should be included in the analysis, when available, in order to correct possible biases.

This analysis allowed improving the understanding of the causes of the SGSI onset and end, ~~and could~~and could be important
385 for projections of the frequency of extreme GSI events in a changing climate.

Data availability. ERA5 reanalysis data are available from <https://cds.climate.copernicus.eu/#/search?text=ERA5&type=dataset> (Accessed on 05-Aug-2022). The AMSU-B T_B values were retrieved at <https://www.ncei.noaa.gov/access/metadata/landing-page/bin/iso?id=gov.noaa.ncdc:C00981> (NCEI DSI 3702_01 dataset). AAO index were retrieved from https://www.cpc.ncep.noaa.gov/products/precip/CWlink/daily_ao_index/aao/aao.shtml.

390 *Author contributions.* CS led the study and performed the analysis, GM, MLL, GP, VF and BP supervised the discussion. MLL and GP helped provide the brightness temperature data and VF provided the atmospheric river datasets. All authors contributed to revise manuscript.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. Hersbach et al. (2018a, b) was downloaded from the Copernicus Climate Change Service (C3S) Climate Data Store. Vincent Favier acknowledge the support from Agence Nationale de la Recherche, project ANR-20-CE01-0013 (ARCA).

- Boone, A.: Description du schema de neige ISBA-ES (Explicit Snow), Note de Centre, Meteo-France/CNRM, 70, 53 pp, 2002.
- Casado, M., Landais, A., Picard, G., Arnaud, L., Dreossi, G., Stenni, B., and Prié, F.: Water Isotopic Signature of Surface Snow Metamorphism in Antarctica, *Geophysical Research Letters*, 48, <https://doi.org/10.1029/2021GL093382>, 2021.
- Champollion, N., Picard, G., Arnaud, L., Lefebvre, E., and Fily, M.: Hoar crystal development and disappearance at Dome C, Antarctica: observation by near-infrared photography and passive microwave satellite, *The Cryosphere*, 7, 1247–1262, <https://doi.org/10.5194/tc-7-1247-2013>, 2013.
- Colbeck, S. C.: An overview of seasonal snow metamorphism, *Reviews of Geophysics*, 20, 45, <https://doi.org/https://doi.org/10.1029/RG020i001p00045>, 1982.
- Colbeck, S. C.: The vapor diffusion coefficient for snow, *Water Resources Research*, 29, 109–115, <https://doi.org/10.1029/92wr02301>, 1993.
- Collow, A. B. M., Shields, C. A., Guan, B., Kim, S., Lora, J. M., McClenny, E. E., Nardi, K., Payne, A., Reid, K., Shearer, E. J., Tomé, R., Wille, J. D., Ramos, A. M., Gorodetskaya, I. V., Leung, L. R., O'Brien, T. A., Ralph, F. M., Rutz, J., Ullrich, P. A., and Wehner, M.: An Overview of ARTMIP's Tier 2 Reanalysis Intercomparison: Uncertainty in the Detection of Atmospheric Rivers and Their Associated Precipitation, *Journal of Geophysical Research: Atmospheres*, 127, <https://doi.org/doi:10.1029/2021jd036155>, 2022.
- Dittmann, A., Schlosser, E., Masson-Delmotte, V., Powers, J. G., Manning, K. W., Werner, M., and Fujita, K.: Precipitation regime and stable isotopes at Dome Fuji, East Antarctica, *Atmos. Chem. Phys.*, <https://doi.org/10.5194/acp-2015-1012>, 2016.
- Dombrovsky, L. A., Kokhanovsky, A. A., and Randrianalisoa, J. H.: On snowpack heating by solar radiation: A computational model, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 227, 72–85, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2019.02.004>, 2019.
- Domine, F., Salvatori, R., Legagneux, L., Salzano, R., Fily, M., and Casacchia, R.: Correlation between the specific surface area and the short wave infrared (SWIR) reflectance of snow, *Cold Regions Science and Technology*, 46, 60–68, <https://doi.org/10.1016/j.coldregions.2006.06.002>, 2006.
- Domine, F., Taillandier, A.-S., and Simpson, W. R.: A parameterization of the specific surface area of seasonal snow for field use and for models of snowpack evolution, *Journal of Geophysical Research*, 112, <https://doi.org/https://doi.org/10.1029/2006JF000512>, 2007.
- Domine, F., Picard, G., Morin, S., Barrere, M., Madore, J.-B., and Langlois, A.: Major Issues in Simulating Some Arctic Snowpack Properties Using Current Detailed Snow Physics Models: Consequences for the Thermal Regime and Water Budget of Permafrost, *Journal of Advances in Modeling Earth Systems*, 11, 34–44, <https://doi.org/https://doi.org/10.1029/2018MS001445>, 2019.
- Essery, R.: Parameter sensitivity in simulations of snowmelt, *Journal of Geophysical Research*, 109, <https://doi.org/doi:10.1029/2004JD005036>, 2004.
- Flanner, M. G. and Zender, C. S.: Linking snowpack microphysics and albedo evolution, *Journal of Geophysical Research*, 111, <https://doi.org/doi:10.1029/2005JD006834>, 2006.
- Gallet, J.-C., Domine, F., Savarino, J., Dumont, M., and Brun, E.: The growth of sublimation crystals and surface hoar on the Antarctic plateau, *The Cryosphere*, 8, 1205–1215, <https://doi.org/doi:10.5194/tc-8-1205-2014>, 2014.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *Journal of Climate*, 30, 5419–5454, <https://doi.org/https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.

- González, S., Vasallo, F., Sanz, P., Quesada, A., and Justel, A.: Characterization of the summer surface mesoscale dynamics at Dome F, Antarctica, *Atmospheric Research*, 259, 105–699, <https://doi.org/10.1016/j.atmosres.2021.105699>, 2021.
- 435 Gordon, A. E., Cavallo, S. M., and Novak, A. K.: Evaluating Common Characteristics of Antarctic Tropopause Polar Vortices, *Journal of the Atmospheric Sciences*, <https://doi.org/https://doi.org/10.1175/JAS-D-22-0091.1>, 2022.
- Grenfell, T. C., Warren, S. G., and Mullen, P. C.: Reflection of solar radiation by the Antarctic snow surface at ultraviolet, visible, and near-infrared wavelengths, *Journal of Geophysical Research*, 99, 18 669, <https://doi.org/10.1029/94JD01484>, 1994.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Sabater, J. M., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on pressure levels from 1979 to present, 440 <https://doi.org/10.24381/cds.bd0915c6>, 2018a.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Sabater, J. M., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on single levels from 1979 to present, <https://doi.org/10.24381/cds.adbb2d47>, 2018b.
- Hirasawa, N., Nakamura, H., Motoyama, H., Hayashi, M., and Yamanouchi, T.: The role of synoptic-scale features and advection in prolonged 445 warming and generation of different forms of precipitation at Dome Fuji station, Antarctica, following a prominent blocking event, *Journal of Geophysical Research: Atmospheres*, 118, 6916–6928, <https://doi.org/10.1002/jgrd.50532>, 2013.
- Jin, Z., Charlock, T. P., Yang, P., Xie, Y., and Miller, W.: Snow optical properties for different particle shapes with application to snow grain size retrieval and MODIS/CERES radiance comparison over Antarctica, *Remote Sensing of Environment*, 112, 3563–3581, <https://doi.org/doi:10.1016/j.rse.2008.04.011>, 2008.
- 450 Keenan, E., Wever, N., Dattler, M., Lenaerts, J. T. M., Medley, B., Munneke, P. K., and Reijmer, C.: Physics-based SNOWPACK model improves representation of near-surface Antarctic snow and firn density, *The Cryosphere*, 15, 1065–1085, <https://doi.org/https://doi.org/10.5194/tc-15-1065-2021>, 2021.
- Kingslake, J., Skarbek, R., Case, E., and McCarthy, C.: Grain-size evolution controls the accumulation dependence of modelled firn thickness, *The Cryosphere*, 16, 3413–3430, <https://doi.org/https://doi.org/10.5194/tc-16-3413-2022>, 2022.
- 455 Kwon, H., Choi, H., Kim, B.-M., Kim, S.-W., and Kim, S.-J.: Recent weakening of the southern stratospheric polar vortex and its impact on the surface climate over Antarctica, *Environmental Research Letters*, 15, 094 072, <https://doi.org/10.1088/1748-9326/ab9d3d>, 2020.
- Leduc-Leballeur, M., Picard, G., Mialon, A., Arnaud, L., Lefebvre, E., Possenti, P., and Kerr, Y.: Modeling L-Band Brightness Temperature at Dome C in Antarctica and Comparison With SMOS Observations, *IEEE Transactions on Geoscience and Remote Sensing*, 53, 4022–4032, <https://doi.org/10.1109/tgrs.2015.2388790>, 2015.
- 460 Leduc-Leballeur, M., Picard, G., Macelloni, G., Arnaud, L., Brogioni, M., Mialon, A., and Kerr, Y.: Influence of snow surface properties on L-band brightness temperature at Dome C, Antarctica, *Remote Sensing of Environment*, 199, 427–436, <https://doi.org/http://dx.doi.org/10.1016/j.rse.2017.07.035>, 2017.
- Lim, E.-P., Hendon, H. H., Butler, A. H., Thompson, D. W. J., Lawrence, Z. D., Scaife, A. A., Shepherd, T. G., Polichtchouk, I., Nakamura, H., Kobayashi, C., Comer, R., Coy, L., Dowdy, A., Garreaud, R. D., Newman, P. A., and Wang, G.: The 2019 Southern Hemisphere Stratospheric Polar Vortex Weakening and Its Impacts, *Bulletin of the American Meteorological Society*, 102, E1150–E1171, 465 <https://doi.org/10.1175/bams-d-20-0112.1>, 2021.
- Palermo, C., Kay, J. E., Genthon, C., L'Ecuyer, T., Wood, N. B., and Claud, C.: How much snow falls on the Antarctic ice sheet?, *The Cryosphere*, 8, 1577–1587, <https://doi.org/doi:10.5194/tc-8-1577-2014>, 2014.

- Parish, T. R. and Bromwich, D. H.: Reexamination of the Near-Surface Airflow over the Antarctic Continent and Implications on Atmospheric
470 Circulations at High Southern Latitudes, *Monthly Weather Review*, 135, 1961–1973, <https://doi.org/10.1175/mwr3374.1>, 2007.
- Picard, G., Fily, M., and Gallée, H.: Surface melting derived from microwave radiometers: a climatic indicator in Antarctica, *Annals of
Glaciology*, 46, 29–34, <https://doi.org/doi:10.3189/172756407782871684>, 2007.
- Picard, G., Brucker, L., Fily, M., Gallée, H., and Krinner, G.: Modeling time series of microwave brightness temperature in Antarctica,
Journal of Glaciology, 55, 537–551, <https://doi.org/doi:10.3189/002214309788816678>, 2009.
- 475 Picard, G., Domine, F., Krinner, G., Arnaud, L., and Lefebvre, E.: Inhibition of the positive snow-albedo feedback by precipitation in interior
Antarctica, *Nature Climate Change*, 2, 795–798, <https://doi.org/10.1038/nclimate1590>, 2012.
- Pohl, B., Favier, V., Wille, J., Udy, D. G., Vance, T. R., Pergaud, J., Dutrievoz, N., Blanchet, J., Kittel, C., Amory, C., Krinner, G., and Codron,
F.: Relationship Between Weather Regimes and Atmospheric Rivers in East Antarctica, *Journal of Geophysical Research: Atmospheres*,
126, <https://doi.org/10.1029/2021jd035294>, 2021.
- 480 Rees, G., Gerrish, L., Fox, A., and Barnes, R.: Finding Antarctica’s Pole of Inaccessibility, *Polar Record*, 57,
<https://doi.org/doi:10.1017/S0032247421000620>, 2021.
- Scambos, T. A., Campbell, G. G., Pope, A., Haran, T., Muto, A., Lazzara, M., Reijmer, C. H., and Broeke, M. R.: Ultralow Surface Tem-
peratures in East Antarctica From Satellite Thermal Infrared Mapping: The Coldest Places on Earth, *Geophysical Research Letters*, 45,
6124–6133, <https://doi.org/10.1029/2018gl078133>, 2018.
- 485 Shen, X., Wang, L., and Osprey, S.: Tropospheric Forcing of the 2019 Antarctic Sudden Stratospheric Warming, *Geophysical Research
Letters*, 47, <https://doi.org/10.1029/2020gl089343>, 2020.
- Simmons, A., Uppala, S., Dee, D., and Kobayashi, S.: ERA-Interim: New ECMWF reanalysis products from 1989 onwards, ECMWF
Newsletter, <https://doi.org/10.21957/pocnex23c6>, 2007.
- Slater, T., Shepherd, A., McMillan, M., Muir, A., Gilbert, L., Hogg, A. E., Konrad, H., and Parrinello, T.: A new digital elevation model of
490 Antarctica derived from CryoSat-2 altimetry, *The Cryosphere*, 12, 1551–1562, <https://doi.org/10.5194/tc-12-1551-2018>, 2018.
- Sturm, M. and Benson, C. S.: Vapor transport, grain growth and depth-hoar development in the subarctic snow, *Journal of Glaciology*, 43,
42–59, <https://doi.org/10.3189/s0022143000002793>, 1997.
- Thompson, D. W. J. and Wallace, J. M.: Annular Modes in the Extratropical Circulation. Part I: Month-to-Month Variability, *Journal of
Climate*, 13, 1000–1016, [https://doi.org/10.1175/1520-0442\(2000\)013<1000:AMITEC>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<1000:AMITEC>2.0.CO;2), 2000.
- 495 Tian, Y., Zhang, S., Du, W., Chen, J., Xie, H., Tong, X., and Li, R.: Surface snow density of East Antarctica derived from in-situ
situ, *The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, XLII-3, 1657–1660,
<https://doi.org/10.5194/isprs-archives-xlii-3-1657-2018>, 2018.
- Town, M. S., Waddington, E. D., Walden, V. P., and Warren, S. G.: Temperatures, heating rates and vapour pressures in near-surface snow at
the South Pole, *Journal of Glaciology*, 54, 487–498, <https://doi.org/10.3189/002214308785837075>, 2008.
- 500 Turner, J., Phillips, T., Hosking, J. S., Marshall, G. J., and Orr, A.: The Amundsen Sea low, *International Journal of Climatology*, 33, 1818–
1829, <https://doi.org/https://doi.org/10.1002/joc.3558>, 2012.
- Turner, J., Phillips, T., Thamban, M., Rahaman, W., Marshall, G. J., Wille, J. D., Favier, V., Winton, V. H. L., Thomas, E., Wang, Z.,
Broeke, M., Hosking, J. S., and Lachlan-Cope, T.: The Dominant Role of Extreme Precipitation Events in Antarctic Snowfall Variability,
Geophysical Research Letters, 46, 3502–3511, <https://doi.org/10.1029/2018gl081517>, 2019.
- 505 Wang, L., Hardiman, S. C., Bett, P. E., Comer, R. E., Kent, C., and Scaife, A. A.: What chance of a sudden stratospheric warming in the
southern hemisphere?, *Environmental Research Letters*, 15, 104 038, <https://doi.org/10.1088/1748-9326/aba8c1>, 2020.

- Wang, Y. and Hou, S.: Spatial distribution of 10 m firn temperature in the Antarctic ice sheet, *Science China Earth Sciences*, 54, 655–666, <https://doi.org/DOI: 10.1007/s11430-010-4066-0>, 2010.
- 510 Wille, J. D., Favier, V., Gorodetskaya, I. V., Agosta, C., Kittel, C., Beeman, J. C., Jourdain, N. C., Lenaerts, J. T. M., and Co-dron, F.: Antarctic Atmospheric River Climatology and Precipitation Impacts, *Journal of Geophysical Research: Atmospheres*, 126, <https://doi.org/10.1029/2020jd033788>, 2021.
- Yamanouchi, T., Hirasawa, N., Hayashi, M., Takahashi, S., and Kaneto, S.: Meteorological characteristics of Antarctic inland station, Dome Fuji, *Memoirs of National Institute of Polar Research. Special issue*, 57, 94–104, 2003.
- 515 Zhao, K., Wulder, M. A., Hu, T., Bright, R., Wu, Q., Qin, H., Li, Y., Toman, E., Mallick, B., Zhang, X., and Brown, M.: Detecting change-point, trend, and seasonality in satellite time series data to track abrupt changes and nonlinear dynamics: A Bayesian ensemble algorithm, *Remote Sensing of Environment*, 232, 111 181, <https://doi.org/10.1016/j.rse.2019.04.034>, 2019.