

Coupled ice/ocean interactions during the future retreat of West Antarctic ice streams in the Amundsen Sea Sector

David T. Bett¹, Alexander T. Bradley^{1,2}, C. Rosie Williams¹, Paul R. Holland¹, Robert J. Arthern¹ and
5 Daniel N. Goldberg³

¹British Antarctic Survey, Cambridge, UK

²Cambridge Zero, Cambridge, UK

³University of Edinburgh, Edinburgh, UK

Correspondence to: David T. Bett (d.bett@bas.ac.uk)

10 **Abstract.**

The Amundsen Sea sector has some of the fastest-thinning ice shelves in Antarctica, caused by high, ocean-driven basal melt rates, which can lead to increased ice stream flow, causing increased sea level rise (SLR) contributions. In this study, we present the results of a new synchronously coupled ice-sheet/ocean model of the Amundsen Sea sector. We use the WAVI ice sheet model to solve for ice velocities and the MITgcm to solve for ice thickness and three-dimensional ocean
15 properties, allowing for full mass conservation in the coupled ice/ocean system. The coupled model is initialised in the present day and run forward under idealised warm and cold ocean conditions with a fixed ice front. We find that Thwaites Glacier dominates the future SLR from the Amundsen Sea sector, with a SLR that is approximately quadratic in time. The future evolution of Thwaites Glacier depends on the life-span of small pinning points that form during the retreat. The rate of melting around these pinning points provides the link between future ocean conditions and the SLR from this sector and will be difficult
20 to capture without a coupled ice/ocean model. Grounding-line retreat leads to a progressively larger Thwaites ice-shelf cavity, leading to a positive trend in total melting, resulting from the increased ice basal surface area. Despite these important sensitivities, Thwaites Glacier retreats even in a scenario with zero ocean-driven melting. This demonstrates that a tipping point may have been passed in these simulations and some SLR from this sector is now committed.

1 Introduction

25 The West Antarctic Ice Sheet (WAIS) is a marine ice sheet of particular importance for future sea level rise (SLR), contributing ~6.5 mm to global SLR between 1992 to 2021 (Otosaka et al., 2023). Within the WAIS, the Amundsen Sea sector has seen the largest SLR contribution over the satellite era (Shepherd et al., 2019). The region experienced a 77% ice mass flux increase from 1973 to 2013, with Thwaites Ice Shelf specifically observing a 33% speedup during this period (Mouginot et al., 2014).

30 The Amundsen Sea sector's ice sheet is grounded on bathymetry below sea level, and the presence of retrograde slopes in the bathymetry may make the region susceptible to rapid and sustained retreat in the future (Favier et al., 2014). Any reduction in ice shelf buttressing via pinning points and side drag can lead to acceleration in ice shelf speed and retreat of the grounding line (Thomas, 1979). Ocean melting can reduce buttressing by thinning ice shelves, reducing side drag and aiding the ungrounding of pinning points. Of particular concern is Thwaites Glacier, which has a unconfined ice shelf, whose current
35 pinning point on its eastern ice shelf appears to be weakening, and has been predicted to unground within decades (Wild et al., 2022).

In the Amundsen Sea, warm modified Circumpolar Deep Water (CDW) resides below a colder and fresher layer of Winter Water (Jacobs et al., 1996), and flows through bathymetric troughs to the base of ice shelves (Walker et al., 2007), causing
40 high melt rates. The region experiences large decadal variability in its ocean conditions, most notably in the thickness of the modified CDW layer (Jenkins et al., 2018; Dutrieux et al., 2014). In addition, it has been suggested that there is an average anthropogenic warming trend superimposed on this internal variability (Holland et al., 2022; Naughten et al., 2022). Future anthropogenic warming of the Amundsen Sea is a key mechanism by which human activities may influence SLR from the Antarctic Ice Sheet (Holland et al., 2022; Jourdain et al., 2022; Holland et al., 2019).

45 Previous studies have used uncoupled ice-only simulations to simulate the future evolution and retreat of the Amundsen Sea sector and WAIS (e.g. Yu et al., 2018; Alevropoulos-Borrill et al., 2020; Feldmann and Levermann, 2015; Reese et al., 2020). However, these studies use ocean melting parametrisations, that contain simplifications of important ocean physics such as, Coriolis force, ocean mixing parameterisations, barotropic ocean flow, and lateral variation in the direction parallel to the
50 grounding line and hence lack spatial variation in melt rates caused by differences in ocean velocity and temperature. Therefore, these models incompletely represent the complex interactions between ice shelf geometry, ocean dynamics and melt rates, potentially leading to overestimations in rates of grounding line retreat and mass loss (Seroussi et al., 2017; De Rydt and Gudmundsson, 2016). To accurately simulate ice evolution, a coupled ice/ocean model must be used.

55 Previous coupled modelling studies have used different approaches to the coupling using either an 'asynchronous' (e.g. Seroussi et al., 2017; De Rydt and Gudmundsson, 2016; Naughten et al., 2021) or a 'synchronous' coupling (e.g., Goldberg and Holland, 2022; Goldberg et al., 2018; Jordan et al., 2018). Synchronous coupling involves continuously changing the ice geometry during the ocean simulation at the ocean model timestep, while asynchronous coupling involves information being exchanged every one or few ice model timesteps, with the ice geometry and ocean state modified at each coupling period.
60 Regional coupled models have been used to simulate parts of the Amundsen Sea sector, for example simulating Thwaites Glacier over 50 years (Seroussi et al., 2017), and the Pope, Smith and Kohler glaciers (Goldberg and Holland, 2022).

In this study we use a new synchronously coupled ice/ocean model of the Amundsen Sea sector to simulate the evolution of its ice streams over the next 180 years. We consider three different idealised forcing scenarios: no basal melting, and cold and warm Amundsen Sea conditions. We use these simulations to explore both the future evolution of the ice sheet in this sector, and the physical processes that determine the speed of the retreat. This allows us to better understand the mechanisms by which the ice loss is sensitive to future ocean conditions, and thus to anthropogenic forcing.

2 Methods

2.1 Ice model

In the coupled ice/ocean model, ice velocities are calculated using the Wavelet-based, Adaptive-grid, Vertically Integrated ice sheet model, WAVI (Arthern et al., 2015; Arthern and Williams, 2017). WAVI is a finite volume ice sheet model including a treatment of both membrane and simplified vertical shear stresses as described by (Goldberg, 2011). We use a numerical model with a time step of 20 days and a 2 km horizontal resolution covering the whole Amundsen Sea sector domain shown in Figure 1a.. This horizontal resolution is found to be appropriate in a recent study, currently under review, testing the impact of resolution on grounding line retreat and SLR contributions, with a nearly identical configuration of the Amundsen Sea sector (Williams et al., PREPRINT) (<https://doi.org/10.21203/rs.3.rs-3405435/v1>). The ice rheology is described using Glen's flow law, with an exponent of $n = 3$. A Weertman' sliding law is used (Weertman, 1964), for which basal sliding drag scales with the cube root of sliding velocity, multiplied by a basal sliding drag coefficient C . As described in Arthern et al. (2015), WAVI uses a data assimilation method to match modelled ice velocities and rates of thickness changes with observations of the Measures 2014/15 surface velocities (Mouginot et al., 2017a; Mouginot et al., 2017b) and rate of change of surface elevation from Smith et al. (2020), resulting in an initial state representing conditions in approximately 2015. Partially grounded cells are utilised to better represent the grounding line, where the grounding fraction is used to proportionally apply the Weertman sliding drag coefficient. An ice model relaxation is then run for a set period of time (4000 years). During this relaxation the grounding line and the thickness of ice shelves remains fixed, but the grounded ice thickness is allowed to change (see Arthern et al. 2015 for full details). This brings the flux divergence into much better agreement with observations of accumulation and rates of ice thickness change but at the cost of the surface elevation and ice velocities agreeing less well with observations. Ice thickness prior to relaxation and bathymetry fields are from BedmachineV3 (Morlighem et al., 2020; Morlighem, 2022), where a minimum thickness of 50 m is applied to determine the initial ice extent. The ice front and outer catchment boundaries are kept fixed throughout the simulations. Accumulation and englacial temperatures are kept constant in the forward simulations, using data sets from Arthern et al. (2006) and Pattyn (2010) respectively.

2.2 Ocean model

We use the Massachusetts Institute of Technology general circulation model (MITgcm; (Marshall et al., 1997)) to simulate the ocean circulation in the Amundsen Sea. The model grid uses polar stereographic Cartesian coordinates with horizontal resolution of 2 km and vertical resolution of 20 m, and we use a time step of 200 seconds. The ocean model domain extent is shown by the black box in Figure 1a. Note that this does not include the whole continental shelf, so we apply ocean boundary conditions and velocities to the northern and western open boundaries of the domain to impose prescribed Amundsen Sea conditions in idealised ‘warm’ and ‘cold’ experiments. We also examine a third ‘no melting’ case in which no ocean-driven melting is applied to the ice shelves. This study focuses solely on the ice sheet, ice shelf and ocean interactions driven by wider Amundsen Sea conditions, so sea ice and other freshwater sources/sinks are not included in the simulations, and no atmospheric forcing is applied over the model domain.

The boundary conditions in warm and cold experiments are as follows. Following previous studies, an idealised, piecewise linear, vertical profile is applied to replicate the warm CDW layer below cold Winter Water, with a thermocline 400 m thick (De Rydt et al., 2014). The base of the thermocline is placed at 600 m depth in the warm scenario and lowered to 800 m depth in the cold scenario (Figure 1c, d). These profiles correspond to the warmest and coldest observed Amundsen Sea conditions, 2009 and 2012 respectively (Dutrieux et al., 2014). On the northern boundary the warm CDW layer has temperature of 1.2°C and a salinity of 34.7 (PSU), whereas the western boundary is forced with a more modified CDW layer, with temperature 0.6°C and salinity 34.6 (PSU) (Figure 1c, d). In all simulations, at the boundaries, we apply average ocean velocities (1965-2015) from a larger regional model (Holland et al., 2022; Naughten et al., 2022), and we restore sea level to zero.

Ocean only simulations, with fixed initial ice geometry, are simulated for 2 years, as a spin up before coupling. The resultant conditions in the ocean model are designed to match the spatial distribution of present day observed maximum subsurface temperatures over the region (Dutrieux et al., 2014), as shown in Figure 2a,b. The warm and cold oceanic forcing cases enable different amounts of CDW to reach the base of the present day Thwaites and Pine Island Glacier (PIG) ice shelves (Figure 2e, f, i, j). The initial melt rates, after the 2 year ocean only spin-up, of the Thwaites and PIG ice shelves for the warm and cold simulations are shown in Figure 2c, d, g, h.

Ice shelf melting is represented by a standard three-equation formulation (Holland and Jenkins, 1999). We tune the dimensionless ice shelf melting drag coefficient in this parametrization to 0.008. This drag coefficient parameterises the ocean stress on the ice base as a function of the ocean model’s mixed layer velocities, in order to calculate turbulent ocean heat and salt fluxes for use in the melting calculation (Jenkins et al., 2010). For PIG and Thwaites ice shelves combined, this value produces the closest average match between the initial MITgcm ice shelf melt rates and those that are implicit in the WAVI initialisation (see Appendix A). This minimises the ‘coupling shock’ - the response of the ice model to any mismatch between

these two fields -, which occurs when the coupled model simulation commences. Without this calibration, the ice sheet trajectory could be impacted, potentially for many decades (Goldberg and Holland, 2022), by the adjustment of the ice due to the transition from implicit initialised melt rates to arbitrarily different ocean model melt rates at the start of the simulation. Note also that this tuned value is close to a 0.01 value derived from observations (Jenkins et al., 2010).

130

Bedmachine version 3 is used as the basis for the bed/seabed geometry throughout the model. However, we found that without modification this dataset led to rapid grounding and advance of PIG. Closer inspection revealed that the seabed bathymetry may have been underestimated near the grounding line of this glacier in Bedmachine version 3 (Appendix B). Therefore, throughout the domain we deepen the seabed wherever needed to achieve a minimum water column thickness of 280 m on the staggered Arakawa C-grid's velocity grid points, located on the grid faces, tapering towards grounded ice over 6 km down to a minimum water column thickness of 140 m. This procedure is only applied to cells with no ice basal sliding drag in the initial state and is only done once at the start of the simulation before the WAVI ice sheet model is initialized and relaxed, rather than being an ongoing process. This edited bathymetry is used in both MITgcm and WAVI.

135

140 **2.3 Ice/Ocean coupling**

This study uses a new, synchronously coupled ice/ocean model, which combines the WAVI ice sheet model with the MITgcm. Crucially, this coupling occurs through the MITgcm ice sheet package STREAMICE (Goldberg and Heimbach, 2013), making use of previous synchronous coupling developments (Jordan et al., 2018; Goldberg et al., 2018), which enables the STREAMICE ice thickness to evolve in the ocean model. The basic concept of the new model is that WAVI solves for the ice velocities, MITgcm STREAMICE solves for the ice thickness, and the MITgcm ocean model solves for all ocean properties, including melting and is also where the ice thickness evolves. Dividing up computations in this way allows for a fully synchronous coupling because both the ice thickness and ocean properties are solved within the MITgcm on the ocean timestep. This has two advantages: firstly, the ice thickness and ocean free surface equations are solved simultaneously, allowing full conservation of mass in the ice and ocean coupled system. Secondly, the melt rate responds instantly to changes in ice thickness.

145

150

In principle, this approach is no more expensive than other coupling approaches because the two-dimensional ice thickness equations can be solved on the ocean timestep with negligible computational expense compared to that incurred in solving three-dimensional ocean equations. However, the approach does require the MITgcm grid to exist wherever ice may go afloat during the simulation, since grounding-line retreat in MITgcm is accomplished naturally. MITgcm solves the ocean free surface equation every ocean timestep in order to conserve mass, which naturally inflates the water column in the MITgcm ocean model wherever the pressure loading of the ice decreases below floatation (Goldberg et al., 2018). In regions of ice which are not floating, a thin subglacial layer exists is specified in order to enable the expansion of the ocean column during

155

grounding line retreat (Goldberg et al., 2018; Jordan et al., 2018). We set it to be 4 m thick but could have been set to any
160 relatively small thickness compared to the ocean model vertical resolution. This small value has been previously demonstrated
to have no impact on the evolution of the coupled system (Goldberg et al., 2018). This layer, which has no effect on basal
sliding drag computed by the ice model, is treated as a porous medium, with Darcy flow used to connect the subglacial cells
to each other (Goldberg et al., 2018).


165 Using an MITgcm domain covering the entire catchment of Amundsen Sea sector would be very inefficient, since it would
include large areas of grounded ice that would never go afloat in the ocean/STREAMICE grid. To avoid this, we introduce the
concept of ‘ice-only’ and ‘coupled’ domains (Figure 1a): In the ice-only domain, WAVI is used to solve for both ice sheet
velocities and thickness, while in the coupled domain, WAVI solves for ice velocities and MITgcm solves for ice thickness
and ocean properties, as described above. The coupled domain only needs to extend far enough inland to accommodate the
170 grounding-line retreat occurring during a projection, which for this study was determined using test simulations.

The coupled model solution procedure is split into coupling periods, chosen to equal the WAVI timestep of 20 days, meaning
that the WAVI model state is fixed during each coupling period. Over each coupling period (Figure 1b) the following takes
place: (1) WAVI calculates ice velocities over the whole ice domain for one 20-day timestep; (2) WAVI steps forward the ice
175 thickness in the ice-only domain for one 20-day time step; (3) MITgcm receives the ice velocity for the coupled domain and
the updated ice thickness on its boundaries, and sub-cycles the ice thickness, grounding line and ocean properties within the
coupled domain using the 200 s ocean timestep for the full 20 day coupling period; (4) MITgcm passes the new ice thickness
and therefore the new ice grounding line for the coupled domain back to WAVI and the next coupling period commences.
Therefore, the choice of the length of the coupling period determines the fastest response time for which the ice velocities can
180 respond to changes in buttressing in the coupled domain, where the ice thickness changes with the ocean timestep. The WAVI
boundary cells outside of the coupled domain are passed to the MITgcm domain and held fixed while MITgcm runs, and then
the thickness throughout the coupled domain is passed from MITgcm to WAVI after each coupling step. This procedure, in
addition to WAVI being fixed during each coupling period, keeps the two models/domains ice thicknesses from diverging and
ensures a smooth transition between the two domains.


185

For consistency, the same advection scheme that is used in WAVI (Arthern et al., 2015), has been coded into STREAMICE.
Ice divergences are updated with the changing ice thickness in STREAMICE every ocean timestep, while the ice velocity
remains fixed over each coupling period. We adopt the principle that the ocean may only melt ice that has no basal sliding
190 drag applied (Arthern and Williams, 2017), so the WAVI ice basal sliding drag field is passed to MITgcm to decide where
melting can occur during each coupling period. This means that no melting can occur on partially grounded cells, as
recommended by Seroussi and Morlighem (2018). In addition, this means that if a cell becomes fully ungrounded in MITgcm,

ice shelf melting only occurs once the coupling period finishes and a new one starts, which updates the ice thickness in WAVI, and subsequently passes back a new basal sliding drag field. However, if, during a coupling period, a grid cell becomes grounded in MITgcm, melting on this cell is immediately switched off. In both the WAVI and STREAMICE models a minimum ice thickness of 50 m is applied.



200 **FIGURE 1:** (a) Outline of the whole Amundsen Sea sector domain. The black box shows the outline of the coupled domain; the ice only domain is defined as everywhere outside of this box. The coloured regions represent the three regions used in analysis: PIG (purple), Thwaites (red), and Smith (cyan). These areas are edited by hand from the Zwally et al. (2012) basins. (b) A schematic diagram of the three grouped steps of the coupled model applied during one coupling period, which are repeated during the coupled simulations. (c) Potential temperature boundary conditions for the Northern (solid line) and Western (dashed line) boundaries for the warm (red) and cold (blue) cases. (d) as in 205 (c) but for salinity.



210 **FIGURE 2:** (a) Initial maximum subsurface potential temperature over the ice/ocean domain for cold case. (c) Initial melt rate under PIG Ice Shelf in the cold forcing case over the green box in (b). The label shows the total initial meltwater flux. (e) Cross-section through PIG Ice Shelf taken along the west-east green line in (a) showing initial potential temperature in the cold case. (g) Initial melt rate over Thwaites Ice Shelf in the cold forcing case over the red box in (b). (i) Cross-section through Thwaites Ice Shelf taken along the south-north red line in (a), showing initial potential temperature in the cold case. (b), (d), (f), (h), (j), as in (a), (c), (e), (g), (i) respectively, but for the warm forcing case.

3 Results

215 3.1 Amundsen Sea sector: 180 years simulation




FIGURE 3: Warm forcing case evolution over 180 years. Panels show snapshots of (a) ice thickness, (b) ice speed and (c) melt rates plotted on a logarithmic scale, taken every 36 years throughout the simulation. Starting (black) and final (red) grounding lines are included.

220

Following initialisation, the coupled ice/ocean model is run forward for 180 years under both forcing cases, as well as a zero-melting case. The evolution of the warm simulation is shown in Figure 3. The most prominent feature of the simulation is the acceleration (figure 3b) and grounding-line retreat of Thwaites Glacier (figure 3a). At the start of the simulation, Thwaites Ice Shelf hosts the observed fast-flowing western shelf and slower eastern ice shelf (figure 3b), where a pinning point restrains the flow of ice. However, as the simulation progresses, both sides of the ice shelf accelerate, reaching speeds of up to ~ 10 km/yr, but then shows signs of deceleration between 144 years and 180 years. In addition, Thwaites Glacier experiences substantial grounding-line retreat during this simulation, approximately 130 km of north-south retreat, leading to a reduction in the total mass of grounded ice and the formation of new larger ice shelf cavity, which features rapid melting rates near the grounding line (figure 3c). The fixed ice front leads to a large and thin future Thwaites Ice Shelf, which may lead to artificially elevated melting overall, though only low ice shelf melt rates occur on the thinner ice (figure 3a,c). However, such trends in ice speed and grounding line retreat are restricted to Thwaites: neither PIG nor Smith Glacier experiences the same level of acceleration or retreat, although we do observe an initial acceleration of the PIG (which is later reversed), and some grounding line retreat in both PIG and Smith.

225

230




FIGURE 4: Timeseries shown for the three cases no melt (black line), cold (blue line), and warm (red line). (a) Cumulative global sea level contribution from the Thwaites area. (b) Rate of change of global sea level contribution for Thwaites area. (c) Total ice shelf melt rate from

235

the Thwaites area, with a 2-year running average applied. (d) Cumulative grounded area loss from the Thwaites area (e) – (h) Same as above but for the PIG area. (i) – (l) Same as above but for the Smith area.

240 In order to analyse SLR contributions, melting and grounded ice area loss from different regions, and to compare the three
forcing cases, we divide the domain into three areas: Thwaites, PIG and Smith (Figure 1a). These areas are edited from the
(Zwally et al., 2012) basins, where the Smith area has been separated out. In addition, the boundary between the PIG and
Thwaites areas is edited by hand to address the otherwise mis-assignment of SLR contributions, melting and grounding ice
245 melting case for the Thwaites and Smith areas, which has important implications. This result demonstrates that the initialised
ice-sheet state in WAMI is intrinsically unsteady and will continue the present-day ice loss for some period of time irrespective
of future climatic forcing. This implies that a tipping point may have been passed at some point for this particular initialised
ice sheet, likely during the 20th century (Mouginot et al., 2014), and some amount of SLR (at least ~40 mm in our simulations)
is now committed. Zero melting can never occur in the real world, because even seawater at the surface freezing point will
250 drive melting at depth due to the pressure decrease in the freezing temperature, so this is a very conservative test of the presence
of committed ice change. However, the contribution from PIG is dependent on the particular choice of bathymetry deepening
that is implemented (Appendix B). Notably, we observe the largest zero-melting SLR contribution from the Thwaites area,
leading to ~45 mm in 180 years, with a near constant SLR rate. For the Thwaites area we also obtain some grounding line
retreat, though the rate of loss reduces during the simulation, suggesting that the SLR contribution would stabilise after a
255 further period of simulation.


The cold and warm cases are far more realistic since they are based on observed Amundsen Sea conditions. Both scenarios
contribute much more SLR than the zero-melting case (figure 4a, e, i). In both cases, the Thwaites area dominates SLR
contributions from the sector, providing approximately two-thirds of the total, and responds differently to melting than the PIG
260 and Smith areas. In PIG and Smith areas, the rate of SLR with melting is approximately constant, but obtains higher rates of
SLR than in the zero-melting case, increasing to ~0.3 mm/yr and ~0.1 mm/yr, in the warm case, respectively, (figure 4f, j).
However, in the Thwaites area with melting a trend emerges in the SLR rate (figure 4b), increasing from ~0.2 mm/yr at the
start of the simulation to ~1.1 mm/yr at the end, in the warm case. In addition, we observe rapid jumps in the SLR rate for
Thwaites that are not present in the other areas. The increase in the SLR rate in the Thwaites results in an approximately
265 quadratic SLR contribution (figure 4a), rather than the more linear SLR contributions obtained for PIG and Smith. In all three
areas ‘noise’ is present in the SLR rates, though this is harder to see for the Thwaites area due to the larger y-axis scale. In
addition, we observe an increase in the total melting that occurs from the Thwaites Ice Shelf as the simulations progresses:
over the 180 years of the simulation, the total melt flux from Thwaites increases by an order of magnitude, from ~45 (~25,
respectively) Gt/yr at the start of the simulation to over ~440 (~210) Gt/yr at the end in the warm (cold) case (figure 4c), which
270 occurs despite constant oceanic boundary forcings during the simulation. This is explored further in Section 3.3. In the PIG

and Smith areas, the total melt flux is approximately constant (figure 4g, k). However, we do observe a strong correlation between the cumulative integral of melting and the SLR contribution individually for each region (Appendix C).

275 For all regions, we obtain consistently higher SLR rates and grounded ice area loss in the warm forcing case compared to the cold forcing case (figure 4b, f, j). However, as a fraction of the total SLR in these models, the difference between warm and cold scenarios is remarkably small, only ~27% of total SLR, by the end of the simulations. Given that these scenarios bracket the coldest and warmest ocean conditions on record in the Amundsen Sea, this suggests that the future SLR from this region is only weakly influenced by variations within the observed range of present day ocean conditions: while melting is important to the SLR, its typical climatic variations are less so. Despite this, in the Thwaites area we observe different timings and strengths in the jumps in the SLR rate between the cold and warm cases (figure 4b). We also observe an increasing difference between the cold and warm total melt rates for Thwaites Ice Shelf (figure 4c), while for the PIG and Smith areas this difference remains approximately constant (figure 4g, k). Additionally, there is a larger relative difference between the melting and no melting cases for the PIG area compared to Thwaites. In the PIG area, ice shelf melting prevents the ice shelf from thickening and re-grounding on the bathymetric ridge below it. In the no melt case, the ice shelf regrounds on this ridge; the buttressing provided by this re-grounding leads to the large differences between the no melt and melting cases for the PIG area (though this is dependent on bathymetry deepening). While there is a smaller relative difference for the Thwaites area, the presence of melting still has a significant impact, changing the SLR contribution from linearly increasing to quadratic.

280 Overall, we conclude that melting and SLR have a fundamentally different response in the Thwaites area than the other areas, leading to an increasing SLR rate and total melting rate. In addition, the Thwaites area dominates the SLR contribution for the Amundsen Sea sector over the 180 years simulated. Therefore, the remainder of this study focusses upon the processes underlying this behaviour in the Thwaites area.

3.2 Thwaites Glacier retreat and ice shelf pinning points



295

FIGURE 5: (a-b) Grounding line retreat rates over the 180 year simulations in the cold (a) and warm (b) indicated by colours, with colour bar shown in (b). In each the purple contours show the presence of isolated pinning points during the simulation. (c) Percentage change in grounding line retreat rates between the cold and warm cases (i.e., the percentage difference between those retreat rates shown in (a) and (b) respectively). (d-e) Bathymetry (d) and Weertman C coefficient (e) shown under initially grounded ice with final extents of the grounding line retreat in the warm (red) and cold (cyan) cases. (f) The difference in grounding line retreat rates between the cold and warm cases (colours) as well as final grounding line area (grey area). The area shown in each panel corresponds to that displayed as a black box in Figure 3.


Figure 5 shows quantities related to Thwaites grounding line retreat, which is one of the key features that differentiates this area from the PIG and Smith areas. Over the simulation, the warm case has a larger area of grounding-line retreat than the cold case, with a faster rate on average (figure 5 a, b, c, f). Grounding line retreat rates are calculated from the discrete migration of the grounding line across grid cells. Specifically, the retreat rate in each grid cell is calculated as $\Delta x_{GL} / \Delta t_{GL}$, where Δt_{GL} is the time between when the subject cell first becomes a grounding line cell (any of the 8 adjacent cells are floating) to when that cell ungrounds, and Δx_{GL} is the grid cell width or diagonal extent depending on which adjacent cell was first floating.

305

310 Figures 5d and 5e show the bed depth and ‘Weertman C’ basal sliding drag coefficient over the area of initially grounded ice..
A lower Weertman coefficient corresponds to a more slippery bed. In addition, the bathymetry in this area generally deepens
inland, which could promote grounding line retreat (Weertman, 1974; Schoof, 2007). In both warm and cold cases, highly
heterogeneous retreat rates are observed (figure 5a, b), with areas of fast retreat as high as ~10 km/y in proximity to areas of
much slower retreat. Towards the end of the warm case simulation, the grounding line experiences rapid retreat (Figure 5b)
315 across a deep and slippery bed section (Figure 5d-e) before slowing down as it encounters a ridge of shallower and less slippery
bed. These features explain the large variations in the Thwaites area SLR rate in the last ~50 years of the simulation (Figure
4b).

Comparing the cold and warm scenarios we see the effect of increased melting on grounding-line retreat rates. More
320 specifically, we observe areas of elevated grounding line retreat rate in the northern part of the ungrounding area, below which
an area with minimal sensitivity to melting scenario, and again by increased rates in the south (Figure 5f). There are very large
percentage increases in grounding-line retreat rates for these two clear bands on the retreated area (Figure 5c).

In addition, we observe the formation of many pinning points as Thwaites Glacier retreats, shown by purple contours in Figure
325 5 for the different cases. An ice grid cell is flagged as a ‘pinning point’ if it is grounded but separated from the main grounded
ice sheet by floating ice. The purple contour is then drawn around all cells that are flagged as pinning points in any one of the
outputs. The pinning points are generally located on areas of shallower bathymetry that are downstream of areas of deeper
bathymetry (Figure 5d). The pinning points typically feature slower retreat rates, and inshore of these pinning points we
generally observe areas of faster retreat (Figure 5a, b). These pinning points are crucial for the future evolution of Thwaites
330 Glacier, as they provide drag and buttressing as the ice retreats. The importance of these pinning points is determined by their
size, position, and duration, so we examine these features and their relationship to SLR rates from the Thwaites area.



335

FIGURE 6: (a) SLR contribution rates for Thwaites area for warm (red), cold (blue) and no melt (black) cases. The background colour indicates the time in the simulation according to bands of 25 years, with a longer starting (ending) band of 45 (35) years. (b) Colours indicate the band of the final year that pinning point cells exists for cold melt case, according to the colours shown in (a). Note that only pinning points that exist for over 1 year are shown. Red labelled boxes refer to groups of pinning points discussed in the text. (c) Same as (b) but for warm case. (d-e) Pinning point duration, taken as the time from isolation to ungrounding, for cold (d) and warm (e) cases. (f) Difference in pinning point duration between cold and warm cases (i.e. the difference the data shown in d and e), where pinning point locations match. In (b)-(e), grey regions indicate areas which do not become ungrounded during the simulation. The area shown in Figures b-f is shown with a black box in Figure 3.


345

Figure 6b-c shows the ungrounding time of pinning points in the cold and warm cases, respectively. By examining these times, we can compare how the ungrounding of pinning points relate to jumps in SLR rate that we observe from the Thwaites area. The labelled red boxes in Figure 6b show the key groups of pinning points in the simulations.

350 Within the first 60 years, the SLR rates of the warm and cold forced cases diverge (Figure 6a). In the warm case, pinning point group 'a' completely ungrounds between years 25 and 60 (Figure 6c), with the final ungrounding of the last pinning point coinciding with the ungrounding of group 'b'. The removal of the combined associate buttressing, leads to a large jump in SLR rate in the warm simulation at ~60 years. However, in the cold simulation, some of these pinning points in group 'a' remain grounded until beyond 95 years (Figure 6b), leading to a period of relatively steady SLR rate in this simulation (Figure
355 6a). In the zero-melting case, part of pinning point group 'a' and all of 'b' remain grounded throughout the simulation (not shown), and SLR rates remain approximately constant as a result (Figure 6a). In the warm simulation, group 'c' becomes ungrounded by ~130 years, causing a rapid increase in SLR rate, and the loss of group 'd' at ~145 years leads to another large jump. The cold simulation loses group 'c' at ~170 years, leading to a small jump in SLR rate, while group 'd' remains grounded at the end of the simulation.

360

Figure 6d-e shows the duration of the pinning points- the time between separation from the main body of grounded ice to ungrounding. There is a large variation in pinning point durations, with some lasting less than a year while others persist for decades. In general, the duration of pinning points is lower in the warm case (Figure 6d-f), with an average pinning point duration over warm/cold matching pinning locations of ~6 years, compared to ~10 years in the cold case, leading to the
365 different ungrounding times and SLR rates as described above. Reducing the duration of pinning points increases the intensity of periods of rapid ice acceleration and grounding line retreat, leading to the differences in the spatial map of grounding line retreat rates between the cold and warm cases shown in Figure 5 c, f.



370

FIGURE 7: Evolution of conditions as grounding line retreats over pinning points. Snapshots every 9 years starting from the date of group ‘c’ pinning point formation in the warm and cold cases (118 and 144 years respectively). (a-c) Cross section through the Thwaites Ice Shelf, taken along the red line in (g) extended to ice front, for warm case. (d-f) Same as (a-c) but for cold case. (g-i) Melt rates for warm case, with

arrows showing ice shelf boundary layer ocean velocities (j-l) Same as (g-i) but for cold case. (m) Ice speed snapshots at start of pinning
375 point formation and (n, o) show differences for warm case. (p-r) Same as (m-o) but for cold case. Area shown in (i-x) is shown in figure 3
as a small green box.

To illustrate the key role of pinning points, we now focus on the ungrounding of group 'c', which causes the rapid jump in
SLR rate at year 130 in the warm simulation, and the smaller jump at year 170 in the cold simulation. Figure 7 shows the
380 evolution of the two simulations, starting from their individual dates of formation of the group 'c' pinning points. Figures 7a-
l show the ice geometry and ocean conditions throughout the subsequent evolution. To begin with, the grounding line is located
in shallower bathymetry at the top of a retrograde slope (figure 7a, d). Ungrounding then occurs laterally around this point,
including upstream, encircling the pinning point and leaving it isolated from the rest of the grounded ice (figure 7h, k). Note
that the pinning point is grounded on the side of the shallower bathymetry, rather than on top of the bathymetric feature. With
385 the grounding line now in a deeper bed an acceleration is expected, although the pinning point continues to provide basal
sliding drag that resists the flow. The advection of thicker ice from the deeper bed upstream enables the ice to remain grounded
on the shallower bed beneath the pinning point. Therefore, only a small acceleration is observed at 9 years, while the ice
remains grounded on the pinning point (Figure 7m-r).

390 However, melting now occurs in the newly-opened cavity upstream of the pinning point. This thins the ice, enlarging this
cavity, enabling greater oceanic connection, and leading to higher ocean velocities and melt rates (figure 7h, k). This melting
and thinning feedback eventually leads to sufficient thinning upstream of the pinning point for it to unground completely
(Figure 7i). The resulting loss in buttressing leads to a large increase in ice speed (figure 7o), further rapid grounding line
retreat and a jump in SLR rate. Therefore, the distribution and strength of localised melt rate patterns strongly determines the
395 duration of these pinning points and thus the overall ice retreat. This is shown clearly in Figure 7g-l, with the warm case leading
to higher melt rates and a much faster ungrounding of the pinning point. The melt rates around these pinning points are highly
heterogeneous (Figure 7g-l), with elevated melt rates typically occurring on their eastern side, where rapid ocean currents drive
high melt rates. These rapid currents occur where buoyancy-driven meltwater flow is trapped against the grounding line by
Coriolis force (Holland and Feltham, 2006).

400




FIGURE 8: Cross sections of potential temperature beneath Thwaites Ice Shelf for cold (left) and warm (right) cases, taken every 36 years, along the same section as shown in Figure 7 and Figure 2i, j, which is, represented in Figure 2a as the red line.

410 An increasing trend in the total ice shelf melting occurs over the evolving Thwaites Ice Shelf, during both the warm and cold forcing simulations (Figure 4c). The wider oceanic forcings are fixed, so these trends must be driven by geometric changes in the ice shelf cavity. In Figure 8 we show how the Thwaites cavity geometry and ocean conditions evolve in the two cases. The warm case has a higher thermocline, so the warm CDW is more easily able to flow over ridges and flood the new cavity areas. In the cold case, the access of the warmest CDW is blocked by seabed highs, with only more heavily-modified CDW reaching

415 the ice base. However, sufficiently warm water is still able to drive melting close to the grounding line. In both cases the ice base near the grounding line remains steeply sloped throughout the retreat.

Figure 9a shows the evolution of total melt flux from the Thwaites Ice Shelf in the warm case, for both the entire ice shelf and for ice below 600 m depth only, which is the thermocline depth in this case. Most of the trend in ice shelf melting occurs in
420 the deeper ice, with melting below 600 m peaking at an increase of ~ 30 times its initial value, which suggests that as the grounding line retreats, a strong increase in ice shelf base area below the thermocline controls total melting (Figure 9b). This confirms that the trend in melting does not result from the increasing ice shelf area associated with an artificially fixed ice front. The increase in deep ice area occurs because the grounding line retreats into deeper bathymetry, but also because the slope of the ice shelf base, below 600 m, gets shallower during the simulation, as shown for example in Figure 8. Without an
425 increasing trend in melt flux, the thicker ice advected across the grounding line as the ice sheet retreats into deeper bathymetry would result in re-grounding on pinning points further downstream. The close correspondence in the increase between total melting (10 to 197 Gt/yr) and ice base area (180 to 3560 km²) implies that the average melt rate (m/yr) beneath deep ice, below 600 m is approximately constant at ~ 45 m/yr (Figure 9c). However, we do find some temporal variability in this average melt rate, with it varying up to 50 % and importantly the local melt rates are highly spatially variable, with a spatial standard
430 deviation close to the mean value. This spatial variability is important for the effect of melting on small scale pinning points, as described above.




FIGURE 9: (a) Total melt flux from the full Thwaites Ice Shelf area (black) and from grid cells located below 600 m depth only (blue). (b)
435 Total melt flux from grid cells below 600 m and total ice area below 600 m. (c) Average melt rate on grid cells located below 600 m depth and spatial standard deviation of these melt rate values.

4 Discussion

This study considers the future evolution of ice streams in the Amundsen Sea sector using a coupled ice/ocean model. Under
440 both warm and cold forcing we find large grounding line retreat and ice acceleration from Thwaites Glacier, with the Thwaites
area dominating future SLR contributions from the region. The SLR from the Thwaites area is consistent with a previous
coupled model study (Seroussi et al., 2017), which considered a shorter time span of 50 years. In this study however, we show
a larger sensitivity to a realistic range of ocean forcings, in the shorter and longer term, and an increasing SLR rate, which
continues over 160 years of the time period, from the Thwaites area. This was not observed in previous coupled modelling of
445 Thwaites Glacier (Seroussi et al., 2017), although it was observed in some ice-only simulations, depending on uncertainty in
ice dynamics (Nias et al., 2019) and uncertainty in melt rates (Arthern and Williams, 2017). A previous ice-only study has set
1 mm/yr SLR rate to be the threshold that implies rapid retreat and collapse of Thwaites glacier (Joughin et al., 2014), and this
is exceeded sooner in our simulations, within 125 years. However, our simulations are not long enough to see how the full
implications play out over multiple centuries.

450

The SLR contributions in this study of 70 - 89 mm after 100 years for the Amundsen Sea sector are within the uncertainties of
previous ensemble studies (Nias et al., 2019; Edwards et al., 2021). However, the SLR and stability suggestions in this study
do differ from recent studies, which found that the present-day geometry is not inherently unstable when starting from a stable
starting position (Hill et al., 2023), and that the Amundsen Sea sector has not tipped yet (Reese et al., 2023). We suggest that
455 these differences arise primarily through the different ice sheet model initialisation strategies adopted, which variously use a
spin up period (Reese et al., 2023), data assimilation of ice velocities into a steady state (Hill et al., 2023), or assimilation of
ice velocities and observations of unsteady thinning (present study). Also, the date of initialisation and differences in
resolutions, datasets used and model physics may also play an important role. Larger coupled-model ensembles are needed to
assess these aspects. Without this there is high uncertainty, for example, in SLR contributions and the timings of pinning point
460 ungroundings. This study is designed to provide a small number of physically-advanced coupled simulations focusing on
ice/ocean processes, rather than providing a larger but uncoupled set of predictions of future SLR contributions from the region.
In our simulations, the increase in SLR rate from the Thwaites area is caused by ocean-driven melting, and the magnitude of
this increase is sensitive to different rates of melting in warm and cold scenarios. The SLR rate is governed by a balance
between retreat of the grounding line into deeper bed regions, and the formation and duration of pinning points during this
465 retreat. Crucially, this study shows the importance of ocean-driven melting in ungrounding these pinning points, reducing ice-
shelf buttressing and enhancing grounding-line retreat. It should be noted that the modelled rates of ungrounding upstream of
pinning points are high enough to explain recent observations (Graham et al., 2022). Other studies have hypothesised about
the mechanics and importance of pinning points (Thomas, 1979), and have shown the effect of pinning points in an idealised
ice/ocean coupled model (De Rydt and Gudmundsson, 2016). In this study we clearly show the importance of these
470 mechanisms in a synchronous coupled model of the future Thwaites Ice Shelf, with strong spatial variability in ice shelf melting

determining the duration of these pinning points and therefore their influence on SLR. Our use of a synchronously coupled model means variations in the ice thickness are instantly felt in the ocean model's melting calculation. This could impact the speed of the evolution of these pinning points, though further work is required to determine the impact of this coupling on the ice dynamics. The importance of future pinning points suggests the need for more accurate knowledge of bathymetry and bed
475 properties in the grounded portion of Thwaites Glacier, as well as highlighting the importance of accurately modelling the effects of pinning points in both ice sheet and ocean models. We have shown that high resolution coupled ice-ocean models are required to investigate the effect of pinning points on ice dynamics, as these small features need to be resolved in the ice model, and the strong spatial variability in ice shelf melting around them needs to be recreated.

480 One important question raised by our study is why Thwaites Glacier appears to behave differently from the other glaciers in the region. We speculate that this is caused by the wide trunk of this glacier. As the grounding line of the Thwaites Glacier retreats it forms an extremely short and wide ice shelf, unlike the other glaciers in the region, whose ice shelves are confined within embayments. As a result, the buttressing provided by lateral ice shelf margins is very weak for Thwaites Glacier; this also explains the high sensitivity to pinning points which therefore provide the majority of the buttressing of Thwaites.
485 Therefore, while a recent study found that Thwaites Ice Shelf provides limited buttressing in the present day (Gudmundsson et al., 2023), we find that future configurations of Thwaites Ice Shelf will provide important buttressing, via future pinning points, as its grounding line retreats. The dependence of this future buttressing on ocean forcings will determine the future SLR from this sector.

490
The simulations show how the geometric changes in the Thwaites Ice Shelf and its cavity can lead to an increasing ice area at depth, meaning that more of the ice base is exposed to deep warm waters and leading to an increasing trend in total melt flux. Although this general depth dependence on melting is often captured by parametrizations of melting (Asay-Davis et al., 2017), we have shown that there is strong spatial variability in melting, and this spatial variability has important consequences for
495 pinning point duration. Therefore, as parametrizations of basal melting generally lack a strong physical basis and perform poorly in spatial detail (Burgard et al., 2022), they may be expected to struggle to capture this spatial variability, affecting the evolution of these pinning points and the resultant SLR contribution rate from Thwaites. However, future work is required, running sets of ice only simulations, to perform an extensive comparison of the wide range of basal melt rate parametrizations. Observations of ice shelf melt rates around such features with sufficient spatial coverage are currently lacking but are essential
500 for improving future modelling efforts. The increasing trend in total melt rates is needed to maintain the ice retreat, as it counteracts increased ice thickness advection across the grounding line, which would otherwise ground the ice on bathymetry further downstream.

One of the limitations of this study is that steady, idealized, ocean forcings are applied to the northern and western boundaries. However, ocean conditions in the region are known to have strong decadal variability (Jenkins et al., 2018; Dutrieux et al., 2014). Some of the pinning points only exist for ~10 years, so decadal variability in oceanic forcings may have an important role in how quickly these pinning points are ungrounded. In addition, a superimposed anthropogenic warming trend in ocean forcing may be expected in the Amundsen Sea (Holland et al., 2022; Naughten et al., 2022), which could decrease the duration of future pinning points and speed up the retreat of Thwaites Glacier. However, while the forcings used in this study are idealised, they are based on the best available information of present-day extremes in observations (Dutrieux et al., 2014). These forcings also agree approximately with the time average of recent projections of future warming in the region (Naughten et al., 2023), which found linearly rising trends of ocean temperature in all tested climatic scenarios. The steady warm and cold forcings used in this study have an average temperature between 200-700 m depth of ~0.35 °C and ~-0.45 °C respectively. The warm case approximates the time-averaged temperatures at this depth over the similar trends projected in 100 year simulations of the Paris 1.5 °C, Paris 2 °C and RCP 4.5 climatic conditions (Naughten et al., 2023), while the cold case approximates a cold historical state (Naughten et al., 2023). Another limitation of this study is the use of an ocean model that does not represent the evolution of oceanic conditions and sea ice on the wider Amundsen Sea continental shelf. As such, our model might lack important feedbacks such as increased ocean currents driven by ice-shelf meltwater bringing more CDW onto to shelf and driving higher melt rates (Kimura et al., 2017; Jourdain et al., 2017; Donat-Magnin et al., 2017). With an order of magnitude increase in ice shelf melting in our projections, such feedbacks would be substantial. In addition, the ocean simulation lacks the physical presence and effect of some freshwater sources, like sea ice and icebergs, which impacts the stratification of the water column, oceanic currents and the delivery of warm CDW to the base of ice shelves in the region (Bett et al., 2020). The lack of sea ice in particular could increase the heat content of CDW reaching the ice shelf bases, due to the lack of sea ice driven convective processes cooling the CDW layer (St-Laurent et al., 2015; Webber et al., 2017). However, it remains unknown how this cooling influence might vary over the coming centuries. These simulations additionally lack subglacial freshwater discharge, which at the grounding line has been found to increase ice shelf melting locally in previous ocean modelling studies (Nakayama et al., 2021), but overall, its affect is small in this region (Holland et al., 2023).

A key limitation of the model used in this study is the lack of ice shelf calving-front retreat, which could impact ice dynamics, ocean conditions and total melting (e.g. Bradley et al., 2022; Joughin et al., 2021). As well as using a fixed temperature field, the model lacks an evolving damage field (Lhermitte et al., 2020). Therefore, the model may be sensitive to the time of initialisation, as this will determine level of the damage that is applied for the entire forward simulation. However, we believe that these limitations cause our simulations to be a conservative estimate of Thwaites Glacier retreat for the cold and warm cases, as calving and evolving damage are only expected to enhance the retreat. Barring the existence of major calving- or damage-driven ice retreat over the next 180 years, our results suggest that the ocean-driven ungrounding of pinning points will dictate the future SLR from the Amundsen Sea sector. However, strong sensitively of ice-sheet projections to basal friction

laws provides uncertainty to modelled ice retreat rates and mass loss, with the Weertman sliding law used in this study found to systematically predict the lowest ice mass losses (Brondex et al., 2019; Cornford et al., 2020). It should also be noted that uncertainties in the accumulation field could potentially affect the modelled ice dynamics, including modelled SLR and grounding line retreat rates. In addition, the accumulation field used in this study is held steady and therefore any effects of future trends in this field are omitted, which could mitigate ocean driven ice loss to some extent (Edwards et al., 2021). Thus, this study focusses solely on dynamical ice loss driven by ocean melting.

All these ice dynamical limitations affect the no melt case results in this study. However, some of these limitations lead to specific additional caveats to the no melt case, where they may have the greatest impact. For example, the fixed ice front mask, includes the current gaps between the east and west of Thwaites Ice Shelf, and these gaps cannot recover during the simulation. This may have the greatest impact in the no melt case, where the ice shelf thickens and recovery of this damage should be possible, and this could lead to an overestimation of SLR contributions from this hypothetical case.

550

5 Conclusions

This study presents, for the first time, 180-years of ice evolution in the Amundsen Sea sector of the West Antarctic Ice Sheet using a new synchronously coupled ice/ocean model, which includes full mass conservation in the coupled ice/ocean system, and an instantaneous response of melt rates to the evolving ice geometry. The coupled simulations were forced with idealised warm and cold ocean conditions in the wider Amundsen Sea, and compared to each other, as well as to a zero ice-shelf melting case.

Even in the zero-melting case, the model predicts that the Thwaites and Smith areas in the region lose ice mass during the simulations. This implies that the ice sheet model is initialised into an intrinsically unsteady state, so that in these simulations a tipping point may have occurred in the past and we are now committed to further sea-level rise from this sector. However, when melting is activated in the coupled model, the rates of ice loss are much higher. This implies that ocean melting plays an important role in the future SLR contribution from this sector, though the difference between warm and cold scenarios is relatively modest, at only ~27% of the total SLR.

For Pine Island and Smith Glaciers, the rate of SLR remains relatively constant, at approximately 0.3 mm/yr and 0.1 mm/yr respectively, throughout the projections, leading to a linearly-increasing sea-level contribution. The Thwaites Glacier area provides a much larger sea-level contribution, and features an increasing SLR rate, which causes its sea-level contribution to increase approximately quadratically with time. The rate of SLR from Thwaites Glacier is closely controlled by the formation and duration of isolated pinning points during the retreat of its grounding line. Ocean-driven melting is crucial in driving the

570 ungrounding of these pinning points, by thinning the ice upstream, and this is the key mechanism by which future ocean conditions affect the SLR from this sector.


The coupled simulations show a large geometry-induced increase in total ocean-driven melting as Thwaites Glacier retreats and its ice shelf enlarges. This increased melting counteracts ice shelf thickening associated with thicker ice being advected
575 across the deeper grounding line, which would otherwise cause the ice to ground downstream and arrest the retreat. Our simulations indicate large spatial and temporal variability in the melt rates at depth below 600 m. This variability will not appear within simple melting parameterisations.

Our results also suggest that accurate modelling of ocean-driven melting and ice response around pinning points, and accurate
580 characterisation of the bed geometry and properties that lead to the formation of pinning points must be future research priorities. In addition, the further development and application of coupled ice/ocean models must be a priority, as it is difficult to envision how many of these results could have been achieved with parameterised ocean-driven melting, though further work is required to symmetrically compare the ice/ocean coupled model to the wide selection of melting parameterisations.

Appendices

585 **Appendix A: Ice shelf melt rate tuning**

In the initial setup of the ice model, we can calculate the implicit melt rate (IMR), which is the melt rate required to recreate observed present day surface elevation changes in the ice model, given initial ice model velocities and geometry (Arthern and Williams, 2017). In this model, we attempt to minimise initial coupling shock by reducing the mismatch between the ice model IMR and the initial ice shelf melting field calculated from the MITgcm ocean model. To do so, we tune the dimensionless ice
590 shelf melting drag coefficient in the three-equation formulation of melting parametrisation used in the MITgcm (Jenkins et al., 2010). We found that a choice of drag coefficient of 0.008 minimises the combined PIG and Thwaites ice shelves mismatch in total melt flux (Figure A1), while average ocean forcings are applied, with the thermocline placed at 700 m depth.



595 **FIGURE A1:** Area shown for PIG and Thwaites ice shelves. a) Initial implicit melt rate from the WAVI ice sheet model. b) Initial ice shelf melt rate from the MITgcm model for different values of the ice shelf melting drag coefficient as follows (and labelled): 0.008 (b), 0.01 (c), and 0.012 (d).. (e-g) Difference between initial implicit melt rate and initial ice shelf melting for the MITgcm model melt rates shown in panels (e)-(g) respectively.

Appendix B: Initial bathymetry deepening

600 To ensure the bathymetry field from BedmachineV3 (Morlighem et al., 2020; Morlighem, 2022) has a specified minimum water column thickness under the initially floating ice area on the staggered Arakawa C-grid's velocity grid points, located on grid faces, we performed a 'deepening' procedure, in which the bathymetry in areas with a water column thinner than a specific value is artificially deepened. This procedure was applied only to grid cells in which no ice basal sliding drag is applied in the initial state and occurs only once in a simulation before the WAVI ice sheet model is initialised and then relaxed. This

605 is because the bathymetry in ice shelf cavities is poorly known, while beneath the grounded ice sheet the bed is better known from radar soundings. We found this step is necessary because, without it, the initial PIG Ice Shelf cavity has an excessively thin water column near the grounding line. This leads to minimal ice shelf melting in this region (Figure B1c), which leads to a large mismatch with the calculated initial IMR (Figure A1). This, combined with the shallow bathymetry, leads PIG to reground immediately at the start of the simulation, resulting in unrealistically low SLR rates from this region (Figure B1d).

610

Therefore, an initial deepening step was applied in order to prevent re-grounding and minimise any changes in SLR rate at the start of the simulation (Figure B1d). However, enforcing a uniform minimum water-column thickness everywhere would create a sudden step-change in the bathymetry at the initial grounding line location. Therefore, a taper is applied to the water column thickness used in the deepening procedure, increasing the minimum thickness from a low value at the grounding line to its standard value over 6 km distance. After applying this initial deepening procedure, PIG does not reground at the start of the simulation. We note that the SLR rate from the Thwaites region (Figure B1a) is only minimally affected by the deepening procedure. While the deepening of the PIG water column is somewhat subjective, the deeper water column better reflects the sparsely available observations of the cavity near the grounding line (Dutrieux et al., 2014). Overall, the necessity of deepening emphasises the importance of obtaining more detailed observations of the geometry of ice shelf cavities, beneath PIG in

620

particular.





FIGURE B1: a) SLR rate of change for the Thwaites area for: zero deepening, 140 m deepening, 280 m deepening, 280 m deepening taper to 140 m over 6 km.. b) Cross-section through PIG Ice Shelf (taken along the green line shown in Figure 2a) showing initial potential

625 temperature in the warm case in the zero-deepening setup. c) Initial melt rate over PIG Ice Shelf in the warm forcing case for the zero-deepening setup. d) Same as a) but for the PIG area. e) Same as (b), but for the 280 m taper 140 m deepening setup. f) Same as (c), but for the 280m taper 140 m deepening setup.

Appendix C: SLR compared against integrated melt

630 A strong correlation is found when comparing the sea level contribution against the integrated melt for both forcing cases in the areas of Thwaites and PIG (Figure C1). However, no variation to the oceanic boundary forcing is applied during the simulations and hence only geometric induced changes in the ice shelf melt rates can occur. These strong correlations are due to the melt rate and rate of change of SLR being approximately constant from PIG's area and having an approximately linear trend from Thwaites's area. The regression between the SLR contribution and the integrated melt for both the Thwaites and
635 PIG areas are different for the two oceanographic forcings applied, which suggests that the ratio of mass loss between ice shelf melting and calving is different between the cases, with the warm case having a higher relative ice shelf melting.



640 **FIGURE C1:** (a) Yearly points of sea level contribution plotted against integrated melt from the Thwaites area for the warm (red) and cold (blue) forced cases. (b) as in (a), but for the PIG area. In both, correlation coefficients are shown for the warm and cold cases.

Code and data availability

The version of MITgcm used in this study is available at https://github.com/David-Bett4/MITgcm/tree/Coupling_wHoriz_lat_Pchild_new_divout. The version of WAVI used in this study is available at https://github.com/RJArthern/WAVI.jl/tree/MITgcm_coupling. Coupling scripts and input text files are available at https://github.com/David-Bett4/MITgcm_WAVI_coupling. The model output underlying the figures and calculations in this paper is available at https://github.com/David-Bett4/Amundsen_coupling_paper.

Author contribution

DTB implemented coupling, set up ocean initial state, ran coupled simulations, performed analysis and led the manuscript. ATB and CRW set up initial WAVI ice model states and performed WAVI ice model relaxation simulations. RJA, PRH, CRW and DNG designed and supervised the project. All of the co-authors contributed to coupling design, experiment design and edited the manuscript.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

DTB, ATB, CRW, PRH, RJA were supported by the NERC Grant NE/S010475/1. CRW was partly funded by the MELT project, a component of the International Thwaites Glacier Collaboration (ITGC), with support from National Science Foundation (NSF: Grant no. 1739003) and Natural Environment Research Council (NERC: Grant no. NE/S006656/1). ITGC Contribution No. ITGC-108. This work used the ARCHER2 UK National Supercomputing Service (<https://www.archer2.ac.uk>).

References

- Alevropoulos-Borrill, A. V., Nias, I. J., Payne, A. J., Golledge, N. R., and Bingham, R. J.: Ocean-forced evolution of the Amundsen Sea catchment, West Antarctica, by 2100, *The Cryosphere*, 14, 1245-1258, 10.5194/tc-14-1245-2020, 2020.
- Arthern, R. J. and Williams, C. R.: The sensitivity of West Antarctica to the submarine melting feedback, *Geophysical Research Letters*, 44, 2352-2359, <https://doi.org/10.1002/2017GL072514>, 2017.
- Arthern, R. J., Hindmarsh, R. C. A., and Williams, C. R.: Flow speed within the Antarctic ice sheet and its controls inferred from satellite observations, *Journal of Geophysical Research: Earth Surface*, 120, 1171-1188, <https://doi.org/10.1002/2014JF003239>, 2015.
- Arthern, R. J., Winebrenner, D. P., and Vaughan, D. G.: Antarctic snow accumulation mapped using polarization of 4.3-cm wavelength microwave emission, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2004JD005667>, 2006.
- Asay-Davis, X. S., Jourdain, N. C., and Nakayama, Y.: Developments in Simulating and Parameterizing Interactions Between the Southern Ocean and the Antarctic Ice Sheet, *Current Climate Change Reports*, 3, 316-329, 10.1007/s40641-017-0071-0, 2017.

- Bett, D. T., Holland, P. R., Naveira Garabato, A. C., Jenkins, A., Dutrieux, P., Kimura, S., and Fleming, A.: The Impact of the Amundsen Sea Freshwater Balance on Ocean Melting of the West Antarctic Ice Sheet, *Journal of Geophysical Research: Oceans*, 125, e2020JC016305, <https://doi.org/10.1029/2020JC016305>, 2020.
- 675 Bradley, A. T., Bett, D. T., Dutrieux, P., De Rydt, J., and Holland, P. R.: The Influence of Pine Island Ice Shelf Calving on Basal Melting, *Journal of Geophysical Research: Oceans*, 127, e2022JC018621, <https://doi.org/10.1029/2022JC018621>, 2022.
- Brondex, J., Gillet-Chaulet, F., and Gagliardini, O.: Sensitivity of centennial mass loss projections of the Amundsen basin to the friction law, *The Cryosphere*, 13, 177-195, 10.5194/tc-13-177-2019, 2019.
- Burgard, C., Jourdain, N. C., Reese, R., Jenkins, A., and Mathiot, P.: An assessment of basal melt parameterisations for Antarctic ice shelves, *The Cryosphere*, 16, 4931-4975, 10.5194/tc-16-4931-2022, 2022.
- 680 Cornford, S. L., Seroussi, H., Asay-Davis, X. S., Gudmundsson, G. H., Arthern, R., Borstad, C., Christmann, J., Dias dos Santos, T., Feldmann, J., Goldberg, D., Hoffman, M. J., Humbert, A., Kleiner, T., Leguy, G., Lipscomb, W. H., Merino, N., Durand, G., Morlighem, M., Pollard, D., Rückamp, M., Williams, C. R., and Yu, H.: Results of the third Marine Ice Sheet Model Intercomparison Project (MISMIP+), *The Cryosphere*, 14, 2283-2301, 10.5194/tc-14-2283-2020, 2020.
- De Rydt, J., Holland, P. R., Dutrieux, P., and Jenkins, A.: Geometric and oceanographic controls on melting beneath Pine Island Glacier, *Journal of Geophysical Research: Oceans*, 119, 2420-2438, <https://doi.org/10.1002/2013JC009513>, 2014.
- 685 De Rydt, J. and Gudmundsson, G. H.: Coupled ice shelf-ocean modeling and complex grounding line retreat from a seabed ridge, *Journal of Geophysical Research: Earth Surface*, 121, 865-880, <https://doi.org/10.1002/2015JF003791>, 2016.
- Donat-Magnin, M., Jourdain, N. C., Spence, P., Le Sommer, J., Gallée, H., and Durand, G.: Ice-Shelf Melt Response to Changing Winds and Glacier Dynamics in the Amundsen Sea Sector, Antarctica, *Journal of Geophysical Research: Oceans*, 122, 10206-10224, <https://doi.org/10.1002/2017JC013059>, 2017.
- 690 Dutrieux, P., De Rydt, J., Jenkins, A., Holland, P. R., Ha, H. K., Lee, S. H., Steig, E. J., Ding, Q., Abrahamsen, E. P., and Schröder, M.: Strong Sensitivity of Pine Island Ice-Shelf Melting to Climatic Variability, *Science*, 343, 174-178, 10.1126/science.1244341, 2014.
- Edwards, T. L., Nowicki, S., Marzeion, B., Hock, R., Goelzer, H., Seroussi, H., Jourdain, N. C., Slater, D. A., Turner, F. E., Smith, C. J., McKenna, C. M., Simon, E., Abe-Ouchi, A., Gregory, J. M., Larour, E., Lipscomb, W. H., Payne, A. J., Shepherd, A., Agosta, C., Alexander, P., Albrecht, T., Anderson, B., Asay-Davis, X., Aschwanden, A., Barthel, A., Bliss, A., Calov, R., Chambers, C., Champollion, N., Choi, Y., Cullather, R., Cuzzone, J., Dumas, C., Felikson, D., Fettweis, X., Fujita, K., Galton-Fenzi, B. K., Gladstone, R., Golledge, N. R., Greve, R., Hattermann, T., Hoffman, M. J., Humbert, A., Huss, M., Huybrechts, P., Immerzeel, W., Kleiner, T., Kraaijenbrink, P., Le clec'h, S., Lee, V., Leguy, G. R., Little, C. M., Lowry, D. P., Malles, J.-H., Martin, D. F., Maussion, F., Morlighem, M., O'Neill, J. F., Nias, I., Pattyn, F., Pelle, T., Price, S. F., Quiquet, A., Radić, V., Reese, R., Rounce, D. R., Rückamp, M., Sakai, A., Shafer, C., Schlegel, N.-J., Shannon, S., Smith, R. S., Straneo, F., Sun, S., Tarasov, L., Trusel, L. D., Van Breedam, J., van de Wal, R., van den Broeke, M., Winkelmann, R., Zekollari, H., Zhao, C., Zhang, T., and Zwinger, T.: Projected land ice contributions to twenty-first-century sea level rise, *Nature*, 593, 74-82, 10.1038/s41586-021-03302-y, 2021.
- 700 Favier, L., Durand, G., Cornford, S. L., Gudmundsson, G. H., Gagliardini, O., Gillet-Chaulet, F., Zwinger, T., Payne, A. J., and Le Brocq, A. M.: Retreat of Pine Island Glacier controlled by marine ice-sheet instability, *Nature Climate Change*, 4, 117-121, 10.1038/nclimate2094, 2014.
- 705 Feldmann, J. and Levermann, A.: Collapse of the West Antarctic Ice Sheet after local destabilization of the Amundsen Basin, *Proceedings of the National Academy of Sciences*, 112, 14191-14196, doi:10.1073/pnas.1512482112, 2015.
- Goldberg, D. N.: A variationally derived, depth-integrated approximation to a higher-order glaciological flow model, *Journal of Glaciology*, 57, 157-170, 10.3189/002214311795306763, 2011.
- 710 Goldberg, D. N. and Heimbach, P.: Parameter and state estimation with a time-dependent adjoint marine ice sheet model, *The Cryosphere*, 7, 1659-1678, 10.5194/tc-7-1659-2013, 2013.
- Goldberg, D. N. and Holland, P. R.: The Relative Impacts of Initialization and Climate Forcing in Coupled Ice Sheet-Ocean Modeling: Application to Pope, Smith, and Kohler Glaciers, *Journal of Geophysical Research: Earth Surface*, 127, e2021JF006570, <https://doi.org/10.1029/2021JF006570>, 2022.
- 715 Goldberg, D. N., Snow, K., Holland, P., Jordan, J. R., Campin, J. M., Heimbach, P., Arthern, R., and Jenkins, A.: Representing grounding line migration in synchronous coupling between a marine ice sheet model and a z-coordinate ocean model, *Ocean Modelling*, 125, 45-60, <https://doi.org/10.1016/j.ocemod.2018.03.005>, 2018.
- Graham, A. G. C., Wählin, A., Hogan, K. A., Nitsche, F. O., Heywood, K. J., Totten, R. L., Smith, J. A., Hillenbrand, C.-D., Simkins, L. M., Anderson, J. B., Wellner, J. S., and Larter, R. D.: Rapid retreat of Thwaites Glacier in the pre-satellite era, *Nature Geoscience*, 15, 706-713, 10.1038/s41561-022-01019-9, 2022.
- 720 Gudmundsson, G. H., Barnes, J. M., Goldberg, D. N., and Morlighem, M.: Limited Impact of Thwaites Ice Shelf on Future Ice Loss From Antarctica, *Geophysical Research Letters*, 50, e2023GL102880, <https://doi.org/10.1029/2023GL102880>, 2023.
- Hill, E. A., Urruty, B., Reese, R., Garbe, J., Gagliardini, O., Durand, G., Gillet-Chaulet, F., Gudmundsson, G. H., Winkelmann, R., Chekki, M., Chandler, D., and Langebroek, P. M.: The stability of present-day Antarctic grounding lines – Part 1: No indication of marine ice sheet instability in the current geometry, *The Cryosphere*, 17, 3739-3759, 10.5194/tc-17-3739-2023, 2023.
- 725

- Holland, D. M. and Jenkins, A.: Modeling Thermodynamic Ice–Ocean Interactions at the Base of an Ice Shelf, *Journal of Physical Oceanography*, 29, 1787-1800, [https://doi.org/10.1175/1520-0485\(1999\)029<1787:MTIOIA>2.0.CO;2](https://doi.org/10.1175/1520-0485(1999)029<1787:MTIOIA>2.0.CO;2), 1999.
- Holland, P. R. and Feltham, D. L.: The Effects of Rotation and Ice Shelf Topography on Frazil-Laden Ice Shelf Water Plumes, *Journal of Physical Oceanography*, 36, 2312-2327, <https://doi.org/10.1175/JPO2970.1>, 2006.
- 730 Holland, P. R., Bevan, S. L., and Luckman, A. J.: Strong Ocean Melting Feedback During the Recent Retreat of Thwaites Glacier, *Geophysical Research Letters*, 50, e2023GL103088, <https://doi.org/10.1029/2023GL103088>, 2023.
- Holland, P. R., Bracegirdle, T. J., Dutrieux, P., Jenkins, A., and Steig, E. J.: West Antarctic ice loss influenced by internal climate variability and anthropogenic forcing, *Nature Geoscience*, 12, 718-724, 10.1038/s41561-019-0420-9, 2019.
- 735 Holland, P. R., O'Connor, G. K., Bracegirdle, T. J., Dutrieux, P., Naughten, K. A., Steig, E. J., Schneider, D. P., Jenkins, A., and Smith, J. A.: Anthropogenic and internal drivers of wind changes over the Amundsen Sea, West Antarctica, during the 20th and 21st centuries, *The Cryosphere*, 16, 5085-5105, 10.5194/tc-16-5085-2022, 2022.
- Jacobs, S. S., Hellmer, H. H., and Jenkins, A.: Antarctic ice sheet melting in the Southeast Pacific, *Geophysical Research Letters*, 23, 957-960, Doi 10.1029/96gl00723, 1996.
- Jenkins, A., Nicholls, K. W., and Corr, H. F. J.: Observation and Parameterization of Ablation at the Base of Ronne Ice Shelf, Antarctica, *Journal of Physical Oceanography*, 40, 2298-2312, <https://doi.org/10.1175/2010JPO4317.1>, 2010.
- 740 Jenkins, A., Shoosmith, D., Dutrieux, P., Jacobs, S., Kim, T. W., Lee, S. H., Ha, H. K., and Stammerjohn, S.: West Antarctic Ice Sheet retreat in the Amundsen Sea driven by decadal oceanic variability, *Nature Geoscience*, 11, 733-738, 10.1038/s41561-018-0207-4, 2018.
- Jordan, J. R., Holland, P. R., Goldberg, D., Snow, K., Arthern, R., Campin, J.-M., Heimbach, P., and Jenkins, A.: Ocean-Forced Ice-Shelf Thinning in a Synchronously Coupled Ice-Ocean Model, *Journal of Geophysical Research: Oceans*, 123, 864-882, <https://doi.org/10.1002/2017JC013251>, 2018.
- 745 Joughin, I., Smith, B. E., and Medley, B.: Marine Ice Sheet Collapse Potentially Under Way for the Thwaites Glacier Basin, West Antarctica, *Science*, 344, 735-738, 10.1126/science.1249055, 2014.
- Joughin, I., Shapero, D., Smith, B., Dutrieux, P., and Barham, M.: Ice-shelf retreat drives recent Pine Island Glacier speedup, *Science Advances*, 7, eabg3080, 10.1126/sciadv.abg3080, 2021.
- 750 Jourdain, N. C., Mathiot, P., Burgard, C., Caillet, J., and Kittel, C.: Ice Shelf Basal Melt Rates in the Amundsen Sea at the End of the 21st Century, *Geophysical Research Letters*, 49, e2022GL100629, <https://doi.org/10.1029/2022GL100629>, 2022.
- Jourdain, N. C., Mathiot, P., Merino, N., Durand, G., Le Sommer, J., Spence, P., Dutrieux, P., and Madec, G.: Ocean circulation and sea-ice thinning induced by melting ice shelves in the Amundsen Sea, *Journal of Geophysical Research: Oceans*, 122, 2550-2573, <https://doi.org/10.1002/2016JC012509>, 2017.
- 755 Kimura, S., Jenkins, A., Regan, H., Holland, P. R., Assmann, K. M., Whitt, D. B., Van Wessem, M., van de Berg, W. J., Reijmer, C. H., and Dutrieux, P.: Oceanographic Controls on the Variability of Ice-Shelf Basal Melting and Circulation of Glacial Meltwater in the Amundsen Sea Embayment, Antarctica, *Journal of Geophysical Research: Oceans*, 122, 10131-10155, <https://doi.org/10.1002/2017JC012926>, 2017.
- Lhermitte, S., Sun, S., Shuman, C., Wouters, B., Pattyn, F., Wuite, J., Berthier, E., and Nagler, T.: Damage accelerates ice shelf instability and mass loss in Amundsen Sea Embayment, *Proceedings of the National Academy of Sciences*, 117, 24735-24741, doi:10.1073/pnas.1912890117, 2020.
- 760 Marshall, J., Hill, C., Perelman, L., and Adcroft, A.: Hydrostatic, quasi-hydrostatic, and nonhydrostatic ocean modeling, *Journal of Geophysical Research: Oceans*, 102, 5733-5752, <https://doi.org/10.1029/96JC02776>, 1997.
- Morlighem, M.: MEaSUREs BedMachine Antarctica, Version 3 [dataset], <https://doi.org/10.5067/FPSU0V1MWUB6>, 2022.
- Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., Eisen, O., Ferraccioli, F., Forsberg, R., Fretwell, P., Goel, V., Greenbaum, J. S., Gudmundsson, H., Guo, J., Helm, V., Hofstede, C., Howat, I., Humbert, A., Jokat, W., Karlsson, N. B., Lee, W. S., Matsuoka, K., Millan, R., Mougino, J., Paden, J., Pattyn, F., Roberts, J., Rosier, S., Ruppel, A., Seroussi, H., Smith, E. C., Steinhage, D., Sun, B., Broeke, M. R. v. d., Ommen, T. D. v., Wessem, M. v., and Young, D. A.: Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet, *Nature Geoscience*, 13, 132-137, 10.1038/s41561-019-0510-8, 2020.
- 765 Mougino, J., Rignot, E., and Scheuchl, B.: Sustained increase in ice discharge from the Amundsen Sea Embayment, West Antarctica, from 1973 to 2013, *Geophysical Research Letters*, 41, 1576-1584, <https://doi.org/10.1002/2013GL059069>, 2014.
- Mougino, J., Scheuchl, B., and Rignot, E.: MEaSUREs Annual Antarctic Ice Velocity Maps, Version 1 [dataset], <https://doi.org/10.5067/9T4EPOXTJYW9>, 2017a.
- Mougino, J., Rignot, E., Scheuchl, B., and Millan, R.: Comprehensive Annual Ice Sheet Velocity Mapping Using Landsat-8, Sentinel-1, and RADARSAT-2 Data, 10.3390/rs9040364, 2017b.
- 770 Nakayama, Y., Cai, C., and Seroussi, H.: Impact of Subglacial Freshwater Discharge on Pine Island Ice Shelf, *Geophysical Research Letters*, 48, e2021GL093923, <https://doi.org/10.1029/2021GL093923>, 2021.
- Naughten, K. A., Holland, P. R., and De Rydt, J.: Unavoidable future increase in West Antarctic ice-shelf melting over the twenty-first century, *Nature Climate Change*, 13, 1222-1228, 10.1038/s41558-023-01818-x, 2023.
- 775 Naughten, K. A., De Rydt, J., Rosier, S. H. R., Jenkins, A., Holland, P. R., and Ridley, J. K.: Two-timescale response of a large Antarctic ice shelf to climate change, *Nature Communications*, 12, 1991, 10.1038/s41467-021-22259-0, 2021.
- 780

- Naughten, K. A., Holland, P. R., Dutrieux, P., Kimura, S., Bett, D. T., and Jenkins, A.: Simulated Twentieth-Century Ocean Warming in the Amundsen Sea, West Antarctica, *Geophysical Research Letters*, 49, e2021GL094566, <https://doi.org/10.1029/2021GL094566>, 2022.
- 785 Nias, I. J., Cornford, S. L., Edwards, T. L., Gourmelen, N., and Payne, A. J.: Assessing Uncertainty in the Dynamical Ice Response to Ocean Warming in the Amundsen Sea Embayment, West Antarctica, *Geophysical Research Letters*, 46, 11253-11260, <https://doi.org/10.1029/2019GL084941>, 2019.
- Otosaka, I. N., Shepherd, A., Ivins, E. R., Schlegel, N. J., Amory, C., van den Broeke, M. R., Horwath, M., Joughin, I., King, M. D., Krinner, G., Nowicki, S., Payne, A. J., Rignot, E., Scambos, T., Simon, K. M., Smith, B. E., Sørensen, L. S., Velicogna, I., Whitehouse, P. L., A. G., Agosta, C., Ahlstrøm, A. P., Blazquez, A., Colgan, W., Engdahl, M. E., Fettweis, X., Forsberg, R., Gallée, H., Gardner, A., Gilbert, L., Gourmelen, N., Groh, A., Gunter, B. C., Harig, C., Helm, V., Khan, S. A., Kittel, C., Konrad, H., Langen, P. L., Lecavalier, B. S., Liang, C. C., Loomis, B. D., McMillan, M., Melini, D., Mernild, S. H., Mottram, R., Mouginit, J., Nilsson, J., Noël, B., Pattle, M. E., Peltier, W. R., Pie, N., Roca, M., Sasgen, I., Save, H. V., Seo, K. W., Scheuchl, B., Schrama, E. J. O., Schröder, L., Simonsen, S. B., Slater, T., Spada, G., Sutterley, T. C., Vishwakarma, B. D., van Wessem, J. M., Wiese, D., van der Wal, W., and Wouters, B.: Mass balance of the Greenland and Antarctic ice sheets from 1992 to 2020, *Earth Syst. Sci. Data*, 15, 1597-1616, 10.5194/essd-15-1597-2023, 2023.
- 790 Pattyn, F.: Antarctic subglacial conditions inferred from a hybrid ice sheet/ice stream model, *Earth and Planetary Science Letters*, 295, 451-461, <https://doi.org/10.1016/j.epsl.2010.04.025>, 2010.
- 795 Reese, R., Levermann, A., Albrecht, T., Seroussi, H., and Winkelmann, R.: The role of history and strength of the oceanic forcing in sea level projections from Antarctica with the Parallel Ice Sheet Model, *The Cryosphere*, 14, 3097-3110, 10.5194/tc-14-3097-2020, 2020.
- Reese, R., Garbe, J., Hill, E. A., Urruty, B., Naughten, K. A., Gagliardini, O., Durand, G., Gillet-Chaulet, F., Gudmundsson, G. H., Chandler, D., Langebroek, P. M., and Winkelmann, R.: The stability of present-day Antarctic grounding lines – Part 2: Onset of irreversible retreat of Amundsen Sea glaciers under current climate on centennial timescales cannot be excluded, *The Cryosphere*, 17, 3761-3783, 10.5194/tc-17-3761-2023, 2023.
- 800 Schoof, C.: Ice sheet grounding line dynamics: Steady states, stability, and hysteresis, *Journal of Geophysical Research: Earth Surface*, 112, <https://doi.org/10.1029/2006JF000664>, 2007.
- Seroussi, H. and Morlighem, M.: Representation of basal melting at the grounding line in ice flow models, *The Cryosphere*, 12, 3085-3096, 10.5194/tc-12-3085-2018, 2018.
- 805 Seroussi, H., Nakayama, Y., Larour, E., Menemenlis, D., Morlighem, M., Rignot, E., and Khazendar, A.: Continued retreat of Thwaites Glacier, West Antarctica, controlled by bed topography and ocean circulation, *Geophysical Research Letters*, 44, 6191-6199, <https://doi.org/10.1002/2017GL072910>, 2017.
- Shepherd, A., Gilbert, L., Muir, A. S., Konrad, H., McMillan, M., Slater, T., Briggs, K. H., Sundal, A. V., Hogg, A. E., and Engdahl, M. E.: Trends in Antarctic Ice Sheet Elevation and Mass, *Geophysical Research Letters*, 46, 8174-8183, <https://doi.org/10.1029/2019GL082182>, 2019.
- 810 Smith, B., Fricker, H. A., Gardner, A. S., Medley, B., Nilsson, J., Paolo, F. S., Holschuh, N., Adusumilli, S., Brunt, K., Csatho, B., Harbeck, K., Markus, T., Neumann, T., Siegfried, M. R., and Zwally, H. J.: Pervasive ice sheet mass loss reflects competing ocean and atmosphere processes, *Science*, 368, 1239-1242, 10.1126/science.aaz5845, 2020.
- 815 St-Laurent, P., Klinck, J. M., and Dinniman, M. S.: Impact of local winter cooling on the melt of Pine Island Glacier, Antarctica, *Journal of Geophysical Research: Oceans*, 120, 6718-6732, <https://doi.org/10.1002/2015JC010709>, 2015.
- Thomas, R. H.: The Dynamics of Marine Ice Sheets, *Journal of Glaciology*, 24, 167-177, 10.3189/S0022143000014726, 1979.
- Walker, D. P., Brandon, M. A., Jenkins, A., Allen, J. T., Dowdeswell, J. A., and Evans, J.: Oceanic heat transport onto the Amundsen Sea shelf through a submarine glacial trough, *Geophysical Research Letters*, 34, Artn L02602, <https://doi.org/10.1029/2006gl028154>, 2007.
- 820 Webber, B. G. M., Heywood, K. J., Stevens, D. P., Dutrieux, P., Abrahamsen, E. P., Jenkins, A., Jacobs, S. S., Ha, H. K., Lee, S. H., and Kim, T. W.: Mechanisms driving variability in the ocean forcing of Pine Island Glacier, *Nature Communications*, 8, 14507, 10.1038/ncomms14507, 2017.
- Weertman, J.: The Theory of Glacier Sliding, *Journal of Glaciology*, 5, 287-303, 10.3189/S0022143000029038, 1964.
- 825 Weertman, J.: Stability of the Junction of an Ice Sheet and an Ice Shelf, *Journal of Glaciology*, 13, 3-11, 10.3189/S0022143000023327, 1974.
- Wild, C. T., Alley, K. E., Muto, A., Truffer, M., Scambos, T. A., and Pettit, E. C.: Weakening of the pinning point buttressing Thwaites Glacier, West Antarctica, *The Cryosphere*, 16, 397-417, 10.5194/tc-16-397-2022, 2022.
- Williams, C., Thodoroff, P., Arthern, R., Byrne, J., Hosking, J. S., Kaiser, M., Lawrence, N., and Kazlauskaitė, I.: Calculating exposure to extreme sea level risk will require high resolution ice sheet models, *Research Square* [<https://doi.org/10.21203/rs.3.rs-3405435/v1>], PREPRINT.
- 830 Yu, H., Rignot, E., Seroussi, H., and Morlighem, M.: Retreat of Thwaites Glacier, West Antarctica, over the next 100 years using various ice flow models, ice shelf melt scenarios and basal friction laws, *The Cryosphere*, 12, 3861-3876, 10.5194/tc-12-3861-2018, 2018.
- Zwally, H. J., Giovinetto, M. B., Matthew, A. B., and Jack, L. S.: Antarctic and Greenland Drainage Systems, GSFC Cryospheric Sciences Laboratory, at http://icesat4.gsfc.nasa.gov/cryo_data/ant_grm_drainage_systems.php, 2012.
- 835